The density of rational points on a certain singular cubic surface

T.D. Browning

Mathematical Institute, 24–29 St. Giles', Oxford OX1 3LB

browning@maths.ox.ac.uk

Abstract

We show that the number of non-trivial rational points of height at most B, which lie on the cubic surface $x_1x_2x_3 = x_4(x_1 + x_2 + x_3)^2$, has order of magnitude $B(\log B)^6$. This agrees with Manin's conjecture.

1 Introduction

The purpose of this paper is to investigate the distribution of rational points on the singular cubic surface $X \subset \mathbb{P}^3$, given by the equation

$$x_1 x_2 x_3 = x_4 (x_1 + x_2 + x_3)^2.$$

This surface has a unique singular point [0, 0, 0, 1] which is of type D_4 , and contains precisely 6 lines [3, Lemma 4]. These lines are all defined over \mathbb{Q} and are given by

 $x_i = x_4 = 0, \quad x_i = x_j + x_k = 0,$

for $\{i, j, k\} = \{1, 2, 3\}$. We shall denote by $U \subset X$ the open subset formed by deleting the lines from X.

Given a rational point $x = [x_1, x_2, x_3, x_4] \in \mathbb{P}^3(\mathbb{Q})$ such that x_1, x_2, x_3, x_4 are relatively prime integers, let $H(x) = \max_{1 \leq i \leq 4} |x_i|$ denote its anticanonical height, metrized by the choice of norm $\max_{1 \leq i \leq 4} |x_i|$. Then for any $B \geq 1$, we shall be concerned with estimating the quantity

$$N_{U,H}(B) = \#\{x \in U \cap \mathbb{P}^3(\mathbb{Q}) : H(x) \leqslant B\}.$$

Manin [8] has provided a very general conjecture concerning the distribution of rational points on Fano varieties. In our case it predicts that there exists a positive constant $c_{X,H}$ such that

$$N_{U,H}(B) \sim c_{X,H} B(\log B)^6$$

as $B \to \infty$. Here the exponent of log *B* is one less than the rank of $\operatorname{Pic}(X)$, where \tilde{X} denotes the minimal desingularisation of *X*. In fact this sort of asymptotic formula is conjectured to hold for any cubic surface with canonical singular locus. Although there has been increasing interest in Manin's conjecture for

cubic surfaces, it has only been completely settled in particularly simple cases such as the toric variety

$$x_1 x_2 x_3 = x_4^3.$$

Such results can be found in the work of Batyrev and Tschinkel [1], de la Bretèche [2], Chambert-Loir and Tschinkel [4], Fouvry [7], Heath-Brown and Moroz [13], and Salberger [17].

More recently, Heath-Brown [14] has established upper and lower bounds for the density of non-trivial rational points on the Cayley cubic surface

$$\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} + \frac{1}{x_4} = 0,$$

which agree with Manin's conjecture. This is a cubic surface containing four A_1 singularities, which is the maximal number of singularities that a non-ruled cubic surface can have. The principal tool in Heath-Brown's work is a passage to the universal torsor above the minimal desingularisation of the Cayley cubic. Originally introduced by Colliot-Thélène and Sansuc [5, 6] to aid in the study of the Hasse principle and Weak approximation, universal torsors were first used by Peyre [16] and Salberger [17] in the context of counting rational points of bounded height. After establishing a bijection between the rational points on the Cayley cubic and the integer points on the universal torsor, which in this setting is given explicitly by nine equations in thirteen variables, Heath-Brown proceeds by applying methods from the geometry of numbers to count integer solutions to certain ternary linear equations.

Our present work is largely inspired by Heath-Brown's treatment of the Cayley cubic surface. We are now in a position to state our main result.

Theorem. We have

$$B(\log B)^6 \ll N_{U,H}(B) \ll B(\log B)^6.$$
 (1.1)

Of the two bounds in our theorem, the lower bound $N_{U,H}(B) \gg B(\log B)^6$ is routine. It will follow from relatively minor adjustments to Heath-Brown's treatment of the Cayley cubic. This is in part due to the fact that both surfaces contain a pair of skew rational lines, which are crucial to the proof. Establishing the upper bound $N_{U,H}(B) \ll B(\log B)^6$ however, is by far the most challenging component of this paper. As in the case of the Cayley cubic, the proof has two fundamental ingredients. The first is a translation of the problem to the universal torsor above \tilde{X} , which in this setting has a natural embedding in \mathbb{A}^{10} , given by

$$s_0 s_1 s_2 s_3 u_1 u_2 u_3 = y_1 u_1 s_1^2 + y_2 u_2 s_2^2 + y_3 u_3 s_3^2.$$

$$(1.2)$$

This has been calculated by Hassett and Tschinkel [9, §4], although we shall present our own deduction of this equation in §4 below. The universal torsor can be thought of as serving to encode factorisation information about the integer solutions to the original equation. In practical terms, it allows us to work with a larger number of variables, all of which are smaller in modulus than the original variables x_1, x_2, x_3, x_4 .

The second main ingredient in our proof of the upper bound involves studying the distribution of integer solutions to the equation obtained by setting $s_0 = 0$ in (1.2). This is the focus of §3 and relies upon lattice methods to count integer solutions to ternary linear and quadratic forms. It seems worthwhile highlighting the fact that this need to consider the contribution from quadratic equations marks a significant departure from Heath-Brown's treatment of the Cayley cubic. In particular, we shall need to pay careful attention to the fact that a given ternary quadratic form does not always have an integer solution.

Our work draws upon a diverse range of techniques. In addition to the geometry of numbers used to study linear and quadratic forms in §2.1, we make use of the large sieve inequality in §2.2, and real character sum estimates due to Heath-Brown and Pólya–Vinogradov in §3.

Acknowledgements. This problem was posed by Yuri Tschinkel during the course of the American Institute of Mathematics workshop "Rational and integral points on higher-dimensional varieties". The author is very grateful to Professor Heath-Brown and Professor Tschinkel for several useful conversations relating to the subject of this work, and to the anonymous referee for his careful reading of the manuscript and numerous pertinent suggestions. While working on this paper, the author was supported by EPSRC grant number GR/R93155/01.

2 Preliminary estimates

We begin by introducing some conventions regarding our choice of notation. Throughout this paper the letters i, j, k will denote generic distinct indices from the set $\{1, 2, 3\}$. We shall use \mathbb{N} to denote the set of positive integers, and for any $n \ge 2$ it will be convenient to let \mathbb{Z}^n denote the set of primitive vectors in \mathbb{Z}^n , where $\mathbf{v} \in \mathbb{Z}^n$ is said to be primitive if h.c.f. $(v_1, \ldots, v_n) = 1$. Similarly, we let N^n denote the set of primitive vectors in \mathbb{N}^n . Furthermore, we let \mathbb{Z}^n_* denote the subset of $\mathbf{v} \in \mathbb{Z}^n$ for which $v_1 \cdots v_n \neq 0$. Upon writing

$$F(\mathbf{x}) = x_1 x_2 x_3 - x_4 (x_1 + x_2 + x_3)^2,$$

it therefore follows that

$$N_{U,H}(B) = \frac{1}{2} \# \Big\{ \mathbf{x} \in Z_*^4 : \max_{1 \le i \le 4} |x_i| \le B, \ F(\mathbf{x}) = 0 \Big\},$$
(2.1)

since \mathbf{x} and $-\mathbf{x}$ represent the same point in \mathbb{P}^3 . It will be convenient to collect together some technical results that will be useful to us.

2.1 The geometry of numbers and ternary forms

Several of our arguments will involve estimating the number of primitive integer solutions to certain ternary homogeneous polynomial equations, which lie in lopsided regions. In the case of linear equations, such an estimate is provided by the following result of Heath-Brown [10, Lemma 3].

Lemma 1. Let $\mathbf{h} \in Z^3$ and let $W_i > 0$. Then the number of $\mathbf{w} \in Z^3$ for which $\sum_{i=1}^3 h_i w_i = 0$, and $|w_i| \leq W_i$, is

$$\leqslant 4 + 12\pi \frac{W_1 W_2 W_3}{\max|h_i|W_i|}$$

We shall also need a result which handles the corresponding problem for diagonal quadratic equations. For this we turn to the following result, in which $\omega(n)$ denotes the number of distinct prime factors of $n \in \mathbb{N}$.

Lemma 2. Let $\mathbf{g}, \mathbf{h} \in Z^3_*$, with $g_1g_2g_3$ square-free, and let $W_i > 0$. Then the number of $\mathbf{w} \in Z^3$ for which $\sum_{i=1}^3 g_i h_i w_i^2 = 0$, and $|w_i| \leq W_i$, is

$$\ll \left(1 + \sqrt{\frac{W_1 W_2 W_3 D_{\mathbf{g}, \mathbf{h}}^{3/2}}{|h_1 h_2 h_3|}}\right) 2^{\omega(h_1 h_2 h_3)},$$

where $D_{\mathbf{g},\mathbf{h}}$ is the product of highest common factors

h.c.f. (h_1h_2, h_1h_3, h_2h_3) h.c.f. (g_1, h_2h_3) h.c.f. (g_2, h_1h_3) h.c.f. (g_3, h_1h_2) .

Lemma 2 will follow from a rather straightforward modification to the proof of Heath-Brown's [12, Theorem 2]. In fact Heath-Brown establishes a version of Lemma 2 with $\mathbf{g} = (1, 1, 1)$ and $d_3(|h_1h_2h_3|)$ in place of $2^{\omega(h_1h_2h_3)}$, where $d_k(n)$ denotes the number of representations of n as a product of k positive integers, for any $k, n \in \mathbb{N}$. It is perhaps worth pointing out that whereas $d_3(n)$ has average order $\frac{1}{2}(\log n)^2$, the function $2^{\omega(n)}$ has average order $\zeta(2)^{-1}\log n$. This saving plays an important role in our work.

In order to prove Lemma 2 we recall that the original idea behind the proof of [12, Theorem 2] is to view the equation $\sum_{i=1}^{3} g_i h_i w_i^2 = 0$ as a collection of lattice conditions upon the solutions $\mathbf{w} \in \mathbb{Z}^3$. Let p be any prime divisor of $h_1 h_2 h_3$, and assume without loss of generality that

$$0 \leqslant \nu_p(h_1) \leqslant \nu_p(h_2) \leqslant \nu_p(h_3),$$

where $\nu_p(n)$ denotes the *p*-adic order of any $n \in \mathbb{N}$. In particular it follows that $\nu_p(h_1) = 0$, since **h** is primitive. We shall only consider here the case in which *p* is an odd prime. The case p = 2 is handled along similar lines. Since $g_1g_2g_3$ is square-free, we may write

$$g_1 = p^{\alpha_1}g'_1, \quad g_2 = p^{\alpha_2}g'_2, \quad g_3 = p^{\alpha_3}g'_3,$$

for $(\alpha_1, \alpha_2, \alpha_3) \in \{(0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ and $p \nmid g'_i$. Similarly, we write

$$h_2 = p^{\beta_2} h'_2, \quad h_3 = p^{\beta_3} h'_3,$$

for $p \nmid h'_2 h'_3$ and $\beta_3 \ge \beta_2 \ge 1$. Then one proceeds by considering solutions to the equation

$$p^{\alpha_1}g'_1h_1w_1^2 + p^{\alpha_2+\beta_2}g'_2h'_2w_2^2 + p^{\alpha_3+\beta_3}g'_3h'_3w_3^2 = 0.$$
(2.2)

Suppose for the moment that we are examining solutions $(u, v, w) \in \mathbb{Z}^3$ to the equation

$$au^{2} + p^{\sigma}bv^{2} + p^{\tau}cw^{2} = 0, \qquad (2.3)$$

for $0 \leq \sigma \leq \tau$ and $p \nmid abc$. Then arguing along similar lines to the proof of [12, Theorem 2], we sketch how this implies that (u, v, w) lies on one of at most 2 sublattices of \mathbb{Z}^3 , each of determinant $p^{\delta(\sigma,\tau)}$, where

$$\delta(\sigma,\tau) = \begin{cases} (\sigma+\tau) - 3\sigma/2, & \sigma \text{ even,} \\ (\sigma+\tau) - [3\sigma/2] + 1, & \sigma \text{ odd.} \end{cases}$$
(2.4)

Suppose first that $\sigma = 2s$ is even. Then (2.3) implies that $p^s \mid u$. By writing $u = p^s u'$, and considering the corresponding congruence $au'^2 + bv^2 \equiv 0 \pmod{p^{\tau-\sigma}}$, we therefore deduce that (u, v, w) lies on one of at most 2 integer lattices, each of determinant $p^{s+\tau-\sigma} = p^{\delta(\sigma,\tau)}$. Suppose now that $\sigma = 2s+1$ is odd. In view of (2.3) we may again write $u = p^s u'$, and consider the corresponding congruence $au'^2 + pbv^2 \equiv 0 \pmod{p^{\tau-2s}}$. Since $\tau - 2s \ge 1$, we may clearly write u' = pu'', and so consider solutions to the equation

$$pa(u'')^2 + bv^2 + p^{\tau - \sigma}cw^2 = 0.$$

Now either $\tau - \sigma = 0$, or else we may write v = pv' and consider the equation $a(u'')^2 + pbv'^2 + p^{\tau - \sigma - 1}cw^2 = 0$. In the former case we conclude that (v, w) lies on one of at most two integer sublattices of determinant p. But then (u, v, w) lies on one of at most 2 integer lattices, each of determinant $p^{s+2} = p^{\delta(\sigma,\tau)}$. In the latter case we have $\tau - \sigma \ge 1$, and we proceed inductively. Thus either $\tau - \sigma = 1$, in which case we deduce that (u, v, w) lies on one of 2 integer lattices of determinant $p^{s+3} = p^{\delta(\sigma,\tau)}$, or else $\tau - \sigma \ge 2$ and we can repeat the process. Since this process clearly terminates we therefore deduce that whenever σ is odd, (u, v, w) lies on one of at most 2 integer lattices, each of determinant $p^{\delta(\sigma,\tau)}$.

Returning to (2.2), our goal is to show that \mathbf{w} lies on one of at most 2 integer sublattices of \mathbb{Z}^3 , each of determinant

$$\geq p^{\beta_2 + \beta_3 - [3(\alpha_1 + \alpha_2 + \alpha_3 + \beta_2)/2]}.$$
(2.5)

In view of the existing proof of [12, Theorem 2], this will suffice to establish Lemma 2 since the inequalities $\beta_3 \ge \beta_2 \ge 1$ imply that

$$\nu_p(h_1h_2h_2) = \beta_2 + \beta_3, \quad \nu_p(D_{\mathbf{g},\mathbf{h}}) = \alpha_1 + \alpha_2 + \alpha_3 + \beta_2.$$

Suppose first that $(\alpha_1, \alpha_2) = (0, 0)$. Then our work above shows that **w** lies on one of at most 2 integer sublattices of \mathbb{Z}^3 , each of determinant

$$p^{\delta(\beta_2,\alpha_3+\beta_3)} \ge p^{\delta(\beta_2,\beta_3)}$$

This is plainly satisfactory for (2.5), by (2.4). Suppose now that $(\alpha_1, \alpha_2, \alpha_3) = (1, 0, 0)$. If $\beta_2 = 0$, then it is not hard to conclude that **w** lies on one of at most 2 lattices of determinant p^{β_3} , which is also satisfactory. If now $\beta_2 \ge 1$ we obtain an equation of the shape (2.3), with $\sigma = \beta_2 - 1$ and $\tau = \beta_3 - 1$. Thus we obtain at most 2 integer lattices, each of determinant $p^{\delta(\beta_2-1,\beta_3-1)}$. It is easily checked that this quantity is bounded below by (2.5). Finally we suppose that $(\alpha_1, \alpha_2, \alpha_3) = (0, 1, 0)$. In this case we again obtain an equation of the form (2.3). Suppose first that $\beta_2 < \beta_3$, so that we may take $\sigma = \beta_2 + 1$ and $\tau = \beta_3$ in (2.3). But then it easily follows that **w** lies on one of at most 2 integer lattices, each of determinant

$$p^{\delta(\beta_2+1,\beta_3)} \ge p^{\beta_2+\beta_3-[3(1+\beta_2)/2]}.$$

Alternatively, if $\beta_2 = \beta_3 = \beta$ say, then we take $\sigma = \beta$ and $\tau = \beta + 1$ in (2.3), thereby deducing that **w** lies on one of at most 2 lattices, each of determinant

$$p^{\delta(\beta,\beta+1)} \ge p^{2\beta - [3(1+\beta)/2]}.$$

This completes the proof of Lemma 2.

2.2 The large sieve and quadratic forms

In addition to considering the density of integer solutions to diagonal quadratic equations, as in the previous section, we shall also need to consider how often such an equation has at least one non-trivial integer solution. We begin by recalling the following version of the large sieve inequality [15].

Lemma 3. For any $N, Q \ge 1$, suppose that $T \subseteq \mathbb{Z} \cap [1, N]$, and that for every prime $p \le Q$ there exists $\tau(p) \in [0, p)$ such that the image of T in $\mathbb{Z}/p\mathbb{Z}$ has $p - \tau(p)$ elements. Then we have

$$\#T \ll \frac{N+Q^2}{L(Q)},$$

where

$$L(Q) = \sum_{q \leq Q} |\mu(q)| \prod_{p|q} \frac{\tau(p)}{p - \tau(p)}.$$

For any $Y \ge 1$ and non-zero $a, b, c \in \mathbb{Z}$, let T(Y; a, b, c) be the set of $y \in \mathbb{Z}$ such that $0 < |y| \le Y$, and the equation

$$ax_1^2 + bx_2^2 + cyx_3^2 = 0$$

has a non-zero solution $\mathbf{x} \in \mathbb{Z}^3$ with h.c.f. $(x_i, x_j) = 1$. A necessary condition for y to belong to T(Y; a, b, c) is that

$$\left(\frac{-ab}{p}\right) \neq -1 \quad \text{for all odd } p \mid cy,$$
 (2.6)

where $(\frac{n}{p})$ is the Legendre symbol for any $n \in \mathbb{Z}$ and prime p. Similarly we may assume that

$$\left(\frac{-acy}{p}\right) \neq -1 \quad \text{for all odd } p \mid b, \quad \left(\frac{-bcy}{p}\right) \neq -1 \quad \text{for all odd } p \mid a.$$
 (2.7)

In particular we take T(Y; a, b, c) to be empty if $\left(\frac{-ab}{p}\right) = -1$ for any odd prime $p \mid c$. For any $n \in \mathbb{N}$ and $z \ge 2$, let

$$\varphi(n;z) = \prod_{\substack{p \le z \\ p|n}} \frac{2}{(1+1/p)^2}.$$
(2.8)

Then with this definition in mind, the remainder of this section is devoted to the proof of the following result.

Lemma 4. For any $\varepsilon > 0$, we have

$$\#T(Y; a, b, c) \ll_{\varepsilon} \frac{(\operatorname{h.c.f.}(a, b)\operatorname{h.c.f.}(a, c)\operatorname{h.c.f.}(b, c))^{\varepsilon}}{\varphi(ab; Y^{1/2})} \frac{Y}{\log Y} \sum_{x \in \mathcal{X}} \frac{|\mu(x)|}{x},$$

where

$$\mathcal{X} = \mathcal{X}(Y; a, b) = \left\{ x \in \mathbb{N} : x \leqslant Y^{1/2}, \ \left(\frac{-ab}{p}\right) = 1 \text{ for all odd } p \mid x \right\}.$$

It is interesting to compare Lemma 4 with work of Serre [18]. In the simpler setting a = b = c = 1, he establishes the upper bound

$$#T(Y; 1, 1, 1) \ll Y(\log Y)^{-1/2}.$$

By an elementary consideration of the Dirichlet series $L(1, \chi)$, where χ is the non-principal character modulo 4, one may retrieve this bound from Lemma 4.

In order to establish Lemma 4 we shall employ Lemma 3. Let p be an odd prime. Then we must calculate the size of the image T_p of T(Y; a, b, c) in $\mathbb{Z}/p\mathbb{Z}$. Suppose first that $p \nmid abc$. Then y = 0 is contained in T_p if and only if

$$\left(\frac{-ab}{p}\right) = 1,\tag{2.9}$$

by (2.6). Now the congruence $ax_1^2 + bx_2^2 + cyx_3^2 \equiv 0 \pmod{p}$ is always soluble whenever $p \nmid abcy$, by the Chevalley–Warning theorem. Hence we conclude that

$$#T_p = \begin{cases} p-1, & \left(\frac{-ab}{p}\right) = -1, \\ p, & \text{otherwise,} \end{cases}$$

whenever $p \nmid abc$. Turning to the size of T_p in the case $p \mid abc$, we suppose first that $p \mid a$ and $p \nmid bc$. Either y = 0 or $y \neq 0$ and (2.7) implies that

$$\left(\frac{-bcy}{p}\right) = 1.$$

For fixed values of b, c there are clearly $\frac{1}{2}(p+1)$ possible values of y over all. Similarly one finds that $\#T_p = \frac{1}{2}(p+1)$ if $p \mid b$ and $p \nmid ac$. If $p \mid c$ and $p \nmid ab$ however, then $\#T_p = p$ since (2.9) must automatically hold for any such p. Finally if $p \mid \text{h.c.f.}(a, b)\text{h.c.f.}(a, c)\text{h.c.f.}(b, c)$, then $\#T_p = p$.

Taking $\tau(p)$ to be $p - \#T_p$, we have therefore shown that

$$\tau(p) = \begin{cases} \frac{1}{2}(p-1), & p \mid a, p \nmid bc, \\ \frac{1}{2}(p-1), & p \mid b, p \nmid ac, \\ 1, & p \nmid abc, (\frac{-ab}{p}) = -1, \\ 0, & \text{otherwise.} \end{cases}$$

Inserting this into Lemma 3, we deduce that

$$\#T(Y;a,b,c) \ll \frac{Y+Q^2}{L(Q)}$$
(2.10)

for any $Q \ge 1$, where

$$\begin{split} L(Q) &= \sum_{q \leqslant Q} |\mu(q)| \prod_{p|q} \frac{\tau(p)}{p - \tau(p)} \\ &= \prod_{p \leqslant Q} \left(1 + \frac{\tau(p)}{p - \tau(p)} \right) \\ &= \frac{1}{f(Q; a, b, c)} \prod_{\substack{p \leqslant Q \\ p \mid ab}} \left(1 + \frac{p - 1}{p + 1} \right) \prod_{\substack{p \leqslant Q, \ p \nmid abc \\ (\frac{-ab}{p}) = -1}} \left(1 + \frac{1}{p - 1} \right). \end{split}$$

Here f(Q; a, b, c) is given by the product

$$\prod_{\substack{p \leq Q\\ p \mid \text{h.c.f.}(a,b) \text{h.c.f.}(a,c) \text{h.c.f.}(b,c)}} \left(1 + \frac{p-1}{p+1}\right).$$

In particular we clearly have

$$f(Q; a, b, c) \ll_{\varepsilon} (\text{h.c.f.}(a, b)\text{h.c.f.}(a, c)\text{h.c.f.}(b, c))^{\varepsilon}, \tag{2.11}$$

Recall that $\left(\frac{-ab}{p}\right) \neq -1$ for any odd prime $p \mid c$. Hence Merten's formula implies that

$$\prod_{\substack{p \leqslant Q, \ p \nmid abc \\ (\frac{-ab}{p}) = -1}} \left(1 + \frac{1}{p-1}\right) \geqslant \prod_{\substack{p \leqslant Q \\ (\frac{-ab}{p}) = -1}} \left(1 + \frac{1}{p}\right) \gg \log Q \prod_{\substack{p \leqslant Q \\ (\frac{-ab}{p}) \neq -1}} \left(1 + \frac{1}{p}\right)^{-1}.$$

Moreover, it is not hard to see that

$$\prod_{\substack{p \leqslant Q \\ p \mid ab}} \left(1 + \frac{p-1}{p+1} \right) \left(1 + \frac{1}{p} \right)^{-1} = \prod_{\substack{p \leqslant Q \\ p \mid ab}} \frac{2}{(1+1/p)^2} = \varphi(ab;Q),$$

in the notation of (2.8). Hence we have the lower bound

$$L(Q) \gg \frac{\varphi(ab;Q)\log Q}{f(Q;a,b,c)} \prod_{\substack{p \leqslant Q \\ (\frac{-ab}{p})=1}} \left(1 + \frac{1}{p}\right)^{-1}.$$

Upon inserting this into (2.10), with the choice $Q = Y^{1/2}$, we conclude that

$$\#T(Y;a,b,c) \ll \frac{f(Y^{1/2};a,b,c)}{\varphi(ab;Y^{1/2})} \frac{Y}{\log Y} \prod_{\substack{p \leqslant Y^{1/2} \\ (\frac{-ab}{p}) = 1}} \left(1 + \frac{1}{p}\right).$$

We complete the proof of Lemma 4 via an application of the upper bound (2.11).

The equation $a_1b_1c_1^2 + a_2b_2c_2^2 + a_3b_3c_3^2 = 0$ 3

The purpose of this section is to bring together the results in $\S2.1$ and $\S2.2$, in order to make a study of the density of integer solutions to the equation

$$a_1b_1c_1^2 + a_2b_2c_2^2 + a_3b_3c_3^2 = 0, (3.1)$$

that appears in the title. Thus for any $A_i, B_i, C_i \ge 1$ we let $\mathcal{M}(A_i, B_i, C_i)$ denote the number of $\mathbf{a}, \mathbf{b}, \mathbf{c} \in Z^3_*$ such that (3.1) holds and

$$|a_i| \leq A_i, \quad |b_i| \leq B_i, \quad |c_i| \leq C_i,$$

with

h.c.f.
$$(a_i, c_j) = \text{h.c.f.}(c_i, c_j) = 1$$
 (3.2)

and

$$|\mu(a_1a_2a_3)| = 1$$
, h.c.f. $(a_i, b_j, b_k) = 1$. (3.3)

It will be convenient to set

$$A = A_1 A_2 A_3, \quad B = B_1 B_2 B_3, \quad C = C_1 C_2 C_3.$$

With this notation in mind, the bulk of the work in this section will be taken up with establishing the following result.

Proposition 1. For any $\varepsilon > 0$, we have

$$\mathcal{M}(A_i, B_i, C_i) \ll_{\varepsilon} A^{2/3} B^{2/3} C^{1/3} + \sigma \tau A B^{1/2} C^{1/2},$$

where

$$\sigma = 1 + \frac{\min\{A, B\}^{\varepsilon}}{\min\{B_i B_j\}^{3/32 - \varepsilon}}, \quad \tau = 1 + \frac{\log B}{\min\{B_i B_j\}^{3/32 - \varepsilon}}.$$

It turns out that we shall need an alternative estimate for $\mathcal{M}(A_i, B_i, C_i)$ to handle the case in which B_1, B_2, B_3 have particularly awkward sizes. The following result will be established in §3.2.

Proposition 2. We have

$$\mathcal{M}(A_i, B_i, C_i) \ll AB_i B_j (C_k + C_i C_j A_k^{-1}) (\log AC)^2,$$

for any permutation $\{i, j, k\}$ of the set $\{1, 2, 3\}$.

Although we shall not need to do so here, it is worth pointing out that with more work it is possible to remove the term $(\log AC)^2$ from the statement of Proposition 2.

3.1 **Proof of Proposition 1**

In this section we shall prove Proposition 1. Our approach is based upon suitable applications of Lemma 1 and Lemma 2. Thus we begin by fixing choices of $\mathbf{a}, \mathbf{c} \in Z^3_*$, and count the corresponding number of $\mathbf{b} \in Z^3_*$ satisfying (3.1) and $|b_i| \leq B$. Applying Lemma 1 with

$$\mathbf{h} = (a_1 c_1^2, a_2 c_2^2, a_3 c_3^2),$$

we easily obtain the upper bound

$$\mathcal{M}(A_i, B_i, C_i) \ll \sum_{\mathbf{a}, \mathbf{c}} \left(1 + \frac{B}{\max |a_i| c_i^2 B_i} \right)$$
$$\leqslant \sum_{\mathbf{a}, \mathbf{c}} \left(1 + B^{2/3} |a_1 a_2 a_3 c_1^2 c_2^2 c_3^2|^{-1/3} \right)$$
$$\ll AC + A^{2/3} B^{2/3} C^{1/3}.$$

We shall use this bound whenever $C \leq B$, under which hypothesis the estimate in Proposition 1 clearly holds.

It remains to handle the case in which C > B. For this we fix choices of $\mathbf{a}, \mathbf{b} \in Z^3_*$ for which (3.3) holds, and count the corresponding number of $\mathbf{c} \in Z^3_*$

satisfying (3.1), (3.2) and $|c_i| \leq C_i$. Thus we are in a position to apply Lemma 2 with

$$\mathbf{g} = (a_1, a_2, a_3), \quad \mathbf{h} = (b_1, b_2, b_3).$$

In particular it follows from (3.1)—(3.3) that

$$\begin{aligned} D_{\mathbf{a},\mathbf{b}} &= \text{h.c.f.}(b_1b_2, b_1b_3, b_2b_3)\text{h.c.f.}(a_1, b_2b_3)\text{h.c.f.}(a_2, b_1b_3)\text{h.c.f.}(a_3, b_1b_2) \\ &= \text{h.c.f.}(b_1b_2, b_1b_3, b_2b_3) \\ &\leqslant \text{h.c.f.}(b_1, b_2)\text{h.c.f.}(b_1, b_3)\text{h.c.f.}(b_2, b_3) = E_{\mathbf{b}}, \end{aligned}$$

say. Moreover, since $|b_1b_2b_3| \leq B$ and C > B, we also have

$$\frac{C}{|b_1b_2b_3|} \ge \frac{C}{B} > 1.$$

Thus under the assumption C > B we may conclude from Lemma 2 that

$$\mathcal{M}(A_i, B_i, C_i) \ll C^{1/2} \sum_{\mathbf{a}, \mathbf{b}}^* \frac{E_{\mathbf{b}}^{3/4}}{|b_1 b_2 b_3|^{1/2}} 2^{\omega(b_1 b_2 b_3)},$$

where $\sum_{\mathbf{a},\mathbf{b}}^{*}$ indicates a summation over $\mathbf{a}, \mathbf{b} \in Z_{*}^{3}$ for which $|a_{i}| \leq A_{i}, |b_{i}| \leq B_{i}$, (3.3) holds, and the equation (3.1) has a solution $\mathbf{c} \in Z_{*}^{3}$ with (3.2) holding.

In order to handle the term $E_{\mathbf{b}}^{3/4}$ in our estimate for $\mathcal{M}(A_i, B_i, C_i)$, we write

$$b_i = h_{ij} h_{ik} b'_i,$$

for fixed $h_{12}, h_{13}, h_{23} \in \mathbb{N}$ such that $h_{ij} = h_{ji}$. Then

$$|b_i'| \leqslant \frac{B_i}{h_{ij}h_{ik}}.\tag{3.4}$$

Since **b** is primitive, it follows that h.c.f. $(h_{ij}, h_{ik}) = 1$. Moreover, for fixed values of h_{12}, h_{13}, h_{23} , it suffices to sum over **a**, **b**' $\in Z^3_*$ for which

h.c.f.
$$(h_{ij}, a_k b'_k) = \text{h.c.f.}(b'_i, b'_j) = 1,$$
 (3.5)

by (3.3) and the fact that **b** is primitive. With this change of variables, the equation (3.1) clearly becomes

$$\beta_1 b_1' c_1^2 + \beta_2 b_2' c_2^2 + \beta_3 b_3' c_3^2 = 0, \qquad (3.6)$$

where we have written

$$\beta_i = a_i h_{ij} h_{ik}$$

for fixed values of a_i, h_{ij} . We shall need to record the equality

$$h.c.f.(\beta_i b'_i, \beta_j b'_j) = h_{ij}, \qquad (3.7)$$

which easily follows from combining the coprimality conditions (3.2),(3.3),(3.5) and h.c.f. $(h_{ij}, h_{ik}) = 1$, with the equation (3.6).

Let $h = h_{12}h_{13}h_{23}$. Then upon collecting our work together, we have therefore established the upper bound

$$\mathcal{M}(A_i, B_i, C_i) \ll C^{1/2} \sum_{h_{ij}} \sum_{\mathbf{a}} \sum_{\mathbf{b}' \in \mathcal{B}} \frac{2^{\omega(h^2 b_1' b_2' b_3')}}{h^{1/4} |b_1' b_2' b_3'|^{1/2}},$$

where $\mathcal{B} = \mathcal{B}(B_i; h_{ij}, a_i)$ denotes the set of $\mathbf{b}' \in Z^3_*$ with pairwise coprime components, for which (3.4) and (3.7) hold, and (3.6) has a solution $\mathbf{c} \in Z^3_*$ with h.c.f. $(c_i, c_j) = 1$. Using the trivial upper bound $2^{\omega(n)} \ll n^{1/32}$, we obtain

$$\mathcal{M}(A_i, B_i, C_i) \ll_{\varepsilon} C^{1/2} \sum_{h_{ij}} h^{-7/32} \sum_{\mathbf{a}} \sum_{\mathbf{b}' \in \mathcal{B}} \frac{2^{\omega(b_1' b_2' b_3')}}{|b_1' b_2' b_3'|^{1/2}}.$$
 (3.8)

Write

$$B_i' = \frac{B_i}{h_{ij}h_{ik}} \tag{3.9}$$

in (3.4), and suppose without loss of generality that $B'_1 \leq B'_2 \leq B'_3$. Then we proceed by using Lemma 4 to establish the following result.

Lemma 5. Let $\varepsilon > 0$. Then we have

$$\sum_{\mathbf{b}' \in \mathcal{B}} 2^{\omega(b_1'b_2'b_3')} \ll_{\varepsilon} h^{1/32} B_1' B_2' B_3' \Big(1 + \frac{\min\{A, B\}^{\varepsilon} + \log B}{(B_1'B_2')^{1/2 - \varepsilon}} \Big).$$

Before beginning the task of proving Lemma 5 we first show how it is sufficient to complete the proof of Proposition 1. Still under the assumption that $B'_1 \leq B'_2 \leq B'_3$, we use partial summation in (3.8). Thus it follows from Lemma 5 that

$$\mathcal{M}(A_i, B_i, C_i) \ll_{\varepsilon} A C^{1/2} \sum_{h_{ij}} \frac{(B_1' B_2' B_3')^{1/2}}{h^{3/16}} \Big(1 + \frac{\min\{A, B\}^{\varepsilon} + \log B}{(B_1' B_2')^{1/2 - \varepsilon}} \Big),$$

for any $\varepsilon > 0$. On recalling the definition (3.9) of B'_i , we see that

$$\sum_{h_{ij}} \frac{(B_1' B_2' B_3')^{1/2}}{h^{3/16}} = B^{1/2} \sum_{h_{ij}} \frac{1}{h^{19/16}} \ll B^{1/2}.$$

Similarly, using the inequalities

$$h_{12} \leq (B_1 B_2)^{1/2}, \quad h_{13} \leq B_1/h_{12}, \quad h_{23} \leq B_2/h_{12},$$

we find that

$$\begin{split} \sum_{h_{ij}} h^{-3/16} (B_1' B_2')^{\varepsilon} B_3'^{1/2} \ll_{\varepsilon} (B_1 B_2)^{\varepsilon} B_3^{1/2} \sum_{h_{ij}} h_{12}^{-3/16} h_{13}^{-11/16} h_{23}^{-11/16} \\ \ll_{\varepsilon} (B_1 B_2)^{5/16+\varepsilon} B_3^{1/2} \sum_{h_{12}} h_{12}^{-13/16} \\ \ll_{\varepsilon} (B_1 B_2)^{13/32+\varepsilon} B_3^{1/2} \ll B^{1/2}. \end{split}$$

It therefore follows that the estimate in Proposition 1 holds in the case C > B, and so holds unconditionally.

Thus in order to complete the proof of Proposition 1 it suffices to establish Lemma 5. In doing so our first step is to note that $2^{\omega(n)} \leq d(n)$ for any $n \in \mathbb{N}$, where d(n) is the ordinary divisor function. It follows that

$$\begin{split} \sum_{\mathbf{b}' \in \mathcal{B}} 2^{\omega(b_1'b_2'b_3')} &= \sum_{|b_1'| \leqslant B_1'} \sum_{|b_2'| \leqslant B_2'} 2^{\omega(b_1'b_2')} \sum_{b_3': \mathbf{b}' \in \mathcal{B}} \sum_{x|b_3'} 1 \\ &\leqslant \sum_{b_1', b_2'} 2^{\omega(b_1'b_2')} \sum_{|x| \leqslant B_3'^{1/2}} \#T(B_3'/|x|; \beta_1 b_1', \beta_2 b_2', \beta_3 x), \end{split}$$

in the notation of Lemma 4. Here the summation over $|x| \leq {B'_3}^{1/2}$ is only over x such that

$$\left(\frac{-\beta_1\beta_2b_1'b_2'}{p}\right) \neq -1 \quad \text{for all odd } p \mid x, \tag{3.10}$$

and

h.c.f.
$$(\beta_1 \beta_2 b'_1 b'_2, x) \mid h.$$
 (3.11)

This latter condition follows from (3.7) and the fact that x is a divisor of b'_3 . Hence (3.7) and Lemma 4 yield the upper bound

$$\#T(B'_3/|x|;\beta_1b'_1,\beta_2b'_2,\beta_3x) \ll_{\varepsilon} \frac{h^{\varepsilon}B'_3}{\varphi(\beta_1\beta_2b'_1b'_2;(B'_3/|x|)^{1/2})|x|\log B'_3} \sum_{y\in\mathcal{Y}} \frac{1}{y},$$

for any $\varepsilon > 0$, where \mathcal{Y} denotes the set

$$\mathcal{Y}(B_3';\beta_1b_1',\beta_2b_2') = \Big\{ y \in \mathbb{N} : y \leqslant {B_3'}^{1/2}, \ \left(\frac{-\beta_1\beta_2b_1'b_2'}{p}\right) = 1 \text{ for all odd } p \mid y \Big\}.$$

Here we have used the trivial inequality $|\mu(n)| \leq 1$ for any $n \in \mathbb{N}$. For any $n \in \mathbb{N}$ and $z \geq 2$, recall the definition (2.8) of the function $\varphi(n; z)$, and let $\omega(n; z)$ denote the number of distinct prime factors $p \mid n$ such that $p \leq z$. Then since the inequality $|x| \leq B'_3^{1/2}$ implies that $B'_3/|x| \geq B'_3^{1/2}$, it easily follows that

$$\begin{aligned} \varphi(\beta_1\beta_2b_1'b_2';(B_3'/|x|)^{1/2}) &\geqslant \prod_{\substack{p \leqslant B_3'^{1/4} \\ p|b_1'b_2'}} \frac{2}{(1+1/p)^2} \prod_{\substack{p \leqslant B_3'^{1/4} \\ p|\beta_1\beta_2,p \nmid b_1'b_2'}} \frac{2}{(1+1/p)^2} \\ &\geqslant \frac{2^{\omega(b_1'b_2';B_3'^{1/4})}}{\vartheta(b_1'b_2')^2}, \end{aligned}$$

where

$$\vartheta(n) = \prod_{p|n} \left(1 + \frac{1}{p} \right), \tag{3.12}$$

At this point it is convenient to collect together some basic facts about the arithmetic function ϑ .

Lemma 6. Let $m, n \in \mathbb{N}$ and let x > 1. Then we have

- (i) $\vartheta(n) = \sum_{d|n} |\mu(d)| d^{-1}$.
- (*ii*) $\vartheta(mn) \leq \vartheta(m)\vartheta(n)$.
- (iii) $\sum_{n \le x} \vartheta(n)^2 \ll x.$

Proof. The first two parts of the lemma are self-evident, and so it remains to establish part (iii). For this we employ part (i) to deduce that

$$\sum_{n \leqslant x} \vartheta(n)^2 = \sum_{n \leqslant x} \left(\sum_{d|n} \frac{|\mu(d)|}{d} \right)^2 = \sum_{n \leqslant x} \sum_{[d_1, d_2]|n} \frac{|\mu(d_1)\mu(d_2)|}{d_1 d_2},$$

where $[d_1, d_2] = d_1 d_2 / \text{h.c.f.}(d_1, d_2)$ denotes the least common multiple of d_1 and d_2 . On writing $n = [d_1, d_2]e$ we deduce that

$$\sum_{n \leqslant x} \vartheta(n)^2 \leqslant \sum_{d_1, d_2 = 1}^{\infty} \sum_{e \leqslant x/[d_1, d_2]} \frac{1}{d_1 d_2} \leqslant x \sum_{d_1, d_2 = 1}^{\infty} \frac{\text{h.c.f.}(d_1, d_2)}{d_1^2 d_2^2} \ll x,$$

since

$$\sum_{d_1,d_2=1}^{\infty} \frac{\text{h.c.f.}(d_1,d_2)}{d_1^2 d_2^2} \leqslant \sum_{d_1,d_2=1}^{\infty} \sum_{k|d_1,d_2} \frac{k}{d_1^2 d_2^2} \leqslant \sum_{k,d_1',d_2'=1}^{\infty} \frac{1}{k^3 {d_1'}^2 {d_2'}^2} \ll 1.$$

This completes the proof of Lemma 6.

Suppose now that p is any prime divisor of $b'_1b'_2$, with $p > {B'_3}^{1/4}$. Then since $|b'_1b'_2| \leq B'_1B'_2 \leq {B'_3}^2$, there can only be O(1) such distinct prime divisors. Moreover, we clearly have

$$\sum_{\substack{x \in B_3^{\prime 1/2} \\ (3.10), (3.11) \text{ hold}}} \frac{1}{x} \ll_{\varepsilon} h^{\varepsilon} \sum_{y \in \mathcal{Y}} \frac{1}{y},$$

with \mathcal{Y} as above. Combining all of this together therefore leads to the estimate

$$\sum_{\mathbf{b}'\in\mathcal{B}} 2^{\omega(b_1'b_2'b_3')} \ll_{\varepsilon} \frac{h^{\varepsilon}B_3'}{\log B_3'} \sum_{|b_1'|\leqslant B_1'} \sum_{|b_2'|\leqslant B_2'} \vartheta(b_1')^2 \vartheta(b_2')^2 \Big(\sum_{y\in\mathcal{Y}} \frac{1}{y}\Big)^2$$
$$\ll_{\varepsilon} \frac{h^{\varepsilon}B_3'}{\log B_3'} \sum_{|b_1'|\leqslant B_1'} \sum_{|b_2'|\leqslant B_2'} \vartheta(b_1')^2 \vartheta(b_2')^2 \sum_{y_1,y_2\in\mathcal{Y}} \frac{1}{y_1y_2}.$$

For any $n \in \mathbb{N}$ and $z \ge 2$, let $d(n; z) = \#\{a, b \leqslant z : ab = n\}$. Then we have

$$d(n;z) \leqslant d(n) \leqslant 2^{\Omega(n)},$$

where $\Omega(n)$ is the total number of prime factors of any $n \in \mathbb{N}$. On writing $\mathcal{Y}' = \mathcal{Y}(B'_3^2; \beta_1 b'_1, \beta_2 b'_2)$, it therefore follows that

$$\sum_{y_1,y_2\in\mathcal{Y}}\frac{1}{y_1y_2} = \sum_{y\in\mathcal{Y}'}\frac{d(y;B_3'^{1/2})}{y} \leqslant \sum_{y\in\mathcal{Y}'}\frac{2^{\Omega(y)}}{y},$$

for fixed choices of $\beta_1, \beta_2, b'_1, b'_2$.

In order to proceed further, we observe that

$$1 + \left(\frac{m}{p}\right) = \begin{cases} 2, & \left(\frac{m}{p}\right) = 1, \\ 0, & \text{otherwise,} \end{cases}$$

for any integer m not divisible by the odd prime p. Thus we have the estimate

$$\sum_{\mathbf{b}' \in \mathcal{B}} 2^{\omega(b_1'b_2'b_3')} \ll_{\varepsilon} \frac{h^{\varepsilon}B_3'}{\log B_3'} \sum_{b_1', b_2'} \vartheta(b_1')^2 \vartheta(b_2')^2 \sum_{y \leqslant B_3'} \frac{1}{y} \prod_{p|y} \left(1 + \left(\frac{-\beta_1 \beta_2 b_1' b_2'}{p}\right) \right)$$
$$\ll_{\varepsilon} \frac{h^{\varepsilon}B_3'}{\log B_3'} \sum_{b_1', b_2'} \vartheta(b_1')^2 \vartheta(b_2')^2 \sum_{y \leqslant B_3'} \frac{1}{y} \sum_{d|y} \left(\frac{-\beta_1 \beta_2 b_1' b_2'}{d}\right), \quad (3.13)$$

where ϑ is given by (3.12) and $\left(\frac{-\beta_1\beta_2b'_1b'_2}{d}\right)$ is the Jacobi symbol for d. We begin by considering the contribution from the terms for which $d = k^2$ is a square. Writing $y = jk^2$, we obtain the contribution

$$\ll_{\varepsilon} \frac{h^{\varepsilon} B'_{3}}{\log B'_{3}} \sum_{b'_{1}, b'_{2}} \vartheta(b'_{1})^{2} \vartheta(b'_{2})^{2} \sum_{k \leqslant B'_{3}^{1/2}} \sum_{j \leqslant B'_{3}/k^{2}} \frac{1}{jk^{2}} \\ \ll_{\varepsilon} \frac{h^{\varepsilon} B'_{3}}{\log B'_{3}} \sum_{j \leqslant B'_{3}} \frac{1}{j} \sum_{b'_{1}, b'_{2}} \vartheta(b'_{1})^{2} \vartheta(b'_{2})^{2} \\ \ll_{\varepsilon} h^{\varepsilon} B'_{1} B'_{2} B'_{3},$$

by Lemma 6. This is plainly satisfactory for Lemma 5, provided that $\varepsilon > 0$ is taken to be sufficiently small.

In order to handle the contribution from the remaining divisors, we define the characteristic function

$$\delta(n) = \begin{cases} 0, & n = k^2 \text{ for some } k \in \mathbb{N}, \\ 1, & \text{otherwise,} \end{cases}$$

for any $n \in \mathbb{N}$. Note that in particular we have $\delta(1) = 0$. Writing y = de, it easily follows that

$$\sum_{b_1',b_2'} \vartheta(b_1')^2 \vartheta(b_2')^2 \sum_{y \leqslant B_3'} \frac{1}{y} \sum_{\substack{d \mid y \\ \delta(d) = 1}} \left(\frac{-\beta_1 \beta_2 b_1' b_2'}{d}\right) = \sum_{e \leqslant B_3'} \frac{S_e}{e},$$
(3.14)

where

$$S_e = \sum_{d \leqslant B'_3/e} \frac{\delta(d)}{d} \sum_{|b'_1| \leqslant B'_1} \sum_{|b'_2| \leqslant B'_2} \vartheta(b'_1)^2 \vartheta(b'_2)^2 \Big(\frac{-\beta_1 \beta_2 b'_1 b'_2}{d}\Big).$$

Our next task is to establish the following inequality.

Lemma 7. Let $\varepsilon > 0$. Then we have

$$S_e \ll_{\varepsilon} B_1' B_2' + h^{\varepsilon} (B_1' B_2')^{1/2+\varepsilon} (\min\{A, B\}^{\varepsilon} + \log B).$$

Proof. Let us consider the contribution $S_e(D_1, D_2)$ to S_e from d contained in the interval $D_1 \ll d \ll D_2$, for various choices of

$$1 \leqslant D_1 \leqslant D_2 \leqslant B_3'/e. \tag{3.15}$$

Suppose that $N \in \mathbb{Z}$ is not a square, and that $D \ge 1$. Then an application of the Pólya–Vinogradov inequality yields

$$\sum_{d \leqslant D} \delta(d) \left(\frac{N}{d}\right) = \sum_{d \leqslant D} \left(\frac{N}{d}\right) - \sum_{\substack{d \leqslant D \\ d = k^2}} 1 \ll N^{1/2} \log N + D^{1/2}.$$

Thus for any D_1, D_2 in the range (3.15), we may combine partial summation with Lemma 6 to deduce that the contribution to $S_e(D_1, D_2)$ from those b'_1, b'_2 for which $-\beta_1\beta_2b_1'b_2'$ is not a square, is

$$\ll \sum_{b_1', b_2'} \delta(-\beta_1 \beta_2 b_1' b_2') \vartheta(b_1')^2 \vartheta(b_2')^2 \left| \sum_{D_1 \ll d \ll D_2} \frac{\delta(d)}{d} \left(\frac{-\beta_1 \beta_2 b_1' b_2'}{d} \right) \right| \\ \ll_{\varepsilon} \sum_{b_1', b_2'} \vartheta(b_1')^2 \vartheta(b_2')^2 \left(D_1^{-1/2} + D_1^{-1} |\beta_1 \beta_2 b_1' b_2'|^{1/2 + \varepsilon} \right) \\ \ll_{\varepsilon} B_1' B_2' + D_1^{-1} |\beta_1 \beta_2|^{1/2 + \varepsilon} (B_1' B_2')^{3/2 + \varepsilon} \\ \ll B_1' B_2' + D_1^{-1} |\beta_1 \beta_2|^{3/4} (B_1' B_2')^{7/4},$$

on taking $\varepsilon = 1/4$. Clearly there are $O((B'_1B'_2)^{1/2})$ values of b'_1, b'_2 for which $-\beta_1\beta_2b'_1b'_2$ is a square. In view of the trivial inequalities $\vartheta(n) \leq 2^{\omega(n)} \ll_{\varepsilon} n^{\varepsilon}$ for any $\varepsilon > 0$, it follows that the total contribution to $S_e(D_1, D_2)$ from such b'_1, b'_2 is

$$\ll_{\varepsilon} (B_1'B_2')^{1/2+\varepsilon}\log D_2$$

for any $\varepsilon > 0$. On noting that $B'_3/e \leq B_3 \leq B$, we have therefore established the bound

$$S_e(D_1, D_2) \ll_{\varepsilon} B_1' B_2' + D_1^{-1} |\beta_1 \beta_2|^{3/4} (B_1' B_2')^{7/4} + (B_1' B_2')^{1/2 + \varepsilon} \log B, \quad (3.16)$$

for any $\varepsilon > 0$ and any D_1, D_2 in the range (3.15).

In order to obtain a second estimate for $S_e(D_1, D_2)$ we begin by focusing on the contribution from small values of d. Let $[d_1, d_2]$ denote the least common multiple of d_1 and d_2 , as usual. Then it follows from a combination of the Pólya–Vinogradov inequality and the proof of Lemma 6 that

$$\begin{split} \left| \sum_{|b_2'| \leqslant B_2'} \vartheta(b_2')^2 \left(\frac{b_2'}{d}\right) \right| &= \left| \sum_{[d_1, d_2]|e| \leqslant B_2'} \frac{|\mu(d_1)\mu(d_2)|}{d_1 d_2} \left(\frac{[d_1, d_2]e}{d}\right) \right| \\ &\leqslant \sum_{d_1, d_2 \leqslant B_2'} \frac{1}{d_1 d_2} \left| \sum_{|e| \leqslant B_2'/[d_1, d_2]} \left(\frac{e}{d}\right) \right| \\ &\ll_{\varepsilon} d^{1/2+\varepsilon} \sum_{d_1, d_2 \leqslant B_2'} \frac{1}{d_1 d_2} \\ &\ll_{\varepsilon} d^{1/2+\varepsilon} (\log B_2')^2. \end{split}$$

Hence we see that

$$S_{e}(1, (B_{1}'B_{2}')^{1/2}) \ll \sum_{|b_{1}'| \leqslant B_{1}'} \vartheta(b_{1}')^{2} \sum_{d \leqslant (B_{1}'B_{2}')^{1/2}} \frac{\delta(d)}{d} \left| \sum_{|b_{2}'| \leqslant B_{2}'} \vartheta(b_{2}')^{2} \left(\frac{b_{2}'}{d}\right) \right|$$
$$\ll_{\varepsilon} \sum_{|b_{1}'| \leqslant B_{1}'} \vartheta(b_{1}')^{2} \sum_{d \leqslant (B_{1}'B_{2}')^{1/2}} d^{-1/2+\varepsilon} (\log B_{2}')^{2}$$
$$\ll_{\varepsilon} \sum_{|b_{1}'| \leqslant B_{1}'} \vartheta(b_{1}')^{2} (B_{1}'B_{2}')^{1/4+3\varepsilon}$$
$$\ll B_{1}'B_{2}', \qquad (3.17)$$

since $B'_1 \leq B'_2$.

For larger d, we employ Heath-Brown's estimate [11, Corollary 4] for real character sums. Suppose that D satisfies $(B'_1B'_2)^{1/2} \leq D \leq B'_3/e$, and write $a_d = \delta(d)(\frac{-\beta_1\beta_2}{d})$. Then proceeding as above we see that

$$S_e(D,D) = \sum_{D \ll d \ll D} \frac{a_d}{d} \sum_{\substack{[d_1,e_1]|f_1| \leqslant B'_1\\[d_2,e_2]|f_2| \leqslant B'_2}} \frac{|\mu(d_1)\mu(d_2)\mu(e_1)\mu(e_2)|}{d_1d_2e_1e_2} \Big(\frac{[d_1,e_1][d_2,e_2]f_1f_2}{d}\Big)$$

Let $b_d = a_d(\frac{[d_1, e_1][d_2, e_2]}{d})$, so that in particular $|b_d| \leq 1$. On recalling the fact that $B'_1 B'_2 \leq D^2$, we therefore deduce from [11, Corollary 4] that $S_e(D, D)$ is

$$\ll \sum_{\substack{d_1, e_1 \leqslant B'_1 \\ d_2, e_2 \leqslant B'_2}} \frac{1}{d_1 d_2 e_1 e_2} \left| \sum_{D \ll d \ll D} \frac{b_d}{d} \sum_{|f_1| \leqslant B'_1/[d_1, e_1]} \sum_{|f_2| \leqslant B'_2/[d_2, e_2]} \left(\frac{f_1 f_2}{d}\right) \right|$$

$$\ll_{\varepsilon} \frac{(DB'_1 B'_2)^{\varepsilon}}{D} \sum_{\substack{d_1, e_1 \leqslant B'_1 \\ d_2, e_2 \leqslant B'_2}} \frac{1}{d_1 d_2 e_1 e_2} \left(\frac{D^{1/2} B'_1 B'_2}{[d_1, e_1][d_2, e_2]} + \frac{D(B'_1 B'_2)^{1/2}}{([d_1, e_1][d_2, e_2])^{1/2}}\right)$$

$$\ll_{\varepsilon} D^{3\varepsilon - 1/2} B'_1 B'_2 + D^{3\varepsilon} (B'_1 B'_2)^{1/2},$$

for any $\varepsilon > 0$. Summing over dyadic intervals for $(B'_1B'_2)^{1/2} \leq D \leq D_2$, for any choice of $D_2 \leq B'_3/e$, we therefore obtain the estimate

$$S_e((B'_1B'_2)^{1/2}, D_2) \ll_{\varepsilon} B'_1B'_2 + D^{\varepsilon}_2(B'_1B'_2)^{1/2}.$$

Once combined with (3.17) this yields the bound

$$S_e(1, D_2) \ll_{\varepsilon} B'_1 B'_2 + D^{\varepsilon}_2 (B'_1 B'_2)^{1/2},$$
 (3.18)

for any $(B'_1B'_2)^{1/2} \leq D_2 \leq B'_3/e$.

Taking $D_2 = B'_3/e$ in (3.18) we obtain the estimate

$$S_e \ll_{\varepsilon} B_1' B_2' + B^{\varepsilon} (B_1' B_2')^{1/2},$$
 (3.19)

for any $\varepsilon > 0$. In order to obtain a second estimate, and so complete the proof of Lemma 7, we apply (3.18) with $D_2 = |\beta_1\beta_2|^{3/4} (B'_1B'_2)^{3/4}$ and (3.16) with $D_1 = |\beta_1\beta_2|^{3/4} (B'_1B'_2)^{3/4}$ and $D_2 = B'_3/e$. This produces the estimate

$$S_e \ll_{\varepsilon} B'_1 B'_2 + (B'_1 B'_2)^{1/2+\varepsilon} \Big(|\beta_1 \beta_2|^{\varepsilon} + \log B \Big),$$
 (3.20)

for any $\varepsilon > 0$. Since $\beta_i = a_i h_{ij} h_{ik}$, we clearly have the upper bound $|\beta_1 \beta_2| \leq h^2 A$. Thus we may combine this inequality with (3.19) and (3.20) in order to complete the proof of Lemma 7.

It remains to substitute Lemma 7 into (3.14), and then insert the resulting estimate into (3.13). But on taking $\varepsilon > 0$ to be sufficiently small it therefore follows that

$$\sum_{\mathbf{b}' \in \mathcal{B}} 2^{\omega(b_1'b_2'b_3')} \ll_{\varepsilon} \frac{h^{\varepsilon}B_3'}{\log B_3'} \sum_{e \leqslant B_3'} \frac{S_e}{e} \\ \ll_{\varepsilon} h^{1/32} B_3' \Big(B_1'B_2' + (B_1'B_2')^{1/2+\varepsilon} (\min\{A, B\}^{\varepsilon} + \log B) \Big).$$

which thereby establishes Lemma 5. This completes the proof of Proposition 1.

3.2 **Proof of Proposition 2**

We now turn to the proof of Proposition 2. Our approach is based upon Heath-Brown's treatment [14, Lemma 4] of the equation $n_1^2 n_2 n_3 + n_4^2 n_5 n_6 = n_7 n_8$. For fixed integers a, b, q we let $\rho(q; a, b)$ denote the number of solutions to the congruence $at^2 + b \equiv 0 \pmod{q}$. For any value of q, we then have

$$\rho(q;a,b) \leqslant \sum_{d|q} |\mu(d)| \left(\frac{-ab}{d}\right). \tag{3.21}$$

We shall establish Proposition 2 in the case (i, j, k) = (1, 2, 3), say. The other cases will follow by symmetry. Now it follows from (3.1) that for given a_i, b_1, b_2, c_3 , and each corresponding solution t of the congruence

$$a_1b_1t^2 + a_2b_2 \equiv 0 \pmod{a_3c_3^2},$$

we must have $c_1 \equiv tc_2 \pmod{a_3c_3^2}$. This gives rise to an equation of the form $\mathbf{h}.\mathbf{w} = 0$, with $\mathbf{h} = (1, -t, a_3c_3^2)$ and $\mathbf{w} = (c_1, c_2, k)$. Upon recalling that h.c.f. $(c_1, c_2) = 1$ from (3.2), an application of Lemma 1 therefore yields the bound

$$\ll \rho(a_3c_3^2; a_1b_2, a_2b_2) \Big(1 + \frac{C_1C_2}{|a_3c_3^2|}\Big).$$

for the number of possible b_3, c_1, c_2 given fixed choices for a_i, b_1, b_2 and c_3 . On employing (3.21) we therefore have

$$\mathcal{M}(A_i, B_i, C_i) \ll \sum_{a_i, b_1, b_2, c_3} \rho(a_3 c_3^2; a_1 b_2, a_2 b_2) \left(1 + \frac{C_1 C_2}{|a_3 c_3^2|}\right)$$
$$\ll \sum_{a_i, b_1, b_2, c_3} \sum_{d \mid a_3 c_3} |\mu(d)| \left(\frac{-a_1 a_2 b_1 b_2}{d}\right) \left(1 + \frac{C_1 C_2}{|a_3 c_3^2|}\right)$$
$$\ll \sum_{a_i, b_1, b_2, c_3} d(a_3) d(c_3) \left(1 + \frac{C_1 C_2}{|a_3 c_3^2|}\right),$$

since the sum over square-free divisors of $a_3c_3^2$ is the same as the sum over square-free divisors of a_3c_3 . But a simple application of partial summation now reveals that

$$\mathcal{M}(A_i, B_i, C_i) \ll \sum_{a_i, b_1, b_2, c_3} d(a_3) d(c_3) + C_1 C_2 \sum_{a_i, b_1, b_2, c_3} \frac{d(a_3) d(c_3)}{|a_3 c_3^2|} \\ \ll \left(AB_1 B_2 C_3 + A_1 A_2 B_1 B_2 C_1 C_2\right) (\log AC)^2,$$

which thereby completes the proof of Proposition 2.

4 Passage to the universal torsor

Our goal in this section is to equate the quantity (2.1) to the cardinality of a certain subset of integral points on the universal torsor above \tilde{X} . In fact

our approach to the universal torsor rests upon an entirely elementary analysis of the equation defining the surface X, and we shall not prove here that the resulting parametrisation is actually the universal torsor above \tilde{X} . This fact will be supplied for us by the work of Hassett and Tschinkel [9].

In any solution $\mathbf{x} \in Z_*^4$ to the equation $F(\mathbf{x}) = 0$ we see that x_4 divides $x_1x_2x_3$. Hence we may write $x_4 = y_1y_2y_3$ and $x_i = y_iz_i$, for some $\mathbf{y}, \mathbf{z} \in \mathbb{Z}^3$ with $y_iz_i \neq 0$. Suppose that $z_i = \varepsilon_i z'_i$ for $\varepsilon_i = \pm 1$ and $z'_i \in \mathbb{N}$. Then one easily employs the equation $F(\mathbf{x}) = 0$ to deduce that $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$. Hence, upon relabeling variables we may assume that

$$x_i = y_i z_i, \quad x_4 = y_1 y_2 y_3,$$

for $(\mathbf{y}, \mathbf{z}) \in \mathbb{Z}^3 \times \mathbb{N}^3$ with $y_i \neq 0$.

Under this substitution, the equation $F(\mathbf{x}) = 0$ becomes

$$z_1 z_2 z_3 = (y_1 z_1 + y_2 z_2 + y_3 z_3)^2.$$
(4.1)

Since \mathbf{x} is assumed to be primitive, it follows that \mathbf{y} is primitive. Moreover, if p is any prime divisor of h.c.f. (z_i, y_j) , then (4.1) implies that p divides $y_k z_k$. But this contradicts the primitivity of \mathbf{x} , whence

h.c.f.
$$(z_i, y_j) =$$
 h.c.f. $(y_1, y_2, y_3) = 1.$ (4.2)

We now write $z_i = w_i t_i^2$, for square-free $w_i \in \mathbb{N}$ and non-zero $t_i \in \mathbb{Z}$. In fact we may assume that $t_i \in \mathbb{N}$, since t_i and $-t_i$ produce the same value of z_i .

Then it follows from (4.2) that

h.c.f.
$$(w_i, y_j) = \text{h.c.f.}(t_i, y_j) = 1,$$
 (4.3)

and from (4.1) that $w_1 w_2 w_3$ is a square. Hence we can write

$$w_1 = u_2 u_3, \quad w_2 = u_1 u_3, \quad w_3 = u_1 u_2,$$

for square-free $u_i \in \mathbb{N}$, satisfying

h.c.f.
$$(u_i, u_j) =$$
 h.c.f. $(u_i, y_i) = 1.$ (4.4)

Indeed w_i is square-free, and any prime divisor of h.c.f. (u_i, y_i) must also divide h.c.f. $(w_i w_k, y_i)$, contrary to (4.3).

Substituting the quantities $w_i = u_j u_k$ into (4.1) therefore yields the expressions

$$\varepsilon t_1 t_2 t_3 u_1 u_2 u_3 = y_1 u_2 u_3 t_1^2 + y_2 u_1 u_3 t_2^2 + y_3 u_1 u_2 t_3^2, \tag{4.5}$$

where $\varepsilon = \pm 1$. It is clear that u_i must divide $y_i u_j u_k t_i^2$. But then u_i divides t_i , since u_i is square-free and h.c.f. $(u_i, y_i u_j u_k) = 1$, by (4.4). We proceed by writing

$$s_0 = \text{h.c.f.}(t_1/u_1, t_2/u_2, t_3/u_3),$$

and $s_i = t_i/(s_0 u_i)$. Plainly $s_0, s_i \in \mathbb{N}$, and $\mathbf{s} = (s_1, s_2, s_3)$ is primitive. Moreover, (4.3) yields

h.c.f.
$$(u_i, y_j) =$$
 h.c.f. $(s_i, y_j) = 1.$ (4.6)

Substituting $t_i = s_0 s_i u_i$ into (4.5), we therefore obtain the equations

$$\varepsilon s_0 s_1 s_2 s_3 u_1 u_2 u_3 = y_1 u_1 s_1^2 + y_2 u_2 s_2^2 + y_3 u_3 s_3^2, \tag{4.7}$$

where $\varepsilon = \pm 1$, and h.c.f. $(s_0, y_i) = 1$ by (4.3).

We proceed by using this equation, together with the fact that ${\bf s}$ is primitive, to establish that

h.c.f.
$$(s_i, u_j) = 1$$
.

If p is any prime divisor of h.c.f. (s_i, u_j) then it follows from (4.7), in conjunction with the coprimality conditions (4.4) and (4.6), that p divides s_k . Considering the corresponding p-adic order of each of the terms in (4.7), one is easily led to the conclusion that p divides $y_j s_j^2$, since u_j is square-free. But h.c.f. $(u_j, y_j) = 1$ by (4.4), and so p divides s_j , which is impossible. In fact we may go further and deduce that the components of **s** satisfy the relation

h.c.f.
$$(s_i, s_j) = 1$$
.

This follows immediately from (4.6), (4.7) and the fact that h.c.f. $(s_i, u_k) = 1$. Let $\mathcal{T} \subset \mathbb{A}^{10}$ denote the set of $(s_0, \mathbf{s}, \mathbf{u}, \mathbf{y}) \in \mathbb{N} \times N^3 \times N^3 \times Z^3_*$ such that

$$s_0 s_1 s_2 s_3 u_1 u_2 u_3 = y_1 u_1 s_1^2 + y_2 u_2 s_2^2 + y_3 u_3 s_3^2,$$
(4.8)

with

$$|\mu(u_1 u_2 u_3)| = \text{h.c.f.}(s_i, s_j) = \text{h.c.f.}(s_i, u_j) = 1,$$
(4.9)

and

h.c.f.
$$(s_0, y_i) =$$
 h.c.f. $(s_i, y_j) =$ h.c.f. $(u_i, y_1 y_2 y_3) = 1.$ (4.10)

Now let $\mathbf{x} \in Z^4_*$ be any solution to the equation $F(\mathbf{x}) = 0$. Then tracing back through our argument, we deduce that there exists $(\pm s_0, \mathbf{s}, \mathbf{u}, \mathbf{y}) \in \mathcal{T}$ such that

$$x_i = y_i u_i^2 u_j u_k s_0^2 s_i^2, \quad x_4 = y_1 y_2 y_3.$$
(4.11)

Conversely, given any $(\pm s_0, \mathbf{s}, \mathbf{u}, \mathbf{y}) \in \mathcal{T}$, the point given by (4.11) will be a primitive integer solution of the equation $F(\mathbf{x}) = 0$, with $x_1 x_2 x_3 x_4 \neq 0$. Indeed if p is any prime divisor of x_1, x_2, x_3, x_4 then we may assume that

$$p \mid y_i, p \mid s_0^2 u_1 u_2 u_3 \text{h.c.f.}(y_j u_j s_j^2, y_k u_k s_k^2)$$

But then (4.10) implies that $p \mid \text{h.c.f.}(y_1, y_2, y_3)$, which is impossible. We have therefore established the following result.

Lemma 8. We have

$$N_{U,H}(B) = \frac{1}{4} \# \Big\{ (s_0, \mathbf{s}, \mathbf{u}, \mathbf{y}) \in \mathcal{T} : \max\{ |y_i u_i^2 u_j u_k s_0^2 s_i^2|, |y_1 y_2 y_3| \} \leqslant B \Big\}.$$

The equation (4.8) is an affine embedding of the universal torsor above the minimal desingularisation \tilde{X} of X. As already mentioned, it has been calculated by Hassett and Tschinkel [9, §4] by computing generators for the Cox ring $Cox(\tilde{X})$ of \tilde{X} .

5 The lower bound

Our method of establishing the lower bound closely follows Heath-Brown's treatment of the Cayley cubic. Consequently we shall adopt similar notation throughout this section. Let $P_1, P_2 \leq B^{\delta}$, for some suitably small choice of $\delta > 0$. This choice will be specified in (5.13), below. The idea is to fix choices of $\mathbf{s}, \mathbf{u} \in N^3$ such that (4.9) holds and

$$u_1 u_2 u_3 = P_1, \quad s_1 s_2 s_3 = P_2.$$

In fact we shall insist upon the stronger condition that P_1P_2 is square-free. This is clearly permissable for the purposes of a lower bound. We then count the number of comparatively large non-zero solutions s_0, y_1, y_2, y_3 to the linear equation (4.8) subject to certain constraints.

Thus for $Y_0, Y_i \ge 1$, we let

$$\mathcal{N} = \mathcal{N}(\mathbf{s}, \mathbf{u}; Y_0, Y_1, Y_2, Y_3)$$

denote the number of $(s_0, y_1, y_2, y_3) \in \mathbb{N} \times \mathbb{Z}^3$ constrained by (4.8) and

h.c.f.
$$(s_0, y_i) = \text{h.c.f.}(y_i, P_1 P_2) = 1,$$
 (5.1)

for which

$$Y_0 \leqslant s_0 < 2Y_0, \quad Y_i \leqslant |y_i| < 2Y_i. \tag{5.2}$$

It should be clear that whenever (4.8) and (5.1) both hold, we automatically have (4.10) and h.c.f. $(y_1, y_2, y_3) = 1$. It will be convenient to define the quantities

$$A_0 = P_1 P_2, \quad A_i = u_i s_i^2,$$

so that (4.8) may be written

$$A_0 s_0 = A_1 y_1 + A_2 y_2 + A_3 y_3. (5.3)$$

Now it follows from Lemma 8 that we are only interested in values of s_0, y_i for which

$$A_0^2 A_i |s_0^2 y_i| \leq BP_1 P_2^2, \quad A_1 A_2 A_3 |y_1 y_2 y_3| \leq BP_1 P_2^2.$$

Hence we shall choose

$$Y_0 = \left[\frac{(BP_1P_2^2)^{1/3}}{2A_0}\right], \quad Y_i = \left[\frac{(BP_1P_2^2)^{1/3}}{2A_i}\right].$$
 (5.4)

Much as in Heath-Brown's treatment, the main difficulty arises from having to keep track of the coprimality conditions (5.1). Let

$$Q = P_1 P_2 \prod_{p \leqslant \sqrt{\log B}} p.$$

Following [14, Equation (3.6)], we write

$$\mathcal{N} \geqslant \mathcal{N}_1 - \mathcal{N}_2,\tag{5.5}$$

where \mathcal{N}_1 is the number of solutions in which the condition h.c.f. $(s_0, y_i) = 1$ is replaced by the weaker condition

h.c.f.
$$(s_0, y_i, Q) = 1$$
,

and \mathcal{N}_2 is the number of solutions in which some y_i shares a prime factor p with s_0 , such that $p \nmid Q$.

We proceed by estimating \mathcal{N}_1 , for which we use the Möbius function to pick out the coprimality conditions. Let $\mathcal{N}_3(\mathbf{d}; \mathbf{e}) = \mathcal{N}_3(d_1, d_2, d_3; e_1, e_2, e_3)$ denote the number of solutions of the equation (5.3) in the range (5.2) with $d_i \mid y_i$ and $e_i \mid s_0, y_i$. Then

$$\mathcal{N}_1 = \sum_{d_i \mid P_1 P_2} \mu(d_1) \mu(d_2) \mu(d_3) \sum_{e_i \mid Q} \mu(e_1) \mu(e_2) \mu(e_3) \mathcal{N}_3(\mathbf{d}; \mathbf{e}).$$
(5.6)

Our task is to estimate $\mathcal{N}_3(\mathbf{d}; \mathbf{e})$. Define the least common multiples

$$h_0 = [e_1, e_2, e_3], \quad h_i = [d_i, e_i]$$

and the lattice

$$\Lambda = \{ (n_1, n_2, n_3) \in \mathbb{Z}^3 : A_i h_i \mid n_i, \ A_0 h_0 \mid n_1 + n_2 + n_3 \}.$$

Then upon defining the region

$$\mathcal{R} = \{ \mathbf{r} \in \mathbb{R}^3 : A_i Y_i \leqslant |r_i| < 2A_i Y_i, \ A_0 Y_0 \leqslant |r_1 + r_2 + r_3| < 2A_0 Y_0 \},\$$

one follows the lines of Heath-Brown's argument in order to deduce that

$$\mathcal{N}_3(\mathbf{d}; \mathbf{e}) = \frac{\operatorname{vol}(\mathcal{R})}{\det \Lambda} + O((\det \Lambda)^2 \max\{Y_i, Y_0\}^2),$$

where

$$\det \Lambda = \frac{A_0 h_0 \prod_i A_i h_i}{\text{h.c.f.}(A_0 h_0, A_i h_i)}.$$

Since $d_i \mid P_1P_2$ and $e_i \mid Q$, we deduce that $A_0h_0 \leq P_1P_2Q^3$ and $A_ih_i \leq P_1^2P_2^3Q$. Hence we have

$$\det \Lambda \leqslant P_1^7 P_2^{10} Q^6 \ll P_1^{13} P_2^{16} \exp(O(\sqrt{\log B})) \ll B^{30\delta}.$$

It follows that the error term in our estimate for $\mathcal{N}_3(\mathbf{d}; \mathbf{e})$ is $O(B^{2/3+62\delta})$, and so (5.6) becomes

$$\mathcal{N}_{1} = \operatorname{vol}(\mathcal{R}) \sum_{d_{i}, e_{i}} \mu(d_{1}) \cdots \mu(e_{3}) \frac{\operatorname{h.c.f.}(A_{0}h_{0}, A_{i}h_{i})}{A_{0}h_{0} \prod_{i} A_{i}h_{i}} + O(B^{2/3 + 63\delta}), \quad (5.7)$$

since there are at most $O(B^{\delta})$ divisors of P_1P_2Q .

We now investigate the sum

$$\sum_{\substack{d_i|P_1P_2\\e_i|Q}} \mu(d_1)\cdots\mu(e_3)\frac{\text{h.c.f.}(A_0h_0,A_ih_i)}{A_0h_0\prod_i A_ih_i} = \prod_{p|Q} E_p,$$
(5.8)

say. When $p \mid Q$, but $p \nmid P_1P_2$, we see that

$$E_p = \sum_{\varepsilon_i \ge 0} \mu(p^{\varepsilon_1}) \mu(p^{\varepsilon_2}) \mu(p^{\varepsilon_3}) \frac{\text{h.c.f.}(p^{\max\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}}, p^{\varepsilon_i})}{p^{\max\{\varepsilon_1, \varepsilon_2, \varepsilon_3\} + \varepsilon_1 + \varepsilon_2 + \varepsilon_3}},$$

from which it easily follows that

$$E_p = 1 - \frac{3}{p^2} + \frac{2}{p^3}.$$
(5.9)

In this calculation we have used the fact that $p \nmid A_0A_i$ whenever $p \nmid P_1P_2$. Next, when $p \mid P_1$ we may assume that p divides precisely one factor, u_1 say. Since P_1P_2 is square-free it follows that $p \nmid u_2u_3P_2$, and that $p^2 \nmid u_1$. Let $A'_0 = p^{-1}A_0, A'_1 = p^{-1}A_1, A'_2 = A_2$ and $A'_3 = A_3$, so that

$$\frac{\text{h.c.f.}(A_0h_0, A_ih_i)}{A_0h_0\prod_i A_ih_i} = \frac{\text{h.c.f.}(pA'_0h_0, pA'_1h_1, A'_2h_2, A'_3h_3)}{p^2A'_0h_0\prod_i A'_ih_i},$$

with $p \nmid A'_0 A'_i$. Then in this setting we see that

$$E_p = \frac{1}{p^2} \sum_{\delta_i, \varepsilon_i \ge 0} \mu(p^{\delta_1}) \mu(p^{\delta_2}) \cdots \mu(p^{\varepsilon_3}) \frac{\text{h.c.f.}(p, p^{\max\{\delta_2, \varepsilon_2\}}, p^{\max\{\delta_3, \varepsilon_3\}})}{p^{\max\{\varepsilon_1, \varepsilon_2, \varepsilon_3\} + \max\{\delta_1, \varepsilon_1\} + \cdots + \max\{\delta_3, \varepsilon_3\}}},$$

whence a straightforward calculation yields

$$E_p = \frac{1}{p^2} \left(1 - \frac{1}{p} - \frac{1}{p^2} + \frac{1}{p^3} \right).$$
 (5.10)

Finally we consider the case $p \mid P_2$, so that p divides precisely one factor, s_1 say. Since P_1P_2 is square-free it follows that $p \nmid P_1s_2s_3$, and that $p^2 \nmid s_1$. Let $A'_0 = p^{-1}A_0, A'_1 = p^{-2}A_1, A'_2 = A_2$ and $A'_3 = A_3$. Then arguing as above we now have

$$E_p = \frac{1}{p^3} \sum_{\delta_i, \varepsilon_i \ge 0} \mu(p^{\delta_1}) \mu(p^{\delta_2}) \cdots \mu(p^{\varepsilon_3}) \frac{\text{h.c.f.}(p, p^{\max\{\delta_2, \varepsilon_2\}}, p^{\max\{\delta_3, \varepsilon_3\}})}{p^{\max\{\varepsilon_1, \varepsilon_2, \varepsilon_3\} + \max\{\delta_1, \varepsilon_1\} + \cdots + \max\{\delta_3, \varepsilon_3\}}}.$$

In view of our calculation for (5.10) we immediately deduce that

$$E_p = \frac{1}{p^3} \left(1 - \frac{1}{p} - \frac{1}{p^2} + \frac{1}{p^3} \right).$$
(5.11)

Taking (5.9)- (5.11) together in (5.8), it therefore follows that

$$\sum_{d_i, e_i} \mu(d_1) \cdots \mu(e_3) \frac{\text{h.c.f.}(A_0 h_0, A_i h_i)}{A_0 h_0 \prod_i A_i h_i} \gg \frac{1}{P_1^2 P_2^3} \frac{\phi(P_1 P_2)}{P_1 P_2} = \frac{\phi(P_1 P_2)}{P_1^3 P_2^4},$$

since $\phi(n) = n \prod_{p|n} (1 - 1/p)$ for any $n \in \mathbb{N}$. Our choices (5.4) for Y_i, Y_0 clearly imply that $\operatorname{vol}(\mathcal{R}) \gg BP_1P_2^2$. We claim that

$$\mathcal{N}_1 \gg \frac{B}{P_1 P_2} \frac{\phi(P_1 P_2)}{P_1 P_2},$$
(5.12)

provided that we take

$$\delta = 1/201. \tag{5.13}$$

In order to establish the claim, it clearly suffices to check that the lower bound in (5.12) is larger than the error term in (5.7) when δ is taken to be 1/201. But on using the trivial lower bound $\phi(n) \ge 1$ for any $n \in \mathbb{N}$, we see that

$$\frac{B}{P_1 P_2} \frac{\phi(P_1 P_2)}{P_1 P_2} \geqslant \frac{B}{(P_1 P_2)^2} \gg B^{1-4\delta}.$$

Since $B^{1-4\delta} \gg B^{2/3+63\delta}$ for $\delta = 1/201$, the claim follows.

Next we must produce an upper bound for \mathcal{N}_2 , for which we may ignore any coprimality conditions whenever we wish to. Suppose that $p \mid s_0, y_1$, for some prime $p \nmid Q$ lying in the range $R \leq p < 2R$. In particular we may assume that $R \ll Y_1$. There are O(R) such primes, and we fix one particular choice. Following Heath-Brown's treatment, we write $s_0 = pt_0$ and $y_1 = pt_1$ and count solutions of the linear equation

$$pA_0t_0 = pA_1t_1 + A_2y_2 + A_3y_3. (5.14)$$

In particular t_0, t_1 are contained in the ranges

$$Y_0/R \ll |t_0| \ll Y_0/R, \quad Y_1/R \ll |t_1| \ll Y_1/R$$

Since P_1P_2 is square-free, it follows that h.c.f. $(A_i, A_0) = u_i s_i$. Hence we may deduce from (5.14) that

$$A_2 y_2 \equiv -A_3 y_3 \pmod{p u_1 s_1}.$$

We may assume by symmetry that $A_2Y_2 \ge A_3Y_3$. Upon noting that pu_1s_1 is coprime to A_2 , since h.c.f. $(y_i, P_1P_2) = 1$, it follows that for each choice of y_3 , there are $O(1 + Y_2/(Ru_1s_1))$ possibilities for y_2 . Now (5.14) implies that

$$Ru_1s_1 \ll \max\{A_2Y_2, A_3Y_3\} = A_2Y_2. \tag{5.15}$$

Moreover, it follows from (5.4) that

$$A_2 u_1 s_1 \ll Y_2,$$
 (5.16)

provided that $\delta \leq 1/5$. Together (5.15) and (5.16) imply that

$$1 \ll \left(\frac{A_2 Y_2}{R u_1 s_1}\right)^{1/2} \left(\frac{Y_2}{A_2 u_1 s_1}\right)^{1/2} = \frac{Y_2}{R^{1/2} u_1 s_1},$$

whence we deduce that there are $O(Y_2Y_3/(R^{1/2}u_1s_1))$ choices for y_2, y_3 . We fix such a choice and write $A_2y_2 + A_3y_3 = pu_1s_1k$. Then it remains to count values of t_0, t_1 for which

$$u_2 u_3 s_2 s_3 t_0 = s_1 t_1 + k. (5.17)$$

Now we have already seen that $R \ll Y_1$. Moreover, as in (5.16), we can use (5.4) to show that $u_2^3 u_3^3 s_2^3 s_3^3 \ll Y_1$ provided that $\delta \leq 1/15$. Together these inequalities imply that

$$1 \ll \left(\frac{Y_1}{R}\right)^{2/3} \left(\frac{Y_1}{u_2^3 u_3^3 s_2^3 s_3^3}\right)^{1/3} = \frac{Y_1}{R^{2/3} u_2 u_3 s_2 s_3}$$

Viewing (5.17) as a congruence modulo $u_2u_3s_2s_3$, one easily concludes that there are $O(Y_1/(R^{2/3}u_2u_3s_2s_3))$ possibilities for t_0, t_1 .

In conclusion we have therefore shown that the total number of admissible p, y_2, y_3, t_0, t_1 , for which $R \leq p < 2R$, is

$$\ll R \cdot \frac{Y_2 Y_3}{R^{1/2} u_1 s_1} \cdot \frac{Y_1}{R^{2/3} u_2 u_3 s_2 s_3} \ll \frac{B P_1 P_2^2}{R^{1/6} P_1^2 P_2^3} = \frac{B}{R^{1/6} P_1 P_2}$$

by (5.4). Summing $R \gg \sqrt{\log B}$ over dyadic intervals, we deduce that

$$\mathcal{N}_2 \ll \frac{B}{P_1 P_2} (\log B)^{-1/12},$$

provided that $\delta \leq 1/15$. It follows from (5.12) and (5.13) that $\mathcal{N}_2 = o(\mathcal{N}_1)$, and so (5.5) implies that

$$\mathcal{N} \gg \frac{B}{P_1 P_2} \frac{\phi(P_1 P_2)}{P_1 P_2}$$

Finally, in order to complete the proof of the lower bound in (1.1), we note that any square-free value of P will factorise into values $u_1, u_2, u_3, s_1, s_2, s_3$ satisfying (4.9), in precisely $d_6(P)$ ways. It therefore follows that

$$N_{U,H}(B) \gg \sum_{P \leqslant B^{2/201}} |\mu(P)| d_6(P) \frac{B}{P} \frac{\phi(P)}{P}.$$
(5.18)

To handle this quantity we define the sum

$$S(x) = \sum_{n \leqslant x} \frac{|\mu(n)| d_6(n)\phi(n)}{n},$$

for any x > 1, and proceed by establishing the following simple bound.

Lemma 9. For any x > 1 we have

$$S(x) \gg x(\log x)^5.$$

Proof. To establish the lemma we shall apply Perron's formula to the corresponding Dirichlet series

$$F(s) = \sum_{n=1}^{\infty} \frac{|\mu(n)| d_6(n) \phi(n)/n}{n^s},$$

defined for $\Re e(s) > 1$. It is a trivial matter to calculate the Euler product

$$F(s) = \prod_{p} \left(1 + \frac{6(1 - 1/p)}{p^s} \right) = \zeta(s)^6 G(s),$$

for some function G(s) that is holomorphic and bounded on the half-plane $\Re e(s) > 0$. Let $\varepsilon > 0$ and let $T \in [1, x]$. Then Perron's formula yields

$$S(x) = \frac{1}{2\pi i} \int_{1+\varepsilon-iT}^{1+\varepsilon+iT} \zeta(s)^6 G(s) \frac{x^s}{s} \mathrm{d}s + O_\varepsilon \left(\frac{x^{1+\varepsilon}}{T}\right).$$

We apply Cauchy's residue theorem to the rectangular contour joining the points 1/2 - iT, 1/2 + iT, $1 + \varepsilon + iT$ and $1 + \varepsilon - iT$, which therefore leads to the conclusion that there exists a polynomial f of degree 5 such that

$$S(x) - xf(\log x) \ll_{\varepsilon} \frac{x^{1+\varepsilon}}{T} + \Big(\int_{1/2-iT}^{1/2+iT} + \int_{1/2-iT}^{1+\varepsilon-iT} + \int_{1+\varepsilon+iT}^{1/2+iT}\Big) \Big|\zeta(s)^{6} \frac{x^{s}}{s}\Big| \mathrm{d}s.$$

Here we have used the fact that G(s) is bounded for $\Re e(s) > 0$. To estimate this error term we apply the well known convexity bound $\zeta(\sigma + it) \ll_{\varepsilon} |t|^{(1-\sigma)/3+\varepsilon}$, valid for any $\sigma \in [1/2, 1]$ and $|t| \ge 1$. Thus it follows that

$$S(x) - xf(\log x) \ll_{\varepsilon} \frac{x^{1+\varepsilon}}{T} + x^{1/2+\varepsilon}T^{1+\varepsilon}.$$

Selecting $T = x^{1/4}$ therefore completes the proof of Lemma 9.

On combining Lemma 9 with an application of partial summation, and then inserting the resulting estimate into (5.18), we therefore deduce that

$$N_{U,H}(B) \gg B(\log B)^6.$$

This completes the proof of the lower bound in (1.1).

6 The upper bound

Fix a choice of $X_1, \ldots, X_4, S_0, S_i, U_i, Y_i \ge 1$. We shall write

$$\mathcal{N} = \mathcal{N}(X_1, \dots, X_4; S_0; S_1, S_2, S_3; U_1, U_2, U_3; Y_1, Y_2, Y_3)$$

for the total contribution to $N_{U,H}(B)$ from **x** contained in the intervals

$$X_{\xi} \leqslant |x_{\xi}| < 2X_{\xi}, \quad (1 \leqslant \xi \leqslant 4), \tag{6.1}$$

and such that the variables s_0 , \mathbf{s} , \mathbf{u} , \mathbf{y} appearing in Lemma 8 satisfy

$$S_0 \leq s_0 < 2S_0, \quad S_i \leq |s_i| < 2S_i, \quad U_i \leq |u_i| < 2U_i, \quad Y_i \leq |y_i| < 2Y_i.$$
 (6.2)

It will be convenient to relable the indices so that

$$X_1 \leqslant X_2 \leqslant X_3. \tag{6.3}$$

Suppose that $\mathbf{x} \in Z^4_*$ is a solution of $F(\mathbf{x}) = 0$, with $|x_1|, \ldots, |x_4| \leq B$. Then (6.1) implies that

$$X_1, X_2, X_3, X_4 \leqslant B. \tag{6.4}$$

If $\mathcal{N} = 0$ there is nothing to prove, and so we assume henceforth that the dyadic ranges in (6.1) and (6.2) produce a non-zero value of \mathcal{N} .

We proceed by showing that under the assumption that $\mathcal{N} \neq 0$, certain choices of dyadic ranges in (6.1) and (6.2) force certain other ranges to have fixed order of magnitude. It will be convenient to write

$$S = S_1 S_2 S_3, \quad U = U_1 U_2 U_3, \quad Y = Y_1 Y_2 Y_3.$$

Hence it follows from (4.11) that

$$X_i \ll Y_i U_i U S_0^2 S_i^2 \ll X_i, \tag{6.5}$$

and that

$$X_4 \ll Y \ll X_4. \tag{6.6}$$

Together, (6.5) and (6.6) imply that

$$\left(\frac{X_1 X_2 X_3}{X_4}\right)^{1/2} \ll S_0^3 S U^2 \ll \left(\frac{X_1 X_2 X_3}{X_4}\right)^{1/2}.$$
(6.7)

We take a moment to record two further inequalities satisfied by the quantities S_0, S_i, U_i, Y_i , which will be crucial in our final analysis. First we deduce from (6.5), (6.6) and (6.7) that

$$S_0 S^{1/3} U^{2/3} Y^{2/3} = (S_0^3 S U^2)^{1/3} Y^{2/3} \ll (X_1 X_2 X_3)^{1/6} X_4^{1/2}.$$
(6.8)

Similarly, we may deduce that

$$S_0 S^{1/2} U Y^{1/2} \leqslant (S_0^3 S U^2)^{1/2} Y^{1/2} \ll (X_1 X_2 X_3 X_4)^{1/4}.$$
(6.9)

It is clear that \mathcal{N} is bounded above by the number of $s_0 \in \mathbb{N}$ and $\mathbf{s}, \mathbf{u}, \mathbf{y} \in Z^3_*$ contained in the ranges (6.2), for which (4.8), (4.9) and (4.10) all hold. Ultimately we shall sum over suitable dyadic intervals for X_1, X_2, X_3, X_4 and S_0, S_i, U_i, Y_i , in order to establish the upper bound in (1.1), and so complete the proof of the theorem.

For any fixed choice of $s_0 \in \mathbb{N}$ and $\mathbf{s}, \mathbf{u} \in \mathbb{Z}^3_*$ in the region (6.2), with (4.9) holding, we let

$$N(s_0, \mathbf{s}, \mathbf{u}) = N(Y_1, Y_2, Y_3; s_0, \mathbf{s}, \mathbf{u})$$

denote the corresponding contribution to \mathcal{N} from the $\mathbf{y} \in Z^3_*$. Clearly we are only interested in values of $s_0, \mathbf{s}, \mathbf{u}$ for which $N(s_0, \mathbf{s}, \mathbf{u})$ is non-zero. Considering $s_0, \mathbf{s}, \mathbf{u}$ to be fixed, we select any vector

$$\hat{\mathbf{y}} = (\hat{y}_1, \hat{y}_2, \hat{y}_3) \tag{6.10}$$

for which the Euclidean norm $|\hat{\mathbf{y}}|$ is least. Following the convention that this vector too is fixed, for fixed values of $s_0, \mathbf{s}, \mathbf{u}$, we define the change of variables

$$z_i = y_i - \hat{y}_i.$$
 (6.11)

We shall let $N_1(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}})$ denote the overall contribution to $N(s_0, \mathbf{s}, \mathbf{u})$ from those \mathbf{y} for which $z_1 z_2 z_3 \neq 0$, and we let $N_2(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}})$ denote the remaining contribution to $N(s_0, \mathbf{s}, \mathbf{u})$ from those \mathbf{y} for which $z_1 z_2 z_3 = 0$. With this notation we therefore have

$$\mathcal{N} = \sum_{s_0, \mathbf{s}, \mathbf{u}} N(s_0, \mathbf{s}, \mathbf{u}) = \sum_{s_0, \mathbf{s}, \mathbf{u}} N_1(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}}) + \sum_{s_0, \mathbf{s}, \mathbf{u}} N_2(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}})$$
$$= \mathcal{N}_1 + \mathcal{N}_2, \tag{6.12}$$

say. Here the summations are over all $(s_0, \mathbf{s}, \mathbf{u}) \in \mathbb{N} \times Z^3_* \times Z^3_*$ in the region (6.2), with (4.9) holding. It will be necessary to investigate the quantities \mathcal{N}_1 and \mathcal{N}_2 separately.

6.1 Estimating \mathcal{N}_1

For any fixed choice of $s_0 \in \mathbb{N}$ and $\mathbf{s}, \mathbf{u} \in Z^3_*$ in the region (6.2), with (4.9) holding, we let $\hat{\mathbf{y}}$ be the corresponding vector (6.10) that was selected above. On recalling the change of variables (6.11), it therefore follows from (4.8) that

$$z_1 u_1 s_1^2 + z_2 u_2 s_2^2 + z_3 u_3 s_3^2 = 0, (6.13)$$

and from (6.2) that $|z_i| < 4Y_i$. Hence we deduce that

$$\mathcal{N}_1 \ll S_0 \sum_{\mathbf{s}, \mathbf{u}} \# \{ \mathbf{z} \in \mathbb{Z}^3 : z_1 z_2 z_3 \neq 0, \ |z_i| < 4Y_i, \ (6.13) \text{ holds} \}.$$

Recall that $u_1u_2u_3$ is square-free, so that h.c.f. $(u_i, u_j) = 1$. It is apparent that the **z** appearing in the summand need not be primitive. Moreover we no longer

necessarily have coprimality conditions corresponding to (4.10). In order to recover a weaker set of coprimality relations, we shall write

$$u_i = d_{jk}u'_i, \quad z_i = d_{ij}d_{ik}ez'_i,$$

say, for any $d_{ij}, e \in \mathbb{N}$ with the convention that $d_{ij} = d_{ji}$. Let

$$U_i' = \frac{U_i}{d_{jk}}, \quad Y_i' = \frac{Y_i}{d_{ij}d_{ik}e},$$

and

$$U' = U_1' U_2' U_3', \quad Y' = Y_1' Y_2' Y_3'.$$

In particular it follows from (4.9) that

$$|\mu(u'_1u'_2u'_3)| = \text{h.c.f.}(s_i, s_j) = \text{h.c.f.}(s_i, u'_j) = 1,$$
(6.14)

for any \mathbf{s} and \mathbf{u}' .

For fixed values of $d_{ij}, e \in \mathbb{N}$, our task is to estimate the number of $\mathbf{s}, \mathbf{u}', \mathbf{z}' \in \mathbb{Z}^3_*$ such that

h.c.f.
$$(u'_i, z'_i, z'_k) = 1$$
,

(6.14) holds,

$$S_i \leq |s_i| < 2S_i, \quad U'_i \leq |u'_i| < 2U'_i, \quad |z'_i| < 4Y'_i,$$

and

$$z_1'u_1's_1^2 + z_2'u_2's_2^2 + z_3'u_3's_3^2 = 0.$$

But this quantity is clearly bounded above by $\mathcal{M} = \mathcal{M}(2U'_i, 4Y'_i, 2S_i)$ in the notation of §3. Thus it follows that

$$\mathcal{N}_1 \ll S_0 \sum_{d_{ij}, e \ll Y} \mathcal{M},\tag{6.15}$$

and Proposition 1 yields

$$\mathcal{M} \ll_{\varepsilon} S^{1/3} U'^{2/3} Y'^{2/3} + \sigma \tau S^{1/2} U' Y'^{1/2}, \tag{6.16}$$

for any $\varepsilon > 0$, where

$$\sigma = 1 + \frac{\min\{U, Y\}^{\varepsilon}}{\min\{Y'_i Y'_j\}^{3/32 - \varepsilon}}, \quad \tau = 1 + \frac{\log Y}{\min\{Y'_i Y'_j\}^{3/32 - \varepsilon}}$$

On applying Proposition 2 we obtain the alternative estimate

$$\mathcal{M} \ll U' Y_i' Y_j' \left(S_k + S_i S_j U_k'^{-1} \right) (\log S U')^2, \tag{6.17}$$

for any permutation $\{i, j, k\}$ of the set $\{1, 2, 3\}$.

We may now use (6.16) and (6.17) to obtain a pair of estimates for \mathcal{N}_1 . Recall the inequality (6.15) for \mathcal{N}_1 , and note that

$$U' = (d_{12}d_{13}d_{23})^{-1}U, \quad Y' = (d_{12}d_{13}d_{23})^{-2}e^{-3}Y.$$

Beginning with an application of (6.16), we deduce that

$$\sum_{d_{ij}, e \ll Y} \mathcal{M} \ll_{\varepsilon} S^{1/3} U^{2/3} Y^{2/3} + \sigma \tau S^{1/2} U Y^{1/2},$$

where

$$\sigma = 1 + \frac{\min\{U, Y\}^{\varepsilon}}{\min\{Y_i Y_j\}^{3/32 - \varepsilon}}, \quad \tau = 1 + \frac{\log B}{\min\{Y_i Y_j\}^{3/32 - \varepsilon}}.$$
 (6.18)

Thus it follows that

$$\mathcal{N}_1 \ll_{\varepsilon} S_0 S^{1/3} U^{2/3} Y^{2/3} + \sigma \tau S_0 S^{1/2} U Y^{1/2}, \tag{6.19}$$

for any $\varepsilon > 0$. Similarly, since (6.4) and (6.5) imply that $\log(SU) \ll \log B$, an application (6.17) yields the estimate

$$\mathcal{N}_1 \ll S_0 U Y_i Y_j (S_k + S_i S_j S_k^{-1} U_k^{-1}) (\log B)^2, \tag{6.20}$$

for any permutation $\{i, j, k\}$ of the set $\{1, 2, 3\}$.

6.2 Estimating \mathcal{N}_2

In this section we estimate the size of \mathcal{N}_2 . For any fixed choice of $s_0 \in \mathbb{N}$ and $\mathbf{s}, \mathbf{u} \in \mathbb{Z}^3_*$ in the region (6.2), with (4.9) holding, let $\hat{\mathbf{y}}$ be the vector (6.10) counted by $N(s_0, \mathbf{s}, \mathbf{u})$ that was selected at the start of §6. Then (6.12) implies that

$$\mathcal{N}_2 = \sum_{s_0, \mathbf{s}, \mathbf{u}} N_2(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}}),$$

where $N_2(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}})$ denotes the contribution to $N(s_0, \mathbf{s}, \mathbf{u})$ from those \mathbf{y} for which

$$\prod_{1\leqslant i\leqslant 3} (y_i - \hat{y}_i) = 0.$$

Let $N_2^{(i)}$ denote the total contribution to $N_2(s_0, \mathbf{s}, \mathbf{u}; \hat{\mathbf{y}})$ from those \mathbf{y} for which $y_i = \hat{y}_i$ is fixed. It therefore follows that

$$\mathcal{N}_2 \leqslant \sum_{s_0, \mathbf{s}, \mathbf{u}} \left(N_2^{(1)} + N_2^{(2)} + N_2^{(3)} \right) = \mathcal{N}_2^{(1)} + \mathcal{N}_2^{(2)} + \mathcal{N}_2^{(3)}, \tag{6.21}$$

say. In order to estimate $N_2^{(i)}$ for fixed values of $s_0 \in \mathbb{N}$ and $\mathbf{s}, \mathbf{u} \in Z^3_*$, it suffices to count non-zero integer solutions y_j, y_k to the equation

$$y_j u_j s_j^2 + y_k u_k s_k^2 = n, (6.22)$$

where $n = s_0 s_1 s_2 s_3 u_1 u_2 u_3 - \hat{y}_i u_i s_i^2$ is fixed. Our first step is to deduce from (4.9) that

h.c.f.
$$(u_i s_i^2, u_j s_j^2) = 1.$$

Noting that $|y_j| < 2Y_j$ and $|y_k| < 2Y_k$, we proceed by applying Lemma 1 to (6.22). Taking

$$\mathbf{h} = (u_j s_j^2, u_k s_k^2, n), \quad \mathbf{w} = (y_j, y_k, 1),$$

we therefore deduce that

$$N_2^{(i)} \ll 1 + \frac{Y_j Y_k}{\max\{Y_j U_j S_j^2, Y_k U_k S_k^2, |n|\}}.$$
(6.23)

Since $Y_i U_i S_i^2 \ll \hat{y}_i u_i s_i^2 \ll Y_i U_i S_i^2$ and $S_0 SU \ll s_0 s_1 s_2 s_3 u_1 u_2 u_3 \ll S_0 SU$, by (6.2), it is easy to see that

$$\begin{aligned} |n| &= |s_0 s_1 s_2 s_3 u_1 u_2 u_3 - \hat{y_i} u_i s_i^2| \\ &\geqslant \left| |\hat{y_i} u_i s_i^2| - |s_0 s_1 s_2 s_3 u_1 u_2 u_3| \right| \\ &\gg Y_i U_i S_i^2, \end{aligned}$$

if $Y_i U_i S_i^2 \gg S_0 SU$. Upon summing (6.23) over all $s_0, \mathbf{s}, \mathbf{u}$, and then inserting the resulting bound into (6.21), we therefore obtain

$$\mathcal{N}_2 \ll S_0 SU + \max_{\{i,j,k\}} \Big\{ \frac{S_0 SU Y_j Y_k}{\max\{Y_j U_j S_j^2, Y_k U_k S_k^2, \theta_i\}} \Big\},$$
(6.24)

where the first maximum is over all permutations $\{i, j, k\}$ of the set $\{1, 2, 3\}$, and

$$\theta_i = \begin{cases} Y_i U_i S_i^2, & Y_i U_i S_i^2 \gg S_0 S U, \\ 1, & \text{otherwise.} \end{cases}$$
(6.25)

6.3 Completion of the upper bound

We begin by treating the estimates (6.19) and (6.20) for \mathcal{N}_1 . Now it follows from the inequalities (6.2), (6.3) and (6.5) that

$$Y_1 U_1 S_1^2 \ll Y_2 U_2 S_2^2 \ll Y_3 U_3 S_3^2. \tag{6.26}$$

In particular (4.8) implies that

$$S_0 SU \ll Y_3 U_3 S_3^2.$$
 (6.27)

Multiplying both sides of this inequality by $S_0^2 U$, and recalling (6.4) and (6.5), we deduce that

$$S_0^3 S U^2 \ll B.$$
 (6.28)

It will also be useful to deduce an inequality involving the maximum size of the U_i . Suppose temporarily that $U_1 \leq U_2 \leq U_3$, so that $U_1U_2 \leq U_3^2$. Then in view of (6.4) and (6.5) it clearly follows that $U_1U_2U_3^2 \ll B$, whence $U_1U_2 \ll B^{1/2}$. Using this sort of argument it is not hard to deduce that in general

$$U_i U_j \ll B^{1/2}.$$
 (6.29)

Throughout the treatment of \mathcal{N}_1 we shall make the assumption that

$$Y_i \leqslant Y_j \leqslant Y_\kappa$$

for some permutation $\{i, j, \kappa\} = \{1, 2, 3\}$. Our plan is to use (6.19) whenever

$$Y_{\kappa} \leqslant (Y_i Y_j)^9 \quad \text{or} \quad U \leqslant (Y_i Y_j)^3, \tag{6.30}$$

and (6.20) in the alternative case

$$Y_i Y_j \leq \min\{Y_{\kappa}^{1/9}, U^{1/3}\}.$$
 (6.31)

Let us consider the case (6.30) first. Recalling the definition (6.18) of σ , it follows that

$$\sigma \ll 1 + (Y_i Y_j)^{11\varepsilon - 3/32}$$

Hence we may take $\sigma \ll 1$ in (6.19), provided that we take $\varepsilon > 0$ to be sufficiently small. It therefore follows from (6.8), (6.19) and the choice $\varepsilon = 1/128$, that

$$\mathcal{N}_1 \ll (X_1 X_2 X_3)^{1/6} X_4^{1/2} + S_0 S^{1/2} U Y^{1/2} \Big(1 + \frac{\log B}{(Y_i Y_j)^{11/128}} \Big), \tag{6.32}$$

whenever (6.30) holds. We now sum over the various dyadic intervals for the X_1, X_2, X_3, X_4 , subject to (6.4), and also the dyadic intervals for S_0, S_i, U_i, Y_i , subject to (6.5), (6.6) and (6.30). Suppose for the moment that we are interested in summing over all possible dyadic intervals $X \leq |x| < 2X$, for which $|x| \leq \mathcal{X}$. Then there are plainly $O(\log \mathcal{X})$ possible choices for X. In addition to this basic estimate, we shall make frequent use of the estimates

$$\sum_{X} X^{\delta} \ll_{\delta} \begin{cases} 1, & \delta < 0, \\ \mathcal{X}^{\delta}, & \delta > 0. \end{cases}$$

Returning to (6.32), we may deduce from (6.5) and (6.7) that values of S_0, Y_1, Y_2, Y_3 are determined by the choices of X_1, X_2, X_3, X_4 and S_i, U_i . Now there are clearly $\ll (\log B)^6$ possible sets of values for S_i, U_i . In view of (6.4), we therefore obtain the estimate

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} (X_1 X_2 X_3)^{1/6} X_4^{1/2} \ll B(\log B)^6.$$
(6.33)

Employing (6.9), we find similarly that

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} S_0 S^{1/2} U Y^{1/2} \ll \sum_{S_0, S_i, U_i, Y_i, X_i, X_4} (X_1 X_2 X_3 X_4)^{1/4} \\ \ll B(\log B)^6.$$
(6.34)

Finally we turn to the term $S_0 S^{1/2} U(Y_i Y_j)^{53/128} Y_{\kappa}^{1/2} \log B$ in (6.32). We shall sum over dyadic intervals subject to the two inequalities

$$Y_{\kappa} \ll \frac{B}{Y_i Y_j}, \quad S_1 \ll \frac{B}{S_0^3 S_2 S_3 U^2}.$$

The first of these follows from (6.4) and (6.6), whereas the second is just (6.28). We therefore obtain the estimate

$$\sum_{S_0, S_i, U_i, Y_i} S_0 S^{1/2} U(Y_i Y_j)^{53/128} Y_{\kappa}^{1/2} \ll B^{1/2} \sum_{S_0, S_i, U_i, Y_i, Y_j} \frac{S_0 S^{1/2} U}{(Y_i Y_j)^{11/128}} \\ \ll B \sum_{S_0, S_2, S_3, U_i, Y_i, Y_j} S_0^{-1/2} (Y_i Y_j)^{-11/128} \\ \ll B(\log B)^5.$$

Since values of X_1, X_2, X_3, X_4 are determined by choices of S_0, S_i, U_i, Y_i , we may combine this latter estimate with (6.33) and (6.34) in (6.32), in order to conclude that

$$\sum_{\substack{S_0, S_i, U_i, Y_i, X_i, X_4 \\ (6.30) \text{ holds}}} \mathcal{N}_1 \ll B(\log B)^6.$$
(6.35)

Next we handle the case in which (6.31) holds. For this we employ the alternative estimate (6.20) for \mathcal{N}_1 , and so deduce that

$$\mathcal{N}_1 \ll (S_0 S_\kappa U Y_i Y_j + S_0 S_i S_j U_i U_j Y_i Y_j) (\log B)^2.$$

We shall again need to sum over dyadic intervals for $S_0, S_i, U_i, Y_i, X_i, X_4$, this time subject to (6.4), (6.5), (6.6) and (6.31).

Let us consider the term $S_0 S_{\kappa} U Y_i Y_j$. But then (6.5), (6.7), (6.29) and (6.31) together imply that

$$\begin{split} S_0 S_{\kappa} U Y_i Y_j &= S_0^3 S U^2 \frac{Y_i Y_j}{S_0^2 S_i S_j U} \\ &\ll (X_1 X_2 X_3)^{1/2} \frac{(Y_i Y_j)^{3/2} (U_i U_j)^{1/2}}{(X_i X_j)^{1/2}} \\ &\ll X_{\kappa}^{1/2} Y_{\kappa}^{1/6} (U_i U_j)^{1/2} \\ &\ll B^{11/12}. \end{split}$$

Since there are at most $O_{\varepsilon}(B^{\varepsilon})$ dyadic intervals for S_0, S_i, U_i, Y_i , which in turn determine values of X_1, X_2, X_3, X_4 , this therefore leads to the conclusion that

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} S_0 S_\kappa U Y_i Y_j (\log B)^2 \ll B,$$
(6.36)

whenever (6.31) holds. Lastly we consider the term $S_0S_iS_jU_iU_jY_iY_j$. Now there are $O(\log B)$ dyadic intervals for Y_{κ} , and (6.31) implies that $Y_i, Y_j \leq U^{1/3}$. Employing the upper bound $S_i \ll B/(S_0^3S_jS_{\kappa}U^2)$, we therefore deduce that

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} S_0 S_i S_j U_i U_j Y_i Y_j \ll \log B \sum_{S_0, S_i, U_i} S_0 S_i S_j U_i U_j U^{2/3}$$
$$\ll B \log B \sum_{S_0, S_j, S_\kappa, U_i} S_0^{-2} S_\kappa^{-1} U^{-1/3}$$
$$\ll B (\log B)^2,$$

whenever (6.31) holds. Once combined with (6.36) this yields the overall contribution

$$\sum_{\substack{S_0, S_i, U_i, Y_i, X_i, X_4 \\ (6.31) \text{ holds}}} \mathcal{N}_1 \ll B(\log B)^4.$$
(6.37)

Finally we turn to the contribution from \mathcal{N}_2 , as estimated in (6.24). Recall the inequality (6.27). Then we proceed by establishing that

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} Y_3 U_3 S_3^2 \ll B(\log B)^4.$$
(6.38)

In order to do so we observe that values of X_1, X_2, X_3, X_4 are determined by the choices of S_0, S_i, U_i, Y_i . Recall the inequalities (6.26). We have two basic cases to consider, according to whether or not $Y_3U_3S_3^2$ is sufficiently large compared with $Y_2U_2S_2^2$.

Suppose first that $Y_3U_3S_3^2 \gg Y_2U_2S_2^2$. Then the ranges (6.2) imply that

$$|y_1u_1s_1^2 + y_2u_2s_2^2 + y_3u_3s_3^2| \ge ||y_3u_3s_3^2| - |y_1u_1s_1^2 + y_2u_2s_2^2|| \gg Y_3U_3S_3^2,$$

in any solution. Since we obviously have $|y_1u_1s_1^2 + y_2u_2s_2^2 + y_3u_3s_3^2| \ll Y_3U_3S_3^2$, the basic equation (4.8) implies that $Y_3U_3S_3^2 \ll S_0SU \ll Y_3U_3S_3^2$, whence

$$S_0 S_1 S_2 U_1 U_2 \ll Y_3 S_3 \ll S_0 S_1 S_2 U_1 U_2. \tag{6.39}$$

Summing over $Y_3 \ll S_0 S_1 S_2 S_3^{-1} U_1 U_2$, we therefore obtain

$$\sum_{S_0,S_i,U_i,Y_i,X_i,X_4} Y_3 U_3 S_3^2 \ll \sum_{S_0,S_i,U_i,Y_1,Y_2} S_0 SU,$$

where the last sum is subject to the inequality (6.28). Since there are $\ll (\log B)^2$ choices for Y_1, Y_2 , we therefore see that this sum is at most

$$\ll (\log B)^2 \sum_{S_0, S_1, S_2, U_i} S_0 S_1 S_2 U \sum_{S_3 \ll B/(S_0^3 S_1 S_2 U^2)} S_3$$
$$\ll B (\log B)^2 \sum_{S_0, S_1, S_2, U_i} S_0^{-2} U_1^{-1} U_2^{-1} U_3^{-1}$$
$$\ll B (\log B)^4,$$

as required for (6.38).

Next, if $Y_2 U_2 S_2^2 \ll Y_3 U_3 S_3^2 \ll Y_2 U_2 S_2^2$, then it follows that any choice of Y_3, U_2, U_3, S_2, S_3 determines a choice of Y_2 . Proceeding in a similar fashion to above, we deduce from (6.4) and (6.5) that $Y_3 \ll B/(U_3 U S_0^2 S_3^2)$. Hence we obtain the estimate

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} Y_3 U_3 S_3^2 \ll B \sum_{S_0, S_i, U_i, Y_1} S_0^{-2} U_1^{-1} U_2^{-1} U_3^{-1} \ll B (\log B)^4.$$

This completes the proof of (6.38).

Recall our estimate (6.24) for \mathcal{N}_2 , and note that values of X_1, X_2, X_3, X_4 are determined by the choices of S_0, S_i, U_i, Y_i , as above. In view of (6.38), it suffices to estimate

$$N^{(i)}(B) = \sum_{S_0, S_i, U_i, Y_i} \frac{S_0 S U Y_j Y_k}{\max\{Y_j U_j S_j^2, Y_k U_k S_k^2, \theta_i\}},$$
(6.40)

for each permutation $\{i, j, k\}$ of $\{1, 2, 3\}$, and where θ_i is given by (6.25). We begin by handling the case i = 3. Suppose first that $Y_3U_3S_3^2 \gg Y_2U_2S_2^2$, so that (6.39) holds and we may take $\theta_3 = Y_3U_3S_3^2$. Then

$$\frac{S_0 S U Y_1 Y_2}{\max\{Y_1 U_1 S_1^2, Y_2 U_2 S_2^2, \theta_3\}} = \frac{S_0 S_1 S_2 U_1 U_2 Y_1 Y_2}{S_3 Y_3}.$$

Moreover, we recall the inequalities

$$Y_2 \ll \frac{B}{U_2 U S_0^2 S_2^2}, \quad Y_1 \ll \frac{Y_3 U_3 S_3^2}{U_1 S_1^2}, \quad S_3 \ll \frac{S_0 S_1 S_2 U_1 U_2}{Y_3},$$

which follow from (6.5), (6.26) and (6.39), respectively. But then it follows that

$$N^{(3)}(B) \ll B \sum_{S_0, S_i, U_i, Y_1, Y_3} \frac{S_1 Y_1}{S_0 S_2 S_3 U_2 U_3 Y_3}$$
$$\ll B \sum_{S_0, S_i, U_i, Y_3} \frac{S_3}{S_0 S_1 S_2 U_1 U_2}$$
$$\ll B \sum_{S_0, S_1, S_2, U_i, Y_3} Y_3^{-1},$$

whence $N^{(3)}(B) \ll B(\log B)^6$ if $Y_3U_3S_3^2 \gg Y_2U_2S_2^2$. Next we suppose that $Y_3U_3S_3^2 \ll Y_2U_2S_2^2$, and take $\theta_3 = 1$ in (6.40). Observe that

$$S_0 \ll \frac{Y_3 S_3}{S_1 S_2 U_1 U_2}, \quad U_3 \ll \frac{Y_2 U_2 S_2^2}{Y_3 S_3^2}, \quad Y_1 \ll \frac{B}{Y_2 Y_3},$$

which follow from (6.27), the inequality $Y_3U_3S_3^2 \ll Y_2U_2S_2^2$ and (6.6), respectively. We then argue as above to deduce that

$$N^{(3)}(B) = \sum_{S_0, S_i, U_i, Y_i} S_0 S_1 S_2^{-1} S_3 U_1 U_3 Y_1$$

$$\ll \sum_{S_i, U_i, Y_i} \frac{Y_1 Y_3 U_3 S_3^2}{U_2 S_2^2}$$

$$\ll \sum_{S_i, U_1, U_2, Y_i} Y_1 Y_2$$

$$\ll B \sum_{S_i, U_1, U_2, Y_2, Y_3} Y_3^{-1}.$$

Hence $N^{(3)}(B) \ll B(\log B)^6$ in this case also.

Finally, we must estimate (6.40), whenever $i \neq 3$. Suppose that i = 1, so that

$$N^{(1)}(B) = \sum_{S_0, S_i, U_i, Y_i} \frac{S_0 S U Y_2 Y_3}{Y_3 U_3 S_3^2} = \sum_{S_0, S_i, U_i, Y_i} S_0 S_1 S_2 S_3^{-1} U_1 U_2 Y_2.$$

Once again we separate our arguments according to the size of $Y_3U_3S_3^2$. Suppose that $Y_3U_3S_3^2 \gg Y_2U_2S_2^2$, so that (6.39) holds. Then Y_3 is fixed by the choices of S_0, S_i, U_1, U_2 . Moreover, we have the inequalities

$$Y_2 \ll \frac{B}{U_2 U S_0^2 S_2^2}, \quad S_1 \ll \frac{S_0 S_2 S_3 U_2 U_3}{Y_1},$$

which follow from (6.5) and the fact that $Y_1U_1S_1^2 \ll Y_3U_3S_3^2 \ll S_0SU$, respec-

tively. Summing over Y_2 , and then over S_1 , we therefore deduce that

$$N^{(1)}(B) \ll B \sum_{S_0, S_i, U_i, Y_1} \frac{S_1}{S_0 S_2 S_3 U_2 U_3}$$
$$\ll B \sum_{S_0, S_2, S_3, U_i, Y_1} Y_1^{-1}$$
$$\ll B(\log B)^6,$$

in this case.

Alternatively, if $Y_3U_3S_3^2 \ll Y_2U_2S_2^2$, then U_3 is determined by choices of S_2, S_3, U_2, Y_2, Y_3 , and it follows that

$$N^{(1)}(B) \ll \sum_{S_0, S_i, U_1, U_2, Y_i} S_0 S_1 S_2 S_3^{-1} U_1 U_2 Y_2.$$

Upon summing over $S_0 \ll Y_3 S_3/(S_1 S_2 U_1 U_2)$, and then over $Y_2 \ll B/(Y_1 Y_3)$, we derive the estimate

$$N^{(1)}(B) \ll B \sum_{S_i, U_1, U_2, Y_1, Y_3} Y_1^{-1} \ll B(\log B)^6,$$

in this case.

An entirely similar argument handles the case i = 2. Upon combining our various estimates we therefore obtain

$$\sum_{S_0, S_i, U_i, Y_i, X_i, X_4} \mathcal{N}_2 \ll B(\log B)^6.$$

Once taken together with (6.35) and (6.37) in (6.12), we conclude that

$$N_{U,H}(B) \ll \sum_{S_0, S_i, U_i, Y_i, X_i, X_4} \mathcal{N} \ll B(\log B)^6.$$

This completes the proof of the upper bound in (1.1), and therefore completes the proof of the theorem.

References

- V. Batyrev and Y. Tschinkel. Manin's conjecture for toric varieties. J. Alg. Geom., 7 (1998), 15–53.
- [2] R. de la Bretèche. Sur le nombre de points de hauteur bornée d'une certaine surface cubique singulière. *Astérisque*, 251 (1998), 51–77.
- [3] J.W. Bruce and C.T. Wall. On the classification of cubic surfaces. J. London Math. Soc., 19 (1979), 257–267.
- [4] A. Chambert-Loir and Y. Tschinkel. On the distribution of points of bounded height on equivariant compactifications of vector groups. *Invent. Math.*, 148 (2002), no. 2, 421–452.

- [5] J.-L. Colliot-Thélène and J.-J. Sansuc. Torseurs sous des groupes de type multiplicatif; applications à l'étude des points rationnels de certaines variétés algébriques, C. R. Acad. Sci. Paris Sér. A-B, 282 (1976), 1113– 1116.
- [6] J.-L. Colliot-Thélène and J.-J. Sansuc. La descente sur les variétés rationnelles. II, Duke Math. J., 54 (1987), 375–492.
- [7] E. Fouvry. Sur la hauteur des points d'une certaine surface cubique singulière. Astérisque, 251 (1998), 31–49.
- [8] J. Franke, Y.I. Manin and Y. Tschinkel. Rational points of bounded height on Fano varieties. *Invent. Math.*, 95 (1989), 421–435.
- B. Hassett and Y. Tschinkel. Universal torsors and Cox rings, 149–173. Arithmetic of higher-dimensional algebraic varieties, Progress in Mathematics 226, Birkhäuser (2003).
- [10] D.R. Heath-Brown. Diophantine approximation with square-free numbers. Math. Zeit., 187 (1984), 335–344.
- [11] D.R. Heath-Brown. A mean value estimate for real character sums. Acta Arith., 72 (1995), 235–275.
- [12] D.R. Heath-Brown. The density of rational points on cubic surfaces. Acta Arith., 79 (1997), 17–30.
- [13] D.R. Heath-Brown and B.Z. Moroz. The density of rational points on the cubic surface $X_0^3 = X_1 X_2 X_3$. Math. Proc. Camb. Phil. Soc.. 125 (1999), 385–395.
- [14] D.R. Heath-Brown, The density of rational points on Cayley's cubic surface. Proceedings of the session in analytic number theory and Diophantine equations, Bonner Math. Schriften 360 (2003).
- [15] H.L. Montgomery, A note on the large sieve. J. London Math. Soc., 43 (1968), 93–98.
- [16] E. Peyre. Terme principal de la fonction zêta des hauteurs et torseurs universels, Astérisque, 251, (1998), 259–298.
- [17] P. Salberger. Tamagawa measures on universal torsors and points of bounded height on Fano varieties. Astérisque, 251, (1998), 91–258.
- [18] J.- P. Serre. Spécialisation des éléments de $\operatorname{Br}_2(\mathbb{Q}(T_1,\ldots,T_n))$. C. R. Acad. Sci. Paris, 311, (1990), 397–402.