

On Effective Non-vanishing of Weil Divisors on Algebraic Surfaces *

Qihong Xie

Abstract

We give a counterexample and some conclusions for effective non-vanishing of Weil divisors on algebraic surfaces.

The basepoint-free theorem plays a crucial role in the Minimal Model Program since it yields a basepoint-free complete linear system, hence an extremal contraction morphism. More precisely, it says that if D is a nef Cartier divisor such that the difference between D and the log canonical divisor is nef and big, then the linear system $|mD|$ is basepoint-free for sufficiently large integer m (cf. [6, 7]). Obviously, it is natural and interesting to investigate the non-emptiness of the linear system $|D|$. Therefore, the following so-called Effective Non-vanishing Conjecture has been put forward formally by Yujiro Kawamata(cf. [5]).

Conjecture 1. *Let X be a complete normal variety, B an effective \mathbb{R} -divisor on X such that the pair (X, B) is Kawamata log terminal, and D a Cartier divisor on X . Assume that D is nef and that $D - (K_X + B)$ is nef and big. Then $H^0(X, D) \neq 0$.*

By the Kawamata-Viehweg vanishing theorem(cf. [6]), we have $H^p(X, D) = 0$ for any positive integer p . Thus the condition $H^0(X, D) \neq 0$ is equivalent to saying that $\chi(X, D) \neq 0$. Since the Riemann-Roch formula of the line bundle $\mathcal{O}_X(D)$ depends only on the intersections of Chern classes of X and D , we can say that the Effective Non-vanishing Conjecture is indeed a topological problem in a sense of complex geometry. This conjecture is of great importance by at least two reasons. At first, it provides a good tool to study the geometry of Fano varieties(cf. [1]). Secondly, it is closely related to the general elephant problem(cf. [10]).

Kawamata has proven that the Effective Non-vanishing Conjecture holds for all log surfaces with only Kawamata log terminal singularities(cf. [5]). In dimension three, this conjecture is more complicated, only a few results are known.

It is also interesting to consider a new problem by replacing Cartier by \mathbb{Q} -Cartier.

*2000 Mathematics Subject Classification Primary 14J25; Secondary 14J45, 14E30.

Problem 2. *Notation are the same as in Conjecture 1. Assume that D is a nef \mathbb{Q} -Cartier Weil divisor on X , such that $D - (K_X + B)$ is nef and big. Does $H^0(X, D) \neq 0$ hold?*

Unfortunately, this problem does not hold generally. In dimension three, Fletcher has given the following famous example(cf. [4]).

Example 3. A general weighted complete intersection $X_{12,14}$ in $\mathbb{P}(2, 3, 4, 5, 6, 7)$ is a terminal \mathbb{Q} -Fano threefold of Picard number one with $H^0(-K_X) = 0$. This variety has the following isolated singularities: one of type $\frac{1}{5}(4, 1, 2)$, two of type $\frac{1}{3}(2, 1, 1)$, and seven of type $\frac{1}{2}(1, 1, 1)$.

In dimension two, such counterexample of Problem 2 also exists, which is the Example 7 given later in this paper. At first, we give some notation and definitions for convenience.

Definition 4. A normal surface X is said to be log terminal(resp. canonical), if the following conditions are satisfied.

- (1) K_X is a \mathbb{Q} -Cartier divisor, i.e., there exists some positive integer r such that rK_X is Cartier, and
- (2) For any resolution $f : Y \rightarrow X$, if we write $K_Y = f^*K_X + \sum \alpha_i E_i$, where E_i are irreducible exceptional curves of f , then $\alpha_i > -1$ (resp. $\alpha_i \geq 0$) for all i .

The index of X is, by definition, the smallest positive integer r such that rK_X is Cartier. If the index of X is one, then X is said to be Gorenstein. If the resolution f is minimal, i.e., K_Y is f -nef, then we have that $\alpha_i \leq 0$ for all i since the intersection matrix $(E_i \cdot E_j)$ is negative definite.

Definition 5. A normal surface X is called a log del Pezzo surface, if X has only log terminal singularities and $-K_X$ is ample.

For a \mathbb{Q} -divisor $D = \sum a_i D_i$, the round up(resp. round down) of D is defined to be $\lceil D \rceil = \sum \lceil a_i \rceil D_i$ (resp. $\lfloor D \rfloor = \sum \lfloor a_i \rfloor D_i$), where $\lceil a \rceil$ (resp. $\lfloor a \rfloor$) is the smallest integer not less(resp. the largest integer not greater) than a .

Proposition 6. *Let X be a log terminal surface, $f : Y \rightarrow X$ the minimal resolution of X . Assume that $K_Y = f^*K_X + \sum \alpha_i E_i$, where $-1 < \alpha_i \leq 0$ for all i . Then for each integer $n \geq 0$, $h^0(-nK_X) := \dim H^0(X, \mathcal{O}_X(-nK_X))$ is equal to*

$$\frac{n(n+1)}{2} K_Y^2 - \frac{2n+1}{2} K_Y \cdot \left(\sum \lceil (n+1)\alpha_i \rceil E_i \right) + \frac{1}{2} \left(\sum \lceil (n+1)\alpha_i \rceil E_i \right)^2 + \chi(\mathcal{O}_Y).$$

Proof. Let $D = -nK_X$. Then we have

$$\begin{aligned} f^*(D - K_X) &= -(n+1)f^*K_X = -(n+1)(K_Y - \sum \alpha_i E_i) \\ &= (-nK_Y + (n+1) \sum \alpha_i E_i) - K_Y \end{aligned}$$

We define \tilde{D}, \tilde{B} as follows.

$$\begin{aligned}\tilde{D} &= [-nK_Y + (n+1) \sum \alpha_i E_i] = -nK_Y + \sum [(n+1)\alpha_i] E_i, \\ \tilde{B} &= \sum ([(n+1)\alpha_i] - (n+1)\alpha_i) E_i.\end{aligned}$$

It is easy to see that $f^*(D - K_X) = \tilde{D} - (K_Y + \tilde{B})$ is an f -nef and f -big divisor, and \tilde{B} has simple normal crossing support. By the Kawamata-Viehweg vanishing theorem, we have $R^j f_* \mathcal{O}_Y([\tilde{D} - \tilde{B}]) = R^j f_* \mathcal{O}_Y(\tilde{D}) = 0$ for any positive integer j .

Since we have

$$\begin{aligned}f^*D &= -nK_Y + \sum n\alpha_i E_i = \tilde{D} + \sum (n\alpha_i - [(n+1)\alpha_i]) E_i, \\ n\alpha_i - [(n+1)\alpha_i] &< (n+1)\alpha_i - [(n+1)\alpha_i] + 1 \leq 1,\end{aligned}$$

there exists a \mathbb{Q} -divisor $H = \sum h_i E_i$ on Y such that $0 \leq h_i < 1$ and $h_i - (n\alpha_i - [(n+1)\alpha_i]) \geq 0$ for each i . For a \mathbb{Q} -divisor M on Y , the sheaf $\mathcal{O}_Y(M)$ is defined to be $\mathcal{O}_Y(\lfloor M \rfloor)$. Thus $f_* \mathcal{O}_Y(\tilde{D}) = f_* \mathcal{O}_Y(\tilde{D} + H) = \mathcal{O}_X(D)$ by the projection formula. It follows from [3] that we have the Leray spectral sequence

$$E_2^{i,j} = H^i(X, R^j f_* \mathcal{O}_Y(\tilde{D})) \Rightarrow H^{i+j}(Y, \mathcal{O}_Y(\tilde{D})).$$

Since $E_2^{i,j} = 0$ for all $i \geq 0$ and $j > 0$, $E_2^{i,0} \cong E_\infty^{i,0}$ implies that $h^i(X, \mathcal{O}_X(D)) = h^i(Y, \mathcal{O}_Y(\tilde{D}))$ for all $i \geq 0$, hence $\chi(X, \mathcal{O}_X(D)) = \chi(Y, \mathcal{O}_Y(\tilde{D}))$. Therefore we have

$$\begin{aligned}& h^0(-nK_X) \\ &= \chi(X, \mathcal{O}_X(D)) = \chi(Y, \mathcal{O}_Y(\tilde{D})) = \frac{1}{2} \tilde{D}(\tilde{D} - K_Y) + \chi(\mathcal{O}_Y) \\ &= \frac{1}{2} (-nK_Y + \sum [(n+1)\alpha_i] E_i) (-nK_Y + \sum [(n+1)\alpha_i] E_i) + \chi(\mathcal{O}_Y) \\ &= \frac{n(n+1)}{2} K_Y^2 - \frac{2n+1}{2} K_Y \cdot (\sum [(n+1)\alpha_i] E_i) + \frac{1}{2} (\sum [(n+1)\alpha_i] E_i)^2 + \chi(\mathcal{O}_Y).\end{aligned}$$

□

Example 7. There exists a log del Pezzo surface S of Picard number one such that $H^0(S, -K_S) = 0$.

Let B be a smooth conic in \mathbb{P}^2 , D a tangent line to B at d , and A a secant line intersecting B at $\{a, b\}$. Let $g_1 : S_1 \rightarrow \mathbb{P}^2$ be the blow-up of \mathbb{P}^2 along d on D with an exceptional curve L_1 . By abuse of notation, we always denote the strict transform of a curve by the same symbol. Let $g_i : S_{i+1} \rightarrow S_i$ be the blow-up of S_i along d on D with exceptional curves L_{i+1} for $i = 1, 2$. It is easy to see that $L_1^2 = L_2^2 = -2, L_3^2 = -1$. In a similar way, let $g_4 : S_4 \rightarrow S_3$ be the blow-up of S_3 5 time along b on B .

Assume that the exceptional curves of g_4 are denoted by M_1, M_2, M_3, M_4, M_5 respectively with $M_1^2 = M_2^2 = M_3^2 = M_4^2 = -2$, and $M_5^2 = -1$. Let $g_5 : \tilde{S} \rightarrow S_4$ be the blow-up of S_4 5 time along a on A . Assume that the exceptional curves of g_5 are denoted by N_1, N_2, N_3, N_4, N_5 respectively with $N_1^2 = N_2^2 = N_3^2 = N_4^2 = -2$, and $N_5^2 = -1$. Denote by $g : \tilde{S} \rightarrow \mathbb{P}^2$ the composite of all g_i .

We consider two reduced connected curves $X_1 := D \cup A \cup M_1 \cup M_2 \cup M_3 \cup M_4$ and $X_2 := L_1 \cup L_2 \cup B \cup N_1 \cup N_2 \cup N_3 \cup N_4$ on \tilde{S} . It is easy to verify that the intersection matrix associated with X_i is negative definite, and the arithmetic genus of any positive cycle with support on X_i is non-positive for $i = 1, 2$. Thus $X_1 \cup X_2$ is contractible, and there is a contraction $f : \tilde{S} \rightarrow S$ such that S is a normal projective surface of Picard number one with only two singular points x_1, x_2 , which is the contraction point of X_1, X_2 respectively (cf. [2]).

It is easy to see that \tilde{S} is just the minimal resolution of S , and the dual graph Γ consists of two connected components corresponding to x_1, x_2 respectively.

$$\Gamma: \begin{array}{cccccccccccccccc} 2^D & 5^A & 2^{M_1} & 2^{M_2} & 2^{M_3} & 2^{M_4} & 2^{L_1} & 2^{L_2} & 4^B & 2^{N_1} & 2^{N_2} & 2^{N_3} & 2^{N_4} \\ \bullet & \bullet \end{array}$$

where 2^D denotes the (-2) -curve on \tilde{S} which is the strict transform of D , and so on.

By an elementary calculation, we have

$$\begin{aligned} K_{\tilde{S}} &= f^*K_S - \frac{15}{37}D - \frac{30}{37}A - \frac{24}{37}M_1 - \frac{18}{37}M_2 - \frac{12}{37}M_3 - \frac{6}{37}M_4 \\ &\quad - \frac{5}{19}L_1 - \frac{10}{19}L_2 - \frac{15}{19}B - \frac{12}{19}N_1 - \frac{9}{19}N_2 - \frac{6}{19}N_3 - \frac{3}{19}N_4 \end{aligned}$$

It follows easily that $K_{\tilde{S}}^2 = 8/(37 \times 19) > 0$, then $-K_S$ is ample since S is rational and $\rho(S) = 1$. Furthermore, we have $H^0(S, -nK_S) = 0$ for $1 \leq n \leq 5$, but $H^0(S, -6K_S) \cong \mathbb{C}$ by Proposition 6.

Remark 8. For a log del Pezzo surface X , we define $\tau(X)$ to be the smallest positive integer τ such that $H^0(X, -\tau K_X) \neq 0$. It is easy to see that the index $r(X)$ of X is not less than $\tau(X)$. The study of complements on surfaces is helpful to solve the non-vanishing of $H^0(X, -nK_X)$ for log del Pezzo surfaces. Shokurov proved that there exists a positive integer N such that $\tau(X) \leq N$ for all log del Pezzo surfaces (cf. [11]). In particular, Prokhorov used the same method to yield that $H^0(X, -nK_X) \neq 0$ for some $n \in \mathcal{R}_2 := \{1, 2, 3, 4, 6\}$, if $K_X^2 > 4$ (cf. [9]).

Example 7 illustrates that Problem 2 does not hold generally for surfaces. But, when some reasonable condition is assumed, we hope that some effective non-vanishing of Weil divisors is available. In particular, we will prove that

effective non-vanishing of Weil divisors holds for Gorenstein log del Pezzo surfaces of Picard number 1.

Theorem 9. *Let X be a Gorenstein log del Pezzo surface with Picard number $\rho(X) = 1$, D a nef and big Weil divisor on X . Then $H^0(X, D) \neq 0$.*

It is easy to see that X has only canonical singularities. The condition $\rho(X) = 1$ enable us to list all type of its singularities.

Lemma 10. *Let X be a Gorenstein log del Pezzo surface with $\rho(X) = 1$. Then its singularity type is one of the following*

$A_1, A_1 + A_2, A_4, 2A_1 + A_3, D_5, A_1 + A_5, 3A_2, E_6, 3A_1 + D_4, A_7, A_1 + D_6, E_7, A_1 + 2A_3, A_2 + A_5, D_8, 2A_1 + D_6, E_8, A_1 + E_7, A_1 + A_7, 2A_4, A_8, A_1 + A_2 + A_5, A_2 + E_6, A_3 + D_5, 4A_2, 2A_1 + 2A_3, 2D_4$.

Proof. X is said to be of type, for example, $2A_1 + A_3$, if X has three singular points, two of type A_1 , and one of type A_3 . We refer to [8] for this proof. \square

Definition 11. The quotient $X = \mathbb{A}^n / \mu_r$ is said to be of cyclic quotient singularity of type $\frac{1}{r}(a_1, \dots, a_n)$, if μ_r acts on \mathbb{A}^n by

$$\mu_r \ni \epsilon : (x_1, \dots, x_n) \mapsto (\epsilon^{a_1} x_1, \dots, \epsilon^{a_n} x_n).$$

Let $\pi : \mathbb{A}^n \rightarrow X$ be the quotient morphism. Then the group μ_r acts on $\pi_* \mathcal{O}_{\mathbb{A}^n}$, and so decomposes it into r eigensheaves $\mathcal{L}_i = \{f | \epsilon(f) = \epsilon^i \cdot f \text{ for all } \epsilon \in \mu_r\}$ for $i = 0, \dots, r-1$. A singularity $p \in X$ with a Weil divisor D is said to be of cyclic quotient singularity of type $\frac{1}{r}(a_1, \dots, a_n)$, if locally, $p \in X$ is isomorphic to a point of type $\frac{1}{r}(a_1, \dots, a_n)$, and $\mathcal{O}_X(D) \cong \mathcal{L}_i$.

There are two keypoints in the proof of Theorem 9. First, one canonical singularity can be considered as the cyclic quotient of another canonical singularity by using the local cyclic cover. Second, we can calculate the Euler characteristic of a Weil divisor, thanks to the Singular Riemann-Roch Formula.

The following lemma is helpful, which can be found in [10].

Lemma 12. *Let $Q \in Y : (f = 0) \subset \mathbb{A}^3$ be a surface germ. Assume that there is an action of μ_r on Y free outside Q . Let $p \in X = (Q \in Y) / \mu_r$ be the surface germ and $\pi : Y \rightarrow X$ the corresponding cyclic cover. If $p \in X$ is canonical, then $Q \in Y$ is also canonical and $\pi : Y \rightarrow X$ is exactly one of the following types.*

	r	Type	f	Description
(1)	any	$\frac{1}{r}(1, -1, 0)$	$xy + z^n$	$A_{n-1} \xrightarrow{r:1} A_{rn-1}$
(2)	4	$\frac{1}{4}(1, 3, 2)$	$x^2 + y^2 + z^{2n-1}$	$A_{2n-2} \xrightarrow{4:1} D_{2n+1}$
(3)	2	$\frac{1}{2}(0, 1, 1)$	$x^2 + y^2 + z^{2n}$	$A_{2n-1} \xrightarrow{2:1} D_{n+2}$
(4)	3	$\frac{1}{3}(0, 1, 2)$	$x^2 + y^3 + z^3$	$D_4 \xrightarrow{3:1} E_6$
(5)	2	$\frac{1}{2}(1, 1, 0)$	$x^2 + y^2 z + z^n$	$D_{n+1} \xrightarrow{2:1} D_{2n}$
(6)	2	$\frac{1}{2}(1, 0, 1)$	$x^2 + y^3 + z^4$	$E_6 \xrightarrow{2:1} E_7$

Proof of Theorem 9. If D is a Cartier divisor, then the non-vanishing follows from [5]. Now assume that D is not Cartier. Since any normal projective surface with log terminal singularities is \mathbb{Q} -factorial(cf. [7]), we can define the index $r(D)$ of Weil divisor D to be the smallest positive integer r such that rD is Cartier. For $p \in X$, we define the local index $r_p(D)$ to be the smallest positive integer r such that rD is Cartier locally around p . It is easy to see that $r(D) = \text{l.c.d.}\{r_p(D)\}_{p \in \text{Sing}X}$.

For the reflexive sheaf $\mathcal{O}_X(D)$, we have the following Singular Riemann-Roch Formula(cf. [10]):

$$\chi(X, \mathcal{O}_X(D)) = \frac{1}{2}D \cdot (D - K_X) + \chi(\mathcal{O}_X) + \sum_{p \in X} C_p(D),$$

where $c_p(D) \in \mathbb{Q}$ is a contribution due to the singularities of $\mathcal{O}_X(D)$ at p , the sum is taken over the points $p \in X$ where D is not Cartier.

If $p \in X$ and D is of type $i(\frac{1}{r}(1, -1))$, then $c_p(D) = -i(r - i)/2r$. For other cases, we use the suitable \mathbb{Q} -smoothing of X , then there is a flat deformation $\{X_\lambda, D_\lambda\}$ of $\{X, D\}$ such that X_λ has only cyclic quotient singularities. Thus $C_p(X)$ is equal to the sum of the contributions of the cyclic quotient singularities of X_λ . The concrete conclusions are listed below.

	X_λ	\mathbb{Q} -smoothing points
(1)	$f + \lambda z$	n
(2)	$f + \lambda z$	2n+1
(3)	$f + \lambda x$	2
(4)	$f + \lambda x$	2
(5)	$f + \lambda z$	n
(6)	$f + \lambda y$	3

Since $-K_X$ is an ample Cartier divisor on X , it follows that $\chi(\mathcal{O}_X) = 1$ and $D \cdot (-K_X) \geq 1$. Thus $h^0(X, D) \geq D^2/2 + 3/2 + \sum_{p \in X} C_p(D)$. Next we will estimate $C_p(D)$ for a given germ $p \in X$. Since $C_p(D)$ depends only on the local analytic type of $p \in X$ and D , we can take $\pi : Y \rightarrow X$ to be the cyclic cover of order $r_p(D)$ corresponding to D locally around $p \in X$. By Lemma 12, $Q \in Y$ has only canonical singularity, and we can calculate the contribution as follows.

Let $p \in X$ be of type A_{n-1} . Then $p \in X$ is a cyclic quotient singularity of type $\frac{1}{n}(1, -1)$. By the definition of the contribution, we have $C_p(D) = -i(n - i)/2n$. If $n = 2k$, then $C_p(D) \geq -k/4$. If $n = 2k + 1$, then $C_p(D) \geq -k(k + 1)/2(2k + 1)$.

Let $p \in X$ be of type D_{n+2} . We may consider $p \in X$ as a cyclic quotient of type (3). Then $C_p(D) \geq (-i(2 - i)/4) \times 2 \geq -1/2$.

Let $p \in X$ be of type E_6 . We may consider $p \in X$ as a cyclic quotient of type (4). Then $C_p(D) \geq (-i(3 - i)/6) \times 2 \geq -2/3$.

Let $p \in X$ be of type E_7 . We may consider $p \in X$ as a cyclic quotient of type (6). Then $C_p(D) \geq (-i(2-i)/4) \times 3 \geq -3/4$.

Let $p \in X$ be of type E_8 . Then $r_p(D) = 1$ for any Weil divisor D , and $C_p(D) = 0$.

It is easy to check for all cases in Lemma 10 that $\sum_{p \in X} C_p(D) \geq -3/2$. Hence $h^0(X, D) \geq D^2/2 > 0$. \square

Acknowledgment I would like to express my gratitude to Professor Yujiro Kawamata for his valuable advice and warm encouragement. I also thank Professor S. A. Kudryavtsev for informing me some known results in Remark 8, and Dr. Hokuto Uehara for stimulating discussions on this paper.

References

- [1] F. Ambro, Ladders on Fano varieties, *Algebraic geometry, 9. J. Math. Sci. (New York)*, **94**(1999), no. 1, 1126-1135.
- [2] M. Artin, Some numerical criteria for contractability of curves on algebraic surfaces, *Amer. J. of Math.*, **84**(1962), 485-496.
- [3] G. Bredon, *Sheaf theory*, Springer, 1997.
- [4] A. R. Fletcher, Working with weighted complete intersections, *Explicit birational geometry of 3-folds*, Lond. Math. Soc. Lect. Note Ser., vol. 281(2000), 101-173.
- [5] Y. Kawamata, On effective non-vanishing and base-point-freeness, *Asian J. Math.*, **4**(2000), 173-182.
- [6] Y. Kawamata, K. Matsuda, K. Matsuki, Introduction to the minimal model problem, *Alg. Geom. Sendai 1985, Adv. Stud. in Pure Math.*, **10**(1987), Kinokuniya, Tokyo, 283-360.
- [7] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math, vol. 134(1998).
- [8] M. Miyanishi, D. Q. Zhang, Gorenstein log del Pezzo surfaces of rank one, *Journal of Algebra*, **118**(1988), 63-84.
- [9] Y. G. Prokhorov, *Lectures on complements on log surfaces*, Math. Soc. of Japan Memoirs, vol. 10(2001).
- [10] M. Reid, Young person's guide to canonical singularities, *Proc. Symp. Pure Math.* **46**(1987), 345-416.

- [11] V. V. Shokurov, Complements on surfaces, *Algebraic geometry, 10. J. Math. Sci. (New York)*, **102**(2000), no. 2, 3876–3932.

QIHONG XIE xqh@ms.u-tokyo.ac.jp
GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO,
KOMABA, MEGURO, TOKYO 153-8914, JAPAN