Geometric stability of the cotangent bundle and the universal cover of a projective manifold

Frédéric Campana and Thomas Peternell

may 1, 2004

Introduction

Let X_n be a complex projective *n*-dimensional manifold and \tilde{X} its universal cover. The Shafarevitch conjecture asserts that \tilde{X} is holomorphically convex, i.e. admits a proper holomorphic map onto a Stein space. There are two extremal cases, namely that this map is constant, i.e. \tilde{X} is compact. This means that $\pi_1(X)$ is finite and not much can be said further. Or the map is a modification, i.e through the general point of \tilde{X} there is no positive-dimensional compact subvariety, e.g. \tilde{X} is Stein. This happens in particular for X an Abelian variety or a quotient of a bounded domain. It is conjectured (see [Ko93], and [CZ04] for the Kähler case) that X should then have a holomorphic submersion onto a variety of general type with Abelian varieties as fibres, after a suitable finite étale cover and birational modification. This follows up to dimension 3 from the solutions of the conjectures of the Minimal Model Program. We prove here a weaker statement in every dimension:

Theorem 0.1 Let X_n be a projective manifold and suppose that its universal cover is not covered by its positive-dimensional compact subvarieties. Then either X is of general type, or $\chi(\mathcal{O}_X) = 0$. In particular, if \tilde{X} is Stein (or does not contain a compact subvariety of positive dimension), then either K_X is ample, or: K_X is nef, $K_X^n = 0$, and $\chi(\mathcal{O}_X) = 0$.

This theorem is deduced (via the comparison theorem [Ca95], which relates the geometric positivity of subsheaves in the cotangent bundle to the geometry of \tilde{X}) from the following more general:

Theorem 0.2 Let X_n be a projective manifold. Suppose that Ω_X^p contains for some p a subsheaf whose determinant is big, i.e. has maximal Kodaira dimension n. Then K_X is big, i.e. $\kappa(X) = n$.

This in turn follows from a slight generalization of Miyaoka's theorem that the cotangent bundle of a projective manifold is "generically nef" unless the manifold is uniruled, and from a characterization of pseudo-effective line bundles by moving curves [BDPP04]. We show indeed that quotients of Ω_X^p have a pseudo-effective

determinant if X is not uniruled. This uniruledness criterion has also other applications, e.g. one can prove that a variety admitting a section in a tensor power of the tangent bundle with a zero, must be uniruled.

Theorem 0.2 is actually a piece in a larger framework. To explain this, we consider subsheaves $\mathcal{F} \subset \Omega_X^p$ for some p > 0. Then one can form $\kappa(\det \mathcal{F})$ and take the supremum over all \mathcal{F} . This gives a modified Kodaira dimension $\kappa^+(X)$, introduced in [Ca95]. Conjecturally

$$\kappa^+(X) = \kappa(X)$$

unless X is uniruled. Theorem 0.2 is nothing than this conjecture in case $\kappa^+(X) = \dim X$. We shall prove the conjecture in several cases, e.g. in dimension 4. It is very much related to (actually a consequence of) the following:

Conjecture: Suppose X is a projective manifold, and suppose a decomposition

$$NK_X = A + B$$

with some positive integer N and a pseudo-effective line bundle B (one may assume A spanned). Then

$$\kappa(X) \ge \kappa(A).$$

The special case $A = \mathcal{O}_X$ implies that $\kappa(X) \ge 0$ if X is not uniruled, using the preceding result, and the pseudo-effectiveness of K_X when X is not uniruled ([BDPP04]).

In another direction we obtain the special case in which B is numerically trivial:

Theorem 0.3 Let X be a projective complex manifold, and $L \in Pic(X)$ be numerically trivial. Then:

- 1. $\kappa(X, K_X + L) \leq \kappa(X)$.
- 2. If $\kappa(X) = 0$, and if $\kappa(X, K_X + L) = \kappa(X)$, then L is a torsion element in the group $\operatorname{Pic}^0(X)$.

In particular, if mK_X is numerically equivalent to an effective divisor, then $\kappa(X) \ge 0$.

Acknowledgement. Our collaboration has been made possible by the priority program "Global methods in complex geometry" of the Deutsche Forschungsgemeinschaft, which we gratefully acknowledge.

Contents

1	Uniruledness Criteria	3
2	 A characterization of varieties of general type 2.1 Refined Kodaira Dimension	7
3	Numerical properties of the Kodaira dimension	11
4	The Universal Cover	15

1 Uniruledness Criteria

Definition 1.1 Let X_n be a projective manifold. A vector bundle E over X is generically nef, if for all ample line bundles H_1, \ldots, H_{n-1} , for all m_i sufficiently large and for general curves C cut out by $m_1H_1, \ldots, m_{n-1}H_{n-1}$, the bundle E|C is nef.

The importance of this notion lies in the following result due to Miyaoka [Mi87], with a short proof by [SB92].

Theorem 1.2 The cotangent bundle bundle of a projective manifold is generically nef unless X is uniruled.

We shall need the following generalization.

Theorem 1.3 Let X be a projective manifold and suppose that X is not uniruled. Let $\pi : \tilde{X} \to X$ be a birational map from a projective manifold \tilde{X} . Then $\pi^*(\Omega^1_X)$ is generically nef.

Proof. We follow the line of argumentation in [SB92]. Let $n = \dim X$. Let H_1, \ldots, H_{n-1} be ample divisors on \tilde{X} , furthermore m_i sufficiently large integers and $\tilde{C} \subset \tilde{X}$ a general curve cut out by the $m_i H_i$. Fix furthermore an ample line bundle H on X.

Now suppose the theorem does not hold for H_1, \ldots, H_{n-1} . Then we find a reflexive subsheaf $\tilde{\mathcal{E}} \subset \pi^*(T_X)$ such that

$$c_1(\tilde{\mathcal{E}}) \cdot C > 0.$$

Then choose as in [SB92,9.1.5] a subsheaf $\tilde{\mathcal{F}} \subset \pi^*(T_X)$ which is a part of the Harder-Narasimhan filtration with minimal slope $\mu(\pi^*(T_X)|C)$. Let

$$\tilde{\mathcal{G}} = \pi^*(T_X)/\tilde{\mathcal{F}}.$$

This is a torsion free sheaf on \tilde{X} . Furthermore we introduce

$$\mathcal{F} = \pi_* (\tilde{\mathcal{F}})^{**}.$$

Since $\pi_*(\tilde{\mathcal{F}}) \subset \pi_*\pi^*(T_X) = T_X$, it is clear by Riemann's extension theorem that

$$\mathcal{F} \subset T_X.$$

Finally let $\mathcal{G} = T_X/\mathcal{F}$. Now [SB92,9.1.3.1] gives

$$\operatorname{Hom}(\bigwedge^2 \tilde{\mathcal{F}}, \tilde{\mathcal{G}}) = 0.$$

Let $U = X \setminus \text{center of } \pi$. Then we conclude

$$\operatorname{Hom}(\bigwedge^2 \mathcal{F}|U,\mathcal{G}|U) = 0$$

and thus

$$\operatorname{Hom}(\bigwedge^2 \mathcal{F}, \mathcal{G}) = 0$$

As in [SB92] we now reduce to char p. Let $F: X \to X$ and $\tilde{F}: \tilde{X} \to \tilde{X}$ be the absolute Frobenius maps. Then for $p \gg 0$, we see as in [SB92,9.1.3.5] that

$$\operatorname{Hom}(\tilde{F}^*(\tilde{\mathcal{F}}),\tilde{\mathcal{G}})=0$$

and therefore, as before

 $\operatorname{Hom}(F^*(\mathcal{F}),\mathcal{G})=0$

so that \mathcal{F} is a 1-foliation in the terminology of [SB92]. Hence one can form the quotient $\rho: X \to Y = X/\mathcal{F}$ and obtains a line bundle G on Y with $H = \rho^*(G)$. Furthermore, introducing $C = \pi(\tilde{C})$, we have

$$C \cdot \rho^*(-K_Y) = p(C \cdot c_1(\mathcal{F})) + (C \cdot c_1(\mathcal{G}))$$

Then we proceed verbatim as in [SB92] to produce rational curves through the general point of X with a bound on the G-degree independent of p. Thus X is also uniruled in char 0, which produces the contradiction we are looking for. Q.E.D.

Now we can strengthen the preceeding result, using [BDPP04] (and answering a question asked in that paper).

First recall that a line bundle L on a projective manifold is called *pseudo-effective* iff $c_1(L)$ is in the closure of the cone generated by the (numerical equivalence classes of the) effective divisors on X.

We will need the following result from [BDPP04] which will also be crucial for Theorem 2.3.

Theorem 1.4 Let X_n be a projective manifold and L a line bundle on X. Then L is pseudo-effective if and only if the following holds. Let $\pi : \hat{X} \to X$ be a birational map from a projective manifold X. Let H_1, \ldots, H_{n-1} be very ample. Then

$$L \cdot \pi_*(H_1 \cap \ldots \cap H_{n-1}) \ge 0.$$

Together with theorem (1.3), this implies:

Theorem 1.5 Let X be a projective manifold and suppose that X is not uniruled. Let Q be a torsion free quotient of $(\Omega_X^1)^{\otimes m}$ for some m > 0. Then det Q is pseudoeffective.

Here we made use of the following

Notation 1.6 Let \mathcal{F} be a coherent sheaf of rank r on the connected manifold X. We define its determinant - a line bundle - to be

$$\det \mathcal{F} = (\bigwedge^r \mathcal{F})^{**}.$$

Proof. In order to show the pseudo-effectivity of det Q, it suffices by (1.4) to verify the following. Let $\pi : \tilde{X} \to X$ be birational from the projective manifold \tilde{X} . Let H_1, \ldots, H_{n-1} be very ample on \tilde{X} (and general in their linear systems). Then

$$\det Q \cdot \pi_*(H_1 \cap \ldots \cap H_{n-1}) \ge 0. \tag{(*)}$$

So let us verify (*). Eventually after substituting H_i by large multiples and by setting

$$\tilde{C} = H_1 \cap \ldots \cap H_{n-1},$$

Theorem 1.3 applies and $\pi^*\Omega^1_X|\tilde{C}$ is nef. Hence $\pi^*(Q)|C$ is nef, therefore also $(\pi^*(Q)/\text{torsion})|\tilde{C}$ is nef, so that $(\det \pi^*(Q))|C$ is nef. Denoting r the (generic) rank of Q, we have $\wedge^r(\pi^*(Q)) = \pi^*(\wedge^r Q)$, hence we obtain a canonical map

$$\bigwedge' \pi^*(Q) \to \pi^*(\det Q).$$

This yields an inclusion

$$\det \pi^*(Q) \subset \pi^*(\det Q).$$

Hence $\pi^*(\det Q)|\tilde{C}$ is nef, too, and (*) is verified. This finishes the proof.

Q.E.D.

Now a pseudo-effective line bundle is nef on moving curves; here "moving" means that the deformations of the curve cover the variety. Actually by [BDPP04] the closed cone generated by by numerical equivalence classes of movable curves coincides with the cone generated by classes of "strongly movable" curves. These are just the curves of the form $\pi(\hat{C})$, where $\pi: \hat{X} \to X$ is a modification, and $\hat{C} \subset \hat{X}$ is a generic intersection of very ample divisors $m_i H_i, 1 \leq i \leq n-1$ on \hat{X} . So we can state: **Corollary 1.7** Let X_n be a projective manifold and suppose that X is not uniruled. Let $(C_t)_{t\in T}$ be an algebraic family of curves, parametrised by the irreducible projective variety T. Assume this family is covering (i.e.: the union of the C_t 's is X, and its generic member is irreducible).

Let \mathcal{F} be a torsion free quotient of $(\Omega^1_X)^{\otimes m}$ for some m > 0. Then $c_1(\mathcal{F}).C_t \ge 0$.

A classical result in group actions on a projective manifold X says that if X carries a holomorphic vector field with zeroes, then X is uniruled. Using Theorem 1.3 we are able to generalize this to arbitrary tensor powers of the tangent bundle, and we may also allow a twist with a topologically trivial line bundle.

Theorem 1.8 Let X be a projective manifold and L a topologically trivial line bundle on X. Let m be a positive integer and

$$v \in H^0(T_X^{\otimes m} \otimes L)$$

a section with zeroes. Then X is uniruled.

Proof. Choose $p \in X$ such that v(p) = 0. Let $\pi : \hat{X} \to X$ be the blow-up of X at p. Assume that X is not uniruled. Then by Theorem 1.3, $\pi^*(\Omega^1_X)$ is generically nef. Hence if \hat{C} is the curve cut out by sufficiently general very ample divisors, then $\pi^*(\Omega^1_X)|\hat{C}$ is nef. Thus $\Omega^1_X|C$ is nef, where $C = \pi(\hat{C})$. Now \hat{C} meets the exceptional divisor of π in a finite set, hence $p \in C$. In total, $(\Omega^1_X)^{\otimes m} \otimes L^*|C$ is nef, but its dual has a section with zeroes. This is impossible. So X is uniruled. Q.E.D.

Now observe that the same proof as in (1.7) works if L^* is assumed to be pseudoeffective instead of L being topologically trivial. More generally, if $\mathcal{F} \subset T_X^{\otimes m}$ is a coherent subsheaf of rank r such that det $\mathcal{F} \to \bigwedge^r(T_X^{\otimes m})$ has a zero and if $(\det \mathcal{F})^*$ is pseudo-effective, then we can view the image of det $\mathcal{F} \to \bigwedge^r(T_X^{\otimes m})$ as a subsheaf of some $(T_X)^{\otimes m}$ and then apply (1.7) to obtain

Corollary 1.9 Let X be a projective manifold, $\mathcal{F} \subset T_X^{\otimes m}$ a subsheaf of rank r such that

$$\det \mathcal{F} \to \bigwedge^r (T_X^{\otimes m})$$

has a zero. Assume that det \mathcal{F} is pseudo-effective. Then X is uniruled.

2 A characterization of varieties of general type

2.1 Refined Kodaira Dimension

The following "refined Kodaira dimension" was introduced in [Ca95]. It measures the geometric positivity of the cotangent bundle, and not only that of the canonical bundle. (Its definition is justified in the next subsection). **Definition 2.1** Let X be a compact (or projective) manifold. Then $\kappa^+(X)$ is the maximal number $\kappa(\det \mathcal{F})$, where $\mathcal{F} \subset \Omega_X^p$ for $1 \leq p \leq \dim X$ is a (saturated) coherent subsheaf.

Obviously we have $\kappa^+(X) \ge \kappa(X)$ for any X.

Assuming the standard conjectures of the Minimal Model Program, one can easily describe $\kappa^+(X)$ as follows (see [Ca95] for details, where the following conjecture was formulated):

Conjecture 2.2 Let X be a projective manifold. If X is not uniruled (or if $\kappa(X) \ge 0$), then $\kappa^+(X) = \kappa(X)$.

When X is uniruled, one has

$$\kappa^+(X) = \kappa^+(R(X)),$$

where R(X) is the so-called "rational quotient" of X; see [Ca95]. This rational quotient is not uniruled, and so should be either one point or have $\kappa^+(R(X)) = \kappa(R(X)) \ge 0$. Thus if X is uniruled, one has $\kappa(X) = -\infty$ but $\kappa^+(X) \ge 0$, unless R(X) is one point, which means that X is rationally connected. In this latter case $\kappa^+(X) = -\infty$. Conversely, if $\kappa^+(X) = -\infty$, then X should be rationally connected. Notice that $\chi(\mathcal{O}_X) = 1$ if $\kappa^+(X) = -\infty$, because $h^0(X, \Omega_X^p) = 0$ for p > 0. In [Ca95] it is shown that X is simply connected if $\kappa^+(X) = -\infty$ which of course is also true for X rationally connected.

The above conjecture is a geometric version of the stability of the cotangent bundle of X when X is not uniruled. It is a version in which positivity of subsheaves is measured by the Kodaira dimension of the determinant bundle, and not by the slope after restricting to "strongly movable curves".

2.2 A Characterisation of Varieties of General Type

As a consequence of the preceeding criteria for uniruledness, we first solve the above conjecture in the extremal case when $\kappa^+(X) = n$ (we shall study in the next section below the intermediate cases):

Theorem 2.3 Let X_n be a projective manifold and suppose $\kappa^+(X) = n$, i.e. some Ω^p_X contains a subsheaf \mathcal{F} with $\kappa(\det \mathcal{F}) = n$. Then $\kappa(X) = n$.

Proof. First let us see that X is not uniruled. In fact, otherwise take a covering family of rational curves and select a general member C so that $T_X|C$ is nef. Hence the dual of $\Omega_X^p|C$ is nef and therefore $\mathcal{F}|C$ cannot have ample determinant. So X cannot be uniruled.

Of course, we may assume that \mathcal{F} saturated, hence $Q = \Omega_X^p / \mathcal{F}$ is torsion free. By taking determinants we get

$$mK_X = \det \mathcal{F} + \det Q$$

for some positive integer m. We learn from (1.6) above that det Q is pseudo-effective. Thus K_X is big, as a sum of a big and a pseudo-effective divisor.

Q.E.D.

2.3 The intermediate case

In this section we want to study the above conjecture 2.2 in the intermediate case $n > \kappa(X_n) \ge 0$.

We shall reduce Conjecture 2.2 to (special cases of) a seemingly considerably simpler:

Conjecture 2.4 Let X be a projective manifold. Let $NK_X = A + B$ with some positive integer N > 0, A effective and B pseudo-effective. Then $\kappa(X) \ge \kappa(A)$.

Remark 2.5 (1) By suitably blowing up, it is easily seen that Conjecture 2.4 is equivalent to the analogous conjecture with A always assumed to be spanned.

(2) If $\nu(L)$ denotes the numerical dimension of an arbitrary pseudo-effective line bundle as introduced by Boucksom [Bo02], then the generalised abundance conjecture states

$$\kappa(K_X) = \nu(K_X).$$

If this generalised abundance conjecture holds, then Conjecture 2.4 holds when $\kappa(X) = 0$, a case sufficient to imply conjecture 2.2 (see below). In fact, if $\kappa(K_X) = 0$ and $NK_X = A + B$ with A spanned and B pseudo-effective, then $\nu(A + B) = 0$, hence $\nu(A) = 0$ and therefore A = 0, A being spanned.

We start with an immediate observation:

Proposition 2.6 Conjecture 2.4 implies Conjecture 2.2, when X is not uniruled (and so when $\kappa(X) \ge 0$).

Proof. Let \mathcal{F} be a saturated subsheaf of Ω_X^p such that $\kappa(X, \det(\mathcal{F})) = \kappa^+(X) \ge 0$, then $Q = \Omega_X^p / \mathcal{F}$ is torsion free. By taking determinants we get

$$mK_X = \det \mathcal{F} + \det Q$$

for some positive integer m. We know that det Q is pseudo-effective, because X is not uniruled. By Conjecture 2.4, we get the claim, since $A := \det(\mathcal{F})$ is Q-effective. Q.E.D.

We now show that Conjecture 2.4 (in case $\kappa(X) \ge 0$), and so 2.2, is implied by the special case $\kappa(X) = 0$ of Conjecture 2.4. More precisely:

Proposition 2.7 Let X_n be a projective manifold with $\kappa(X) \ge 0$. Let $d = n - \kappa(X) \ge 0$. If Conjecture 2.4 holds for all manifolds G of dimension d and with $\kappa(G) = 0$, then Conjecture 2.4 (and so also Conjecture 2.2) holds for X.

Proof. By blowing up we may assume that the Iitaka fibration $g : X \to W$ is holomorphic. Let G be a general fiber of g. Thus $\kappa(G) = 0$. Let A be effective and B pseudo-effective on X such that

$$NK_X = A + B$$

for some positive integer N. Then A_G is effective, B_G is pseudo-effective and

$$NK_G = A_G + B_G.$$

Thus by Conjecture 2.4 applied to G, we conclude that $\kappa(G, A_{|G} \leq 0)$. By the easy additivity theorem for Kodaira dimension, we obtain that

$$\kappa(X, A) \le \dim(W) + \kappa(G, A_{|G}) \le \dim(W) = \kappa(X).$$

Q.E.D.

Q.E.D.

The preceding observation shows that the only two crucial cases of Conjecture 2.4 are when $\kappa(X)$ is either 0, or $-\infty$.

We now give some cases in which Conjecture 2.4 can be solved, so that 2.7 can be applied.

We first recall a notion from Mori theory. Let X be a projective manifold. A variety X' with at most terminal singularities is said to be a good minimal model for X, if X' is birational to X and some $mK_{X'}$ is (locally free and) spanned. Good minimal models are predicted to exist for every X with $\kappa(X) \ge 0$ but this known only in dimension up to 3.

Proposition 2.8 Let G be a projective manifold with $\kappa(G) = 0$. Suppose G has a good minimal model and that

$$NK_G = A + B$$

with A spanned and B pseudo-effective. Then A = 0.

Proof. Let G' be a good minimal model for G. Then $K_{G'} \equiv 0$ and actually $K_{G'}$ is torsion. Let A' and B' be the "strict transforms" of A and B in G'. Then A' is effective and B' is pseudo-effective with

$$NK_{G'} = A' + B'.$$

It follows A' = B' = 0 so that A = 0.

Since good minimal models exist in dimension up to 3, Prop. 2.7 gives in particular:

Theorem 2.9 Let X_n be a projective manifold, $\kappa(X) \ge 0$. Suppose $\kappa^+(X) \ge n-3$. Then $\kappa^+(X) = \kappa(X)$.

Therefore we can solve Conjecture 2.2 in dimension 4:

Corollary 2.10 Let X be a complex projective manifold of dimension 4 with $\kappa(X) \ge 0$.

1. $\kappa^+(X) = \kappa(X)$.

2. If $\kappa(X) = 0$, then Ueno's Conjecture holds for X, i.e.

$$h^0(X, \Omega^p_X) \le {4 \choose p},$$

for all $p \geq 0$.

Proof. For the first assertion, notice that we have either $\kappa^+(X) \ge 4-3=1$, or $\kappa^+(X) = 0$. Hence the conclusion follows from (2.9), since $\kappa^+(X) \ge \kappa(X) \ge 0$. For the second assertion, recall that Ueno's conjecture asserts that $h^0(X, \Omega_X^p) \le {n \choose p}$, for all $p \ge 0$, if $\kappa(X_n) = 0$. This is an immediate consequence of conjecture 2.2, as observed in [Ca95,5.10]. Q.E.D.

In the rest of this section we discuss Conjecture 2.4. When $B \in \text{Pic}^{0}(X)$ and A effective, this conjecture says that if K_X is numerically equivalent to an effective divisor, then $\kappa(X) \geq 0$. We will solve this case in section 4. For some other result towards (2.4) we state

Proposition 2.11 Let X_n be a projective manifold, $NK_X = A + B$ with A spanned and B pseudo-effective. Let $f : X \to Y$ be the fibration determined by |A|. Let F be the general fiber of f. If B|F is big, then K_X is big, i.e. $\kappa(X) = n$.

Proof. [CP04]

Corollary 2.12 Let X_n be a projective manifold, $NK_X = A + B$ with A spanned and B pseudo-effective. If $\kappa(A) = n - 1$, then $\kappa(X) \ge n - 1$.

Proof. Let $f : X \to Y$ be the fibration associated with A and let F denote the general fiber. Since dim F = 1, either B_F is ample or $B_F \equiv 0$.

In the first case we simply apply (2.11). In the second we notice $NK_F = B_F \equiv 0$ so that F is elliptic and $B_F = 0$. Then we can write

$$mB = f^*(L) + \sum d_i D_i$$

with L a line bundle on Y, with d_i integers, not necessarily positive, and with D_i irreducible divisors with dim $f(D_i) \leq n-2$, but not pull-backs of divisors on Y. Intersecting with movable curves in Y, it is easy to see that L is pseudo-effective. Writing $A = f^*(A')$, it follows that A' + L is big. Hence

$$\kappa(NmK_X + \sum (-d'_i D_i)) = n - 1,$$

where d'_i are just the negative d_i . Then however

$$\kappa(X) = n - 1$$

too.

Q.E.D.

Q.E.D.

3 Numerical properties of the Kodaira dimension

We solve here Conjecture 2.4 in the special case where B is numerically trivial.

Theorem 3.1 Let X be a projective complex manifold, and $L \in \text{Pic}^{0}(X)$ be numerically trivial. Then:

- 1. $\kappa(X, mK_X + L) \leq \kappa(X)$.
- 2. If $\kappa(X) = 0$, and if $\kappa(X, mK_X + L) = \kappa(X)$, then L is a torsion element in the group $\operatorname{Pic}^0(X)$.

Remark 3.2 The conclusion of (2) above does no longer hold when $\kappa(X) \ge 1$, as shown by curves (or even arbitrary manifolds) of general type. Another point not shown by our arguments is the behaviour of the modified plurigenera $p_m^+(X) :=$ $\sup\{h^0(X, mK_X + L), L \equiv 0\}$, as *m* is large and divisible. One may expect that then $p_m^+(X) = p_m(X)$, and that the maximum is attained at a torsion point, for every m > 0 (this is true for m = 1, by the arguments below).

Proof. We proceed in two steps:

A. We prove the result in the special case where $\kappa(X) \leq 0$. This will be done below in the two next propositions.

B. We now reduce the general case where $\kappa(X) > 0$ to the special case $\kappa(X) = 0$, as in 2.7 above.

Observe first that the statements involved are preserved by birational transformations of X. We can thus assume that both f, g are holomorphic, where $g: X \to W$ is the Iitaka-Moishezon fibration of X defined by some $|mK_X|$, and $f: X \to Y$ is the Iitaka fibration defined by some $|m(K_X + L)|$. If G is a general fibre of G, then it is sufficient to show that f(G) is a single point of Y. But then $f_{|G|}$ is nothing, but the Iitaka fibration on G defined by $(K_X + L)_{|G|}$. Because $\kappa(G) = 0$, we obtain the conclusion from the first step.

Remark 3.3 We see moreover that, in order to have equality $\kappa(K_X + L) = \kappa(X)$, it is necessary that $L_{|G}$ be torsion. This is Step B for the claim (2).

To conclude the proof of the preceding theorem, we still need to solve the case $\kappa(X) \leq 0$. This is the content of the next two propositions (in which additive and multiplicative notations for line bundles are mixed). We first deal with the case m = 1.

Proposition 3.4 Let X be a projective manifold, $L \in \text{Pic}^{0}(X)$. Suppose that $h^{0}(K_{X} \otimes L) \geq r$ for r = 1 or r = 2. Then

1. There exists a finite étale abelian cover $\tilde{X} \to X$ such that $h^0(2K_{\tilde{X}}) \ge r$. (In particular, $\kappa(X) \ge r - 1 \ge 0$)

2. If $\kappa(X) = 0$, and if $h^0(K_X \otimes L) = 1$, L is a torsion element in the group $\operatorname{Pic}^0(X)$.

Proof. Fix r = 1 or r = 2. Our aim is to prove that there is a finite étale cover $\tilde{X} \to X$ such that

$$h^0(K_{\tilde{X}} \otimes L^*) \ge r,$$

Then the canonical morphism

$$H^0(K_{\tilde{X}} \otimes L) \otimes H^0(K_{\tilde{X}} \otimes L^*) \to H^0(2K_{\tilde{X}})$$

will give the first claim $h^0(2K_{\tilde{X}}) \ge r$. Let

$$S = \{ G \in \operatorname{Pic}^{0}(X) \mid h^{0}(K_{X} \otimes G) \ge r \}.$$

If $S = \text{Pic}^{0}(X)$, then we are clearly done. Otherwise Simpson [Si93] gives the structure of S:

$$S = \bigcup \{A_i + T_i\}$$

with A_i torsion elements and T_i subtori of $\operatorname{Pic}^0(X)$. Let

$$S^* = \{ G^* \mid G \in S \}.$$

Then $S^* = \bigcup \{-A_i + T_i\}$. Hence we can write $L^* = -A_i + \tilde{L}$ with some $\tilde{L} \in T_i$. Consequently $\tilde{L} = \hat{L} - A_j$ with $\hat{L} \in S$, so that in total

$$L^* = -A_i - A_j + \hat{L}.$$

Now choose a finite étale cover $f : \tilde{X} \to X$ such that $f^*(A_i + A_j) = \mathcal{O}$. Then we conclude

$$h^0(K_{\tilde{X}} \otimes f^*(L^*)) \ge r$$

and we are done for assertion 1.

(Notice that if L is unitary flat, then by Hodge theory it is obvious that $h^0(K_X \otimes L) = h^0(K_X \otimes L^*)$, without using [Si93]).

Let us now prove statement (2). Replacing X by \tilde{X} as above, we can write: $L^* = \tilde{L}$, with $\tilde{L} \in T_i \subset S$. If T_i is not trivial, we get a one-parameter subgroup $L_t, t \in \mathbb{C}$ contained in $T_i \subset S$. The canonical morphisms

$$H^0(K_{\tilde{X}} \otimes L_t) \otimes H^0(K_{\tilde{X}} \otimes L_t^*) \to H^0(2K_{\tilde{X}})$$

show that $h^0(2K_{\tilde{X}}) \geq 2$, contradicting our assumption that $\kappa(X) = 0$.

Q.E.D.

We shall now reduce the general case of $m \ge 2$ to the special case m = 1, by means of cyclic covers.

Theorem 3.5 Let X be a projective manifold and L a line bundle with $c_1(L) = 0$ in $H^2(X, \mathbb{Z})$.

- 1. Suppose that there is a positive integer m such that $h^0(mK_X \otimes L) \geq 2$. Then $\kappa(X) \geq 1$.
- 2. Suppose that there is a positive integer m such that $h^0(mK_X \otimes L) \neq 0$. Then $\kappa(X) \geq 0$.
- 3. Suppose that $\kappa(X) = 0$, and that $h^0(mK_X \otimes L) \neq 0$. Then L is torsion in $\operatorname{Pic}^0(X)$.

Proof. We only prove (1), the proof of (2) being identical, simply omitting the divisor D in the arguments below. Since our claim is invariant by finite étale covers, we can pass to such covers as we like. In particular, we may assume that $L \in \text{Pic}^{0}(X)$. If m = 1, then our claim is Proposition 4.1, hence we shall assume $m \geq 2$. Furthermore we may assume that L = mL', so that

$$h^0(m(K_X \otimes L')) \ge 2.$$

Let $\sum b_i B_i$ be the fixed part of $|m(K_X \otimes L')|$, so that we can write

$$m(K_X \otimes L') = \sum b_i B_i + D$$

with D reduced and movable. By possibly blowing up we may assume that the support of $\sum b_i B_i + D$ has normal crossings. Now take the m-th root, normalize and desingularize to obtain $f : Y \to X$. We have to compute $f_*(K_Y)$, following [Es82,Vi83]. In fact, introduce the line bundles

$$H_j = j(K_X \otimes L') - \sum [jb_i m^{-1}]B_i.$$

Here $[\mathbf{x}]$ denotes the integral part of x. Then

$$f_*(K_Y) = K_X \otimes \bigoplus_{j=0}^{m-1} H_j.$$

Hence the direct summand of $f_*(K_Y) \otimes L'$ corresponding to j = m - 1 is just

$$D + \sum_{i} (b_i - [b_i(m-1)m^{-1}])B_i.$$

Since D moves, we obtain

$$h^0(f_*(K_Y) \otimes L') \ge 2,$$

hence

$$h^0(K_Y \otimes f^*(L')) \ge 2$$

so that $\kappa(Y) \ge 1$. We still need to prove $\kappa(X) \ge 1$. In to order to do that we must look more carefully at $f: Y \to X$. This map decomposes as follows. First we take the cyclic covering $h_0: Y_0 \to X$ determined by $m(K_X \otimes L') = \mathcal{O}_X(D)$. Then we have the normalisation $h_1: Y_1 \to Y_0$, and finally $h_2: Y \to Y_1$ is a desingularisation. Then Y_0 is Gorenstein and

$$K_{Y_0} = h_0^*(mK_X \otimes (m-1)L');$$

furthermore

$$K_{Y_1} \subset h^*(K_{Y_0})$$

via the trace map $(h_1)_*(K_{Y_1}) \to \mathcal{I} \otimes K_{Y_0}$ (with \mathcal{I} the conductor ideal) and finally

$$(h_2)_*(K_Y) = K_{Y_1} \tag{(+)}$$

since Y_1 has rational singularities [Es82,Vi83]. In total

$$K_Y \subset f^*(mK_X \otimes (m-1)L')) + \sum a_i E_i$$

where E_i are the exceptional components for h_2 and a_i are integers and we have

$$(h_2)_*(\mathcal{O}_Y(\sum a_i E_i)) = \mathcal{O}_{Y_1}$$

by (+). Thus

$$K_Y \otimes f^*((1-m)L') \subset f^*(mK_X) + \sum a_i E_i$$

We claim that there is a finite étale cover $g: \tilde{Y} \to Y$ such that

$$h^{0}(K_{\tilde{Y}} \otimes g^{*}f^{*}((1-m)L')) = h^{0}(g^{*}(K_{Y}) \otimes g^{*}f^{*}((1-m)L')) \ge 2.$$
(*)

This will prove $\kappa(g^*(f^*(K_X) + \sum a_i E_i)) \ge 1$, hence $\kappa(X) \ge 1$.

Now (*) is an easy application of Simpson's theorem, this time on Y. In fact, we introduce S_Y , the analogue of S on X. Then $S_Y = \bigcup_j \{B_j + T'_j\}$ with B_j torsion and T'_j tori. We already know that $f^*(L) \in S_Y$ so that

$$f^*(L) = B + M$$

with $M \in T'_j$, and $B = B_j$ for some j. Then $M' = (1 - m)f^*(L) \in T'_j$ and we find a finite étale cover $g: \tilde{Y} \to Y$ such that $g^*((1 - m)f^*(L)) = g^*(M')$, i.e. we choose g such that $g^*(B) = \mathcal{O}$. Now

$$h^0(K_Y + B + M') \ge 2$$

since $M' \in T'_j$. Hence $h^0(K_{\tilde{Y}} + g^*(M')) \ge 2$ which gives (*).

As already said, the proof of (2) is essentially the same, omitting the movable part. The proof of assertion (3) is the same as that of assertion (2) of the preceeding Proposition 3.4.

Q.E.D.

Remark 3.6 The preceeding result makes plausible the expectation that the generalised Green-Lazarsfeld sets

$$S_{m,p,r} = \{ L \in \operatorname{Pic}^{0}(X) \mid h^{p}(mK_{X} \otimes L) \ge r \}$$

might have the same structure as in [Si93] (finite union of translates of subtori by torsion elements).

Corollary 3.7 Let X be a projective manifold, A effective and B pseudo-effective divisors on X. Assume that $mK_X = A + B$ for some positive integer m. Suppose also that $\nu(B) = 0$, in the sense of [Bo02]. Then $\kappa(X) \ge 0$.

Proof. By [Bo02], we can write $B \equiv \sum b_i B_i$ with positive rational numbers b_i . Now apply (3.1).

Q.E.D.

4 The Universal Cover

Another invariant of X is defined via the universal cover \tilde{X} of a compact Kähler or projective manifold X. By identifying points in \tilde{X} which can be joined by a compact connected analytic set, one obtains an almost holomorphic meromorphic map $\tilde{X} \to \Gamma(\tilde{X})$. Here "almost holomorphic" is to say that the degeneracy locus does not project onto the image. If \tilde{X} is holomorphically convex (which is expected to be always true by the so-called Shafarevitch conjecture), then this map is holomorphic and is just the usual Remmert holomorphic reduction. In any case it induces the so-called Shafarevich map $\gamma_X : X \to \Gamma(X) = \Gamma(\tilde{X})/\pi_1(\tilde{X})$.

Definition 4.1 $\gamma d(X) = \dim \Gamma(X)$ is the Γ -dimension of X.

Notice that $\gamma d(X) = 0$ iff $\pi_1(X)$ is finite and that $\gamma d(X) = \dim X$ iff through the general point of \tilde{X} there is no positive dimensional compact subvariety, i.e. \tilde{X} geometrically seems as a modification of a Stein space.

The following result [Ca95,(4.1)] gives a relation between $\kappa^+(X)$ and $\gamma d(X)$.

Theorem 4.2 Let X be a compact Kähler manifold. If $\chi(X, \mathcal{O}_X) \neq 0$, then either

- 1. $\kappa^+(X) \ge \gamma d(X)$, or
- 2. $\kappa^+(X) = -\infty$, and so X is simply connected.

By (2.9) we obtain

Corollary 4.3 Let X_n be a projective manifold, $\kappa(X) \ge 0$. Suppose $\gamma d(X) \ge n-3$ and $\chi(\mathcal{O}_X) \ne 0$. Then $\kappa(X) \ge \gamma d(X) \ge n-3$. In particular, if n = 4, $\kappa(X) \ge 0$, $\pi_1(X)$ is infinite and $\chi(\mathcal{O}_X) \ne 0$, then $\kappa(X) \ge 1$. In other words, if X is a projective 4-fold with $\kappa(X) = 0$ and $\pi_1(X)$ is not finite, then $\chi(\mathcal{O}_X) = 0$; so there is either a holomorphic 1-form, or a holomorphic 3-form. Hence as in [Ca95, 5.9], we conclude:

Corollary 4.4 Let X be a projective manifold of dimension 4 such that $\kappa(X) = 0$, and $\chi(\mathcal{O}_X) \neq 0$. Then $\pi_1(X)$ is finite and has at most 8 elements.

This result should hold in arbitrary dimension n, with 8 replaced by 2^{n-1} , as a consequence of the standard conjecture that $\pi_1(X)$ should be almost abelian if $\kappa(X) = 0$. From Theorem 2.3 we deduce

Corollary 4.5 Let X be a projective manifold and suppose that its universal cover is not covered by its positive-dimensional compact subvarieties. Then either X is of general type, or $\chi(\mathcal{O}_X) = 0$.

Proof. By our assumption and [Ca95, (4.1)], we have $\kappa^+(X) = \dim X$ or $\chi(\mathcal{O}_X) = 0$. Now theorem (2.3) gives the claim. Q.E.D.

Corollary 4.6 Let X_n be a projective manifold whose universal cover is Stein (or has no positive-dimensional subvariety). Then either K_X is ample or $\chi(\mathcal{O}_X) = 0$, K_X is nef and $K_X^n = 0$.

Proof. This is immediate from (4.4) by observing that X does not have any rational curve, so that K_X must be nef by Mori theory. Moreover if K_X is big, then K_X is ample by Kawamata [Ka92].

Q.E.D.

We are lead to ask for the structure of projective manifolds X_n whose universal cover is Stein and with $K_X^n = 0$.

Conjecture 4.7 Let X_n be a projective manifold whose universal cover \tilde{X} is Stein. Assume $K_X^n = 0$. Then up to finite étale cover of X, the manifold X has a torus submersion over a projective manifold Y with K_Y ample and universal cover again Stein.

If the universal cover is only assumed not to admit a positive-dimensional subvariety through the general point, then one expects a birational version of 4.7, which is actually proved in [Ko93,5.8]. Here is the "Stein version" of this result which does not follow immediately from Kollár's result since we make a biholomorphic statement. The main point is to explain that we must have a holomorphic Iitaka fibration which is "almost smooth" and then apply Kollár's techniques to make it smooth.

Proposition 4.8 Conjecture 4.7 holds if $\kappa(X) \ge n-3$.

Proof. (1) Since the case $\kappa(X) = n - 1$ is the simplest, we do it first. Here the numerical dimension $\nu(X) = \kappa(X)$, so that K_X is good, i.e. some multiple is spanned [Ka85b]. Therefore we have a holomorphic litaka fibration $f: X \to Y$. The general fiber is an elliptic curve. Since X does not contain rational curves, it follows easily that all fibers are elliptic, sometimes multiple. Now [Ko93,sect.6] yields a finite étale cover such that the induced map is smooth; see below for some details.

(2) In the other case we consider the normalized graph $p : \mathcal{C} \to X$ of the family determined by the general fibers of the meromorphic Iitaka fibration. Let $q : \mathcal{C} \to T$ denote the parameter space. All irreducible fibers of q have dimension 2 (resp. 3) and every such fiber is an étale quotient of a torus by Lemma 4.9 below. Now we have a formula (via the trace map)

$$K_{\mathcal{C}} = p^*(K_X) + E$$

with an effective (Weil) divisor E. Restricting to a general (normal, hence smooth by (4.8) below) fiber F of q, we get

$$0 \equiv p^*(K_X)|F + E|F.$$

Hence $p^*(K_X)|F \equiv 0 = E|F$. Now consider the reduction F_0 of a component of a singular fiber (or rather its normalization) and use the conservation law (and the nefness of $p^*(K_X)$) to deduce $p^*(K_X)|F_0 \equiv 0$. Thus $p^*(K_X)$ is "q-numerically trivial". This proves immediately $\nu(X) = n - 2$ (resp. $\nu(X) = n - 3$) and again mK_X is spanned for a suitable m.

Now let again F_0 be the reduction of a component of a singular fiber F, this time of the Iitaka fibration $f: X \to Y$.

We claim that actually $F = aF_0$ and that f is equidimensional.

If dim T = 2, this is easy and well-known of course (take a general curve through $f(F_0)$ and observe that singular non-multiple fibers produce rational curves).

So suppose dim T = 1. Take μ maximal such that $\mu F_0 \subset F$. Then $N_{F_0}^{*\mu}$ has a section which has a zero, since F is reducible. Hence $K_{F_0} = -D$ with $D \neq \mathbb{Q}$ -effective divisor by the adjunction formula. Now normalize and then desingularize. The result \hat{F}_0 has $\kappa(\hat{F}_0) = -\infty$ (use formula (*) below), so that F_0 is uniruled. Since this is forbidden by the universal cover, we obtain $F = aF_0$.

Then $K_{F_0} \equiv 0$, so that its normalization F_0 has

$$K_{\tilde{F}_0} = \nu^*(K_{F_0}) - \tilde{N} \tag{(*)}$$

with \tilde{N} the preimage of the non-normal locus. Since $K_{\tilde{F}_0} \equiv 0$ by (2.9), we conclude that F_0 must have been normal, hence smooth.

Now we apply [Ko93,5.8] to obtain a finite étale cover X' of X which is birational to a torus submersion. But since X' does not contain rational curves, we obtain a holomorphic birational map from a torus submersion to X'. Since multiple fibers cannot be resolved by birational transformations on the base, we conclude that X' is a torus submersion itself. Q.E.D.

It remains to prove the following lemma of independent interest.

Lemma 4.9 Let X be an irreducible reduced variety of dimension at most 3. Assume that the universal cover of X is Stein (or does not contain compact subvarieties). Let $\tilde{X} \to X$ be the normalization and $\pi : \hat{X} \to \tilde{X}$ be a desingularization. Suppose $\kappa(\hat{X}) = 0$. Then \tilde{X} is an étale quotient of a torus.

Proof. We only treat the case dim X = 3, the surface case being easier and left to the reader. By [NS95], \hat{X} admits a finite étale cover $h : X' \to \hat{X}$ which is birational to a product of a simply connected manifold and an abelian variety. By our assumption on the universal cover, the simply connected part does not appear. It follows that the Albanese map $\alpha : X' \to A$ is birational. Now all irreducible components of all non-trivial fibers α are filled up by rational curves (α factors via Mori contractions). Since \tilde{X} does not contain rational curves, the map $X' \to \hat{X} \to \tilde{X}$ therefore factors over α , i.e. we obtain a finite map $g : A \to \tilde{X}$.

This map is étale in codimension 1. In fact otherwise by the ramification formula $K_A = g^*(K_{\tilde{X}}) + R$ (as Weil divisors). Thus $-K_{\tilde{X}}$ is non-zero effective and therefore $\kappa(\hat{X}) = -\infty$, contradiction.

We want to see that \tilde{X} is actually smooth and an étale quotient of A. First notice that \tilde{X} is \mathbb{Q} -Gorenstein (if g has degree d, then $dK_{\tilde{X}} = \mathcal{O}$ on the regular part of \tilde{X} , hence everywhere). Now we can compare the formulas

$$K_{X'} = h^* \pi^*(K_{\tilde{X}}) + \sum a_i E_i$$

and

$$K_{X'} = \sum b_j F_j$$

where E_i are the preimages of the π -exceptional components and F_j are the α -exceptional components; notice $b_j > 0$. Then both sets of exceptional divisors are equal, and thus all $a_i > 0$. Therefore \tilde{X} has only terminal singularities. We also notice that $\pi_1(\tilde{X})$ is almost abelian, i.e. abelian up to finite index. Therefore $\pi_1(\tilde{X})$ is abelian after finite étale cover. Then [Ka85] applies and \tilde{X} is an étale quotient of an abelian threefold. Here of course we use again that the universal cover of \tilde{X} is Stein. Q.E.D.

References

- [BDPP04] Boucksom,S.;Demailly,J.P.;Paun,M.;Peternell,Th.: The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. Preprint 2004
- [Bo02] Boucksom,S.: Cônes positifs des variétés complexes compactes. Thesis, Grenoble 2002
- [Ca95] Campana, F.: Fundamental group and positivity properties of cotangent bundles of compact Kähler manifolds. J. Alg. Geom. 4, 487-502 (1995)

- [CP04] Campana, F.; Peternell, Th.: Strictly nef divisors. Preprint in preparation (2004)
- [CQ04] Campana, F.; Zhang, Q.: Compact Kähler threefolds of π_1 -general type. math. AG/0402241
- [Es82] Esnault,H.: Fibre de Milnor d'un cône sur une courbe plane singuliére. Inv. math. 68, 477-496 (1982)
- [Ka85] Kawamata,Y.: Minimal models and the Kodaira dimension of algebraic fiber spaces. J. f. r. u. a. Math. 363, 1-46 (1985)
- [Ka85b] Kawamata, Y.: Pluricanonical systems on minimal algebraic varieties. Inv. math. 79, 567-588 (1985)
- [Ka92] Kawamata, Y.: Moderate degenerations of algebraic surfaces. Lecture Notes in Math. 1507, 113-133. Springer 1992
- [Ko93] Kollár, J.: Shafarevitch maps and plurigenera of algebraic varieties. Inv.math. 113, 177-215 (1993)
- [Mi87] Miyaoka,Y.: Deformations of a morphism along a foliation. Proc. Symp. Pure Math. vol. 46, 245-268 (1987)
- [NS95] Namikawa,Y.; Steenbrinck,J.H.M.: Global smoothings of Calabi-Yau 3folds. Inv. math. 122, 403-419 (1995)
- [SB92] Shepherd-Barron, N.: Miyaoka's theorem on the seminegativity of T_X . In: Flips an abundance for algebraic threefolds; ed. J. Kollár; Astérisque 211, 103-114 (1992)
- [Si93] Simpson, C.: Subspaces of Moduli Spaces of Rank One Local Systems. Ann.Sc. ENS. 26, 361-401 (1993).
- [Vi83] Viehweg,E.: Weak positivity and the additivity of the Kodaira dimension for certain fiber spaces. Ad. Stud. Pure Math. 1, 329-353 (1983)

Frédéric Campana Département de Mathématiques Université de Nancy F-54506 Vandoeuvre-les-Nancy, France frederic.campana@iecn.u-nancy.fr Thomas Peternell Mathematisches Institut Universität Bayreuth D-95440 Bayreuth, Germany thomas.peternell@uni-bayreuth.de