Gröbner bases and syzygies on bimodules

Manuel García Román and Socorro García Román

Departamento de Matemática Fundamental, Universidad de La Laguna, E-38271 La Laguna, Spain

Abstract

A new more efficient method for the computation of two-sided Gröbner bases of ideals and bimodules shifting the problem to the enveloping algebra is proposed. Arising from the ideas this method involves, we introduce the notion of two-sided syzygy, which reveals to be useful in the computation of the intersection of bimodules. Further applications are left for a sequel.

1 Introduction

Though first developed in the ring of polynomials, the methods based on Gröbner bases also work in some noncommutative rings, e.g. the Weyl algebras or, more generally, the so-called Poincaré-Birkhoff-Witt rings (PBW, for short), including some classical quantum groups. After first results in the Weyl algebra (Galligo, 1982) and in tensor algebras of finite-dimensional Lie algebras (Apel and Lassner, 1985), Kandri-Rodi and Weispfenning were the first to introduce Gröbner bases in the more general class of algebras where the degree of a skew-commutator $p_{ij} = x_j x_i - c_{ij} x_i x_j$ is bounded by the degree of the product of generators $x_i x_j$, for $1 \leq i < j \leq n$ (see Kandri-Rody and Weispfenning, 1990). The theory has recently been surveyed in Bueso et.al. (1998). Algorithms to compute the Gelfand-Kirillov dimension, to check whether a two-sided ideal is prime or not and to compute the projective dimension of a module have also been developed (see Bueso et.al., 1996; Lobillo, 1998; Bueso et.al., 1999; Gago-Vargas, 2003).

On these generalizations, authors were mainly interested in one-sided ideals and modules, whereas methods for the two-sided counterparts are merely patches in order to cope with the two-sided input data (cf. Pesch, 1998; Bueso et.al., 2003)

¹ Research supported by PI2002/075, Gobierno de Canarias.

In this note we show that those mends are not necessary, due to the very well known fact that two-sided ideals and bimodules may be seen as left modules on the enveloping algebra. First, we show that the enveloping algebra of a PBW algebra is another PBW algebra. Second, we find a method to shift the data back and forth through the morphism

$$\mathfrak{m}^s: (R^{\mathrm{env}})^s \longrightarrow R^s; \quad (f_i \otimes g_i)_{i=1}^s \longmapsto (f_i g_i)_{i=1}^s$$

in order to carry out the computations on the enveloping algebra using onesided techniques.

This philosophy allows, for example, to compute Gröbner bases for bimodules with only one call to the left Buchberger algorithm, instead of the a priori unknown number of calls typical of the aforementioned methods.

The techniques we use have led us to study the syzygy bimodule, which is the two-sided counterpart of the left syzygy module. Amongst its applications, we show that it can be used in the computation of intersections of bimodules when one starts, as usual, from two-sided input data.

Throughout this paper we will use the following notation. We will denote by ϵ_i the element $(0, ..., \overset{i}{1}, ..., 0) \in \mathbb{N}^n$. The symbol x^{α} will denote the monomial $x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ in the free algebra $k\langle x_1, \ldots, x_n \rangle$ or in any of its epimorphic images, k being a field. R will be a PBW k-algebra, R^{op} its opposite algebra and R^{env} its enveloping algebra $R \otimes_k R^{\text{op}}$. Finally, for any subset F of the free left R-module R^s , we will denote by $_R\langle F \rangle$, resp. $_R\langle F \rangle_R$, the left R-module, resp. the R-bimodule, generated by F.

The computations of the examples shown in this paper were done using a library of procedures built by the authors using the package of symbolic computation Maple 6. The computation times correspond to a Pentium III 700 Mhz personal computer with 256 Mb RAM.

2 The enveloping algebra of a PBW algebra

A subset $Q = \{x_j x_i - q_{ji} x_i x_j - p_{ji}; 1 \leq i < j \leq n\}$ of the free algebra $k \langle x_1, \ldots, x_n \rangle$ is a set of quantum relations bounded by the admissible order " \preceq " on \mathbb{N}^n if $q_{ji} \in k^*$ and p_{ji} is a finite k-linear combination of standard monomials x^{α} ($\alpha \in \mathbb{N}^n$) such that $\exp(p_{ji}) \prec \epsilon_i + \epsilon_j$, for all i < j, where $\exp(f)$ denotes the exponent of the leading term of the element f. As may be found in the literature, (see e.g. Kandri-Rody and Weispfenning, 1990; Bueso et.al., 1998, 2003), a *Poincaré-Birkhoff-Witt algebra* (PBW algebra, for short) is a k-algebra R where the set of standard monomials $\{x^{\alpha}; \alpha \in \mathbb{N}^n\}$ is a k-basis and

such that there exists a set of quantum relations Q bounded by an admissible order " \preceq " satisfying

$$R = \frac{k \langle x_1, \dots, x_n \rangle}{k \langle x_1, \dots, x_n \rangle \langle Q \rangle_{k \langle x_1, \dots, x_n \rangle}}$$

This algebra is usually denoted by $k\{x_1, \ldots, x_n; Q, \preceq\}$.

Amongst the examples of PBW algebras, we find the commutative polynomial ring $k[x_1, ..., x_n]$, some iterated Ore extensions or a pretty large class of quantum groups just as the multiparameter *n*-dimensional quantum space $\mathcal{O}_q(\mathbb{A}^n)$, the bialgebra of quantum matrices $M_q(2)$, the Weyl algebra $A_n(k)$, the enveloping algebra of traceless matrices $U(\mathfrak{sl}(2))$, etc.

The tensor product of PBW algebras is a new PBW algebra:

Proposition 1 If $R = k\{x_1, \ldots, x_m; Q_R, \preceq_R\}$ and $S = k\{y_1, \ldots, y_n; Q_S, \preceq_S\}$ are PBW algebras with quantum relations

$$Q_R = \{ x_j x_i - q_{ji} x_i x_j - p_{ji}; \ 1 \le i < j \le m \},\$$
$$Q_S = \{ y_j y_i - q'_{ji} y_i y_j - p'_{ji}; \ 1 \le i < j \le n \},\$$

then $R \otimes_k S$ is the PBW algebra denoted by

$$k\{x_1 \otimes 1, \dots, x_m \otimes 1, 1 \otimes y_1, \dots, 1 \otimes y_n; Q, \preceq\}, \text{ where}$$

$$Q = \begin{cases} (x_j \otimes 1)(x_i \otimes 1) - q_{ji}(x_i \otimes 1)(x_j \otimes 1) - p_{ji} \otimes 1; & 1 \le i < j \le m \\ (1 \otimes y_j)(x_i \otimes 1) - (x_i \otimes 1)(1 \otimes y_j); & 1 \le i \le m, \ 1 \le j \le n \\ (1 \otimes y_j)(1 \otimes y_i) - q'_{ji}(1 \otimes y_i)(1 \otimes y_j) - 1 \otimes p'_{ji}; & 1 \le i < j \le n. \end{cases}$$

and " \leq " is one amongst the elimination orders (as defined, e.g., in Cox et.al., 1992) arising from " \leq_R " and " \leq_S ".

As a first example, note that $A_{n+m}(k)$ is the PBW algebra $A_n(k) \otimes A_m(k)$ constructed in the proposition. Another example of this construction is the enveloping algebra R^{env} of $R = k\{x_1, \ldots, x_n; Q, \leq\}$. Before we describe it, let us define the composition orders. For any $\alpha = (\alpha_1, \ldots, \alpha_1) \in \mathbb{N}^n$, denote by α^{op} the *n*-tuple $(\alpha_n, \ldots, \alpha_1)$.

Definition 2 Let " \leq " be an order on \mathbb{N}^n . The up-component composition

order in \mathbb{N}^{2n} , denoted " \preceq^{c} ", is defined by

$$(\alpha,\beta) \prec^{c} (\gamma,\delta) \Leftrightarrow \begin{cases} \alpha + \beta^{\mathrm{op}} \prec \gamma + \delta^{\mathrm{op}}, \text{ or} \\ \alpha + \beta^{\mathrm{op}} = \gamma + \delta^{\mathrm{op}} \text{ and } \beta^{\mathrm{op}} \prec \delta^{\mathrm{op}} \end{cases}$$

The down-compontent composition order " \preceq_c " is defined by

$$(\alpha,\beta) \prec_c (\gamma,\delta) \Leftrightarrow \begin{cases} \alpha + \beta^{\mathrm{op}} \prec \gamma + \delta^{\mathrm{op}}, \text{ or} \\ \alpha + \beta^{\mathrm{op}} = \gamma + \delta^{\mathrm{op}} \text{ and } \alpha \prec \gamma \end{cases}$$

If " \preceq " is an admissible order on \mathbb{N}^n , then both composition orders " \preceq^c " and " \preceq_c " are admissible orders on \mathbb{N}^{2n} .

Note that the opposite algebra R^{op} is the PBW algebra $k\{x_n, \ldots, x_1; Q^{\text{op}}, \preceq^{op}\}$, where the elements of Q^{op} are those of Q written oppositely and " \preceq^{op} " is the order in \mathbb{N}^n given by $\alpha \preceq^{\text{op}} \beta \iff \alpha^{\text{op}} \preceq \beta^{\text{op}}$.

Proposition 3 If $R = k\{x_1, \ldots, x_n; Q, \preceq\}$ is a PBW algebra with quantum relations $Q = \{x_jx_i - q_{ji}x_ix_j - p_{ji}; 1 \leq i < j \leq n\}$, then R^{env} is the PBW algebra $k\{x_1 \otimes 1, \ldots, x_n \otimes 1, 1 \otimes x_n, \ldots, 1 \otimes x_1; Q^*, \preccurlyeq\}$, where

$$Q^* = \begin{cases} (x_j \otimes 1)(x_i \otimes 1) - q_{ji}(x_i \otimes 1)(x_j \otimes 1) - p_{ji} \otimes 1; & 1 \le i < j \le n \\ (1 \otimes x_j)(x_i \otimes 1) - (x_i \otimes 1)(1 \otimes x_j); & 1 \le i, j \le n \\ (1 \otimes x_i)(1 \otimes x_j) - q_{ji}(1 \otimes x_j)(1 \otimes x_i) - 1 \otimes p_{ji}; & 1 \le i < j \le n. \end{cases}$$

and " \preccurlyeq " is, either any of the elimination orders " \preceq *" or " \preceq *" in \mathbb{N}^{2n} corresponding to " \preceq " and " \preceq °", or any of the composition orders " \preceq " or " \preceq c" on \mathbb{N}^{2n} corresponding to " \preceq ".

In what follows, we will work on the free *R*-bimodule R^s , being *s* a positive integer and *R* a PBW algebra, and we will use the *R*-module basis $\{e_i\}_{i=1}^s$ consisting of $e_i = (0, \dots \overset{i}{1}, \dots, 0) \in R^s$ for all $1 \leq i \leq s$.

The notion and some applications of left Gröbner bases in PBW algebras and left modules may be found, e.g., in Bueso et.al. (2003). For convenience, just recall that if $M \subset \mathbb{R}^s$ is an *R*-bimodule, then $G = \{g_1, ..., g_r\} \subset M \setminus \{0\}$ is a two-sided Gröbner basis if one of the following equivalent statements holds

(1) $M = {}_{R}\langle G \rangle_{R}$ and $\operatorname{Exp}(M) = \bigcup_{k=1}^{r} (\mathbb{N}^{n} + \operatorname{exp}(\boldsymbol{g}_{k}));$

(2) G is a left Gröbner basis and $M = {}_{R}\langle G \rangle_{R} = {}_{R}\langle G \rangle;$

(3) G is a left Gröbner basis, $M = {}_{R}\langle G \rangle_{R}$ and $g_{k}x_{i} \in {}_{R}\langle G \rangle$, for all $k \in \{1, ..., r\}$ and $i \in \{1, ..., n\}$.

A set $G \subset R^s$ is said to be a *two-sided Gröbner basis* if so it is for the *R*-bimodule $_R\langle G \rangle_R$.

3 Computing two-sided Gröbner Bases

This section is devoted to the methods for computation of two-sided Gröbner bases for *R*-subbimodules of R^s . We denote by $\boldsymbol{f} \otimes \boldsymbol{g}$ the element $(f_1 \otimes g_1, ..., f_s \otimes g_s) \in (R^{\text{env}})^s$, where $\boldsymbol{f} = (f_1, ..., f_s), \boldsymbol{g} = (g_1, ..., g_s) \in R^s$.

As a consequence of the third caracterization of two-sided Gröbner bases above, some authors have proposed an algorithm to compute them (see Bueso et.al., 2003). Alternatively, we propose a new algorithm which improves that one, since it calls only once the left Buchberger Algorithm, although it uses more variables and input elements. The philosophy is to transform the problem into computing a left Gröbner basis in the free module $(R^{env})^s$. This may be done since, just as we saw in the previous section, R^{env} has a PBW structure.

It is known that *R*-bimodules are exactly left R^{env} -modules. Notice that, in particular, the free module R^s is a left R^{env} -module with the action $(r \otimes r') \mathbf{f} = (rf_1r', ..., rf_sr')$, and $(R^{\text{env}})^s$ possesses an *R*-bimodule structure whose multiplications are given by $r(\mathbf{f} \otimes \mathbf{g})r' = (rf_1 \otimes g_1r', ..., rf_s \otimes g_sr')$, where $r, r' \in R$ and $\mathbf{f} = (f_1, ..., f_s), \ \mathbf{g} = (g_1, ..., g_s) \in R^s$.

Likewise, the map $\mathfrak{m}^s = \mathfrak{m} \times \stackrel{s}{\cdots} \times \mathfrak{m} : (R^{\text{env}})^s \to R^s$, where $\mathfrak{m}(r \otimes r') = rr'$, for $r, r' \in R$, is an epimorphism of left R^{env} -modules. Thus there exists a bijection

$$\{N \subseteq (R^{\text{env}})^s; \operatorname{Ker}(\mathfrak{m}^s) \subseteq N \in R^{\text{env}} - \operatorname{Mod}\} \longrightarrow \{M \subseteq R^s; M \in R - \operatorname{Bimod}\}$$
$$N \to M_N := \mathfrak{m}^s(N),$$
$$N_M := (\mathfrak{m}^s)^{-1}(M) \leftarrow M$$

Using this bijection, for each *R*-bimodule $M \subset R^s$, we have a left R^{env} -module $N_M \subset (R^{\text{env}})^s$. Moreover, from a finite generator system for *M* it is possible to obtain one for N_M , just as the following results show.

Lemma 4 Let R be a k-algebra.

(1) If $M = {}_{R}\langle \boldsymbol{f_1}, ..., \boldsymbol{f_t} \rangle_R \in R^s$, then $N_M = {}_{R^{\mathrm{env}}}\langle \boldsymbol{f_1} \otimes \boldsymbol{1}, ..., \boldsymbol{f_t} \otimes \boldsymbol{1} \rangle + \mathrm{Ker}(\mathfrak{m}^s);$ (2) $\mathrm{Ker}(\mathfrak{m}^s) = {}_{R^{\mathrm{env}}}\langle \boldsymbol{f} \otimes \boldsymbol{1} - \boldsymbol{1} \otimes \boldsymbol{f}; \boldsymbol{f} \in R^s \rangle;$

(3) If
$$R = k\{x_1, ..., x_n; Q; \leq\}$$
 is a PBW algebra, then $_{R^{env}}\langle \boldsymbol{f} \otimes \boldsymbol{1} - \boldsymbol{1} \otimes \boldsymbol{f}; \boldsymbol{f} \in R^s \rangle = _{R^{env}}\langle \boldsymbol{x}^{(\boldsymbol{\epsilon}_j, \boldsymbol{k})} \otimes \boldsymbol{1} - \boldsymbol{1} \otimes \boldsymbol{x}^{(\boldsymbol{\epsilon}_j, \boldsymbol{k})}; 1 \leq j \leq n, 1 \leq k \leq s \rangle.$

Corollary 5 If $R = k\{x_1, ..., x_n; Q; \leq\}$ is a PBW algebra and $M = {}_R\langle f_1, ..., f_t \rangle_R \subseteq R^s$ then

$$N_M = {}_{R^{\mathrm{env}}} \langle \ \{ \boldsymbol{f_i} \otimes \boldsymbol{1} \}_{i=1}^t, \{ \boldsymbol{x^{(\epsilon_j, k)}} \otimes \boldsymbol{1} - \boldsymbol{1} \otimes \boldsymbol{x^{(\epsilon_j, k)}} \}_{1 \leq j \leq n, \ 1 \leq k \leq s} \ \rangle.$$

Given an admissible order in \mathbb{N}^n , from here on we will call TOP (term over position), resp. POT (position over term) the orders in $\mathbb{N}^{n,(s)}$ given by

$$(\alpha, i) \prec (\beta, j) \iff \begin{cases} \alpha \prec \beta, \text{ or} \\ \alpha = \beta \text{ and } i > j \end{cases} \text{ resp. } \begin{cases} i > j, \text{ or} \\ i = j \text{ and } \alpha \prec \beta \end{cases}$$

Lemma 6 Let $R = k\{x_1, ..., x_n; Q; \leq\}$ be a PBW algebra and consider the order TOP (or POT) on both R^s and $(R^{env})^s$.

- Taking \preceq^* or \preceq^c on R^{env} , if $\mathbf{h} \in (R^{\text{env}})^s$ is such that $\exp_{(R^{\text{env}})^s}(\mathbf{h}) = ((\alpha, 0), i) \in \mathbb{N}^{2n, (s)}$, then $\mathbf{h} \notin \operatorname{Ker}(\mathfrak{m}^s)$ and $\exp_{R^s}(\mathfrak{m}^s(\mathbf{h})) = (\alpha, i);$
- Taking \preceq_* or \preceq_c on R^{env} , if $\mathbf{h} \in (R^{\text{env}})^s$ is such that $\exp_{(R^{\text{env}})^s}(\mathbf{h}) = ((0, \alpha), i) \in \mathbb{N}^{2n,(s)}$, then $\mathbf{h} \notin \operatorname{Ker}(\mathfrak{m}^s)$ and $\exp_{R^s}(\mathfrak{m}^s(\mathbf{h})) = (\alpha^{\operatorname{op}}, i)$.

Using these results we have:

Theorem 7 Let $R = k\{x_1, ..., x_n; Q, \preceq\}$ be a PBW algebra, $M \subset R^s$ be an R-bimodule and consider in R^{env} the PBW structure given in the proposition 3 (where the order is one of \preceq^* , \preceq^c , \preceq_* or \preceq_c).

If G is a left Gröbner basis for $N_M = (\mathfrak{m}^s)^{-1}(M)$ with TOP (resp. POT), then the set $\mathfrak{m}^s(G) \setminus \{0\}$ is a two-sided Gröbner basis for M with TOP (resp. POT).

The theorem provides a method of construction of two-sided Gröbner bases for bimodules. For convenience we write explicitly under the name of Algorithm 1.

Algorithm 1 Two-sided Gröbner bases

Require: $F = \{f_1, ..., f_t\} \subseteq R^s \setminus \{0\};$ **Ensure:** $G = \{g_1, ..., g_{t'}\}, a \text{ two-sided Gröbner basis for } {}_R\langle F \rangle_R \text{ such that } F \subseteq G;$

INITIALIZATION: $B := \{f_i \otimes 1\}_{i=1}^t \cup \{x^{(\epsilon_j, k)} \otimes 1 - 1 \otimes x^{(\epsilon_j, k)}\}_{1 \le j \le n, 1 \le k \le s};$ Using the Left Buchberger Algorithm, compute a left Gröbner basis G' in the PBW algebra $(R^{env})^s$ for the input data B;

If $G' = \{ \boldsymbol{g'_1}, ..., \boldsymbol{g'_r} \}$ with $\boldsymbol{g'_i} = (\sum_{j \in \mathfrak{I}_i} p_{ij}^1 \otimes q_{ij}^1, ..., \sum_{j \in \mathfrak{I}_i} p_{ij}^s \otimes q_{ij}^s)$, take

$$g_{i} := (\sum_{j \in \mathfrak{I}_{i}} p_{ij}^{1} q_{ij}^{1}, ..., \sum_{j \in \mathfrak{I}_{i}} p_{ij}^{s} q_{ij}^{s});$$

$$G := \emptyset;$$
for all $i = 1$ to r do
$$if g_{i} \neq 0 \text{ then}$$

$$G := G \cup \{g_{i}\};$$
end if
end for

The advantage offered by this algorithm is that only one call to the left Buchberger Algorithm is done, whereas the one shown in (Bueso et.al., 2003) makes an, a priori, unknown number of calls.

Example 8 Let R be the quantum plane, i.e., $R = \mathbb{C}\{x, y; \{yx - qxy\}, \leq_{(1,3)}\}$ (take $q = i\sqrt{2}$) where $\leq_{(1,3)}$ is the (1,3)-weighted lexicographical order. Let $F = \{(2x, x^2y, xy^2 + y^2), (xy, 0, -x^2y^2), (x^2, 2, 0)\} \subset \mathbb{R}^3$ and consider the order TOP in \mathbb{R}^3 .

The old algorithm (see Bueso et.al., 2003) takes 66.371 seconds to compute the two-sided Gröbner basis, consisting of 17 elements:

$$\begin{split} &G = \{ (2x, x^2y, xy^2 + y^2), \, (xy, 0, -x^2y^2), \, (x^2, 2, 0), \, (xy - 2x, x^3y - x^2y - 4, -y^2), \\ &(xy, x^4y + y - 4x, 0), \, (0, (i\sqrt{2} + 2)x^3y - 12, 0), \, (0, (i\sqrt{2} + 2)y, 0), \\ &((-i\sqrt{2} - 2)xy, (-i\sqrt{2} - 2)x^3y + (-4 + 4i\sqrt{2})x + 12, 0), \, (0, (-4 + 4i\sqrt{2})x^2, 0), \\ &((i\sqrt{2} - 1)xy^2 + (2 - 2i\sqrt{2})xy, (2 - 2i\sqrt{2})x^3y^2 + 3x^2y^2 + (-4 - 2i\sqrt{2})y, 0), \\ &((1 - i\sqrt{2})xy^2 + (-2 + 2i\sqrt{2})xy, (1 + 2i\sqrt{2})x^3y^2 - 3x^2y^2, 0), \\ &((1 + i\sqrt{2}/2)xy^2, 0, 0), \, (0, 3y, 0), \, ((1 - i\sqrt{2}/4)xy^2, 3/4y^2 + (-4 + i\sqrt{2})xy, 0), \\ &(xy, -2i\sqrt{2}x, 0), \, (0, (-4 + 2i\sqrt{2})x, 0), \, (0, -4 + 2i\sqrt{2}, 0) \}. \end{split}$$

The algorithm 1 takes 46.972 seconds to compute a two-sided Gröbner basis with 12 elements (considering the order \leq^{c} (see remark 2) in \mathbb{R}^{env}):

$$\begin{split} &G = \{ (2x, x^2y, xy^2 + y^2), \ (xy, 0, -x^2y^2), \ (x^2, 2), \ (xy - 2x, x^3y - x^2y - 4, -y^2), \\ &(0, (i\sqrt{2} + 2)x^3y - 12, 0), \ ((i\sqrt{2} - 1)xy^2 + (2 - 2i\sqrt{2})xy, 3x^2y^2 + (2 - 2i\sqrt{2})y, \\ &0), \ (2xy, -i\sqrt{2}y - 4i\sqrt{2}x, 0), \ (0, (-i\sqrt{(2)}/4 - 1/2)y^2 + (2 - 2i\sqrt{2})xy, 0), \\ &(0, 3y, 0), \ (0, -2i\sqrt{2}x^2, 0), \ (0, -6i\sqrt{2}x, 0), \ (0, -16i\sqrt{2} - 16, 0) \}. \end{split}$$

Example 9 Let R be the PBW algebra $M_q(2) = \mathbb{C}\{x, y, z, t; Q, \preceq_{glex}\}$ of quantum matrices (with $q = i\sqrt{2}$), where $Q = \{yx - qxy, ty - qyt, zx - qxz, tz - qzt, zy - yz, tx - xt - (q^{-1} - q)yz\}$ and consider the order "POT" in \mathbb{R}^2 . Let $F = \{(-xzt + 1, 2y^3), (x^2t, y^2)\} \subset \mathbb{R}^2$.

The old algorithm takes 216.433 seconds to compute the two-sided Gröbner basis, consisting of 26 elements:

$$\begin{split} G &= \{ \left(-xzt + 1, 2y^3 \right), \left(x^2t, y^2 \right), \left(-x, -2xy^3 + 1/2y^2z \right), \left(-3/2yz^2 - 1, 4i\sqrt{2}xy^3zt - 3y^4z^2 - y^2z^2t - 2y^3 \right), \left(0, 4i\sqrt{2}x^2y^3t - i\sqrt{2}xy^2zt + 3/8i\sqrt{2}y^3z^2 + y^2 \right), \\ (0, 3i\sqrt{2}xy^4z^2 + 3/8y^3z^3 - (4i\sqrt{2} + 2)xy^3 + (1/2 - 1/2i\sqrt{2})y^2z), \\ (0, -3i\sqrt{2}x^2y^4z - 3/8i\sqrt{2}xy^3z^2 - 3xy^2), \left(0, (4i\sqrt{2} + 2)x^2y^3 - (i\sqrt{2} + 1/2)xy^2z), \left(0, (12 - 6i\sqrt{2}xy^4z^2 - 3/4(i\sqrt{2} + 1)y^3z^3 + (2 + 4i\sqrt{2})xy^3 + 3/2y^2z), \left(0, -12x^2y^4z - 3/2xy^3z^2 + (-2 + 2i\sqrt{2})xy^2 \right), \\ (0, (1 + 1/2i\sqrt{2})y^3), \left(0, (1/2 + 1/4i\sqrt{2})y^3z), \left(0, (1 - i\sqrt{2})y^3z^2t \right), \left(0, (i\sqrt{2} + 2)xy^3zt + (3/8i\sqrt{2} - 3/8)y^4z^2 + (1 + 1/2i\sqrt{2})y^3 \right), \left(0, (1 + 1/2i\sqrt{2})y^2z \right), \\ (0, (1/2 + 1/4i\sqrt{2})y^2z^2), \left(0, (1 - i\sqrt{2})y^2z^3t \right), \left(0, (i\sqrt{2} + 2)xy^2z^2t + (3/8i\sqrt{2} - 3/8)y^3z^3 + (1 + 1/2i\sqrt{2})y^2z), \left((1 + 1/4i\sqrt{2})t, -8xy^3zt^2 + 3i\sqrt{2}y^4z^2t - i\sqrt{2}y^2z^2t^2 + 4y^3t \right), \left(3/2y, 3y^4 + 3y^2t \right), \left(3/2i\sqrt{2}yz, (-4i\sqrt{2} - 2)xy^3t + 3i\sqrt{2}y^4z + (1/2 + i\sqrt{2})y^2zt), \left((-1 + 1/4i\sqrt{2})t, (4i\sqrt{2} + 8)xy^3zt^2 + (-3 - 3i\sqrt{2})y^4z^2t + (i\sqrt{2} - 1)y^2z^2t^2), \left(0, (-3i\sqrt{2}xy^4zt + 3/8i\sqrt{2}y^3z^2t + (1 + 1/2i\sqrt{2})y^2t), \left(-1, 0 \right), \left(0, -i\sqrt{2}y^2t), \left(0, (7/9 - 4/9i\sqrt{2})y^2 \right) \}. \end{split}$$

The algorithm 1 takes 59.909 seconds to compute the two-sided Gröbner basis (considering the order \preceq^* in \mathbb{R}^{env}), consisting of 11 elements:

$$\begin{split} G &= \{ (-xzt+1,2y^3), (x^2t,y^2), (-x,-2xy^3+1/2y^2z), (0,-3i\sqrt{2}xy^4z^2-3/8y^3z^3 \\ &+ (4i\sqrt{2}+2)xy^3 + (1/2i\sqrt{2}-1/2)y^2z), (-3/2yz^2t+(-1/2i\sqrt{2}-1)t,-6y^3t), \\ &(0,-3i\sqrt{2}x^2y^4z-3/8i\sqrt{2}xy^3z^2-3xy^2), (0,(1+1/2i\sqrt{2})y^3), \\ &(0,(1+1/2i\sqrt{2})y^2z), (3/2y,3y^2t), (-1,0), (0,y^2) \}. \end{split}$$

4 Syzygy Bimodules

In this section we study the notion of syzygy bimodule of a subset of R^s , R being a PBW algebra and $s \in \mathbb{N}^*$. This notion can be viewed as the analogous one of *left syzygy module* for left modules, since it presents some similar properties.

There exists an algorithm (see again Bueso et.al., 2003) which computes a generator system of the left syzygy module $Syz^l(F)$, provided a finite set of input data $F \subset R^s$ is given. This algorithm is shown below and will be used within Algorithm 3.

Algorithm 2 Left Syzygy Module

Require: $F = \{f_1, ..., f_t\} \subseteq R^s \setminus \{0\};$ **Ensure:** H, a finite left generator system of $Syz^l(F)$;

$\label{eq:INITIALIZATION: Run the Left Buchberger Algorithm for the input data$

F in order to compute: $- a \text{ left Gröbner basis } G = \{g_1, ..., g_r\} \subset R^s \text{ for }_R \langle F \rangle,$ $- \text{ the elements } h_{ij}^k \in R \text{ such that } SP(g_i, g_j) = \sum_{k=1}^r h_{ij}^k g_k \text{ for all } 1 \leq i < j \leq r, \text{ and}$ $- \text{ the matrix } Q \in M_{r \times t}(R) \text{ such that } (g_1, ..., g_r) = (f_1, ..., f_t)Q^t;$ $for \text{ all } 1 \leq i < j \leq r \text{ do}$ $if \text{ level } (\exp(g_i)) = \text{ level } (\exp(g_j)) \text{ then}$ $Compute \ r_{ij}, r_{ji} \text{ such that } SP(g_i, g_j) = r_{ij}g_i - r_{ji}g_j;$ $\text{Let } p_{ij} := (0, ..., r_{ij}^i, ..., 0) - (0, ..., r_{ji}^j, ..., 0) - (h_{ij}^1, ..., h_{ij}^r);$ $end \ if$

Let $H := \{ p_{ij}Q \text{ such that } 1 \leq i < j \leq r, \text{ and level } (\exp(\mathbf{g}_i)) = \text{level } (\exp(\mathbf{g}_j)) \}.$

Definition 10 Let $f_1, ..., f_t \in \mathbb{R}^s$. The syzygy bimodule of the matrix

$$F = \begin{bmatrix} \boldsymbol{f_1} \\ \vdots \\ \boldsymbol{f_t} \end{bmatrix} \in M_{t \times s}(R),$$

denoted by Syz(F) or $Syz(f_1, ..., f_t)$, is the kernel of the homomorphism of left R^{env} -modules $(R^{\text{env}})^t \longrightarrow R^s$; $(h_1, ..., h_t) \longmapsto \sum_{i=1}^t h_i f_i$.

We can compute the syzygy bimodule of a matrix F using again the techniques showed in section 3, that is, we will *move* the problem to the context of the enveloping algebra in order to use the methods on the left side.

Proposition 11 Let $M \subseteq R^s$ be a *R*-bimodule and $N = (\mathfrak{m}^s)^{-1}(M)$. Let $\{h_1, ..., h_r\} \subset (R^{\text{env}})^{t+sn}$ be a generator system of $Syz^l(\{f_i \otimes 1\}_{i=1}^t, \{x^{(\epsilon_j, k)} \otimes 1 - 1 \otimes x^{(\epsilon_j, k)}\}_{1 \leq j \leq n, 1 \leq k \leq s})$ as a left R^{env} -module.

Then $Syz(f_1, ..., f_t) = {}_R\langle \pi(h_1), ..., \pi(h_r) \rangle_R$, where π is the projection homomorphism $\pi : (R^{env})^t \times (R^{env})^{sn} \longrightarrow (R^{env})^t$.

PROOF. Notice that $\sum_{i=1}^{t} g_i(f_i \otimes 1) \in \operatorname{Ker}(\mathfrak{m}^s)$ for any $g = (g_1, ..., g_t) \in$

 $\begin{array}{l} Syz(\boldsymbol{f_1},...,\ \boldsymbol{f_t}). \text{ So, there exists an element } \boldsymbol{g'} = (g'_{11},...,g'_{1s},...,g'_{n1},...,g'_{sn}) \\ \in (R^{\text{env}})^{sn} \text{ such that } \sum_i g_i(\boldsymbol{f_i} \otimes \mathbf{1}) = \sum_{j,k} g'_{jk}(\boldsymbol{x^{(\epsilon_j,k)}} \otimes \mathbf{1} - \mathbf{1} \otimes \boldsymbol{x^{(\epsilon_j,k)}}). \text{ Hence,} \\ (\boldsymbol{g}, -\boldsymbol{g'}) \text{ is in } Syz^l(\{\boldsymbol{f_i} \otimes \mathbf{1}\}_{i=1}^t, \{\boldsymbol{x^{(\epsilon_j,k)}} \otimes \mathbf{1} - \mathbf{1} \otimes \boldsymbol{x^{(\epsilon_j,k)}}\}_{1 \leq j \leq n, \ 1 \leq k \leq s}). \text{ From this point, the proof may easily be finished.} \quad \Box \end{array}$

Algorithm 3 Syzygy Bimodule

Require: $F = \{f_1, ..., f_t\} \subseteq R^s \setminus \{0\};$ **Ensure:** H, a finite generator system of Syz(F) as an R-bimodule;

INITIALIZATION: $B := \{f_i \otimes \mathbf{1}\}_{i=1}^t \cup \{x^{(\epsilon_j, \mathbf{k})} \otimes \mathbf{1} - \mathbf{1} \otimes x^{(\epsilon_j, \mathbf{k})}\}_{1 \leq j \leq n, 1 \leq k \leq s};$ Using the Left Syzygy Module Algorithm, compute in the PBW algebra $(R^{\text{env}})^s$ a generator system $H = \{h_1, ..., h_r\}$ of $Syz^l(B)$ as a left R^{env} -module;

If $\mathbf{h}_{i} = (\mathbf{h}_{i}', \mathbf{h}_{i}'')$ where $\mathbf{h}_{i}' \in (\mathbb{R}^{\text{env}})^{t}$ and $\mathbf{h}_{i}'' \in (\mathbb{R}^{\text{env}})^{sn}$ for $1 \leq i \leq r$, take $H := \{\mathbf{h}_{1}', ..., \mathbf{h}_{r}'\}.$

Example 12 Let R be the quantum plane with the PBW algebra structure $\mathbb{C}\{x, y; \{yx - qxy\}, \leq_{(2,1)}\}$, (where $q = i\sqrt{2}$), and consider the order POT in \mathbb{R}^2 . Let $F = \{(x + 1, y), (xy, 0)\} \subset \mathbb{R}^2$.

Algorithm 3 takes 23.622 seconds to compute the R-bimodule generator system H of Syz(F) consisting of 8 elements:

$$\begin{split} H &= \{ (1 \otimes y - y \otimes 1, (-1 + i\sqrt{2})1 \otimes 1), \\ ((\frac{i\sqrt{2}}{6} - \frac{1}{3})y \otimes x + (-\frac{2}{3} - \frac{i\sqrt{2}}{3})xy \otimes 1, 1 \otimes x + 1 \otimes 1), (0, 1 \otimes y + \frac{i\sqrt{2}}{2}y \otimes 1), \\ ((\frac{i\sqrt{2}}{6} - \frac{1}{3})y \otimes x + (-\frac{2}{3} + \frac{i\sqrt{2}}{3})xy \otimes 1, i\sqrt{2}x \otimes 1 + 1 \otimes 1), \\ ((-\frac{1}{3} - \frac{i\sqrt{2}}{3})y \otimes x + (-\frac{2i\sqrt{2}}{3} - \frac{2}{3})xy \otimes 1, i\sqrt{2} \otimes x + i\sqrt{2} \otimes 1), \\ (-y \otimes y + y^2 \otimes 1 + i\sqrt{2} \otimes y, -i\sqrt{2}y \otimes 1), \\ ((\frac{1}{3} + \frac{i\sqrt{2}}{3})1 \otimes x^2 + (\frac{1}{3} - \frac{2i\sqrt{2}}{3})x \otimes x + (-\frac{2}{3} + \frac{i\sqrt{2}}{3})x^2 \otimes 1, 0), \\ ((\frac{1}{3} + \frac{i\sqrt{2}}{3})1 \otimes xy + (\frac{i\sqrt{2}}{6} - \frac{1}{3})y \otimes x + (-\frac{1}{3} - \frac{i\sqrt{2}}{3})x \otimes y + (\frac{1}{3} + \frac{i\sqrt{2}}{3})xy \otimes 1, 0) \}. \end{split}$$

Although elimination techniques are useful at solving several problems in Module Theory, they are computationally inefficient, mainly because elimination orders are unavoidably used.

On the other hand, it has been proved, first in the commutative case and then using left syzygy R-modules being R a non-commutative ring (such as a PBW algebra), that syzygies provide a much more efficient treatment, for example, in the computation of the intersection of left R-submodules of R^s , ideal quotients, kernels of homomorphisms of left R-submodules, etc. (see Bueso et.al., 2003).

In what follows, we will see that some applications of left syzygies can be generalized using the new definition of syzygy bimodules, so that, for example, it is possible to give an algorithm to compute a finite intersection of Rsubbimodules of R^s when, as natural, two-sided input data are given. Further applications will be studied in the sequels.

The following result states a general property which the above-mentioned algorithm will be based on.

Lemma 13 Let M be an R-subbimodule of R^s such that there exist $p, q \ge 1$ and $H = \begin{bmatrix} H_1 \\ H_2 \end{bmatrix} \in M_{(s+p)\times q}(R)$, where $H_1 \in M_{s\times q}(R)$ and $H_2 \in M_{p\times q}(R)$, satisfying the following two conditions:

i)
$$(\mathfrak{m}^{s}(\boldsymbol{h})\otimes \mathbf{1})H_{1} = \boldsymbol{h}H_{1}, \ \forall \boldsymbol{h} \in (R^{\mathrm{env}})^{s};$$

ii)
$$M = \{ \boldsymbol{h} \in R^s ; \exists \boldsymbol{h}'' \in (R^{env})^p \text{ such that } (\boldsymbol{h} \otimes \boldsymbol{1}, \boldsymbol{h}'') \in Syz(H) \}.$$

Let us split up each element $\mathbf{h} \in (\mathbb{R}^{\text{env}})^{s+p}$ into $\mathbf{h} = (\mathbf{h}', \mathbf{h}'')$ with $\mathbf{h}' \in (\mathbb{R}^{\text{env}})^s$ and $\mathbf{h}'' \in (\mathbb{R}^{\text{env}})^p$.

- (1) If $\{\mathbf{h_1}, ..., \mathbf{h_t}\} \subseteq (R^{env})^{s+p}$ is an R-bimodule generator system of Syz(H)then $M = {}_R \langle \mathfrak{m}^s(\mathbf{h_1}'), ..., \mathfrak{m}^s(\mathbf{h_t}') \rangle_R$.
- (2) Furthermore, if {h₁, ..., h_t} is a left Gröbner basis of Syz(H) (as a left R^{env}-module) for the order POT in (R^{env})^{s+p} and any of ≤*, ≤^c, ≤_{*}, ≤_c in R^{env}, then {m^s(h₁'), ..., m^s(h_t')} \ {0} is a two-sided Gröbner basis of M for POT.

Theorem 14 Let $\{M_i\}_{i=1}^r$ be a family of *R*-subbimodules of R^s and suppose that $M_i = {}_R \langle f_1^i, ..., f_{t_i}^i \rangle_R \subseteq R^s$. Then

$$\bigcap_{i=1}^{r} M_{i} = \{ \boldsymbol{h} \in R^{s} ; \exists \boldsymbol{h}'' \in (R^{\text{env}})^{\sum_{j=1}^{r} t_{j}} \text{ such that } (\boldsymbol{h} \otimes 1, \boldsymbol{h}'') \in Syz(H) \},\$$

where

$$H = \begin{pmatrix} I_s & \cdots & I_s \\ f_1^1 & \cdots & \mathbf{0} \\ \vdots & & \vdots \\ f_{t_1}^1 & \cdots & \mathbf{0} \\ \vdots & & \vdots \\ \mathbf{0} & \cdots & f_1^r \\ \vdots & & \vdots \\ \mathbf{0} & \cdots & f_{t_r}^r \end{pmatrix} \in M_{(s + \sum_{i=1}^r t_i) \times rs}(R)$$

From 13 and 14, an algorithm to compute finite intersections of R-subbimodules of R^s may be formulated (see Algorithm 4).

Corollary 15 Let M_i and H be as in 14. If $Syz(H) = {}_R\langle g_1, ..., g_t \rangle_R$ with $g_k = (g_k', g_k'') \in (R^{env})^s \times (R^{env})^{\sum_{j=1}^r t_j}$ for all $1 \le k \le t$, then

$$\bigcap_{i=1}^{r} M_{i} = {}_{R} \langle \mathfrak{m}^{s}(\boldsymbol{g_{1}}'), ..., \mathfrak{m}^{s}(\boldsymbol{g_{t}}') \rangle_{R}.$$

If $G = \{\mathbf{g_1}, ..., \mathbf{g_t}\}$ is also a left Gröbner basis of Syz(H) (as left R^{env} -module) with POT in $(R^{env})^{s+\sum_{j=1}^r t_j}$, then $\{\mathfrak{m}^s(\mathbf{g_1}'), ..., \mathfrak{m}^s(\mathbf{g_t}')\} \setminus \{0\}$ is a two-sided Gröbner basis of $\bigcap_{i=1}^r M_i$ with POT in R^s .

Algorithm 4 Intersection of R-subbimodules of R^s

Require: $\{M_i\}_{i=1}^r$, a family of *R*-subbimodules of R^s with $M_i = {}_R\langle f_1^i, ..., f_{t_i}^i \rangle_R$; **Ensure:** M, a finite generator system of $\bigcap_{i=1}^r M_i$ as an *R*-bimodule;

INITIALIZATION:

$$H := \begin{pmatrix} I_s & \cdots & I_s \\ f_1^1 & \cdots & \mathbf{0} \\ \vdots & & \vdots \\ f_{t_1}^1 & \cdots & \mathbf{0} \\ \vdots & & \vdots \\ \mathbf{0} & \cdots & f_1^r \\ \vdots & & \vdots \\ \mathbf{0} & \cdots & f_{t_r}^r \end{pmatrix} \in M_{(s + \sum_{j=1}^r t_j) \times rs}(R);$$

Using the Syzygy Bimodule Algorithm compute a generator system $G = \{g_1, ..., g_t\}$ of Syz(H) as an R-bimodule; If $g_k = (g_k', g_k'')$ where $g_k' \in (R^{env})^s$ and $g_k'' \in (R^{env})^{\sum_{j=1}^r t_j}$ for $1 \le k \le t$, take $M := \{\mathfrak{m}^s(g_1'), ..., \mathfrak{m}^s(g_t')\}.$

Example 16 Let R be the quantum plane (as in example 12) and consider the order POT in \mathbb{R}^2 . Let M_1 and M_2 be the R-subbimodules of \mathbb{R}^2 generated by $\{(2x^2 + 2x, -y), (0, -8), (-3xy, 0)\}$ and $\{(x + 2, 0), (1, -y)\}$, respectively.

Algorithm 4 takes 111.418 seconds to compute the R-bimodule generator system M of $M_1 \cap M_2$, consisting of 8 elements:

$$\begin{split} M &= \{ \left(\frac{4i\sqrt{2}}{3}x^2y + \frac{7}{3}xy, \frac{i\sqrt{2}}{3}y^2\right), \left(-\frac{2i\sqrt{2}}{3}x^3 + \left(-\frac{2i\sqrt{2}}{3} - \frac{4}{3}\right)x^2 - \frac{4}{3}x, \frac{i\sqrt{2}}{3}xy + \frac{2}{3}y\right), \\ &\left(-\frac{5}{3}x^2y + \frac{4i\sqrt{2}}{3}xy, -\frac{2}{3}y^2\right), \left(\left(-1 - \frac{5i\sqrt{2}}{3}\right)x^3 - \left(\frac{19}{3} + \frac{5i\sqrt{2}}{3}\right)x^2 - \frac{16}{3}x, \\ &\frac{4i\sqrt{2}}{3}xy + \frac{8}{3}y\right), \left(5x^2y + \left(-1 - 3i\sqrt{2}\right)xy, 2y^2\right), \left(\frac{2}{3}x^2 + \frac{2}{3}x, -\frac{4}{3}y\right), \\ &\left(\frac{5i\sqrt{2}}{3}x^2y + \frac{8}{3}xy, \frac{2i\sqrt{2}}{3}y^2\right), \left(\frac{8}{3}x^2y^2 - \frac{7}{3}xy^2, -\frac{1}{3}y^3\right), \left(-\frac{2}{3}x^3 - 2x^2 - \frac{4}{3}x, \frac{1}{3}xy + \frac{2}{3}y\right), \left(\left(\frac{4i\sqrt{2}}{3} + \frac{8}{3}\right)x^2y + \left(\frac{4}{3} - \frac{4i\sqrt{2}}{3}\right)xy, \left(\frac{i\sqrt{2}}{3} + \frac{2}{3}\right)y^2\right), \\ &\left(-\frac{4i\sqrt{2}}{3}x^3y - 4x^2y + \frac{4i\sqrt{2}}{3}xy, -\frac{i\sqrt{2}}{3}xy^2 - \frac{2}{3}y^2\right), \left(\left(-\frac{2}{3} + \frac{2i\sqrt{2}}{3}\right)x^3 + \left(-\frac{2}{3} + \frac{2i\sqrt{2}}{3}\right)x^2, 0\right), \left(-2x^2y + \left(-\frac{2}{3} + \frac{2i\sqrt{2}}{3}\right)xy, 0\right), \left(\frac{2}{3}x^2 + \frac{2}{3}x, -\frac{4}{3}y\right) \}. \end{split}$$

Acknowledgements

Authors are grateful to Prof. Francisco Castro for his motivation, and to Prof. Alain Verschoren for his corrections and suggestions.

References

- Apel, J. and Lassner, W., 1985. An algorithm for calculations in enveloping fields of Lie algebras, Proc. Conf. Comp. Alg. Dubna, 231–241.
- Bueso, J.L., Castro, F., Gómez Torrecillas, J. and Lobillo, F.J., 1996. Computing the Gelfand-Kirillov dimension, SAC Newsleter, 1, 39–52.
- Bueso, J.L., Castro, F., Gómez Torrecillas, J. and Lobillo, F.J., 1998. An introduction to effective calculus in quantum groups, Lect. Notes Pure App. Math. 197, 55-83.
- Bueso, J.L., Castro, F., Gómez Torrecillas, J. and Lobillo, F.J., 1999. Test de primalité dans une extension de Ore itérée, C. R. Acad. Sci. Paris Sr. I Math. 328, 459-462.
- Bueso, J.L., Gómez Torrecillas, J. and Verschoren, A., 2003. Algorithmic Methods in Noncommutative Algebra: Applications to Quantum Groups, Kluwer.
- Cox, D., Little, J. and O'Shea, D., 1992. *Ideals, Varieties, and Algorithms*, Undergraduate Texts in Mathematics, Springer.
- Gago-Vargas, J., 2003. Bases for projective modules in $A_n(k)$, J. Symb. Comput. **36**, 845–853.
- Galligo, A., 1982. Algorithmes de cacul de bases standard, preprint Niza.
- García Román, M.D. and García Román, M.S., 2001. Un nuevo algoritmo para el cálculo de bases de Gröbner biláteras (in spanish), Proc. EACA'2001 held in La Rioja-Spain (there is an accessible abstract on http://csimbolico.rediris.es/eaca01/20garcia.pdf).
- Kandri-Rody, A. and Weispfenning, V., 1990. Non-commutative Gröbner bases in algebras of solvable type, J. Symb. Comput. 9, 1–26.
- Lobillo, F.J., 1998. *Métodos algebraicos y efectivos en grupos cuánticos* (in spanish), PhD thesis, Universidad de Granada.
- Pesch, M., 1998. Two-sided Gröbner bases in iterated Ore extensions, Symbolic rewriting techniques. Papers from the workshop held in Ascona, Switzerland, April 30 - May 4, 1995. Prog. Comput. Sci. Appl. Log. 15, 225-243 (1998).