

A COMPUTER-BASED APPROACH TO THE CLASSIFICATION OF NILPOTENT LIE ALGEBRAS

CSABA SCHNEIDER

ABSTRACT. We adopt the p -group generation algorithm to classify small-dimensional nilpotent Lie algebras over small fields. Using an implementation of this algorithm, we list the nilpotent Lie algebras of dimension at most 9 over \mathbb{F}_2 and those of dimension at most 7 over \mathbb{F}_3 and \mathbb{F}_5 .

1. INTRODUCTION

The classification of n -dimensional nilpotent Lie algebras over a given field \mathbb{F} is very difficult problem even for a relatively small n . The aim of this article is to present a series of computer calculations that led to the following theorem.

Theorem 1.1. *The number of isomorphism types of 6-dimensional nilpotent Lie algebras is 36 over \mathbb{F}_2 , and 34 over \mathbb{F}_3 and \mathbb{F}_5 . The number of isomorphism types of 7-dimensional nilpotent Lie algebras is 202 over \mathbb{F}_2 , 199 over \mathbb{F}_3 , and 211 over \mathbb{F}_5 . The number of isomorphism types of nilpotent Lie algebras with dimension 8 and 9 over \mathbb{F}_2 is 1836 and 26711, respectively.*

The classifications in Theorem 1.1 were obtained via a series of computer calculations using a GAP 4 [The GAP Group 2004] implementation of a nilpotent Lie algebra generation algorithm. The ideas used in these calculations are the same as those used in the classification of finite 2-groups with order at most 2^9 ; see [O'Brien 1990, Eick & O'Brien 1999]. Let $\gamma_i(L)$ denote the i -th term of the lower central series of a Lie algebra L , so that $\gamma_1(L) = L$, $\gamma_2(L) = L'$, etc. If L is a finitely generated nilpotent Lie algebra with nilpotency class c then L is an immediate descendant of $L/\gamma_c(L)$ (see Section 2 for definitions). Further, $L/\gamma_c(L)$ is an immediate descendant of $L/\gamma_{c-1}(L)$. Continuing this way, we can see that every finitely generated nilpotent Lie algebra can be obtained after finitely many steps from a finite-dimensional abelian Lie algebra by computing immediate descendants. This suggests that a theoretical algorithm to generate all n -dimensional, nilpotent Lie algebras can be designed once we can efficiently compute immediate descendants. We will see that every immediate descendant of L is a quotient of another nilpotent Lie algebra, which is referred to as the *cover*. It is shown in this paper that, for a finite-dimensional, nilpotent \mathbb{F}_p -Lie algebra L , it is possible to effectively compute the cover,

Date: 18 June 2004.

2000 Mathematics Subject Classification. 17B05, 17B30, 17-08.

Key words and phrases. nilpotent Lie algebras, immediate descendants, covers, Lie algebra generation algorithm.

and then to compute a complete and irredundant list of the isomorphism types of the immediate descendants of L . Repeating the immediate descendant calculation finitely many times, it is, in theory, possible to obtain a complete and irredundant list of all isomorphism types of the nilpotent Lie algebras with a given dimension over a finite field. In practice, this calculation quickly becomes unfeasible as n grows. Nevertheless, using this approach, it is possible to obtain classifications of Lie algebras that would otherwise be beyond hope; see Theorem 1.1.

The structure of this paper is as follows. In Section 2 we develop the theory of a Lie algebra generation algorithm. An application of the algorithm to prove Theorem 1.1 will be presented in Section 3. The final Section 4 will discuss an implementation of the algorithm.

2. A NILPOTENT LIE ALGEBRA GENERATION ALGORITHM

Our nilpotent Lie algebra generation algorithm is an adaptation of O'Brien's p -group generation algorithm, whose details can be found in [O'Brien 1990]. The Lie algebra generation algorithm is described without proofs in this section. Another variation on this theme is presented in [O'Brien et al. 2004] where the authors classified groups and nilpotent Lie rings of order p^6 . Recently Michael Vaughan-Lee used the same approach to extend this result to p^7 .

Throughout this section L is a finite-dimensional, nilpotent Lie algebra. Let $Z(L)$ denote the center of L . A nilpotent Lie algebra K is said to be a *central extension* of L if K has an ideal I such that $I \leq K' \cap Z(K)$ and $K/I \cong L$. In the terminology of [Batten et al. 1996, Batten & Stitzinger 1996], (K, I) is said to be a defining pair for L . The algebra K is said to be an *immediate descendant* of L if $L \cong K/\gamma_c(K)$ where c is the nilpotency class of K . Hence an immediate descendant is a special kind of central extension. The *cover* L^* of a finitely generated nilpotent Lie algebra L is defined as follows. Suppose that $\dim L/L' = d$. Then L is a d -generator Lie algebra, and so the free Lie algebra F_d with rank d has an ideal I such that $F_d/I \cong L$. The cover L^* is defined as the Lie algebra $F_d/[I, F_d]$. The *multiplicator* of L^* is the ideal $I/[I, F_d]$. The cover L^* is also a finite-dimensional nilpotent Lie algebra. Moreover, if L has nilpotency class c then the class of L^* is at most $c + 1$, and $\gamma_{c+1}(L^*)$ is referred to as the *nucleus* of L^* .

Suppose now without loss of generality that $L = F_d/I$ as in the previous paragraph. Let L^* be the cover of L with multiplicator M and nucleus N . Then K is a central extension of L if and only if $K \cong L^*/J$ for some ideal $J \leq M$. Further, in this case, K is an immediate descendant of L if and only if $J \neq M$ and $J + N = M$. A proper subspace J of M with $J + N = M$ is said to be *allowable*. Thus it is possible to obtain a complete list of immediate descendants of L by listing all quotients L^*/J where J runs through the allowable subspaces of the multiplicator M . Unfortunately, two different allowable subspaces may lead to isomorphic Lie algebras. This problem can, however, be tackled using the automorphism group of L . If α is an automorphism, then α can be lifted to an automorphism α^* of the cover L^* as follows. Let $\psi : L^* \rightarrow L$ denote the natural epimorphism with kernel M . Suppose that b_1, \dots, b_d is a minimal generating set for L^* ; then $b_1\psi, \dots, b_d\psi$ is a minimal generating set for L . Suppose that $y_1, \dots, y_d \in L^*$ are chosen so that $b_i\psi\alpha = y_i\psi$ for all $i \in \{1, \dots, d\}$.

Then the map $b_i \mapsto y_i$, for $i = 1, \dots, d$, can uniquely be extended to an automorphism of L^* . This automorphism is denoted α^* , even though it is not uniquely determined by α . On the other hand the restriction of α^* to $M = I/[I, F_d]$ depends only on α . This defines a linear representation

$$(1) \quad \varrho : \text{Aut}(L) \rightarrow \text{GL}(M) \quad \text{given by} \quad \alpha \mapsto \alpha^*|_M.$$

Using a familiar argument, it is not hard to see that two allowable subspaces J_1 and J_2 give isomorphic Lie algebras L^*/J_1 and L^*/J_2 if and only if J_1 and J_2 are in the same orbit under the action $\text{Aut}(L)\varrho$.

If J is an allowable subspace of the multiplier then the automorphism group of $K = L^*/J$ can also be computed using $\text{Aut}(L)$. Let S denote the stabiliser in $\text{Aut}(L)$ of J under the representation ϱ . Let X denote a generating set for S . For each $\alpha \in X$ choose $\alpha^* \in \text{Aut}(L^*)$ as in the previous paragraph and let $X^* = \{\alpha^* \mid \alpha \in X\}$. Suppose that $\{b_1, \dots, b_d\}$ is a minimal generating set for K and that $\{c_1, \dots, c_l\}$ is a basis for the last non-trivial term of the lower central series of K . For $i \in \{1, \dots, d\}$ and $j \in \{1, \dots, l\}$ let $\psi_{i,j}$ denote the automorphism that maps b_i to $b_i + c_j$ and fixes $b_1, \dots, b_{i-1}, b_{i+1}, \dots, b_d$. Then $X^* \cup \{\psi_{i,j} \mid i = 1, \dots, d, j = 1, \dots, l\}$ is a generating set for $\text{Aut}(K)$.

The cover of a finite-dimensional nilpotent Lie algebra L can be constructed in a way that is very similar to the construction of the p -covering group of a finite p -group. The good description of this procedure can be found in [Newman et al. 1998]. Suppose that L has class c , and hence the lower central series is as follows:

$$L = \gamma_1(L) > \gamma_2(L) = L' > \gamma_3(L) > \dots > \gamma_c(L) > \gamma_{c+1}(L) = 0.$$

We say that a basis $\mathcal{B} = \{b_1, \dots, b_n\}$ for L is *compatible with the lower central series* if there are indices $1 = i_1 < i_2 < \dots < i_{c-1} < i_c \leq n$ such that $\{b_{i_k}, \dots, b_n\}$ is a basis of $\gamma_k(L)$ for $k \in \{1, \dots, c\}$.

Suppose that $b_i \in \gamma_j(L) \setminus \gamma_{j+1}(L)$. Then we say that the number j is the *weight* of b_i . We call a basis \mathcal{B} a *nilpotent basis* if the following hold.

- (i) The basis \mathcal{B} is compatible with the lower central series;
- (ii) for each $b_i \in \mathcal{B}$ with weight $w \geq 2$ there are $b_{j_1}, b_{j_2} \in \mathcal{B}$ with weight 1 and $w - 1$, respectively, such that $b_i = [b_{j_1}, b_{j_2}]$. The product $[b_{j_1}, b_{j_2}]$ is called the *definition* of b_i .

If $\{b_1, \dots, b_n\}$ is a nilpotent basis for a Lie algebra L , then there are coefficients $\alpha_{i,j}^k$ for $i < j < k$ such that

$$(2) \quad [b_i, b_j] = \sum_{k=j+1}^n \alpha_{i,j}^k b_k.$$

It is routine to see that every finitely generated nilpotent Lie algebra has a nilpotent basis.

Suppose that $\mathcal{B} = \{b_1, \dots, b_n\}$ is a nilpotent basis for a d -generator, nilpotent Lie algebra and the $\alpha_{i,j}^k$ are as in (2). We build a presentation for the Lie algebra L^* as follows. The set

$\{b_{d+1}, \dots, b_n\}$ is a basis for L' . If, for some $i < j$, the product $[b_i, b_j]$ is not a definition and $w(b_i) + w(b_j) \leq c + 1$, then we modify the product in (2) by introducing a central basis element $b_{i,j}$ and set

$$[b_i, b_j] = \sum_{k=j+1}^n \alpha_{i,j}^k b_k + b_{i,j}.$$

We introduce the new basis elements so that different non-defining products $[b_i, b_j]$ are augmented with different basis elements $b_{i,j}$. We also ensure that the newly introduced basis elements $b_{i,j}$ are central. If a product $[b_i, b_j]$ is a definition of b_k , say, then the product $[b_i, b_j] = b_k$ is not modified. Similarly if $w(b_i) + w(b_j) > c + 1$ then $[b_i, b_j]$ is left untouched. This way we obtain an anti-commutative algebra \hat{L} with basis $\{b_1, \dots, b_d\} \cup \{b_{i,j}\}$ where the product of two basis elements is defined using the rules above. We compute the ideal J in \hat{L} generated by the set of elements

$$\{[b_i, b_j, b_k] + [b_j, b_k, b_i] + [b_k, b_i, b_j] \mid i, j, k \in \{1, \dots, n\}\}.$$

Then we obtain the cover L^* as \hat{L}/J .

It is possible to make this basic algorithm to compute the cover more effective. In practice we only introduce a new basis element for products of the form $[b_i, b_j]$ where $w(b_i) = 1$ and compute products $[b_i, b_j]$ with $w(b_i) > 1$ using the Jacobi identity. We also use the result in [Havas et al. 1990] that J is already generated by the set of element

$$\{[b_i, b_j, b_k] + [b_j, b_k, b_i] + [b_k, b_i, b_j] \mid i \in \{1, \dots, d\}, i < j < k \leq n\}.$$

The proof that the resulting Lie algebra is isomorphic to L^* is completely analogous to that in the p -group case; see [Newman et al. 1998] for details.

3. SOME CLASSIFICATIONS OF SMALL LIE ALGEBRAS

In theory it is possible to use the procedures described in Section 2 to classify nilpotent \mathbb{F}_q -Lie algebras of a given dimension using a recursion. It is clear that there is a unique 1-dimensional nilpotent Lie algebra over each field \mathbb{F}_q ; the automorphism group of this algebra is naturally isomorphic to the multiplicative group \mathbb{F}_q^* . Suppose that we have a complete and irredundant list of nilpotent \mathbb{F}_q -Lie algebras of dimension $1, \dots, n-1$ for some $n \geq 2$ and we are also given the automorphism groups of these algebras. Up to isomorphism, there is exactly one n -dimensional abelian \mathbb{F}_q -Lie algebra. Each non-abelian nilpotent Lie algebra with dimension n is an immediate descendant of a smaller-dimensional Lie algebra. Hence, for each algebra L with dimension m in the precomputed list we construct the Lie cover L^* , the multiplier M , and $\text{Aut}(L)_\varrho$ where ϱ is the representation in (1). Then, using that M is finite, we construct the orbits of the $(m + \dim M - n)$ -dimensional allowable subspaces under the finite linear group $\text{Aut}(L)_\varrho$. For each orbit representative U we construct the quotient L^*/U and the stabiliser of U in $\text{Aut}(L)$ under the representation ϱ . The automorphism group of L^*/U can also be constructed as described in Section 2. The collection of all Lie algebras

L^*/U so obtained is a complete and irredundant list of the isomorphism types of the non-abelian nilpotent Lie algebras with dimension n .

Suppose that L is a finite-dimensional nilpotent Lie algebra. Let c denote the class of L . Then the *type* of the Lie algebra L is the symbol

$$[\dim L/L', \dim L'/\gamma_3(L), \dots, \dim \gamma_c(L)][\dim Z(L)].$$

It is well known that, over an arbitrary field, there is just one nilpotent Lie algebra with dimension 1 and 2. There are two nilpotent Lie algebras with dimension 3 (the types are $[3][3]$ and $[2, 1][1]$), and 3 nilpotent Lie algebras with dimension 4 (the types are $[4][4]$, $[3, 1][2]$, $[2, 1, 1][1]$). The number of isomorphism types of 5-dimensional nilpotent Lie algebras is 9 over all fields; see [Goze & Khakimjanov 1996]. Up to isomorphism, there is exactly one Lie algebra with each of the following types: $[5][5]$, $[4, 1][3]$, $[4, 1][1]$, $[3, 2][2]$, $[3, 1, 1][2]$, $[3, 1, 1][1]$, $[2, 1, 2][2]$; there are two Lie algebras with type $[2, 1, 1, 1][1]$.

The number of 6-dimensional nilpotent Lie algebras depends on the underlying field. Using the **GAP 4** package described in Section 4, I obtained 36 isomorphism classes of 6-dimensional nilpotent Lie algebras over \mathbb{F}_2 , and 34 such classes over \mathbb{F}_3 and \mathbb{F}_5 . It is mentioned in Wilkinson's paper [Wilkinson 1988] that the number of isomorphism classes of finite p -groups with order p^6 and exponent p is 34 whenever $p \geq 7$. Though there are several mistakes in the main part of Wilkinson's paper, this particular claim has independently been checked by several people and is widely considered to be true. Using the Lazard correspondence [O'Brien et al. 2004, Section 4] we obtain that, for $p \geq 7$, there are 34 pairwise non-isomorphic 6-dimensional nilpotent \mathbb{F}_p -Lie algebras. In fact, the computation referred to above implies that this claim holds already for $p = 3, 5$. The number of 6-dimensional, nilpotent \mathbb{F}_2 -Lie algebras for each possible type can be found in Table 1, while Table 2 contains the same information over \mathbb{F}_3 and \mathbb{F}_5 . One can read off, for instance, from these tables that there are 6 pairwise non-isomorphic nilpotent Lie algebras with type $[2, 1, 1, 1, 1][1]$ over \mathbb{F}_2 and there are only 5 such Lie algebras over \mathbb{F}_3 and \mathbb{F}_5 .

There exist classifications of 6-dimensional nilpotent Lie algebras over infinite fields; see for instance [Goze & Khakimjanov 1996].

A classification of finite p -groups with exponent 7 and order p^7 was obtained by Wilkinson [Wilkinson 1988]. If $p \geq 7$ then, by the Lazard correspondence, the number of finite p -groups with exponent p and order p^7 coincides with the number of 7-dimensional nilpotent \mathbb{F}_p -Lie algebras. According to Wilkinson this number is $173 + 7p + 2 \gcd(p - 1, 3)$, but Michael Vaughan-Lee pointed out in private communication that there are several mistakes in Wilkinson's calculations and the correct number is

$$(3) \quad 174 + 7p + 2 \gcd(p - 1, 3).$$

Computer calculations with the **GAP 4** package described in Section 4 shows that the number of 7-dimensional nilpotent Lie algebras over \mathbb{F}_2 , \mathbb{F}_3 , and \mathbb{F}_5 is 202, 199, 211, respectively; the number of Lie algebras for each possible type is presented in Tables 3–5 in Section 5. This

calculation also shows that (3) is valid over \mathbb{F}_5 . Michael Vaughan-Lee independently obtained a classification of nilpotent Lie rings with order p^7 , and the numbers above were also confirmed by his computation.

For some classifications of 7-dimensional nilpotent Lie algebras over infinite fields we refer to [Ancochéa et al. 1989, Romdhani 1989, Gong 1998, Goze & Remm]

The author's **GAP 4** program was also used to obtain a classification of nilpotent \mathbb{F}_2 -Lie algebras with dimension 8 and 9. The total number of such Lie algebras is 1836 and 26711. More detailed information about the possible types can be found in Tables 6–8 of Section 5.

The classifications of nilpotent Lie algebras in Theorem 1.1 are available in **GAP 4** format on the author's web site (www.sztaki.hu/~schneider/Research/SmallLie/).

4. IMPLEMENTATION OF THE ALGORITHMS

Implementations of all procedures described in Section 2 are available in the **GAP 4** computer algebra package **Sophus**. This program can freely be downloaded from the author's web page (www.sztaki.hu/~schneider/Research/Sophus). The current version of **Sophus** contains

- (i) a program to compute the cover of a nilpotent Lie algebra;
- (ii) a program to compute the automorphism group of a nilpotent Lie algebra;
- (iii) a program to compute the set of immediate descendants of a nilpotent Lie algebra;
- (iv) a program to check if two nilpotent Lie algebras are isomorphic.

The full implementation of these procedures is nearly 4000 lines long.

The classifications presented in the previous section were computed on several Pentium 4 computers between 1.7 and 2.5 GHz CPU speed and 1-2 GB memory. The computation of the list of the \mathbb{F}_2 -Lie algebras with dimension at most 6 takes only a few seconds, while those of dimension 7 takes about 3 minutes.

Determining the remaining classes of nilpotent Lie algebras in Theorem 1.1 is more complicated and requires human intervention. Most of the descendant computations for the 8 and 9-dimensional Lie algebras over \mathbb{F}_2 could easily be carried out. However, computing the 8-dimensional descendants of the 6-dimensional abelian Lie algebra requires finding representatives of the $\mathrm{GL}(6, 2)$ -orbits on the set of 178,940,587 allowable subspaces under the action in (1). In the computation of the 9-dimensional descendants of the 7-dimensional abelian Lie algebra, the number of allowable subspaces is 733,006,703,275. In such cases I applied the Cauchy-Frobenius Lemma (see [Eick & O'Brien 1999, Section 4]) to predict the number of descendants. Then I used either the ideas of O'Brien's extended algorithm presented in [O'Brien 1991, Section 2], or the existing classification of 2-groups of order at most 2^9 . In the latter case I constructed Lie algebras associated with the 2-central series filtration of the groups, tested them for isomorphism, and eliminated the duplicates.

For computing the 7-dimensional descendants of the 5-dimensional abelian Lie algebras over \mathbb{F}_3 and \mathbb{F}_5 , I used the result of the corresponding computation over \mathbb{F}_2 . The Cauchy-Frobenius Lemma implies that the number of these Lie algebras is the same over \mathbb{F}_2 , \mathbb{F}_3 , and \mathbb{F}_5 . It is possible to interpret the structure constants table of the \mathbb{F}_2 -Lie algebras over \mathbb{F}_3 and \mathbb{F}_5 and obtain the required lists. Then the algebras in these lists were tested for non-isomorphism.

The most difficult problem when computing the immediate descendants of a nilpotent Lie algebra is computing the orbits of the allowable subspaces under the representation (1). Further, for computing the automorphism group of an immediate descendant, the stabiliser of an allowable subspace must also be calculated; see Section 2. These orbit-stabiliser computations were carried out adopting the procedures described in [Eick et al. 2002].

5. THE TABLES

$[6][6]$	$[5, 1][4]$	$[5, 1][2]$	$[4, 2][3]$
1	1	1	1
$[4, 2][2]$	$[4, 1, 1][3]$	$[4, 1, 1][2]$	$[4, 1, 1][1]$
3	1	1	1
$[3, 3][3]$	$[3, 2, 1][2]$	$[3, 2, 1][1]$	$[3, 1, 2][3]$
1	3	3	1
$[3, 1, 2][2]$	$[3, 1, 1, 1][2]$	$[3, 1, 1, 1][1]$	$[2, 1, 2, 1][2]$
3	2	4	1
$[2, 1, 2, 1][1]$	$[2, 1, 1, 1, 1][1]$		
2	6		

Table 1: The nilpotent Lie algebras of dimension 6 over \mathbb{F}_2

$[6][6]$	$[5, 1][4]$	$[5, 1][2]$	$[4, 2][3]$
1	1	1	1
$[4, 2][2]$	$[4, 1, 1][3]$	$[4, 1, 1][2]$	$[4, 1, 1][1]$
3	1	1	1
$[3, 3][3]$	$[3, 2, 1][2]$	$[3, 2, 1][1]$	$[3, 1, 2][3]$
1	3	3	1
$[3, 1, 2][2]$	$[3, 1, 1, 1][2]$	$[3, 1, 1, 1][1]$	$[2, 1, 2, 1][2]$
3	2	3	1
$[2, 1, 2, 1][1]$	$[2, 1, 1, 1, 1][1]$		
2	5		

Table 2: The nilpotent Lie algebras with dimension 6 over \mathbb{F}_3 and \mathbb{F}_5

$[7][7]$ 1	$[6, 1][5]$ 1	$[6, 1][3]$ 1	$[6, 1][1]$ 1
$[5, 2][4]$ 1	$[5, 2][3]$ 3	$[5, 2][2]$ 2	$[5, 1, 1][4]$ 1
$[5, 1, 1][3]$ 1	$[5, 1, 1][2]$ 1	$[5, 1, 1][1]$ 1	$[4, 3][4]$ 1
$[4, 3][3]$ 5	$[4, 2, 1][3]$ 3	$[4, 2, 1][2]$ 12	$[4, 2, 1][1]$ 9
$[4, 1, 2][4]$ 1	$[4, 1, 2][3]$ 3	$[4, 1, 2][2]$ 5	$[4, 1, 1, 1][3]$ 2
$[4, 1, 1, 1][2]$ 4	$[4, 1, 1, 1][1]$ 5	$[3, 3, 1][3]$ 1	$[3, 3, 1][2]$ 3
$[3, 3, 1][1]$ 2	$[3, 2, 2][3]$ 2	$[3, 2, 2][2]$ 21	$[3, 2, 1, 1][2]$ 9
$[3, 2, 1, 1][1]$ 13	$[3, 1, 2, 1][3]$ 1	$[3, 1, 2, 1][2]$ 11	$[3, 1, 2, 1][1]$ 8
$[3, 1, 1, 1, 1][2]$ 6	$[3, 1, 1, 1, 1][1]$ 21	$[2, 1, 2, 2][2]$ 3	$[2, 1, 2, 1, 1][2]$ 4
$[2, 1, 2, 1, 1][1]$ 14	$[2, 1, 1, 1, 2][2]$ 4	$[2, 1, 1, 1, 1, 1][1]$ 15	

Table 3: The nilpotent Lie algebras of dimension 7 over \mathbb{F}_2

$[7][7]$ 1	$[6, 1][5]$ 1	$[6, 1][3]$ 1	$[6, 1][1]$ 1
$[5, 2][4]$ 1	$[5, 2][3]$ 3	$[5, 2][2]$ 2	$[5, 1, 1][4]$ 1
$[5, 1, 1][3]$ 1	$[5, 1, 1][2]$ 1	$[5, 1, 1][1]$ 1	$[4, 3][4]$ 1
$[4, 3][3]$ 5	$[4, 2, 1][3]$ 3	$[4, 2, 1][2]$ 12	$[4, 2, 1][1]$ 9
$[4, 1, 2][4]$ 1	$[4, 1, 2][3]$ 3	$[4, 1, 2][2]$ 5	$[4, 1, 1, 1][3]$ 2
$[4, 1, 1, 1][2]$ 3	$[4, 1, 1, 1][1]$ 5	$[3, 3, 1][3]$ 1	$[3, 3, 1][2]$ 3
$[3, 3, 1][1]$ 5	$[3, 2, 2][3]$ 2	$[3, 2, 2][2]$ 21	$[3, 2, 1, 1][2]$ 8
$[3, 2, 1, 1][1]$ 14	$[3, 1, 2, 1][3]$ 1	$[3, 1, 2, 1][2]$ 10	$[3, 1, 2, 1][1]$ 12
$[3, 1, 1, 1, 1][2]$ 5	$[3, 1, 1, 1, 1][1]$ 17	$[2, 1, 2, 2][2]$ 3	$[2, 1, 2, 1, 1][2]$ 4
$[2, 1, 2, 1, 1][1]$ 16	$[2, 1, 1, 1, 2][2]$ 3	$[2, 1, 1, 1, 1, 1][1]$ 11	

Table 4: The nilpotent Lie algebras with dimension 7 over \mathbb{F}_3

$[7][7]$ 1	$[6, 1][5]$ 1	$[6, 1][3]$ 1	$[6, 1][1]$ 1
$[5, 2][4]$ 1	$[5, 2][3]$ 3	$[5, 2][2]$ 2	$[5, 1, 1][4]$ 1
$[5, 1, 1][3]$ 1	$[5, 1, 1][2]$ 1	$[5, 1, 1][1]$ 1	$[4, 3][4]$ 1
$[4, 3][3]$ 5	$[4, 2, 1][3]$ 3	$[4, 2, 1][2]$ 12	$[4, 2, 1][1]$ 9
$[4, 1, 2][4]$ 1	$[4, 1, 2][3]$ 3	$[4, 1, 2][2]$ 5	$[4, 1, 1, 1][3]$ 2
$[4, 1, 1, 1][2]$ 3	$[4, 1, 1, 1][1]$ 5	$[3, 3, 1][3]$ 1	$[3, 3, 1][2]$ 3
$[3, 3, 1][1]$ 6	$[3, 2, 2][3]$ 2	$[3, 2, 2][2]$ 21	$[3, 2, 1, 1][2]$ 8
$[3, 2, 1, 1][1]$ 18	$[3, 1, 2, 1][3]$ 1	$[3, 1, 2, 1][2]$ 10	$[3, 1, 2, 1][1]$ 16
$[3, 1, 1, 1, 1][2]$ 5	$[3, 1, 1, 1, 1][1]$ 16	$[2, 1, 2, 2][2]$ 3	$[2, 1, 2, 1, 1][2]$ 4
$[2, 1, 2, 1, 1][1]$ 18	$[2, 1, 1, 1, 2][2]$ 3	$[2, 1, 1, 1, 1, 1][1]$ 13	

Table 5: The nilpotent Lie algebras with dimension 7 over \mathbb{F}_5

$[8][8]$ 1	$[7, 1][6]$ 1	$[7, 1][4]$ 1	$[7, 1][2]$ 1
$[6, 2][5]$ 1	$[6, 2][4]$ 3	$[6, 2][3]$ 2	$[6, 2][2]$ 8
$[6, 1, 1][5]$ 1	$[6, 1, 1][4]$ 1	$[6, 1, 1][3]$ 1	$[6, 1, 1][2]$ 1
$[6, 1, 1][1]$ 1	$[5, 3][5]$ 1	$[5, 3][4]$ 5	$[5, 3][3]$ 16
$[5, 2, 1][4]$ 3	$[5, 2, 1][3]$ 13	$[5, 2, 1][2]$ 35	$[5, 2, 1][1]$ 12
$[5, 1, 2][5]$ 1	$[5, 1, 2][4]$ 3	$[5, 1, 2][3]$ 5	$[5, 1, 2][2]$ 14
$[5, 1, 1, 1][4]$ 2	$[5, 1, 1, 1][3]$ 4	$[5, 1, 1, 1][2]$ 5	$[5, 1, 1, 1][1]$ 5
$[4, 4][4]$ 4	$[4, 3, 1][4]$ 1	$[4, 3, 1][3]$ 20	$[4, 3, 1][2]$ 51
$[4, 3, 1][1]$ 34	$[4, 2, 2][4]$ 2	$[4, 2, 2][3]$ 40	$[4, 2, 2][2]$ 217
$[4, 2, 1, 1][3]$ 9	$[4, 2, 1, 1][2]$ 62	$[4, 2, 1, 1][1]$ 56	$[4, 1, 2, 1][4]$ 1
$[4, 1, 2, 1][3]$ 11	$[4, 1, 2, 1][2]$ 48	$[4, 1, 2, 1][1]$ 26	$[4, 1, 1, 1, 1][3]$ 6
$[4, 1, 1, 1, 1][2]$ 21	$[4, 1, 1, 1, 1][1]$ 39	$[3, 3, 2][4]$ 1	$[3, 3, 2][3]$ 15
$[3, 3, 2][2]$ 77	$[3, 3, 1, 1][3]$ 3	$[3, 3, 1, 1][2]$ 13	$[3, 3, 1, 1][1]$ 6
$[3, 2, 3][3]$ 28	$[3, 2, 2, 1][3]$ 11	$[3, 2, 2, 1][2]$ 164	$[3, 2, 2, 1][1]$ 84
$[3, 2, 1, 1, 1][2]$ 49	$[3, 2, 1, 1, 1][1]$ 88	$[3, 1, 2, 2][3]$ 3	$[3, 1, 2, 2][2]$ 37
$[3, 1, 2, 1, 1][3]$ 4	$[3, 1, 2, 1, 1][2]$ 69	$[3, 1, 2, 1, 1][1]$ 84	$[3, 1, 1, 1, 2][3]$ 4
$[3, 1, 1, 1, 2][2]$ 39	$[3, 1, 1, 1, 1, 1][2]$ 15	$[3, 1, 1, 1, 1, 1][1]$ 80	$[2, 1, 2, 3][3]$ 1
$[2, 1, 2, 2, 1][2]$ 26	$[2, 1, 2, 2, 1][1]$ 20	$[2, 1, 2, 1, 2][3]$ 2	$[2, 1, 2, 1, 2][2]$ 24
$[2, 1, 2, 1, 1, 1][2]$ 12	$[2, 1, 2, 1, 1, 1][1]$ 24	$[2, 1, 1, 1, 2, 1][2]$ 11	$[2, 1, 1, 1, 1, 1, 1][1]$ 47

Table 6: The nilpotent Lie algebras with dimension 8 over \mathbb{F}_2

$[9][9]$ 1	$[8, 1][7]$ 1	$[8, 1][5]$ 1	$[8, 1][3]$ 1
$[8, 1][1]$ 1	$[7, 2][6]$ 1	$[7, 2][5]$ 3	$[7, 2][4]$ 2
$[7, 2][3]$ 8	$[7, 2][2]$ 6	$[7, 1, 1][6]$ 1	$[7, 1, 1][5]$ 1
$[7, 1, 1][4]$ 1	$[7, 1, 1][3]$ 1	$[7, 1, 1][2]$ 1	$[7, 1, 1][1]$ 1
$[6, 3][6]$ 1	$[6, 3][5]$ 5	$[6, 3][4]$ 16	$[6, 3][3]$ 122
$[6, 2, 1][5]$ 3	$[6, 2, 1][4]$ 12	$[6, 2, 1][3]$ 35	$[6, 2, 1][2]$ 70
$[6, 2, 1][1]$ 18	$[6, 1, 2][6]$ 1	$[6, 1, 2][5]$ 3	$[6, 1, 2][4]$ 5
$[6, 1, 2][3]$ 14	$[6, 1, 2][2]$ 25	$[6, 1, 1, 1][5]$ 2	$[6, 1, 1, 1][4]$ 4
$[6, 1, 1, 1][3]$ 5	$[6, 1, 1, 1][2]$ 5	$[6, 1, 1, 1][1]$ 5	$[5, 4][5]$ 4
$[5, 4][4]$ 53	$[5, 3, 1][5]$ 1	$[5, 3, 1][4]$ 21	$[5, 3, 1][3]$ 210
$[5, 3, 1][2]$ 376	$[5, 3, 1][1]$ 200	$[5, 2, 2][5]$ 2	$[5, 2, 2][4]$ 40
$[5, 2, 2][3]$ 475	$[5, 2, 2][2]$ 834	$[5, 2, 1, 1][4]$ 9	$[5, 2, 1, 1][3]$ 64
$[5, 2, 1, 1][2]$ 233	$[5, 2, 1, 1][1]$ 133	$[5, 1, 2, 1][5]$ 1	$[5, 1, 2, 1][4]$ 11
$[5, 1, 2, 1][3]$ 48	$[5, 1, 2, 1][2]$ 180	$[5, 1, 2, 1][1]$ 37	$[5, 1, 1, 1, 1][4]$ 6
$[5, 1, 1, 1, 1][3]$ 21	$[5, 1, 1, 1, 1][2]$ 39	$[5, 1, 1, 1, 1][1]$ 47	$[4, 5][5]$ 2
$[4, 4, 1][4]$ 16	$[4, 4, 1][3]$ 74	$[4, 4, 1][2]$ 134	$[4, 4, 1][1]$ 53
$[4, 3, 2][5]$ 1	$[4, 3, 2][4]$ 37	$[4, 3, 2][3]$ 615	$[4, 3, 2][2]$ 2727
$[4, 3, 1, 1][4]$ 3	$[4, 3, 1, 1][3]$ 82	$[4, 3, 1, 1][2]$ 307	$[4, 3, 1, 1][1]$ 306
$[4, 2, 3][4]$ 28	$[4, 2, 3][3]$ 1377	$[4, 2, 2, 1][4]$ 11	$[4, 2, 2, 1][3]$ 382

Table 7: The nilpotent Lie algebras with dimension 9 over \mathbb{F}_2 (continued on the next page)

$[4, 2, 2, 1][2]$ 2679	$[4, 2, 2, 1][1]$ 719	$[4, 2, 1, 1, 1][3]$ 49	$[4, 2, 1, 1, 1][2]$ 496
$[4, 2, 1, 1, 1][1]$ 576	$[4, 1, 2, 2][4]$ 3	$[4, 1, 2, 2][3]$ 37	$[4, 1, 2, 2][2]$ 258
$[4, 1, 2, 1, 1][4]$ 4	$[4, 1, 2, 1, 1][3]$ 69	$[4, 1, 2, 1, 1][2]$ 470	$[4, 1, 2, 1, 1][1]$ 313
$[4, 1, 1, 1, 2][4]$ 4	$[4, 1, 1, 1, 2][3]$ 39	$[4, 1, 1, 1, 2][2]$ 191	$[4, 1, 1, 1, 1, 1][3]$ 15
$[4, 1, 1, 1, 1, 1][2]$ 80	$[4, 1, 1, 1, 1, 1][1]$ 213	$[3, 3, 3][4]$ 16	$[3, 3, 3][3]$ 642
$[3, 3, 2, 1][4]$ 2	$[3, 3, 2, 1][3]$ 115	$[3, 3, 2, 1][2]$ 787	$[3, 3, 2, 1][1]$ 326
$[3, 3, 1, 1, 1][3]$ 16	$[3, 3, 1, 1, 1][2]$ 86	$[3, 3, 1, 1, 1][1]$ 76	$[3, 2, 4][4]$ 12
$[3, 2, 3, 1][3]$ 248	$[3, 2, 3, 1][2]$ 424	$[3, 2, 3, 1][1]$ 218	$[3, 2, 2, 2][3]$ 44
$[3, 2, 2, 2][2]$ 908	$[3, 2, 2, 1, 1][3]$ 73	$[3, 2, 2, 1, 1][2]$ 1264	$[3, 2, 2, 1, 1][1]$ 1240
$[3, 2, 1, 1, 2][3]$ 33	$[3, 2, 1, 1, 2][2]$ 325	$[3, 2, 1, 1, 1, 1][2]$ 163	$[3, 2, 1, 1, 1, 1][1]$ 435
$[3, 1, 2, 3][4]$ 1	$[3, 1, 2, 3][3]$ 21	$[3, 1, 2, 2, 1][3]$ 26	$[3, 1, 2, 2, 1][2]$ 622
$[3, 1, 2, 2, 1][1]$ 302	$[3, 1, 2, 1, 2][4]$ 2	$[3, 1, 2, 1, 2][3]$ 73	$[3, 1, 2, 1, 2][2]$ 359
$[3, 1, 2, 1, 1, 1][3]$ 12	$[3, 1, 2, 1, 1, 1][2]$ 228	$[3, 1, 2, 1, 1, 1][1]$ 317	$[3, 1, 1, 1, 2, 1][3]$ 11
$[3, 1, 1, 1, 2, 1][2]$ 181	$[3, 1, 1, 1, 1, 1, 1][2]$ 47	$[3, 1, 1, 1, 1, 1, 1][1]$ 423	$[2, 1, 2, 3, 1][3]$ 5
$[2, 1, 2, 3, 1][2]$ 10	$[2, 1, 2, 2, 2][3]$ 19	$[2, 1, 2, 2, 2][2]$ 170	$[2, 1, 2, 2, 1, 1][2]$ 60
$[2, 1, 2, 2, 1, 1][1]$ 98	$[2, 1, 2, 1, 2, 1][3]$ 6	$[2, 1, 2, 1, 2, 1][2]$ 62	$[2, 1, 2, 1, 2, 1][1]$ 16
$[2, 1, 2, 1, 1, 1, 1][2]$ 40	$[2, 1, 2, 1, 1, 1, 1][1]$ 124	$[2, 1, 1, 1, 2, 2][2]$ 7	$[2, 1, 1, 1, 2, 1, 1][2]$ 45
$[2, 1, 1, 1, 2, 1, 1][1]$ 18	$[2, 1, 1, 1, 1, 1, 2][2]$ 32	$[2, 1, 1, 1, 1, 1, 1, 1][1]$ 124	

Table 8: The nilpotent Lie algebras with dimension 9 over \mathbb{F}_2 (started on the previous page)

6. ACKNOWLEDGMENTS

A large part of the research presented in this paper was carried out at the Technische Universität Braunschweig. I am grateful for Bettina Eick for her interest in this project. I am

also indebted to Michael Vaughan-Lee for sharing his results concerning the 7-dimensional Lie algebras.

REFERENCES

- [Ancochéa et al. 1989] J. M. Ancochéa-Bermúdez, and M. Goze, Classification des algèbres de Lie nilpotentes complexes de dimension 7, *Arch. Math. (Basel)* **52** (1989), no. 2, 175–185.
- [Batten et al. 1996] P. Batten, K. Moneyhun, and E. Stitzinger, On characterizing nilpotent Lie algebras by their multipliers, *Comm. Algebra* **24** (1996), no. 14, 4319–4330.
- [Batten & Stitzinger 1996] P. Batten and E. Stitzinger, On covers of Lie algebras, *Comm. Algebra* **24** (1996), no. 14, 4301–4317.
- [Eick & O’Brien 1999] B. Eick and E. A. O’Brien, Enumerating p -groups, *J. Austral. Math. Soc. Ser. A* **67** (1999), no. 2, 191–205.
- [Eick et al. 2002] Bettina Eick, C. R. Leedham-Green, and E. A. O’Brien, Constructing automorphism groups of p -groups, *Comm. Algebra* **30** (2002), no. 5, 2271–2295.
- [The GAP Group 2004] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4*, 2004, (www.gap-system.org).
- [Gong 1998] Ming-Peng Gong, *Classification of nilpotent Lie algebras of dimension 7 (over algebraically closed fields and \mathbb{R})*, PhD Thesis, University of Waterloo, 1998. Available at etd.uwaterloo.ca/etd/mpgong1998.pdf.
- [Goze & Khakimdjanoz 1996] Michel Goze and Yusupdjanoz Khakimdjanoz, *Nilpotent Lie algebras*, Kluwer Acad. Publ., Dordrecht, 1996.
- [Goze & Remm] Michel Goze and Elisabeth Remm, *Classifications of complex nilpotent Lie algebras*, (www.math.uha.fr/~algebre/goze/LIE/Lie.html).
- [Havas et al. 1990] G. Havas, M. F. Newman, and M. R. Vaughan-Lee, A nilpotent quotient algorithm for graded Lie rings, *J. Symbolic Comput.* **9** (1990), no. 5-6, 653–664.
- [Romdhani 1989] M. Romdhani, Classification of real and complex nilpotent Lie algebras of dimension 7, *Linear and Multilinear Algebra* **24** (1989), no. 3, 167–189.
- [O’Brien 1991] E. A. O’Brien, The groups of order 256, *J. Algebra* **143** (1991), no. 1, 219–235.
- [O’Brien 1990] E. A. O’Brien, The p -group generation algorithm, *J. Symbolic Comput.* **9** (1990), no. 5-6, 677–698.
- [O’Brien et al. 2004] M. F. Newman, E. A. O’Brien, and M. R. Vaughan-Lee, Groups and nilpotent Lie rings whose order is the sixth power of a prime, *J. Algebra* **278** (2004), no. 1, 383–401.
- [Newman et al. 1998] M. F. Newman, W. Nickel, and A. C. Niemeyer, Descriptions of groups of prime-power order, *J. Symbolic Comput.* **25** (1998), no. 5, 665–682.
- [Wilkinson 1988] D. Wilkinson, The groups of exponent p and order p^7 (p any prime), *J. Algebra* **118** (1988), no. 1, 109–119.

INFORMATICS LABORATORY, COMPUTER AND AUTOMATION RESEARCH INSTITUTE OF THE HUNGARIAN ACADEMY OF SCIENCES, 1111 BUDAPEST LÁGYMÁNYOSI U. 11., HUNGARY

E-mail address: csaba.schneider@sztaki.hu
 WWW: www.sztaki.hu/~schneider