SOME EXTREMAL CONTRACTIONS BETWEEN SMOOTH VARIETIES ARISING FROM PROJECTIVE GEOMETRY

ALBERTO ALZATI AND FRANCESCO RUSSO

INTRODUCTION

Let $\psi: X \to Y$ be a proper morphism with connected fibers from a smooth projective variety X onto a normal variety Y, i.e. a contraction. If $-K_X$ is ψ -ample, then ψ is said to be an extremal contraction and if moreover $Pic(X)/\psi^*(Pic(Y)) \simeq \mathbb{Z}$ then ψ is said to be an elementary extremal contraction or a Fano-Mori contraction. Usually contractions are divided into two types: birational and of fiber type, i.e. with dim $(X) > \dim(Y)$.

The importance of this study and the parallel development of the notion of extremal ray and of the possible contractions of these rays was established by Mori in [18]. In dimension 2 and over an arbitrary field only three types of Fano-Mori contractions exist: the inverse of the blow-up of a reduced and geometrically irreducible set of points defined over the field (birational), conic bundle structure and del Pezzo surfaces (of fiber type) and this classification is essentially equivalent to the theory of minimal models in dimension 2. In dimension 3 the theory is more delicate but there is a complete description for algebraically closed fields given by Mori in characteristic zero and by Kollár for arbitrary algebraically closed fields, see for example [17], theorem 1.32 or [18]. The common feature of extremal elementary contractions in dimension 3 (and 2) is that if $E \subseteq X$ is the exceptional locus of the contraction, i.e. the locus of positive dimensional fibers, then $\phi_{|E} : E \to \phi(E) \subseteq Y$ is equidimensional. Moreover in the birational case (and over an algebraically closed field) these contractions are always the inverse of the blow-up of a (possible singular) point or of a smooth curve, see *loc. cit*.

In dimension greater than three the situation is more complicated as it is shown by simple examples and no general result is known. Here we furnish some examples of Fano-Mori contractions $\psi : X \to Y$ between smooth projective varieties with $\dim(X) \ge 4$ defined over "sufficiently" small fields and in arbitrary characteristic, having some exceptional fibers of unexpected dimension. All these contractions come from classical projective geometry, or more precisely they are associated to the projective geometry of remarkable classes of varieties: varieties with one apparent double, triple, quadruple point or varieties defining *special* Cremona transformations. Surely many examples of the phenomenon of non-equidimensionality of the fibers of $\phi_{|E}$ were constructed and are well known at least in characteristic zero or

¹⁹⁹¹ Mathematics Subject Classification. Primary 14E05; Secondary 14N05.

Key words and phrases. birational map, elementary extremal contraction, multisecant lines. First author partially supported by the M.I.U.R. of the Italian Government in the framework

of the national Research Project "Geometry of algebraic varieties" (Cofin 2002).

Second author partially supported by CNPq (Centro Nacional de Pesquisa), grant 300961/2003-0 and by PRONEX-Algebra Comutativa e Geometria Algebrica.

over algebraically closed fields (see for example [8], pg. 35, or [17], pg. 44, and also [3] or [4]). In the examples presented below the accent is on the geometric nature of these constructions and on the close relations with classical results. Moreover, they are also the first instances, from a historical point of view, of a class of morphisms with isolated exceptional fibers, a phenomenon studied in detail, in a local setting, in [3], [4]. In these last two papers some examples of these phenomena were constructed over the complex field by using vector bundles and geometrical constructions.

In section 2 we construct interesting Fano-Mori contractions by considering codimension 2 determinantal smooth subvarieties of \mathbb{P}^m , $3 \leq m \leq 5$, see proposition 2.1. Then we furnish an example of a Fano-Mori birational contraction between smooth varieties not having equidimensional exceptional fibers (so that it is not the inverse of a blow-up with smooth center) by considering a very particular special quartoquartic Cremona transformation of \mathbb{P}^4 studied classically in the general case by Todd and Room, see [30] and [21]. The existence of this exceptional quarto-quartic special Cremona transformation seemed not to be noticed before, see example 2. We remark that the general quarto-quartic transformation does not give this exceptional phenomenon. The particular contraction is related to the construction of a determinantal degree 10 smooth surface containing plane quartic curves, whose existence was essentially suggested by Room, see example *iii*) on page 80 of [21].

An example of not equidimensional fiber type Fano-Mori contraction can be constructed from the congruence of trisecant lines to a Bordiga surface in \mathbb{P}^4 , a surface with one apparent triple point according to Severi ([29]). The double-ten of pointsplanes associated to it (see page 71 of [21]) shows that the associated contraction, which is of fiber type in dimension 4, has 10 exceptional two dimensional fibers, the focal locus of the congruence outside the Bordiga surface, see example 3. This example is in a certain sense also more classical and its origins are harder to trace.

In dimension 5 the determinantal construction furnishes an example of fiber type Fano-Mori contraction onto \mathbb{P}^4 coming from a degree 10 determinantal 3-fold $Z \subset \mathbb{P}^5$ having one apparent quadruple point. In the general case the congruence of its 4-secant lines has a smooth curve of exceptional 2 dimensional fibers. In a particular constructed example it has a curve of exceptional 2 dimensional fibers and exactly one 3 dimensional fiber corresponding to the unique singular point of the curve, see example 5. A general hyperplane section of this particular example furnishes a family of surfaces (and hence of interesting Fano-Mori contraction) of the type discussed in example 2.

From some very particular special quinto-quintic Cremona transformation of \mathbb{P}^5 we construct some Fano-Mori birational contractions between smooth 5-folds and \mathbb{P}^5 having general exceptional fiber of dimension 1 and a finite number of isolated exceptional fibers of dimension 2 or 3, example 4.

Once the path has been cleared we continued the study and furnished examples of different nature, whose interest appears also in other important problems. Continuing the study in dimension 4 we showed how the analysis given by Fano in [12] of general smooth cubic hypersurfaces in \mathbb{P}^5 passing through a del Pezzo quintic surface can be extended to arbitrary (infinite) fields and also to every smooth cubic hypersurface through a del Pezzo surface. Some particular smooth cubic hypersurfaces generate Fano-Mori birational contractions between these smooth 4-folds and \mathbb{P}^4 with general exceptional fiber of dimension 1 and with 1,2,3 or 4 exceptional fibers isomorphic to \mathbb{P}^2 . These fibers contract to singular points of the fundamental locus of the inverse map, which is the projection of a degree 9 normal (K3) surface of genus 8, see 4.1. This result and the analogous for cubic hypersurfaces through a degree 4 rational normal scroll in \mathbb{P}^5 are probably well known and were rediscovered many times. We take the opportunity of giving the exact classical reference and the extension to arbitrary fields.

In dimension 6 an interesting Fano-Mori birational contraction can be constructed from Semple and Tyrrell example of a quadro-quartic special Cremona transformation in \mathbb{P}^6 ([27]), given by quadrics through a general rational octic surface $S \subset \mathbb{P}^6$. In this case the associated Fano-Mori contraction $\psi : Bl_S(\mathbb{P}^6) \to \mathbb{P}^6$ is very interesting and has a rather intricate behaviour for the exceptional fibers of dimension 2: there are 28 isolated *planes* and a two dimensional family of nonisolated fibers which are *quadrics*, parametrised by a smooth quadric surface, see 5.1. Therefore there are isolated two dimensional fibers and a two dimensional family of exceptional two dimensional fibers in the same Fano-Mori contraction and therefore the local behaviour and the global behaviour of exceptional fibers can be completely different for the same elementary extremal contraction. Probably this is the most intriguing example of Fano-Mori birational contraction together with the next examples related to it. Indeed, one constructs a 3-fold $X \subset \mathbb{P}^7$ of degree 8 having one apparent double point and having S as its general hyperplane section; this 3-fold produces a Fano-Mori contraction of fiber type $\psi: Bl_X(\mathbb{P}^7) \to \mathbb{P}^6$ for which the behaviour of the exceptional fibers of dimension greater or equal to 2 is very interesting and it is treated completely by geometrical methods, see 6.1. We conclude by constructing a Fano-Mori birational contraction between smooth cubic hypersurfaces $Y \subset \mathbb{P}^7$ through the above mentioned 3-fold X and \mathbb{P}^6 and we study in detail the inverse map and all the exceptional fibers. In particular we show that $\phi^{-1}: \mathbb{P}^6 \dashrightarrow Y$ is given by quintic forms vanishing along a 4-fold $Z \subset \mathbb{P}^6$ of degree 13 and having at least a two dimensional locus of double points, over which the fiber of the contraction $\widetilde{\phi}: Bl_X(Y) \to \mathbb{P}^6$ consists of (non-isolated) quadric surfaces, see 6.2.

This list can be considered as a natural continuation of the classical study of examples of interesting geometrical phenomena. The guiding principle was the beginning of Semple and Tyrrell paper [27]: Any Cremona transformation defined by a homaloidal system of primals with a single irreducible non-singular base variety is a rare enough phenomenon to merit special study. We were deeply influenced by the classical papers and books [29], [27], [26], [28], [25], [21], [30], [12] and the more recent [11].

We conclude by remarking that even if these examples could have some interest for higher dimensional geometry from the point of view of contractions, the methods we used are completely elementary and geometric and all the results can be considered classical or generalizations of them. Last but not least, we claim no originality for the geometrical interpretations of these classical examples.

Acknowledgements. The second author would like to thank Professor Shepherd-Barron for asking him about the classical study of the general contractions in 4.1. This was the starting point of this list of examples.

1. Definitions and preliminary results

Let us recall the following definitions.

Definition 1. Let $\phi : \mathbb{P}^n \to \mathbb{P}^n$ be a birational map, which is not an isomorphism. Then the linear system $\phi_*^{-1}|\mathcal{O}(1)|$ is called a homaloidal system. If the map ϕ is given by the homogeneous forms F_0, \ldots, F_n of degree $d \geq 2$ defining a smooth, geometrically irreducible variety $X^1 = V(F_0, \ldots, F_n) \subset \mathbb{P}^n$, then ϕ is said a special Cremona transformation and the associated linear system a special homaloidal system. More generally, a linear system of hypersurfaces of \mathbb{P}^r is said to be subhomaloidal if the (closure of a) general fiber of the associated rational map $\phi: \mathbb{P}^r \to \mathbb{P}^n, r \geq n$, is a linear projective space. It is said to be *special* if the base locus scheme of the linear system is smooth and irreducible; it is said to be completely subhomaloidal if (the closure of) every fiber is a linear projective space. These definitions can also be extended to linear systems of hypersurfaces such that the associated map $\phi: \mathbb{P}^r \dashrightarrow Z \subseteq \mathbb{P}^m, Z = \overline{Im(\phi)}$, is birational onto the image, respectively has a linear projective space as general fiber.

Thus, let $\phi : \mathbb{P}^n \dashrightarrow \mathbb{P}^n$ be a special Cremona transformation of type (d_1, d_2) , which means that ϕ is given by a linear subsystem of $|\mathcal{O}(d_1)|$ and ϕ^{-1} by a linear subsystem of $|\mathcal{O}(d_2)|$. Let X¹ be the base locus scheme of ϕ , which is smooth and geometrically irreducible by definition, and let X^2 be the base locus scheme of ϕ^{-1} . Let $r_i = \dim(X^i)$ for i = 1, 2. Let $\pi : Bl_{X^1}(\mathbb{P}^n) \to \mathbb{P}^n$ stand for the structural map. We can consider $Bl_{X^1}(\mathbb{P}^n)$ contained in $\mathbb{P}^n \times \mathbb{P}^n$ so that there is a canonical diagram



where ψ is the projection onto the second factor.

Put $E_1 = \pi^{-1}(X^1)$ and $E_2 = \psi^{-1}(X^2)$. Take $H_1 \in |\pi^*(\mathcal{O}(1))|, H_2 \in |\psi^*(\mathcal{O}(1))|$ and let T_1 and T_2 stand, respectively, for the strict transform on $Bl_{X^1}(\mathbb{P}^n)$ = $Bl_{X^2}(\mathbb{P}^n)$ of a general line contained in \mathbb{P}^n and of a general line of the target \mathbb{P}^n .

Let us recall that, given a subvariety $W \subset \mathbb{P}^n$ and an integer $s \geq 2$, an *s*-secant line to W is a line in \mathbb{P}^n not contained in W and intersecting W in at least s points (counted with multiplicities over \overline{k}). The closure $\operatorname{Sec}_{s}(W) \subset \mathbb{P}^{n}$ of the union of all s-secant lines to W is the variety of s-secant lines. Of course, $Sec_2(W)$ is the ordinary secant variety of W.

The following lemma contains the preliminary results of [11] and it will be a fundamental tool for our investigation (\sim stands for linear equivalence).

Lemma 1.1. ([11], section 1) Let notations be as above.

- (i) $H_2 \sim d_1 H_1 E_1, H_1 \sim d_2 H_2 E_2, E_2 = (E_2)_{\text{red}} \text{ and } T_1 \cdot H_2 = d_1, T_2 \cdot H_1 =$ $d_2, E_1 \cdot T_2 = E_2 \cdot T_1 = d_1 d_2 - 1.$ Moreover, $E_2 \sim (d_1 d_2 - 1) H_1 - d_2 E_1.$
- (ii) $\psi(E_1) = \operatorname{Sec}_{d_2}(X^2)$ is a hypersurface of degree $d_1d_2 1$ and $\pi(E_2) =$ $\operatorname{Sec}_{d_1}(X^1)$ is a hypersurface of degree $d_1d_2 - 1$.
- (iii) $2 + r_1 = d_2[(n+1) d_1(n r_1 1)].$
- $\begin{aligned} & (\text{iv}) \quad 2 + r_2 = d_1 [(n+1) d_2 (n-r_2-1)]. \\ & (\text{v}) \quad h^0(\mathcal{I}_{X^1}(d_1)) = n+1, \ h^0(\mathcal{I}_{X^1}(d_1-1)) = 0, \ h^0(\mathcal{I}_{X^1}(1)) = 0 \ and \ X^1 \ is \ not \end{aligned}$ a complete intersection.

(vi) If the base field k is algebraically closed, then $\psi^{-1}(U) \to U$ is the blow-up of $U := \mathbb{P}^n \setminus \text{Sing}((X^2)_{\text{red}})$ along $(X^2)_{\text{red}} \cap U$. In U, $(X^2)_{\text{red}} = X^2$ so that X^2 is generically smooth.

We recall the following theorem of Danilov.

Theorem 1.2. ([9]) Let $f : X \to Y$ be a projective birational morphism of smooth algebraic spaces defined over an arbitrary field k such that $\dim(f^{-1}(y)) \leq 1$ for every $y \in Y$. Then f can be decomposed into a sequence of monoidal transformations with smooth centers of codimension 2.

The next remark collects some general facts for further reference.

Remark 1.3. Let $X \subset \mathbb{P}^m$, $m \ge n+2$, be an irreducible non-degenerate variety of dimension n. It is well known that $n+2 \le \dim(\operatorname{Sec}_2(X)) \le \min\{m, 2n+1\}$. We recall that, if the base field has characteristic zero, the main result of [20] assures $\dim(\operatorname{Sec}_{n+2}(X)) \le n+1$. This last property is true over an arbitrary field for smooth irreducible curves by the Trisecant Lemma, see for example [15] IV.3, and also for the examples of varieties $X \subset \mathbb{P}^{n+2}$ of section 2 by a direct verification, using the following elementary observation.

Suppose that for a general point $p \in \operatorname{Sec}_s(X) \subseteq \mathbb{P}^m$, $s \geq 2$, the union of the *s*-secant lines through p is a projective space $\mathbb{P}_p^r \subseteq \mathbb{P}^m$, $r \geq 2$, cutting scheme theoretically X along a hypersurface in \mathbb{P}_p^r . Let $X' \subset \mathbb{P}^{m-1}$ be a general hyperplane section of X. Then for a general point $p \in \operatorname{Sec}_s(X')$, the union of the *s*-secant lines through p is a projective space $\mathbb{P}_p^{r-1} \subseteq \mathbb{P}^{m-1}$, cutting scheme theoretically X' along a hypersurface in \mathbb{P}_p^{r-1} .

We conclude the section by combining all the above result in the following proposition.

Proposition 1.4. Let $\phi : \mathbb{P}^n \dashrightarrow \mathbb{P}^n$ be a special Cremona transformation of type (d_1, d_2) and let notations be as above. Then $\psi : \widetilde{\mathbb{P}^n} = Bl_{X^1}(\mathbb{P}^n) \to \mathbb{P}^n$ is an extremal elementary contraction between smooth varieties. The morphism ψ is the inverse of a blow-up with smooth center if and only if X^2 is smooth, i.e. if and only if ϕ^{-1} is special. If ϕ^{-1} is not special and if the base field is algebraically closed, then $\psi : \psi^{-1}(U) \to U$ is the blow-up of $U = \mathbb{P}^n \setminus Sing((X^2)_{red})$ along $X^2 \setminus Sing((X^2)_{red})$. If ϕ^{-1} is not special and $codim(X^2) = 2$, then $\psi : \psi^{-1}(V) \to V$ is the blow-up of $V := \mathbb{P}^n \setminus \{x \in X^2 : \dim(\psi^{-1}(x)) \ge 2\}$ along $X^2 \setminus \{x \in X^2 : \dim(\psi^{-1}(x)) \ge 2\}$.

Proof. By Zariski's Main theorem and by Zariski's connectedness theorem, the morphism ψ has connected fibers and it is elementary by lemma 1.1. Now we show that $-K_{\widetilde{\mathbb{P}^n}}$ is ψ -ample. Since ψ is given by $|d_1H_1 - E_1|$, on an arbitrary fiber F of ψ we have $d_1H_{1|F} = E_{1|F}$. Then from lemma 1.1 part iv) we get $-K_{\widetilde{\mathbb{P}^n}|F} = (n+1)H_{1|F} - (n-r_1-1)E_{1|F} = (n+1-d_1(n-r_1-1))H_{1|F} = \frac{2+r_1}{d_2}H_{1|F}$. The last parts follow from lemma 1.1 part vi), if $k = \overline{k}$, or by theorem 1.2 in the remaining case; recall that in the last case by the theorem of the dimension of the fibers $X^2 \setminus \{x \in X^2 : \dim(\psi^{-1}(x)) \ge 2\}$ is not empty.

2. Determinantal examples

In this section we will construct some examples of Fano-Mori contractions both birational and of fiber type by considering some codimension 2 determinantal smooth subvarieties in \mathbb{P}^4 and in \mathbb{P}^5 having an interesting and rich extrinsic geometry.

Let $A(\mathbf{x})$ be a $n + 1 \times n$ matrix of linear forms on \mathbb{P}^m with coordinates $\mathbf{x} = (x_0 : \ldots : x_m), m \ge n$. Suppose the n + 1 minors of maximal order n of $A(\mathbf{x}), F_0, \ldots, F_n$ define a smooth geometrically irreducible codimension 2 subvariety $X_{m-2} \subset \mathbb{P}^m$. By definition, the variety X_{m-2} is scheme theoretically defined by the forms F_i 's and, for example by using the Hilbert-Burch theorem (or the Eagon-Northcott complex), the ideal sheaf $\mathcal{I}_{X_{m-2}}$ has the minimal resolution

$$0 \to \mathcal{O}(-n-1)^n \stackrel{A(\mathbf{x})}{\to} \mathcal{O}(-n)^{n+1} \to \mathcal{I}_{X_{m-2}} \to 0.$$

In particular X_{m-2} is an arithmetically Cohen-Macaulay variety. The homogeneous forms F_0, \ldots, F_n define a rational map $\phi : \mathbb{P}^m \dashrightarrow \mathbb{P}^n$.

Let $\mathbf{y} = (y_0 : \ldots : y_n)$ be coordinates on the target \mathbb{P}^n . The closure of the graph of ϕ , Γ_{ϕ} , in $\mathbb{P}^m \times \mathbb{P}^n$ is $Bl_{X_{m-2}}(\mathbb{P}^m) = \widetilde{\mathbb{P}^m}$, which coincides outside the exceptional divisor E with the scheme of equation

$$\sum a_{i,j}(\mathbf{x})y_j = A(\mathbf{x})^t \cdot \mathbf{y}^t = \mathbf{0}.$$

In fact for $\mathbf{x} \in \mathbb{P}^m \setminus X_{m-2}$ by elementary linear algebra we get:

$$A(\mathbf{x})^t \cdot \mathbf{y}^t = \mathbf{0} \Leftrightarrow \mathbf{y} = (F_0(\mathbf{x}) : \ldots : F_n(\mathbf{x})) \Leftrightarrow (\mathbf{x}, \mathbf{y}) \in \Gamma_{\phi}.$$

There exists a $n \times (m+1)$ matrix of linear forms $B(\mathbf{y})$ such that

$$A(\mathbf{x})^t \cdot \mathbf{y}^t = B(\mathbf{y}) \cdot \mathbf{x}^t$$

Let $\widetilde{\phi} : \widetilde{\mathbb{P}^m} \to \mathbb{P}^n$ be the resolution of the rational map ϕ . Let $\mathbf{y} = \phi(\mathbf{x}), \mathbf{x} \in \mathbb{P}^m \setminus X_{m-2}$. The above discussion yields

$$\phi^{-1}(\mathbf{y}) = \{ \mathbf{x} \in \mathbb{P}^m \setminus X_{m-2} : B(\mathbf{y}) \cdot \mathbf{x}^t = \mathbf{0} \},\$$

whose closure in \mathbb{P}^m is clearly a linear space of dimension $m - \operatorname{rk}(B(\mathbf{y}))$, let us say $\mathbb{P}^{m-\operatorname{rk}(B(\mathbf{y}))}_{\mathbf{y}}$. In particular we deduce that the fibers of ϕ are all irreducible and that the rank of $B(\mathbf{y})$ at the general point of $Im(\phi)$ is equal to $\dim(Im(\phi))$ by the theorem of the dimension of the fibers.

It is also clear that, when $\mathbb{P}_{\mathbf{y}}^{m-\operatorname{rk}(B(\mathbf{y}))}$ is positive dimensional, it cuts X_{m-2} along a determinantal hypersurface of degree n. We leave the details to the reader. The geometrical interpretation is that if through a point $p \in \mathbb{P}^m \setminus X_{m-2}$ there passes a n-secant line to X_{m-2} , since this line is contracted by ϕ , the locus of n-secant lines to X_{m-2} through p is the projective space $\mathbb{P}_{\phi(p)}^{m-\operatorname{rk}(B(\phi(p)))}$ cutting X_{m-2} along a hypersurface whose equation is determinantal in this projective space. Moreover this geometric description implies that, by looking at ϕ , we can say that every fiber of ϕ is either a linear space, eventually of varying dimension, or it is isomorphic to a subvariety of a linear space of the form $\mathbb{P}^{r(q)} \times q$, $q \in \mathbb{P}^n$, contained in the exceptional divisor E. We will show below that ϕ is always dominant. If m = n, then ϕ is birational and it is a Fano-Mori contraction by proposition 1.4. If m > nthe same argument used in the proof of proposition 1.4 easily gives that $-K_{\widetilde{\mathbb{P}m}}$ is ϕ -ample. In any case, ϕ is a Fano-Mori elementary contraction. If $3 \leq m \leq 5$ it is known that examples of smooth varieties X_{m-2} as above exist in characteristic zero by Bertini's theorem for spanned vector bundles. It is not difficult to construct smooth, geometrically irreducible examples over suitable "small" fields, where the above construction applies as we see below. If $m \geq 6$ the above arguments apply but the variety X_{m-2} will be singular. For m = n and for $A(\mathbf{x})$ sufficiently general (for example of rank n and with good Fitting ideals), one obtains a series of Cremona transformations of type (n, n) investigated by classical geometers and in particular by Godeaux, see [13] and also [23], example 2.4.

The above discussion about the locus of *n*-secant lines through a general point of \mathbb{P}^m is a particular version of proposition 2.9.1 of [21]. This result of Room can be considered as the first case of the general situation described in proposition 3.1, see also [16], [33] and [23]. In fact Room's result is true for more general situations and determinantal varieties, see *loc. cit.*

If m = n (and if $3 \le n \le 5$), it is easy to see that $\operatorname{Sec}_n(X_{n-2}) \subsetneq \mathbb{P}^n$, for example by applying 1.3 to X_{n-2} or to its general hyperplane section. Therefore ϕ will be dominant and birational, i.e. $\phi : \mathbb{P}^n \dashrightarrow \mathbb{P}^n$ will be a special Cremona transformation of type (n, n), by lemma 1.1 iii), whose base locus is exactly X_{n-2} . The general fiber is necessarily a point because $\operatorname{Sec}_n(X_{n-2}) \subsetneq \mathbb{P}^n$, so that $\operatorname{rk}(B(\mathbf{y})) = n$ for a general point $\mathbf{y} \in \operatorname{Im}(\phi)$. By applying the same argument to the matrix of linear forms $B(\mathbf{y})$ and from the fact that ϕ^{-1} is given by forms of degree n, we get that the inverse map is given by the maximal minors of $B(\mathbf{y})$. The inverse map is not defined along a determinantal scheme X^2 of the same type, which in general will not be smooth but only generically smooth. From this description it is also clear that $\operatorname{Sing}(X^2)$ will be given by the vanishing of the $(n-1) \times (n-1)$ minors of $B(\mathbf{y})$ so that it is easily computable. One can look at remark 2.3 of [23] for an example of non-special Cremona transformation of similar type for which the maximal minors of $B(\mathbf{y})$ have a common factor so that the corresponding ϕ is not of type (n, n).

The proper transform of $\operatorname{Sec}_n(X_{n-2})$ on \mathbb{P}^n will be the geometrically irreducible divisor E_2 by lemma 1.1. The general fiber of $\widetilde{\phi}_{|E_2} : E_2 \to X^2$ has dimension 1 by the theorem of the fibers and it is an *n*-secant line to $X^1 = X_{n-2}$. By lemma 1.4, $\widetilde{\phi} : \widetilde{\phi}^{-1}(V) \to V$ is the blow-up of $V = \mathbb{P}^n \setminus \{x \in X^2 : \dim(\psi^{-1}(x)) \ge 2\}$ along $X^2 \setminus \{x \in X^2 : \dim(\psi^{-1}(x)) \ge 2\}$. By the above description in this case $\{x \in X^2 : \dim(\psi^{-1}(x)) \ge 2\} = \operatorname{Sing}(X^2)$.

If m = n + 1 (and to have smooth $X_{n-1} = X^1 \subset \mathbb{P}^{n+1}$, $2 \leq n \leq 4$), the description of the fibers of ϕ yields $\operatorname{Sec}_n(X^1) = \mathbb{P}^{n+1}$ and the fact that the general fiber F of the associated rational map $\phi : \mathbb{P}^{n+1} \dashrightarrow \mathbb{P}^n$ is a *n*-secant line, see remark 1.3. The contraction $\tilde{\phi}$ is easily seen to be extremal and elementary by arguing as in proposition 1.4. Moreover, if F is a one dimensional fiber as above, $-K_{\widetilde{\mathbb{P}^m}|_F} \simeq \mathcal{O}_{\mathbb{P}^1}(2)$. With the classical terminology, the variety X^1 has one apparent *n*-ple point. Take now an arbitrary hypersurface $Y = V(G) \subset \mathbb{P}^{n+1}$ of degree n+1 through $X_{n-2} \subset \mathbb{P}^{n+1}$. The restriction of ϕ is a birational isomorphism $\phi|_Y : Y \dashrightarrow \mathbb{P}^n$ because a general fiber of ϕ is a *n*-secant line to X_{n-2} not contained in Y, so that this general fiber residually cuts Y in a point outside X_{n-2} . Moreover the equation of an Y as above can be written as the determinant of a $(n+1) \times (n+1)$ matrix of linear forms on \mathbb{P}^{n+1} . In fact since $G(\mathbf{x}) = \sum_{i=0}^n l_i(\mathbf{x})F_i(\mathbf{x})$, the graph of $\phi|_Y$ inside $\mathbb{P}^{n+1} \times \mathbb{P}^n$ coincides outside $E \cap Y$ with the scheme given by the n+1bihomogeneous (1, 1) forms

$$A(\mathbf{x})^t \cdot \mathbf{y}^t = \mathbf{0},$$

$$\sum_{i=0}^{n} l_i(\mathbf{x}) y_i = 0,$$

which can be written as

(1) $\widetilde{A}(\mathbf{x})^t \cdot \mathbf{y}^t = \mathbf{0},$

where $\widetilde{A}(\mathbf{x})$ is the desired $(n + 1) \times (n + 1)$ matrix of linear forms on \mathbb{P}^{n+1} . The conclusion easily follows since $\det(\widetilde{A}(\mathbf{x}))$ is a homogeneous polynomial of degree n + 1 vanishing at the general point of Y. We remark that the same construction applies also for a geometrically reduced, not necessarily geometrically irreducible, determinantal variety $X_{n-1} \subset \mathbb{P}^{n+1}$, as soon as $\operatorname{Sec}(X_{n-1}) = \mathbb{P}^{n+1}$. By reverting the construction we have that a hypersurface of degree m in \mathbb{P}^m containing a codimension 2 geometrically reduced variety of the above type has an equation given by the determinant of a $m \times m$ matrix of linear forms on \mathbb{P}^m .

We collect the above discussion in the following proposition.

Proposition 2.1. Let $A(\mathbf{x})$ be a $n + 1 \times n$ matrix of linear forms on \mathbb{P}_k^m , k an arbitrary filed, with coordinates $\mathbf{x} = (x_0 : \ldots : x_m)$, $m \ge n$. Suppose the n + 1 minors of maximal order n of $A(\mathbf{x})$, F_0, \ldots, F_n define a smooth geometrically irreducible codimension 2 subvariety $X_{m-2} \subset \mathbb{P}^m$. The homogeneous forms F_0, \ldots, F_n define a rational map $\phi : \mathbb{P}^m \dashrightarrow \mathbb{P}^n$, which resolves to a Fano-Mori contraction $\widetilde{\phi} : \widetilde{\mathbb{P}^m} := Bl_{X_{m-2}}\mathbb{P}^m \to \mathbb{P}^n$.

If $3 \leq m = n \leq 5$, then $\phi : \mathbb{P}^n \to \mathbb{P}^n$ is a special Cremona transformation of type (n,n) and $\phi_{\widetilde{\phi}^{-1}(V)} : \widetilde{\phi}^{-1}(V) \to V$ is the blow-up of $V = \mathbb{P}^n \setminus \operatorname{Sing}(X^2)$ along $X^2 \setminus \operatorname{Sing}(X^2)$.

If $3 \leq m \leq 5$, n = m - 1, then $\phi : Bl_{X_{m-2}}\mathbb{P}^m \to \mathbb{P}^{m-1}$ is generically a conicbundle, i.e. a one dimensional fiber F of ϕ is an m-1-secant line to X_{m-2} and $-K_{\widetilde{\mathbb{P}^m}|_F} \simeq \mathcal{O}_{\mathbb{P}^1}(2)$. Moreover any hypersurface $Y \subset \mathbb{P}_k^m$ of degree m through X_{m-2} has an equation given by the determinant of a $m \times m$ matrix of linear forms on \mathbb{P}_k^m (and the same is true for an arbitrary determinantal $X_{m-2} \subset \mathbb{P}^m$) and the restriction of ϕ to Y is birational.

Remark 2.2. We remark that the result concerning the rationality over the base field and the determinantal nature of arbitrary hypersurfaces through some determinantal reduced codimension 2 varieties, whose proof is elementary and geometric, can be considered as a suitable generalization of the following fact. Over a field k, a cubic surface $S \subset \mathbb{P}^3_k$ is rational over k and it has an equation given by the determinant of a 3×3 matrix of linear forms if it contains a degree 3 non-degenerate reduced connected curve defined over k, which is easily seen to have ideal generated by minors of a 2×3 matrix of linear forms on \mathbb{P}^3_k . Usually over an algebraically closed field one takes as the above curve either a twisted cubic or the union of two skew lines and a line intersecting them or the union of a line and a conic intersecting in one point or three concurrent lines. Since a smooth cubic surface over an algebraically closed field is easily seen to contain a degree 3 curve of the first three kinds, the above proposition applies to this class of hypersurfaces, a classical result known to Grassmann and Clebsch, see [14], [7], [25], pg. 123 and also [5], 6.4. The extension to arbitrary fields was considered in [24] and also in [5], 6.5. Surely for smooth cubic surfaces the point is to show geometrically the existence of the curve without knowing a priori the plane representation of the cubic surface. The morphism to \mathbb{P}^2 , i.e. the plane representation for the ones which are not cones, is obtained by solving the linear system (1).

We are now ready to begin the construction of examples of Fano-Mori contractions with exceptional behaviour. We begin by discussing the following special completely subhomaloidal linear system only as an elementary application of the above methods and also because we need this result in the last sections. As we see below, all the other examples presented later will be only suitable generalizations of this case.

Example 1. Quadrics through the Segre 3-fold in \mathbb{P}^5 or its general linear section and associated elementary contractions. Let n = 2 and m = 5 and let notations be as above. Let $\phi : \mathbb{P}^5 \dashrightarrow \mathbb{P}^2$ be the associated completely subhomaloidal linear system, by abusing language, of quadrics through $X^1 = \mathbb{P}^1 \times \mathbb{P}^2$. Modulo projective transformations we can suppose that $A(\mathbf{x})$ is the following 3×2 matrix of linear forms on \mathbb{P}^5 , whose minors give rise to X^1 :

$$\left(egin{array}{ccc} x_0 & x_3 \ x_1 & x_4 \ x_2 & x_5 \end{array}
ight)$$
 .

The 2×6 matrix $B(\mathbf{y})$ is

 $\left(\begin{array}{ccccc} y_0 & y_1 & y_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & y_0 & y_1 & y_2 \end{array}\right),$

so that every fiber of $\widetilde{\phi}: \widetilde{\mathbb{P}^5} \to \mathbb{P}^2$ is a \mathbb{P}^3 cutting X^1 along a quadric surface.

Example 2. (Todd-Room contraction). A Fano-Mori birational contraction between a smooth projective 4-fold and \mathbb{P}^4 having an isolated exceptional fiber of dimension 2. Let m = n = 4 in the above construction with $A(\mathbf{x}) \in M_{5\times 4}(k[\mathbf{x}])$, k to be chosen during the construction. If it exists, $X^1 \subset \mathbb{P}^4$ is a smooth degree 10 and genus 11 surface defined over k, whose ideal is generated by 5 quartic forms having the first syzygies generated by the linear ones and defining a special Cremona transformation. A general $A(\mathbf{x})$ will provide such an example but we aim at discussing all the cases defining a smooth X^1 .

Let $\tilde{\phi} : \mathbb{P}^4 = Bl_{X_1}(\mathbb{P}^4) \to \mathbb{P}^4$ be the resolution of the associated special Cremona transformation $\phi : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$. By proposition 2.1 ϕ is of type (4, 4) and $\tilde{\phi}^{-1}(V) \to V$ is the blow-up of $V = \mathbb{P}^4 \setminus \operatorname{Sing}(X^2)$ along $X^2 \setminus \operatorname{Sing}(X^2)$. Recall that $\operatorname{Sing}(X^2) =$ $\{y \in \mathbb{P}^4 : \operatorname{rk}(B(\mathbf{y})) \leq 2\}$. If $\mathbf{y} \in \operatorname{Sing}(X^2)$, then $\tilde{\phi}^{-1}(\mathbf{y})$ is a plane in $\mathbb{P}^4 \times \mathbf{y}$ cutting X^1 scheme theoretically along a plane quartic determinantal curve. The singular points of the base-locus scheme X^2 are then in bijective correspondence with quartic plane curves contained in X^1 . This number is necessarily finite as it is easy to see, eventually passing to \overline{k} .

Now we show that the singular points of X^2 are of multiplicity 2 for a general quartic hypersurface containing X^2 and of multiplicity 3 for X^2 . In fact, the image of a general plane passing through a singular point $p_0 \in \text{Sing}(X^2)$ under the rational map induced by the linear system of quartics through X^2 , restricted to the plane, is linked (4, 4) to the union of X^1 and the corresponding plane over p_0 , so that it has degree 5. It follows that the base locus of the plane system of quartics consists of

16-5=11 points: p_0 with multiplicity h and other t points with multiplicity 1 such that $11 = h^2 + t$. As the dimension of that system of plane quartics is 5, it is easy to see that h = 2 and t = 7 so that p_0 is a point of multiplicity $\deg(X^2) - t = 10-7=3$ for X^2 .

The above analysis also shows that the exceptional planes will be points of multiplicity 2 for E_2 , the blow-up of $\operatorname{Sec}_4(X^1)$ along X^1 , and that the tangent cone at a singular point $p \in \operatorname{Sing}(X^2)$ is a cone over a twisted cubic, in fact it has dimension 2 and degree 3. This is in accordance with the main-theorem of [3], pg. 256. Note that the tangent cone is linked (2, 2) to the tangent plane at some point of a smooth Bordiga surface passing through p and linked (4, 4) to X^2 .

By proposition 2.1, $\phi : \widetilde{\mathbb{P}^4} \to \mathbb{P}^4$ is a Fano-Mori contraction, which is the inverse of a blow-up with smooth center if and only if $\operatorname{Sing}(X^2) = \emptyset$.

We proceed by constructing an example of a smooth surface X^1 such that $\operatorname{Sing}(X^2) = (0 : 0 : 0 : 0 : 1)$ and defined over "sufficiently" large (or small, depending on the point of view) fields. To this aim it is sufficient to construct X^1 as given by the matrix of linear forms $A(\mathbf{x})$

$$\begin{pmatrix} -2x_1 + x_0 & -2x_2 + x_0 & 2x_0 & -x_1 - x_4 \\ x_3 + x_0 & -x_1 + x_2 & x_1 - 2x_3 & 2x_2 - x_3 \\ -x_1 - x_3 & -x_3 + 2x_4 & -2x_4 + x_0 & x_1 - x_0 \\ -2x_1 + x_4 & -x_2 - x_0 & x_2 + x_3 & x_2 + x_4 \\ x_3 & x_4 & x_3 & x_4 \end{pmatrix}$$

One verifies that X^1 is smooth and that the matrix $B(\mathbf{y})$ is

$$\begin{pmatrix} -2y_0 - y_2 - 2y_3 & 0 & y_1 - y_2 + y_4 & y_3 & y_0 + y_1 \\ -y_1 & -2y_0 + y_1 - y_3 & -y_2 & 2y_2 + y_4 & y_0 - y_3 \\ y_1 & y_3 & -2y_1 + y_3 + y_4 & -2y_2 & 2y_0 + y_2 \\ -y_0 + y_2 & 2y_1 + y_3 & -y_1 & -y_0 + y_3 + y_4 & -y_2 \end{pmatrix}$$

If we fix a point $p \in \mathbb{P}^4$ and solve the system of linear equations

$$B(p) \cdot \mathbf{x}^t = 0$$

we get the fiber $\phi^{-1}(p)$. In particular, it is immediate to see that $\phi^{-1}(0:0:0:0:1)$ is the plane $x_3 = x_4 = 0$ which cuts X^1 along a smooth quartic plane curve. An easy computation on $B(\mathbf{y})$ shows that $\operatorname{Sing}(X^2) = (0:0:0:0:1)$ so that the are no other 2-dimensional exceptional fibers.

We remark that the general case of this quarto-quartic transformation is studied in [30], while the existence of a surface X^1 of the above type containing plane quartic curves is suggested by Room in example iii) of page 80 of [21]. Room does not give an explicit example of such an X^1 containing plane quartic curves. For this reason we called the above example(s) *Todd-Room contraction*.

Example 3. (Bordiga conic-bundle). A Fano-Mori contraction of fiber type between a smooth projective 4-fold and \mathbb{P}^3 with 10 isolated 2-dimensional fibers. Take m = 4 and n = 3 in the above construction and let us consider a general matrix of linear forms $A(\mathbf{x}) \in M_{4\times 3}(k[\mathbf{x}])$, k an arbitrary field. If it defines a smooth surface $X^1 \subset \mathbb{P}^4$, then X^1 is a degree 6 and genus 3 Bordiga surface, which is arithmetically Cohen-Macaulay. By Bertini's theorem in characteristic zero and over an algebraically closed field a general $A(\mathbf{x})$ will define such a surface. A random choice over a suitable field will also give such examples.

Let $\phi : \mathbb{P}^4 \longrightarrow \mathbb{P}^3$ be the associated map. By the general discussion in remark 1.3 $\operatorname{Sec}_3(X^1) = \mathbb{P}^4$ and through the general point of \mathbb{P}^4 there passes a unique trisecant line to X^1 , i.e. the Bordiga surface has one apparent triple point, see [29]. By proposition 2.1 ϕ is dominant and for general $\mathbf{y} \in \mathbb{P}^3$ we have $\operatorname{rk}(B(\mathbf{y})) = 3$. If $\operatorname{rk}(B(\mathbf{y})) \leq 2$ for some $\mathbf{y} \in \mathbb{P}^3$, then necessarily $\operatorname{rk}(B(\mathbf{y})) = 2$ and the corresponding plane $\mathbb{P}^2_{\mathbf{y}}$ cuts X^1 along a determinantal cubic plane curve. To compute the number of these planes, we can calculate the degree of the locus of points $\mathbf{y} \in \mathbb{P}^3$ such that $\operatorname{rk}(B(\mathbf{y})) \leq 2$. If we assume that $B(\mathbf{y})$ is a general 3×5 matrix of linear forms on \mathbb{P}^3 then the above locus consists of exactly 10 points by well known formulas, see for example [21], p. 42. This gives the existence of 10 exceptional two dimensional fibers.

We remark that the above description could be also obtained by looking at the plane representation of a general Bordiga surface, eventually passing to \overline{k} . One easily verifies that there are exactly 10 cubics on such a surface, forming a double 10 with the ten chosen points for the plane representation. In fact a general Bordiga surface is the blow-up of \mathbb{P}^2 at 10 general points which is embedded by the linear system of plane quartics through these points.

By proposition 2.1 a quartic hypersurface through a Bordiga surface has equation given by the determinant of a 4×4 matrix of linear forms on \mathbb{P}^4 and it is rational over k.

The study of this examples is very classical and attracted the attention of classical geometers for its remarkable geometry (congruence of lines, etc, etc.).

Example 4. A Fano-Mori birational contraction between a smooth projective 5-fold and \mathbb{P}^5 having one dimensional exceptional fibers except for an isolated 2 dimensional (or 3 dimensional) fiber.

Take m = n = 5 in the general construction and let $X^1 \subset \mathbb{P}^5$ be a degree 15 and genus 26 smooth 3-fold defined by a matrix of linear forms $A(\mathbf{x}) \in M_{6\times 5}(k[\mathbf{x}])$ for a suitable field k.

Since $\operatorname{Sec}_5(X^1) \subsetneq \mathbb{P}^5$ by remark 1.3, we have a special Cremona transformation $\phi : \mathbb{P}^5 \dashrightarrow \mathbb{P}^5$. By lemma 1.1 $\operatorname{Sec}_5(X^1)$ is a geometrically irreducible hypersurface (of degree 24). The inverse of ϕ is not defined along a codimension 2 determinantal scheme X^2 , which is generically smooth and defined by the maximal minors of $B(\mathbf{y})$. The exceptional locus E_2 of $\tilde{\phi} : Bl_{X_1}(\mathbb{P}^5) = \widetilde{\mathbb{P}^5} \to \mathbb{P}^5$ is then irreducible of dimension 4 and it is smooth outside $\tilde{\phi}^{-1}(\operatorname{Sing}(X^2))$.

From the above geometrical description it follows that, for a general choice of $A(\mathbf{x})$, X^2 is smooth. In fact, for example by counting parameters we have that a general X^1 does not contain any plane quintic determinantal curve, neither a quintic determinantal surface spanning a \mathbb{P}^3 .

Now we furnish examples of matrices $A(\mathbf{x})$, defined over suitable fields k, such that X^2 has only one singular point p and such that $\tilde{\phi}^{-1}(p) \simeq \mathbb{P}^2$, respectively

 $\widetilde{\phi}^{-1}(p) \simeq \mathbb{P}^3$. The singularities of both X_2 and E_2 can be described explicitly as in the previous example, but we leave the details to the interested reader.

Let us construct an example such that X^2 has one singular point such that the corresponding fibre of $\tilde{\phi}$ will be isomorphic to an "isolated" \mathbb{P}^2 . We can take $A(\mathbf{x})$ equal to

1	$-2x_1 - 2x_4$	$-2x_1 + x_5 - x_0$	$-x_2 + x_5 - x_0$	$x_3 - 2x_4 + 2x_0$	$-2x_2 + 2x_4$	١
	$2x_2 + x_0$	$2x_2 - 2x_0$	$x_2 - x_3 - x_4$	$-x_1 + x_2$	$-2x_3 + 2x_4$	
	$x_4 + x_5$	$-x_4 + x_5 + x_0$	$x_1 + 2x_0$	$-x_2 - 2x_5 + x_0$	$x_1 + x_5$	
	$-x_3 - x_0$	$-2x_1 + x_3$	$-2x_1 + x_4$	$-x_2 + x_4$	$-x_1 - x_3 - 2x_5$	1
	$x_1 + x_3 - 2x_5$	$x_3 - 2x_5$	$-2x_3 + 2x_5$	$x_1 + 2x_5$	$x_2 - x_0$	
1	x_4	x_5	x_4	x_5	x_4	/

To have an example with an isolated exceptional fiber of dimension 3, in fact isomorphic to a \mathbb{P}^3 , it is sufficient to take $A(\mathbf{x})$ equal to

1	$-2x_1 - 2x_4$	$-2x_1 + x_5 - x_0$	$-x_2 + x_5 - x_0$	$x_3 - 2x_4 + 2x_0$	$-2x_2 + 2x_4$	١
l	$2x_2 + x_0$	$2x_2 - 2x_0$	$x_2 - x_3 - x_4$	$-x_1 + x_2$	$-2x_3 + 2x_4$	
	$x_4 + x_5$	$-x_4 + x_5 + x_0$	$x_1 + 2x_0$	$-x_2 - 2x_5 + x_0$	$x_1 + x_5$	
l	$-x_3 - x_0$	$-2x_1 + x_3$	$-2x_1 + x_4$	$-x_2 + x_4$	$-x_1 - x_3 - 2x_5$	·
l	$x_1 + x_3 - 2x_5$	$x_3 - 2x_5$	$-2x_3 + 2x_5$	$x_1 + 2x_5$	$x_2 - x_0$	
1	x_4	x_5	x_3	x_5	x_4	/

It is clear that the plane $x_3 = x_4 = x_5 = 0$, respectively the 3-plane of equation $x_4 = x_5 = 0$, is mapped to (0:0:0:0:1) and it is easy to verify that this is the only singular point of X_2 , i.e. that the corresponding linear space is the unique exceptional fiber of the unexpected dimension.

We now discuss the last example of this kind, a smooth 3-fold $X^1 \subset \mathbb{P}^5$ of degree 10 and genus 11 having one apparent quadruple point. Its general hyperplane section is the surface considered in example 2. Firstly we discuss the general case and then we study an exceptional phenomenon for the fibers

Example 5. (Determinantal 5-dimensional conic bundle) A Fano-Mori birational contraction between a smooth 5-fold and \mathbb{P}^4 having an isolated 3-dimensional fiber in a one dimensional locus of 2-dimensional fibers. Take m = 5 and n = 4 in the general construction. Let us discuss firstly the general case to construct a conic bundle $\tilde{\phi}: \mathbb{P}^5 \to \mathbb{P}^4$ such that the general fiber is a 4-secant line to X^1 and having a smooth curve $C \subset \mathbb{P}^4$ of 2 dimensional fibers isomorphic to \mathbb{P}^2 . To this aim let us take $A(\mathbf{x})$ equal to

1	$x_1 + x_4$	$x_1 + x_5 - x_0$	$-x_2 + x_5 - x_0$	$x_3 + x_4 - x_0$	١
I	$-x_2 + x_0$	$-x_2 + x_0$	$x_2 - x_3 - x_4$	$-x_1 + x_2$	
l	$x_4 + x_5$	$-x_4 + x_5 + x_0$	$x_1 - x_0$	$-x_2 + x_5 + x_0$.
I	$-x_3 - x_0$	$x_1 + x_3$	$x_1 + x_4$	$-x_2 + x_4$	
($x_1 + x_3 + x_5$	$x_3 + x_5$	$x_3 - x_5$	$x_1 - x_5$	/

The corresponding matrix $B(\mathbf{y})$ is equal to

12

1	$y_0 + y_4$	$-y_1$	$-y_3 + y_4$	$y_0 + y_2$	$y_2 + y_4$	$y_1 - y_3$	\
	$y_0 + y_3$	$-y_1$	$y_3 + y_4$	$-y_2$	$y_0 + y_2 + y_4$	$-y_0 + y_1 + y_2$	
	$y_2 + y_3$	$-y_0 + y_1$	$-y_1 + y_4$	$-y_1 + y_3$	$y_0 - y_4$	$-y_0 - y_2$	·
	$-y_1 + y_4$	$y_1 - y_2 - y_3$	y_0	$y_0 + y_3$	$y_2 - y_4$	$-y_0 + y_2$	/

The locus of points \mathbf{y} of \mathbb{P}^4 such that $\operatorname{rk}(B(\mathbf{y})) = 3$ is a smooth curve C of degree 20 and genus 26.

Now we can force the existence of an exceptional isolated fiber of greater dimension by taking the following matrix $A(\mathbf{x})$ defining a smooth 3-fold $X^1 \subset \mathbb{P}^5$ as above.

1	$x_1 + x_2 + x_4 + x_5$	$x_1 - x_5 - x_0$	$-x_2 + x_5 - x_0$	$x_3 + x_4 - x_0$)
1	$-x_2 + x_0$	$-x_2 + x_3 + x_0$	$x_2 - x_3 - x_4$	$-x_1 + x_2 - x_5$	
	$x_1 + x_3 + x_4 + x_5$	$-x_4 + x_5 + x_0$	$x_1 + x_3 - x_0$	$x_1 - x_2 + x_5 + x_0$	
	$-x_3 - x_0$	$x_1 + x_3$	$x_1 + x_4 - x_5$	$-x_2 + x_4$	
/	x_4	x_5	x_4	x_5	/

In this case the resulting curve $C \subset \mathbb{P}^4$ will have exactly one singular point at (0:0:0:0:1); the fibre over (0:0:0:1) is isomorphic to \mathbb{P}^3 . We leave the details to the reader.

We remark that a general hyperplane section of this smooth 3-fold $X^1 \subset \mathbb{P}^5$, would give a whole class of examples of surfaces $X^1 \subset \mathbb{P}^4$ originating examples of Fano-Mori birational contractions of the type described in example 2. In fact all these surfaces will contain a plane quartic curve, the intersection of the hyperplane with the quartic surface cut on X^1 by $\tilde{\phi}^{-1}((0:0:0:0:1)) \simeq \mathbb{P}^3$.

By proposition 2.1 a quintic hypersurface through a variety $X^1 \subset \mathbb{P}^5$ as above has equation given by the determinant of a 5 × 5 matrix of linear forms on \mathbb{P}^5 and it is rational over k.

3. More preliminary results

Let us recall that a set of homogeneous forms of fixed degree $d_1 \geq 2$ is said to verify *condition* K_{d_1} , see [33], if the trivial (or Koszul) syzygies of these forms are generated by the linear ones. This is a weakening of the condition of having the module of first syzygies generated by the linear ones and it is sufficient for many applications. Particular forms of this result were known to Room, see [21] proposition 2.9.1 for example, as we explained in the previous section. About *condition* K_{d_1} we have the following:

Proposition 3.1. ([33], [16]) Let $X^1 \subset \mathbb{P}^m$ be a smooth variety of dimension r_1 scheme theoretically defined by forms F_0, \ldots, F_n of degree d_1 satisfying condition K_{d_1} . Then the rational map $\phi : \mathbb{P}^m \dashrightarrow \mathbb{P}^n$ defined by the F_i 's is an embedding off $\operatorname{Sec}_{d_1}(X)$. Moreover the closure of a positive dimensional fiber of ϕ is a linear space in \mathbb{P}^m intersecting X^1 in a hypersurface of degree d_1 . In particular, if $\operatorname{Sec}_{d_1}(X_1) \subsetneq$ \mathbb{P}^n and if m = n, then ϕ is a special Cremona transformation.

In the sequel we will use the above proposition to study the case of birational maps between a projective space and a cubic hypersurface in \mathbb{P}^5 . About the proof of the proposition we recall that it relies on the fact that, outside the exceptional

divisor of the blow up of \mathbb{P}^m at X^1 , the graph $\overline{\Gamma_{\phi}} \subset \mathbb{P}^m \times \mathbb{P}^n$ coincides with the scheme given by the equations $y_i F_j - y_j F_i$. By conditions K_{d_1} it can be described by bihomogeneous forms of degree (1, 1).

Now let $Y \subset \mathbb{P}^{2n+1}$, $n \geq 2$, be a smooth hypersurface defined over a field k. Let $\phi: Y \dashrightarrow \mathbb{P}^{2n}$ be a birational map of type (d_1, d_2) . This means that ϕ is given by a linear subsystem of $|\mathcal{O}_Y(d_1)|$ and ϕ^{-1} by a linear subsystem of $|\mathcal{O}_{\mathbb{P}^{2n}}(d_2)|$. Let X be the base locus scheme of ϕ . We will assume that that $X \subset Y$ is smooth and connected, i.e. that ϕ is special.

Let $\pi : \widetilde{Y} = Bl_X(Y) \to Y$ stand for the structural map. As we can consider \widetilde{Y} contained in $\mathbb{P}^{2n+1} \times \mathbb{P}^{2n}$, there is a canonical diagram as follows:



where $\tilde{\phi}$ is the projection onto the second factor.

Let $S = \{u \in \mathbb{P}^{2n} : \dim(\widetilde{\phi}^{-1}(u)) \geq 1\}$. Since \mathbb{P}^{2n} is smooth, $E =: \widetilde{\phi}^{-1}(S)$ is a divisor, which is irreducible because $Pic(\widetilde{Y}) = \mathbb{Z} \oplus \mathbb{Z}$, see for example [11], proposition 1.3. Let Z denote the base locus scheme of $\widetilde{\phi}^{-1}$, so that $S = Z_{\text{red}}$. Let $E_1 = \pi^{-1}(X)$ and $E_2 = \widetilde{\phi}^{-1}(Z)$. Take $H_1 \in |\pi^*(\mathcal{O}_Y(1))|, H_2 \in |\widetilde{\phi}^*(\mathcal{O}_{\mathbb{P}^{2n}}(1))|$ and let T_1 and T_2 stand, respectively, for the strict transform on \widetilde{Y} of a general line contained in Y and of a general line of \mathbb{P}^{2n} .

Then we have the following generalization of lemma 1.1.

Lemma 3.2. Let notations be as above.

- (i) $H_2 \sim d_1 H_1 E_1$, $H_1 \sim d_2 H_2 E_2$, $E_2 = (E_2)_{\text{red}}$, $T_2 \cdot H_1 = d_2$ and $E_1 \cdot T_2 = d_1 d_2 1$. Moreover, $E_2 \sim (d_1 d_2 1) H_1 d_2 E_1$.
- (ii) φ(E₁) = Sec_{d2}(Z) is a hypersurface of degree d₁d₂ − 1; π(E₂) is a divisor in |O_Y(d₁d₂ − 1)|, which is the locus of the d₁-secant lines to X contained in Y.
- (iii) $2 + r_1 = d_2[2n 1 + d_1(r_1 + 1 2n)]$ so that $\widetilde{\phi} : \widetilde{Y} \to \mathbb{P}^{2n}$ is a Fano-Mori contraction.
- (iv) Suppose $\operatorname{codim}(Z) = 2$ and let $V = \mathbb{P}^{2n} \setminus \{z \in Z : \dim(\widetilde{\phi}^{-1}(z)) \geq 2\}$. Then $\widetilde{\phi}^{-1}(V)$ is the blowing up of V along $Z \setminus \{z \in Z : \dim(\widetilde{\phi}^{-1}(z)) \geq 2\}$, which is a smooth variety.
- (v) Suppose $k = \overline{k}$ and let $U = \mathbb{P}^{2n} \setminus \operatorname{Sing}(Z_{\operatorname{red}})$. Then $\phi^{-1}(U)$ is the blowing up of U along $Z_{\operatorname{red}} \cap U$. In U, $Z_{\operatorname{red}} = Z$ so that Z is generically smooth.

Some results of this type were used in [23] to classify quadro-quadric special birational maps from \mathbb{P}^{2m-2} and $\mathbb{G}(1,m)$. Clearly they can be used for the study of arbitrary birational map from \mathbb{P}^q onto projectively normal smooth varieties with Picard group isomorphic to \mathbb{Z} , generated by the hyperplane section, and containing a large family of lines.

In the sequel we will need the following geometric result, which relies strongly on the fact that a del Pezzo quintic surface $X \subset \mathbb{P}^5$ is a variety with one apparent double point. Let us recall that a smooth irreducible *n*-dimensional variety $X \subset \mathbb{P}^{2n+1}$ is said to be a variety with one apparent double point if through a general point of \mathbb{P}^{2n+1} there passes a unique secant line to X. This means that by projecting X from a general point of \mathbb{P}^{2n+1} into \mathbb{P}^{2n} the variety X acquires only one double point as its singularities.

Lemma 3.3. Let $X \subset \mathbb{P}^5$ a smooth del Pezzo surface of degree 5 and let $Y \subset \mathbb{P}^5$ be a smooth cubic hypersurface containing X. Then Y contains at most a finite number of planes spanned by the conics contained in X.

Proof. The conics contained in X are divided into 5 pencils, at least over \overline{k} . The union of the planes generated by one pencil of conics is a rational normal scroll, which is necessarily a Segre 3-fold $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$, see [22] and [1]. Hence X is contained in five Segre 3-folds defined over \overline{k} . If Y would contain infinitely many planes spanned by conics contained in X, then it necessarily would contain a Segre 3-fold, but this would force Y to be singular by the Grothendieck-Lefschetz theorem.

4. Fano's extremal elementary contractions from some smooth cubic hypersurface in \mathbb{P}^5 onto \mathbb{P}^4 or onto a smooth quadric $Q \subset \mathbb{P}^5$

In this section we prove the following results, which were originally proved by Fano in [12] for general cubic hypersurfaces of the mentioned types.

Theorem 4.1. ([12]) Let $Y \subset \mathbb{P}^5$ be a smooth cubic hypersurface defined over a field k and containing a quintic smooth del Pezzo surface X defined over k. Then the linear systems of quadrics through X restricted to Y defines an elementary extremal birational contraction $\tilde{\phi}: \tilde{Y} = Bl_X(Y) \to \mathbb{P}^4$. The inverse map $\tilde{\phi}^{-1}: \mathbb{P}^4 \dashrightarrow Y$ is given by quartic hypersurfaces passing through a normal surface $Z \subset \mathbb{P}^4$ of degree 9 and genus 8. For a general choice of Y through X, the surface Z is smooth, and it is a (non-minimal) K3 surface of degree 9 and genus 8, the blow-up at 5 points of a general degree 14 and genus 8 K3 surface, and $\tilde{\phi}$ is the inverse of the blow-up with center Z. The fiber over an isolated singular point of Z, if any, is a plane contained in Y and spanned by a conic $C \subset X \subset \mathbb{P}^5$. There exist smooth Y containing a finite number of planes spanned by conics in X; these Y furnish examples of extremal elementary birational contractions $\tilde{\phi}: \tilde{Y} \to \mathbb{P}^4$ having isolated 2 dimensional exceptional fibers isomorphic to \mathbb{P}^2 .

Remark 4.2. It is well known that smooth cubic hypersurfaces Y in \mathbb{P}^5 containing a del Pezzo quintic surface X are Pfaffian and describe an irreducible divisor in the moduli space of smooth cubic hypersurfaces, see for example [5]. This divisor coincides with the locus of smooth cubic hypersurfces containing a rational normal scroll X of degree 4, see [12]. In fact this was the main result and the main motivation of Fano's paper. The theorems, proved by Fano in the second part of [12], describe every smooth cubic hypersurface through such X's. The locus of smooth cubic hypersurfaces containing a plane forms a divisor too. The smooth cubic hypersurfaces giving Fano-Mori contractions, not inverse of a blow-up, are particular smooth cubic hypersurfaces in the intersection of the above divisors in the moduli space.

The general cubic hypersurfaces through such X's were later studied by various authors by using Riemann-Roch, Serre's duality, Kodaira's vanishing (so restricting to characteristic zero and algebraically closed fields) and similar or more advanced techniques, see for example [31] and [32]. Our approach will be completely elementary and geometric and will follow Fano's original argument. We use basics of liaison, the theorem of Danilov and the above generalizations of the preliminary results of [11].

The proof of theorem 4.1 will occupy the rest of this section. We recall that, by the same techniques, we could also prove the following theorem.

Theorem 4.3. ([12]) Let $Y \subset \mathbb{P}^5$ be a smooth cubic hypersurface defined over a field k and containing a rational normal scroll X of degree 4 defined over k. Then the linear system of quadrics through X, restricted to Y, defines an elementary extremal birational contraction $\tilde{\phi}: \tilde{Y} = Bl_X(Y) \to Q \subset \mathbb{P}^5$, where Q is a smooth quadric hypersurface. The inverse map $\tilde{\phi}^{-1}: Q \dashrightarrow Y$ is given by the restriction to Q of the linear system of cubic hypersurfaces passing through a normal surface $Z \subset Q$ of degree 10 and genus 7. For a general choice of Y through X, the surface Z is smooth and it is a (non-minimal) K3 surface of degree 10 and genus 7, the blow-up in a point of a general degree 14 and genus 8 K3 surface, and $\tilde{\phi}$ is the inverse of the blow-up with center Z. The fiber over a singular isolated point of Z, if any, is a plane contained in Y and spanned by a conic $C \subset X \subset \mathbb{P}^5$. There exist smooth Y containing a finite number of planes spanned by conics in X; these Y furnish examples of extremal elementary birational contractions $\tilde{\phi}: \tilde{Y} \to Q \subset \mathbb{P}^5$ having isolated 2 dimensional exceptional fibers isomorphic to \mathbb{P}^2 .

We will not give the proof of theorem 4.3 here, but we will state all the needed lemmas and propositions after the corresponding (and proved) statements needed for the proof of theorem 4.1. Let us begin with the following Lemma.

Lemma 4.4. Let $X \subset \mathbb{P}^5$ be a smooth quintic del Pezzo surface defined over a field k. Let H be the class of a hyperplane divisor of \mathbb{P}^5 . The linear system $|2H-X| = |H^0(I_X(2))|$ on \mathbb{P}^5 has X as base locus scheme and it defines a morphism $\tilde{\phi} : Bl_X(\mathbb{P}^5) \to \mathbb{P}^4$ such that every fiber is either isomorphic to a secant line to X or to a plane spanned by a conic contained in X.

Let $X' \subset \mathbb{P}^5$ be a smooth rational normal scroll of degree 4 defined over a field k. The linear system $|2H - X'| = |H^0(I'_X(2))|$ on \mathbb{P}^5 has X' as base locus scheme and it defines a morphism $\tilde{\phi}' : Bl_{X'}(\mathbb{P}^5) \to \mathbb{P}^5$, whose image is a smooth quadric hypersurface Q, such that every fiber is either isomorphic to a secant line to X' or to a plane spanned by a conic contained in X'.

Proof. It is well known that the ideal of X is generated by 5 quadratic forms having the first syzygies generated by the linear ones. Moreover, by applying proposition 3.1 to the general hyperplane section of X, we have that $\tilde{\phi}$ is surjective and the general fiber of $\tilde{\phi}$ is a secant line to X, so that X has one apparent double point (see also [1]). The conclusion now follows taking into account that all the fibers of $\tilde{\phi}$ are positive dimensional.

Let now $Y \subset \mathbb{P}^5$ be a smooth cubic hypersurface defined over k and containing X. Let $Y' \subset \mathbb{P}^5$ be a smooth cubic hypersurface defined over k and containing X', Let notations be as in the previous section and by abuse of language let us also denote by $\tilde{\phi} : \tilde{Y} = Bl_X(Y) \to \mathbb{P}^4$ the restriction of $\tilde{\phi}$ to \tilde{Y} , respectively by $\tilde{\phi}' : \tilde{Y}' = Bl_{X'}(Y) \to Q \subset \mathbb{P}^5$ the restriction of $\tilde{\phi}'$ to \tilde{Y}' .

Lemma 4.5. The morphism $\tilde{\phi} : \tilde{Y} = Bl_X(Y) \to \mathbb{P}^4$ is birational and its exceptional fibers are isomorphic either to a secant line to X contained in Y, general case, or to a plane contained in Y and spanned by a conic contained in X. The base locus Z of $\tilde{\phi}^{-1}$ is a 2-dimensional irreducible scheme with at most a finite number of singular points, which are exactly the images of the planes contained in Y, spanned by conics contained in X.

The morphism $\widetilde{\phi}': \widetilde{Y} = Bl_{X'}(Y) \to Q \subset \mathbb{P}^5$ is birational and its exceptional fibers are isomorphic either to a secant line to X', general case, or to a plane spanned by a conic contained in X'. The base locus Z' of $\widetilde{\phi}'^{-1}$ is a 2-dimensional irreducible scheme with at most a finite number of singular points, which are exactly the images of the planes contained in Y', spanned by conics contained in X'.

Proof. By the analysis of the fibers of $\tilde{\phi}: \widetilde{\mathbb{P}^5} \to \mathbb{P}^4$ given in the previous lemma, it follows that $\tilde{\phi}: \tilde{Y} \to \mathbb{P}^4$ is a birational morphism: a general secant line to X is not contained in Y and cuts Y only in one point outside X. Hence the general fiber of $\tilde{\phi}$ consists of one point. The other fibers can be only lines (in $\mathbb{P}^5 \times \tilde{\phi}^{-1}(p)$, the residual intersection of the plane of a conic in X, outside X) or eventually planes (these are the planes of the conics contained in X which are also planes contained in Y). By lemma 3.3 there are only a finite number of these planes contained in Y and the conclusion follows from lemma 3.2 part iv). In fact the general fiber of $\tilde{\phi}_{|E_1}: E_1 \to Z$ is a line, so that that Z is 2-dimensional.

Remark 4.6. The birationality of the restriction of ϕ to \tilde{Y} has the following algebraic interpretation. The graph of $\phi : \mathbb{P}^5 \dashrightarrow \mathbb{P}^4$ is $\widetilde{\mathbb{P}^5} = Bl_X(\mathbb{P}^5) \subset \mathbb{P}^5 \times \mathbb{P}^4$, which is given by forms of bidegree (1,1), outside the exceptional divisor E, by proposition 3.1. Points $(\mathbf{x}, \mathbf{y}) \in (\widetilde{\mathbb{P}^5} \setminus E) \times \mathbb{P}^4$ are of the following form $\mathbf{y} = (Q_0(\mathbf{x}) :$ $\dots : Q_4(\mathbf{x}))$, where the Q_i are the quadratic forms defining X. Let $F(\mathbf{x}) = 0$ be the equation of Y. The graph of $\phi : Y \dashrightarrow \mathbb{P}^4$ is $\tilde{Y} = Bl_X(Y)$. Outside the exceptional divisor this graph can be described by the above (1,1) forms together with the equation of Y "restricted" to $\widetilde{\mathbb{P}^5} \setminus E$. Since $F = \sum L_i Q_i$, with $L_i(\mathbf{x})$ linear forms, the restriction of $F(\mathbf{x})$ to $\widetilde{\mathbb{P}^5}$ outside E is defined by one more (1,1) form: $\sum L_i(\mathbf{x})y_i = 0$; so that the general fiber of ϕ becomes a point. This means that the cubic hypersurface gives rise to a linear syzygy among the equations defining X.

By the previous lemmas it follows that the 2-dimensional scheme Z parameterizes the secant lines to X contained in Y. Hence these secant lines form an irreducible 2-dimensional family. The inverse map $\tilde{\phi}^{-1} : \mathbb{P}^4 \dashrightarrow Y$ will be given by forms of degree d_2 having the base locus supported on Z.

Proposition 4.7. ([12]) Let notations be as above. Then $\tilde{\phi}^{-1} : \mathbb{P}^4 \dashrightarrow Y$ is given by quartics vanishing on Z, i.e. $d_2 = 4$, and $\pi(E_2) \in |\mathcal{O}_Y(7)|$. Moreover, $\tilde{\phi}$ is the blow-down of E_2 outside $\tilde{\phi}^{-1}(\operatorname{Sing}(Z))$ and the quintic del Pezzo surface is a locus of singular points of multiplicity 4 for $\pi(E_2)$.

The map $\widetilde{\phi}'^{-1}: Q \dashrightarrow Y'$ is given by cubics vanishing on Z', i.e. by $|\mathcal{O}_Q(3) - Z'|$, and $\pi(E_2) \in |\mathcal{O}_{Y'}(5)|$. Moreover, $\widetilde{\phi}'$ is the blow-down of E_2 outside $\widetilde{\phi}'^{-1}(\operatorname{Sing}(Z'))$ and the rational normal scroll X' is a locus of singular points of multiplicity 3 for $\pi(E_2)$. Proof. Since $r_1 = 2$ and n = 2 by lemma 3.2 part iii) we get $d_2 = 4$ and that, by lemma 3.2 part i), $E_2 \sim 7H_1 - 4E_1$, i.e. $le \propto^2$ rette del sistema Γ ricoprono una $M^{7\cdot3}$ per la quale X é superficie quadrupla, see [12], pg. 79.

We cannot resists in furnishing Fano's original geometric argument on page 79 of [12] proving that $d_2 = 4$, also to show the richness of his geometrical arguments and how it works perfectly. Let $\Gamma := \pi(E_2)$ be the 2-dimensional family of secant lines to X contained in Y. Fano shows that through a general point $p \in X$ there passes 4 lines of Γ , i.e. if $\Gamma \sim \alpha H_1 - \beta E_1$, then $\beta = 4$ and apparently he knows that $\beta = d_2$. Indeed, $E_2 \cdot \pi^{-1}(p) = (\alpha H_1 - \beta E_1) \cdot \pi^{-1}(p) = -\beta E_1 \cdot \pi^{-1}(p) = \beta$. The β lines of Γ passing through p are contained in $T_p(Y)$. Let $C = T_p(Y) \cap X$. Obviously C has degree 5 and it is a tangent hyperplane section of X, so that it has a double point in p. If we project C from p to a general \mathbb{P}^3 inside $T_p(Y)$ we get a 2-dimensional cone over a twisted cubic. The lines of Γ passing through p are the intersections of Y with this cone, out of C. As the intersection scheme has degree 9 we get: $\beta = 9 - 5 = 4$. From this he deduces $d_2 = 4$

Perhaps Fano was not aware that $d_2 = 4$ implies $\Gamma \in |\mathcal{O}_Y(7)|$, so he shows this last fact geometrically in the following way, giving another different proof of the above assertion, since by lemma 3.2 part i) we have $7 = 2d_2 - 1$. Let us recall that the 3-dimensional Γ is the family of the secant lines to X contained in Y. If $\Gamma \in |\mathcal{O}_Y(d)|$, then Γ intersects a general line $l \subset Y$ in d points. Then d is also the number of lines of Γ intersecting *l* since through a general point of *Y* there passes a unique secant line to X. By projecting X from l onto a disjoint \mathbb{P}^3 we get a degree 5 singular surface $\widetilde{X} := \pi_l(X)$, as l does not intersect X. One immediately sees that the double curve D of \widetilde{X} has degree 5, since a general plane section of \widetilde{X} is a plane quintic of geometric genus 1, the same sectional genus of X, so that it has 5 double points. Now, as X has one apparent double point, we can define Σ_l , the scroll of secant lines to X passing through (a general point of) l. A general \mathbb{P}^4 through l cuts Σ_l along a curve which is the union of l and the 5 secant lines to X which are contracted by π_l , hence deg $(\Sigma_l) = 6$. The secant lines of Γ intersecting l are the intersection of Y and Σ_l outside l and outside the curve $\pi_l^{-1}(D)$. Note that $\pi_l^{-1}(D)$ has degree 10 because π_l is a degree 2 map onto D. Therefore $d = 6 \cdot 3 - 1 - 10 = 7$, see [12], pg. 79.

The following lemma will conclude the proof of theorem 4.1.

Lemma 4.8. The base locus scheme $Z \subset \mathbb{P}^4$ of ϕ^{-1} is a degree 9 and sectional genus 8 normal surface containing 5 skew lines. The ideal sheaf of Z has the following minimal resolution:

$$0 \to \mathcal{O}(-6) \to \mathcal{O}(-5)^{\oplus 6} \to \mathcal{O}(-4)^{\oplus 6} \to \mathcal{I}_Z \to 0.$$

If Z is smooth, then the 5 skew lines are (-1)-curve and Z is the blow-up at 5 points of a genus 8 and degree 14 K3 surface $\widetilde{Z} \subset \mathbb{P}^8$; moreover, the surface Z is a linear projection of \widetilde{Z} by a suitable 3-plane cutting \widetilde{Z} in 5 points. If Z is singular, its singular points lie on these skew lines, they are of multiplicity 3 and their tangent cones are cones over a twisted cubic curve; there are at most 2 singular points on each of the 5 lines. In particular there are at most 10 singular points on Z.

The base locus scheme $Z' \subset Q \subset \mathbb{P}^5$ of ϕ'^{-1} is a degree 10 and sectional genus 7 normal surface containing a conic γ . If Z' is smooth, then the conic γ is a (-1)-curve and Z' is the blow-up at a point of a genus 8 and degree 14 K3 surface

 $\widetilde{Z} \subset \mathbb{P}^8$; moreover, the surface Z is the projection of \widetilde{Z} from a general tangent plane. If Z' is singular, its singular points lie on γ ; they are of multiplicity 3 and their tangent cones are cones over a twisted cubic curve.

Proof. By the previous lemma we know that Z has at most a finite number of singular points and we calculate its degree. Let us recall that $\tilde{\phi}$ is given by the linear system $|2H_1 - E_1|$ and let us take two general divisors $D_1, D_2 \in |2H_1 - E_1|$. Then $D_1 \cap D_2 := F$ is a smooth surface of degree 7 and sectional genus 3. By restricting the map $\tilde{\phi}$ to F, we obtain a morphism $\tilde{\phi} : F \to \mathbb{P}^2$. Its inverse is given by the linear system of plane quartic curves passing through deg(Z) points by the explicit description of $\tilde{\phi}^{-1}$. These points are smooth points for Z, otherwise the linear system of plane quartics would have some fixed singular point and F could not have sectional genus 3 (by the way this gives a direct proof that Z is generically reduced and hence generically smooth). Since F has degree 7, we have that deg(Z) = 4 \cdot 4 - 7 = 9 (see the footnote 25 on page 79 of [12]). Moreover lemma 3.2 yields that $\tilde{\phi}(E_1) = \operatorname{Sec}_4(Z)$ is a hypersurface of degree 7.

Let us now show that Z is locally Cohen-Macaulay. To get the resolution of the ideal sheaf of Z we will use some basic facts of liaison.

Let us take two general elements in $|4H_2 - E_2|$, they give rise to two general quartic hypersurfaces V_1 and V_2 in \mathbb{P}^4 , containing Z, such that $V_1 \cap V_2 = Z \cup Z^*$ as schemes, with Z" a degree 7 subscheme of \mathbb{P}^4 . Let us recall that, by lemma 3.2, we know that $|4H_2 - E_2|$ is the linear system on \widetilde{Y} corresponding to the hyperplane sections of Y. Hence, via $\widetilde{\phi}^{-1}$, Z" will correspond to the intersection of Y with a general \mathbb{P}^3 ; i.e. Z" is the image by $\widetilde{\phi}$ of the blow-up at 5 (= deg(X)) points of a smooth cubic surface G obtained by intersecting Y with the \mathbb{P}^3 corresponding to $V_1 \cap V_2$. The surface G is smooth by the general choice of V_1 and V_2 . The restriction of $\widetilde{\phi}$ to $G, \widetilde{\phi} : G \to Z$ ", is given by the linear system of quadric surfaces in \mathbb{P}^3 passing through the five points of intersection of the above \mathbb{P}^3 with X. By this description we get that Z" is reduced, irreducible and normal. We can also suppose that Z" is smooth at $\operatorname{Sing}(Z)$ because a general \mathbb{P}^3 will cut any plane $\widetilde{\phi}^{-1}(z), z \in \operatorname{Sing}(Z)$, at one point only. In any case Z" is locally Cohen-Macaulay and normal, so that Z is locally Cohen-Macaulay too by liaison and since it is non-singular in codimension 1 it is also normal. In conclusion Z is a reduced, normal, irreducible surface of degree 9.

If $\operatorname{Sing}(Z)$ is not empty a general quartic hypersurface V in \mathbb{P}^4 passing through $z \in \operatorname{Sing}(Z)$ will have a double point at z. In fact the multiplicity of such a V at $z \in \operatorname{Sing}(Z)$ will be the multiplicity at z of the quartic curve obtained by cutting V with a general plane of \mathbb{P}^4 passing through z. Let us consider the restriction of $\tilde{\phi}^{-1}$ to a generic plane of \mathbb{P}^4 passing through z. The image of this plane, by the restriction of $\tilde{\phi}^{-1}$, is a surface in \mathbb{P}^5 which is linked (4,4) to the union of X and the corresponding plane over z, so that it has degree 6. The map is given by the linear systems of plane quartic passing through z with multiplicity h and other t points with multiplicity 1 such that $10 = h^2 + t$. As the projective dimension of that system of plane quartics is 5, it is easy to see that h = 2 and t = 6 so that z is a point of multiplicity $\deg(Z) - t = 3$ for Z.

Now, let V_1, V_2 be two generic quartic hypersurfaces in \mathbb{P}^4 passing through Z, such that $V_1 \cap V_2 = Z \cup Z^n$. By the above discussion we have that Z^n is smooth at any $z \in \text{Sing}(Z)$. The tangent cone of Z at any singular point z is a cone over

a twisted cubic, in fact V_1 and V_2 have multiplicity 2 at z and their tangent cones at z are cones over quadrics of \mathbb{P}^3 , so that the tangent cone of Z at z is a degree 3 cone over a degree 3 curve which is contained in the intersection of two smooth quadrics and residual to a line, i.e. a twisted cubic. By a generalization of [19], proposition 4.1 (see also remark 1.12 of [2]), we deduce that, for general V_1, V_2, Z " is smooth.

In conclusion the surface Z is linked (4,4) to a smooth surface Z" having degree 7 and sectional genus 4; for this last fact recall that Z" is birational to the above mentioned G via the restriction of $\tilde{\phi}$. By looking at the resolution of the ideal sheaf of such a surface in [10] and by using liaison we get that the ideal sheaf of Z has minimal resolution

$$0 \to \mathcal{O}(-6) \to \mathcal{O}(-5)^{\oplus 6} \to \mathcal{O}(-4)^{\oplus 6} \to \mathcal{I}_Z \to 0.$$

Let us assume that Z is smooth. By liaison we also get that the sectional genus of Z is 8, that $q(Z) = h^1(\mathcal{O}_Z) = 0$ and that $h^0(\mathcal{O}_Z(K_Z)) = 1$, since $h^0(\mathcal{I}_{Z^n}(3)) = 1$. Let us recall that Z contains five skew lines $l_1, ..., l_5$, corresponding to the five pencils of conics contained in X (see lemma 3.3). Let H be a hyperplane section of Z so that $14 = H^2 + H \cdot K_Z = 9 + H \cdot K_Z$ implies $H \cdot K_Z = 5$. One gets $(K_Z - \sum l_i) \cdot H = 0$ so that $K_Z \sim \sum l_i$ and the 5 lines are the only (-1)-curves of Z. By blowing down these 5 exceptional (-1)-curves, we obtain a degree 14 and genus 8 smooth surface \widetilde{Z} such that $K_{\widetilde{Z}}$ is trivial and $q(\widetilde{Z}) = 0$, i.e. a genus 8 smooth K3 surface.

Let us assume that Z is singular. We know that the singular points of Z lie on the five lines quoted above. By lemma 3.3 we also know that X is a divisor of type (1,2) in five Segre 3-folds of \mathbb{P}^5 and that the singular points of Z lying on each line are in one to one correspondence with the planes of type (1,0) contained in the intersection of each Segre 3-fold with the cubic hypersurface Y. This hypersurface cuts a divisor of type (3,3) on each Segre 3-fold. As $X \subset Y$ we have a residual divisor of type (2,1) on each Segre 3-fold, and such divisors contain at most two planes of type (1,0). So that there are at most 10 singular points on Z.

The previous lemma concludes the proof of theorem 4.1. We can also analyze the singularities of E_2 along a plane Π , corresponding to $z \in \text{Sing}(Z)$, respectively $z' \in \text{Sing}(Z')$, [12], pg. 79.

Proposition 4.9. Let notations be as above. Then the general point of $\tilde{\phi}^{-1}(z)$, $z \in \text{Sing}(Z)$, is a point of multiplicity 2 for E_2 .

The general point of $\tilde{\phi}'^{-1}(z')$, $z' \in \operatorname{Sing}(Z')$, is a point of multiplicity 2 for E_2 .

Proof. Let notations be as in the proof of proposition 4.7. Let Π be any plane, contained in Y, and spanned by a conic C contained in X. Let us take a general line l in Y and let q be the intersection of Π with l. The degree 6 scroll Σ_l described in proposition 4.7 degenerates into the union of Π and a degree 5 scroll Σ'_l with the same directrix l. Since we can suppose that through every point of $l \setminus q$ there passes a unique secant line to X, the scroll Σ'_l is smooth outside $\Pi \cap \Sigma'_l$ and there passes only one line of this scroll through q. The double curve D degenerates into the union of a quartic curve and a line, which is the projection of the conic $C = \Pi \cap X$ from l (see the proof of proposition 4.7). The intersection with Σ'_l of a general hyperplane of \mathbb{P}^5 passing through l consists of four lines outside Π and a secant line l' to C through q. Then the number of lines of $\Gamma = \pi(E_2)$ intersecting l outside q is equal to $3 \cdot 5 - 8 - 1 - 1 = 5$ where 8 is the degree of the curve projecting itself onto D from l and the last two -1 are given by the existence of l and l'. This finally yields $\operatorname{mult}_q(E_2) = 2$ because $\Gamma \in |\mathcal{O}_Y(7)|$.

The fact that there exist smooth cubic hypersurfaces Y containing a smooth del Pezzo surface and some planes of the conics on the surface is part of the construction of a del Pezzo surface, or of a rational normal scroll of degree 4.

Proposition 4.10. There exist smooth cubic hypersurfaces $Y \subset \mathbb{P}^5$ containing a smooth quintic del Pezzo surface X and some of the planes spanned by a conic contained in X, respectively containing a rational normal scroll of degree 4 X' and some of the planes spanned by conics contained in X'.

Proof. Take the plane Π : $x_0 = x_1 = x_2 = 0$ in \mathbb{P}^5 and consider the Segre variety $M = \mathbb{P}^1 \times \mathbb{P}^2$ given by the vanishing of the 2×2 minors Q_i , i = 0, 1, 2 of the matrix of linear forms

$$\left(\begin{array}{rrrr} x_0 & x_1 & x_2 \\ x_3 & x_4 & x_5 \end{array}\right).$$

Take a general smooth quadric hypersurface Q_3 containing the plane Π . Then $Q_3 \cap M = \Pi \cup X$ with X a smooth del Pezzo surface of degree 5, realized as a divisor of type (1,2) in M. Take $Y = l_0Q_0 + \ldots + l_3Q_3$ with $l_i = 0$ general hyperplanes in \mathbb{P}^5 . Then by Bertini theorem, Y is non singular outside $\Pi \cup X$ and a direct computation shows that it is also non-singular along $\Pi \cup X$. The ideal of X can be generated by taking a fifth quadric Q_4 through X, not containing M and passing through another general plane of M. From this it is clear that the map $\phi : \mathbb{P}^5 \dashrightarrow \mathbb{P}^4$ contracts the variety M onto a line corresponding, for example, to the pencil $\langle Q_3, Q_4 \rangle$. The fiber of $\phi_{|M}$ is the unique \mathbb{P}^2 of the Segre residual to X obtained by intersecting the corresponding quadric with M.

One easily constructs examples of smooth cubic hypersurfaces Y through X, defined over "sufficiently" small fields and containing only one plane spanned by the conics on X, or two planes, belonging either to the same pencil or to different pencils. We leave the computational details to the interested reader. As we saw, 10 is the maximal number of planes of the above type contained in a smooth cubic hypersurface, at most two planes for each pencil, but we do not know if this maximal case really occurs.

We end the section by remarking some amusing consequences in terms of bilinear forms defining the graph of ϕ inside $\mathbb{P}^5 \times \mathbb{P}^4$, or better $\tilde{Y} \times \mathbb{P}^4$. We change a little the notations introduced in section 2.

From the resolution of \mathcal{I}_Z we know that there exists a 6×6 matrix $B(\mathbf{y})$ of linear forms given by the linear syzygies of six degree 4 generators of I_Z . By the above description and from the fact that $\tilde{\phi}^{-1}$ is given by the linear system of these quartics, we get some restrictions which $B(\mathbf{y})$ has to satisfy and we get equations for Y. Let us remark that $\tilde{Y} = Bl_X(Y) = Bl_Z(\mathbb{P}^4)$ is clearly the graph of $\tilde{\phi}$ inside $\mathbb{P}^5 \times \mathbb{P}^4$. Outside E_2 this graph is given by the equations $B(\mathbf{y}) \cdot \mathbf{x}^t = \mathbf{0}$ which are forms of bidegree (1, 1), where $\mathbf{x} = (x_0, \ldots, x_5)$. Take a (general) point $\mathbf{p} \in \mathbb{P}^5$. To get the coordinates of $\tilde{\phi}(p)$ one has to resolve the system of linear equations in $\mathbf{y} = (y_0, \ldots, y_4)$ given by $\mathbf{0} = B(\mathbf{y}) \cdot \mathbf{p}^t = A(\mathbf{p})^t \cdot \mathbf{y}^t$. The matrix $A(\mathbf{x})$ can be obtained, as always, by derivation with respect to the variables \mathbf{y} of the matrix of homogeneous (1, 1) forms $B(\mathbf{y}) \cdot \mathbf{x}^t$. We have that there will be no solutions for a general \mathbf{p} because $\tilde{\phi}^{-1}(\mathbb{P}^4)$ is the cubic hypersurface Y. This means that the 6×5 matrix $A(\mathbf{x})$ has rank 5 at the general point of \mathbb{P}^5 . On the contrary, at the general point of Y there will be a unique solution, translating into the fact that the six 5×5 minor of $A(\mathbf{x})$ contain a common irreducible factor of degree 3 which is an equation of Y. The six residual quadratic forms define a (proper) closed subscheme of Y, which in fact has to be a subscheme of $\pi(E_2) = \Gamma$. By a computer algebra system one easily verifies that this scheme is the original (or new, depending on the point of view) quintic del Pezzo surface.

5. Semple and Tyrrell extremal elementary contraction from a smooth 6-dimensional variety to \mathbb{P}^6

Let X^1 be the embedding in \mathbb{P}^6 of the blow-up $\tilde{\mathbb{P}}^2$ of \mathbb{P}^2 at eight points p_i , i = 1, ...8, in general position, given by the linear system of quartics passing through the p'_i s, (actually it is sufficient that they do not belong to a smooth conic and that no four of them are collinear). The surface X^1 is arithmetically Cohen-Macaulay and it is cut out by 7 quadrics since it is linearly normal, regular and it has degree $8 = 2 \operatorname{codim}(X^1)$, see [1]. During the rest of this section we will prove the following result.

Theorem 5.1. (Semple and Tyrrell birational Fano-Mori contraction, [27]) Let notations be as above. The map $\phi: \mathbb{P}^6 \dashrightarrow \mathbb{P}^6$ given by $|H^0(\mathcal{I}_{X^1}(2))|$ is a special Cremona transformation of type (2, 4). The inverse transformation is not defined along a subvariety $X^2 \subset \mathbb{P}^6$ of dimension 4 and degree 8, which is linked (4,4) to a general projection of the Veronese embedding $\nu_2(\mathbb{P}^4)$ from the \mathbb{P}^7 spanned by 8 general points on it. The variety X^2 has a singular locus consisting of 28 isolated triple points $p_{i,j}$, i < j, i = 1, ..., 8, (corresponding, via ϕ , to the planes of the 28 conics contained in X^1 which are given by the $\binom{8}{2}$ lines joining any couple of the 8 points in \mathbb{P}^2) and of a quadric surface of double points. The fibre over any double point of X^2 is a quadric surface. The triple points on X^2 have a cone over a Segre 3-fold $\mathbb{P}^1 \times \mathbb{P}^2$ as tangent cone. The map ϕ gives a birational extremal elementary contraction $\widetilde{\phi}: \widetilde{\mathbb{P}^6} = Bl_{X^1}(\mathbb{P}^6) \to \mathbb{P}^6$. The fibers of $\widetilde{\phi}_{|E_2} \to X^2$ are either proper transforms of secant lines to X^1 or proper transforms of the 28 planes of the conics contained in X^1 , mapping to the 28 isolated triple points $p_{i,j}$ of X^2 or quadric surfaces mapping to the points of a smooth quadric \tilde{Q} contained in X^2 (these being 2-dimensional non-isolated fibers). On $\mathbb{P}^6 \setminus (\widetilde{Q} \cup (\cup_{i,j} p_{i,j}))$ the morphism $\widetilde{\phi}$ is the blow-up of the smooth variety $X^2 \setminus (\widetilde{Q} \cup (\cup_{i,j} p_{i,j})).$

Let us briefly show geometrically, as indicated in [27], why the quadrics through X^1 define a Cremona transformation $\phi : \mathbb{P}^6 \dashrightarrow \mathbb{P}^6$ (see [27], [16] and also [1], section 6). At the same time we furnish the complete description of the fibers of ϕ .

Lemma 5.2. ([27], [16]) The surface $X^1 \subset \mathbb{P}^6$ is the complete intersection of two divisors of type (1,2) on a cone $W \subset \mathbb{P}^6$ over a Segre 3-fold $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5 \subset \mathbb{P}^6$. The vertex of W is $P := f(p_9)$ where p_9 is the ninth associated point to p_1, \ldots, p_8 and $f : \widetilde{\mathbb{P}^2} \hookrightarrow \mathbb{P}^6$ is the embedding the surface $\widetilde{\mathbb{P}^2}$. *Proof.* As $f: \mathbb{P}^2 \hookrightarrow \mathbb{P}^6$ is given by the linear system $|4H - p_1 - \ldots - p_8|$, where H is the class of a line in \mathbb{P}^2 , each member of the pencil of cubics $\{D_\lambda\}_{\lambda \in \mathbb{P}^1}$ through p_1, \ldots, p_9 is embedded by f as a quartic normal curve, and the general one will be elliptic. The linear spans $\mathbb{P}^3_\lambda = \langle D_\lambda \rangle, \lambda \in \mathbb{P}^1$, generate a cone $W \subset \mathbb{P}^6$ of vertex P over a Segre 3-fold $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$, which is the only smooth (rational) 3-dimensional scroll in \mathbb{P}^5 . The cone W contains X^1 and this surface is the complete intersection of two divisors of type (1, 2) on W (see [27], pg. 214 or [16], pg. 434), where a divisor D of type (1, 2) on W is the intersection of W with a hyperquadric $Q \subset \mathbb{P}^6$ containing a $\mathbb{P}^3 = R$ of the ruling, i.e. $Q \cap W = D \cup R$. Also from this description one can deduce that the ideal of X^1 is generated by 7 quadratic forms, three coming from the cone over the Segre 3-fold and the remaining four coming from the fact that each divisor of type (1, 2) on W is given, on W, by the restriction of two quadratic forms, see [27], pg. 214 or [16], pg. 434. □

Lemma 5.3. ([27]) For any point $p \in \text{Sec}(X^1) \setminus W$ let L_p be the locus of secant lines to X^1 passing through p. Then L_p consists either of a unique secant (or tangent) line \mathbb{P}_p^1 or of the whole plane $\Pi_{i,j}$ if p belongs to one of the 28 planes $\Pi_{i,j} = \langle C_{i,j} \rangle$ spanned by the 28 conics on X^1 .

Proof. Let us take a generic point $p \in \mathbb{P}^6 \setminus W = \operatorname{Sec}(W) \setminus W$. The locus of secant lines to W passing through p is a linear \mathbb{P}_p^4 cutting W along a rank 4 quadric Q_p which is singular at P, the vertex of W, by an obvious adaptation of example 1. Hence L_p is given by the secant lines to the intersection of Q_p with two divisors of type (1,2) on Q_p , $\widehat{Q_1}$, $\widehat{Q_2}$, i.e. by secant lines through p to a degree 4 curve $C_p \subset \mathbb{P}_p^4$ (see [27] pg. 206). Let us remark that $Q_p \cap \widehat{Q_1} \cap \widehat{Q_2} = C_p \cup P$. The degree 4 curve C_p can be:

- (i) a smooth quartic rational normal curve F, the embedding by f of the blow up of a line of \mathbb{P}^2 not passing through any of the points p_i ;
- (ii) the union of a *twisted cubic* T_j and a line E_j intersecting at one point, the embedding by f of the blow up of a line passing through a point p_j ;
- (iii) the union of a *conic* $C_{i,j}$ and two lines cutting the conic at two distinct points, the embedding by f of the blow up of a line $\langle p_i, p_j \rangle$.

Moreover we have that the locus of secant lines to C_p passing through a point $p \in \text{Sec}(X^1) \setminus W$ consists either of a unique secant or tangent line to one of the above reduced curves or of the whole plane $\prod_{i,j}$, if p belongs to one of the 28 planes $\prod_{i,j} = \langle C_{i,j} \rangle$. Recall that a quartic curve in \mathbb{P}^4 has a 4-secant plane only if it is the union of four lines; see also the next proposition where it is shown that systems of quadrics defining these curves satisfy condition K_2 yielding a(nother) proof of this geometrical fact by proposition 3.1.

We study the following more general situation.

Lemma 5.4. Let $C \subset \mathbb{P}^4$ be a reduced non-degenerated quartic curve as in one of the cases (i), (ii), (iii) of the previous lemma. Suppose also that C is the complete intersection of two divisors of type (1,2) of a rank 4 hyperquadric $Q \subset \mathbb{P}^4$, whose vertex will be indicated by $O, O \notin C$. Then the rational map defined by the linear system of quadrics passing through C defines a birational map $\mu : \mathbb{P}^4 \dashrightarrow \widetilde{Q} \subset \mathbb{P}^5$, where \widetilde{Q} is an irreducible quadric. The subsystem of quadrics passing also through O gives a birational map $\mu' : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$ not defined along $C \cup O$ and which is the composition of μ with the projection of \widetilde{Q} from $\mu(O)$. The map μ' is an isomorphism outside $\operatorname{Sec}(C) \cup \mu^{-1}[T_{\mu(O)}(\widetilde{Q}) \cap \widetilde{Q}] = \operatorname{Sec}(C) \cup Q$.

Proof. The curve C is a hyperplane section of one of the two rational scrolls of degree 4 in \mathbb{P}^5 , S(2,2) or S(1,3), which are scheme theoretically defined by 6 quadrics satisfying condition K_2 so that C is scheme theoretically defined by 6 quadrics satisfying condition K_2 . The rational map defined by these quadrics is a birational map, $\mu : \mathbb{P}^4 \dashrightarrow \widetilde{Q} \subset \mathbb{P}^5$, where \widetilde{Q} is an irreducible hyperquadric and μ is an isomorphism outside $\operatorname{Sec}(C)$, (see for example [27] or apply proposition 3.1 and a simple calculation with Chern classes to deduce that \widetilde{Q} is a quadric hypersurface). The subsystem of quadrics passing also through O gives a birational map $\mu' : \mathbb{P}^4 \dashrightarrow \mathbb{P}^4$ not defined along $C \cup O$ and which is the composition of μ with the projection of \widetilde{Q} from $\mu(O)$. The map μ' is an isomorphism outside $\operatorname{Sec}(C) \cup \mu^{-1}(T_{\mu(O)}(\widetilde{Q}) \cap \widetilde{Q}) = \operatorname{Sec}(C) \cup Q$. In fact $\mu^{-1}(T_{\mu(O)}(\widetilde{Q}) \cap \widetilde{Q})$ is a quadric hypersurface in \mathbb{P}^4 passing through O and C, i.e. an element of the linear system defining μ' . Since the exceptional locus of μ' consists only of this quadric hypersurface and $\operatorname{Sec}(C)$ and, by the definition of C, Q is contracted by μ' , the conclusion easily follows.

Now we can describe the fibers of ϕ and we can give a geometrical idea of the birationality of $\phi : \mathbb{P}^6 \dashrightarrow \mathbb{P}^6$.

Lemma 5.5. Let notations be as above. The map $\phi : \mathbb{P}^6 \dashrightarrow \mathbb{P}^6$ is birational and it is an isomorphism outside $\operatorname{Sec}(X^1)$. Let $\tilde{\phi} : \widetilde{\mathbb{P}^6} = Bl_{X^1}\mathbb{P}^6 \to \mathbb{P}^6$ its resolution. Moreover, for any point $p \in \operatorname{Sec}(X^1) \setminus W$ we have that $\tilde{\phi}^{-1}(\tilde{\phi}(p))$ is equal to the locus of secant lines to X^1 passing through p, i.e. is isomorphic to a secant or tangent line to X^1 or to one of the 28 planes $\prod_{i,j}$ if $p \in \prod_{i,j} \setminus X^1$. If \widetilde{W} is the strict transform of W on $Bl_{X^1}(\mathbb{P}^6)$, then for every $p \in \widetilde{W}$ we have that $\tilde{\phi}^{-1}(\tilde{\phi}(p))$ is isomorphic to a quadric surface.

Proof. We can suppose that, among the seven quadrics defining X^1 , Q_0, Q_1, Q_2 define the cone W. Let $\psi : \mathbb{P}^6 \setminus W \to \mathbb{P}^2$ be the rational map defined by the above net of quadrics. The closure of every fiber $\psi^{-1}(\psi(p)), p \in \mathbb{P}^6 \setminus W$ is a linear space equal to \mathbb{P}^4_p , see example 1. Hence the \mathbb{P}^4_p 's vary in a two dimensional family such that through every point of $\mathbb{P}^6 \setminus W$ there passes only one member of the family. Then the closure of any fiber $\phi^{-1}(\phi(p)), p \in \mathbb{P}^6 \setminus W$, is clearly contained in \mathbb{P}^4_p , so that it is equal to the closure in \mathbb{P}^4_p of the restriction of ϕ to \mathbb{P}^4_p . For any $p \in \mathbb{P}^6 \setminus W$, the map $\phi_p := \phi_{|\mathbb{P}^4_p}$ is not defined along the quartic curve C_p and in P and it is given by the linear system of quadrics scheme theoretically defining $C_p \cup P$, see lemmas 5.3 and 5.4. By lemma 5.4 the map $\phi_p := \mathbb{P}^4_p \subset \mathbb{P}^6$ is birational so that ϕ is birational.

Moreover, we can now describe all the positive dimensional fibers $\phi^{-1}(\phi(p))$ for every $p \in \text{Sec}(X^1) \setminus W$ by describing the fibers of the corresponding ϕ_p .

Putting all the above geometric descriptions together we finally obtain that the closure of the fiber $\phi^{-1}(\phi(p)), p \in \mathbb{P}^6 \setminus X^1$ is one of the following types: it reduces to p, if $p \notin \operatorname{Sec}(X^1)$; it is equal to a tangent line or a secant line to C_p , if $p \in \operatorname{Sec}(X^1) \setminus W$ and C_p has at most two components; if C_p contains one of the 28 conics $C_{i,j}$ of X^1 , then it is a secant or tangent line to C_p if $p \in \operatorname{Sec}(X^1) \setminus (\prod_{i,j} \cup W)$ and it is the whole plane $\prod_{i,j}$ if $p \in \prod_{i,j} \setminus C_{i,j}$; it is a hyperquadric if $p \in W \setminus X^1$, see [27]. In fact the cone W is contracted by ϕ onto a smooth quadric surface

 $\widetilde{Q} \subset \mathbb{P}(\langle Q_3, Q_4, Q_5, Q_6 \rangle)$. The closure of a fiber of $\phi_{|W} : W \dashrightarrow \widetilde{Q}$ is easily seen to be a quadric surface, see also [27].

Now we are able to conclude the proof of theorem 5.1 by describing the birational map $\widetilde{\phi} : Bl_{X_1}(\mathbb{P}^6) \to \mathbb{P}^6$.

Proof. (of theorem 5.1) By the previous lemma we have that the strict transform of $Sec(X^1)$ is 5-dimensional and it is contracted onto a 4 dimensional variety Z = $\phi(Sec(X^1)) = (X^2)_{red}$. The planes $\Pi_{i,j}$ are contracted into the points $\phi(\Pi_{i,j}) =$ $p_{i,j} \in Z \setminus \widetilde{Q}$. Every fiber of $\widetilde{\phi}$ is of dimension less or equal to 1 on $V = \mathbb{P}^6 \setminus$ $(\tilde{Q} \cup (\cup_{i,j} p_{i,j}))$ so that by theorem 1.2 the restriction of $\tilde{\phi}$ to $U = \tilde{\phi}^{-1}(V)$ is the blowing-up of \mathbb{P}^6 along the smooth variety $Z \setminus (\widetilde{Q} \cup (\bigcup_{i,j} p_{i,j}))$. By theorem 1.1 of [11], we also know that the points $p_{i,j}$ and the surface \widetilde{Q} are the singular locus of Z, so that $Z = X^2$. By reasoning as in the case discussed in section 4, one immediately deduces that X^2 is the base locus scheme of ϕ^{-1} and that ϕ^{-1} is the rational map given by the linear systems of quartics containing X^2 , by lemma 1.1 (see [27] pg. 207). Moreover we also get that the points $p_{i,j}$ are isolated triple points for X^2 , that \widetilde{Q} is the locus of double points for X^2 , and that the planes $\prod_{i,j}$ are loci of points of multiplicity 2 of the strict transform of $Sec(X^1)$, $E_2 = Sec(X^1)$, i.e. of the exceptional divisor of ϕ . The tangent cone of X^2 at the points $p_{i,j}$ is easily seen to be a cone over a Segre 3-fold $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$, see also [27]. Now, by taking two general quartic hypersurfaces Y_1 and Y_2 passing through X^2 , we have that $Y_1 \cap Y_2 = X^2 \cup (X^2)'$, with $(X^2)'$ the image of the restriction of ϕ to a general \mathbb{P}^4 inside the original \mathbb{P}^6 . Since X^1 has degree 8, the linear system giving the restriction is a linear system of quadrics in \mathbb{P}^4 passing through 8 points in general linear position, so that $(X^2)'$ is the projection of $\nu_2(\mathbb{P}^4)$ from the linear span of 8 general points on it and $\deg(X^2) = \deg((X^2)') = 8$.

We remark that, see [27], one can also show that X^2 is isomorphic to the projection of $\nu_2(\mathbb{P}^4)$ from the linear span of 8 suitable points on it; it is in fact a degree 8 variety in \mathbb{P}^6 having exactly the same singularities as X^2 . Then the singular double points of X^2 on the quadric surface $\tilde{Q} \subset X^2$ correspond to pairs of points in \mathbb{P}^4 imposing only one linear condition to the linear system of hyperquadrics in \mathbb{P}^4 passing through the 8 points spanning the center of the projection of $\nu_2(\mathbb{P}^4)$.

6. Cubic hypersurfaces through a degree 8 3-fold with one apparent double point

Once we have done the effort of studying in detail the geometry of the secant variety of the octic surface in the previous example, at least outside the cone over the Segre 3-fold, we can do the same for another interesting smooth variety $X \subset \mathbb{P}^7$ with one apparent double point and whose general hyperplane section is the above mentioned surface.

The construction is the following. Take a line $l \subset \mathbb{P}^7$ and a skew \mathbb{P}^5 inside the fixed \mathbb{P}^7 . Let $M \subset \mathbb{P}^7$ be the cone with vertex l above a Segre 3-fold $\mathbb{P}^1 \times \mathbb{P}^2$ inside the fixed \mathbb{P}^5 . Take the complete intersection of two general divisors of type (1,2) on M (see [6]). Notice that a divisor of type (1,2) on M is once again the intersection of M with a quadric hypersurface containing a \mathbb{P}^4 of the ruling. The resulting variety is a smooth 3-fold $X \subset \mathbb{P}^7$ of degree 8 such that $Sec(X) = \mathbb{P}^7$

and such that through the general point of \mathbb{P}^7 there passes a unique secant line to X. A general hyperplane section is clearly an octic surface of the type discussed in Semple and Tyrrell example of the previous section, see also [6]. The variety X is arithmetically Cohen-Macaulay and its ideal is generated by 7 degree 2 forms giving rise to, a rational map $\phi : \mathbb{P}^7 \dashrightarrow \mathbb{P}^6$ not defined exactly along X. By restricting to a general hyperplane section, it immediately follows that the general fiber of ϕ is a line, necessarily a secant line to X, i.e. $|H^0(\mathcal{I}_X(2))|$ is a special subhomaloidal system.

Proposition 6.1. (Semple and Tyrrell conic fibration) Let notations be as above, let $\phi : \mathbb{P}^7 \dashrightarrow \mathbb{P}^6$ be the rational map defined by the linear system of quadrics vanishing along X and let $\tilde{\phi} : Bl_X(\mathbb{P}^7) \to \mathbb{P}^6$ be the resolution of the indeterminacy of ϕ . Let \widetilde{M} be the strict transform of M in $Bl_X(\mathbb{P}^7)$. Then for every $p \in Bl_X(\mathbb{P}^7) \setminus \widetilde{M}$ the fiber $\tilde{\phi}^{-1}(\tilde{\phi}(p))$ is isomorphic to a linear space \mathbb{P}_p^k cutting X along a quadric hypersurface $Q_p = \mathbb{P}_p^k \cap X \subset \mathbb{P}_p^k$. There are three possibilities:

- (i) k = 1, \mathbb{P}_p^1 is a secant (or tangent) line to X and Q_p is a couple of points;
- (ii) k = 2, p belongs to one of the \mathbb{P}^4 's generated by a $S(1,2) \subset X$ and Q_p is the conic cut along S(1,2) by the plane \mathbb{P}_p^2 ;
- (iii) k = 3, p belongs to one of the 28 linear spaces $\Pi_{i,j} \simeq \mathbb{P}^3$, i < j, $i = 1, \ldots, 8$, generated by the 28 quadrics $Q_{i,j}$ contained in X, $\mathbb{P}_p^3 = \Pi_{i,j}$ and $Q_p = Q_{i,j}$.

For any $p \in \widetilde{M}$, the fiber consists of a three dimensional quadric hypersurface and every \mathbb{P}^4 of the ruling cuts the variety X along a degree 4 cone over an elliptic normal curve. In particular, $\widetilde{\phi} : Bl_X(\mathbb{P}^7) \to \mathbb{P}^6$ is a Fano-Mori contraction of fiber type having 28 isolated 3-dimensional fibers isomorphic to \mathbb{P}^3 , eight 3-dimensional families of 2-dimensional fibers isomorphic to \mathbb{P}^2 , each of them parameterized by a Segre 3-fold, and a 2-dimensional family of 3-dimensional fibers isomorphic to quadric hypersurfaces and parameterized by a smooth quadric surface.

Proof. By cutting with a general hyperplane section we get a Cremona transformation of the type described in Semple and Tyrrell example. The analysis of the previous section yields that each fiber of $\tilde{\phi}$ is either linear and of dimension 1, 2 or 3 or a 3 dimensional quadric hypersurfaces.

The locus of the secant lines through every point $p \in \mathbb{P}^7 \setminus M$ is a \mathbb{P}_p^5 cutting M along a rank 4 quadric Q_p . The secant lines to X passing through p are the secant lines through p of the intersection of Q_p with two divisors of type (1,2) on M, i.e. a 2-dimensional degree 4 (reduced) variety. For general p it will be a smooth rational normal scroll of degree 4, which is known to be a surface with one apparent double point. For special p the variety can degenerate into the union of a plane and a smooth degree 3 rational normal scroll S(1,2) intersecting along a line, or into the union of a quadric surface and two skew planes intersecting the quadric along a line. There are 28 quadric surface of this kind generating 28 \mathbb{P}^3 's as it is seen by cutting with a general hyperplane and recalling that a general hyperplane section contains 28 conics (or directly by representing X as the \mathbb{P}^1 -bundle $\mathbb{P}(\mathcal{E})$ over \mathbb{P}^2 , where \mathcal{E} is the very ample vector bundle given by the extension $0 \to \mathcal{O}_{\mathbb{P}^2} \to \mathcal{E} \to \mathcal{I}_{\{p_1,\dots,p_8\},\mathbb{P}^2}(4) \to 0, p_1,\dots,p_8$ in general position, see [6]). Moreover, the surfaces $S(1,2) \subset X$ are divided into 8 pencils corresponding to the pencils of lines passing through the 8 points in the above representation of X as a \mathbb{P}^1 -bundle. The last claims follow in the same way as in the previous section. \Box

Now we can produce our last example of Fano-Mori birational contraction. By abuse of notation we also call ϕ this contraction.

Theorem 6.2. Let $X \subset \mathbb{P}^7$ be as above and let $Y \subset \mathbb{P}^7$ be a smooth cubic hypersurface through X. Then $\phi_{|Y} : Y \dashrightarrow \mathbb{P}^6$ is birational. The morphism $\widetilde{\phi}: \widetilde{Y} = Bl_X(Y) \to \mathbb{P}^6$ is a Fano-Mori birational contraction. The general fiber of $\widetilde{\phi}_{|E_2}: E_2 \to Z := \widetilde{\phi}(E_2)$ corresponds to a secant line to X contained in Y, $Z \subset \mathbb{P}^6$ is a 4-fold of degree 13 and $\phi^{-1}: \mathbb{P}^6 \dashrightarrow Y$ is given by the linear system of quintics vanishing along Z. The cubic hypersurface Y contains at most a finite number of \mathbb{P}^4 's spanned by the cubic scrolls S(1,2) contained in X, at most a finite number of \mathbb{P}^4 's generating the cone M and clearly at most a finite number of the $\Pi_{i,j}$. By ϕ every \mathbb{P}^4 of the ruling of M contained in Y goes into a line in Z and this line is a locus of singular points for Z; moreover for each point p in the strict transform of this \mathbb{P}^4 the fiber $\phi^{-1}(\phi(p))$ is isomorphic to a 3-dimensional quadric in \mathbb{P}^4 ; in particular these 3-dimensional fibers vary in a one-dimensional family. By ϕ any \mathbb{P}^4 spanned by a cubic scroll S(1,2) and contained in Y goes into a plane contained in Z, and this plane is a locus of singular points for Z; moreover for any point p in the strict transform of this \mathbb{P}^4 the fiber $\phi^{-1}(\phi(p))$ is isomorphic to a plane. Every $\Pi_{i,j}$ contained in Y contracts into a triple point for Z. The remaining positive dimensional fibers of $\tilde{\phi}$ come from secant lines to X contained in Y or from quadric surfaces, which are the residual intersection, outside X, of Y with a 3-dimensional quadric in a \mathbb{P}^4 of the ruling of M not contained in Y; moreover these quadric surfaces vary in a two dimensional family parameterized by an open subset of a quadric surface. Hence Z has a two dimensional locus of singular points at least, even if Y does not contain any \mathbb{P}^4 of the ruling of M or any \mathbb{P}^4 spanned by a cubic scroll contained in X.

Proof. If Y contains infinitely many's \mathbb{P}^4 of the above mentioned families then it contains a 5 dimensional rational and singular scroll in \mathbb{P}^4 's of degree 3. In fact, in the first case Y would contain M, in the second case note that anyone of the eight families of degree 3 scrolls S(1,2) is contained in such a scroll. On the other hand, for instance by the Grothendieck-Lefchetz theorem, it is clear that Y cannot contain such a scroll, necessarily as a hyperplane section. The assertions about fibers of ϕ are then easily proved since we have only to restrict $\phi: Bl_X(\mathbb{P}^7) \to \mathbb{P}^6$ to $Bl_X(Y)$ and the fibers of ϕ are completely described in proposition 6.1. About the singular points of Z we can argue as in the previous section. About the last assertions we recall that the intersection of X with any \mathbb{P}^4 of the ruling of M is a cone over an elliptic quartic curve which is given by the intersection of two quadric cones having l as vertex. The quartic surface is the base locus of a pencil of quadrics and, generically, Y cuts any member of the pencil, out of the base locus, along a residue quadric surface giving rise to the quoted 2-dimensional fibres of ϕ . So we have only to prove that Z has degree 13 since it is clearly of dimension 4. Lemma 3.2 yields that the inverse map is given by quintic forms through Z. Take two general quintics forms W_1, W_2 through Z and let $W_1 \cap W_2 = Z \cup Z'$. Then Z' is a 4-dimensional variety which is the image via ϕ of a general linear section of Y with a general \mathbb{P}^5 , let us say L. Let $Y' = L \cap Y \subset Y$ and let $C = L \cap X$. Then C is a smooth degree 8 and genus 3 curve in \mathbb{P}^5 and $\phi_{|Y'}$ resolves to a birational morphism $\widetilde{\sigma}: Bl_C(Y') \to Z' \subset \mathbb{P}^6$ given by the restriction to Y' of the linear system of hyperquadrics passing through C. Let H' be the pull back in $Bl_C(Y')$ of the hyperplane class of Y' and let E' be the exceptional divisor so that |2H' - E'| is the linear system giving rise to $\tilde{\sigma}$. Then we have that $\deg(Z') = (2H' - E')^4 = 16(H')^4 - 4(2H') \cdot (E')^3 + (E')^4 = 48 - 8 \deg(C) + 3 \deg(C) + 2g(C) - 2 = 48 - 64 + 28 = 12$ and the claim $\deg(Z) = 13$ follows.

References

- A. Alzati, F. Russo, Special subhomaloidal systems of quadrics and varieties with one apparent double point, Math. Proc. Camb. Phil. Soc., 134 (2003), 65-82.
- [2] A. B. Aure, K. Ranestad, The Smooth Surfaces of Degree 9 in P⁴, Complex Projective Geometry Trieste 1989/Bergen 1989 Lect. Notes in Math. Lond. Math. Soc. 179, 1992.
- [3] M. Andreatta, J. Wisniewski, On contractions of smooth varieties, J. Alg. Geometry 7 (1998), 253–312.
- [4] M. Andreatta, J. Wisniewski, Contractions of smooth varieties II. Computations and applications., Boll. Unione Mat. Ital. B (8) 1 (1998), 343–360.
- [5] A. Beauville, Determinantal hypersurfaces, Mich. Math. J. 48 (2000), 39–64.
- C. Ciliberto, M. Mella, F. Russo, Varieties with one apparent double point, Journ. of Alg. Geom. 13 (2004), 475–512.
- [7] A. Clebsch, Die Geometrie auf den Flächen dritter Ordnung, J. Reine Angew. Math. 65 (1866), 359–380.
- [8] H. Clemens, J. Kollár, S. Mori, *Higher dimensional complex Geometry*, Astérisque 166, 1988.
- [9] V. I. Danilov, Decomposition of certain birational morphisms, Math. USSR Izv. 16 (1981), 419–429.
- [10] W. Decker, L. Ein, F. O. Schreyer, Construction of surfaces in P⁴, J. of Alg. Geom. 2 (1993), 185-237.
- [11] L. Ein, N. Shepherd-Barron, Some special Cremona transformations, Amer. J. Math. 11 (1989), 783–800.
- [12] G. Fano, Sulle forme cubiche dello spazio a cinque dimensioni contenenti rigate razionali del 4^o ordine, Comm. Math. Helv. 15 (1942), 71–80.
- [13] L. Godeaux, Sur une correspondance Crémonienne entre deux espaces á n dimensions, Rend. Ist. Lombardo 43 (1910), 116–119.
- H. Grassmann, Die stereometrischen Gleichungen dritten grades, und die dadurch erzeugten Oberflächen, J. Reine Angew. Math. 49 (1855), 47–65.
- [15] R. Hartshorne, Algebraic Geometry, GTM 52, Springer-Verlag, New York, U.S.A., 1977.
- [16] K. Hulek, S. Katz, F.O. Schreyer, Cremona transformations and syzygies, Math. Z. 209 (1992), 419–443.
- [17] J. Kollár, S. Mori, Birational Geometry of Algebraic Varieties, Camb. Tracts in Math. 134, Camb. University Press, Cambridge, U. K., 1998.
- [18] S. Mori, Threefolds whose canonical bundles are not numerically effective, Ann. Math. 116 (1982), 133–176.
- [19] C. Peskine, L. Szpiro, Liason des variétés algébriques I, Inv. Math. 26 (1974), 271–302.
- [20] Z. Ran, The (dimension+2)-secant Lemma, Inv. Math. 106 (1991), 65-71.
- [21] T. G. Room, *The Geometry of Determinantal Loci*, Cambridge University Press, Cambridge, U. K., 1938.
- [22] F. Russo, On a theorem of Severi, Math. Ann. 316 (2000), 1-17.
- [23] F. Russo, A. Simis, On birational maps and Jacobian matrices, Comp. Math. 126 (2001), 335–358.
- [24] B. Segre, On the rational solutions of homogeneous cubic equations in four variables, Math. Notae 11 (1951), 1–68.
- [25] J. G. Semple, L. Roth, Introduction to Algebraic Geometry, Oxford University Press, 1949 and 1986.
- [26] J. G. Semple, J. A. Tyrrell, The Cremona transformation of S₆ by quadrics through a normal elliptic septimic scroll ¹R⁷, Mathematika 16 (1969), 88–97.
- [27] J. G. Semple, J. A. Tyrrell, The T_{2,4} of S₆ defined by a rational surface ³F⁸, Proc. Lond. Math. Soc. 20 (1970), 205–221.
- [28] J. G. Semple, J. A. Tyrrell, Specialization of Cremona transformations, Mathematika 15 (1968), 171–177.

- [29] F. Severi, Intorno ai punti doppi impropri di una superficie generale dello spazio a quattro dimensioni e ai suoi punti tripli apparenti, Rend. Circ. Mat. Palermo 15 (1901), 33–51.
- [30] J. A. Todd, The quarto-quartic transformation of four-dimensional space associated with certain projectively generated loci, Proc. Camb. Phil. Soc. 26 (1929), 323–333.
- [31] S. L. Tregub, Three constructions of rationality of cubic fourfold, Moscow Univ. Math. Bull. 39 (3) (1984), 8–16.
- [32] S. L. Tregub, Two remarks on four dimensional cubics, Russian Math. Surveys, 48 (2) (1993), 206–208.
- P. Vermeire, Some results on secant varieties leading to a geometric flip construction, Comp. Math. 125 (2001), 263–282.

DIPARTIMENTO DI MATEMATICA "F. ENRIQUES", UNIVERSITÀ DEGLI STUDI DI MILANO, VIA C. SALDINI 50, 20133 MILANO, ITALIA

 $E\text{-}mail \ address: \texttt{alzatiQmat.unimi.it}$

Departamento de Matematica, Universidade Federal de Pernambuco, Cidade Universitaria, 50670-901 Recife-PE, Brasil

E-mail address: frusso@dmat.ufpe.br