# Frobenius Rational Loop Algebra

David Chataur and Jean-Claude Thomas

November 13, 2018

#### Abstract

Recently R. Cohen and V. Godin have proved that the loop homology  $H_*(LM; \mathbf{k})$ of a closed oriented manifold M with coefficients in a field  $\mathbf{k}$  has the structure of a unital Frobenius algebra without counit. In this paper we describe explicitly the dual of the coproduct  $H_*(LM; \mathbb{Q}) \to (H_*(LM; \mathbb{Q}) \otimes H_*(LM; \mathbb{Q}))_{*-m}$  and prove it respects the Hodge decomposition.

**AMS Classification**: 55P35, 54N45,55N33, 17A65, 81T30, 17B55

**Key words** : Frobenius algebra, topological quantum field theory, free loop space, loop homology, rational homotopy, Hochschild cohomology, Hodge decomposition.

#### Introduction

Let  $\mathbb{I}_k$  be a field with a unit element denoted 1. Singular homology (respectively cohomology) of a space X with coefficients in  $\mathbb{I}_k$  is denoted  $H_*(X)$  (respectively  $H^*(X)$ ). For simplicity we identify  $H^*(X^{\times 2})$  with  $H^*(X)^{\otimes 2}$  and  $H^*(X)$  with the graded dual of the homology  $(H_*(X))^{\vee}$ .

Let M be a 1-connected closed oriented m-manifold and let  $M^{S^1}$  be the space of free loops. The space  $M^{S^1}$  can be replaced by a Hilbert manifold denoted LM (see [8] for more details) so that the evaluation map  $p_0: LM \to M$ ,  $\gamma \mapsto \gamma(0)$  is a smooth map.

Chas and Sullivan [6] have constructed a natural product, called the *loop product* 

 $P: H_*(LM \times LM) \to H_{*-m}(LM)$ 

so that  $I\!H_*(LM) := H_{*+m}(LM)$  is a commutative graded algebra. Later on Chas and Sullivan, [7], and Sullivan, [20], have introduced "higher string structures". Cohen-Godin, [9], have proved that some of these higher string structures can be defined in terms of two dimensional positive boundary TQFT (topological quantum field theory). In, particular they prove:

**Theorem CG** ([9]-Theorem 1) There exists a loop coproduct

$$\Phi: H_*(LM) \to H_{*-m}(LM \times LM)$$

such that P and  $\Phi$  define a structure of a Frobenius algebra without counit on  $I\!H_*(LM)$ .

Recall that a *Frobenius algebra* is equivalent to a two dimensional positive boundary TQFT, [9]. In particular, it is a unital associative and commutative graded algebra A together with a (non zero) coassosiative coproduct  $\Phi : A \to A \otimes A$  which is a (degree d) A-linear map. Here the coproduct  $\Phi$  is assumed to be without counit, compare with [1], [2].

It results from the very definition of the loop coproduct and of the loop product that the Frobenius algebra structure on  $H_*(LM)$  is natural with respect to any orientation preserving smooth map between connected closed oriented *m*-manifolds (see §3-Proposition 1). Roughly speaking the above theorem asserts that the functor  $H_*(-) \circ L$  carries the Poincaré duality structure on  $H_*(M)$  to a Frobenius algebra structure on  $H_*(LM)$ . Let us also notice that the orientation preserving condition is necessary because the image of the fundamental class of M in  $H_*(LM)$  is the unit of the loop product.

Our first result is :

**Theorem A** If k is of characteristic zero, the coproduct of the Frobenius algebra  $H_*(LM)$ is explicitly described in terms of Sullivan models and with the Euler class of the diagonal embedding  $\Delta: M \to M \times M$ . (See Theorem 10 in §4 for a more precise statement.)

Moreover, any orientation preserving smooth map  $\varphi$  between 1-connected closed oriented m-manifolds such that  $H^*(\varphi)$  is an isomorphism induces an isomorphism  $H_*(L\varphi)$ which respects the loop products and the loop coproducts.

The loop product has already been described in terms of Sullivan models in [15]-Theorem A. For any field of coefficients the loop product can be identified to the cup product of the Hochschild cohomology  $HH^*(C^*M; C^*M)$  when  $C^*M$  denotes the singular cochaîns on M, [10] ([18] when  $\mathbf{k} = \mathbb{R}$  and [15] when  $\mathbf{k} = \mathbb{Q}$ ). While the Hochschild cohomology of a differential graded algebra,  $HH^*(A; A)$ , is not natural with respect to graded algebra homomorphisms, we know that if  $\varphi : A \to B$  is a quasi-isomophism of differential graded algebras then there is an isomorphism of graded algebras  $HH(A; A) \to$ HH(B; B), [13]. Since the identification of  $H_*(LM)$  with  $HH^*(C^*M; C^*M)$  depends a priori of the Poincaré duality of  $H_*(M)$  it is not possible to conclude from these results that the abstract algebra structure on  $H_*(LM)$  is a homotopy invariant. (The difficulty here is to obtain a convenient chain level representative of the cap product with the orientation class, see [14]-Theorem 13 and [15]). After a preliminary version has been posted, we have learned that R. Cohen, J. Klein and D. Sullivan have announced at several summer conferences in 2003 that the string topology operations are integral homotopy invariant. (A preprint entitled "On the homotopy invariance of string topology" is in preparation.)

The tools involved in the proof of theorem A are Sullivan models [19]. In this context we naturally work with the cohomology of LM instead of homology. Therefore we are interested by the dual of the loop copoduct  $\Phi^{\vee} : H^*(LM \times LM) \to H^{*+m}(LM)$ , (respectively the dual of the loop product  $P^{\vee} : H^*(LM) \to H^{*+m}(LM \times LM)$ ) rather than the loop coproduct  $\Phi$  (respectively the loop product P).

Theorem A allows to do explicit examples. For instance we have established in § 5 the formula for  $\Phi^{\vee}$  when  $M = \mathbb{C}P^n$  and we show that the composition  $\Phi \circ \text{loop product} :$  $(\mathbb{H}(LM) \otimes \mathbb{H}(LM))_* \to (\mathbb{H}(LM) \otimes \mathbb{H}(LM))_*$  is not the trivial map.

A second application of Theorem A concerns the "Hodge decomposition of  $H^*(LM)$ ". Recall that LM is equipped with a natural  $S^1$ -action. Power maps  $\varphi_N : LM \to LM$ are then induced by the covering maps  $e^{i\theta} \mapsto e^{ni\theta}$ . The power operations  $H^*(\varphi_n) :$  $H^*(LM) \to H^*(LM)$  define, if  $\mathbf{k}$  is of characteristic zero, a natural Hodge decomposition, [5]:

$$H^*(LM) = \oplus_{i \ge 0} H^*_{(i)} , \text{ where } \begin{cases} H^*_{(0)} &= \tilde{H}^*(M) \\ H^n_{(i)} &= 0 , \text{ if } i > n \\ H^n_{(i)} &\text{ is an eingenspace of } H^n(\varphi_n) \\ &\text{ for the eigenvalue } n^i, 0 < i \le n \end{cases}$$

**Theorem B.** If k is of characteristic zero the dual of the loop coproduct  $\Phi^{\vee}$  respects the Hodge decomposition in  $H^*(LM)$ .

The paper is organized as follows;

- 1 Definition of the loop coproduct and of the loop product.
- 2 Shriek map of a k-embedding.
- 3 Sullivan representative of  $\delta_{in}$  and of  $\delta_{out}$ .
- 4 Proof of theorem A.
- 5 Example  $M = \mathbb{C}P^n$ .
- 6 Proof of theorem B.

Acknowledgments. We would like to thanks R. Cohen, Y. Felix and J. Stasheff for helpful comments.

 $\S1$  - Definition of the loop coproduct and of the loop product.

The purpose of this section is to provide us with an elementary definition of the dual of the loop coproduct and of the dual of the loop product in algebraic topology.

Following [20] and [9] the loop product (respectively the loop coproduct) is a particular case of an "operation"

$$\mu_{\Sigma}: H_*(LM)^{\otimes p} \to H_*(LM)^{\otimes q}$$

with p = 2, q = 1 (respectively p = 1, q = 2) that will define a positive boundary TQFT. Here  $\Sigma$  denotes an oriented surface of genus 0 with a fixed parameterization of the p + q boundary components:

$$(S^1)^{\coprod p} \stackrel{in}{\to} \partial_{in} \Sigma \hookrightarrow \Sigma \iff \partial_{out} \Sigma \stackrel{out}{\leftarrow} (S^1)^{\coprod q}.$$

Applying the functor Map(-, M) on gets the diagram

(1) 
$$(LM)^{\times p} \stackrel{in\#}{\leftarrow} Map(\Sigma, M) \stackrel{out\#}{\longrightarrow} (LM)^{\times q}$$
.

Diagram (1) is homotopically equivalent to the diagram

(2)  $(LM)^{\times p} \xleftarrow{\rho_{in}} Map(c, M) \xrightarrow{\rho_{out}} (LM)^{\times q}$ .

where c denotes a reduced Sullivan chord diagram with markings which is associated to  $\Sigma$  [9]-§2.

In order to define the loop coproduct we restrict to the case p = 1 and q = 2. Then  $\Sigma$  is the oriented surface of genus 0 with one incoming and two outcoming components of the boundary. The associated Sullivan chord diagram with markings is determined by the pushout diagram

$$\begin{array}{cccc} *\coprod * & \to & * \\ \downarrow & & \downarrow \\ S^1 & \stackrel{in}{\to} & c \end{array}$$

In particular,  $c \simeq S^1 \vee S^1$ ,  $S^1 \xrightarrow{in} c$  is homotopic to the folding map  $\nabla : S^1 \to S^1 \vee S^1$  and  $S^1 \coprod S^1 \xrightarrow{out} c$  is homotopic to the natural projection  $S^1 \coprod S^1 \to S^1 \vee S^1$ .

Diagram (2) then reduces to a commutative diagram of fibrations

$$(3) \begin{array}{cccc} LM^{\times 2} & \stackrel{\delta_{out}}{\longleftarrow} & LM \times_M LM & \stackrel{\delta_{in}}{\longrightarrow} & LM \\ p_0^{\times 2} \downarrow & p_0 \downarrow & & \downarrow (p_0, p_{\frac{1}{2}}) \\ M^{\times 2} & \xleftarrow{\Delta} & M & \stackrel{\Delta}{\to} & M^{\times 2} \end{array}$$

where i)  $p_t(\gamma) = \gamma(t), \ \gamma \in M^{(a,b)}, \ t \in (a,b) \subset I := (0,1)$ ii)  $\Delta(x) = (x,x)$ 

iii) The fibration  $LM \stackrel{(p_0,p_{\frac{1}{2}})}{\to} M^{\times 2}$  is defined by the pullback diagram

$$\begin{array}{cccc} LM & \stackrel{\delta'}{\to} & M^{[0,\frac{1}{2}]} \times M^{[\frac{1}{2},1]} \\ (4) & (p_0, p_{\frac{1}{2}}) \downarrow & \downarrow p_0 \times p_{\frac{1}{2}} \times p_{\frac{1}{2}} \times p_1 \\ & M^{\times 2} & \stackrel{\Delta'}{\to} & M^{\times 4} \end{array}$$

and  $\Delta'(x,y) = (x,y,y,x), x, y \in M$ 

iv) the squares are pullback diagrams of fibrations and  $\delta_{in}$ ,  $\delta_{out}$  are the natural inclusion.

Diagram (3) converts into the commutative diagram of k-linear maps

$$\begin{array}{cccc} H^*(LM^{\times 2}) & \stackrel{H^*(\delta_{out})}{\to} & H_*(LM \times_M LM) & \stackrel{\delta_{in}!}{\to} & H^*(LM) \\ H^*(p_0^{\times 2}) \uparrow & & H^*(p_0) \uparrow & & \uparrow H^*(p_0, p_{\frac{1}{2}}) \\ H^*(M^{\times 2}) & \stackrel{H^*(\Delta)}{\to} & H^*(M) & \stackrel{\Delta!}{\to} & H^*(M^{\times 2}) \end{array}$$

where  $\delta_{in}^{!}$  (respectively  $\Delta^{!}$ ) denotes the shrick map of the *m*-embedding  $\delta_{in}$  (respectively of the diagonal embedding of M) (See section 2 for a precise definition). By definition, ([9]- Definition 3), the dual of the loop coproduct is

$$\Phi^{\vee} := \delta_{in}^! \circ H^*(\delta_{out}) : (H^*(LM))^{\otimes 2} \to H^*(LM) \,.$$

To define the dual of the loop product one consider the diagram

where *Comp* denotes composition of free loops and  $\delta_{out}$  is a fibrewize embedding. Diagram (7) converts into the commutative diagram of  $\mathbf{k}$ -linear maps

$$\begin{array}{cccc} H^*(LM^{\times 2}) & \stackrel{\delta^!_{out}}{\leftarrow} & H_*(LM \times_M LM) & \stackrel{H^*(Comp)}{\leftarrow} & H^*(LM) \\ H^*(p_0^{\times 2}) \uparrow & & H^*(p_0) \uparrow & & \uparrow H^*(p_0) \\ H^*(M^{\times 2}) & \stackrel{\Delta^!}{\leftarrow} & & H^*(M) & = & H^*(M) \end{array}$$

where  $\delta_{out}$  denotes the shrink map of the *m*-embedding  $\delta_{out}$  (See section 2 for a precise definition). Following [10] or [9], the dual of the loop product is defined by:

$$P^{\vee} = \delta_{out}^! \circ H^*(Comp) : H^*(LM) \to (H^*(LM))^{\otimes 2}$$

 $\S 2$  - Shriek map of a k-embedding.

A fiber bundle pair

$$(p, p'): (F, F') \xrightarrow{i} (E, E') \xrightarrow{p} B$$

is such that F' is subspace of F and local trivializations of E' obtained by restricting local trivialization of p. If is well known (see [16]-Theorem 4D8 for example) that for

any ring of coefficients  $\boldsymbol{k}$  that if there exist cohomology classes  $c_i \in H^*(E, E'; \boldsymbol{k})$  whose restrictions  $\{H^*(i, i'; \boldsymbol{k})(c_i)$  freely generate the  $\boldsymbol{k}$ -module  $H^*(F, F'; \boldsymbol{k})$  then the  $H^*(B; \boldsymbol{k})$ module  $H^*(E, E'; \boldsymbol{k})$  is freely generated by the  $c_i$ 's. In the case of fiber bundle pair (p, p')with fiber  $(D^k, S^{k-1})$   $(D^k$  denotes the k-disk in  $\mathbb{R}^k$  and its boundary  $S^{k-1}$ ) an element  $\tau \in H^k(E, E')$  whose restriction to  $H^k(D^k, S^{k-1})$  is the fundamental class is called the Thom class of the fiber bundle pair (p, p'). If the fiber bundle pair (p, p') with fiber  $(D^k, S^{k-1})$  admits a Thom class  $\tau^{(p,p')}$  then the  $H^*(B)$ -linear map

$$H^*(p)(-) \cup \tau^{(p,p')} : H^*(B) \to H^{*+k}(E, E'), \quad x \mapsto H(p)(x) \cup \tau$$

is an isomorphism, called the Thom isomorphism, and  $H^{l}(E, E') = 0$  if l < k.

Recall, see for example [16]-Theorem 4D10), if  $\mathbf{k}$  is a field of characteristic  $\neq 2$  then every orientable fiber bundle pair (p, p') with fiber  $(D^k, S^{k-1})$  admits a Thom class. In this case the Gysin sequence of the sphere bundle p' can be deduced from the long exact of the pair (E, E') and the Thom isomorphism. In particular, one has the following commutative diagram

where  $e^{p'} = (H^*(p))^{-1} \circ j^{(E,E')}(\tau) \in H^k(B)$  is usually called the *Euler class of the sphere* bundle p'.

Let M and N be (smooth Banach and without boundary) connected manifolds and  $f: M \to N$  be a (smooth) closed embedding, [17]-II.§2. Then we have the exact sequence of fiber bundles

$$0 \to TM \xrightarrow{Tf} TN_{|_M} \to \nu_f \to 0$$

where TM and TN are the tangent bundles and  $\nu_f$  is the normal fiber bundle of f. By definition of an immersion, [17]-II Proposition 2.3, this exact sequence split. Hereafter we will identify  $\nu_f$  with a factor of  $TN_{|_M}$ .

When the fiber of  $\nu_f$  is of finite dimension k the embedding is called a k-embedding. In this case the normal fiber bundle pair  $(\nu_f^D, \nu_f^S)$  is defined and the Thom class of the oriented normal fiber bundle pair  $(\pi^D, \pi^S)$ , denoted  $\tau_f$ , is called the Thom class of the oriented embedding f.

For any embedding f, the exponential map  $D \subset TN \to N$  restricted to  $\nu_f$  is a local isomorphism on the zero cross section of the bundle  $TN \to N$ . Thus if we assume that f is a closed embedding and that N admits a partition of unity then, by [17]-IV-Theorem 5.1, there exists

a) an open neighborhood Z of the zero section of  $\nu_f$ ,

b) an open neighborhood U of f(M) in N,

c) an isomorphism  $\Theta: Z \to U$  which identifies the zero section of  $\nu_f$  with f(M).

If moreover, we assume that f is a k-embedding we identify Z with  $\nu_f^D$  and the isomorphism

$$\nu_f^D \xrightarrow{\theta} U := \operatorname{tube} f$$

restricts to an isomorphism

$$\nu_f^S \cong \theta(\nu_f^S) := \partial \text{tube } f.$$

It results from, [17]-VII-Corollary 4.2, that  $\nu_f \cong \nu_f^D$ . The above discussion is summarized in the next commutative diagram

$$\begin{array}{cccc} H^*(N, N - f(M)) & \stackrel{j^{(N, N - f(N))}}{\longrightarrow} & H^*(N) \\ \cong \downarrow & \text{Excision} & \downarrow \\ H^*(\text{tube } f, \partial \text{tube } f) & \stackrel{j^{(\text{tube } f, \partial \text{tube } f)}}{\longrightarrow} & H^*(\text{tube } f) \\ H^*(\text{tube } f, \partial \text{tube } f) & \stackrel{j^{(\mu_f^S, \nu_f^S)}}{\longrightarrow} & H^*(\text{tube } f) \\ H^*(\nu_f^D, \nu_f^S) & \stackrel{j^{(\nu_f^S, \nu_f^S)}}{\longrightarrow} & H^*(\nu_f^D) \\ - \cup \tau_f \uparrow \cong & \uparrow \pi^D \circ f_{|_{f(M)}}^{-1} \\ H^{*-k}(M) & \stackrel{- \cap e^{\pi^S}}{\longrightarrow} & H^*(M) \end{array}$$

The composition

$$TP^*: H^*(\nu_f^D, \nu_f^S) = H^*(\text{tube } f, \partial \text{tube} f) \stackrel{Excision}{\cong} H^*(N, N - f(M)) \stackrel{j^{N, N - f(M)}}{\to} H^*(N)$$

is called the cohomology Thom-Pontryagin collapse map.

The  $H^*(N)$ -linear map

$$f^{!}: H^{*}(M) \xrightarrow{H^{*}(\pi^{D} \circ f)(-) \cup \tau_{f}} H^{*+k}(\nu_{f}^{D}, \nu_{f}^{S}) \xrightarrow{TP^{*}} H^{*+k}(N)$$

is called the *cohomology shriek map* and  $f'(1) = e_f \in H^k(N)$  is called the *Euler class* of the embedding f.

Proposition 1. Let us consider the commutative diagram

$$\begin{array}{cccc} M & \stackrel{f}{\to} & N \\ \varphi \downarrow & & \downarrow \psi \\ M' & \stackrel{g}{\to} & N' \end{array}$$

where

 $\left\{ \begin{array}{l} M\,,N\,,M'\,,N' \ are \ smooth \ Hilbert \ manifolds \ (without \ boundary) \\ f\,,g\,, \ are \ two \ finite \ codimensionnal \ closed \ oriented \ embeddings \\ \varphi\,,\psi \ are \ two \ smooth \ maps \ which \ preserve \ orientations \\ of \ the \ associated \ normal \ pairs \end{array} \right.$ 

Then we have the following commutative diagram

$$\begin{array}{ccc} H^*(M) & \xrightarrow{f^{:}} & H^{*+k}(N) \\ H^*(\varphi) \uparrow & & \uparrow H^*(\psi) \\ H^*(M') & \xrightarrow{g^{!}} & H^{*+k}(N') \end{array}$$

**Proof.** The maps  $\varphi$  and  $\psi$  preserve orientations of the normal pairs implies that they have the same rank and that the Euler classes of the embeddings are preserved. By, [17]-IV-Theorem 6.2, there exists a vector bundle isomorphism  $\lambda$  between  $\nu_f^D$  and the pullback  $\varphi^*\nu_g^D$  of  $\nu_g^D$  via  $\varphi$  and an isotopy  $h: \nu_f^D \times \mathbb{R} \to \varphi^*\nu_g^D$  such that h(-,0) = f and  $h(-,1) = g \circ \lambda$ . Therefore the pair  $(\psi, \varphi)$  induces a well defined homomorphism

$$H^*(\operatorname{tube} g, \partial \operatorname{tube} g) \to H^*(\operatorname{tube} f, \partial \operatorname{tube} f)$$

The remaining of the proof is now straightforward.

We have the very useful result

**Proposition 2.** Let M and N be (smooth Hilbert and without boundary) connected manifolds and  $f: M \to N$  be a closed oriented k-embedding with Euler class  $e_f \in H^k(N)$ . The composite

$$H^*(N) \xrightarrow{H^*(f)} H^*(M) \xrightarrow{f!} H^{*+k}(N)$$

is the right multiplication by  $e_f$ . Moreover, the restriction of  $f^!$  to the image of  $H^*(f)$  is well defined by the formula:

$$f^! \circ H^*(f) = - \cup e_f.$$

In particular, if  $H_*(f)$  is an isomorphism then  $f! = (H^*(f))^{-1}$ .

**Proof.** Observe that, by construction,  $f^{!}$  is a  $H^{*}(N)$ -module homomorphism (here  $H^{*}(M)$  is a  $H^{*}(N)$ -module via  $H^{*}(f)$ ). Then for any  $x \in H^{l}(N)$  and any  $y \in H^{*}(M)$ 

$$f^!(H^*(f)(x) \cup y) = x \cup f^!(y).$$

If  $\alpha = H^*(f)(\beta)$ , for some  $\beta \in H^*(N)$  and if 1 denotes the unit in  $H^0(M)$  then

$$f^{!}(\alpha) = f^{!}(H^{*}(f)(\beta) \cup 1) = \beta \cup f^{!}(1) = \beta \cup e_{f}.$$

Since  $f^!$  is well defined the right-hand part of the above formula is independent of the choice of  $\beta$ . This can be also established in the following way. From diagram (6) one deduces the commutative diagram

$$\begin{array}{cccc} H^*(\nu_f^D,\nu_f^S) & \stackrel{\cong}{\leftarrow} & H^*(N,N-f(M)) & \stackrel{j^{N,N-f(M)}}{\to} & H^*(N) \\ & -\cup\tau_f \uparrow & & \uparrow -\cup\tau_f^{N,N-f(M)} \\ H^{*-k}(\nu_f^D) = H^{*-k}(M) & \stackrel{H^*(f)}{\leftarrow} & H^{*-k}(N) \end{array}$$

where  $\tau_f^{N,N-f(M)}$  denotes the image of the Thom class  $\tau_f$  via the isomorphism  $H^*(\nu_f^D, \nu_f^S) \cong H^*(N, N - f(M))$ . Now let  $\beta, \beta' \in H^*(N)$  be such that  $H^*(f)(\beta) = \alpha = H^*(f)(\beta')$ . Thus  $\beta - \beta' \in \ker H^*(f)$  and then  $(\beta - \beta') \cup \tau_f^{N,N-f(M)} = 0$ . Therefore  $(\beta - \beta') \cup e_f = 0$ .

Let M be a simply connected compact and oriented m-manifold. A slight modification in the proof of Lemma 5.4.1 in [4] shows that the smooth map  $p_0: LM \to M$  is a fiber bundle. Thus each fibration apparing in the two pullback diagrams

$$(3) \begin{array}{cccc} LM^{\times 2} & \stackrel{\delta_{out}}{\longleftarrow} & LM \times_M LM & \stackrel{\delta_{in}}{\longrightarrow} & LM \\ p_0 \times p_0 \downarrow & p_0 \downarrow & & \downarrow (p_0, p_{\frac{1}{2}}) \\ M^{\times 2} & \stackrel{\Delta}{\leftarrow} & M & \stackrel{\Delta}{\rightarrow} & M^{\times 2} \end{array}$$

are smooth fiber bundles.

**Theorem 3** ([10]-§1 and [9]-§2)  $\delta_{in}$  and  $\delta_{out}$  are closed m-embeddings.

It results from Proposition 1 that we have the following commutative diagram

(7) 
$$\begin{array}{cccc} H^{*+k}(LM)^{\otimes 2} & \stackrel{\delta^{l}_{out}}{\longleftarrow} & H^{*}(LM \times_{M} LM) & \stackrel{\delta^{l}_{in}}{\longrightarrow} & H^{*+k}(LM) \\ H^{*}(p_{0})^{\otimes 2} \uparrow & H^{*}(p_{0}) \uparrow & \uparrow H^{*}(p_{0}, p_{\frac{1}{2}}) \\ H^{*+k}(M)^{\otimes 2} & \stackrel{\Delta^{l}}{\longleftarrow} & H^{*}(M) & \stackrel{\Delta^{l}}{\longrightarrow} & H^{*+k}(M)^{\otimes 2} \end{array}$$

In particular

$$e_{\delta_{in}} = H^*(p_0, p_{\frac{1}{2}})(e_{\Delta}) \in H^k(LM) \text{ and } e_{\delta_{out}} = H^*(p_0, p_1)(e_{\Delta}) \in H^k(LM \times LM)$$

where  $e_{\Delta}$  denotes the Euler class of the diagonal embedding of M.

Moreover, by Proposition 1, diagram (7) is natural with respect to orientation preserving smooth maps between *m*-dimensional simply connected compact and oriented manifolds.

### §3 - Sullivan representative of $\delta_{in}$ and of $\delta_{out}$ .

In this section  $\mathbf{k}$  is a field of characteristic zero. We refer systematically the reader to [12] for notation, terminology and needed results. Nonetheless, we precise here some notation and recall some relevant results. If  $V = \{V^i\}_{i \in \geq 0}$  is a graded  $\mathbf{k}$ -vector space  $\overline{V}$ denotes the suspension of V,  $(\overline{V})^n = V^{n+1}$ . We denote by  $\bigwedge V$  the free graded commutative algebra generated by V. A Sullivan algebra is a cochain algebra of the form  $(\bigwedge V, d)$ . It is minimal if  $d(V) \subset \bigwedge^{\geq 2}(V)$ .

Any path connected space X admits a Sullivan model,

$$\mathfrak{M}_X := (\bigwedge V, d) \xrightarrow{\simeq} A_{PL}(X)$$

where  $A_{PL}$  denotes the contravariant functor of piece linear differential forms, [12]-§10 and §12. If X is 1-connected  $\mathfrak{M}_X$  can be chosen minimal and in this case it is unique up to isomorphism.

Let X and Y be two path-connected spaces. Any continuous map  $f: X \to Y$  admits a Sullivan representative  $\mathfrak{M}_f: \mathfrak{M}_Y \to \mathfrak{M}_X$ : there exists a diagram

$$\begin{array}{cccc} A_{PL}(Y) & \stackrel{A_{PL}(f)}{\longrightarrow} & A_{PL}(X) \\ \rho_Y \uparrow \simeq & \simeq \uparrow \rho_X \\ \mathfrak{M}_Y & \stackrel{\mathfrak{M}_f}{\longrightarrow} & \mathfrak{M}_X \end{array}$$

such that  $\rho_X \circ \mathfrak{M}_f$  and  $A_{PL}(f) \circ \rho_Y$  are homotopic homomorphisms of commutative differential graded algebras. Moreover, if  $f, g: X \to Y$  are homotopic then  $\mathfrak{M}_f, \mathfrak{M}_g: \mathfrak{M}_Y \to \mathfrak{M}_X$ are homotopic.

Here after we will make the following identifications:

$$H^{*}(X) = H(A_{PL}(X)) = H(\mathfrak{M}_{X}), \ H^{*}(f) = H(A_{PL}(f)) = H(\mathfrak{M}_{f}).$$

It is convenient to consider "cofibrant" Sullivan representative of f called *relative* Sullivan model of f. The map  $f: X \to Y$  admits also a relative Sullivan model, [12]-§14:

$$\begin{array}{cccc} A_{PL}(Y) & \stackrel{A_{PL}(f)}{\longrightarrow} & A_{PL}(X) \\ \rho_{Y} \uparrow \simeq & \simeq \uparrow \rho_{X} \\ \mathfrak{M}_{Y} & \stackrel{\lambda_{f}}{\longrightarrow} & \mathfrak{M}_{X} = (\mathfrak{M}_{Y} \otimes \bigwedge W, d) \end{array}$$

where  $\lambda_f$  denotes the natural inclusion of  $\mathfrak{M}_Y$  into  $\mathfrak{M}_Y \otimes \bigwedge W$  and d is such that  $de_i \in \mathfrak{M}_Y \otimes \bigwedge (e_1, e_2, \dots, e_{i-1})$  for some linear basis  $e_1, e_2, \dots, e_n, \dots$  of W.

It important to observe here that :

- 1) the relative Sullivan model of f is unique up to isomorphism [12]-Theorem 14.12.
- 2) all the rational homotopy informations on f are hidden in the differential d.

**Theorem 4.** Let M be a 1-connected compact oriented manifold and let  $\mathfrak{M}_M = (\bigwedge V, d)$ be a minimal model of M whose product is denoted by  $\mu : \bigwedge V \otimes \bigwedge V \to \bigwedge V$ . a) A Sullivan representative of  $\delta_{out}$  is explicitly given by:

b) A Sullivan representative of  $\delta_{in}$  is explicitly given by:

$$\begin{array}{cccc} ((\bigwedge V)^{\otimes 2} \otimes (\bigwedge \bar{V})^{\otimes 2}, d') & \stackrel{\mu \otimes id \otimes id}{\longrightarrow} & (\bigwedge V \otimes (\bigwedge \bar{V})^{\otimes 2}, d) \\ & & & & & \\ & & & & & \\ & & & & & \\ \mathfrak{M}'_{LM} & \stackrel{\mathfrak{M}_{\delta_{in}}}{\longrightarrow} & \mathfrak{M}_{LM \times_M LM} \end{array}$$

The differentials appearing in the above statement will be precisely described during the proof.

Observe here that while the homomorphisms of graded algebras  $\mathfrak{M}_{\delta_{in}}$  and  $\mathfrak{M}_{\delta_{out}}$  are equal the homomorphism  $H(\mathfrak{M}_{\delta_{in}})$  and  $H(\mathfrak{M}_{\delta_{out}})$  are different. The remaining of this section is devoted to the proof of Theorem 4. The proof follows from a sequence of lemmas which rely on the "preservation under pushout" property of the relative Sullivan models:

**Lemma 5.** [12]-Proposition 15.8 A pullback diagram in the category of topological spaces where p is a fibration :

$$\begin{array}{ccccc}
f^*E & \stackrel{f'}{\to} & E \\
p' \downarrow & & \downarrow p \\
B' & \stackrel{f}{\to} & B
\end{array}$$

converts into the pushout diagram in the category of commutative differential graded algebras

$$\mathfrak{M}_{B} \xrightarrow{\mathfrak{M}_{f}} \mathfrak{M}_{B'} \xrightarrow{\mathfrak{M}_{f}} \mathfrak{M}_{B'} \\
\lambda_{p} \downarrow \qquad \qquad \downarrow \lambda_{p'} \\
\mathfrak{M}_{E} = (\mathfrak{M}_{B} \otimes \bigwedge W, d) \xrightarrow{\mathfrak{M}_{f} \otimes id} (\mathfrak{M}_{B'} \otimes \bigwedge W, d') = \mathfrak{M}_{E'}$$

Let us just precise here that the differential  $d' = \varphi \circ d \circ \varphi^{-1}$  where  $\varphi$  is the canonical isomorphism of commutative graded algebras such that the next diagram commutes

$$\begin{array}{cccc} \mathfrak{M}_{B'}\otimes_{\mathfrak{M}_{B}}(\mathfrak{M}_{B}\otimes\wedge W) & \xrightarrow{\varphi} & \mathfrak{M}_{B'}\otimes\wedge W & a\otimes b\otimes c\mapsto a\mathfrak{M}_{f}(b)\otimes c\\ \mathfrak{M}_{f}\otimes id\otimes id\uparrow & \uparrow & \mathfrak{M}_{f}\otimes id\\ (\mathfrak{M}_{B}\otimes_{\mathfrak{M}_{B}}\mathfrak{M}_{B})\otimes\wedge W & \xrightarrow{\mu\otimes id} & \mathfrak{M}_{B}\otimes\wedge W & a\otimes b\otimes c\mapsto ab\otimes c \end{array}$$

**Lemma 6.** [12]-§15-Example 1 The relative Sullivan model of the fibration  $p_0 \times p_1$ :  $M^I \to M \times M$  is of the form

$$\mathfrak{M}_{M}^{\otimes 2} = (\bigwedge V, d)^{\otimes 2} \stackrel{\lambda_{p_{0} \times p_{1}}}{\hookrightarrow} ((\bigwedge V)^{\otimes 2} \otimes \bigwedge \bar{V}, d) = \mathfrak{M}_{M^{I}}.$$

Here the differential is precisely defined as follows.

 $\begin{cases} d(v \otimes 1 \otimes 1) &= dv \otimes 1 \otimes \bar{1} \\ d(1 \otimes v \otimes \bar{1}) &= 1 \otimes dv \otimes \bar{1} \\ d(1 \otimes 1 \otimes \bar{v}) &= (v \otimes 1 - 1 \otimes v) \otimes \bar{1} - \sum_{i=1}^{\infty} \frac{(sd)^i}{i!} (v \otimes 1 \otimes \bar{1}) \end{cases}, \quad v \in V, \bar{v} \in \bar{V} \text{ and } s \\ \text{being the unique degree -1 derivation of } \bigwedge V \otimes \bigwedge V \otimes \bigwedge \bar{V} \text{ defined by} \end{cases}$ 

$$\begin{cases} s(v \otimes 1 \otimes \bar{1}) = 1 \otimes 1 \otimes \bar{v} = s(1 \otimes v \otimes \bar{1}) \\ s(1 \otimes 1 \otimes \bar{v}) = 0 \end{cases}, \quad v \in V, \bar{v} \in \bar{V}$$

We denote by 1 (respectively  $\overline{1}$ ) the unit of  $\bigwedge V$  (respectively  $\bigwedge \overline{V}$ ).

**Lemma 7.** The relative Sullivan model for  $p_0 : LM \to M$  is  $\lambda_{p_0} : \mathfrak{M}_M = (\bigwedge V, d) \hookrightarrow (\bigwedge V \otimes \bigwedge \overline{V}, d) = \mathfrak{M}_{LM}$ .

**Proof** Since multiplication  $\mu : (\bigwedge V, d)^{\otimes 2} \to (\bigwedge V, d)$  is a Sullivan representative of  $\Delta$ , by Lemma 6 and Lemma 7 the pullback diagram

$$\begin{array}{rcl} LM & \rightarrow & M^{I} \\ p_{0} \downarrow & & \downarrow p_{0} \times p_{1} \\ M & \stackrel{\Delta}{\rightarrow} & M^{\times 2} \end{array}$$

yields the pushout diagram in the category of commutative differential graded algebras

$$\begin{array}{cccc} (\bigwedge V, d)^{\otimes 2} & \xrightarrow{\mu} & (\bigwedge V, d) \\ \lambda_{p_0 \times p_1} \downarrow & & \downarrow \lambda_{p_0} \\ ((\bigwedge V)^{\otimes 2} \otimes \bigwedge \bar{V}, d) & \xrightarrow{\mu \otimes id} & (\bigwedge V \otimes \bigwedge \bar{V}, \bar{d}) \end{array}$$

The relative Sullivan model for  $p_0: LM \to M$  is  $\lambda_{p_0}$  and  $\begin{cases} d(1 \otimes v) &= dv \otimes \bar{1} \\ d(1 \otimes \bar{v}) &= -s(dv \otimes \bar{1}) \end{cases}$  with s being the unique degree -1 derivation of  $\bigwedge V \otimes \bigwedge \bar{V}$  defined by  $\begin{cases} s(v \otimes \bar{1}) &= 1 \otimes \bar{v} \\ s(1 \otimes \bar{v}) &= 0 \end{cases}$ ,  $v \in V, \bar{v} \in \bar{V}.$ 

**Proof of part a) of theorem 4.** By Lemma 5 and Lemma 7 the pullback diagram of fibrations

$$\begin{array}{cccc} LM \times_M LM & \stackrel{o_{out}}{\longrightarrow} & LM \times LM \\ p_0 \downarrow & & \downarrow p_0 \times p_0 \\ M & \stackrel{\Delta}{\rightarrow} & M^{\times 2} \end{array}$$

converts into the pushout diagram

where

$$\begin{cases} d(v \otimes \overline{1} \otimes \overline{1}) &= dv \otimes \overline{1} \otimes \overline{1} \\ d(1 \otimes \overline{v} \otimes \overline{1}) &= -s(dv \otimes \overline{1} \otimes \overline{1}) \\ d(1 \otimes \overline{1} \otimes \overline{v}) &= -s'(dv \otimes \overline{1} \otimes \overline{1}) \end{cases}, v \in V, \overline{v} \in \overline{V}.$$

Here s and s' are the unique degree -1 derivation of  $\bigwedge V \otimes (\bigwedge \overline{V})^{\otimes 2}$  defined by

$$\begin{cases} s(v \otimes \overline{1} \otimes \overline{1}) &= 1 \otimes \overline{v} \otimes \overline{1} \\ s'(v \otimes \overline{1} \otimes \overline{1}) &= 1 \otimes \overline{1} \otimes \overline{v} \\ s(1 \otimes \overline{v} \otimes \overline{1}) &= 0 = s'(1 \otimes \overline{v} \otimes \overline{1}) \\ s(1 \otimes \overline{1} \otimes \overline{v}) &= 0 = s'(1 \otimes \overline{1} \otimes \overline{v}) \end{cases}, \qquad v \in V, \overline{v} \in \overline{V}.$$

Thus  $\mathfrak{M}_{\delta_{out}} = \mu \otimes id \otimes id.$ 

**Lemma 8.** A relative Sullivan model for  $(p_0, p_{\frac{1}{2}}) : LM \to M \times M$  is

$$\lambda_{p_0 \times p_{\frac{1}{2}}} : (\bigwedge V, d)^{\otimes 2} \hookrightarrow ((\bigwedge V)^{\otimes 2} \otimes (\bigwedge \bar{V})^{\otimes 2}, d') = \mathfrak{M}'_{LM}.$$

Proof. By Lemma 6 and Lemma 7 the pullback diagram

(4) 
$$\begin{array}{ccc} LM & \stackrel{\delta'}{\to} & M^{[0,\frac{1}{2}]} \times M^{[\frac{1}{2},1]} \\ (p_0, p_{\frac{1}{2}}) \downarrow & \qquad \downarrow p_0 \times p_{\frac{1}{2}} \times p_{\frac{1}{2}} \times p_1 \\ M^{\times 2} & \stackrel{\Delta'}{\to} & M^{\times 4} \end{array}$$

converts into the pushout diagram

$$\begin{array}{cccc} (\bigwedge V, d)^{\otimes 4} & \stackrel{\mathfrak{M}_{\Lambda'}}{\to} & (\bigwedge V, d)^{\otimes 2} \\ \lambda_{p_0 \times p_{\frac{1}{2}}}^{\otimes 2} \downarrow & & \downarrow \lambda_{p_0 \times p_{\frac{1}{2}}} \\ (\bigwedge V \otimes \bigwedge V \otimes \bigwedge \bar{V}, d)^{\otimes 2} & \stackrel{\mathfrak{M}_{\delta'}}{\to} & ((\bigwedge V)^{\otimes 2} \otimes (\bigwedge \bar{V})^{\otimes 2}, d') \end{array}$$

where  $\mathfrak{M}_{\Delta'}(a \otimes b \otimes a' \otimes b') = (-1)^{|b'|(|a'|+|b|)}ab' \otimes ba'$ , with  $\lambda_{p_0 \times p_{\frac{1}{2}}}$  defined by

$$\begin{cases} v \otimes 1 \mapsto v \otimes 1 \otimes \overline{1} \otimes \overline{1} \otimes \overline{1} \\ 1 \otimes v \mapsto 1 \otimes v \otimes \overline{1} \otimes \overline{1} \otimes \overline{1} \end{cases}, \text{ and } d' \text{ defined by} \\ \begin{cases} d'(v \otimes 1 \otimes \overline{1} \otimes \overline{1}) &= dv \otimes 1 \otimes \overline{1} \otimes \overline{1} \\ d'(1 \otimes v \otimes \overline{1} \otimes \overline{1}) &= 1 \otimes dv \otimes \overline{1} \otimes \overline{1} \\ d'(1 \otimes 1 \otimes \overline{v} \otimes \overline{1}) &= (v \otimes 1 - 1 \otimes v) \otimes \overline{1} \otimes \overline{1} - \sum_{n \ge 1} \frac{(sd')^n}{n!} (v \otimes 1 \otimes \overline{1} \otimes \overline{1}) \\ d'(1 \otimes 1 \otimes \overline{1} \otimes \overline{v}) &= (1 \otimes v - v \otimes 1) \otimes \overline{1} \otimes \overline{1} - \sum_{n \ge 1} \frac{(sd')^n}{n!} (1 \otimes v \otimes \overline{1} \otimes \overline{1}) \end{cases}$$

where s' denotes the unique degree -1 derivation of  $(\bigwedge V)^{\otimes 2} \otimes (\bigwedge \bar{V})^{\otimes 2}$  defined by

$$\begin{cases} s'(v \otimes 1 \otimes \overline{1} \otimes \overline{1}) &= 1 \otimes 1 \otimes \overline{v} \otimes \overline{1} \\ s'(1 \otimes v \otimes \overline{1} \otimes \overline{1}) &= 1 \otimes 1 \otimes \overline{1} \otimes \overline{v} \\ s'(1 \otimes 1 \otimes \overline{v} \otimes \overline{1}) &= 0 \\ s'(1 \otimes 1 \otimes \overline{1} \otimes \overline{v}) &= 0 \end{cases}, \quad v \in V, \overline{v} \in \overline{V}.$$

**Lemma 9.** Let  $\mu$  (respectively  $\bar{\mu}$ ) be the product of  $\bigwedge V$  (respectively of  $\bigwedge \bar{V}$ ). The surjective homomorphism of differential graded algebras

$$\mu \otimes \bar{\mu} : \mathfrak{M}'_{LM} = ((\bigwedge V)^{\otimes 2} \otimes (\bigwedge \bar{V})^{\otimes 2}, d') \to (\bigwedge V \otimes \bigwedge \bar{V}, d) = \mathfrak{M}_{LM},.$$

is a quasi-isomorphism.

$$\mathbf{Proof.} \text{ Formulae} \begin{cases} v \otimes 1 \otimes \overline{1} \otimes \overline{1} \mapsto (v \otimes 1 + 1 \otimes v) \otimes \overline{1} \otimes \overline{1} \\ 1 \otimes v \otimes \overline{1} \otimes \overline{1} \mapsto (v \otimes 1 - 1 \otimes v) \otimes \overline{1} \otimes \overline{1} \\ 1 \otimes 1 \otimes \overline{v} \otimes \overline{1} \mapsto 1 \otimes 1 \otimes (\overline{v} \otimes 1 + 1 \otimes \overline{v}) \\ 1 \otimes 1 \otimes \overline{1} \otimes \overline{v} \mapsto 1 \otimes 1 \otimes (\overline{v} \otimes 1 - 1 \otimes \overline{v}) \end{cases}, \qquad v \in V, \overline{v} \in \overline{V} \text{ define}$$

an automorphism of commutative graded algebras of  $(\bigwedge V)^{\otimes 2} \otimes (\bigwedge \overline{V})^{\otimes 2}$ . This automorphism exhibits the differential graded algebra  $((\bigwedge V)^{\otimes 2} \otimes (\bigwedge \overline{V})^{\otimes 2}, d')$  as the tensor product of  $\mathfrak{M}_{LM}$  with an acyclic commutative differential graded algebra. Therefore the surjective homomorphism of differential graded algebras

$$\rho:\mathfrak{M}'_{LM}=((\bigwedge V)^{\otimes 2}\otimes (\bigwedge \bar{V})^{\otimes 2},d')\to (\bigwedge V\otimes \bigwedge \bar{V},d)=\mathfrak{M}_{LM},.$$

uniquely defined by formulae 
$$\begin{cases} v \otimes 1 \otimes \bar{1} \otimes \bar{1} \mapsto v \otimes \bar{1} \\ 1 \otimes v \otimes \bar{1} \otimes \bar{1} \mapsto v \otimes \bar{1} \\ 1 \otimes 1 \otimes \bar{v} \otimes \bar{1} \mapsto 1 \otimes \bar{v} \\ 1 \otimes 1 \otimes \bar{1} \otimes \bar{v} \mapsto 1 \otimes \bar{v} \end{cases}, \quad v \in V, \bar{v} \in \bar{V} \text{ is a quasi-}$$
isomorphism. By construction  $\rho = \mu \otimes \bar{\mu}.$ 

Proof of part b) of theorem 4. By Lemma 5 and Lemma 8 the pullback diagram of fibrations

$$\begin{array}{cccc} LM \times_M LM & \stackrel{o_{i_{\mathcal{D}}}}{\to} & & LM \\ p_0 \downarrow & & \downarrow p_0 \times p_{\frac{1}{2}} \\ M & \stackrel{\Delta}{\to} & M^{\times 2} \end{array}$$

converts into the pushout diagram

Thus  $\mathfrak{M}_{\delta_{in}} = \mu \otimes id \otimes id$ 

## $\S 4$ - Proof of Theorem A.

In this section  $\boldsymbol{k}$  is a field of characteristic zero and we use notation introduced in section  $\S4$ .

First recall, [3]-VI-Theorem 12.4, that the Euler class of the diagonal embedding  $\Delta$ :  $M \to M \times M$  (also called the diagonal class) is the cohomology class

$$e_{\Delta} = \sum_{l} (-1)^{|\beta_j|} \hat{\beta}_l \otimes \beta_l \in H^m(M \times M) = (H^*(M) \otimes H^*(M))^m$$

where  $\{\beta_l\}$  denotes a homogeneous linear basis of  $H^*(M)$  and  $\{\hat{\beta}_l\}$  its Poincaré dual basis.

Let  $b_l$  (respectively  $\hat{b}_l$ ) be a cocycle of  $(\bigwedge V, d)$  representing  $\beta_l \in H^*(\bigwedge V, d) = H^*(M)$ (respectively  $\hat{\beta}_l \in H^*(\bigwedge V, d) = H^*(M)$ ). Since,  $e_\Delta = [\sum_l (-1)^{|b_l|} \hat{b}_l \otimes b_l] \in H^m(M \times M)$  it results from Theorem 3 and the definition of  $\lambda_{p_0,p_{\frac{1}{4}}}$  given in lemma 9, that

$$e_{\delta_{in}} = [\lambda_{p_0 \times p_{\frac{1}{2}}} (\sum_l (-1)^{|b_l|} \hat{b}_l \otimes b_l)] = [\sum_l (-1)^{|b_l|} \hat{b}_l \otimes b_l \otimes \bar{1} \otimes \bar{1}] \in H^m L M \,.$$

We consider the **degree m linear map** 

$$\begin{aligned} \mathfrak{M}_{LM\times_{M}LM} &= (\bigwedge V \otimes (\bigwedge \bar{V})^{\otimes 2}, d) \xrightarrow{\mathfrak{M}_{\delta_{in}}^{!}} ((\bigwedge V)^{\otimes 2} \otimes (\bigwedge V)^{\otimes 2}, d') = \mathfrak{M}_{LM}' \\ (a \otimes \bar{x} \otimes \bar{y}) \mapsto & \left(\frac{1}{2}(a \otimes 1 + 1 \otimes a) \otimes \bar{x} \otimes \bar{y}\right) \sum_{l} (-1)^{|b_{l}|} \hat{b}_{l} \otimes b_{l} \otimes \bar{1} \otimes \bar{1} \end{aligned}$$

It is straightforward to check that  $d' \circ \mathfrak{M}^!_{\delta_{in}} = (-1)^m \mathfrak{M}^!_{\delta_{in}} \circ d.$ 

Now we can state precisely the first part of theorem A.

**Theorem 10.** With notation introduced above, the composition

$$\mathfrak{M}_{LM}^{\otimes 2} \cong \mathfrak{M}_{LM \times LM} \xrightarrow{\mathfrak{M}_{\delta_{i_n}}^! \circ \mathfrak{M}_{\delta_{out}}} \mathfrak{M}'_{LM} \xrightarrow{\mu \otimes \bar{\mu}} \mathfrak{M}_{LM} ,$$

where  $\mu \otimes \bar{\mu}$  is considered in Lemma 10, induces the dual of the loop coproduct  $\Phi^{\vee}$ :  $H^*(LM) \otimes H^*(LM) \to H^{*+m}(LM).$ 

**Proof.** Let  $\phi : \mathfrak{M}_{LM}^{\otimes 2} \cong \mathfrak{M}_{LM \times LM} \longrightarrow \mathfrak{M}'_{LM}$  be a degree *m* **k**-linear map such that  $H(\phi) = \delta_{in}^!$ . Then by, Proposition 2,  $H(\phi \circ \mathfrak{M}_{\delta_{in}}) = - \cup e_{\delta_{in}}$ . On the other hand, since  $H(\phi \circ \mathfrak{M}_{\delta_{out}}) = \Phi^{\vee}$ , in order to prove theorem 10 it suffices to prove that

$$\operatorname{im}(H^*(\delta_{out})) \subset \operatorname{im}(H^*(\delta_{in}))$$

To prove this inclusion, we consider the commutative diagram

$$\begin{array}{cccc} \mathfrak{M}_{M}^{\otimes 2} & \stackrel{\lambda_{0}, \frac{1}{2}}{\to} & \mathfrak{M}_{LM}' \\ \lambda_{0}^{\otimes 2} \downarrow & & \simeq \downarrow \mu \otimes \bar{\mu} \\ \mathfrak{M}_{LM}^{\otimes 2} & \stackrel{\hat{\mu}}{\to} & \mathfrak{M}_{LM} \end{array}$$

where  $\hat{\mu}$  is the product in the graded algebra  $\mathfrak{M}_{LM}$ . By Lemma 9,  $\mu \otimes \bar{\mu}$  is a surjective quasi-isomorphism, thus Lemma 14.4 in [12] implies that there exists  $\theta : \mathfrak{M}_{LM}^{\otimes 2} \to \mathfrak{M}'_{LM}$  such that  $\theta \circ \lambda_0^{\otimes 2} = \lambda_{0,\frac{1}{2}}$  and  $\mu \otimes \bar{\mu} \circ \theta = \hat{\mu}$ . Therefore if  $\alpha$  is a cocycle in  $\mathfrak{M}_{LM}^{\otimes 2}$  then  $\theta(\alpha)$  is a cocycle of  $\mathfrak{M}'_{LM}$  and  $\mathfrak{M}_{\delta_{out}}(\alpha) = \mathfrak{M}_{\delta_{in}}(\theta(\alpha))$ .

### End of the proof of Theorem A.

The naturality assumptions in Theorem A for the loop coproduct follows from those in Propositions 1 and the fact that two Sullivan representatives  $\varphi$  and  $\psi$  of a map  $f: X \to Y$  are identified, at the homology level via isomorphisms :

$$\begin{array}{cccc} H(\mathfrak{M}_Y) & \stackrel{H(\varphi)}{\to} & H(\mathfrak{M}_X) \\ \| & & \| \\ H(\mathfrak{M}_Y) & \stackrel{H(\psi)}{\to} & H(\mathfrak{M}_X) \end{array}$$

The naturality for the loop product follows, in the same way, directly from the construction performed in [15]-3.5.

§5 - EXAMPLE 
$$M = \mathbb{C}P^n$$
.

One know that  $\mathfrak{M}_{\mathbb{C}P^n} = (\bigwedge(x, y), d)$  with |x| = 2,  $dy = x^{n+1}$  and that  $H^*(\mathbb{C}P^m) = \bigwedge x/(x^{m+1})$ . Moreover there is a quasi-isomorphism  $(\bigwedge(x, y), d) \xrightarrow{\simeq} (\bigwedge x/(x^n + 1), 0) = \overline{\mathfrak{M}}_{\mathbb{C}P^n}$ . For simplicity we will replace  $\mathfrak{M}_{\mathbb{C}P^n}$  by  $\overline{\mathfrak{M}}_{\mathbb{C}P^n}$  so that we obtain:

$$\begin{split} \bar{\mathfrak{M}}_{LM} &= (\bigwedge x/(x^n+1) \otimes \bigwedge(\bar{x},\bar{y}),\bar{d}) \\ \bar{\mathfrak{M}}_{LM \times_M LM} &= (\bigwedge x/(x^n+1) \otimes \bigwedge(\bar{x},\bar{y}) \otimes \bigwedge(\bar{x},\bar{y}),\bar{d}) \\ \bar{\mathfrak{M}}'_{LM} &= (\bigwedge x/(x^n+1) \otimes \bigwedge(\bar{x},\bar{y}) \otimes \bigwedge(\bar{x},\bar{y}) \otimes \bigwedge(\bar{x},\bar{y}),\bar{d}') \\ \bar{\mathfrak{M}}'_{LM} &= (\bigwedge x/(x^n+1) \otimes \bigwedge x/(x^n+1) \otimes \bigwedge(\bar{x},\bar{y}) \otimes \bigwedge(\bar{x},\bar{y}),\bar{d}') \\ \bar{\mathfrak{M}}^{!}_{\delta_{in}}(x^k \otimes \bar{u} \otimes \bar{v}) &= \frac{1}{2} \left( \sum_{i=0}^{n} x^i + k \otimes x^{n-i} + x^i \otimes x^{n-i+k} \right) \otimes \bar{u} \otimes \bar{v} \\ \bar{\mathfrak{M}}_{\delta_{out}}(x^k \otimes x^l \otimes \bar{u} \otimes \bar{v}) &= x^{k+l} \otimes \bar{u} \otimes \bar{v} \\ \rho \circ \bar{\mathfrak{M}}^{!}_{\delta_{in}} \circ \bar{\mathfrak{M}}_{\delta_{out}}(x^k \otimes x^l \otimes \bar{x}^{\epsilon} \bar{y}^s \otimes \bar{x}^{\epsilon'} \bar{y}^{s'}) &= \begin{cases} 0 & \text{if } k+l > 0 \\ x^n \otimes \bar{x}^{\epsilon+\epsilon'} \bar{y}^{s+s'} & \text{if } k=l=0 \,. \end{cases}$$

Where  $\epsilon$  and  $\epsilon'$  is 0 or 1. Since,

 $\{1, [x \otimes \bar{y}^{s}], [x^{2} \otimes \bar{y}^{s}], ..., [x^{n} \otimes \bar{y}^{s}], [x \otimes \bar{x}\bar{y}^{s}], ..., [x^{n-1} \otimes \bar{x}\bar{y}^{s}]\}_{s \ge 0}$ 

is a linear basis of  $H^*(LM)$  we have obtained an explicit formula for  $\Phi^{\vee}$  in this basis.

Let us denote by  $\theta : \overline{\mathfrak{M}}_{LM} \to \overline{\mathfrak{M}}_{LM} \otimes \overline{\mathfrak{M}}_{LM}$  a map inducing the dual product. From the formula obtained for  $\theta$  in [15]-3.6 one deduce also that

$$\begin{aligned} \theta \circ \rho \circ \bar{\mathfrak{M}}_{\delta_{in}}^{!} \circ \bar{\mathfrak{M}}_{\delta_{out}}(x^{k} \otimes x^{l} \otimes \bar{x}^{\epsilon} \bar{y}^{s} \otimes \bar{x}^{\epsilon'} \bar{y}^{s'}) \\ &= \begin{cases} (1 \otimes \bar{x} + \bar{x} \otimes 1) \sum_{j=0}^{s+s'} \binom{s+s'}{j} x^{n} \otimes \bar{y}^{i} \otimes x^{n} \otimes \bar{y}^{s+s'-j} & \text{if } \epsilon + \epsilon' = 1 \\ \\ &= \sum_{j=0}^{s+s'} \binom{s+s'}{j} x^{n} \otimes \bar{y}^{i} \otimes x^{n} \otimes \bar{y}^{s+s'-j} & \text{if } \epsilon = \epsilon' = 0 \,. \end{cases} \end{aligned}$$

This proves in particular that the composition  $\Phi \circ P$  is not trivial.

§6 - Proof of theorem B.

Let us denote by  $\bigwedge V \otimes \bigwedge^p \bar{V}$  the subvector space of  $\bigwedge V \otimes \bigwedge \bar{V}$  generated by the words of length p in  $\bar{V}$ . The differential of  $\mathfrak{M}_{LM} = (\bigwedge V \otimes \bigwedge \bar{V}, d)$  satisfies  $d(\bigwedge V \otimes \bigwedge^p \bar{V}) \subset$  $\bigwedge V \otimes \bigwedge^p \bar{V}$ . M. Vigué has proved, [21]-Theorem 3.3, that  $H^n_{(i)}(LM) = H^n(\bigwedge V \otimes \bigwedge^i V, d)$ .

To achieve the proof of theorem B it remains to prove that the maps  $\mathfrak{M}^{!}_{\delta_{in}}, \mathfrak{M}_{\delta_{out}}$ , and  $\mu \otimes \bar{\mu}$  respect the word length in  $\bar{V}$ . The obvious relation

$$\mu \otimes \bar{\mu} \left( \bigwedge^r V \otimes \bigwedge^t V \otimes \bigwedge^p \bar{V} \otimes \bigwedge^q \bar{V} \right) \subset \bigwedge^{r+t} (V) \otimes \bigwedge^{p+q} \bar{V}.$$

proves that  $\mu \otimes \overline{\mu}$  respects the word length in  $\overline{V}$ . For  $\mathfrak{M}^!_{\delta_{in}}$ ,  $\mathfrak{M}_{\delta_{out}}$  (defined in Theorem 10 and Theorem 4) the same argument works.

# References

- L. Abrams Two dimensional topological quantum field theory and Frobenius algebras, J. Knot theory ramifications 5 (1996) 569-587.
- [2] L. Abrams Modules, comodules and cotensor products over Frobenius algebras, Preprint.

(1961)

- [3] G.E. Bredon Topology and geometry Graduate Texts in Mathematics 139, Springer-Verlag, 1993.
- [4] Brylinski Loop spaces, Characteristic Classes and Geometric Quantification, Progress in Math 107 Birkhäuser Boston 1993.
- [5] D. Burghelea and M. Vigué A model for cyclic homology and algebraic K-theory for 1-connected topological space, J. of differential Geometry 22 (1985) 243-253.
- [6] M. Chas and D. Sullivan, String topology, preprint math.GT/9911159 To appear in Annals of Math.
- [7] M. Chas and D. Sullivan, Closed String Operators in topology leading to Lie bialgebra and higher string algebra, preprint math.GT/9911159
- [8] D. Chataur, A bordism approach to string topology, preprint, Barcelona (2003).
- [9] R.L. Cohen and V. Godin A Polarized view of string topology, math.AT/03030003.

- [10] R.L. Cohen, J.D.S. Jones, A Homotopy theoretic realization of string topology Math. Ann. 324 (2002) 773-798.
- [11] R.L. Cohen, J.D.S. Jones, and Jun Yan, The Loop Homology algebra of Spheres and Projective Spaces. in Categorical Decomposition Techniques in Algebraic Topology. Progress in Mathematics 215 - BirkhaÄuser-Verlag - 2004
- [12] Y. Felix, S. Halperin and J.-C. Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics 205, Springer-Verlag, 2000.
- [13] Y. Felix, L. Menichi and J.-C. Thomas, Duality in Gerstenhaber algebras, To appear in the Journal of Pure and Appl. Algebra (2005).
- [14] Y. Felix, J.-C. Thomas and M. Vigué-Poirrier, The Hochschild cohomology of a closed manifold, Publications of the IHES 99 (2004) 235-248.
- [15] Y. Felix, J.-C. Thomas and M. Vigué-Poirrier, Rational String homology, Preprint AT/0406593
- [16] A. Hatcher Algebraic Topology Cambridge University Press 2002.
- [17] S. Lang, Differential and Riemannian manifolds. Third edition. Graduate text in mathematics, 160 Springer-Verlag, New York 1995.
- [18] S.A. Merkulov De Rham model for string homology. Preprint AT/0209038.
- [19] D. Sullivan, Infinitesimal computations in topology, Inst. Hautes Etudes Sci. Publ. Math. 47 (1977), 269-331.
- [20] D. Sullivan, Open and closed string field theory interpreted in classical algebraic topology, Inst. Hautes Etudes Sci. Publ. Math. 47 (1977), 269-331.
- [21] M. Vigué-Poirrier, Décomposition de l'homologie des algèbres différentielles graduées commutatives, K-Theory 4 (1991) 399-410.

david.chataur@univ-angers.fr and jean-claude.thomas@univ-angers.fr Département de mathématique Faculté des Sciences 2, Boulevard Lavoisier 49045 Angers, France .