THE q-BINOMIAL FORMULA AND THE ROGERS DILOGARITHM IDENTITY

R.M. KASHAEV

ABSTRACT. The q-binomial formula in the limit $q \to 1^-$ is shown to be equivalent to the Rogers five term dilogarithm identity.

1. INTRODUCTION

For any $q, x \in [0, 1]$ define a q-exponential function as an infinite product

$$\phi(x) := 1/(x;q)_{\infty} := 1/\prod_{n \ge 0} (1-q^n x)$$

A finite product

$$(x;q)_k := \prod_{n=0}^{k-1} (1-q^n x), \quad \forall k \in \mathbb{Z}_{\geq 0}$$

can be expressed as a ratio of two q-exponentials:

$$(x;q)_k = \frac{(x;q)_\infty}{(xq^k;q)_\infty} = \frac{\phi(xq^k)}{\phi(x)}$$

The q-binomial formula (see, for example, [2]) is given by the following identity

(1)
$$\sum_{n>0} \frac{(a;q)_n}{(q;q)_n} z^n = \frac{(az;q)_\infty}{(z;q)_\infty}, \quad |z| < 1$$

which, by using the above notation, can also be written entirely in terms of the function $\phi(x)$:

(2)
$$\sum_{n\geq 0} \frac{\phi(aq^n)}{\phi(q^{n+1})} z^n = \frac{\phi(a)\phi(z)}{\phi(q)\phi(az)}$$

The following expansion formulas

(3)
$$\phi(x) = \sum_{n \ge 0} \frac{x^n}{(q;q)_n}$$

and

(4)
$$\frac{1}{\phi(x)} = \sum_{n \ge 0} \frac{(-1)^n q^{n(n-1)/2} x^n}{(q;q)_n}$$

are both particular cases of the q-binomial formula.

The asymptotic formula

$$\phi(x) \sim e^{-\operatorname{Li}_2(x)/\ln q}, \quad q \to 1^-$$

Date: July 2004.

The work is supported in part by the Swiss National Science Foundation.

where

$$\operatorname{Li}_2(x) := \sum_{n=1}^{\infty} \frac{x^n}{n^2}$$

is the Euler dilogarithm function, has been used in [1, 3] to give an interpretation to $\phi(x)$ as a quantum version of the dilogarithm function. In particular, by using a formal reasoning coming from quantum mechanics, it has been shown that the quantum five term identity

(5)
$$\phi(\mathbf{u})\phi(\mathbf{v}) = \phi(\mathbf{v})\phi(-\mathbf{v}\mathbf{u})\phi(\mathbf{u})$$

where $\phi(\mathbf{u})$, $\phi(\mathbf{v})$, and $\phi(-\mathbf{v}\mathbf{u})$ are elements in the algebra $\mathcal{A}_q = \mathbb{C}_q[[\mathbf{u}, \mathbf{v}]]$ of formal power series in two elements \mathbf{u}, \mathbf{v} satisfying the commutation relation $\mathbf{u}\mathbf{v} = q\mathbf{v}\mathbf{u}$, in the limit $q \to 1^-$ reproduces the Rogers pentagonal identity for the dilogarithm (6)

$$\operatorname{Li}_{2}(a) + \operatorname{Li}_{2}(z) = \operatorname{Li}_{2}(az) + \operatorname{Li}_{2}\left(\frac{a-az}{1-az}\right) + \operatorname{Li}_{2}\left(\frac{z-az}{1-az}\right) + \ln\left(\frac{1-z}{1-az}\right) \ln\left(\frac{1-a}{1-az}\right)$$

The purpose of this paper is to make the statement of the paper [1] mathematically rigorous¹. Namely, we first show that the identity (5) is related to the *q*-binomial formula (1) and then derive from the latter the Rogers identity (6) in the limit $q \to 1^-$. The main result follows.

Theorem 1. Let $q, a, z \in]0, 1[$. Then in the limit $q \to 1^-$ the q-binomial identity (2) leads to the Rogers pentagonal identity (6).

The rest of this paper is organized as follows. In Section 2 the equivalence between he q-binomial formula and the quantum pentagonal identity is explained, while Section 3 contains the proof of Theorem 1.

2. The q-binomial formula and the quantum pentagonal identity

The relation between the formulas (1) and (5) can be established by comparing the expansion coefficients of $a^m z^n$ in (1) and $v^n u^m$ in (5), respectively.

Proposition 1. The q-binomial formula is equivalent to the following set of identities

(7)
$$\frac{q^{mn}}{(q;q)_m(q;q)_n} = \sum_{k=0}^{\min(m,n)} \frac{(-1)^k q^{k(k-1)/2}}{(q;q)_{m-k}(q;q)_{n-k}(q;q)_k}, \quad \forall m, n \in \mathbb{Z}_{\ge 0}$$

Proof. Let us write the q-binomial formula in the form

$$\sum_{n \ge 0} \frac{\phi(aq^n)}{(q;q)_n} z^n = \frac{\phi(a)\phi(z)}{\phi(az)}$$

or, using formula (3) in the left hand side, we have

$$\sum_{m,n\geq 0} \frac{q^{mn}a^m z^n}{(q;q)_m(q;q)_n} = \frac{\phi(a)\phi(z)}{\phi(az)}$$

Again, using the expansion formulas (3), (4) in the right hand side, and equating the coefficients of the monomials $a^m z^n$ in both sides of the equality, we arrive at formula (7).

 $^{^{1}}$ I would like to thank Yu. Manin for posing this question.

Proposition 2. The set of identities (7) is equivalent to the quantum five term identity (5).

Proof. We multiply the both sides of (7) by $v^n u^m$ and sum over m and n. The result can be easily written in the form of equation (5) by using the commutation relation uv = qvu, and, in particular, the formula $v^k u^k q^{k(k-1)/2} = (vu)^k$.

3. Proof of Theorem 1

Lemma 1. Let $k, l \in \mathbb{Z}$ be such that $k \leq l$ and $f_{\pm} : [k, l+1] \to \mathbb{R}_{\geq 0}$ be functions, where f_{-} is decreasing and f_{+} is increasing. Then

(8)
$$\sum_{n=k+1}^{l+1} f_{-}(n) \le \int_{k}^{l+1} f_{-}(t) dt \le \sum_{n=k}^{l} f_{-}(n)$$

(9)
$$\sum_{n=k}^{l} f_{+}(n) \leq \int_{k}^{l+1} f_{+}(t) dt \leq \sum_{n=k+1}^{l+1} f_{+}(n)$$

Proof. The inequality

$$f_{-}(n+1) \le f_{-}(x) \le f_{-}(n), \quad \forall n \in \mathbb{Z} \cap [k,l], \ \forall x \in [n, n+1]$$

implies that $f_{-}(n+1) \leq \int_{n}^{n+1} f_{-}(x) dx \leq f_{-}(n)$. Thus, summing over all possible n we arrive at formula (8). The proof of formula (9) is similar. \Box

Remark 1. The variables k and l in Lemma 1 can take infinite values $k = -\infty$ or $l = \infty$.

In what follows, for any function $f \colon \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ we shall use the notation

$$S(f) := \sum_{n \ge 0} f(n), \quad I(f) := \int_0^\infty f(t) dt$$

If a decreasing function $f: \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ is integrable on $\mathbb{R}_{\geq 0}$ then, as a particular case of Lemma 1, we have

$$S(f) - f(0) \le I(f) \le S(f)$$

or equivalently

(10)
$$0 \le S(f) - I(f) \le f(0)$$

Example 1. The function $f(t) = -\ln(1 - q^t x)$ is decreasing and integrable on $\mathbb{R}_{\geq 0}$, and

$$S(f) = \ln \phi(x), \quad I(f) = -\int_0^\infty \ln(1 - q^t x) dt = \frac{1}{\ln q} \int_0^x \ln(1 - z) \frac{dz}{z} = -\frac{\text{Li}_2(x)}{\ln q}$$

Thus, for any $q, x \in]0, 1[$ inequalities (10) imply that

(11)
$$1 \le \phi(x) e^{\operatorname{Li}_2(x)/\ln q} \le \frac{1}{1-x}$$

Lemma 2. Let $g: \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ be an integrable function increasing in the segment $[0, x_0]$ and decreasing on the interval $[x_0, \infty[$. Let also $n_0 \in \mathbb{Z}_{\geq 0}$ be such that $g(n) \leq g(n_0)$ for all $n \in \mathbb{Z}_{\geq 0}$ (n_0 is equal either to $[x_0]$ (the integer part of x_0) or $[x_0] + 1$). Then

(12)
$$g(n_0) \le \sum_{n \ge 0} g(n) \le \int_0^\infty g(x) dx + g(n_0)$$

Proof. The inequality $g(n_0) \leq \sum_{n\geq 0} g(n)$ follows directly from the positivity of g(x). To prove the second part of (12), note that we can apply Lemma 1 to functions $f_+ = g|_{[0,[x_0]]}$ and $f_- = g|_{[[x_0]+1,\infty[}$. Thus, the left hand sides of the inequalities in Lemma 1 take the forms

$$\sum_{n=0}^{[x_0]-1} g(n) \le \int_0^{[x_0]} g(x) dx, \quad \sum_{n=[x_0]+2}^{\infty} g(n) \le \int_{[x_0]+1}^{\infty} g(x) dx$$

Adding these to each other, we obtain

$$\sum_{n=0}^{\infty} g(n) - g([x_0]) - g([x_0] + 1) \le \int_0^{\infty} g(x) dx - \int_{[x_0]}^{[x_0] + 1} g(x) dx$$

which, combined with the inequality

$$\int_{[x_0]}^{[x_0]+1} g(x) dx \ge g(n'_0)$$

where $\{n_0, n'_0\} = \{[x_0], [x_0] + 1\}$, is equivalent to the second part of (12).

Proposition 3. There exists $\epsilon \in [0, 1[$ such that for any $q \in [1 - \epsilon, 1[$ the function

$$g(x) = \frac{\phi(aq^x)}{\phi(q^{1+x})} z^x$$

where $a, z \in]0, 1[$, satisfies the conditions of Lemma 2.

Proof. The integrability of g(x) is evident. We have the following formula for its derivative

$$\frac{g'(x)}{g(x)} = \ln z - \ln(q)(q-a)S(h_x)$$

where

$$h_x(t) = \frac{q^{x+t}}{(1-q^{1+x+t})(1-aq^{x+t})}$$

satisfies the conditions of Lemma 1 so that

$$S(h_x) \ge I(h_x) = -\frac{1}{\ln(q)(q-a)} \ln\left(\frac{z(1-aq^x)}{1-q^{1+x}}\right)$$

Evidently, the function $S(h_x)$ is decreasing in x. Assuming that q > 1 - z(1 - a), we obtain

$$\frac{g'(0)}{g(0)} \ge \ln\left(\frac{z(1-a)}{1-q}\right) > 0$$

Besides, it is easy to see that

$$\lim_{x \to \infty} \frac{g'(x)}{g(x)} = \ln z < 0$$

Thus, we have shown that for $\epsilon = z(1-a)$ and any $q \in [1-\epsilon, 1[$ the continuous function g'(x)/g(x) is decreasing, positive at x = 0 and negative for sufficiently large x, i.e. there exists unique $x_0 \in [0, \infty[$ such that $g'(x_0) = 0$ and all conditions of Lemma 2 are satisfied.

Proposition 4.

(13)
$$\lim_{q \to 1^{-}} \ln(q) \ln S(g) = F(\xi_0), \quad \xi_0 = \frac{1-z}{1-az}$$

where

$$F(\xi) = \operatorname{Li}_2(\xi) - \operatorname{Li}_2(a\xi) + \ln(\xi)\ln(z)$$

Proof. For any $\xi \in]0, 1[$ equation (11) implies that

$$\lim_{q\to 1^-}\ln(q)\ln(g\left(\ln\xi/\ln q\right))=F(\xi)$$

Thus, one has asymptotically

$$g((\ln \xi/\ln q) \sim e^{\frac{F(\xi)}{\ln q}}, \quad q \to 1^-$$

and, by using the steepest decent method, one has also

$$I(g) \sim e^{\frac{F(\xi_0)}{\ln q}}, \quad q \to 1^-$$

where $\xi_0 = (1-z)/(1-az) \in]0, 1[$ is the unique solution of the equation $F'(\xi) = 0$. The asymptotic formula for S(g) follows immediately from Lemma 2 after taking into account the fact that

$$x_0 \sim n_0 \sim \frac{\ln \xi_0}{\ln q}, \quad q \to 1$$

and, correspondingly, $g(n_0) \sim g(x_0) \sim I(g), q \to 1^-$.

Proof of Theorem 1. Using Lemma 1, we have immediately

$$\lim_{q \to 1^{-}} \ln(q) \ln\left(\frac{\phi(a)\phi(z)}{\phi(q)\phi(az)}\right) = \operatorname{Li}_{2}(1) + \operatorname{Li}_{2}(az) - \operatorname{Li}_{2}(a) - \operatorname{Li}_{2}(y)$$

Combining this formula with equation (13), we conclude that the *q*-binomial identity (2) leads to the following identity:

$$F(\xi_0) = \text{Li}_2(1) + \text{Li}_2(az) - \text{Li}_2(a) - \text{Li}_2(z)$$

or explicitly,

$$\operatorname{Li}_{2}(\xi_{0}) - \operatorname{Li}_{2}(a\xi_{0}) + \ln(\xi_{0})\ln(z) = \operatorname{Li}_{2}(1) + \operatorname{Li}_{2}(az) - \operatorname{Li}_{2}(a) - \operatorname{Li}_{2}(z)$$

which we rewrite in the form

$$\operatorname{Li}_{2}(a) + \operatorname{Li}_{2}(z) = \operatorname{Li}_{2}(az) + \operatorname{Li}_{2}(a\xi_{0}) + \operatorname{Li}_{2}(1) - \operatorname{Li}_{2}(\xi_{0}) - \ln(\xi_{0})\ln(z)$$

Using the identity

$$\operatorname{Li}_{2}(x) + \operatorname{Li}_{2}(1-x) = \operatorname{Li}_{2}(1) - \ln(x)\ln(1-x), \quad \forall x \in [0,1]$$

we rewrite it further

$$\text{Li}_2(a) + \text{Li}_2(z) = \text{Li}_2(az) + \text{Li}_2(a\xi_0) + \text{Li}_2(1 - \xi_0) + \ln(\xi_0) \ln((1 - \xi_0)/z)$$

which is exactly the Rogers identity (6).

R.M. KASHAEV

References

[1] L.D. Faddeev, R.M. Kashaev: Quantum dilogarithm, Mod. Phys. Lett. A9 (1994) 427-434.

[2] G. Gasper, M. Rahman: Basic hypergeometric series, Cambridge University Press, 1990.

[3] A.N. Kirillov: *Dilogarithm identities*, Prog. Theor. Phys. Suppl. 118 (1995), 61–142.

Université de Genève, Section de mathématiques, 2-4, rue du Lièvre, CP 240, 1211 Genève 24, Suisse

V.A. Steklov Institute of Mathematics at St. Petersburg , 27, Fontanka, St. Petersburg 191023, Russia

E-mail address: Rinat.Kashaev@math.unige.ch