Gaudin model and opers

Edward Frenkel

Abstract. This is a review of our previous works [FFR, F1, F3] (some of them joint with B. Feigin and N. Reshetikhin) on the Gaudin model and opers. We define a commutative subalgebra in the tensor power of the universal enveloping algebra of a simple Lie algebra \mathfrak{g} . This algebra includes the hamiltonians of the Gaudin model, hence we call it the Gaudin algebra. It is constructed as a quotient of the center of the completed enveloping algebra of the affine Kac-Moody algebra $\hat{\mathfrak{g}}$ at the critical level. We identify the spectrum of the Gaudin algebra with the space of opers associated to the Langlands dual Lie algebra ${}^{L}\mathfrak{g}$ on the projective line with regular singularities at the marked points. Next, we recall the construction of the eigenvectors of the Gaudin algebra using the Wakimoto modules over $\hat{\mathfrak{g}}$ of critical level. The Wakimoto modules are naturally parameterized by Miura opers (or, equivalently, Cartan connections), and the action of the center on them is given by the Miura transformation. This allows us to relate solutions of the Bethe Ansatz equations to Miura opers and ultimately to the flag varieties associated to the Langlands dual Lie algebra ${}^{L}\mathfrak{q}$.

Introduction

Let \mathfrak{g} be a finite-dimensional simple Lie algebra over \mathbb{C} and $U(\mathfrak{g})$ its universal enveloping algebra. Choose a basis $\{J_a\}, a = 1, \ldots, d$, of \mathfrak{g} , and let $\{J^a\}$ the dual basis with respect to a non-degenerate invariant bilinear form on \mathfrak{g} . Let z_1, \ldots, z_N be a collection of distinct complex numbers.

The Gaudin Hamiltonians are the following elements of the algebra $U(\mathfrak{g})^{\otimes N}$:

(0.1)
$$\Xi_i = \sum_{j \neq i} \sum_{a=1}^d \frac{J_a^{(i)} J^{a(j)}}{z_i - z_j}, \qquad i = 1, \dots, N,$$

where for $A \in \mathfrak{g}$ we denote by $A^{(i)}$ the element of $U(\mathfrak{g})^{\otimes N}$ which is the tensor product of A in the *i*th factor and 1 in all other factors. One checks easily that these elements commute with each other and are invariant with respect to the diagonal action of \mathfrak{g} on $U(\mathfrak{g})^{\otimes N}$.

Partially supported by grants from the NSF and DARPA.

For any collection M_1, \ldots, M_N of g-modules, the Gaudin Hamiltonians give rise to commuting linear operators on $\bigotimes_{i=1}^N M_i$. We are interested in the diagonalization of these operators. More specifically, we will consider the following two cases: when all of the M_i 's are Verma modules and when they are finite-dimensional irreducible modules.

It is natural to ask: are there any other elements in $U(\mathfrak{g})^{\otimes N}$ which commute with the Gaudin Hamiltonians? Clearly, the N-fold tensor product $Z(\mathfrak{g})^{\otimes N}$ of the center $Z(\mathfrak{g})$ of $U(\mathfrak{g})$ is the center of $U(\mathfrak{g})^{\otimes N}$, and its elements obviously commute with the Ξ_i 's. As shown in [FFR], if \mathfrak{g} has rank grater than one, then in addition to the Gaudin Hamiltonians and the central elements there are other elements in $U(\mathfrak{g})^{\otimes N}$ of orders higher than two which commute with the Gaudin operators and with each other (but explicit formulas for them are much more complicated and unknown in general). Adjoining these "higher Gaudin Hamiltonians" to the Ξ_i 's together with the center $Z(\mathfrak{g})^{\otimes N}$, we obtain a large commutative subalgebra of $U(\mathfrak{g})^{\otimes N}$. We will call it the *Gaudin algebra* and denote it by $\mathcal{Z}_{(z_i)}(\mathfrak{g})$.

The construction of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ will be recalled in Section 2. The key point is the realization of $U(\mathfrak{g})^{\otimes N}$ as the space of *coinvariants* of induced modules over the affine Kac-Moody algebra $\widehat{\mathfrak{g}}$ on the projective line. Using this realization, we obtain a surjective map from the center of the completed universal enveloping algebra of $\widehat{\mathfrak{g}}$ at the *critical level* onto $\mathcal{Z}_{(z_i)}(\mathfrak{g})$.

The next natural question is what is the spectrum of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, i.e., the set of all maximal ideals of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, or equivalently, algebra homomorphisms $\mathcal{Z}_{(z_i)}(\mathfrak{g}) \to \mathbb{C}$. Knowing the answer is important, because then we will know how to think about the common eigenvalues of the higher Gaudin operators on the tensor products $\bigotimes_{i=1}^{N} M_i$ of \mathfrak{g} -modules. These common eigenvalues correspond to points of the spectrum of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$.

The answer comes from the description of the center of the completed universal enveloping algebra of $\hat{\mathfrak{g}}$ at the critical level. In [FF2, F2] it is shown that the spectrum of this center (more precisely, the center of the corresponding vertex algebra) is canonically identified with the space of the so-called ^LG-opers, where ^LG is the Langlands dual Lie group to \mathfrak{g} (of adjoint type), on the formal disc. This result leads us to the following description of the spectrum of the algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ of higher Gaudin Hamiltonians: it is the space of ^LG-opers on \mathbb{P}^1 with regular singularities at the points z_1, \ldots, z_N and ∞ .

We obtain this description from some basic facts about the spaces of coinvariants from [FB]. Recall that the space of coinvariants $H_V(X; (x_i); (M_i))$ is defined in [FB] for any (quasi-conformal) vertex algebra V, a smooth projective curve X, a collection x_1, \ldots, x_n of distinct points of X and a collection of V-modules M_1, \ldots, M_n attached to those points. Suppose that V is a commutative vertex algebra, and so in particular it is a commutative algebra. Suppose that the spectrum of V is the space S(D) of certain geometric objects, such as ^LG-opers, on the disc $D = \operatorname{Spec} \mathbb{C}[[t]]$. Then a V-module is the same as a smooth module over the complete topological algebra U(V) of functions on $S(D^{\times})$, which is the space of our objects (such as ^LG-opers) on the punctured disc $D^{\times} = \operatorname{Spec} \mathbb{C}((t))$. Suppose in addition that each V-module M_i is the space of functions on a subspace S_i of $S(D^{\times})$ (with its natural Fun $S(D^{\times})$ -module structure). Then the space of coinvariants $H_V(X; (x_i); (M_i))$ is naturally a commutative algebra, and its spectrum is the space of our objects (such as ^LG-opers) on $X \setminus \{x_1, \ldots, x_n\}$ whose restriction to the punctured disc $D_{x_i}^{\times}$ around x_i belongs to $S_i \subset S(D_{x_i}^{\times}), i = 1, \ldots, n$.

For example, if $\mathfrak{g} = \mathfrak{sl}_2$, then ${}^L G = PGL_2$, and PGL_2 -opers are the same as second order differential operators $\partial_t^2 - q(t)$ acting from sections of the line bundle $\Omega^{-1/2}$ to sections of $\Omega^{3/2}$. A PGL_2 -oper on \mathbb{P}^1 with regular singularities at z_1, \ldots, z_N and ∞ may be written as the Fuchsian differential operator of second order with regular singularities at z_1, \ldots, z_N ,

$$\partial_t^2 - \sum_{i=1}^N \frac{c_i}{(t-z_i)^2} - \sum_{i=1}^N \frac{\mu_i}{t-z_i}$$

satisfying the condition $\sum_{i=1}^{N} \mu_i = 0$ that insures that it also has regular singularity at ∞ . Defining such an operator is the same as giving a collection of numbers $c_i, \mu_i, i = 1, \ldots, N$, such that $\sum_{i=1}^{N} \mu_i = 0$. The set

$$\left\{c_i, \mu_i, i=1, \dots, N \mid \sum_{i=1}^N \mu_i = 0\right\},\$$

is then the spectrum of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, which in this case is the polynomial algebra generated by the Casimir operators $C_i = \frac{1}{2} \sum_a J_a^{(i)} J^{a(i)}, i = 1, \ldots, N$, and the Gaudin Hamiltonians Ξ_i , subject to the relation $\sum_{i=1}^N \Xi_i = 0$. In other words, the numbers c_i record the eigenvalues of the C_i 's, while the numbers μ_i record the eigenvalues of the Ξ_i 's.

For a general simple Lie algebra \mathfrak{g} , the Gaudin algebra $\mathfrak{Z}_N(\mathfrak{g})$ has many more generators, and its spectrum does not have such a nice system of coordinates as the c_i 's and the μ_i 's in the above example. Therefore the description of the spectrum as a space of LG -opers is very useful. In particular, we obtain that common eigenvalues of the higher Gaudin Hamiltonians are encoded by LG -opers on \mathbb{P}^1 , with regular singularities at prescribed points. These LG -opers appear as generalizations of the above second order Fuchsian operators.

Next, we ask which points in the spectrum of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ might occur as the common eigenvalues on particular tensor products $\bigotimes_{i=1}^{N} M_i$. We answer this question first in the case when each M_i admits a central character: namely, it turns out that the central character of M_i fixes the residue of the ^LG-oper at the point z_i . We then show that if all \mathfrak{g} -modules M_i are finite-dimensional and irreducible, then the ^LG-opers encoding possible eigenvalues of the higher Gaudin Hamiltonians in $\bigotimes_{i=1}^{N} M_i$ necessarily have trivial monodromy representation.

We conjecture that there is a bijection between the eigenvalues of the Gaudin Hamiltonians on $\bigotimes_{i=1}^{N} M_i$, where the M_i 's are irreducible finite-dimensional g-modules, and ^LG-opers on \mathbb{P}^1 with prescribed singularities at z_1, \ldots, z_N, ∞ and trivial monodromy.

Thus, we obtain a correspondence between two seemingly unrelated objects: the eigenvalues of the generalized Gaudin Hamiltonians and the ${}^{L}G$ -opers on \mathbb{P}^{1} . The connection between the eigenvalues of the Gaudin operators and differential operators of some sort has been observed previously, but it was not until [FFR, F1] that this phenomenon was explained conceptually.

We present a more geometric description the ${}^{L}G$ -opers without monodromy (which occur as the eigenvalues of the Gaudin Hamiltonians) as isomorphism classes of holomorphic maps from \mathbb{P}^{1} to ${}^{L}G/{}^{L}B$, the flag manifold of ${}^{L}G$, satisfying a certain transversality condition. For example, if ${}^{L}G = PGL_2$, they may be described as holomorphic maps $\mathbb{P}^{1} \to \mathbb{P}^{1}$ whose derivative vanishes to prescribed orders at the marked points z_1, \ldots, z_N and ∞ , and does not vanish anywhere else (these orders correspond to the highest weights of the finite-dimensional representations inserted at those points).

If the ^LG-oper is non-degenerate (in the sense explained in Section 5.2), then we can associate to it an eigenvector of the Gaudin Hamiltonians called a *Bethe vector*. The procedure to construct eigenvectors of the Gaudin Hamiltonians that produces these vectors is known as the *Bethe Ansatz*. In [FFR] we explained that this procedure can also be understood in the framework of coinvariants of $\hat{\mathfrak{g}}$ -modules of critical level. We need to use a particular class of $\hat{\mathfrak{g}}$ -modules, called the *Wakimoto modules*.

Let us recall that the Wakimoto modules of critical level are naturally parameterized by objects closely related to opers, which we call *Miura opers*. They may also be described more explicitly as certain connections on a particular ${}^{L}H$ -bundle $\Omega^{-\rho}$ on the punctured disc, where ${}^{L}H$ is the Cartan subgroup of ${}^{L}G$. The center acts on the Wakimoto module corresponding to a Cartan connection by the Miura transformation of this connection (see [F2]). The idea of [FFR] was to use the spaces of coinvariants of the tensor product of the Wakimoto modules to construct eigenvectors of the generalized Gaudin Hamiltonians. We found in [FFR] that the eigenvalues of the Gaudin Hamiltonians on these vectors are encoded by the ${}^{L}G$ -opers which are obtained by applying the Miura transformation to certain Cartan connections on \mathbb{P}^{1} .

More precisely, the Bethe vector depends on an *m*-tuple of complex numbers w_j , where j = 1, ..., m, with an extra datum attached to each of them, $i_j \in I$, where I is the set of nodes of the Dynkin diagram of \mathfrak{g} (or equivalently, the set of simple roots of \mathfrak{g}). These numbers have to be distinct from the z_i 's and satisfy the following system of *Bethe Ansatz equations*:

(0.2)
$$\sum_{i=1}^{N} \frac{\langle \lambda_i, \check{\alpha}_{i_j} \rangle}{w_j - z_i} - \sum_{s \neq j} \frac{\langle \alpha_{i_s}, \check{\alpha}_{i_j} \rangle}{w_j - w_s} = 0, \quad j = 1, \dots, m,$$

where λ_i denotes the highest weight of the finite-dimensional g-module $M_i = V_{\lambda_i}, i = 1, \ldots, N$.

We can compute explicitly the ${}^{L}G$ -oper encoding the eigenvalues of the generalized Gaudin Hamiltonians on this vector. As shown in [FFR], this ${}^{L}G$ -oper is obtained by applying the Miura transformation of the connection

(0.3)
$$\partial_t + \sum_{i=1}^N \frac{\lambda_i}{t - z_i} - \sum_{j=1}^m \frac{\alpha_{i_j}}{t - w_j}$$

on the ^LH-bundle $\Omega^{-\rho}$ on \mathbb{P}^1 . This ^LG-oper automatically has trivial monodromy.

The Bethe vector corresponding to a solution of the system (0.2) is a highest weight vector in $\bigotimes_{i=1}^{N} V_{\lambda_i}$ of weight

$$\mu = \sum_{i=1}^{N} \lambda_i - \sum_{j=1}^{m} \alpha_{i_j},$$

so it can only be non-zero if μ is a dominant integral weight of \mathfrak{g} . But it is still interesting to describe the set of *all* solutions of the Bethe Ansatz equations (0.2), even for non-dominant weights μ .

While the eigenvalues of the Gaudin Hamiltonians are parameterized by ${}^{L}G$ opers, it turns out that the solutions of the Bethe Ansatz equations are parameterized by the (non-degenerate) Miura ${}^{L}G$ -opers. As mentioned above, those may
in turn be related to very simple objects, namely, connections on an ${}^{L}H$ -bundle $\Omega^{-\rho}$ of the kind given above in formula (0.3).

A ${}^{L}G$ -oper on a curve X is by definition a triple $(\mathcal{F}, \nabla, \mathcal{F}_{L_B})$, where \mathcal{F} is a ${}^{L}G$ -bundle on X, ∇ is a connection on \mathcal{F} and \mathcal{F}_{L_B} is a reduction of \mathcal{F} to a Borel subgroup ${}^{L}B$ of ${}^{L}G$, which satisfies a certain transversality condition with ∇ . A Miura ${}^{L}G$ -oper is by definition a quadruple $(\mathcal{F}, \nabla, \mathcal{F}_{L_B}, \mathcal{F}'_{L_B})$ where \mathcal{F}'_{L_B} is another ${}^{L}B$ -reduction of \mathcal{F} , which is preserved by ∇ . The space of Miura opers on a curve X whose underlying oper has regular singularities and trivial monodromy representation (so that \mathcal{F} is isomorphic to the trivial bundle) is isomorphic to the flag manifold ${}^{L}G/{}^{L}B$ of ${}^{L}G$. Indeed, in order to define the ${}^{L}B$ -reduction \mathcal{F}'_{L_B} of such \mathcal{F} everywhere, it is sufficient to define it at one point $x \in X$ and then use the connection to "spread" it around. But choosing a ${}^{L}B$ -reduction at one point means choosing an element of the twist of ${}^{L}G/{}^{L}B$ by \mathcal{F}_x , and so we see that the space of all reductions is isomorphic to the flag manifold of ${}^{L}G$.

The relative position of the two reductions \mathcal{F}_{L_B} and \mathcal{F}'_{L_B} at each point of X is measured by an element w of the Weyl group W of G. The two reductions are in generic relative position (corresponding to w = 1) almost everywhere on X. The mildest possible non-generic relative positions correspond to the simple reflections s_i from W. We call a Miura oper on \mathbb{P}^1 with marked points z_1, \ldots, z_N non-degenerate if the two reductions \mathcal{F}_B and \mathcal{F}'_B are in generic position at z_1, \ldots, z_N , and elsewhere on \mathbb{P}^1 their relative position is either generic or corresponds to a simple reflection. We denote the points where the relative position is not generic

by $w_j, j = 1, ..., m$; each point w_j comes together with a simple reflection s_{i_j} , or equivalently a simple root α_{i_j} attached to it.

It is then easy to see that this collection satisfies the equations (0.2), and conversely any solution of (0.2) corresponds to a non-degenerate Miura oper (or to an ^LH-connection (0.3)). Thus, we obtain that there is a bijection between the set of solutions of (0.2) (for all possible collections $\{i_1, \ldots, i_m\}$) and the set of non-degenerate Miura ^LG-opers such that the underlying ^LG-opers have prescribed residues at the points z_1, \ldots, z_N, ∞ and trivial monodromy.

Now let us fix $\lambda_1, \ldots, \lambda_N$ and μ . Then every LG -oper τ on \mathbb{P}^1 with regular singularities at z_1, \ldots, z_N and ∞ and with prescribed residues corresponding to $\lambda_1, \ldots, \lambda_N$ and μ and trivial monodromy admits a horizontal LB -reduction \mathcal{F}'_{L_B} satisfying the conditions of a non-degenerate Miura oper. Since these are open conditions, we find that for such z_1, \ldots, z_N the non-degenerate Miura oper structures on a particular LG -oper τ on \mathbb{P}^1 form an open dense subset in the set of all Miura oper structures on τ . But the set of all Miura structures on a given LG -oper τ is isomorphic to the flag manifold ${}^LG/{}^LB$. Therefore we find that the set of non-degenerate Miura oper structures on τ is an open dense subset of ${}^LG/{}^LB$!

Recall that the set of all solutions of the Bethe Ansatz equations (0.2) is the union of the sets of non-degenerate Miura oper structures on all ${}^{L}G$ -opers with trivial monodromy. Hence it is naturally a disjoint union of subsets, parameterized by these ${}^{L}G$ -opers. We have now identified each of these sets with an open and dense subset of the flag manifold ${}^{L}G/{}^{L}B$.

Let us summarize our results:

- the eigenvalues of the Hamiltonians of the Gaudin model associated to a simple Lie algebra \mathfrak{g} on the tensor product of finite-dimensional representations are encoded by ${}^{L}G$ -opers on \mathbb{P}^{1} , where ${}^{L}G$ is the Langlands dual group of G, which have regular singularities at the marked points $z_{1}, \ldots, z_{N}, \infty$ and trivial monodromy;
- if such an oper τ is non-degenerate, then we can associate to it a solution of the Bethe Ansatz equations (0.2), which gives rise to the Bethe eigenvector of dominant integral weight whose eigenvalues are encoded by τ ;
- there is a one-to-one correspondence between the set of all solutions of the Bethe Ansatz equations (0.2) and the set of non-degenerate Miura opers corresponding to a fixed ${}^{L}G$ -oper;
- the set of non-degenerate Miura opers corresponding to the same underlying ${}^{L}G$ -oper is an open dense subset of the flag manifold ${}^{L}G/{}^{L}B$ of the Langlands dual group, and therefore the set of all solutions of the Bethe Ansatz equations (0.2) is the union of certain open dense subsets of the flag manifold of the Langlands dual group, one for each ${}^{L}G$ -oper.

Finally, to a degenerate ${}^{L}G$ -oper we can also attach, at least in some cases, an eigenvector of the Gaudin hamiltonians by generalizing the Bethe Ansatz procedure, as explained in Section 5.5.

Gaudin model and opers

The paper is organized as follows. In Section 1 we introduce opers and discuss their basic properties. We define opers with regular singularities and their residues. In Section 2, following [FFR], we define the Gaudin algebra using the coinvariants of the affine Kac-Moody algebra $\hat{\mathfrak{g}}$ of critical level. We recall the results of [FF2, F2] on the isomorphism of the center of the completed universal enveloping algebra of $\hat{\mathfrak{g}}$ at the critical level and the algebra of functions on the space of ^LG-opers on the punctured disc. Using these results and general facts about the spaces of coinvariants from [FB], we describe the spectrum of the Gaudin algebra. In Section 3 we introduce Miura opers, Cartan connections and the Miura transformation and describe their properties, following [F2, F3]. We use these results in the next section, Section 4, to describe the Bethe Ansatz, a construction of eigenvectors of the Gaudin algebra. We introduce the Wakimoto modules of critical level, following [FF1, F2]. The Wakimoto modules are naturally parameterized by the Cartan connections on the punctured disc introduced in Section 3. The action of the center on the Wakimoto modules is given by the Miura transformation. We construct the Bethe vectors, following [FFR], using the coinvariants of the Wakimoto modules. We show that the Bethe Ansatz equations which ensure that this vector is an eigenvector of the Gaudin Hamiltonians coincide with the requirement that the Miura transformation of the Cartan connection on \mathbb{P}^1 encoding the Wakimoto modules has no singularities at the points w_1, \ldots, w_m . Finally, in Section 5 we consider the Gaudin model in the case when all modules M_i finite-dimensional modules. We describe the precise connection between the spectrum of the Gaudin algebra on the tensor product of finite-dimensional modules and the set of ^{L}G -opers with prescribed singularities at z_1, \ldots, z_N, ∞ and trivial monodromy.

1. Opers

1.1. Definition of opers

Let G be a simple algebraic group of adjoint type, B a Borel subgroup and N = [B, B] its unipotent radical, with the corresponding Lie algebras $\mathbf{n} \subset \mathbf{b} \subset \mathbf{g}$. The quotient H = B/N is a torus. Choose a splitting $H \to B$ of the homomorphism $B \to H$ and the corresponding splitting $\mathfrak{h} \to \mathfrak{b}$ at the level of Lie algebras. Then we will have a Cartan decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}$. We will choose generators $\{e_i\}, i = 1, \ldots, \ell$ of \mathfrak{n} and generators $\{f_i\}, i = 1, \ldots, \ell$ of \mathfrak{n}_- corresponding to simple roots, and denote by $\check{\rho} \in \mathfrak{h}$ the sum of the fundamental coweights of \mathfrak{g} . Then we will have the following relations: $[\check{\rho}, e_i] = 1, [\check{\rho}, f_i] = -1$.

A *G*-oper on a smooth curve *X* (or a disc $D \simeq \operatorname{Spec} \mathbb{C}[[t]]$ or a punctured disc $D^{\times} = \operatorname{Spec} \mathbb{C}((t))$) is by definition a triple $(\mathcal{F}, \nabla, \mathcal{F}_B)$, where \mathcal{F} is a principal *G*-bundle \mathcal{F} on *X*, ∇ is a connection on \mathcal{F} and \mathcal{F}_B is a *B*-reduction of \mathcal{F} such that locally on *X* (with respect to a local coordinate *t* and a local trivialization of \mathcal{F}_B)

the connection has the form

(1.1)
$$\nabla = \partial_t + \sum_{i=1}^{t} \psi_i(t) f_i + \mathbf{v}(t),$$

where each $\psi_i(t)$ is a nowhere vanishing function, and $\mathbf{v}(t)$ is a \mathfrak{b} -valued function. The space of *G*-opers on *X* is denoted by $\operatorname{Op}_G(X)$.

This definition is due to A. Beilinson and V. Drinfeld [BD1] (in the case when X is the punctured disc opers were first introduced in [DS]).

In particular, if $U = \operatorname{Spec} R$ is an affine curve with the ring of functions Rand t is a global coordinate on U (for example, if $U = \operatorname{Spec} \mathbb{C}[[t]]$), then $\operatorname{Op}_G(U)$ is isomorphic to the quotient of the space of operators of the form

(1.2)
$$\nabla = \partial_t + \sum_{i=1}^{\ell} f_i + \mathbf{v}(t), \qquad \mathbf{v}(t) \in \mathfrak{b}(R),$$

by the action of the group N(R) (we use the action of H(R) to make all functions $\psi_i(t)$ equal to 1). Recall that the gauge transformation of an operator $\partial_t + A(t)$, where $A(t) \in \mathfrak{g}(R)$ by $g(t) \in G(R)$ is given by the formula

$$g \cdot (\partial_t + A(t)) = \partial_t + gA(t)g^{-1} - \partial_t g \cdot g^{-1}.$$

The operator ad $\check{\rho}$ defines the principal gradation on \mathfrak{b} , with respect to which we have a direct sum decomposition $\mathfrak{b} = \bigoplus_{i>0} \mathfrak{b}_i$. Set

$$p_{-1} = \sum_{i=1}^{\ell} f_i$$

Let p_1 be the unique element of degree 1 in \mathfrak{n} , such that $\{p_{-1}, 2\check{\rho}, p_1\}$ is an \mathfrak{sl}_2 -triple. Let $V_{\operatorname{can}} = \bigoplus_{i \in E} V_{\operatorname{can},i}$ be the space of ad p_1 -invariants in \mathfrak{n} . Then p_1 spans $V_{\operatorname{can},1}$. Choose a linear generator p_j of $V_{\operatorname{can},d_j}$ (if the multiplicity of d_j is greater than one, which happens only in the case $\mathfrak{g} = D_{2n}^{(1)}, d_j = 2n$, then we choose linearly independent vectors in $V_{\operatorname{can},d_j}$). The following result is due to Drinfeld and Sokolov [DS] (the proof is reproduced in Lemma 2.1 of [F3]).

Lemma 1.1. The gauge action of N(R) on $\widetilde{\operatorname{Op}}_G(\operatorname{Spec} R)$ is free, and each gauge equivalence class contains a unique operator of the form $\nabla = \partial_t + p_{-1} + \mathbf{v}(t)$, where $\mathbf{v}(t) \in V_{\operatorname{can}}(R)$, so that we can write

(1.3)
$$\mathbf{v}(t) = \sum_{j=1}^{\ell} v_j(t) \cdot p_j.$$

1.2. PGL_2 -opers

For $\mathfrak{g} = \mathfrak{sl}_2, G = PGL_2$ we obtain an identification of the space of PGL_2 -opers with the space of operators of the form

$$\partial_t + \left(\begin{array}{cc} 0 & v(t) \\ 1 & 0 \end{array} \right).$$

If we make a change of variables $t = \varphi(s)$, then the corresponding connection operator will become

$$\partial_s + \left(\begin{array}{cc} 0 & \varphi'(s)v(\varphi(s)) \\ \varphi'(s) & 0 \end{array} \right)$$

Applying the B-valued gauge transformation with

$$\left(\begin{array}{cc} 1 & \frac{1}{2} \frac{\varphi''(s)}{\varphi'(s)} \\ 0 & 1 \end{array}\right) \left(\begin{array}{cc} (\varphi'(s))^{1/2} & 0 \\ 0 & (\varphi'(s))^{1/2} \end{array}\right),$$

we obtain the operator

$$\partial_s + \left(\begin{array}{cc} 0 & v(\varphi(s))\varphi'(s)^2 - \frac{1}{2}\{\varphi, s\} \\ 1 & 0 \end{array}\right),$$

where

$$\{\varphi,s\} = \frac{\varphi^{\prime\prime\prime}}{\varphi^{\prime}} - \frac{3}{2} \left(\frac{\varphi^{\prime\prime}}{\varphi^{\prime}}\right)^2$$

is the Schwarzian derivative. Thus, under the change of variables $t = \varphi(s)$ we have

$$v(t) \mapsto v(\varphi(s))\varphi'(s)^2 - \frac{1}{2}\{\varphi, s\}$$

this coincides with the transformation properties of the second order differential operators $\partial_t^2 - v(t)$ acting from sections of $\Omega^{-1/2}$ to sections of $\Omega^{3/2}$, where Ω is the canonical line bundle on X. Such operators are known as *projective connections* on X (see, e.g., [FB], Sect. 9.2), and so PGL_2 -opers are the same as projective connections.

For a general \mathfrak{g} , the first coefficient function $v_1(t)$ in (1.3) transforms as a projective connection, and the coefficient $v_i(t)$ with i > 1 transforms as a $(d_i + 1)$ -differential on X. Thus, we obtain an isomorphism

(1.4)
$$\operatorname{Op}_{G}(X) \simeq \operatorname{Proj}(X) \times \bigoplus_{i=2}^{\ell} \Gamma(X, \Omega^{(d_{i}+1)}).$$

1.3. Opers with regular singularities

Let x be a point of a smooth curve X and $D_x = \operatorname{Spec} \mathcal{O}_x, D_x^{\times} = \operatorname{Spec} \mathcal{K}_x$, where \mathcal{O}_x is the completion of the local ring of x and \mathcal{K}_x is the field of fractions of \mathcal{O}_x . Choose a formal coordinate t at x, so that $\mathcal{O}_x \simeq \mathbb{C}[[t]]$ and $\mathcal{K}_x = \mathbb{C}((t))$. Recall that the space $\operatorname{Op}_G(D_x)$ (resp., $\operatorname{Op}_G(D_x^{\times}))$ of G-opers on D_x (resp., $D_x^{\times})$ is the quotient of the space of operators of the form (1.1) where $\psi_i(t)$ and $\mathbf{v}(t)$ take values in \mathcal{O}_x (resp., in \mathcal{K}_x) by the action of $B(\mathcal{O}_x)$ (resp., $B(\mathcal{K}_x)$).

A G-oper on D_x with regular singularity at x is by definition (see [BD1], Sect. 3.8.8) a $B(\mathcal{O}_x)$ -conjugacy class of operators of the form

(1.5)
$$\nabla = \partial_t + t^{-1} \left(\sum_{i=1}^{\ell} \psi_i(t) f_i + \mathbf{v}(t) \right),$$

where $\psi_i(t) \in \mathcal{O}_x, \psi_i(0) \neq 0$, and $\mathbf{v}(t) \in \mathfrak{b}(\mathcal{O}_x)$. Equivalently, it is an $N(\mathcal{O}_x)$ equivalence class of operators

(1.6)
$$\nabla = \partial_t + \frac{1}{t} \left(p_{-1} + \mathbf{v}(t) \right), \qquad \mathbf{v}(t) \in \mathfrak{b}(\mathfrak{O}_x)$$

Denote by $\operatorname{Op}_{G}^{\operatorname{RS}}(D_x)$ the space of opers on D_x with regular singularity. It is easy to see that the natural map $\operatorname{Op}_{G}^{\operatorname{RS}}(D_x) \to \operatorname{Op}_{G}(D_x^{\times})$ is injective. Therefore an oper with regular singularity may be viewed as an oper on the punctured disc. But to an oper with regular singularity one can unambiguously attach a point in

$$\mathfrak{g}/G := \operatorname{Spec} \mathbb{C}[\mathfrak{g}]^G \simeq \mathbb{C}[\mathfrak{h}]^W =: \mathfrak{h}/W$$

its residue, which in our case is equal to $p_{-1} + \mathbf{v}(0)$.

Given $\check{\lambda} \in \mathfrak{h}$, we denote by $\operatorname{Op}_{G}^{\operatorname{RS}}(D_{x})_{\check{\lambda}}$ the subvariety of $\operatorname{Op}_{G}^{\operatorname{RS}}(D_{x})$ which consists of those opers that have residue $\varpi(-\check{\lambda} - \check{\rho}) \in \mathfrak{h}/W$, where ϖ is the projection $\mathfrak{h} \to \mathfrak{h}/W$.

In particular, the residue of a regular oper $\partial_t + p_{-1} + \mathbf{v}(t)$, where $\mathbf{v}(t) \in \mathfrak{b}(\mathcal{O}_x)$, is equal to $\overline{\omega}(-\check{\rho})$ (see [BD1]). Indeed, a regular oper may be brought to the form (1.6) by using the gauge transformation with $\check{\rho}(t) \in B(\mathcal{K}_x)$, after which it takes the form

$$\partial_t + \frac{1}{t} \left(p_{-1} - \check{\rho} + t \cdot \check{\rho}(t) (\mathbf{v}(t)) \check{\rho}(t)^{-1} \right).$$

If $\mathbf{v}(t)$ is regular, then so is $\check{\rho}(t)(\mathbf{v}(t))\check{\rho}(t)^{-1}$. Therefore the residue of this oper in

 \mathfrak{h}/W is equal to $\varpi(-\check{\rho})$, and so $\operatorname{Op}_G(D_x) = \operatorname{Op}_G^{\mathrm{RS}}(D_x)_0$. For $G = PGL_2$ we identify \mathfrak{h} with \mathbb{C} and so $\check{\lambda}$ with a complex number. Then one finds that $\operatorname{Op}_{PGL_2}^{\mathrm{RS}}(D_x)_{\check{\lambda}}$ is the space of second order operators of the form

(1.7)
$$\partial_t^2 - \frac{\lambda(\lambda+2)/4}{t^2} - \sum_{n \ge -1} v_n t^n$$

Now suppose that λ is a dominant integral coweight of \mathfrak{g} . Following Drinfeld, introduce the variety $\operatorname{Op}_G(D_x)_{\check{\lambda}}$ as the quotient of the space of operators of the form

(1.8)
$$\nabla = \partial_t + \sum_{i=1}^{t} \psi_i(t) f_i + \mathbf{v}(t),$$

where

$$\psi_i(t) = t^{\langle \alpha_i, \lambda \rangle}(\kappa_i + t(\dots)) \in \mathcal{O}_x, \qquad \kappa_i \neq 0$$

and $\mathbf{v}(t) \in \mathfrak{b}(\mathcal{O}_x)$, by the gauge action of $B(\mathcal{O}_x)$. Equivalently, $\operatorname{Op}_G(D_x)_{\lambda}$ is the quotient of the space of operators of the form

(1.9)
$$\nabla = \partial_t + \sum_{i=1}^{\ell} t^{\langle \alpha_i, \check{\lambda} \rangle} f_i + \mathbf{v}(t)$$

where $\mathbf{v}(t) \in \mathfrak{b}(\mathcal{O}_x)$, by the gauge action of $N(\mathcal{O}_x)$. Considering the $N(\mathcal{K}_x)$ -class of such an operator, we obtain an oper on D_x^{\times} . Thus, we have a map $\operatorname{Op}_G(D_x)_{\check{\lambda}} \to$ $\operatorname{Op}_G(D_x^{\times}).$

10

Lemma 1.2 ([F3], Lemma 2.4). The map $\operatorname{Op}_G(D_x)_{\check{\lambda}} \to \operatorname{Op}_G(D_x^{\times})$ is injective and its image is contained in the subvariety $\operatorname{Op}_G^{\mathrm{RS}}(D_x)_{\check{\lambda}}$. Moreover, the points of $\operatorname{Op}_G(D_x)_{\check{\lambda}}$ are precisely those G-opers with regular singularity and residue $\check{\lambda}$ which have no monodromy around x.

The space $\operatorname{Op}_{PGL_2}(D_x)_{\check{\lambda}}$ is the subspace of codimension one in $\operatorname{Op}_{PGL_2}(D_x)_{\check{\lambda}}^{\mathrm{RS}}$. In terms of the coefficients $v_n, n \geq -1$, appearing in formula (1.7) the corresponding equation has the form $P_{\lambda}(v_n) = 0$, where P_{λ} is a polynomial of degree $\lambda + 1$, where we set deg $v_n = n + 2$. For example, $P_0 = v_{-1}, P_2 = 2v_{-1}^2 - v_0$, etc. In general, the subspace $\operatorname{Op}_G(D_x)_{\check{\lambda}} \subset \operatorname{Op}_G^{\mathrm{RS}}(D_x)_{\check{\lambda}}$ is defined by dim N polynomial equations, where N is the unipotent subgroup of G.

2. The Gaudin model

Let \mathfrak{g} be a simple Lie algebra. The *Langlands dual Lie algebra* ${}^{L}\mathfrak{g}$ is by definition the Lie algebra whose Cartan matrix is the transpose of that of \mathfrak{g} . We will identify the set of roots of \mathfrak{g} with the set of coroots of ${}^{L}\mathfrak{g}$ and the set of weights of \mathfrak{g} with the set of coweights of ${}^{L}\mathfrak{g}$. The results on opers from the previous sections will be applied here to the Lie algebra ${}^{L}\mathfrak{g}$. Thus, in particular, ${}^{L}G$ will denote the adjoint group of ${}^{L}\mathfrak{g}$.

2.1. The definition of the Gaudin model

Here we recall the definition of the Gaudin model and the realization of the Gaudin Hamiltonians in terms of the spaces of conformal blocks for affine Kac-Moody algebras of critical level. We follow closely the paper [FFR].

Choose a non-degenerate invariant inner product κ_0 on \mathfrak{g} . Let $\{J_a\}, a = 1, \ldots, d$, be a basis of \mathfrak{g} and $\{J^a\}$ the dual basis with respect to κ_0 . Denote by Δ the quadratic Casimir operator from the center of $U(\mathfrak{g})$:

$$\Delta = \frac{1}{2} \sum_{a=1}^{d} J_a J^a.$$

The Gaudin Hamiltonians are the elements

(2.1)
$$\Xi_i = \sum_{j \neq i} \sum_{a=1}^d \frac{J_a^{(i)} J^{a(j)}}{z_i - z_j}, \qquad i = 1, \dots, N,$$

of the algebra $U(\mathfrak{g})^{\otimes N}$. Note that they commute with the diagonal action of \mathfrak{g} on $U(\mathfrak{g})^{\otimes N}$ and that

$$\sum_{i=1}^{N} \Xi_i = 0.$$

2.2. Gaudin model and coinvariants

Let $\widehat{\mathfrak{g}}_{\kappa_c}$ be the affine Kac-Moody algebra corresponding to \mathfrak{g} . It is the extension of the Lie algebra $\mathfrak{g} \otimes \mathbb{C}((t))$ by the one-dimensional center $\mathbb{C}K$. The commutation relations in $\widehat{\mathfrak{g}}_{\kappa_c}$ read

(2.2)
$$[A \otimes f(t), B \otimes g(t)] = [A, B] \otimes fg - \kappa_c(A, B) \operatorname{Res}_{t=0} fdg \cdot K$$

where κ_c is the *critical* invariant inner product on \mathfrak{g} defined by the formula

$$\kappa_c(A, B) = -\frac{1}{2} \operatorname{Tr}_{\mathfrak{g}} \operatorname{ad} A \operatorname{ad} B.$$

Note that $\kappa_c = -h^{\vee}\kappa_0$, where κ_0 is the inner product normalized as in [K] and h^{\vee} is the dual Coxeter number.

Denote by $\widehat{\mathfrak{g}}_+$ the Lie subalgebra $\mathfrak{g} \otimes \mathbb{C}[[t]] \oplus \mathbb{C}K$ of $\widehat{\mathfrak{g}}_{\kappa_c}$. Let M be a \mathfrak{g} -module. We extend the action of \mathfrak{g} on M to $\mathfrak{g} \otimes \mathbb{C}[[t]]$ by using the evaluation at zero homomorphism $\mathfrak{g} \otimes \mathbb{C}[[t]] \to \mathfrak{g}$ and to $\widehat{\mathfrak{g}}_+$ by making K act as the identity. Denote by \mathbb{M} the corresponding induced $\widehat{\mathfrak{g}}_{\kappa_c}$ -module

$$\mathbb{M} = U(\widehat{\mathfrak{g}}_{\kappa_c}) \underset{U(\widehat{\mathfrak{g}}_+)}{\otimes} M.$$

By construction, K acts as the identity on this module. We call such modules the $\hat{\mathfrak{g}}_{\kappa_c}$ -modules of *critical level*. For example, for $\lambda \in \mathfrak{h}^*$ let \mathbb{C}_{λ} be the one-dimensional \mathfrak{b} -module on which \mathfrak{h} acts by the character $\lambda : \mathfrak{h} \to \mathbb{C}$ and \mathfrak{n} acts by 0. Let M_{λ} be the Verma module over \mathfrak{g} of highest weight λ ,

$$M_{\lambda} = U(\mathfrak{g}) U(\mathfrak{b}) \mathbb{C}_{\lambda}.$$

The corresponding induced module \mathbb{M}_{λ} is the Verma module over $\widehat{\mathfrak{g}}_{\kappa_c}$ with highest weight λ .

For a dominant integral weight $\lambda \in \mathfrak{h}^*$ denote by V_{λ} the irreducible finitedimensional \mathfrak{g} -module of highest weight λ . The corresponding induced module \mathbb{V}_{λ} is called the Weyl module over $\widehat{\mathfrak{g}}_{\kappa_c}$ with highest weight λ . Consider the projective line \mathbb{P}^1 with a global coordinate t and N distinct

Consider the projective line \mathbb{P}^1 with a global coordinate t and N distinct finite points $z_1, \ldots, z_N \in \mathbb{P}^1$. In the neighborhood of each point z_i we have the local coordinate $t - z_i$ and in the neighborhood of the point ∞ we have the local coordinate t^{-1} . Set $\tilde{\mathfrak{g}}(z_i) = \mathfrak{g} \otimes \mathbb{C}((t - z_i))$ and $\tilde{\mathfrak{g}}(\infty) = \mathfrak{g} \otimes \mathbb{C}((t^{-1}))$. Let $\hat{\mathfrak{g}}_N$ be the extension of the Lie algebra $\bigoplus_{i=1}^N \tilde{\mathfrak{g}}(z_i) \oplus \tilde{\mathfrak{g}}(\infty)$ by a one-dimensional center $\mathbb{C}K$ whose restriction to each summand $\tilde{\mathfrak{g}}(z_i)$ or $\tilde{\mathfrak{g}}(\infty)$ coincides with the above central extension.

Suppose we are given a collection M_1, \ldots, M_N and M_{∞} of \mathfrak{g} -modules. Then the Lie algebra $\widehat{\mathfrak{g}}_N$ naturally acts on the tensor product $\bigotimes_{i=1}^N \mathbb{M}_i \otimes \mathbb{M}_{\infty}$ (in particular, K acts as the identity).

Let $\mathfrak{g}_{(z_i)} = \mathfrak{g}_{z_1,\ldots,z_N}$ be the Lie algebra of \mathfrak{g} -valued regular functions on $\mathbb{P}^1 \setminus \{z_1,\ldots,z_N,\infty\}$ (i.e., rational functions on \mathbb{P}^1 , which may have poles only at the points z_1,\ldots,z_N and ∞). Clearly, such a function can be expanded into a

Laurent power series in the corresponding local coordinates at each point z_i and at ∞ . Thus, we obtain an embedding

$$\mathfrak{g}_{(z_i)} \hookrightarrow \bigoplus_{i=1}^N \widetilde{\mathfrak{g}}(z_i) \oplus \widetilde{\mathfrak{g}}(\infty).$$

It follows from the residue theorem and formula (2.2) that the restriction of the central extension to the image of this embedding is trivial. Hence this embedding lifts to the embedding $\mathfrak{g}_{(z_i)} \to \widehat{\mathfrak{g}}_N$.

Denote by $H(M_1, \ldots, M_N, M_\infty)$ the space of coinvariants of $\bigotimes_{i=1}^N \mathbb{M}_i \otimes \mathbb{M}_\infty$ with respect to the action of the Lie algebra $\mathfrak{g}_{(z_i)}$. By construction, we have a canonical embedding of a \mathfrak{g} -module M into the induced $\widehat{\mathfrak{g}}_{\kappa_c}$ -module \mathbb{M} :

$$x \in M \to 1 \otimes x \in \mathbb{M},$$

which commutes with the action of \mathfrak{g} on both spaces (where \mathfrak{g} is embedded into $\widehat{\mathfrak{g}}_{\kappa_c}$ as the constant subalgebra). Thus we have an embedding

$$\bigotimes_{i=1}^N M_i \otimes M_\infty \hookrightarrow \bigotimes_{i=1}^N \mathbb{M}_i \otimes \mathbb{M}_\infty$$

The following result is proved in the same way as Lemma 1 in [FFR].

Lemma 2.1. The composition of this embedding and the projection

$$\bigotimes_{i=1}^{N} \mathbb{M}_{i} \otimes \mathbb{M}_{\infty} \twoheadrightarrow H(M_{1}, \ldots, M_{N}, M_{\infty})$$

gives rise to an isomorphism

$$H(M_1,\ldots,M_N,M_\infty)\simeq (\bigotimes_{i=1}^N M_i\otimes M_\infty)/\mathfrak{g}_{\mathrm{diag}}$$

Let \mathbb{V}_0 be the induced $\widehat{\mathfrak{g}}_{\kappa_c}$ -module of critical level, which corresponds to the one-dimensional trivial \mathfrak{g} -module V_0 ; it is called the *vacuum module*. Denote by v_0 the generating vector of \mathbb{V}_0 . We assign the vacuum module to a point $u \in \mathbb{P}^1$ which is different from z_1, \ldots, z_N, ∞ . Denote by $H(M_1, \ldots, M_N, M_\infty, \mathbb{C})$ the space of $\mathfrak{g}_{(z_i),u}$ -invariant functionals on $\bigotimes_{i=1}^N \mathbb{M}_i \otimes \mathbb{M}_\infty \otimes \mathbb{V}_0$ with respect to the Lie algebra $\mathfrak{g}_{(z_i),u}$. Lemma 2.1 tells us that we have a canonical isomorphism

$$H(M_1,\ldots,M_N,M_\infty,\mathbb{C})\simeq H(M_1,\ldots,M_N,M_\infty).$$

Now observe that by functoriality any endomorphism $X \in \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{V}_0$ gives rise to an endomorphism of the space of coinvariants $H(M_1, \ldots, M_N, M_\infty, \mathbb{C})$, and hence of $H(M_1, \ldots, M_N, M_\infty)$. Thus, we obtain a homomorphism of algebras

 $\operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_0}} \mathbb{V}_0 \to \operatorname{End}_{\mathbb{C}} H(M_1, \ldots, M_N, M_\infty).$

Let us compute this homomorphism explicitly.

First of all, we identify the algebra $\operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{V}_0$ with the space

$$\mathfrak{z}(\widehat{\mathfrak{g}}) = \mathbb{V}_0^{\mathfrak{g}[[t]]}$$

of $\mathfrak{g}[[t]]$ -invariant vectors in \mathbb{V}_0 . Indeed, a $\mathfrak{g}[[t]]$ -invariant vector v gives rise to an endomorphism of \mathbb{V}_0 commuting with the action of $\widehat{\mathfrak{g}}_{\kappa_c}$ which sends the generating vector v_0 to v. Conversely, any $\widehat{\mathfrak{g}}_{\kappa_c}$ -endomorphism of V_0 is uniquely determined by the image of v_0 which necessarily belongs to $\mathfrak{z}(\widehat{\mathfrak{g}})$. Thus, we obtain an isomorphism $\mathfrak{z}(\widehat{\mathfrak{g}}) \simeq \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}}(\mathbb{V}_0)$ which gives $\mathfrak{z}(\widehat{\mathfrak{g}})$ an algebra structure. The opposite algebra structure on $\mathfrak{z}(\widehat{\mathfrak{g}})$ coincides with the algebra structure induced by the identification of \mathbb{V}_0 with the algebra $U(\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}])$. But we will see in the next section that the algebra $\mathfrak{z}(\widehat{\mathfrak{g}})$ is commutative and so the two algebra structures on it coincide.

Now let $v \in \mathfrak{z}(\widehat{\mathfrak{g}}) \subset \mathbb{V}_0$. For any

$$x \in (\bigotimes_{i=1}^{N} M_i \otimes M_\infty) / \mathfrak{g}_{\operatorname{diag}} \simeq H(M_1, \dots, M_N, M_\infty)$$

take a lifting \widetilde{x} to $\bigotimes_{i=1}^{N} M_i \otimes M_{\infty}$. By Lemma 2.1, the projection of the vector

$$\widetilde{x} \otimes v \in \bigotimes_{i=1}^{N} \mathbb{M}_{i} \otimes \mathbb{M}_{\infty} \otimes \mathbb{V}_{0}$$

onto

$$H(M_1,\ldots,M_N,M_\infty,\mathbb{C})\simeq (\bigotimes_{i=1}^N M_i\otimes M_\infty)/\mathfrak{g}_{\mathrm{diag}}$$

is equal to the projection of a vector of the form $(\Psi_u(v) \cdot \tilde{x}) \otimes v_0$, where

$$\Psi_u(v) \cdot \widetilde{x} \in (\bigotimes_{i=1}^N M_i \otimes M_\infty)/\mathfrak{g}_{\text{diag}}.$$

For $A \in \mathfrak{g}$ and $n \in \mathbb{Z}$, denote by A_n the element $A \otimes t^n \in \widehat{\mathfrak{g}}_{\kappa_c}$. Then $\mathbb{V}_0 \simeq U(\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}])v_0$ has a basis of lexicographically ordered monomials of the form $J_{n_1}^{a_1} \ldots J_{n_m}^{a_m}v_0$ with $n_i < 0$. Let us set

$$\mathbb{J}_{n}^{a}(u) = -\sum_{i=1}^{N} \frac{J^{a(i)}}{(z_{i}-u)^{n}}.$$

Define an anti-homomorphism

$$\Phi_u: U(\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}]) \to U(\mathfrak{g})^{\otimes N} \otimes \mathbb{C}[(u-z_i)^{-1}]_{i=1,\dots,N}$$

by the formula

(2.3)
$$\Phi_u(J_{n_1}^{a_1}\dots J_{n_m}^{a_m}v_0) = \mathbb{J}_{n_m}^{a_m}(u)\mathbb{J}_{n_{m-1}}^{a_{m-1}}(u)\dots \mathbb{J}_{n_1}^{a_1}(u).$$

According to the computation presented in the proof of Prop. 1 of [FFR], we have

$$\Psi_u(v) \cdot \widetilde{x} = \Phi_u(v) \cdot \widetilde{x}.$$

14

In general, $\Phi_u(v)$ does not commute with the diagonal action of \mathfrak{g} , and so $\Psi_v(u) \cdot \widetilde{x}$ depends on the choice of the lifting \widetilde{x} . But if $v \in \mathfrak{z}(\widehat{\mathfrak{g}}) \subset \mathbb{V}_0$, then $\Phi_u(v)$ commutes with the diagonal action of \mathfrak{g} and hence gives rise to a well-defined endomorphism $(\bigotimes_{i=1}^N M_i \otimes M_\infty)/\mathfrak{g}_{\text{diag}}$.

Thus, we restrict Φ_u to $\mathfrak{z}(\hat{\mathfrak{g}})$. This gives us a homomorphism of algebras

$$\mathfrak{z}(\widehat{\mathfrak{g}}) \to (U(\mathfrak{g})^{\otimes N})^G \otimes \mathbb{C}[(u-z_i)^{-1}]_{i=1,\ldots,N},$$

which we also denote by Φ_u .

For example, consider the Segal-Sugawara vector in \mathbb{V}_0 :

(2.4)
$$S = \frac{1}{2} \sum_{a=1}^{d} J_{a,-1} J_{-1}^{a} v_0.$$

One shows (see, e.g., [FB]) that this vector belongs to $\mathfrak{z}(\hat{\mathfrak{g}})$. Consider the corresponding element $\Phi_u(S)$.

Denote by Δ the Casimir operator $\frac{1}{2} \sum_{a} J_{a} J^{a}$ from $U(\mathfrak{g})$.

Proposition 2.2 ([FFR], Prop. 1). We have

$$\Phi_u(S) = \sum_{i=1}^N \frac{\Xi_i}{u - z_i} + \sum_{i=1}^N \frac{\Delta^{(i)}}{(u - z_i)^2}$$

where the Ξ_i 's are the Gaudin operators (2.1).

We wish to study the algebra generated by the image of the map Φ_u .

Proposition 2.3 ([FFR], Prop. 2). For any $Z_1, Z_2 \in \mathfrak{g}(\widehat{\mathfrak{g}})$ and any points $u_1, u_2 \in \mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$ the linear operators $\Psi_{Z_1}(u_1)$ and $\Psi_{Z_2}(u_2)$ commute.

Let $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ be the span in $U(\mathfrak{g})^{\otimes N}$ of the coefficients in front of the monomials $\prod_{i=1}^{N} (u-z_i)^{n_i}$ of the series $\Phi_u(v), v \in \mathfrak{z}(\widehat{\mathfrak{g}})$. Since Φ_u is an algebra homomorphism, we find that $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ is a subalgebra of $(U(\mathfrak{g})^{\otimes N})^G$, which is commutative by Proposition 2.3. We call it the *Gaudin algebra* associated to \mathfrak{g} and the collection z_1, \ldots, z_N , and its elements the *generalized Gaudin Hamiltonians*.

2.3. The center of \mathbb{V}_0 and LG -opers

In order to describe the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ and its spectrum we need to recall the description of $\mathfrak{z}(\widehat{\mathfrak{g}})$. According to [FF2, F2], $\mathfrak{z}(\widehat{\mathfrak{g}})$ is identified with the algebra Fun $\operatorname{Op}_{L_G}(D)$ of (regular) functions on the space $\operatorname{Op}_{L_G}(D)$ of ^LG-opers on the disc $D = \operatorname{Spec} \mathbb{C}[[t]]$, where ^LG is the Langlands dual group to G. Since we have assumed that G is simply-connected, ^LG may be defined as the adjoint group of the Lie algebra ^L \mathfrak{g} whose Cartan matrix is the transpose of that of \mathfrak{g} .

This isomorphism satisfies various properties, one of which we will now recall. Let $\text{Der } \mathcal{O} = \mathbb{C}[[t]]\partial_t$ be the Lie algebra of continuous derivations of the topological algebra $\mathcal{O} = \mathbb{C}[[t]]$. The action of its Lie subalgebra $\text{Der}_0 \mathcal{O} = t\mathbb{C}[[t]]\partial_t$ on \mathcal{O} exponentiates to an action of the group Aut \mathcal{O} of formal changes of variables. Both Der O and Aut O naturally act on V_0 in a compatible way, and these actions preserve $\mathfrak{z}(\hat{\mathfrak{g}})$. They also act on the space $\operatorname{Op}_{L_G}(D)$.

Denote by Fun $\operatorname{Op}_{L_G}(D)$ the algebra of regular functions on $\operatorname{Op}_{L_G}(D)$. In view of Lemma 1.1, it is isomorphic to the algebra of functions on the space of ℓ -tuples $(v_1(t), \ldots, v_\ell(t))$ of formal Taylor series, i.e., the space $\mathbb{C}[[t]]^{\ell}$. If we write $v_i(t) = \sum_{n>0} v_{i,n}t^n$, then we obtain

(2.5)
$$\operatorname{Fun}\operatorname{Op}_{L_G}(D) \simeq \mathbb{C}[v_{i,n}]_{i \in I, n > 0}.$$

Note that the vector field $-t\partial_t$ acts naturally on $\operatorname{Op}_{L_G}(D)$ and defines a \mathbb{Z} -grading on Fun $\operatorname{Op}_{L_G}(D)$ such that $\operatorname{deg} v_{i,n} = d_i + n + 1$. The vector field $-\partial_t$ acts as a derivation such that $-\partial_t \cdot v_{i,n} = -(d_i + n + 1)v_{i,n+1}$.

Theorem 2.4 ([FF2, F2]). There is a canonical isomorphism

$$\mathfrak{z}(\widehat{\mathfrak{g}}) \simeq \operatorname{Fun} \operatorname{Op}_{L_G}(D)$$

of algebras which is compatible with the action of Der O and Aut O.

We use this result to describe the twist of $\mathfrak{z}(\hat{\mathfrak{g}})$ by the Aut O-torsor Aut_x of formal coordinates at a smooth point x of an algebraic curve X,

$$\mathfrak{z}(\widehat{\mathfrak{g}})_x = \mathcal{A}ut_x \underset{\text{Aut (f)}}{\times} \mathfrak{z}(\widehat{\mathfrak{g}})$$

(see Ch. 6 of [FB] for more details). It follows from the definition that the corresponding twist of Fun $\operatorname{Op}_{L_G}(D)$ by Aut_x is nothing but Fun $\operatorname{Op}_{L_G}(D_x)$, where D_x is the disc around x. Therefore we obtain from Theorem 2.4 an isomorphism

(2.6)
$$\mathfrak{z}(\widehat{\mathfrak{g}})_x \simeq \operatorname{Fun} \operatorname{Op}_{L_G}(D_x).$$

The module \mathbb{V}_0 has a natural \mathbb{Z} -grading defined by the formulas deg $v_0 = 0$, deg $J_n^a = -n$, and it carries a translation operator T defined by the formulas $Tv_0 = 0, [T, J_n^a] = -nJ_{n-1}^a$. Theorem 2.4 and the isomorphism (2.5) imply that there exist non-zero vectors $S_i \in \mathbb{V}_0^{\mathfrak{g}[[t]]}$ of degrees $d_i + 1, i \in I$, such that

$$\mathfrak{z}(\widehat{\mathfrak{g}}) = \mathbb{C}[T^n S_i]_{i \in I, n > 0} v_0.$$

Then under the isomorphism of Theorem 2.4 we have $S_i \mapsto v_{i,0}$, the \mathbb{Z} -gradings on both algebras get identified and the action of T on $\mathfrak{z}(\hat{\mathfrak{g}})$ becomes the action of $-\partial_t$ on Fun $\operatorname{Op}_{L_G}(D)$. Note that the vector S_1 is nothing but the vector (2.4), up to a non-zero scalar.

Recall from [FB] that \mathbb{V}_0 is a vertex algebra, and $\mathfrak{z}(\hat{\mathfrak{g}})$ is its commutative vertex subalgebra; in fact, it is the center of \mathbb{V}_0 . We will also need the center of the completed universal enveloping algebra of $\hat{\mathfrak{g}}$ of critical level. This algebra is defined as follows.

Let $U_{\kappa_c}(\widehat{\mathfrak{g}})$ be the quotient of the universal enveloping algebra $U(\widehat{\mathfrak{g}}_{\kappa_c})$ of $\widehat{\mathfrak{g}}_{\kappa_c}$ by the ideal generated by (K-1). Define its completion $\widetilde{U}_{\kappa_c}(\widehat{\mathfrak{g}})$ as follows:

$$U_{\kappa_c}(\widehat{\mathfrak{g}}) = \lim U_{\kappa_c}(\widehat{\mathfrak{g}})/U_{\kappa_c}(\widehat{\mathfrak{g}}) \cdot (\mathfrak{g} \otimes t^N \mathbb{C}[[t]]).$$

It is clear that $\widetilde{U}_{\kappa_c}(\widehat{\mathfrak{g}})$ is a topological algebra which acts on all smooth $\widehat{\mathfrak{g}}_{\kappa_c}$ -module. By definition, a smooth $\widehat{\mathfrak{g}}_{\kappa_c}$ -module is a $\widehat{\mathfrak{g}}_{\kappa_c}$ -module such that any vector is annihilated by $\mathfrak{g} \otimes t^N \mathbb{C}[[t]]$ for sufficiently large N, and K acts as the identity.

Let $Z(\widehat{\mathfrak{g}})$ be the center of $\widetilde{U}_{\kappa_c}(\widehat{\mathfrak{g}})$.

Denote by Fun $\operatorname{Op}_{L_G}(D^{\times})$ the algebra of regular functions on the space $\operatorname{Op}_{L_G}(D^{\times})$ of ^LG-opers on the punctured disc $D^{\times} = \operatorname{Spec} \mathbb{C}((t))$. In view of Lemma 1.1, it is isomorphic to the algebra of functions on the space of ℓ -tuples $(v_1(t), \ldots, v_{\ell}(t))$ of formal Laurent series, i.e., the ind-affine space $\mathbb{C}((t))^{\ell}$. If we write $v_i(t) = \sum_{n \in \mathbb{Z}} v_{i,n} t^n$, then we obtain that Fun $\operatorname{Op}_{L_G}(D)$ is isomorphic to the completion of the polynomial algebra $\mathbb{C}[v_{i,n}]_{i \in I, n \in \mathbb{Z}}$ with respect to the topology in which the basis of open neighborhoods of zero is formed by the ideals generated by $v_{i,n}, i \in I, n \leq N$, for $N \leq 0$.

Theorem 2.5 ([F2]). There is a canonical isomorphism

$$Z(\widehat{\mathfrak{g}}) \simeq \operatorname{Fun} \operatorname{Op}_{{}^{L}G}(D^{\times})$$

of complete topological algebras which is compatible with the action of Der O and Aut O.

If M is a smooth $\widehat{\mathfrak{g}}_{\kappa_c}$ -module, then the action of $Z(\widehat{\mathfrak{g}})$ on M gives rise to a homomorphism

$$Z(\widehat{\mathfrak{g}}) \to \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa}} M.$$

For example, if $M = \mathbb{V}_0$, then using Theorems 2.4 and 2.5 we identify this homomorphism with the surjection

Fun
$$\operatorname{Op}_{L_G}(D^{\times}) \twoheadrightarrow \operatorname{Fun} \operatorname{Op}_{L_G}(D)$$

induced by the natural embedding

$$\operatorname{Op}_{L_G}(D) \hookrightarrow \operatorname{Op}_{L_G}(D^{\times}).$$

Recall that the Harish-Chandra homomorphism identifies the center $Z(\mathfrak{g})$ of $U(\mathfrak{g})$ with the algebra $(\operatorname{Fun}\mathfrak{h}^*)^W$ of polynomials on \mathfrak{h}^* which are invariant with respect to the action of the Weyl group W. Therefore a character $Z(\mathfrak{g}) \to \mathbb{C}$ is the same as a point in Spec(Fun $\mathfrak{h}^*)^W$ which is the quotient \mathfrak{h}^*/W . For $\lambda \in \mathfrak{h}^*$ we denote by $\varpi(\lambda)$ its projection onto \mathfrak{h}^*/W . In particular, $Z(\mathfrak{g})$ acts on M_{λ} and V_{λ} via its character $\varphi(\lambda + \rho)$. We also denote by I_{λ} the maximal ideal of $Z(\mathfrak{g})$ equal to the kernel of the homomorphism $Z(\mathfrak{g}) \to \mathbb{C}$ corresponding to the character $\varphi(\lambda + \rho)$.

In what follows we will use the canonical identification between \mathfrak{h}^* and the Cartan subalgebra ${}^{L}\mathfrak{h}$ of the Langlands dual Lie algebra ${}^{L}\mathfrak{g}$. Recall that in Section 1.3 we defined the space $\operatorname{Op}_{L_G}^{\mathrm{RS}}(D)$ of ${}^{L}G$ -opers on D^{\times} with regular singularity and its subspace $\operatorname{Op}_{L_G}^{\mathrm{RS}}(D)_{\lambda}$ of opers with residue $\varpi(-\lambda - \rho)$. We also defined the subspace

$$\operatorname{Op}_{{}^{L}G}(D)_{\check{\lambda}} \subset \operatorname{Op}_{{}^{L}G}^{\operatorname{RS}}(D)_{\check{\lambda}} \subset \operatorname{Op}_{{}^{L}G}(D^{\times})$$

of those ${}^{L}G$ -opers which have trivial monodromy. Here we identify the coweights of the group ${}^{L}G$ with the weights of G.

The following result is obtained by combining Theorem 12.4, Lemma 9.4 and Proposition 12.8 of [F2].

Theorem 2.6.

(1) Let \mathbb{U} be the $\widehat{\mathfrak{g}}_{\kappa_c}$ -module induced from the $\mathfrak{g}[[t]] \oplus \mathbb{C}K$ -module $U(\mathfrak{g})$. Then the homomorphism $Z(\widehat{\mathfrak{g}}) \to \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{U}$ factors as

$$Z(\widehat{\mathfrak{g}}) \simeq \operatorname{Fun}\operatorname{Op}_{L_G}(D^{\times}) \twoheadrightarrow \operatorname{Fun}\operatorname{Op}_{L_G}^{\operatorname{RS}}(D) \to \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{U}.$$

(2) Let M be a \mathfrak{g} -module on which the center $Z(\mathfrak{g})$ acts via its character $\varpi(\lambda + \rho)$, and let \mathbb{M} be the induced $\widehat{\mathfrak{g}}_{\kappa_c}$ -module. Then the homomorphism $Z(\widehat{\mathfrak{g}}) \to \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{M}$ factors as follows

$$Z(\widehat{\mathfrak{g}}) \simeq \operatorname{Fun} \operatorname{Op}_{L_G}(D^{\times}) \twoheadrightarrow \operatorname{Fun} \operatorname{Op}_{L_G}^{\operatorname{RS}}(D)_{\lambda} \to \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{M}$$

Moreover, if $M = M_{\lambda}$, then the last map is an isomorphism

$$\operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_{\alpha}}} \mathbb{M} \simeq \operatorname{Fun} \operatorname{Op}_{L_{G}}^{\operatorname{RS}}(D)_{\lambda}.$$

(3) For an integral dominant weight $\lambda \in \mathfrak{h}^*$ the homomorphism

$$\operatorname{Fun}\operatorname{Op}_{{}^{L}G}(D^{\times})\to\operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_{c}}}\mathbb{V}_{\lambda}$$

factors as

$$\operatorname{Fun}\operatorname{Op}_{{}^{L}G}(D^{\times})\to\operatorname{Fun}\operatorname{Op}_{{}^{L}G}(D)_{\lambda}\to\operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_{c}}}\mathbb{V}_{\lambda},$$

and the last map is an isomorphism

$$\operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_{c}}} \mathbb{V}_{\lambda} \simeq \operatorname{Fun} \operatorname{Op}_{L_{G}}(D)_{\lambda}$$

2.4. Example

Let us consider the case $\mathfrak{g} = \mathfrak{sl}_2$ in more detail. Introduce the Segal-Sugawara operators $S_n, n \in \mathbb{Z}$, by the formula

$$S(z) = \sum_{n \in \mathbb{Z}} S_n z^{-n-2} = \frac{1}{2} \sum_a : J^a(z) J_a(z):,$$

where the normal ordering is defined as in [FB]. Then the center $Z(\mathfrak{sl}_2)$ is the completion $\mathbb{C}[S_n]_{n\in\mathbb{Z}}^{\sim}$ of the polynomial algebra $\mathbb{C}[S_n]_{n\in\mathbb{Z}}$ with respect to the topology in which the basis of open neighborhoods of zero is formed by the ideals of $S_n, n > N$, for $N \ge 0$.

We have the following diagram of (vertical) isomorphisms and (horizontal) surjections

where J_{λ} is the ideal generated by $(S_0 - \frac{1}{4}\lambda(\lambda + 2))$ and J'_{λ} is the ideal generated by I_{λ} and the polynomial P_{λ} introduced at the end of Section 1.3.

The space of PGL_2 -opers on D^{\times} is identified with the space of projective connections of the form $\partial_t^2 - \sum_{n \in \mathbb{Z}} v_n t^n$. The isomorphism of Theorem 2.5 sends

 S_n to v_{-n-2} . The relevant spaces of PGL_2 -opers with regular singularities were described at the end of Section 1.3, and these descriptions agree with the above diagram and Theorem 2.6.

2.5. The Gaudin algebra

Now we are ready to identify the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ with the algebra of functions on a certain space of opers on \mathbb{P}^1 .

Let $\operatorname{Op}_{L_G}^{\operatorname{RS}}(\mathbb{P}^1)_{(z_i),\infty}$ be the space of LG -opers on \mathbb{P}^1 with regular singularities at z_1, \ldots, z_N and ∞ . For an arbitrary collection of weights $\lambda_1, \ldots, \lambda_N$ and λ_{∞} , let

$$\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\circ}}$$

be its subspace of those opers whose residue at the point z_i (resp., ∞) is equal to $\varpi(-\lambda_i - \rho), i = 1, \ldots, N$ (resp., $\varpi(-\lambda_{\infty} - \rho)$). Finally, if all of the weights $\lambda_1, \ldots, \lambda_N, \lambda_{\infty}$ are dominant integral, we introduce a subset

$$\operatorname{Op}_{{}^{L}G}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}} \subset \operatorname{Op}_{{}^{L}G}^{\operatorname{RS}}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}}$$

which consists of those ${}^{L}G$ -opers which have trivial monodromy representation.

On the other hand, for each collection of points z_1, \ldots, z_N on $\mathbb{P}^1 \setminus \infty$ we have the Gaudin algebra

$$\mathcal{Z}_{(z_i)}(\mathfrak{g}) \subset \left(U(\mathfrak{g})^{\otimes N} \right)^G \simeq \left(U(\mathfrak{g})^{\otimes (N+1)} / \mathfrak{g}_{\text{diag}} \right)^G,$$

where the second isomorphism is obtained by identifying $U(\mathfrak{g})^{\otimes (N+1)}/\mathfrak{g}_{\text{diag}}$ with $U(\mathfrak{g})^{\otimes N} \otimes 1$.

We have a homomorphism $c_i : Z(\mathfrak{g}) \to U(\mathfrak{g}) \to U(\mathfrak{g})^{\otimes (N+1)}$ corresponding to the *i*th factor, for all $i = 1, \ldots, N$, and a homomorphism $c_{\infty} : Z(\mathfrak{g}) \to U(\mathfrak{g})^{\otimes (N+1)}$ corresponding to the (N+1)st factor. It is easy to see that the images of $c_i, i = 1, \ldots, N$, and c_{∞} belong to $\mathfrak{Z}_{(z_i)}(\mathfrak{g})$. For a collection of weights $\lambda_1, \ldots, \lambda_N$ and λ_{∞} , let $I_{(\lambda_i),\lambda_{\infty}}$ be the ideal of $\mathfrak{Z}_{(z_i)}(\mathfrak{g})$ generated by $c_i(I_{\lambda_i}), i = 1, \ldots, N$, and $c_{\infty}(I_{\lambda_{\infty}})$. Let $\mathfrak{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ be the quotient of $\mathfrak{Z}_{(z_i)}(\mathfrak{g})$ by $I_{(\lambda_i),\lambda_{\infty}}$.

The algebra $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$ acts on the space of \mathfrak{g} -coinvariants in

$$\bigotimes_{i=1}^N M_i \otimes M_\infty$$

where M_i is a g-module with central character $\varpi(\lambda_i + \rho), i = 1, \ldots, N$, and M_{∞} is a g-module with central character $\varpi(\lambda_{\infty} + \rho)$. In particular, if all the weights $\lambda_1, \ldots, \lambda_N, \lambda_{\infty}$ are dominant integral, then we can take as the M_i 's the finitedimensional irreducible modules V_{λ_i} for $i = 1, \ldots, N$, and as M_{∞} the module $V_{\lambda_{\infty}}$. The corresponding space of g-coinvariants is isomorphic to the space

$$\left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_\infty}\right)^G$$

of *G*-invariants in $\bigotimes_{i=1}^{N} V_{\lambda_i} \otimes V_{\lambda_{\infty}}$. Let $\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$ be the image of the algebra $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ in End $\left(\bigotimes_{i=1}^{N} V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)^G$.

We have the following result.

Theorem 2.7.

(1) The algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ is isomorphic to the algebra of functions on the space $\operatorname{Op}_{L_C}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty}$.

(2) The algebra $\mathfrak{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$ is isomorphic to the algebra of functions on the space $\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$.

(3) For a collection of dominant integral weights $\lambda_1, \ldots, \lambda_N, \lambda_\infty$, there is a surjective homomorphism from the algebra of functions $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_\infty}$ to the algebra $\overline{\mathbb{Z}}_{(z_i),\infty;(\lambda_i),\lambda_\infty}(\mathfrak{g})$.

Proof. In [FB] we defined, for any quasi-conformal vertex algebra V, a smooth projective curve X, a set of points $x_1, \ldots, x_N \in X$ and a collection of V-modules M_1, \ldots, M_N , the space of coinvariants $H_V(X, (x_i), (M_i))$, which is the quotient of $\bigotimes_{i=1}^N M_i$ by the action of a certain Lie algebra. This construction (which is recalled in the proof of Theorem 4.7 in [F3]) is functorial in the following sense. Suppose that we are given a homomorphism $W \to V$ of vertex algebras (so that each M_i becomes a V-module), a collection R_1, \ldots, R_N of W-modules and a collection of homomorphisms of W-modules $M_i \to R_i$ for all $i = 1, \ldots, N$. Then the corresponding map $\bigotimes_{i=1}^N R_i \to \bigotimes_{i=1}^N M_i$ gives rise to a map of the corresponding spaces of coinvariants

$$H_W(X, (x_i), (R_i)) \to H_V(X, (x_i), (M_i)).$$

Suppose now that W is the center of V (see [FB]). Then the action of Won any V-module M factors through a homomorphism $\widetilde{U}(W) \to \operatorname{End}_{\mathbb{C}} M$, where $\widetilde{U}(W)$ is the enveloping algebra of W (see [FB], Sect. 4.3). Let W(M) be the image of this homomorphism. Then $H_W(X, (x_i), (W(M_i)))$ is an algebra, and we obtain a natural homomorphism of algebras

$$(2.7) H_W(X,(x_i),(W(M_i))) \to \operatorname{End}_{\mathbb{C}} H_V(X,(x_i),(M_i))$$

If $V = \mathbb{V}_0$, then the center of V is precisely the subspace $\mathfrak{z}(\widehat{\mathfrak{g}})$ of $\mathfrak{g}[[t]]$ -invariant vectors in \mathbb{V}_0 (see [FB]). In particular, $\mathfrak{z}(\widehat{\mathfrak{g}})$ is a commutative vertex subalgebra of \mathbb{V}_0 . A module over the vertex algebra $\mathfrak{z}(\widehat{\mathfrak{g}})$ is the same as a module over the topological algebra $\widetilde{U}(\mathfrak{z}(\widehat{\mathfrak{g}}))$ which is nothing but the center $Z(\widehat{\mathfrak{g}})$ of $\widetilde{U}_{\kappa_c}(\widehat{\mathfrak{g}})$ (see [F2], Sect. 11). The action of $Z(\widehat{\mathfrak{g}})$ on any $\widehat{\mathfrak{g}}_{\kappa_c}$ -module M factors through the homomorphism $Z(\widehat{\mathfrak{g}}) \to \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} M$. Let Z(M) denote the image of this homomorphism. Recall that we have identified $Z(\widehat{\mathfrak{g}})$ with $\operatorname{Fun} \operatorname{Op}_{L_G}(D^{\times})$ in Theorem 2.5. For each $\widehat{\mathfrak{g}}_{\kappa_c}$ -module M, the algebra Z(M) is a quotient of $\operatorname{Fun} \operatorname{Op}_{L_G}(D^{\times})$, and hence $\operatorname{Spec} Z(M)$ is a subscheme in $\operatorname{Op}_{L_G}(D^{\times})$ which we denote by $\operatorname{Op}_{L_G}^M(D^{\times})$. The space of coinvariants $H_{\mathfrak{z}(\hat{\mathfrak{g}})}(X, (x_i), Z(M_i))$ is computed in the same way as in Theorem 4.7 of [F3]:

(2.8)
$$H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(X,(x_i),Z(M_i)) \simeq \operatorname{Fun}\operatorname{Op}_{L_G}(X,(x_i),(M_i))$$

where $\operatorname{Op}_{L_G}(X, (x_i), (M_i))$ is the space of ^LG-opers on X which are regular on $X \setminus \{x_1, \ldots, x_N\}$ and such that their restriction to D_x^{\times} belongs to $\operatorname{Op}_{L_G}^{M_i}(D_{x_i}^{\times})$ for all $i = 1, \ldots, N$.

Let u be an additional point of X, different from x_1, \ldots, x_N , and let us insert $\mathfrak{g}(\hat{\mathfrak{g}}) \simeq Z(\mathbb{V}_0)$ at this point. Then by Theorem 10.3.1 of [FB] we have an isomorphism

$$H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(X,(x_i),(Z(M_i))) \simeq H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(X;(z_i),u;(Z(M_i)),Z(\mathbb{V}_0)).$$

Hence we obtain a homomorphism

(2.9)
$$\mathfrak{z}(\widehat{\mathfrak{g}})_u \simeq Z(\mathbb{V}_0)_u \to H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(X, (x_i), (Z(M_i))).$$

The corresponding homomorphism

Fun
$$\operatorname{Op}_{L_G}(D_u) \to \operatorname{Fun} \operatorname{Op}_{L_G}(X, (x_i), (M_i))$$

(see formula (2.6)) is induced by the embedding

$$\operatorname{Op}_{L_G}(X, (x_i), (M_i)) \hookrightarrow \operatorname{Op}_{L_G}(D_u)$$

obtained by restricting an oper to D_u .

We apply this construction in the case when the curve X is \mathbb{P}^1 , the points are z_1, \ldots, z_N and ∞ , and the modules are $\widehat{\mathfrak{g}}_{\kappa_c}$ -modules M_1, \ldots, M_N and M_{∞} . It is proved in [FB] (see Theorem 9.3.3 and Remark 9.3.10) that the corresponding space of coinvariants is the space of $\mathfrak{g}_{(z_i)}$ -coinvariants of $\bigotimes_{i=1}^N \mathbb{M}_i \otimes \mathbb{M}_{\infty}$, which is the space $H(M_1, \ldots, M_N, M_{\infty})$ that we have computed in Lemma 2.1.

The homomorphism (2.7) specializes to a homomorphism

$$(2.10) \qquad H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(\mathbb{P}^1; (z_i), \infty; (Z(M_i)), Z(M_\infty)) \to \operatorname{End}_{\mathbb{C}} H(M_1, \dots, M_N, M_\infty).$$

Observe that by its very definition the homomorphism

$$\Phi_u : \mathfrak{g}(\widehat{\mathfrak{g}})_u \to \operatorname{End}_{\mathbb{C}} H(M_1, \ldots, M_N, M_\infty)$$

constructed in Section 2.2 factors through the homomorphisms (2.10) and (2.9). Hence the image of Φ_u is a quotient of the algebra

$$H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(\mathbb{P}^{1};(z_{i}),\infty;(Z(M_{i})),Z(M_{\infty}))\simeq \operatorname{Fun}\operatorname{Op}_{L_{G}}(\mathbb{P}^{1};(z_{i}),\infty;(Z(M_{i})),Z(M_{\infty})),$$

according to the isomorphism (2.8).

Let us specialize this result to our setting. First we suppose that all M_i 's and M_{∞} are equal to $U(\mathfrak{g})$. By Theorem 2.6,(1), we have

$$\operatorname{Op}_{L_G}^{U(\mathfrak{g})}(D^{\times}) = \operatorname{Op}_{L_G}^{\operatorname{RS}}(D).$$

Therefore we find that

(2.11)
$$H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(\mathbb{P}^{1};(z_{i}),\infty;(U(\mathfrak{g})),U(\mathfrak{g}))\simeq \operatorname{Fun}\operatorname{Op}_{L_{G}}^{\operatorname{RS}}(\mathbb{P}^{1})_{(z_{i}),\infty}.$$

Next, by Theorem 2.6,(2), we have

$$\operatorname{Op}_{{}^{L}G}^{U(\mathfrak{g})/I_{\lambda}}(D^{\times}) = \operatorname{Op}_{{}^{L}G}^{\operatorname{RS}}(D)_{\lambda}.$$

Therefore

$$(2.12) \quad H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(\mathbb{P}^{1};(z_{i}),\infty;(U(\mathfrak{g})/I_{\lambda_{i}}),U(\mathfrak{g})/I_{\lambda_{\infty}})\simeq \operatorname{Fun}\operatorname{Op}_{L_{G}}^{\operatorname{RS}}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}}.$$

Finally, by Theorem 2.6,(3), we have

$$\operatorname{Op}_{L_G}^{\mathbb{V}_{\lambda}}(D^{\times}) = \operatorname{Op}_{L_G}(D)_{\lambda}.$$

Therefore

(2.13)
$$H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(\mathbb{P}^{1};(z_{i}),\infty;(\mathbb{V}_{\lambda_{i}}),\mathbb{V}_{\lambda_{\infty}})\simeq \operatorname{Fun}\operatorname{Op}_{L_{G}}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}}.$$

Moreover, in all three cases the homomorphism Φ_u is just the natural homomorphism from Fun $\operatorname{Op}_{L_G}(D_u)$ to the above algebras of functions that is induced by the restrictions of the corresponding opers to the disc D_u .

Now consider the homomorphism (2.10) in the case of the space of coinvariants given by the left-hand side of formula (2.11),

Fun
$$\operatorname{Op}_{L_G}^{\operatorname{RS}}(\mathbb{P}^1)_{(z_i),\infty} \to \operatorname{End}_{\mathbb{C}} U(\mathfrak{g})^{\otimes (N+1)}/\mathfrak{g}_{\operatorname{diag}} \simeq \operatorname{End}_{\mathbb{C}} U(\mathfrak{g})^{\otimes N}$$

where the second isomorphism we use the identification of $U(\mathfrak{g})^{\otimes (N+1)}/\mathfrak{g}_{\text{diag}}$ with $U(\mathfrak{g})^{\otimes N}$ corresponding to the first N factors. It follows from the explicit computation of this map given above that its image belongs to

$$(U(\mathfrak{g})^{\otimes N})^G \subset \operatorname{End}_{\mathbb{C}} U(\mathfrak{g})^{\otimes N},$$

and so we have a homomorphism

By definition, the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ is the image of this homomorphism.

Likewise, we obtain from formula (2.12) that the homomorphism (2.10) gives rise to a homomorphism

(2.15)
$$\operatorname{Fun}\operatorname{Op}_{L_{G}}^{\operatorname{RS}}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}} \to \left(U(\mathfrak{g})^{\otimes(N+1)}/\mathfrak{g}_{\operatorname{diag}}\right)^{G}/I_{(\lambda_{i}),\lambda_{\infty}},$$

whose image is the algebra $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$.

Finally, formula (2.13) gives us a homomorphism

Fun
$$\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} \to \operatorname{End}_{\mathbb{C}}\left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right),$$

whose image is the algebra $\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$. This proves part (3) of the theorem.

To prove parts (1) and (2), it remains to show that the homomorphisms (2.14) and (2.15) are injective. It is sufficient to prove that the latter is injective. To see that, we pass to the associate graded spaces on both sides with respect to natural filtrations which we now describe.

22

According to the identification given in formula (1.4), the algebra of functions on $\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is filtered, and the corresponding associated graded algebra is the algebra of functions on the vector space

$$C_{(z_i),\infty}^{\mathrm{RS}} = \bigoplus_{i=1}^{\ell} \Gamma(\mathbb{P}^1, \Omega^{\otimes (d_i+1)}(-d_i z_1 - \ldots - d_i z_N - d_i \infty)),$$

where Ω is the canonical line bundle on \mathbb{P}^1 . The algebra

$$\left(U(\mathfrak{g})^{\otimes (N+1)}/\mathfrak{g}_{\operatorname{diag}}\right)^G/I_{(\lambda_i),\lambda_{\infty}}$$

carries a PBW filtration, and the associated graded is the algebra of functions on the space $\mu^{-1}((T^*G/B)^{N+1})/G$, where $\mu: (T^*G/B)^{N+1} \to \mathfrak{g}^*$ is the moment map corresponding to the diagonal action of G on $(T^*G/B)^{N+1}$. The two filtrations are compatible according to [F2]. The corresponding homomorphism of the associate graded algebras

(2.16)
$$\operatorname{Fun} C^{\mathrm{RS}}_{(z_i),\infty} \to \operatorname{Fun} \mu^{-1}((T^*G/B)^{N+1})/G$$

is induced by a map

$$h_{(z_i),\infty}: \mu^{-1}((T^*G/B)^{N+1})/G \to C^{\mathrm{RS}}_{(z_i),\infty},$$

that we now describe.

Let us identify the tangent space to a point $gB \subset G/B$ with $(\mathfrak{g}/g\mathfrak{b}g^{-1})^* \simeq g\mathfrak{n}g^{-1}$. Then a point in $\mu^{-1}((T^*G/B)^{N+1})/G$ consists of an (N+1)-tuple of points g_iB of G/B and an (N+1)-tuple of vectors (η_i) , where $\eta_i \in g_i\mathfrak{n}g_i^{-1} \subset \mathfrak{g}$ such that $\sum_{i=1}^{N+1} \eta_i = 0$, considered up to simultaneous conjugation by G.

We attach to it the $\mathfrak{g}\text{-valued}$ one-form

$$\eta = \sum_{i=1}^{N} \frac{\eta_i}{t - z_i} dt$$

on \mathbb{P}^1 with poles at z_1, \ldots, z_N, ∞ . Let P_1, \ldots, P_ℓ be generators of the algebra of *G*-invariant polynomials on \mathfrak{g} of degrees $d_i + 1$. Then

$$h_{(z_i),\infty}((g_i),(\eta_i)) = (P_i(\eta))_{i=1}^{\ell} \in C_{(z_i),\infty}^{\mathrm{RS}}$$

The space $\mu^{-1}((T^*G/B)^{N+1})/G$ is identified with the moduli space of Higgs fields on the trivial *G*-bundle with parabolic structures at z_1, \ldots, z_N, ∞ , and the map $h_{(z_i),\infty}$ is nothing but the Hitchin map (see [ER]). The Hitchin map is known to be proper, so in particular it is surjective (see, e.g., [M]). Therefore the corresponding homomorphism (2.16) of algebras of functions is injective. This implies that the homomorphism (2.15) is also injective and completes the proof of the theorem.

This theorem has an important application to the question of simultaneous diagonalization of generalized Gaudin Hamiltonians, or equivalently, of the commutative algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, on the tensor product $\bigotimes_{i=1}^N M_i$ of \mathfrak{g} -modules. Indeed, the joint eigenvalues of the generalized Gaudin Hamiltonians on any eigenvector

in $\bigotimes_{i=1}^{N} M_i$ correspond to a point in the spectrum of the algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, which, according to Theorem 2.7,(1), is a point of the space $\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty}$.

If we assume in addition that each of the modules M_i admits a central character $\varpi(\lambda_i + \rho)$ (for instance, if M_i is the Verma module M_{λ_i}) and we are looking for eigenvectors in the component of $\bigotimes_{i=1}^N M_i$ corresponding to the central character $\varpi(-\lambda_{\infty} - \rho)$ with respect to the diagonal action of \mathfrak{g} , then the joint eigenvalues define a point in the spectrum of the algebra $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$, i.e., a point of $\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$.

Finally, for a collection of dominant integral weights $\lambda_1, \ldots, \lambda_N, \lambda_\infty$, the joint eigenvalues of the generalized Gaudin Hamiltonians on $(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_\infty})^G$ is a point in the spectrum of the algebra $\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_\infty}(\mathfrak{g})$, which is a point of $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_\infty}$.

A natural question is whether, conversely, one can attach to a ${}^{L}G$ -oper on \mathbb{P}^{1} with regular singularities at $z_{1}, \ldots, z_{N}, \infty$ (and satisfying additional conditions as above) an eigenvector in $\bigotimes_{i=1}^{N} M_{i}$ with such eigenvalues. It turns out that for general modules this is not true, but if these modules are finite-dimensional, then we conjecture that it is true. In order to construct the eigenvectors we use the procedure called Bethe Ansatz. As shown in [FFR], this procedure may be cast in the framework of coinvariants that we have discussed in this section, using the Wakimoto modules over $\hat{\mathfrak{g}}_{\kappa_{c}}$. We will explain that in Section 4 and Section 5.5. But first we need to introduce Miura opers and Cartan connections.

3. Miura opers and Cartan connections

By definition (see [F2], Sect. 10.3), a *Miura G-oper* on X (which is a smooth curve or a disc) is a quadruple $(\mathcal{F}, \nabla, \mathcal{F}_B, \mathcal{F}'_B)$, where $(\mathcal{F}, \nabla, \mathcal{F}_B)$ is a *G*-oper on X and \mathcal{F}'_B is another *B*-reduction of \mathcal{F} which is preserved by ∇ .

We denote the space of Miura G-opers on X by $MOp_G(X)$.

3.1. Miura opers and flag manifolds

A *B*-reduction of \mathcal{F} which is preserved by the connection ∇ is uniquely determined by a *B*-reduction of the fiber \mathcal{F}_x of \mathcal{F} at any point $x \in X$ (in the case when U = D, x has to be the origin $0 \in D$). The set of such reductions is the twist

(3.1)
$$(G/B)_{\mathcal{F}_x} = \mathcal{F}_x \underset{G}{\times} G/B = \mathcal{F}'_{B,x} \underset{B}{\times} G/B = (G/B)_{\mathcal{F}'_{B,x}}$$

of the flag manifold G/B. If X is a curve or a disc and the oper connection has a regular singularity and trivial monodromy representation, then this connection gives us a global (algebraic) trivialization of the bundle \mathcal{F} . Then any *B*-reduction of the fiber \mathcal{F}_x gives rise to a global (algebraic) *B*-reduction of \mathcal{F} . Thus, we obtain:

Lemma 3.1. Suppose that we are given an oper τ on a curve X (or on the disc) such that the oper connection has a regular singularity and trivial monodromy.

Then for each $x \in X$ there is a canonical isomorphism between the space of Miura opers with the underlying oper τ and the twist $(G/B)_{\mathcal{F}_{B,x}}$.

Recall that the *B*-orbits in G/B, known as the *Schubert cells*, are parameterized by the Weyl group W of G. Let w_0 be the longest element of the Weyl group of G. Denote the orbit $Bw_0wB \subset G/B$ by S_w (so that S_1 is the open orbit). We obtain from the second description of $(G/B)_{\mathcal{F}_x}$ given in formula (3.1) that $(G/B)_{\mathcal{F}_x}$ decomposes into a union of locally closed subvarieties $S_{w,\mathcal{F}'_{B,x}}$, which are the $\mathcal{F}'_{B,x}$ -twists of the Schubert cells S_w . The *B*-reduction $\mathcal{F}'_{B,x}$ defines a point in $(G/B)_{\mathcal{F}'_{B,x}}$ which is the one point orbit $S_{w_0,\mathcal{F}'_{B,x}}$. We will say that the *B*-reductions $\mathcal{F}_{B,x}$ and $\mathcal{F}'_{B,x}$ are in *relative position* w with if $\mathcal{F}_{B,x}$ belongs to $S_{w,\mathcal{F}'_{B,x}}$. In particular, if it belongs to the open orbit $S_{1,\mathcal{F}'_{B,x}}$, we will say that $\mathcal{F}_{B,x}$ and $\mathcal{F}'_{B,x}$ are in generic position.

A Miura *G*-oper is called *generic* at the point $x \in X$ if the *B*-reductions $\mathcal{F}_{B,x}$ and $\mathcal{F}'_{B,x}$ of \mathcal{F}_x are in generic position. In other words, $\mathcal{F}_{B,x}$ belongs to the stratum $\operatorname{Op}_G(X) \times S_{1,\mathcal{F}'_{B,x}} \subset \operatorname{MOp}_G(X)$. Being generic is an open condition. Therefore if a Miura oper is generic at $x \in X$, then there exists an open neighborhood U of xsuch that it is also generic at all other points of U. We denote the space of generic Miura opers on U by $\operatorname{MOp}_G(U)_{\text{gen}}$.

Lemma 3.2. Suppose we are given a Miura oper on the disc D_x around a point $x \in X$. Then its restriction to the punctured disc D_x^{\times} is generic.

Proof. Since being generic is an open condition, we obtain that if a Miura oper is generic at x, it is also generic on the entire D_x . Hence we only need to consider the situation where the Miura oper is not generic at x, i.e., the two reductions $\mathcal{F}_{B,x}$ and $\mathcal{F}'_{B,x}$ are in relative position $w \neq 1$. Let us trivialize the *B*-bundle \mathcal{F}_B , and hence the *G*-bundle \mathcal{F}_G over D_x . Then ∇ gives us a connection on the trivial *G*-bundle which we can bring to the canonical form

$$\nabla = \partial_t + p_{-1} + \sum_{j=1}^{\ell} v_j(t) \cdot p_j$$

(see Lemma 1.1). It induces a connection on the trivial G/B-bundle. We are given a point gB in the fiber of the latter bundle which lies in the orbit $S_w = Bw_0wB$, where $w \neq 1$. Consider the horizontal section whose value at x is gB, viewed as a map $D_x \to G/B$. We need to show that the image of this map lies in the open B-orbit $S_1 = Bw_0B$ over D_x^{\times} , i.e., it does not lie in the orbit S_y for any $y \neq 1$.

Suppose that this is not so, and the image of the horizontal section actually lies in the orbit S_y for some $y \neq 1$. Since all *B*-orbits are *H*-invariant, we obtain that the same would be true for the horizontal section with respect to the connection $\nabla' = h \nabla h^{-1}$ for any constant element of *H*. Choosing $h = \check{\rho}(a)$ for $a \in \mathbb{C}^{\times}$, we can bring the connection to the form

$$\partial_t + a^{-1}p_{-1} + \sum_{j=1}^{\ell} a^{d_j} v_j(t) \cdot p_j.$$

Changing the variable t to $s = a^{-1}t$, we obtain the connection

$$\partial_s + p_{-1} + \sum_{j=1}^{\ell} a^{d_j + 1} v_j(t),$$

so choosing small a we can make the functions $v_j(t)$ arbitrarily small. Therefore without loss of generality we can consider the case when our connection operator is $\nabla = \partial_t + p_{-1}$.

In this case our assumption that the horizontal section lies in $S_y, y \neq 1$, means that the vector field ξ_{p-1} corresponding to the infinitesimal action of p_{-1} on G/B is tangent to an orbit $S_y, y \neq 1$, in the neighborhood of some point gB of $S_w \subset G/B, w \neq 1$. But then, again because of the *H*-invariance of the *B*-orbits, the vector field $\xi_{hp-1h^{-1}}$ is also tangent to this orbit for any $h \in H$. For any $i = 1 \dots, \ell$, there exists a one-parameter subgroup $h_{\epsilon}^{(i)}, \epsilon \in \mathbb{C}^{\times}$ in *H*, such that $\lim_{\epsilon \to 0} \epsilon p_{-1} \epsilon^{-1} = f_i$. Hence we obtain that each of the vector fields $\xi_{f_i}, i = 1 \dots, \ell$, is tangent to the orbit $S_y, y \neq 1$, in the neighborhood of $gB \in S_w, w \neq 1$. But then all commutators of these vectors fields are also tangent to this orbit. Hence we obtain that all vector fields of the form $\xi_p, p \in \mathfrak{n}_-$, are tangent to S_y in the neighborhood of $gB \in S_w$.

Consider any point of G/B that does not belong to the open dense orbit S_1 . Then the quotient of the tangent space to this point by the tangent space to the *B*-orbit passing through this point is non-zero and the vector fields from the Lie algebra \mathfrak{n}_- map surjectively onto this quotient. Therefore they cannot be tangent to the orbit $S_y, y \neq 1$, in a neighborhood of gB. Therefore our Miura oper is generic on D_x^{\times} .

This lemma shows that any Miura oper on any smooth curve X is generic over an open dense subset.

3.2. Cartan connections

Introduce the *H*-bundle $\Omega^{\check{\rho}}$ on *X* which is uniquely determined by the following property: for any character $\lambda : H \to \mathbb{C}^{\times}$, the line bundle $\Omega^{\check{\rho}} \times \lambda$ associated to the corresponding one-dimensional representation of *H* is $\Omega^{\langle \lambda, \check{\rho} \rangle}$.

Explicitly, connections on $\Omega^{\tilde{\rho}}$ may be described as follows. If we choose a local coordinate t on X, then we trivialize $\Omega^{\tilde{\rho}}$ and represent the connection as an operator $\partial_t + \mathbf{u}(t)$, where $\mathbf{u}(t)$ is an \mathfrak{h} -valued function on X. If s is another coordinate such that $t = \varphi(s)$, then this connection will be represented by the operator

(3.2)
$$\partial_s + \varphi'(s)\mathbf{u}(\varphi(s)) - \check{\rho} \cdot \frac{\varphi''(s)}{\varphi'(s)}.$$

Let $\operatorname{Conn}(\Omega^{\check{\rho}})_X$ be the space of connections on the *H*-bundle $\Omega^{\check{\rho}}$ on *X*. When no confusion can arise, we will simply write Conn_X . We define a map

$$\mathbf{b}_X : \operatorname{Conn}_X \to \operatorname{MOp}_G(X)_{\operatorname{gen}}.$$

Suppose we are given a connection $\overline{\nabla}$ on the *H*-bundle $\Omega^{\check{\rho}}$ on *D*. We associate to it a generic Miura oper as follows. Let us choose a splitting $H \to B$ of the homomorphism $B \to H$ and set $\mathfrak{F} = \Omega^{\check{\rho}} \times G, \mathfrak{F}_B = \Omega^{\check{\rho}} \times B$, where we consider the adjoint action of *H* on *G* and on *B* obtained through the above splitting. The choice of the splitting also gives us the opposite Borel subgroup B_- , which is the unique Borel subgroup in generic position with *B* containing *H*. Let again w_0 be the longest element of the Weyl group of \mathfrak{g} . Then $w_0 B$ is a *B*-torsor equipped with a left action of *H*, so we define the *B*-subbundle \mathfrak{F}'_B of \mathfrak{F} as $\Omega^{\check{\rho}} \times w_0 B$.

Observe that the space of connections on F is isomorphic to the direct product

$$\operatorname{Conn}_X \times \bigoplus_{\alpha \in \Delta} \Gamma(X, \Omega^{\alpha(\check{\rho})+1})$$

Its subspace corresponding to negative simple roots is isomorphic to the tensor product of $\left(\bigoplus_{i=1}^{\ell} \mathfrak{g}_{-\alpha_i}\right)$ and Fun X. Having chosen a basis element f_i of $\mathfrak{g}_{-\alpha_i}$ for each $i = 1, \ldots, \ell$, we now construct an element $p_{-1} = \sum_{i=1}^{\ell} f_i \otimes 1$ of this space. Now we set $\nabla = \overline{\nabla} + p_{-1}$. By construction, ∇ has the correct relative position with the *B*-reduction \mathcal{F}_B and preserves the *B*-reduction \mathcal{F}'_B . Therefore the quadruple $(\mathcal{F}, \nabla, \mathcal{F}_B, \mathcal{F}'_B)$ is a generic Miura oper on X. We define the morphism \mathbf{b}_X by setting $\mathbf{b}_X(\overline{\nabla}) = (\mathcal{F}, \nabla, \mathcal{F}_B, \mathcal{F}'_B)$.

This map is independent of the choice of a splitting $H \to B$ and of the generators $f_i, i = 1, \ldots, \ell$.

Proposition 3.3 ([F2],Prop. 10.4). The map \mathbf{b}_X is an isomorphism of algebraic varieties

$$\operatorname{Conn}_X \to \operatorname{MOp}_G(X)_{\operatorname{gen}}.$$

Thus, generic Miura opers are the same as Cartan connections, which are much simpler objects than opers.

The composition $\overline{\mathbf{b}}_X$ of \mathbf{b}_X and the forgetful map $\mathrm{MOp}_G(X)_{\mathrm{gen}} \to \mathrm{Op}_G(X)$ is called the *Miura transformation*.

For example, in the case of $\mathfrak{g} = \mathfrak{sl}_2$, we have a connection $\overline{\nabla} = \partial_t - u(t)$ on the line bundle $\Omega^{1/2}$ (equivalently, a connection $\overline{\nabla}^t = \partial_t + u(t)$ on $\Omega^{-1/2}$), and the Miura transformation assigns to this connection the PGL_2 -oper

$$\partial_t + \left(\begin{array}{cc} -u(t) & 0 \\ 1 & u(t) \end{array} \right).$$

The oper B reduction \mathcal{F}_B corresponds to the upper triangular matrices, and the Miura B-reduction \mathcal{F}'_B corresponds to the lower triangular matrices. The corresponding projective connection is

$$\partial_t^2 - v(t) = (\partial_t - u(t))(\partial_t + u(t)),$$

i.e.,

$$u(t) \mapsto v(t) = u(t)^2 - u'(t).$$

3.3. Singularities of Cartan connections

Consider the Miura transformation $\overline{\mathbf{b}}_{D_x^{\times}}$ in the case of the punctured disc D_x^{\times} ,

$$\overline{\mathbf{b}}_{D_x^{\times}} : \operatorname{Conn}(\Omega^{\check{\rho}})_{D_x^{\times}} \to \operatorname{Op}_G(D_x^{\times}).$$

Let $\operatorname{Conn}(\Omega^{\check{\rho}})_{D_x}^{\operatorname{RS}} \subset \operatorname{Conn}(\Omega^{\check{\rho}})_{D_x^{\times}}$ be the space of all connections on the *H*-bundle $\Omega^{\check{\rho}}$ on D_x with regular singularity, i.e., those for which the connection operator has the form

$$\overline{\nabla} = \partial_t + \frac{\lambda}{t} + \sum_{n \ge 0} u_n t^n.$$

We define a map

$$\operatorname{res}_{\mathfrak{h}}:\operatorname{Conn}(\Omega^{\check{\rho}})_D^{\operatorname{RS}}\to\mathfrak{h}$$

assigning to such a connection its residue λ .

It follows from the definition of the Miura transformation $\overline{\mathbf{b}}_{D_x^{\times}}$ that its restriction to $\operatorname{Conn}(\Omega^{\check{\rho}})_{D_x}^{\mathrm{RS}} \subset \operatorname{Conn}(\Omega^{\check{\rho}})_{D_x^{\times}}$ takes values in $\operatorname{Op}_G^{\mathrm{RS}}(D_x)$. Hence we obtain a morphism

$$\overline{\mathbf{b}}_x^{\mathrm{RS}} : \mathrm{Conn}(\Omega^{\check{
ho}})_{D_x}^{\mathrm{RS}} \to \mathrm{Op}_G^{\mathrm{RS}}(D_x)$$

Explicitly, after choosing a coordinate t on D_x , we can write $\overline{\nabla}$ as $\partial_t + t^{-1}\mathbf{u}(t)$, where $\mathbf{u}(t) \in \mathfrak{h}[[t]]$. Its residue is $\mathbf{u}(0)$. Then the corresponding oper with regular singularity is by definition the N((t))-equivalence class of the operator

$$\nabla = \partial_t + p_{-1} + t^{-1} \mathbf{u}(t),$$

which is the same as the N[[t]]-equivalence class of the operator

$$\check{p}(t)\nabla\check{\rho}(t)^{-1} = \partial_t + t^{-1}(p_{-1} - \check{\rho} + \mathbf{u}(t)),$$

so it is indeed an oper with regular singularity.

Therefore it follows from the definition that we have a commutative diagram

$$(3.3) \qquad \begin{array}{ccc} \operatorname{Conn}(\Omega^{\check{\rho}})_{D_{x}}^{\operatorname{RS}} & \xrightarrow{\overline{\mathbf{b}}_{x}^{\operatorname{RS}}} & \operatorname{Op}_{G}^{\operatorname{RS}}(D_{x}) \\ & & & & \downarrow^{\operatorname{res}} \\ & & & & \downarrow^{\operatorname{res}} \\ & & & & \mathfrak{h}/W \end{array}$$

where the lower horizontal map is the composition of the map $\check{\lambda} \mapsto \check{\lambda} - \check{\rho}$ and the projection $\varpi : \mathfrak{h} \to \mathfrak{h}/W$.

Now let $\operatorname{Conn}_{D_x,\check{\lambda}}^{\operatorname{reg}}$ be the preimage under $\overline{\mathbf{b}}_x^{\operatorname{RS}}$ of the subspace $\operatorname{Op}_G(D_x)_{\check{\lambda}} \subset \operatorname{Op}_G^{\operatorname{RS}}(D_x)$ of *G*-opers on D_x^{\times} with regular singularity, residue $\varpi(-\check{\lambda}-\check{\rho})$ and trivial monodromy. By the commutativity of the above diagram, a connection in $\operatorname{Conn}_{D_x,\check{\lambda}}^{\operatorname{reg}}$ necessarily has residue of the form $-w(\check{\lambda}+\check{\rho})+\check{\rho}$ for some element w of the Weyl group of *G*, so that $\operatorname{Conn}_{D_x,\check{\lambda}}^{\operatorname{reg}}$ is the disjoint union of its subsets $\operatorname{Conn}_{D_x,\check{\lambda},w}^{\operatorname{reg}}$ consisting of connections with residue $-w(\check{\lambda}+\check{\rho})+\check{\rho}$.

The restriction of $\overline{\mathbf{b}}^{\mathrm{RS}}_x$ to $\mathrm{Conn}^{\mathrm{reg}}_{D_x,\check{\lambda},w}$ is a map

$$\overline{\mathbf{b}}_{\check{\lambda},w}: \operatorname{Conn}_{D_x,\check{\lambda},w}^{\operatorname{reg}} \to \operatorname{Op}_G(D_x)_{\check{\lambda}}.$$

Let us recall that by construction of the Miura transformation $\overline{\mathbf{b}}$, each oper on D_x^{\times} which lies in the image of the map $\overline{\mathbf{b}}$ (hence in particular, in the image of $\overline{\mathbf{b}}_{\tilde{\lambda},w}$) carries a canonical horizontal *B*-reduction

$$\mathfrak{F}'_B = \Omega^{\check{\rho}} \underset{H}{\times} w_0 B$$

(i.e., it carries a canonical structure of Miura oper on D_x^{\times}). But if this oper is in the image of $\overline{\mathbf{b}}_{\check{\lambda},w}$, i.e., belongs to $\operatorname{Op}_G(D_x)_{\check{\lambda}}$, then the oper *B*-reduction \mathcal{F}_B (and hence the oper bundle \mathcal{F}) has a canonical extension to a *B*-bundle on the entire disc D_x , namely, one for which the oper connection has the form (1.8). Therefore the *B*-reduction $\mathcal{F}'_B = \Omega^{\check{\rho}} \times w_0 B$ may also be extended to D_x .

Therefore we can lift $\overline{\mathbf{b}}_{\tilde{\lambda},w}$ to a map

$$\mathbf{b}_{\check{\lambda},w}: \operatorname{Conn}_{D_x,\check{\lambda},w}^{\operatorname{reg}} \to \operatorname{MOp}_G(D_x)_{\check{\lambda}}.$$

Let $\operatorname{MOp}_G(D_x)_{\check{\lambda},w} \subset \operatorname{MOp}_G(D_x)_{\check{\lambda}}$ be the subvariety of those Miura opers of coweight $\check{\lambda}$ which have relative position w at x. Then

$$\operatorname{MOp}_G(D_x)_{\check{\lambda},w} \simeq \operatorname{Op}_G(D_x)_{\check{\lambda}} \times S_{w,\mathcal{F}'_{B,x}}$$

The following result is due to D. Gaitsgory and myself [FG] (see [F3], Proposition 2.9).

Proposition 3.4. For each $w \in W$ the morphism $\mathbf{b}_{\lambda,w}$ is an isomorphism between the varieties $\operatorname{Conn}_{D_x \lambda,w}^{\operatorname{reg}}$ and $\operatorname{MOp}_G(D_x)_{\lambda,w}$.

Proof. First we observe that at the level of points the map defined by $\mathbf{b}_{\bar{\lambda},w}, w \in W$, from the union of $\operatorname{Conn}_{D_x,\bar{\lambda},w}^{\operatorname{reg}}, w \in W$, to $\operatorname{MOp}_G(D_x)_{\bar{\lambda}}$, is a bijection. Indeed, by Proposition 3.3 we have a map taking a Miura oper from $\operatorname{MOp}_G(D_x)_{\bar{\lambda}}$, considered as a Miura oper on the punctured disc D_x^{\times} , to a connection $\overline{\nabla}$ on the *H*-bundle $\Omega^{\check{\rho}}$ over D_x^{\times} . We have shown above that $\overline{\nabla}$ has regular singularity at x and that its residue is of the form $-w(\check{\lambda}+\check{\rho})+\check{\rho}, w\in W$. Thus, we obtain a map from the set of points of $\operatorname{MOp}_G(D_x)_{\bar{\lambda}}$ to the union of $\operatorname{Conn}_{D_x,\bar{\lambda},w}^{\operatorname{reg}}, w \in W$, and by Proposition 3.3 it is a bijection.

It remains to show that if the Miura oper belongs to $\mathrm{MOp}_G(D_x)_{\check{\lambda},w}$, then the corresponding connection has residue precisely $-w(\check{\lambda}+\check{\rho})+\check{\rho}$.

Thus, we are given a *G*-oper $(\mathcal{F}, \nabla, \mathcal{F}_B, \mathcal{F}'_B)$ of coweight $\check{\lambda}$. Let us choose a trivialization of the *B*-bundle \mathcal{F}_B . Then the connection operator reads

(3.4)
$$\nabla = \partial_t + \sum_{i=1}^{\ell} t^{\langle \alpha_i, \check{\lambda} \rangle} f_i + \mathbf{v}(t), \qquad \mathbf{v}(t) \in \mathfrak{b}[[t]].$$

Suppose that the horizontal *B*-reduction \mathcal{F}'_B of our Miura oper has relative position w with \mathcal{F}_B at x (see Section 3 for the definition of relative position). We need to show that the corresponding connection on $\mathcal{F}'_H \simeq \Omega^{\check{\rho}}$ has residue $-w(\check{\lambda} + \check{\rho}) + \check{\rho}$.

This is equivalent to the following statement. Let $\Phi(t)$ be the *G*-valued solution of the equation

(3.5)
$$\left(\partial_t + \sum_{i=1}^{\ell} t^{\langle \alpha_i, \check{\lambda} \rangle} f_i + \mathbf{v}(t)\right) \Phi(t) = 0,$$

such that $\Phi(0) = 1$. Since the connection operator is regular at t = 0, this solution exists and is unique. Then $\Phi(t)w^{-1}w_0$ is the unique solution of the equation (3.5) whose value at t = 0 is equal to $w^{-1}w_0$.

By Lemma 3.2, we have

$$\Phi(t)w^{-1}w_0 = X_w(t)Y_w(t)Z_w(t)w_0,$$

where

$$X_w(t) \in N((t)), \qquad Y_w(t) \in H((t)), \qquad Z_w(t) \in N_-((t)).$$

We can write $Y_w(t) = \check{\mu}_w(t) Y_w(t)$, where $\check{\mu}_w$ is a coweight and $Y_w(t) \in H[[t]]$. Since the connection ∇ preserves

$$\Phi(t)w_0\mathfrak{b}_+w_0\Phi(t)^{-1} = \Phi(t)\mathfrak{b}_-\Phi(t)^{-1},$$

the connection $X(t)_w^{-1}\nabla X_w(t)$ preserves

$$Y_w(t)Z_w(t)\mathfrak{b}_-Z_w(t)^{-1}Y_w(t)^{-1} = \mathfrak{b}_-,$$

and therefore has the form

$$\partial_t + \sum_{i=1}^{\ell} t^{\langle \alpha_i, \check{\lambda} \rangle} f_i - \frac{\check{\mu}_w}{t} + \mathbf{u}(t), \qquad \mathbf{u}(t) \in \mathfrak{h}[[t]].$$

By conjugating it with $\lambda(t)$ we obtain a connection

$$\partial_t + p_{-1} - \frac{\check{\lambda} + \check{\mu}_w}{t} + \mathbf{u}(t), \qquad \mathbf{u}(t) \in \mathfrak{h}[[t]].$$

Therefore we need to show that

(3.6)
$$\check{\mu}_w = w(\lambda + \check{\rho}) - (\lambda + \check{\rho}).$$

To see that, let us apply the identity $\Phi(t)w^{-1} = X_w(t)Y_w(t)Z_w(t)$ to a nonzero vector $v_{w_0(\nu)}$ of weight $w_0(\nu)$ in a finite-dimensional irreducible \mathfrak{g} -module V_{ν} of highest weight ν (so that $v_{w_0(\nu)}$ is a lowest weight vector and hence is unique up to scalar). The right-hand side will then be equal to a $P(t)v_{w_0(\nu)}$ plus the sum of terms of weights greater than $w_0(\nu)$, where $P(t) = ct^{\langle w_0(\nu), \check{\mu}_w \rangle}, c \neq 0$, plus the sum of terms of higher degree in t. Applying the left-hand side to $v_{w_0(\nu)}$, we obtain $\Phi(t)v_{w^{-1}w_0(\nu)}$, where $v_{w^{-1}w_0(\nu)} \in V_{\nu}$ is a non-zero vector of weight $w^{-1}w_0(\nu)$ which is also unique up to a scalar.

Gaudin model and opers

Thus, we need to show that the coefficient with which $v_{w_0(\nu)}$ enters the expression $\Phi(t)v_{w^{-1}w_0(\nu)}$ is a polynomial in t whose lowest degree is equal to

$$\langle w_0(\nu), w(\lambda + \check{
ho}) - (\lambda + \check{
ho}) \rangle,$$

because if this is so for all dominant integral weights ν , then we obtain the desired equality (3.6). But this formula is easy to establish. Indeed, from the form (3.4) of the oper connection ∇ it follows that we can obtain a vector proportional to v_{w_0} by applying the operators $\frac{1}{\langle \alpha_i, \bar{\lambda} \rangle + 1} t^{\langle \alpha_i, \bar{\lambda} \rangle + 1} f_i, i = 1, \ldots, \ell$, to $v_{w^{-1}w_0(\nu)}$ in some order. The linear combination of these monomials appearing in the solution is the term of the lowest degree in t with which $v_{w_0(\nu)}$ enters $\Phi(t)v_{w^{-1}w_0(\nu)}$. It follows from Lemma 3.2 that it is non-zero. The corresponding power of t is nothing but the difference between the $(\check{\lambda} + \check{\rho})$ -degrees of the vectors $v_{w^{-1}w_0}$ and v_{w_0} , i.e.,

$$\langle w^{-1}w_0(\nu), \check{\lambda} + \check{\rho} \rangle - \langle w_0(\nu), \check{\lambda} + \check{\rho} \rangle = \langle w_0(\nu), w(\check{\lambda} + \check{\rho}) - (\check{\lambda} + \check{\rho}) \rangle,$$

as desired. This completes the proof.

Thus, we have identified the space $\operatorname{Conn}_{D_x,\check{\lambda},w}^{\operatorname{reg}}$ of connections on the *H*bundle $\Omega^{\check{\rho}}$ on D_x^{\times} of the form with the space of Miura opers on D_x^{\times} such that the underlying oper belongs to $\operatorname{Op}_G(D_x)_{\check{\lambda}}$ and the corresponding *B*-reductions \mathcal{F}_B and \mathcal{F}'_B have relative position w at x.

The condition that the image under $\mathbf{b}_{\check{\lambda},w}$ of a connection of the form

$$\partial_t - \frac{w(\dot{\lambda} + \check{\rho}) - \check{\rho}}{t} + \mathbf{u}(t), \qquad \mathbf{u}(t) \in \mathfrak{h}[[t]],$$

is an oper without monodromy imposes polynomial equations on the coefficients of the series $\mathbf{u}(t)$. Consider the simplest case when $\check{\lambda} = 0$ and $w = s_i$, the *i*th simple reflection. Then $-w(\check{\lambda} + \check{\rho}) + \check{\rho} = \check{\alpha}_i$, so we write this connection as

(3.7)
$$\overline{\nabla} = \partial_t + \frac{\check{\alpha}_i}{t} + \mathbf{u}(t), \qquad \mathbf{u}(t) \in \mathfrak{h}[[t]].$$

Lemma 3.5 ([F3],Lemma 2.10). A connection of the form (3.7) belongs to $\operatorname{Conn}_{D_x,s_i}^{\operatorname{reg}}$ (*i.e.*, the corresponding *G*-oper is regular at *x*) if and only if $\langle \alpha_i, \mathbf{u}(0) \rangle = 0$.

4. Wakimoto modules and Bethe Ansatz

In this section we explain how to construct Bethe eigenvectors of the generalized Gaudin Hamiltonians. For that we utilize the Wakimoto modules over $\hat{\mathfrak{g}}_{\kappa_c}$ which are parameterized by Cartan connections on the punctured disc. As the result, the eigenvectors will be parameterized by the Cartan connections on \mathbb{P}^1 with regular singularities at z_1, \ldots, z_N, ∞ and some additional points w_1, \ldots, w_m with residues $\lambda_1, \ldots, \lambda_N, \lambda_\infty$ and $-\alpha_{i_1}, \ldots, -\alpha_{i_m}$ and whose Miura transformation is an oper that has no singularities at w_1, \ldots, w_m .

4.1. Definition of Wakimoto modules

We recall some of the results of [FF1, F2] on the construction of the Wakimoto realization.

Let $\mathcal{A}^{\mathfrak{g}}$ be the Weyl algebra with generators $a_{\alpha,n}, a_{\alpha,n}^*, \alpha \in \Delta_+, n \in \mathbb{Z}$, and relations

(4.1)
$$[a_{\alpha,n}, a_{\beta,m}^*] = \delta_{\alpha,\beta} \delta_{n,-m}, \quad [a_{\alpha,n}, a_{\beta,m}] = [a_{\alpha,n}^*, a_{\beta,m}^*] = 0.$$

Introduce the generating functions

(4.2)
$$a_{\alpha}(z) = \sum_{n \in \mathbb{Z}} a_{\alpha,n} z^{-n-1},$$

(4.3)
$$a_{\alpha}^*(z) = \sum_{n \in \mathbb{Z}} a_{\alpha,n}^* z^{-n}.$$

Let $M_{\mathfrak{g}}$ be the Fock representation of $\mathcal{A}^{\mathfrak{g}}$ generated by a vector $|0\rangle$ such that

$$a_{\alpha,n}|0\rangle = 0, \quad n \ge 0; \qquad a_{\alpha,n}^*|0\rangle = 0, \quad n > 0.$$

It carries a vertex algebra structure (see [F2]).

Let π_0 be the commutative algebra $\mathbb{C}[b_{i,n}]_{i=1,...,\ell;n<0}$ with the derivation T given by the formula

$$T \cdot b_{i_1,n_1} \dots b_{i_m,n_m} = -\sum_{j=1}^m n_j b_{i_1,n_1} \dots b_{i_j,n_j-1} \dots b_{i_m,n_m}.$$

Then π_0 is naturally a commutative vertex algebra (see [FB], § 2.3.9). In particular, we have

$$Y(b_{i,-1}, z) = b_i(z) = \sum_{n < 0} b_{i,n} z^{-n-1}.$$

Recall that \mathbb{V}_0 carries the structure of a vertex algebra. Its vacuum vector will be denoted by v_{κ_c} .

In the next theorem describing the Wakimoto realization we will need the formulas defining the action of the Lie algebra \mathfrak{g} on the space of functions on the big cell U of the flag manifold G/B_+ . Here U is the open orbit of the unipotent subgroup $N_+ = [B_+, B_+]$ which is isomorphic to N_+ . Since the exponential map $\mathfrak{n}_+ \to N_+$ is an isomorphism, we obtain a system of coordinates $\{y_\alpha\}_{\alpha \in \Delta_+}$ on U corresponding to a fixed basis of root vectors $\{e_\alpha\}_{\alpha \in \Delta_+}$ in \mathfrak{n}_+ .

For each $\chi \in \mathfrak{h}^*$ we have a homomorphism $\rho_{\chi} : \mathfrak{g} \hookrightarrow \mathcal{D}_{\leq 1}(N_+)$ under which the $\mathcal{D}_{\leq 1}(N_+)$ -module Fun U becomes isomorphic to the \mathfrak{g} -module M_{χ}^* that is contragredient to the Verma module of highest weight χ . Explicitly, ρ_{χ} looks as follows. Let $e_i, h_i, f_i, i = 1 \dots, \ell$, be the generators of \mathfrak{g} . Then

(4.4)
$$\rho_{\chi}(e_i) = \frac{\partial}{\partial y_{\alpha_i}} + \sum_{\beta \in \Delta_+} P^i_{\beta}(y_{\alpha}) \frac{\partial}{\partial y_{\beta}},$$

(4.5)
$$\rho_{\chi}(h_i) = -\sum_{\beta \in \Delta_+} \beta(h_i) y_{\beta} \frac{\partial}{\partial y_{\beta}} + \chi(h_i),$$

(4.6)
$$\rho_{\chi}(f_i) = \sum_{\beta \in \Delta_+} Q^i_{\beta}(y_{\alpha}) \frac{\partial}{\partial y_{\beta}} + \chi(h_i) y_{\alpha_i},$$

for some polynomials P^i_{β}, Q^i_{β} in $y_{\alpha}, \alpha \in \Delta_+$.

In addition, we have a Lie algebra anti-homomorphism $\rho^R : \mathfrak{n}_+ \to \mathcal{D}_{\leq 1}(N_+)$ which corresponds to the *right* action of \mathfrak{n}_+ on N_+ . The differential operators $\rho^R(x), x \in \mathfrak{n}_+$, commute with the differential operators $\rho_{\chi}(x'), x' \in \mathfrak{n}_+$ (though their commutation relations with $\rho_{\chi}(x'), x' \notin \mathfrak{n}_+$, are complicated in general). We have

$$\rho^{R}(e_{i}) = \frac{\partial}{\partial y_{\alpha_{i}}} + \sum_{\beta \in \Delta_{+}} P^{R,i}_{\beta}(y_{\alpha}) \frac{\partial}{\partial y_{\beta}}$$

for some polynomials $P_{\beta}^{R,i}, Q_{\beta}^{i}$ in $y_{\alpha}, \alpha \in \Delta_{+}$. We let

(4.7)
$$e_i^R(z) = \sum_{n \in \mathbb{Z}} e_{i,n}^R z^{-n-1} = a_{\alpha_i}(z) + \sum_{\beta \in \Delta_+} P_{\beta}^{R,i}(a_{\alpha}^*(z)) a_{\beta}(z).$$

Now we can state the main result, due to [FF1, F2], concerning the Wakimoto realization at the critical level.

Theorem 4.1. There exists a homomorphism of vertex algebras

$$w_{\kappa_c}: \mathbb{V}_0 \to M_{\mathfrak{g}} \otimes \pi_0$$

such that

$$\begin{split} e_i(z) &\mapsto a_{\alpha_i}(z) + \sum_{\beta \in \Delta_+} :P^i_\beta(a^*_\alpha(z))a_\beta(z):, \\ h_i(z) &\mapsto -\sum_{\beta \in \Delta_+} \beta(h_i):a^*_\beta(z)a_\beta(z): + b_i(z), \\ f_i(z) &\mapsto \sum_{\beta \in \Delta_+} :Q^i_\beta(a^*_\alpha(z))a_\beta(z): + c_i\partial_z a^*_{\alpha_i}(z) + b_i(z)a^*_{\alpha_i}(z), \end{split}$$

where the polynomials P^i_{β}, Q^i_{β} are introduced in formulas (4.4)–(4.6).

In order to make the homomorphism w_{κ_c} coordinate-independent, we need to define the actions of the group Aut \mathcal{O} on \mathbb{V}_0 , $M_{\mathfrak{g}}$ and π_0 which are intertwined by w_{κ_c} . We already have a natural action of Aut \mathcal{O} on \mathbb{V}_0 which is induced by its action on $\widehat{\mathfrak{g}}_{\kappa_c}$ preserving the subalgebra $\mathfrak{g}[[t]]$. Next, we define the Aut \mathcal{O} -action on

 $M_{\mathfrak{g}}$ by stipulating that for each $\alpha \in \Delta_+$ the generating functions $a_{\alpha}(z)$ and $a_{\alpha}^*(z)$ transform as a one-form and a functions on D^{\times} , respectively.

Finally, we identify π_0 with the algebra Fun $\operatorname{Conn}(\Omega^{-\rho})_D$, where $\operatorname{Conn}(\Omega^{-\rho})_D$ is the space of connections on the ^{*L*}*H*-bundle $\Omega^{-\rho}$ on *D*. Namely, we write a connection from $\operatorname{Conn}(\Omega^{-\rho})_D$ as the first order operator $\partial_t + \chi(t)$, where $\chi(t)$ takes values in ^{*L*} $\mathfrak{h} \simeq \mathfrak{h}^*$. Set

$$b_i(t) = \langle \check{\alpha}_i, \chi(t) \rangle = \sum_{n < 0} b_{i,n} t^{-n-1}.$$

Then we obtain an identification

Fun Conn
$$(\Omega^{\rho})_D \simeq \mathbb{C}[b_{i,n}]_{i=1,\dots,\ell;n<0} = \pi_0.$$

Now the natural action of Aut \mathcal{O} on Fun $\operatorname{Conn}(\Omega^{\rho})_D$ gives rise to an action of Aut \mathcal{O} on π_0 .

To write down this action explicitly, suppose that we have a connection on the ^LH-bundle $\Omega^{-\rho}$ on a curve X which with respect to a local coordinate t is represented by the first order operator $\partial_t + \chi(t)$. If s is another coordinate such that $t = \varphi(s)$, then this connection will be represented by the operator

(4.8)
$$\partial_s + \varphi'(s)\chi(\varphi(s)) + \rho \cdot \frac{\varphi''(s)}{\varphi'(s)}$$

Proposition 4.2 ([F2], Cor. 5.4). The homomorphism w_{κ_c} commutes with the action of Aut \mathfrak{O} .

For a connection $\overline{\nabla}$ on the ^{*L*}*H*-bundle $\Omega^{-\rho}$, let $\overline{\nabla}^*$ be the dual connection on Ω^{ρ} . Explicitly, if $\overline{\nabla}$ is given by the operator $\partial_t + \mathbf{w}(t)$, then $\overline{\nabla}^*$ is given by the operator $\partial_t - \mathbf{w}(t)$. Sending $\overline{\nabla}$ to $\overline{\nabla}^*$, we obtain an isomorphism $\operatorname{Conn}(\Omega^{-\rho})_U \simeq$ $\operatorname{Conn}(\Omega^{\rho})_U$.

Recall the Miura transformation

$$\overline{\mathbf{b}}_{D^{\times}} : \operatorname{Conn}(\Omega^{\rho})_{D^{\times}} \to \operatorname{Op}_{{}^{L}G}(D^{\times})$$

defined in Section 3.3. Using the isomorphism $\operatorname{Conn}(\Omega^{-\rho})_U \simeq \operatorname{Conn}(\Omega^{\rho})_U$, we obtain a map

(4.9)
$$\overline{\mathbf{b}}_{D^{\times}}^* : \operatorname{Conn}(\Omega^{-\rho})_{D^{\times}} \to \operatorname{Op}_{{}^{L}G}(D^{\times})$$

which by abuse of terminology we will also refer to as Miura transformation.

Consider the center $\mathfrak{z}(\hat{\mathfrak{g}})$ of the vertex algebra \mathbb{V}_0 . Recall that we have

$$\mathfrak{z}(\widehat{\mathfrak{g}}) = \mathbb{V}_0^{\mathfrak{g}[[t]]} \simeq \operatorname{End}_{\widehat{\mathfrak{g}}_{\kappa_c}} \mathbb{V}_0.$$

Theorem 4.3 ([F2], Theorem 11.3). Under the homomorphism w_{κ_c} the center $\mathfrak{z}(\widehat{\mathfrak{g}}) \subset \mathbb{V}_0$ gets mapped to the vertex subalgebra π_0 of $M_{\mathfrak{g}} \otimes \pi_0$. Moreover, we have the following commutative diagram



where the right vertical map is induced by the Miura transformation $\overline{\mathbf{b}}_D^*$ given by (4.9).

This theorem implies the following result. Let N be a module over the vertex algebra $M_{\mathfrak{g}}$ and R a module over the vertex algebra $\pi_0 \simeq \operatorname{Fun} \operatorname{Conn}(\Omega^{\rho})_D$. Then the homomorphism w_{κ_c} gives rise to the structure of a \mathbb{V}_0 -module, and hence of a smooth $\widehat{\mathfrak{g}}_{\kappa_c}$ -module on the tensor product $N \otimes R$.

It follows from the general theory of [FB], Ch. 5, that a module over the vertex algebra $M_{\mathfrak{g}}$ is the same as a smooth module over the Weyl algebra $\mathcal{A}^{\mathfrak{g}}$, i.e., such that every vector is annihilated by $a_{\alpha,n}, a_{\alpha,n}^*$ for large enough n. Likewise, a module over the commutative vertex algebra Fun $\operatorname{Conn}(\Omega^{-\rho})_D$ is the same as a smooth module over the commutative topological algebra Fun $\operatorname{Conn}(\Omega^{-\rho})_{D^{\times}}$. Note that

Fun Conn
$$(\Omega^{-\rho})_{D^{\times}} \simeq \lim \mathbb{C}[b_{i,n}]_{i=1,\ldots,\ell;n\in\mathbb{Z}}/I_N,$$

where I_N is the ideal generated by $b_{i,n}, i = 1, \ldots, \ell; n \geq N$. A module over this algebra is called smooth if every vector is annihilated by the ideal I_N for large enough N. Thus, for each choice of $\mathcal{A}^{\mathfrak{g}}$ -module N we obtain a "semi-infinite induction" functor from the category of smooth Fun $\operatorname{Conn}(\Omega^{-\rho})_{D^{\times}}$ -modules to the category of smooth $\widehat{\mathfrak{g}}_{\kappa_c}$ -modules (on which the central element **1** acts as the identity), $R \mapsto N \otimes R$.

Now Theorem 4.3 implies

Corollary 4.4. The action of $Z(\widehat{\mathfrak{g}})$ on the $\widehat{\mathfrak{g}}_{\kappa_c}$ -module $N \otimes R$ is independent of the choice of N and factors through the homomorphism $Z(\widehat{\mathfrak{g}}) \simeq \operatorname{Fun} \operatorname{Op}_{L_G}(D^{\times}) \to \operatorname{Fun} \operatorname{Conn}(\Omega^{-\rho})_{D^{\times}}$ induced by the Miura transformation $\overline{\mathbf{b}}_{D^{\times}}^*$.

For example, for a connection $\overline{\nabla} = \partial_z + \chi(z)$ in $\operatorname{Conn}(\Omega^{-\rho})_{D^{\times}}$, let $\mathbb{C}_{\overline{\nabla}} = \mathbb{C}_{\chi(z)}$ be the corresponding one-dimensional module of Fun $\operatorname{Conn}(\Omega^{-\rho})_{D^{\times}}$. Then $M_{\mathfrak{g}} \otimes \mathbb{C}_{\chi(z)}$ is called the *Wakimoto module* of critical level corresponding to $\partial_z + \chi(z)$, and is denoted by $W_{\chi(z)}$.

We will use below the following result.

Lemma 4.5 ([FFR], Lemma 2). Let

$$\chi(z) = -\frac{\alpha_i}{z} + \sum_{n=0}^{\infty} \chi_n z^n, \qquad \chi_n \in \mathfrak{h}^*.$$

The vector $e_{i,-1}^R |0\rangle \in W_{\chi(z)}$ is $\mathfrak{g}[[t]]$ -invariant if and only if $\langle \check{\alpha}_i, \chi_0 \rangle = 0$.

4.2. Functoriality of coinvariants

We will construct eigenvectors of the generalized Gaudin Hamiltonians using functionals on the space of coinvariants of Wakimoto modules on \mathbb{P}^1 . First we explain some general facts about the compatibility of various spaces of coinvariants.

Following the general construction of [FB], Ch. 10, we define the spaces of coinvariants $H_{M_{\mathfrak{g}}}(X,(x_i),(M_i))$ (resp., $H_{\pi_0}(X,\{x_i\},(R_i))$) attached to a smooth projective curve X, points x_1, \ldots, x_p and $M_{\mathfrak{g}}$ -modules N_1, \ldots, N_p (resp., π_0 -modules R_1, \ldots, R_p). These spaces are defined in elementary terms in [FFR], Sect. 5, and in [FB], Ch. 13–14. We also define spaces of coinvariants for the tensor product vertex algebra $M_{\mathfrak{g}} \otimes \pi_0$. We have a natural identification

 $H_{M_{\mathfrak{g}} \otimes \pi_{0}}(X, (x_{i}), (N_{i} \otimes R_{i})) = H_{M_{\mathfrak{g}}}(X, (x_{i}), (N_{i})) \otimes H_{\pi_{0}}(X, (x_{i}), (R_{i})).$

The vertex algebra homomorphism $w_{\kappa_c} : \mathbb{V}_0 \to M_{\mathfrak{g}} \otimes \pi_0$ of Theorem 4.1 gives rise to a surjective map of the spaces of coinvariants

$$H_{\mathbb{V}_0}(X, (x_i), (N_i \otimes R_i)) \twoheadrightarrow H_{M_{\mathfrak{g}} \otimes \pi_0}(X, (x_i), (N_i \otimes R_i))$$

and hence a surjective map

$$(4.10) \qquad H_{\mathbb{V}_0}(X, (x_i), (N_i \otimes R_i)) \twoheadrightarrow H_{M_{\mathfrak{g}}}(X, (x_i), (N_i)) \otimes H_{\pi_0}(X, (x_i), (R_i)).$$

Now recall that in the proof of Theorem 2.7 we explained that since $\mathfrak{z}(\hat{\mathfrak{g}}) \simeq$ Fun $\operatorname{Op}_{L_G}(D)$ is the center of \mathbb{V}_0 , we have a homomorphism of algebras

$$(4.11) H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(X,(x_i),(Z(M_i))) \to \operatorname{End}_{\mathbb{C}} H_{\mathbb{V}_0}(X,(x_i),(M_i)).$$

In addition, we have an isomorphism

(4.12)
$$H_{\mathfrak{z}(\widehat{\mathfrak{g}})}(X,(x_i),(Z(M_i))) \simeq \operatorname{Fun} \operatorname{Op}_{L_G}(X,(x_i),(M_i)),$$

where $\operatorname{Op}_{L_G}(X, (x_i), (M_i))$ is the space of ^LG-opers on X which are regular on $X \setminus \{x_1, \ldots, x_N\}$ and such that their restriction to D_x^{\times} belongs to the space $\operatorname{Op}_{L_G}^{M_i}(D_{x_i}^{\times}) = \operatorname{Spec} Z(M_i)$ for all $i = 1, \ldots, N$.

We can describe the space of coinvariants $H_{\pi_0}(X, (x_i), (R_i))$ for the commutative vertex algebra $\pi_0 \simeq \operatorname{Fun} \operatorname{Conn}(\Omega^{-\rho})_D$ in similar terms. If R is a π_0 -module, or equivalently, a smooth $\operatorname{Fun} \operatorname{Conn}(\Omega^{-\rho})_{D^{\times}}$ -module, let $\pi_0(R)$ be the image of the corresponding homomorphism

Fun $\operatorname{Conn}(\Omega^{-\rho})_{D^{\times}} \to \operatorname{End}_{\mathbb{C}} R$.

Let $\operatorname{Conn}(\Omega^{-\rho})(X, (x_i), (R_i))$ be the space of connections on $\Omega^{-\rho}$ on $X \setminus \{x_1, \ldots, x_N\}$ whose restriction to $D_{x_i}^{\times}$ belongs to the space $\operatorname{Conn}(\Omega^{-\rho})_{D_{x_i}^{\times}}^{R_i} = \operatorname{Spec} \pi_0(R_i)$. Then we have an isomorphism

$$H_{\pi_0}(X, (x_i), (R_i)) \simeq \operatorname{Fun} \operatorname{Conn}(\Omega^{-\rho})(X, (x_i), (R_i)).$$

It follows from Corollary 4.4 that if $M_i = N_i \otimes R_i$, then $Z(N_i \otimes R_i)$ is the image of $Z(\hat{\mathfrak{g}}) \simeq \operatorname{Fun}_{L_G}(D^{\times})$ under the homomorphism

$$\operatorname{Fun}_{L_G}(D^{\times}) \xrightarrow{\mathbf{b}_{D^{\times}}} \operatorname{Fun}\operatorname{Conn}(\Omega^{-\rho})_{D^{\times}} \to \operatorname{End}_{\mathbb{C}}R_i,$$

where the first map is induced by the Miura transformation.
Recall that in Section 3.2 we introduced the maps

$$\mathbf{b}_U : \operatorname{Conn}(\Omega^{\rho})_U \to \operatorname{MOp}_{L_G}(U)_{\operatorname{gen}}, \quad \overline{\mathbf{b}}_U : \operatorname{Conn}(\Omega^{\rho})_U \to \operatorname{Op}_{L_G}(U)$$

Using the identification $\operatorname{Conn}(\Omega^{\rho})_U \simeq \operatorname{Conn}(\Omega^{-\rho})_U$, we obtain maps

 $\mathbf{b}_U^* : \operatorname{Conn}(\Omega^{-\rho})_U \to \operatorname{MOp}_{L_G}(U)_{\operatorname{gen}}, \qquad \overline{\mathbf{b}}_U^* : \operatorname{Conn}(\Omega^{-\rho})_U \to \operatorname{Op}_{L_G}(U).$

The second map is the Miura transformation. In the case when $U = X \setminus \{x_1, \ldots, x_N\}$ the Miura transformation $\overline{\mathbf{b}}_U^*$ restricts to a map

$$\operatorname{Conn}(\Omega^{-\rho})(X,(x_i),(R_i)) \to \operatorname{Op}_{{}^{L}G}(X,(x_i),(N_i \otimes R_i)),$$

and hence gives rise to a homomorphism

(4.13) $\operatorname{Fun}\operatorname{Op}_{L_G}(X,(x_i),(N_i\otimes R_i))\to\operatorname{Fun}\operatorname{Conn}(\Omega^{-\rho})(X,(x_i),(R_i)),$

which does not depend on the N_i 's.

Now we have an action of the algebra $\operatorname{Op}_{L_G}(X, (x_i), (N_i \otimes R_i))$ on both sides of the map (4.10). The action on the left-hand side comes from the homomorphisms (4.11) and (4.12). The action on the right-hand side comes from the homomorphism (4.13).

The functoriality of the coinvariants implies the following

Lemma 4.6. The map (4.10) commutes with the action on both sides of the algebra $\operatorname{Op}_{L_G}(X, (x_i), (N_i \otimes R_i)).$

Consider the special case where all modules $R_i, i = 1, ..., N$, are one-dimensional, i.e., $R_i = \mathbb{C}_{\overline{\nabla}_i}$, where $\overline{\nabla}_i$ is a point in $\operatorname{Conn}(\Omega^{-\rho})_{D_{x_i}^{\times}}$, and R_i is the representation obtained from the homomorphism $\operatorname{Fun}\operatorname{Conn}(\Omega^{-\rho})_{D_{x_i}^{\times}}$ induced by $\overline{\nabla}_i$. In this case we have the following description of the space $H_{\pi_0}(X, (x_i), (\mathbb{C}_{\overline{\nabla}_i}))$ which follows from the general results on coinvariants of commutative vertex algebras from Sect. 9.4 of [FB] (see also [FFR], Prop. 4, in the case when $X = \mathbb{P}^1$).

Proposition 4.7. The space $H_{\pi_0}(X, (x_i), (\mathbb{C}_{\overline{\nabla}_i}))$ is one-dimensional if and only if there exists a connection $\overline{\nabla}$ on the ^LH-bundle $\Omega^{-\rho}$ on $X \setminus \{x_1, \ldots, x_p, \infty\}$ such that the restriction of $\overline{\nabla}$ to the punctured disc at each x_i is equal to $\overline{\nabla}_i, i = 1, \ldots, N$. Otherwise, $H_{\pi_0}(X, (x_i), (\mathbb{C}_{\overline{\nabla}_i})) = 0$.

Suppose that such a connection $\overline{\nabla}$ exists and the space $H_{\pi_0}(X, (x_i), (\mathbb{C}_{\overline{\nabla}_i}))$ is one-dimensional. Then the map (4.10) reads

$$H_{\mathbb{V}_0}(X,(x_i),(N_i\otimes\mathbb{C}_{\overline{\nabla}_i}))\twoheadrightarrow H_{M_{\mathfrak{g}}}(X,(x_i),(N_i)).$$

Composing this map with any linear functional on $H_{M_{\mathfrak{g}}}(X,(x_i),(N_i))$, we obtain a linear functional

$$\phi: H_{\mathbb{V}_0}(X, (x_i), (N_i \otimes \mathbb{C}_{\overline{\nabla}_i})) \to \mathbb{C}.$$

The algebra $\mathfrak{z}(\widehat{\mathfrak{g}})_u = \operatorname{Fun} \operatorname{Op}_{L_G}(D_u)$ acts on the space $H_{\mathbb{V}_0}(X, (x_i), (N_i \otimes \mathbb{C}_{\overline{\nabla}_i}))$, and its action factors through the homomorphism (2.9)

Fun
$$\operatorname{Op}_{L_G}(D_u) \to \operatorname{Fun} \operatorname{Op}_{L_G}(X, (x_i), (N_i \otimes \mathbb{C}_{\overline{\nabla}_i}))$$

According to Lemma 4.6, this action factors through the homomorphism (4.13). In our case $R_i = \mathbb{C}_{\overline{\nabla}_i}$, and so $\operatorname{Conn}(\Omega^{-\rho})(X, (x_i), (R_i))$ is a point corresponding to the connection $\overline{\nabla}$. It follows from the definition that $\operatorname{Op}_{L_G}(X, (x_i), (N_i \otimes \mathbb{C}_{\overline{\nabla}_i}))$ is also a point, corresponding to the oper which is the image of $\overline{\nabla}$ under the Miura transformation $\overline{\mathbf{b}}_{X \setminus \{x_1, \dots, x_N\}}^*$. Thus, we obtain:

Proposition 4.8. The functional ϕ is an eigenvector of the algebra $\mathfrak{g}(\widehat{\mathfrak{g}})_u \simeq \operatorname{Op}_{L_G}(D_u)$, and the corresponding homomorphism $\operatorname{Fun} \operatorname{Op}_{L_G}(D_u) \to \mathbb{C}$ is defined by the point $\overline{\mathbf{b}}_{D_u}^*(\overline{\nabla}|_{D_u}) \in \operatorname{Op}_{L_G}(D_u)$.

4.3. The case of \mathbb{P}^1

We will use Proposition 4.8 in the case when $X = \mathbb{P}^1$, equipped with a global coordinate t. Consider a collection of distinct points x_1, \ldots, x_p and ∞ on \mathbb{P}^1 . As the $M_{\mathfrak{g}}$ -module N_i attached to $x_i, i = 1, \ldots, p$, we will take $M_{\mathfrak{g}}$. As the $M_{\mathfrak{g}}$ -module N_{∞} attached to the point ∞ we will take another module $M'_{\mathfrak{g}}$ generated by a vector $|0\rangle'$ such that

$$a_{\alpha,n}|0\rangle' = 0, \quad n > 0; \qquad a_{\alpha,n}^*|0\rangle' = 0, \quad n \ge 0$$

As the π_0 -module R_i attached to $x_i, i = 1, \ldots, p$, we will take the one-dimensional module $\mathbb{C}_{\overline{\nabla}_i} = \mathbb{C}_{\chi_i(z)}$, where $\overline{\nabla}_i = \partial_z + \chi_i(z), \ \chi_i(z) \in \mathfrak{h}^*((z))$ and as module R_∞ attached to the point ∞ we will take $\mathbb{C}_{\chi_\infty(z)}$. Thus, $M_i \otimes R_i$ is the Wakimoto module $W_{\chi_i(z)}$ for all $i = 1, \ldots, p$. We denote the module $M'_{\mathfrak{g}} \otimes \mathbb{C}_{\chi_\infty(z)}$ by $W'_{\chi_\infty(z)}$.

We have the following special case of Proposition 4.7:

Proposition 4.9 ([FFR], Prop. 4). The space $H_{M_{\mathfrak{g}}}(\mathbb{P}^1; (x_i), \infty; (M_{\mathfrak{g}}), M'_{\mathfrak{g}})$ is onedimensional and the projection of the vector $|0\rangle^{\otimes N} \otimes |0\rangle'$ on it is non-zero.

The space $H_{\pi_0}(\mathbb{P}^1, (x_i), \infty; (\mathbb{C}_{\chi_i(z)}), \mathbb{C}_{\chi_\infty}(z))$ is one-dimensional if and only if there exists a connection $\overline{\nabla}$ on the ^LH-bundle $\Omega^{-\rho}$ on \mathbb{P}^1 which is regular on $\mathbb{P}^1 \setminus \{x_1, \ldots, x_p, \infty\}$ and such that the restriction of $\overline{\nabla}$ to the punctured disc at each x_i is equal to $\partial_t + \chi_i(t - x_i)$, and its restriction to the punctured disc at ∞ is equal to $\partial_{t^{-1}} + \chi_\infty(t^{-1})$.

Otherwise, $H_{\pi_0}(\mathbb{P}^1, (x_i), \infty; (\mathbb{C}_{\chi_i(z)}), \mathbb{C}_{\chi_\infty}(z)) = 0.$

Formula (4.8) shows that if we have a connection on $\Omega^{-\rho}$ over \mathbb{P}^1 whose restriction to $\mathbb{P}^1 \setminus \infty$ is represented by the operator $\partial_t + \chi(t)$, then its restriction to the punctured disc D_{∞}^{\times} at ∞ reads, with respect to the coordinate $u = t^{-1}$, is represented by the operator

$$\partial_u - u^{-2}\chi(u^{-1}) - 2\rho u^{-1}.$$

We will choose $\chi_i(z)$ to be of the form

$$\chi_i(z) = \chi_i + \sum_{n \ge 0} \chi_{i,n} z^n,$$

and $\chi_{\infty}(z)$ to be of the form

$$\chi_{\infty}(z) = \chi_{\infty} + \sum_{n \ge 0} \chi_{\infty,n} z^n.$$

The condition of the proposition is then equivalent to saying that the restriction of $\overline{\nabla}$ to $\mathbb{P}^1 \setminus \infty$ is represented by the operator

$$\partial_t + \sum_{i=1}^p \frac{\chi_i}{t - z_i},$$

and that $\chi_i(t-z_i)$ is the expansion of $\chi(t)$ at $z_i, i = 1, \ldots, p$, while $\chi_{\infty}(t^{-1})$ is the expansion of $-t^2\chi(t) - 2\rho t$ in powers of t^{-1} . Note that then we have $\chi_{\infty}(u) = \chi_{\infty}/u + \ldots$, where $\chi_{\infty} = -2\rho - \sum_{i=1}^{p} \chi_i$. By Proposition 4.9, the spaces of coinvariants $H_{M_{\mathfrak{g}}}(\mathbb{P}^1; (x_i), \infty; (M_{\mathfrak{g}}), M'_{\mathfrak{g}})$

By Proposition 4.9, the spaces of coinvariants $H_{M_{\mathfrak{g}}}(\mathbb{P}^1; (x_i), \infty; (M_{\mathfrak{g}}), M'_{\mathfrak{g}})$ and $H_{\pi_0}(\mathbb{P}^1; (x_i), \infty; (\mathbb{C}_{\chi_i(z)}), \mathbb{C}_{\chi_\infty}(z))$ are one-dimensional. Hence the map (4.10) gives rise to a non-zero linear functional

(4.14)
$$\tau_{(x_i)}: H_{\mathbb{V}_0}(\mathbb{P}^1; (x_i), \infty; (W_{\chi_i(z)}), W'_{\chi_\infty(z)}) \to \mathbb{C}.$$

We normalize it so that its value on $|0\rangle^{\otimes N} \otimes |0\rangle'$ is equal to 1.

The algebra $\mathfrak{z}(\widehat{\mathfrak{g}})_u = \operatorname{Fun} \operatorname{Op}_{L_G}(D_u)$ acts on this space. By Proposition 4.8, $\tau_{(x_i)}$ is an eigenvector of this algebra and the corresponding homomorphism

Fun
$$\operatorname{Op}_{L_G}(D_u) \to \mathbb{C}$$

is defined by the point $\overline{\mathbf{b}}_{D_u}^*(\overline{\nabla}|_{D_u}) \in \operatorname{Op}_{{}^LG}(D_u).$

Let us fix highest weights of $\mathfrak{g}, \lambda_1, \ldots, \lambda_N$, and a set of simple roots of $\mathfrak{g}, \alpha_{i_1}, \ldots, \alpha_{i_m}$. Consider a connection $\overline{\nabla}$ on $\Omega^{-\rho}$ on \mathbb{P}^1 whose restriction to $\mathbb{P}^1 \setminus \infty$ is equal to $\partial_t + \lambda(t)$, where

(4.15)
$$\lambda(t) = \sum_{i=1}^{N} \frac{\lambda_i}{t - z_i} - \sum_{j=1}^{m} \frac{\alpha_{i_j}}{t - w_j}$$

Denote by $\lambda_i(t-z_i)$ the expansions of $\lambda(t)$ at the points $z_i, i = 1, ..., N$, and by $\mu_j(t-w_j)$ the expansions of $\lambda(t)$ at the points $w_j, j = 1, ..., m$. We have:

$$\lambda_i(z) = \frac{\lambda_i}{z} + \cdots, \qquad \mu_j(z) = -\frac{\alpha_{i_j}}{z} + \mu_{j,0} + \cdots,$$

where

(4.16)
$$\mu_{j,0} = \sum_{i=1}^{N} \frac{\lambda_i}{w_j - z_i} - \sum_{s \neq j} \frac{\alpha_{i_s}}{w_j - w_s}.$$

λī

Finally, let $\lambda_{\infty}(t^{-1})$ be the expansion of $-t^2\lambda(t) - 2\rho t$ in powers of t^{-1} . Thus we have

(4.17)
$$\lambda_{\infty}(z) = z^{-1} \left(-\sum_{i=1}^{N} \lambda_i + \sum_{j=1}^{m} \alpha_{i_j} - 2\rho \right) + \cdots$$

We have a linear functional $\tau_{(z_i),(w_i)}$ on the corresponding space of coinvariants

$$H_{\mathbb{V}_0}(\mathbb{P}^1; (z_i), (w_j), \infty; (W_{\lambda_i(z)}), (W_{\mu_j(z)}), W'_{\lambda_\infty(z)}).$$

In particular, this space is non-zero. The functional $\tau_{(z_i),(w_j)}$ is an eigenvector of $\mathfrak{z}(\widehat{\mathfrak{g}})_u = \operatorname{Fun} \operatorname{Op}_{L_G}(D_u)$, and its eigenvalue is the ^LG-oper on D_u which is the restriction to D_u of the oper in $\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),(w_j),\infty;(\lambda_i),(-\alpha_{i_j}),\lambda_{\infty}}$ given by the Miura transformation of the connection $\partial_t + \lambda(t)$. We will now use it to construct an eigenvector of the Gaudin algebra.

4.4. Bethe vectors

By Lemma 4.5, the vectors $e_{i_j,-1}^R |0\rangle \in W_{\mu_j(z)}$ are $\mathfrak{g}[[t]]$ -invariant if and only if the equations $\langle \check{\alpha}_{i_j}, \mu_{j,0} \rangle = 0$ are satisfied, where $\mu_{j,0}$ is the constant coefficient in the expansion of $\lambda(t)$ at w_j given by formula (4.16), i.e., the following system of equations is satisfied

(4.18)
$$\sum_{i=1}^{N} \frac{\langle \check{\alpha}_{i_j}, \lambda_i \rangle}{w_j - z_i} - \sum_{s \neq j} \frac{\langle \check{\alpha}_{i_j}, \alpha_{i_s} \rangle}{w_j - w_s} = 0, \quad j = 1, \dots, m.$$

They are called the *Bethe Ansatz equations*. This is a system of equations on the complex numbers w_j , j = 1, ..., m, to each of which we attach a simple root α_{i_j} . We have an obvious action of a product of symmetric groups permuting the points w_j corresponding to simple roots of the same kind. In what follows, by a *solution* of the Bethe Ansatz equations we will understand a solution defined up to these permutations. We will adjoin to the set of all solutions associated to all possible collections $\{\alpha_{i_j}\}$ of simple roots of \mathfrak{g} , the unique "empty" solution, corresponding to the empty set of simple roots.

Suppose that these equations are satisfied. Then we obtain a homomorphism of $\widehat{\mathfrak{g}}_{\kappa_c}\text{-modules}$

$$\bigotimes_{i=1}^{N} W_{\lambda_{i}(z)} \otimes \mathbb{V}_{0}^{\otimes m} \otimes W'_{\lambda_{\infty}(z)} \to \bigotimes_{i=1}^{N} W_{\lambda_{i}(z)} \otimes \bigotimes_{j=1}^{m} W_{\mu_{j}(z)} \otimes W'_{\lambda_{\infty}(z)},$$

which sends the vacuum vector in the *j*th copy of \mathbb{V}_0 to $e_{i_j,-1}^H|_0 \in W_{\mu_j(z)}$. Hence we obtain the corresponding map of the spaces of coinvariants. But the insertion of \mathbb{V}_0 does not change the space of coinvariants. Hence the first space of coinvariants is isomorphic to

$$H_{\mathbb{V}_0}(\mathbb{P}^1;(z_i),\infty;(W_{\lambda_i(z)}),W'_{\lambda_\infty(z)}) = H((W_{\lambda_i(z)}),W'_{\lambda_\infty(z)})$$

Therefore, composing this map with $\tau_{(z_i),(w_j)}$, we obtain a linear functional

(4.19)
$$\widetilde{\tau}_{(z_i),(w_j)}: H((W_{\lambda_i(z)}), W'_{\lambda_{\infty}(z)}) \to \mathbb{C}$$

By construction, $\tilde{\tau}_{(z_i),(w_j)}$ is an eigenvector of $\mathfrak{z}(\hat{\mathfrak{g}})_u = \operatorname{Fun} \operatorname{Op}_{L_G}(D_u)$, and its eigenvalue is the ^LG-oper on D_u which is the restriction of an oper on \mathbb{P}^1 with singularities given by the Miura transformation of the connection $\partial_t + \lambda(t)$.

Let $U = \mathbb{P}^1 \setminus \{z_1, \ldots, z_N, w_1, \ldots, w_m, \infty\}$ and $\operatorname{Conn}(\Omega^{-\rho})_{(z_i), \infty; (\lambda_i), \lambda_{\infty}}^{\operatorname{gen}}$ be the subset of $\operatorname{Conn}(\Omega^{-\rho})_U$, consisting of all connections on $\Omega^{-\rho}$ on \mathbb{P}^1 with regular

singularities of the form $\partial_t + \lambda(t)$, where $\lambda(t)$ is given by (4.15), such that their Miura transformation is a ^LG-oper in

(4.20)
$$\operatorname{Op}_{L_{G}}^{\mathrm{RS}}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}} \subset \operatorname{Op}_{L_{G}}^{\mathrm{RS}}(\mathbb{P}^{1})_{(z_{i}),(w_{j}),\infty;(\lambda_{i}),(-\alpha_{i_{j}}),\lambda_{\infty}}.$$

In other words, $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$ consists of those connections $\partial_t + \lambda(t)$, where $\lambda(t)$ is given by (4.15), whose Miura transformation is a ^LG-oper on Uwhose restriction to $D_{w_j}^{\times}$ belongs to $\operatorname{Op}_{L_G}(D_{w_j}) \subset \operatorname{Op}_{L_G}^{\operatorname{RS}}(D_{w_j}^{\times})$.

Thus, we have a map

$$(4.21) \qquad \overline{\mathbf{b}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^* : \operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}} \to \operatorname{Op}_{L_G}^{\operatorname{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$$

Proposition 4.10 ([FFR], Prop. 6). There is a bijection between the set of solutions of the Bethe Ansatz equations (4.18) and the set $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$.

Proof. Applying the Miura transformation to the connection $\partial_t + \lambda(t)$, where $\lambda(t)$ is given by (4.15), we obtain a ^LG-oper on \mathbb{P}^1 which has regular singularities at the points z_1, \ldots, z_N , and ∞ with residues $\varpi(\lambda_1 + \rho), \ldots, \varpi(\lambda_N + \rho)$ and $\varpi(\lambda_\infty + \rho)$, respectively, and at the points w_1, \ldots, w_m with residues $\varpi(-\alpha_{i_j} + \rho) = \varpi(\rho)$. But by Lemma 3.5, this oper is regular at the points w_1, \ldots, w_m if and only if the equations $\langle \check{\alpha}_{i_j}, \mu_{j,0} \rangle = 0$ are satisfied, where $\mu_{j,0}$ is the constant coefficient in the expansion of $\lambda(t)$ at w_j given by formula (4.16). These equations are precisely the Bethe Ansatz equations (4.18).

Now let w_1, \ldots, w_m be a solution of the Bethe Ansatz equations (4.18). Consider the corresponding connection $\partial_t + \lambda(t)$ in $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$, where $\lambda(t)$ is of the form (4.15). Recall that the linear functional $\tilde{\tau}_{(z_i),(w_j)}$ from formula (4.19) is an eigenvector of $\mathfrak{z}(\widehat{\mathfrak{g}})_u = \operatorname{Fun}\operatorname{Op}_{L_G}(D_u)$, and its eigenvalue is the LG -oper on D_u obtained by restriction of the oper in $\operatorname{Op}_{L_G}^{\operatorname{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ given by the Miura transformation of the connection $\partial_t + \lambda(t)$.

Therefore we obtain that the action of $\mathfrak{g}(\widehat{\mathfrak{g}})_u$ on the space $H((W_{\lambda_i(z)}), W'_{\lambda_\infty(z)})$ factors through the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, which acts on it according to the character corresponding to the LG -oper $\overline{\mathbf{b}}^*_{(z_i),\infty;(\lambda_i),\lambda_\infty}(\partial_t + \lambda(t))$.

We wish to use this fact to construct eigenvectors of the algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ on the tensor product of Verma modules $\bigotimes_{i=1}^N M_{\lambda_i}$. For $\lambda(z) \in \mathfrak{h}^* \otimes z^{-1}\mathbb{C}[[z]]$ denote by λ_{-1} the residue of $\lambda(z)dz$ at z = 0. Let

For $\lambda(z) \in \mathfrak{h}^* \otimes z^{-1}\mathbb{C}[[z]]$ denote by λ_{-1} the residue of $\lambda(z)dz$ at z = 0. Let $\widetilde{W}_{\lambda(z)}$ the subspace of a Wakimoto module $W_{\lambda(z)}$, which is generated from the vector $|0\rangle$ by the operators $a_{\alpha,0}^*, \alpha \in \Delta_+$. This space is stable under the action of the constant subalgebra \mathfrak{g} of $\widehat{\mathfrak{g}}_{\kappa_c}$. It follows from the formulas defining the action of $\widehat{\mathfrak{g}}_{\kappa_c}$ on $W_{\lambda(z)}$ that as a \mathfrak{g} -module, $\widetilde{W}_{\lambda(z)}$ is isomorphic to the module $M_{\lambda-1}^*$ contragredient to the Verma module over \mathfrak{g} with highest weight λ_{-1} . Likewise, the subspace $\widetilde{W}'_{\lambda(z)}$ generated from the vector $|0\rangle'$ by the operators $a_{\alpha,0}, \alpha \in \Delta_+$, is stable under the action of the constant subalgebra \mathfrak{g} of $\widehat{\mathfrak{g}}_{\kappa_c}$ and is isomorphic as a \mathfrak{g} -module to the module $M'_{2\rho+\lambda_{-1}}$ the Verma module over \mathfrak{g} with *lowest* weight $2\rho + \lambda_{-1}$.

Let \mathbb{M}^*_{χ} be the $\widehat{\mathfrak{g}}_{\kappa_c}$ -module induced from M^*_{χ} , and let \mathbb{M}'_{χ} be the $\widehat{\mathfrak{g}}_{\kappa_c}$ -module induced from M'_{χ} (see Section 2.2). The embeddings of \mathfrak{g} -modules $M^*_{\lambda_{-1}} \to W_{\lambda(z)}$ and $M'_{2\rho+\lambda_{-1}} \to W'_{\lambda(z)}$ induce homomorphisms of $\widehat{\mathfrak{g}}_{\kappa_c}$ -modules $\mathbb{M}^*_{\lambda_{-1}} \to W_{\lambda(z)}$ and $\mathbb{M}'_{2\rho+\lambda_{-1}} \to W'_{\lambda(z)}$. Thus, we obtain a homomorphism

$$\bigotimes_{i=1}^{N} \mathbb{M}_{\lambda_{i}}^{*} \otimes \mathbb{M}_{2\rho+\lambda_{\infty}}^{\prime} \to \bigotimes_{i=1}^{N} W_{\lambda_{i}(z)} \otimes W_{\lambda_{\infty}(z)}^{\prime},$$

where

$$\lambda_{\infty} = -2\rho - \sum_{i=1}^{N} \lambda_i + \sum_{j=1}^{m} \alpha_{i_j}.$$

This homomorphism gives rise to a map of the corresponding spaces of coinvariants

$$H((\mathbb{M}^*_{\lambda_i}), \mathbb{M}'_{2\rho+\lambda_{\infty}}) \to H((W_{\lambda_i(z)}), W'_{\lambda_{\infty}(z)}).$$

The composition of this map with the functional $\widetilde{\tau}_{(z_i),(w_j)}$ defined above is a linear functional

$$\overline{\tau}_{(z_i),(w_j)}: H((\mathbb{M}^*_{\lambda_i}), \mathbb{M}'_{2\rho+\lambda_{\infty}}) \to \mathbb{C}.$$

But we have an isomorphism

$$H((\mathbb{M}^*_{\lambda_i}), \mathbb{M}'_{2\rho+\lambda_{\infty}}) \simeq (\bigotimes_{i=1}^N M^*_{\lambda_i} \otimes M'_{2\rho+\lambda_{\infty}})/\mathfrak{g} \simeq (\bigotimes_{i=1}^N M^*_{\lambda_i})_{-2\rho-\lambda_{\infty}}/\mathfrak{n}_{-},$$

where the subscript indicates the subspace of weight $-2\rho - \lambda_{\infty} = \sum_{i=1}^{N} \lambda_i - \sum_{i=1}^{m} \alpha_{i_i}$. Thus, we obtain an \mathfrak{n}_{-} -invariant functional

$$\psi(w_1^{i_1},\ldots,w_m^{i_m}):(\bigotimes_{i=1}^N M_{\lambda_i}^*)_{-2\rho-\lambda_\infty}\to\mathbb{C},$$

or equivalently an n_+ -invariant vector

$$\phi(w_1^{i_1},\ldots,w_m^{i_m}) \in \bigotimes_{i=1}^N M_{\lambda_i}$$

of weight $\sum_{i=1}^{N} \lambda_i - \sum_{j=1}^{m} \alpha_{i_j}$.

Recall that $\tilde{\tau}_{(z_i),(w_j)}$ is an eigenvector of the algebra $\mathfrak{z}(\widehat{\mathfrak{g}})_u = \operatorname{Fun} \operatorname{Op}_{L_G}(D_u)$, and its eigenvalue is the LG -oper on \mathbb{P}^1 given by the Miura transformation of the connection $\partial_t + \lambda(t)$, where $\lambda(t)$ is given by (4.15). Hence $\phi(w_1^{i_1}, \ldots, w_m^{i_m})$ is also an eigenvector of $\mathfrak{z}(\widehat{\mathfrak{g}})_u$ with the same eigenvalue. But we know from the proof of Theorem 2.7 that the action of $\mathfrak{z}(\widehat{\mathfrak{g}})_u$ on the space of coinvariants $H((\mathbb{M}^*_{\lambda_i}), \mathbb{M}'_{2\rho+\lambda_\infty})$ factors through the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ and further through $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_\infty}(\mathfrak{g})$. Therefore $\phi(w_1^{i_1},\ldots,w_m^{i_m})$ is an eigenvector of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, and the eigenvalues of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ on this vector are encoded by the LG -oper obtained by the Miura transformation of the connection $\partial_t + \lambda(t)$, where $\lambda(t)$ is given by (4.15).

Let us summarize our results.

Theorem 4.11 ([FFR], Theorem 3). If the Bethe Ansatz equations (4.18) are satisfied, then the vector

$$(w_1^{i_1},\ldots,w_m^{i_m})\in\otimes_{i=1}^N M_\lambda$$

 $\phi(w_1^{i_1}, \dots, w_m^{i_m}) \in \bigotimes_{i=1}^N M_{\lambda_i}$ is an eigenvector of the algebra $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_\infty}(\mathfrak{g})$. Its eigenvalues define a point in

Spec
$$\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} = \operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$$

which is the image of the connection $\partial_t + \lambda(t)$ with $\lambda(t)$ as in formula (4.15), under the Miura transformation $\overline{\mathbf{b}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^*$ given by formula (4.21).

The vector $\phi(w_1^{i_1}, \ldots, w_m^{i_m})$ is called the *Bethe vector* corresponding to a solution of the Bethe Ansatz equations (4.18), or equivalently, an element of the set $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$. An explicit formula for this vector is given in [FFR], Lemma 3 (see also [ATY]):

(4.22)

$$\phi(w_1^{i_1}, \dots, w_m^{i_m}) = (-1)^m \sum_{p=(I^1,\dots,I^N)} \prod_{j=1}^N \frac{F_{i_1^j}^{(j)} F_{i_2^j}^{(j)} \dots F_{i_{a_j}^j}^{(j)}}{(w_{i_1^j} - w_{i_2^j})(w_{i_2^j} - w_{i_3^j}) \dots (w_{i_{a_j}^j} - z_j)} v_{\lambda_1} \otimes \dots v_{\lambda_N}$$

Here the summation is taken over all *ordered* partitions $I^1 \cup I^2 \cup \ldots \cup I^N$ of the set $\{1, \ldots, m\}$, where $I^j = \{i_1^j, i_2^j, \ldots, i_{a_i}^j\}$.

The fact that this vector is an eigenvector of the Gaudin Hamiltonians Ξ_i has also been established by other methods in [BaFl, RV]. But Theorem 4.11 gives us much more: it tells us that this vector is also an eigenvector of all other generalized Gaudin Hamiltonians, and identifies the ${}^{L}G$ -oper encoding their eigenvalues on this vector as the Miura transformation of the connection $\partial_t + \lambda(t)$. We recall that this last result follows from Corollary 4.4 which states that the central characters on the Wakimoto modules are obtained via the Miura transformation.

5. The case of finite-dimensional g-modules

In this section we consider the eigenvectors of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ on the space G

$$H((V_{\lambda_i}), V_{\lambda_{\infty}}) = \left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)$$

(recall that V_{λ} denotes the irreducible finite-dimensional g-module with a dominant integral highest weight λ). Recall that the action of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ on this space factors through the algebra $\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$, so that we have a homomorphism

$$\mathfrak{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} \to \operatorname{End}_{\mathbb{C}} \bigotimes_{i=1}^{N} V_{\lambda_i} \otimes V_{\lambda_{\infty}}.$$

The image of this homomorphism was denoted by $\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$. Then the set of all common eigenvalues of the algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ acting on $\left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)^G$ (without multiplicities) is precisely the spectrum of the algebra $\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$.

By Theorem 2.7,(3), the algebra $\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$ is a quotient of the algebra of functions on the space $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$, so that its spectrum injects into $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. Thus, we have there following result:

Proposition 5.1. There is an injective map

(5.1)
$$\operatorname{Spec} \overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g}) \hookrightarrow \operatorname{Op}_{{}^LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$$

In other words, each common eigenvalue of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ that occurs in the space $\left(\bigotimes_{i=1}^{N} V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)^G$ is encoded by a point of $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$, i.e., by a ^LG-oper on \mathbb{P}^1 with regular singularities at z_1, \ldots, z_N, ∞ , with residues $\lambda_1, \ldots, \lambda_N, \lambda_{\infty}$ at those points and trivial monodromy representation

$$\pi_1(\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}) \to {}^L G.$$

We wish to construct an inverse map from $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ to the spectrum of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ on $\left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)^G$.

We start with a more intrinsic geometric description of $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$.

5.1. Geometric description of opers without monodromy

In order to simplify our notation, in this section we will use the symbol G instead of ${}^{L}G$ to denote our group.

Suppose that we have a *G*-oper τ : $(\mathcal{F}, \nabla, \mathcal{F}_B)$ on a curve *U*. Consider the associated bundle of flag manifolds

$$(G/B)_{\mathcal{F}}=\mathfrak{F}\underset{G}{\times}G/B\simeq\mathfrak{F}_{B}\underset{B}{\times}G/B=(G/B)_{\mathfrak{F}_{B}}$$

The reduction \mathcal{F}_B gives rise to a section of this bundle.

Now suppose that the monodromy representation $\pi_1(U) \to G$ corresponding to this oper is trivial. Picking a point $x \in U$, we may then use the connection ∇ to identify the fibers of the oper bundle \mathcal{F} at all points of U with the fiber \mathcal{F}_x at x. Choosing a trivialization $i_x : G \xrightarrow{\sim} \mathcal{F}_x$ of \mathcal{F}_x , we then obtain a trivialization of the bundle \mathcal{F} , and hence of the bundle $(G/B)_{\mathcal{F}}$. Let us fix i_x and the corresponding trivialization of \mathcal{F} . Then the *B*-reduction \mathcal{F}_B gives rise to a map $\phi_{\tau} : U \to G/B$.

The oper condition may be described as follows. Define an open *B*-orbit $\mathbf{O} \subset [\mathfrak{n}, \mathfrak{n}]^{\perp}/\mathfrak{b} \subset \mathfrak{g}/\mathfrak{b}$, consisting of vectors which are stabilized by $N \subset B$, and such that all of their negative simple root components, with respect to the adjoint action of H = B/N, are non-zero. This orbit may also be described as the *B*-orbit of the sum of the projections of simple root generators f_i of any nilpotent subalgebra \mathfrak{n}_- , which is in generic position with \mathfrak{b} , onto $\mathfrak{g}/\mathfrak{b}$. The torus H acts simply transitively on \mathbf{O} , and so \mathbf{O} is an H-torsor.

Define an ℓ -dimensional distribution $T_{\mathrm{Op}}G/B$ in the tangent bundle to G/B as follows. The tangent space to gB is identified with the quotient $\mathfrak{g}/g\mathfrak{b}g^{-1}$. The fiber of $T_{\mathrm{Op}}G/B$ at gB is by definition its subspace $[\mathfrak{n},\mathfrak{n}]^{\perp}/\mathfrak{b}$. It contains an open dense subset $g\mathbf{O}g^{-1}$. We denote by $\mathring{T}_{\mathrm{Op}}G/B$ the open dense subset of $T_{\mathrm{Op}}G/B$ whose fiber at gB is $g\mathbf{O}g^{-1}$.

A map $\phi: U \to G/B$ is said to satisfy the oper condition if at each point of U the tangent vector to ϕ belongs to $\mathring{T}_{\text{Op}}G/B$.

For example, if $G = PGL_2$, then the flag manifold is \mathbb{P}^1 , and a map $\phi : U \to \mathbb{P}^1$ satisfies the oper condition if and only if it has a non-vanishing differential everywhere on X.

If τ is an oper, then the map $\phi_{\tau} : U \to G/B$ satisfies the oper condition (note that $\overset{\circ}{T}_{Op}G/B$ is *G*-invariant, and therefore this condition is independent of the trivialization ι_x). Conversely, if we are given a map $\phi : U \to G/B$ satisfying the oper condition, then the triple $(\mathcal{F}, \nabla, \mathcal{F}_B)$, where \mathcal{F} is the trivial *G*-bundle on U, ∇ is the trivial connection, and \mathcal{F}_B is the *B*-reduction of \mathcal{F} defined by the map ϕ , is a *G*-oper. Thus, we obtain a bijection between the set of all *G*-opers on U with trivial monodromy representation and the set of equivalence classes, with respect to the action of *G* on G/B, of maps $\phi : U \to G/B$ satisfying the oper condition.

Now let $\check{\lambda}$ be a dominant integral coweight of G. A map $D^{\times} \to H$ is said to vanish at the origin to order $\check{\lambda}$ if it may be obtained as the composition of an embedding $D^{\times} \to C^{\times}$ at the origin and the cocharacter $\check{\lambda} : \mathbb{C}^{\times} \to H$. We will say that a map $\phi : U \to G/B$ satisfies the $\check{\lambda}$ -oper condition at $y \in U$ if the restriction of ϕ to the punctured disc D_y^{\times} satisfies the oper condition and the corresponding map $\phi_{D_y^{\times}} : D_y^{\times} \to \mathbf{O} \simeq H$ vanishes to the order $\check{\lambda}$.

For example, if $G = PGL_2$, then we can identify the dominant integral weights of G with non-negative integers. Then a map $\phi : U \to \mathbb{P}^1$ satisfies the λ -oper condition at $y \in U$ if its differential vanishes to the order λ at y.

Recalling the definition of the space $Op_G(D_x)_{\check{\lambda}}$ given in Section 1.3, we obtain the following result.

Proposition 5.2. There is a bijection between the set $\operatorname{Op}_G(\mathbb{P}^1)_{(z_i),\infty;(\tilde{\lambda}_i),\tilde{\lambda}_{\infty}}$ and the set of equivalence classes, with respect to the action of G on G/B, of the maps $\phi: \mathbb{P}^1 \to G/B$ satisfying the oper condition on $\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$, the $\check{\lambda}_i$ -oper condition at z_i and the $\check{\lambda}_{\infty}$ -oper condition at ∞ .

In particular, the points of $\operatorname{Op}_{PGL_2}(\mathbb{P}^1)_{(z_i),\infty;(\check{\lambda}_i),\check{\lambda}_{\infty}}$ are the same as the equivalence classes of maps $\mathbb{P}^1 \to \mathbb{P}^1$ whose differential does not vanish anywhere on $\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$) and vanishes to order $\check{\lambda}_i$ at $z_i, i = 1, \ldots, N$, and to order $\check{\lambda}_{\infty}$ at ∞ .

5.2. From opers without monodromy to Bethe vectors

Let $\tau = (\mathfrak{F}, \nabla, \mathfrak{F}_{L_B}) \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. By Proposition 5.2, we attach to it a map $\phi_{\tau} : \mathbb{P}^1 \to {}^L G/{}^L B$. A horizontal ${}^L B$ -reduction on τ is completely

determined by the choice of ${}^{L}B$ -reduction at any given point $x \in \mathbb{P}^{1}$, which is a point of $({}^{L}G/{}^{L}B)_{\mathcal{F}_{L_{B,x}}} \simeq {}^{L}G/{}^{L}B$ (see Lemma 3.1). We will choose the point ∞ as our reference point on \mathbb{P}^{1} . Then we obtain

Lemma 5.3. For $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ the set of Miura opers on \mathbb{P}^1 with the underlying oper τ is in bijection with the set of points of $({}^LG/{}^LB)_{\mathcal{F}_{L_{B,\infty}}} \simeq {}^LG/{}^LB$.

Now suppose we are given an oper $\tau \in \operatorname{Op}_{LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. We fix a trivialization of the fiber \mathcal{F}_{∞} of \mathcal{F} at ∞ and consider the corresponding map $\phi_{\tau} : \mathbb{P}^1 \to {}^LG/{}^LB$. Then the value of ϕ_{τ} at $\infty \in \mathbb{P}^1$ is the point ${}^LB \in {}^LG/{}^LB$. Consider the Miura oper structure on τ corresponding to ${}^LB \in {}^LG/{}^LB$, considered as a point in the fiber $({}^LG/{}^LB)_{\mathcal{F}_{L_{B,\infty}}}$. This is the unique Miura oper structure on τ for which the horizontal Borel reduction $\mathcal{F}'_{L_{B,\infty}}$ at the point ∞ coincides with the oper reduction $\mathcal{F}_{L_{B,\infty}}$ (which is the point ${}^LB \in {}^LG/{}^LB$ with respect to our trivialization).

Let us suppose that the oper τ satisfies the following conditions:

- (1) $\phi_{\tau}(z_i)$ is in generic position with ^LB for all $i = 1, \ldots, N$;
- (2) the relative position of $\phi_{\tau}(x)$ and ${}^{L}B$ is either generic or corresponds to a simple reflection $s_i \in W$ for all $x \in \mathbb{A}^1 = \mathbb{P}^1 \setminus \infty$.

Then we will call τ a *non-degenerate* oper.

Let τ be a non-degenerate oper in $\operatorname{Op}_{LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. Consider the unique Miura oper $(\mathcal{F}, \nabla, \mathcal{F}_{LB}, \mathcal{F}'_{LB})$ whose underlying oper is τ and such that $\mathcal{F}_{LB,\infty}$ corresponds to the point ${}^{LB} \in {}^{L}G/{}^{LB} \simeq ({}^{L}G/{}^{LB})_{\mathcal{F}_{LB,\infty}}$. According to the above conditions, the reductions \mathcal{F}_{LB} and \mathcal{F}'_{LB} are in generic position for all but finitely many points of \mathbb{P}^1 , which are distinct from z_1, \ldots, z_N, ∞ . Let us denote these points by w_1, \ldots, w_m . Then at the point w_j the two reductions have relative position s_{i_j} . According to Proposition 3.3, to this Miura oper corresponds a connection on the ${}^{L}H$ -bundle Ω^{ρ} , and hence a connection $\overline{\nabla}$ on the dual bundle $\Omega^{-\rho}$ on $\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, w_1, \ldots, w_m, \infty\}$. Moreover, by Proposition 3.4, the connection $\overline{\nabla}$ extends to a connection with regular singularity on $\Omega^{-\rho}$ on the entire \mathbb{P}^1 , with residues λ_i at $z_i, i = 1, \ldots, N, w_0(\lambda_{\infty} + \rho) - \rho$ at ∞ , and $-\alpha_{i_j}$ at $w_j, j = 1, \ldots, m$. (Note that in Proposition 3.4 we described the residues of connections of the bundle Ω^{ρ} , since now we consider connections on $\Omega^{-\rho}$ which is dual to Ω^{ρ} , when applying Proposition 3.4 we need to change the signs of all residues.)

Therefore, the restriction of $\overline{\nabla}$ to $\mathbb{P}^1 \setminus \infty$ reads

(5.2)
$$\overline{\nabla} = \partial_t + \sum_{i=1}^N \frac{\lambda_i}{t - z_i} - \sum_{j=1}^m \frac{\alpha_{i_j}}{t - w_j}.$$

Using transformation formula (3.2), we find that the residue of $\overline{\nabla}$ at ∞ is equal to

$$-2\rho - \sum_{i=1}^{N} \lambda_i + \sum_{j=1}^{m} \alpha_{i_j}.$$

But by our assumption this residue is equal to $w_0(\lambda_{\infty} + \rho) - \rho$. Therefore we find that

(5.3)
$$-w_0(\lambda_\infty) = \sum_{i=1}^N \lambda_i - \sum_{j=1}^m \alpha_{i_j}$$

In addition, the condition that the oper τ has trivial monodromy representation and Lemma 3.5 imply that if

$$\partial_t - \alpha_{i_j}/(t - w_j) + \mathbf{u}_j(t - w_j), \qquad \mathbf{u}_j(u) \in {}^L \mathfrak{h}[[u]],$$

is the expansion of the connection (5.2) at the point w_j , then $\langle \check{\alpha}_{i_j}, \mathbf{u}_j(0) \rangle = 0$ for all $j = 1, \ldots, m$. When we write them explicitly, we see immediately that these equations are precisely the Bethe Ansatz equations (4.18)!

Thus, we have attached to a non-degenerate oper $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ a connection $\overline{\nabla} = \partial_t + \lambda(t)$ on $\Omega^{-\rho}$ and a solution of the Bethe Ansatz equations. By construction, the Miura transformation of this connection is the oper τ , and so we obtain the following

Proposition 5.4. Suppose that all opers in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ are non-degenerate. Then there is a bijection between the set $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ and the set of solutions of Bethe Ansatz equations (4.18) such that the weight $\sum_{i=1}^N \lambda_i - \sum_{j=1}^m \alpha_{i_j}$ is dominant.

Thus, given a non-degenerate oper $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$, we obtain a solution of Bethe Ansatz equations (4.18). We then follow the procedure of Section 4.4 to assign to the above solution of the Bethe Ansatz equations an eigenvector of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ in $\bigotimes_{i=1}^N M_{\lambda_i}$, which is an \mathfrak{n}_+ -invariant vector $\phi(w_1^{i_1},\ldots,w_m^{i_m})$ of weight $\sum_{i=1}^N \lambda_i - \sum_{j=1}^m \alpha_{i_j}$. Consider its projection onto $\bigotimes_{i=1}^N V_{\lambda_i}$. By formula (5.3), the weight of this vector is

(5.4)
$$\sum_{i=1}^{N} \lambda_i - \sum_{j=1}^{m} \alpha_{i_j} = -w_0(\lambda_\infty),$$

and so it may be viewed as an eigenvector of the Gaudin algebra in the space

$$V_{(\lambda_i),\lambda_{\infty}}^G = \left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)^G.$$

By Theorem 4.11, the eigenvalues of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ on this vector are encoded by the Miura transformation of the connection $\overline{\nabla}$, which is precisely the oper τ with which we have started. Therefore we obtain the following result.

Lemma 5.5. The two opers encoding the eigenvalues of the Gaudin algebra on the Bethe vectors corresponding to two different solutions of the Bethe Ansatz equations are necessarily different.

Suppose that all opers in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ are *non-degenerate*¹ and that each of the Bethe vectors is *non-zero*. Then we obtain a map

$$\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} \to \operatorname{Spec}\overline{\mathcal{Z}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}(\mathfrak{g})$$

that is inverse to the map (5.1) discussed at the beginning of this section.

Let us summarize our results.

Proposition 5.6. Assume that all ^LG-opers in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ are non-degenerate and that all Bethe vectors obtained from solutions of the Bethe Ansatz equations (4.18) satisfying the condition (5.4) are non-zero. Then there is a bijection between the spectrum of the generalized Gaudin Hamiltonians on $V_{(\lambda_i),\lambda_{\infty}}^G$ (not counting multiplicities) and the set $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ of ^LG-opers on \mathbb{P}^1 with regular singularities at z_1,\ldots,z_N,∞ and residues $\varpi(\lambda_1 + \rho),\ldots,\varpi(\lambda_N + \rho)$ and $\varpi(\lambda_{\infty} + \rho)$, respectively, which have trivial monodromy.

Moreover, if in addition the Gaudin Hamiltonians are diagonalizable and have simple spectrum on $V_{(\lambda_i),\lambda_{\infty}}^G$, then the Bethe vectors constitute an eigenbasis of $V_{(\lambda_i),\lambda_{\infty}}^G$.

The last statement of Proposition 5.6 that the Bethe vectors constitute an eigenbasis of $V^G_{(\lambda_i),\lambda_{\infty}}$ is referred to as the completeness of the Bethe Ansatz.

We note that the completeness of the Bethe Ansatz has been previously proved for $\mathfrak{g} = \mathfrak{sl}_2$ and generic values of z_1, \ldots, z_N by Scherbak and Varchenko [SV].

We will discuss the degenerate opers and the corresponding eigenvectors of the Gaudin hamiltonians in Section 5.5.

5.3. The case of \mathfrak{sl}_2

In the case when $\mathfrak{g} = \mathfrak{sl}_2$, we identify the weights with complex numbers. The Bethe vector has a simple form in this case. We use the standard basis $\{e, h, f\}$ of \mathfrak{sl}_2 . Then

(5.5)
$$\phi(w_1,\ldots,w_m) = f(w_1)\ldots f(w_m)v_{\lambda_1}\otimes\ldots v_{\lambda_N},$$

where

$$f(w) = \sum_{i=1}^{N} \frac{f^{(i)}}{w - z_i}$$

The Bethe Ansatz equations read

$$\sum_{i=1}^{N} \frac{\lambda_i}{w_j - z_i} = \sum_{s \neq j} \frac{2}{w_j - w_s}$$

48

¹It follows from the results of Mukhin and Varchenko in [MV2] that for some $\lambda_1, \ldots, \lambda_N, \lambda_\infty$ it is possible that there exist degenerate opers in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_\infty}$ even for generic values of z_1, \ldots, z_N .

If these equations are satisfied, then this vector is a highest weight vector of weight λ_{∞} (with respect to the diagonal action), and it is an eigenvector in $\bigotimes_{i=1}^{N} M_{\lambda_i}$ of the Gaudin algebra

$$\mathcal{Z}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} = \mathbb{C}[\Xi_i]_{i=1,\dots,N} / \left(\sum_{i=1}^N \Xi_i\right),$$

where Ξ_i is the *i*th Gaudin Hamiltonian given by formula (0.1).

Its eigenvalue is represented by the PGL_2 -oper

$$\partial_t^2 - \sum_{i=1}^N \frac{\lambda_i (\lambda_i + 2)/4}{(t - z_i)^2} - \sum_{i=1}^N \frac{c_i}{t - z_i} = (\partial_t - \lambda(t))(\partial_t + \lambda(t)),$$

where

$$\lambda(t) = \sum_{i=1}^{N} \frac{\lambda_i/2}{t - z_i} - \sum_{j=1}^{m} \frac{1}{t - w_j}.$$

In other words, the eigenvalue c_i of Ξ_i on $\phi(w_1, \ldots, w_m)$ is equal to

(5.6)
$$c_i = \lambda_i \left(\sum_{j \neq i} \frac{\lambda_j}{z_i - z_j} - \sum_{j=1}^m \frac{1}{z_i - w_j} \right).$$

For $\mathfrak{g} = \mathfrak{sl}_2$, all opers in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_\infty}$ are non-degenerate for generic values of z_1, \ldots, z_N (for example, this follows from the results of [SV]).

5.4. Solutions of the Bethe Ansatz equations and flag manifolds

In Section 5.2 we considered only those solutions of the Bethe Ansatz equation for which

(5.7)
$$\sum_{i=1}^{N} \lambda_i - \sum_{j=1}^{m} \alpha_{i_j}$$

is a dominant integral weight. The reason for that was that we wanted to construct eigenvectors of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ in $\bigotimes_{i=1}^N V_{\lambda_i}$ of weight (5.7). Since the Bethe vectors are automatically \mathfrak{n}_+ -invariant, we find that such a vector can be non-zero in $\bigotimes_{i=1}^N V_{\lambda_i}$ only if this weight is dominant integral. In this case we may write it in the form $-w_0(\lambda_\infty)$, where λ_∞ is another dominant integral weight. Then the corresponding Bethe vector gives us an eigenvector of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ in the space

$$V_{(\lambda_i),\lambda_{\infty}}^G = \left(\bigotimes_{i=1}^N V_{\lambda_i} \otimes V_{\lambda_{\infty}}\right)^G.$$

We have seen in Section 5.2 that these solutions of the Bethe Ansatz equations are in one-to-one correspondence with the Miura opers on \mathbb{P}^1 such that the corresponding oper in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is non-degenerate, and such that the horizontal LB -reduction \mathcal{F}'_{L_B} coincides with the oper LB -reduction \mathcal{F}_{L_B} at ∞ .

It is natural to consider more general solutions of the Bethe Ansatz equations, i.e., those for which the weight (5.7) is not necessarily dominant integral. The first step is to relate these solutions to Miura opers.

Suppose that we are given an arbitrary solution of the Bethe Ansatz equations (4.18). By Proposition 4.10, we attach to it the connection $\overline{\nabla}$ on the ^{*L*}*H*-bundle $\Omega^{-\rho}$ on \mathbb{P}^1 whose restriction to $\mathbb{P}^1 \setminus \infty$ reads

(5.8)
$$\overline{\nabla} = \partial_t + \sum_{i=1}^N \frac{\lambda_i}{t - z_i} - \sum_{j=1}^m \frac{\alpha_{i_j}}{t - w_j}.$$

Recall the Miura transformation $\overline{\mathbf{b}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^*$ introduced in formula (4.21). Applying this map to $\overline{\nabla}$, we obtain a LG -oper τ . It has regular singularities at z_1, \ldots, z_N and ∞ , and is regular elsewhere on \mathbb{P}^1 (recall that the Bethe Ansatz equations ensure that the oper τ is regular at the points w_1, \ldots, w_m). Around the point z_i the oper connection reads

$$\nabla = \partial_t + p_{-1} + \frac{\lambda_i}{t - z_i} + \operatorname{reg}.$$

and by applying the gauge transformation with $\lambda(t)$ we can bring it to the form (1.9). Therefore, according to Lemma 1.2, τ has no monodromy around $z_i, i = 1, \ldots, N$.

Let us consider the point ∞ . Since τ has no monodromy anywhere on $\mathbb{P}^1 \setminus \infty$, it cannot have monodromy at ∞ either. By Lemma 1.2, the restriction of τ to the disc around ∞ belongs to $\operatorname{Op}_G(D_\infty)_{\lambda_\infty}$ for some integral dominant weight λ_∞ . But then the commutative diagram (3.3) implies that the residue of $\overline{\nabla}$ at ∞ is equal to $w'(\lambda_\infty + \rho) - \rho$ for some $w' \in W$. Writing $w' = ww_0$, we find that the residue is equal to $ww_0(\lambda_\infty + \rho) - \rho$. On the other hand, using transformation formula (3.2), we find that the residue of $\overline{\nabla}$ at ∞ is equal to

$$-2\rho - \sum_{i=1}^{N} \lambda_i + \sum_{j=1}^{m} \alpha_{i_j}.$$

Therefore we must have the equality

(5.9)
$$\sum_{i=1}^{N} \lambda_i - \sum_{j=1}^{m} \alpha_{i_j} = w(-w_0(\lambda_{\infty}) + \rho) - \rho$$

for some dominant integral weight λ_{∞} and $w \in W$. Thus, the oper τ belongs to $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$, and so $\overline{\nabla}$ is an element of $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$.

Recall from Section 4.2 that for any curve U we have maps

 $\mathbf{b}_U^* : \operatorname{Conn}(\Omega^{-\rho})_U \to \operatorname{MOp}_{L_G}(U)_{\operatorname{gen}}, \qquad \overline{\mathbf{b}}_U^* : \operatorname{Conn}(\Omega^{-\rho})_U \to \operatorname{Op}_{L_G}(U).$

Let $U = \mathbb{P}^1 \setminus \{z_1, \ldots, z_N, w_1, \ldots, w_m, \infty\}$. Then for any weights $\lambda_1, \ldots, \lambda_N, \lambda_\infty$ we have a subset $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_\infty}^{\operatorname{gen}}$ of $\operatorname{Conn}(\Omega^{-\rho})_U$ which is mapped by the Miura transformation $\overline{\mathbf{b}}_{(z_i),\infty;(\lambda_i),\lambda_\infty}^{\mathrm{R}}$ into $\operatorname{Op}_{L_G}^{\mathrm{RS}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_\infty}$ (see formula (4.20)). But as we saw above, if the weights $\lambda_1, \ldots, \lambda_N, \lambda_\infty$ are dominant integral, then the image is actually contained in even smaller set $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ of those opers which have trivial monodromy. Therefore we have a map

$$\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}} \to \operatorname{Op}_{{}^{L}G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$$

which we also denote by $\overline{\mathbf{b}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^*$. The restriction of the corresponding map \mathbf{b}_U^* to $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_\infty}^{\operatorname{gen}}$ gives us a map

$$\mathbf{b}^*_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}: \operatorname{Conn}(\Omega^{-\rho})^{\operatorname{gen}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} \to \operatorname{MOp}_{{}^LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$$

where $\mathrm{MOp}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is the space of Miura opers on \mathbb{P}^1 whose underlying opers belong to $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. We will show that the image of this map is open and dense.

Let

$$\mathrm{MOp}_{L_G}(\mathbb{P}^1)^{\mathrm{gen}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} \subset \mathrm{MOp}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$$

be the subvariety of those Miura opers whose horizontal ^{L}B -reduction satisfies the conditions (1) and (2) of Section 5.2. We call such Miura opers non-degenerate.

Recall from Lemma 5.3 that the space of all Miura opers on \mathbb{P}^1 corresponding to a fixed ${}^{L}G$ -oper $\tau \in \operatorname{Op}_{{}^{L}G}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}}$ is isomorphic to the flag variety ${}^{L}G/{}^{L}B$. Non-degenerate Miura opers corresponding to τ form a subvariety of ${}^{L}G/{}^{L}B$ which we denote by $({}^{L}G/{}^{L}B)_{\tau}$. It is clear that $({}^{L}G/{}^{L}B)_{\tau}$ is an open and dense subvariety of ${}^{L}G/{}^{L}B$. Indeed, $({}^{L}G/{}^{L}B)_{\tau}$ is contained in the intersection U_{τ} of finitely many open and dense subsets, namely, the sets of points of ${}^{L}G/{}^{L}B$ which are in generic relative position with $\phi_{\tau}(z_i)$ (each is isomorphic to the big Schubert cell). The complement of $({}^{L}G/{}^{L}B)_{\tau}$ in U_{τ} is a subvariety of codimension one. This subvariety consists of all points in ${}^{L}G/{}^{L}B$ which are in relative position w with $\phi_{\tau}(x), x \in \mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$, where w runs over the subset of W of all elements of length $l(w) \geq 2$. The subvariety of these points for fixed x has codimension two, and therefore their union, as x moves along the curve $\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$, has codimension (at least) one.

It follows from our construction that the image of the map $\mathbf{b}^*_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is contained in $\operatorname{MOp}_{L_G}^{\operatorname{gen}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. Conversely, suppose we are given a Miura oper in $\operatorname{MOp}_{L_G}^{\operatorname{gen}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. Then we obtain a connection on $w_0(\Omega^{\rho}) = \mathcal{F}'_{L_B}/^L N$, and hence a connection on $\Omega^{-\rho}$. It follows from Lemma 3.5 that this connection belongs to $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$. Moreover, in the same way as in the proof of Proposition 3.3 we find that we obtain a map

$$\mathrm{MOp}_{L_G}^{\mathrm{gen}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}} \to \mathrm{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\alpha}}^{\mathrm{gen}}$$

that is inverse to $\mathbf{b}^*_{(z_i),\infty;(\lambda_i),\lambda_\infty}$. Therefore we obtain the following result.

Lemma 5.7. The map $\mathbf{b}^*_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is a bijection between $\operatorname{Conn}(\Omega^{-\rho})^{\operatorname{gen}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ and $\operatorname{MOp}_{L_G}^{\operatorname{gen}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$.

Combining Lemma 5.7 and Proposition 5.2, we obtain

Theorem 5.8. The set of solutions of the Bethe Ansatz equations (4.18) is in bijection with the set of points of $\operatorname{MOp}_{L_G}(\mathbb{P}^1)^{\operatorname{gen}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$, where λ_{∞} satisfies (5.9) for some $w \in W$.

By Proposition 3.4, the points of $\operatorname{MOp}_{LG}(\mathbb{P}^1)^{\operatorname{gen}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ which satisfy formula (5.9) with a fixed $w \in W$ are those Miura opers for which the value of ϕ_{τ} at ∞ belongs to the Schubert cell

$$S_{ww_0} = {}^L B w_0 w w_0 {}^L B \subset {}^L G / {}^L B.$$

Therefore such Miura opers correspond to the points of intersection of $({}^{L}G/{}^{L}B)_{\tau}$ and $S_{ww_{0}}$.

Note that except for the big cell S_1 , the intersection between the Schubert cell S_{ww_0} and the open dense subset $({}^LG/{}^LB)_{\tau} \subset {}^LG/{}^LB$ could be either an open dense subset of S_{ww_0} or empty.² We make the statement of Theorem 5.8 more precise as follows:

Theorem 5.9. The set of solutions of the Bethe Ansatz equations (4.18) decomposes into a union of disjoint subsets labeled by $\operatorname{Op}_{LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. The set of points corresponding to LG -oper $\tau \in \operatorname{Op}_{LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is in bijection with the set of points of an open and dense subset $({}^LG/{}^LB)_{\tau}$ of the flag variety ${}^LG/{}^LB$.

Further, each of these solution must satisfy the equation (5.9) for some $w \in W$, and the solutions which satisfy this equation with a fixed $w \in W$ are in bijection with an open subset $S_{ww_0} \cap ({}^LG/{}^LB)_{\tau}$ of the Schubert cell $S_{ww_0} = {}^LBw_0ww_0{}^LB$.

Remark 5.10. In [MV1] (resp., [BM]), it was shown, by a method different from ours, that in the case when \mathfrak{g} is of types A_n, B_n or C_n (resp., G_2) and z_1, \ldots, z_N are generic, the set of solutions of the Bethe Ansatz equations satisfying (5.9) is in bijection with a disjoint union of open and dense subsets of S_w . The connection between the results of [MV1] and our results is explained in [F3].

In [F3] we obtained a generalization of Theorem 5.9 to the case when \mathfrak{g} is an arbitrary Kac-Moody algebra.

5.5. Degenerate opers

By Lemma 5.7 we have a bijection between the set $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^{\operatorname{gen}}$ and an open subset $\operatorname{MOp}_{L_G}^{\operatorname{gen}}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ of non-degenerate Miura opers in

$$\operatorname{MOp}_{{}^{L}G}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}}$$

Now we wish to extend this bijection to the entire set $\mathrm{MOp}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. To any point of $\mathrm{MOp}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ we can still assign, as before, a connection $\overline{\nabla}$ on $\Omega^{-\rho}$ with regular singularities at z_1, \ldots, z_N, ∞ and some additional points

²For example, it follows from the results of Mukhin and Varchenko in [MV2] that sometimes this open set may not contain the one point Schubert cell $S_{w_0} \subset G/B$ even if we allow z_1, \ldots, z_N to be generic.

 w_1, \ldots, w_m . By Proposition 3.4, the residue of $\overline{\nabla}$ at z_i (resp., ∞, w_j) must be equal to $y_i(\check{\lambda}_i + \check{\rho}) - \check{\rho}$ (resp., $y_{\infty}(\check{\lambda}_{\infty} + \check{\rho}) - \check{\rho}, y'_j(\check{\rho}) - \check{\rho}$) for some elements $y_i, y_{\infty}, y'_j \in W$. Hence this connection has the form

(5.10)
$$\partial_t + \sum_{i=1}^N \frac{y_i(\lambda_i + \rho) - \rho}{t - z_i} + \sum_{j=1}^m \frac{y'_j(\rho) - \rho}{t - w_j}$$

Considering the expansion of this connection at ∞ , we obtain the following relation

(5.11)
$$\sum_{i=1}^{N} (y_i(\check{\lambda}_i + \check{\rho}) - \check{\rho}) + \sum_{j=1}^{m} (y'_j(\check{\rho}) - \check{\rho}) = y_\infty w_0(-w_0(\check{\lambda}_\infty) + \check{\rho}) - \check{\rho}$$

Let $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ be the set of all connections of the form (5.10) whose Miura transformation belongs to $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. Using Lemma 5.3, we then obtain the following generalization of Lemma 5.7 (see [F3]):

Theorem 5.11. There is a bijection between the sets $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ and $\operatorname{MOp}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. In particular, the set of those connections in $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ which correspond to a fixed LG -oper $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\check{\lambda}_i),\check{\lambda}_{\infty}}$ is isomorphic to the set of points of the flag variety ${}^LG/{}^LB$.

Moreover, the residues of these connections must satisfy the relation (5.11) for some $y_{\infty} \in W$. The set of those connections which satisfy this relation is in bijection with the Schubert cell ${}^{L}Bw_{0}y_{\infty}{}^{L}B$ in ${}^{L}G/{}^{L}B$.

Consider, in particular, the unique Miura oper corresponding to an oper

$$\tau \in \operatorname{Op}_{{}^{L}G}(\mathbb{P}^{1})_{(z_{i}),\infty;(\lambda_{i}),\lambda_{\infty}}$$

in which the two Borel reductions coincide at ∞ (it corresponds to the one point Schubert cell ^LB in ^LG/^LB). Let $\overline{\nabla}_{\tau}$ be the corresponding connection in $\operatorname{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. It has the form (5.10) and its residues satisfy the relation (5.11) with $y_{\infty} = w_0$:

$$\sum_{i=1}^{N} (y_i(\lambda_i + \rho) - \rho) + \sum_{j=1}^{m} (y'_j(\rho) - \rho) = -w_0(\lambda_\infty).$$

The oper $\tau \in \operatorname{Op}_{LG}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ is degenerate if and only if either some of the elements y_i are not equal to 1 or some of the elements y'_j have lengths greater than 1 (i.e., are not simple reflections).

Observe that we can write $y_i(\lambda_i + \rho) - \rho$ as λ_i minus a combination of simple roots with non-negative coefficients, and likewise, $y'_j(\rho) - \rho$ is equal to minus a linear combination of the simple roots with non-negative coefficients. It is instructive to think of the connection $\overline{\nabla}_{\tau}$ corresponding to degenerate opers as the limit of a family of connections corresponding to non-degenerate opers, under which some of the points w_j 's either collide with some of the z_i 's or with each other (leading to the degeneracies of types (1) and (2), respectively). The resulting residues of the connection become the sums of the residues of the points that have coalesced

together. Proposition 3.4 places severe restrictions on what kinds of residues (and hence collisions) can occur: namely, at the point z_i the residue must lie in the W-orbit of λ_i , whereas at the additional points w_j the residues must be in the W-orbit of 0 (under the ρ -shifted action of W).

We expect that if z_1, \ldots, z_N are generic, then for any $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ we have $y_i = 1$ for all $i = 1, \ldots, N$ in the connection $\overline{\nabla}_{\tau}$. In other words, this Miura oper satisfies condition (1) from Section 5.2, but may not satisfy condition (2), that is at least one of the y'_i 's is not a simple reflection.

Suppose that τ is such an oper, so in particular we have

(5.12)
$$\sum_{i=1}^{N} \lambda_i + \sum_{j=1}^{m} (y'_j(\rho) - \rho) = -w_0(\lambda_\infty).$$

Then we can still attach to the connection (5.10) an eigenvector of the generalized Gaudin hamiltonians in $V^G_{(\lambda_i),\lambda_{\infty}}$ with eigenvalue τ by generalizing the procedure of Section 4.4.³

More precisely, suppose that the residue of $\lambda(t)$ at w_j is equal to $y'_j(\check{\rho}) - \check{\rho}$ for some $y'_j \in W$. Then the expansion of this connection at w_j is $\mu_j(t-w_j)$, where

$$\mu_j(t) = \frac{y'_j(\check{\rho}) - \check{\rho}}{t} + \sum_{n \ge 0} \mu_{j,n} t^n, \qquad \mu_{j,n} \in \mathfrak{h}^*.$$

The condition that the oper τ has no monodromy at w_j (i.e., that the Miura transformation of the connection $\partial_t + \mu_j(t)$ is regular at t = 0) translates into a system of equations on the coefficients $\mu_{j,n}$ of this expansion. We denote this system by S_j . Recall from Lemma 3.5 that when $y'_j = s_{i_j}$ there is only one equation $\langle \check{\alpha}_{i_j}, \mu_{j,0} \rangle = 0$.

To see what these equations look like, we consider the example when $\mathfrak{g} = \mathfrak{sl}_n$. The PGL_n -opers may be represented by *n*th order differential operators of the form

(5.13)
$$\partial_t^n + v_1(t)\partial_t^{n-2} + \ldots + v_{n-1}(t).$$

Let us identify the dual Cartan subalgebra of \mathfrak{sl}_n with the hyperplane $\sum_{k=1}^n \epsilon_k = 0$ of the vector space span $\{\epsilon_k\}_{k=1,\dots,n}$. Then an ^L \mathfrak{h} -valued connection has the form $\partial_t + \sum_{k=1}^n u_k(t)\epsilon_k$. The Miura transformation of this connection is given by the formula

(5.14)
$$\partial_t^n + v_1(t)\partial_t^{n-2} + \ldots + v_{n-1}(t) = (\partial_t - u_1(t))\ldots(\partial_t - u_n(t)).$$

The system of equations S_j is obtained by writing $\mu_j(t)$ as $\sum_{k=1}^n u_k(t)\epsilon_k$, substituting the functions $u_k(t)$ into formula (5.14) and setting to zero all coefficients in

³the construction outlined below was found by B. Feigin and myself around the time [FFR] was written; it was explained in [F1], Sect. 5.6–5.8 in the case when $\mathfrak{g} = \mathfrak{sl}_3$

front of the negative powers of t in the resulting nth order differential operator. For example, if $y'_i = s_a s_b$, |a - b| > 1, then we have

$$u_m(t) = -(\delta_{m,a} - \delta_{m,a+1} + \delta_{m,b} - \delta_{m,b+1})t^{-1} + \sum_{r \ge 0} u_{m,r}t^r$$

and it is easy to write down the equations on the coefficients $u_{m,r}$ corresponding to the regularity of the operator (5.14) at t = 0.

The system of equations S_j also has a nice interpretation in the theory of the Wakimoto modules. Namely, consider the Wakimoto module $W_{\mu_j(z)}$. Then it contains a $\mathfrak{g}[[t]]$ -invariant vector P_j if and only if the system S_j of equations on $\mu_j(z)$ is satisfied. In the case when $y'_j = s_{ij}$ and $\mu_j(z) = -\alpha_{ij}z^{-1} + \ldots$, this is the statement of Lemma 4.5; this vector is given by the formula $e^R_{ij,-1}|0\rangle$ in this case. In general the formulas for these vectors are more complicated. They can probably be obtained as the limits of the Bethe vectors corresponding to small deformations of the highest weights λ_i .

It follows from the definition that for our connection $\overline{\nabla}_{\tau}$ the systems S_j are satisfied for all $j = 1, \ldots, m$. Therefore each module $W_{\mu_j(z)}$ contains a $\mathfrak{g}[[t]]$ invariant vector P_j . Then, in the same way as in Section 4.4, we obtain a homomorphism of $\hat{\mathfrak{g}}_{\kappa_c}$ -modules

$$\bigotimes_{i=1}^{N} W_{\lambda_{i}(z)} \otimes \mathbb{V}_{0}^{\otimes m} \otimes W'_{\lambda_{\infty}(z)} \to \bigotimes_{i=1}^{N} W_{\lambda_{i}(z)} \otimes \bigotimes_{j=1}^{m} W_{\mu_{j}(z)} \otimes W'_{\lambda_{\infty}(z)},$$

which sends the vacuum vector in the *j*th copy of \mathbb{V}_0 to $P_j \in W_{\mu_j(z)}$. Therefore, composing the corresponding map of the spaces of coinvariants with the functional $\tau_{(z_i),(w_j)}$ introduced at the end of Section 4.3, we obtain a linear functional

$$\widetilde{\tau}_{(z_i),(w_j)}: H((W_{\lambda_i(z)}), W'_{\lambda_\infty(z)}) \to \mathbb{C}$$

By construction, $\tilde{\tau}_{(z_i),(w_j)}$ is an eigenvector of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$, and the corresponding eigenvalue is given by the ^LG-oper $\overline{\mathbf{b}}_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}^*(\overline{\nabla}_{\tau}) = \tau$.

Continuing along the lines of the construction presented in Section 4.4, we obtain an \mathfrak{n}_+ -invariant vector in $\bigotimes_{i=1}^N M_{\lambda_i}$ of weight $\sum_{i=1}^N \lambda_i + \sum_{j=1}^m (y'_j(\rho) - \rho)$, which, according to formula (5.12), is equal to $-w_0(\lambda_\infty)$ which is a dominant integral weight. The projection of this vector onto $\bigotimes_{i=1}^N V_{\lambda_i}$ gives rise to an eigenvector of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ in $V^G_{(\lambda_i),\lambda_\infty}$ with the eigenvalue corresponding to our degenerate oper τ .

Thus, we have assigned to any oper τ in $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ such that the corresponding special connection $\overline{\nabla}_{\tau}$ satisfies condition (1) that $y_i = 1$, for all $i = 1, \ldots, N$, in (5.10), an eigenvector of the Gaudin algebra in $V_{(\lambda_i),\lambda_{\infty}}^G$ with the eigenvalue τ .

The next question is what happens if $\overline{\nabla}_{\tau}$ has $y_i \neq 1$ for some *i*. In this case the above construction gives rise to an eigenvector of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ in the space

$$\left(\otimes_{i=1}^{N} M_{y_i(\lambda_i+\rho)-\rho}\right)_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$$

of the diagonal \mathfrak{n}_+ -invariants of $\bigotimes_{i=1}^N M_{y_i(\lambda_i+\rho)-\rho}$ of weight $-w_0(\lambda_\infty)$.

Now recall that for any dominant integral weight λ and $y \in W$ the Verma module $M_{y(\lambda+\rho)-\rho}$ is a \mathfrak{g} -submodule of M_{λ} . Hence $\bigotimes_{i=1}^{N} M_{y_i(\lambda_i+\rho)-\rho}$ is naturally a subspace of $\bigotimes_{i=1}^{N} M_{\lambda_i}$. It is also a $\mathfrak{g}_{\text{diag}}$ -submodule of $\bigotimes_{i=1}^{N} M_{\lambda_i}$ which is contained in the maximal proper $\mathfrak{g}_{\text{diag}}$ -submodule $I_{(\lambda_i)}$. The quotient of $\bigotimes_{i=1}^{N} M_{\lambda_i}$ by $I_{(\lambda_i)}$ is isomorphic to $\bigotimes_{i=1}^{N} M_{\lambda_i}$ with the eigenvalue τ , but this eigenvector of the Gaudin algebra in $\bigotimes_{i=1}^{N} M_{\lambda_i}$ with the eigenvalue τ , but this eigenvector is inside its maximal proper $\mathfrak{g}_{\text{diag}}$ -submodule $I_{(\lambda_i)}$. Some sample computations that we have made in the case when $\mathfrak{g} = \mathfrak{sl}_2$ suggest that in this case the generalized eigenspace of the Gaudin algebra in $(\bigotimes_{i=1}^{N} M_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}+}$ corresponding to the eigenvalue τ has dimension greater than one and that its projection onto $(\bigotimes_{i=1}^{N} V_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}+} \simeq V_{(\lambda_i),\lambda_{\infty}}^G$ is non-zero. In other words, we expect that in this case there also exists an eigenvector of the Gaudin algebra $\mathfrak{Z}_{(z_i)}(\mathfrak{g})$ in $V_{(\lambda_i),\lambda_{\infty}}^G$ with the same eigenvalue τ .

If true, this would explain the meaning of the degeneracy of the connection $\overline{\nabla}_{\tau}$ at the point z_i . Namely, for some special values of z_i it may happen that some of the eigenvalues of the Gaudin operators in $(\bigotimes_{i=1}^N V_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$ and in $(I_{(\lambda_i)})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$ become equal. Then they may combine into a Jordan block such that the eigenvector in $(\bigotimes_{i=1}^N V_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$ is only a generalized eigenvector in $(\bigotimes_{i=1}^N M_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$. If that happens, we can no longer obtain this eigenvector by projection from $(\bigotimes_{i=1}^N M_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$ (but we could potentially obtain it by considering the family of eigenvectors corresponding to small perturbations of the z_i 's).

Let $\otimes_{i=1}^{N} M_{y_i(\lambda_i+\rho)-\rho}$ be the smallest $\mathfrak{g}^{\otimes N}$ -submodule of $\otimes_{i=1}^{N} M_{\lambda_i}$ in which the corresponding "true" eigenvector in $(\otimes_{i=1}^{N} M_{\lambda_i})_{-w_0(\lambda_{\infty})}^{\mathfrak{n}_+}$ is contained. We believe that this is precisely the situation when the connection $\overline{\nabla}_{\tau}$ develops a singularity with residue $y_i(\lambda_i+\rho)-\rho$ at the point z_i .

Let us summarize the emerging picture. According to Proposition 5.1, we have an injective map from the spectrum of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ occurring in $V_{(\lambda_i),\lambda_{\infty}}^G$ to $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. We wish to construct the inverse map, in other words, to assign to each oper $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ an eigenvector of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ with eigenvalue τ in $V_{(\lambda_i),\lambda_{\infty}}^G$. (Here we will not discuss the question as to whether $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ acts diagonally on $V_{(\lambda_i),\lambda_{\infty}}^G$ and whether there are multiplicities. It is known that Jordan blocks occur even in the case when $\mathfrak{g} = \mathfrak{sl}_2$, though it is expected that for generic z_i 's the action of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ is diagonalizable and has simple spectrum. Here we will only discuss the spectrum as a set without multiplicities.)

First, we associate to τ the Miura oper in which the horizontal Borel reduction coincides with the oper reduction at the point ∞ . This Miura oper gives rise to a connection $\overline{\nabla}_{\tau} \in \text{Conn}(\Omega^{-\rho})_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. It has the form (5.10), and the residues satisfy the condition (5.12). The simplest situation occurs if all elements y_i in (5.10) are equal to 1 and all elements y'_j are simple reflections. This is the situation where Bethe Ansatz is applicable. Namely, as explained in Section 4.4, we assign to $\overline{\nabla}_{\tau}$ a certain coinvariant of the tensor product of the Wakimoto modules, which gives rise to an eigenvector of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ with eigenvalue τ in $V^G_{(\lambda_i),\lambda_{\infty}}$. This is the Bethe vector given by an explicit formula (4.22). It was believed that this was in fact a generic situation, i.e., for fixed $\lambda_1, \ldots, \lambda_N, \lambda_{\infty}$ and generic z_1, \ldots, z_N the connection $\overline{\nabla}_{\tau}$ satisfies these conditions for all opers $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$. But recent results of Mukhin and Varchenko [MV2] show that this is not the case. Hence we analyze the possible degeneracies.

The first type of degeneracy that may occur is the following: while all elements y_i in (5.10) are still equal to 1, some of the elements y'_j are no longer simple reflections. We expect this to be the generic situation. In this case we can still construct an eigenvector of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ with eigenvalue τ in $V^G_{(\lambda_i),\lambda_{\infty}}$, as explained above. A formula for this vector will be more complicated than the formula for a Bethe vector, but in principal such a vector can be obtained by an algorithmic procedure.

The most general and most difficult case is when some of the elements y_i are not equal to 1 and some of the elements y'_j are not simple reflections. In this case, as explained above, we still believe that there exists an eigenvector of $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ with eigenvalue τ in $V^G_{(\lambda_i),\lambda_{\infty}}$. But this eigenvector cannot be constructed directly, by using the Wakimoto modules, as above. Indeed, our construction using the Wakimoto modules gives eigenvectors in $\otimes_{i=1}^N M_{\lambda_i}$, but we expect that in this most degenerate case the eigenvector in $V^G_{(\lambda_i),\lambda_{\infty}}$ cannot be lifted to an eigenvector in $\otimes_{i=1}^N M_{\lambda_i}$ (only to a generalized eigenvector). However, one can probably obtain this eigenvector by considering families of eigenvectors corresponding to small perturbations of the z_i 's or λ_i 's.

Finally, there is a question as to whether all of the eigenvectors constructed by means of the Wakimoto module construction (including the Bethe vectors) are non-zero. It was conjectured by S. Chmutov and I. Scherbak in [CS] that the Bethe vectors are always non-zero for $\mathfrak{g} = \mathfrak{sl}_n$.

Now we propose the following conjecture:

Conjecture 1. For any set of integral dominant weights $\lambda_1, \ldots, \lambda_N, \lambda_\infty$ and an arbitrary collection of distinct complex numbers z_1, \ldots, z_N there is a bijection between the set $\operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_\infty}$ and the spectrum of the Gaudin algebra $\mathcal{Z}_{(z_i)}(\mathfrak{g})$ in $V^G_{(\lambda_i),\lambda_\infty}$ (not counting multiplicities).

We do not know whether one can prove this conjecture by following the strategy outlined above. Rather, one should view the above discussion only as an explanation of the link between this conjecture and the traditional Bethe Ansatz conjectures. What we have in mind is a completely different approach to proving this conjecture that is suggested by the geometric Langlands correspondence. We conclude this section by briefly outlining this approach.

As explained in [F1] (following the work [BD1] of A. Beilinson and V. Drinfeld), to each ${}^{L}G$ -oper $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$ one assigns a D-module $\widetilde{\mathcal{F}}_{\tau}$ on the moduli stack $\operatorname{Bun}_{G}(\mathbb{P}^1)_{(z_i),\infty}$ of G-bundles on \mathbb{P}^1 with B-reductions at the points z_1, \ldots, z_N, ∞ . This D-module is a Hecke eigensheaf whose "eigenvalue" is the flat ${}^{L}G$ -bundle on $\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$ obtained by forgetting the ${}^{L}B$ -reduction of τ . This flat ${}^{L}G$ -bundle has trivial monodromy and hence is isomorphic to the restriction to $\mathbb{P}^1 \setminus \{z_1, \ldots, z_N, \infty\}$ of the trivial flat ${}^{L}G$ -bundle on \mathbb{P}^1 .

On the other hand, one can show that the D-module that we construct on $\operatorname{Bun}_G(\mathbb{P}^1)_{(z_i),\infty}$ "does not depend" on the B-reductions at the points z_i and ∞ , i.e., it is a pull-back from a D-module \mathcal{F}_{τ} on the moduli stack $\operatorname{Bun}_G(\mathbb{P}^1)$ of G-bundles on \mathbb{P}^1 . This D-module is a Hecke eigensheaf whose eigenvalue is the trivial LG -bundle with the trivial connection.

The moduli stack $\operatorname{Bun}_G(\mathbb{P}^1)$ has an open part $\operatorname{Bun}_G^{\circ}(\mathbb{P}^1)$ corresponding to the trivial bundle, whose preimage in $\operatorname{Bun}_G(\mathbb{P}^1)_{(z_i),\infty}$ is isomorphic to $(G/B)^{N+1}/G_{\operatorname{diag}}$.

We expect that the following two properties hold: (a) the Hecke eigensheaf \mathcal{F}_{τ} is non-zero for any $\tau \in \operatorname{Op}_{L_G}(\mathbb{P}^1)_{(z_i),\infty;(\lambda_i),\lambda_{\infty}}$, and (b) the restriction of any non-zero Hecke eigensheaf on $\operatorname{Bun}_G(\mathbb{P}^1)$ to $\operatorname{Bun}_G^{\circ}(\mathbb{P}^1)$ is necessarily also non-zero.

Suppose that these properties hold (in the case of \mathfrak{sl}_2 property (a) can be proved using the Sklyanin separation of variables discussed in [F1]). Then we obtain that the restriction of the D-module $\widetilde{\mathcal{F}}_{\tau}$ on $\operatorname{Bun}_G(\mathbb{P}^1)_{(z_i),\infty}$ to $(G/B)^{N+1}/G_{\operatorname{diag}}$ is a pull-back of a non-zero D-module on $\operatorname{Bun}^o_G(\mathbb{P}^1)$, and hence is isomorphic to a direct sum of copies of its structure sheaf, considered as a D-module. This, in turn, is equivalent to the existence of a non-zero eigenvector of the Gaudin algebra with eigenvalue τ in $V^G_{(\lambda_i),\lambda_{\infty}}$, as explained in [F1]. Hence we obtain a proof of Conjecture 1. We hope to return to this question in a future publication.

Acknowledgements

This paper reviews the results of our previous works [FFR, F1, F3], some of which were obtained jointly with B. Feigin and N. Reshetikhin. I wish to thank I. Scherbak for useful discussions.

I thank the organizers of the Workshop "Infinite-dimensional algebras and quantum integrable systems" in Faro in July of 2003 for their invitation to give a talk on this subject and for encouraging me to write this review.

References

- [ATY] H. Awata, A. Tsuchiya and Y. Yamada, Integral formulas for the WZNW correlation functions, Nucl. Phys. B 365 (1991) 680–698.
- [BaF1] H.M. Babujian and R. Flüme, Off-shell Bethe Ansatz equation for Gaudin magnets and solutions of Knizhnik-Zamolodchikov equations, Mod. Phys. Lett. A 9 (1994) 2029–2039.

- [BD1] A. Beilinson and V. Drinfeld, Quantization of Hitchin's integrable system and Hecke eigensheaves, Preprint, available at www.math.uchicago.edu/~benzvi.
- [BD2] A. Beilinson and V. Drinfeld, Opers, Preprint.
- [BM] L. Borisov and E. Mukhin, Self-self-dual spaces of polynomials, Preprint math. QA/0308128.
- [CS] S. Chmutov and I. Scherbak, On Bethe vectors in the Gaudin model, Preprint math.RT/0407367.
- [DS] V. Drinfeld and V. Sokolov, Lie algebras and KdV type equations, J. Sov. Math. 30 (1985) 1975–2036.
- [ER] B. Enriquez and V. Rubtsov, *Hitchin systems, higher Gaudin operators and R-matrices*, Math. Res. Lett. 3 (1996) 343–357.
- [FF1] B. Feigin, E. Frenkel, Affine Kac-Moody Algebras and semi-infinite flag manifolds, Comm. Math. Phys. 128, 161–189 (1990).
- [FF2] B. Feigin and E. Frenkel, Affine Kac-Moody algebras at the critical level and Gelfand-Dikii algebras, Int. Jour. Mod. Phys. A7, Supplement 1A (1992) 197–215.
- [FFR] B. Feigin, E. Frenkel and N. Reshetikhin, Gaudin model, Bethe Ansatz and critical level, Comm. Math. Phys. 166 (1994) 27–62.
- [F1] E. Frenkel, Affine algebras, Langlands duality and Bethe Ansatz, in Proceedings of the International Congress of Mathematical Physics, Paris, 1994, ed. D. Iagolnitzer, pp. 606–642, International Press, 1995.
- [F2] E. Frenkel, Lectures on Wakimoto modules, opers and the center at the critical level, Preprint math.QA/0210029.
- [F3] E. Frenkel, Opers on the projective line, flag manifolds and Bethe Ansatz, Preprint math.QA/0308269, to appear in Moscow Math. Journal.
- [FB] E. Frenkel and D. Ben-Zvi, Vertex algebras and algebraic curves, Second Edition, Mathematical Surveys and Monographs, vol. 88. AMS 2004.
- [FG] E. Frenkel and D. Gaitsgory, to appear.
- [K] V.G. Kac, Infinite-dimensional Lie Algebras, 3rd Edition, Cambridge University Press, 1990.
- [M] E. Markman, Spectral curves and integrable systems, Compositio Math. 93 (1994) 255–290.
- [MV1] E. Mukhin and A. Varchenko, Critical points of master functions and flag varieties, Preprint math.QA/0209017.
- [MV2] E. Mukhin and A. Varchenko, Multiple orthogonal polynomials and a counterexample to Gaudin Bethe Ansatz Conjecture, Preprint math.QA/0501144.
- [RV] N. Reshetikhin and A. Varchenko, Quasiclassical asymptotics of solutions of the KZ equations, in Geometry, Topology and Physics for Raoul Bott, pp. 293–322, International Press, 1994.
- [S1] I. Scherbak, Rational functions with prescribed critical points, Geom. Anal. Funct. Anal. 12 (2002) 1–16.
- [S2] I. Scherbak, A theorem of Heine-Stieltjes, the Wronski map, and Bethe vectors in the sl_p Gaudin model, Preprint math.AG/0211377.

[SV] I. Scherbak and A. Varchenko, Critical points of functions, sl₂ representations, and Fuchsian differential equations with only univalued solutions, Moscow Math. J. 3 (2003), no. 2, 621–645.

Department of Mathematics, University of California, Berkeley, CA 94720, USA

60