

A SHORT NOTE ON THE NON-NEGATIVITY OF PARTIAL EULER CHARACTERISTICS

TONY J. PUTHENPURAKAL

ABSTRACT. Let (A, \mathfrak{m}) be a Noetherian local ring, M a finite A -module and $x_1, \dots, x_n \in \mathfrak{m}$ such that $\lambda(M/\mathbf{x}M)$ is finite. Serre [2, Appendix 2] proved that all partial Euler characteristics of M with respect to \mathbf{x} is non-negative. This fact is easy to show when A contains a field [1, 4.7.12]. We give an elementary proof of Serre's result when A does not contain a field.

Let (A, \mathfrak{m}) be a Noetherian local ring and M a finite A -module. Let $x_1, \dots, x_n \in \mathfrak{m}$ be a *multiplicity system* of M i.e. $\lambda(M/(\mathbf{x})M)$ is finite. (Here $\lambda(-)$ denotes length). Let $K(\mathbf{x}, M)$ be the Koszul complex of \mathbf{x} with coefficients in M and let $H_\bullet(\mathbf{x}, M)$ be its homology. Note that $H_\bullet(\mathbf{x}, M)$ has finite length. One defines for all $j \geq 0$ the *partial Euler characteristics*

$$\chi_j(\mathbf{x}, M) = \sum_{i \geq j} (-1)^{i-j} \lambda(H_i(\mathbf{x}, M))$$

of M with respect to \mathbf{x} . Serre showed all the partial Euler characteristics are non-negative. It is well known that $\chi_0(\mathbf{x}, M)$ is either zero or the multiplicity of M with respect to the ideal (x_1, \dots, x_n) . It is also easy to see that $\chi_1(\mathbf{x}, M)$ is non-negative, [1, 4.7.10]. The non-negativity of $\chi_j(\mathbf{x}, M)$ for $j \geq 2$ can be easily proved if A contains a field, [1, 4.7.12]. In this short note we give an elementary proof of Serre's Theorem when A does not contain a field.

Theorem 1. *Let (A, \mathfrak{m}) be a Noetherian local ring, A not containing a field. Let M be a finite A -module and $x_1, \dots, x_n \in \mathfrak{m}$ a multiplicity system of M . Then*

$$\chi_j(\mathbf{x}, M) \geq 0 \quad \text{for each } j \geq 0.$$

Proof. We may assume that A is complete. To prove the theorem we construct a local Noetherian ring (B, \mathfrak{m}) with a local homomorphism $\varphi : B \rightarrow A$, and $y_1, \dots, y_n \in \mathfrak{m}$ such that

Date: June 26, 2018.

- (1) $\varphi(y_i) = x_i$.
- (2) M becomes a finite B -module (via φ).
- (3) y_1, \dots, y_n is a regular sequence and a s.o.p of B .

Since $K(\mathbf{y}, M) \simeq K(\mathbf{x}, M)$ (as B -modules), we have $H_\bullet(\mathbf{y}, M) \cong H_\bullet(\mathbf{x}, M)$ and so $\chi_j(\mathbf{y}, M) = \chi_j(\mathbf{x}, M)$ for each $j \geq 0$.

Suppose we have constructed B as above. The result then follows on similar lines as in [1, 4.7.12]. We give the proof here for the readers convenience. We prove the result by induction on j . For $j = 0, 1$ the result is already known. Let $j > 1$ and consider an exact sequence

$$0 \rightarrow U \rightarrow F \rightarrow M \rightarrow 0$$

where F is a finite free B -module. Since \mathbf{y} is B -regular we have $H_i(\mathbf{y}, F) = 0$ for $i > 0$. Therefore $H_i(\mathbf{y}, M) \simeq H_{i-1}(\mathbf{y}, U)$ for all $i > 1$. This yields $\chi_j(\mathbf{y}, M) = \chi_{j-1}(\mathbf{y}, U)$ and the proof is complete by induction hypothesis.

Construction of B

Since A is complete there exists a DVR, (R, ρ) and a ring homomorphism $\varphi : R \rightarrow A$ which induces an isomorphism $R/\rho R \rightarrow A/\mathfrak{m}$. Set $S = R[[X_1, \dots, X_n]]$ and let \mathfrak{q} be its maximal ideal and consider the natural ring map $\phi : S \rightarrow A$, with $\phi(X_i) = x_i$.

We consider M as an S -module via ϕ . Since $M/(\mathbf{X})M = M/(\mathbf{x})M$ is a finite length A -module and so a finite length S -module, since $S/\mathfrak{q}S \cong A/\mathfrak{m}$. So M is a finite S -module. Also note that

$$\mathfrak{q} = \sqrt{\text{ann}_S(M/\mathbf{X}M)} = \sqrt{\text{ann}_S(M) + (\mathbf{X})}.$$

So there exists $\Delta \in \text{ann}_S(M) \setminus (\mathbf{X})$. Observe that Δ, X_1, \dots, X_n is an s.o.p. of S . Since S is regular local ring of dimension $n + 1$, we have that Δ, X_1, \dots, X_n is an S -regular sequence. Set $B = S/\Delta$ and $y_i = \overline{X_i}$ for $i = 1, \dots, n$. Note that B satisfies our requirements. \square

Acknowledgment : The author thanks Prof. W. Bruns and Prof. J. Herzog for helpful discussions.

REFERENCES

- [1] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, vol. 39, Cambridge studies in advanced mathematics, Cambridge, 1993.
- [2] J. P. Serre, *Local Algebra*, Springer Monographs in Mathematics, Springer-Verlag, 2000.

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076
E-mail address: `tputhen@math.iitb.ac.in`