RESOLUTIONS OVER KOSZUL ALGEBRAS

EDWARD L. GREEN, GREGORY HARTMAN, EDUARDO N. MARCOS, AND $\varnothing {\rm YVIND}$ SOLBERG

ABSTRACT. In this paper we show that if $\Lambda = \coprod_{i \ge 0} \Lambda_i$ is a Koszul algebra with Λ_0 isomorphic to a product of copies of a field, then the minimal projective resolution of Λ_0 as a right Λ -module provides all the information necessary to construct both a minimal projective resolution of Λ_0 as a left Λ -module and a minimal projective resolution of Λ as a right module over the enveloping algebra of Λ . The main tool for this is showing that there is a comultiplicative structure on a minimal projective resolution of Λ_0 as a right Λ -module.

INTRODUCTION AND PRELIMINARIES

Let $\Lambda = \coprod_{i\geq 0}\Lambda_i$ be a Koszul algebra over a field k with Λ_0 a product of copies of k, where we recall the definition of Koszul later in this section. Denote by (\mathbb{L}, e) a minimal (graded) projective resolution of Λ_0 as a right Λ -module. We show that (\mathbb{L}, e) contains all the information needed to construct a minimal projective resolution of Λ as a right Λ^e -module, where $\Lambda^e = \Lambda^{\mathrm{op}} \otimes_k \Lambda$. The resolution (\mathbb{L}, e) is shown to have a "comultiplicative structure". This structure is used to prove that one can obtain a minimal projective resolution of Λ_0 over Λ as a left Λ -module from the knowledge of (\mathbb{L}, e) . We apply these results to prove an unpublished result of E. L. Green and D. Zacharia that Λ is a Koszul algebra if and only if Λ is a linear module as a right module over Λ^e . In [2], the comultiplicative structure is applied to give the multiplicative structure of the Hochschild cohomology ring of a Koszul algebra and also the structure constants for a basis for the Koszul dual.

The rest of the section is devoted to recalling definitions, results, and terminology relevant to this paper. Let $\Lambda = \coprod_{i \ge 0} \Lambda_i$ be a graded algebra over a field k. Assume that (i) Λ_0 is a product of copies of k, that (ii) each Λ_i is finite dimensional over k, and that (iii) Λ as an algebra is generated in degrees 0 and 1. Such an algebra Λ is isomorphic to a quotient of the path algebra kQ/I, where kQ is isomorphic to the tensor algebra $T_{\Lambda_0}(\Lambda_1) = \coprod_{i \ge 0} \underbrace{\Lambda_1 \otimes_{\Lambda_0} \cdots \otimes_{\Lambda_0} \Lambda_1}_{i}$. Conversely, if Q is a quotient of

I is an ideal generated by length homogeneous elements in kQ, then $\Lambda = kQ/I$ is a graded algebra over k satisfying the conditions above. Throughout this paper Λ denotes a graded algebra having properties (i)–(iii).

Let $\mathfrak{r} = \coprod_{i \ge 1} \Lambda_i$, which is the graded Jacobson radical of Λ . If (\mathbb{P}, d) :

$$\cdots \to P^2 \xrightarrow{d^2} P^1 \xrightarrow{d^1} P^0 \xrightarrow{d^0} M \to 0$$

Date: October 29, 2018.

The three first authors received financial support from a joint project of CNPq-NSF. The last author thanks the Fapesp-Brasil for financial support for a scientific visit to Brasil where this work was done. Finally the third authour thanks CNPq for a research grant.

is a graded projective resolution of a graded Λ -module M, then it is *minimal* if $\operatorname{Im} d^n \subseteq \mathfrak{r} P^{n-1}$ for $n \geq 1$. It is well known that graded modules over graded algebras have minimal graded projective resolutions. We say that a graded projective resolution

$$\cdots \to P^2 \xrightarrow{d^2} P^1 \xrightarrow{d^1} P^0 \xrightarrow{d^0} M \to 0$$

is linear, and M is a linear module if, for $n \ge 0$, the graded module P^n is generated in degree n. Note that a linear resolution is a minimal projective resolution. A graded algebra Λ is a Koszul algebra if Λ_0 is a linear module; that is, Λ_0 has a linear (graded) projective resolution (\mathbb{L}, e) :

$$\cdots \to L^2 \xrightarrow{e^2} L^1 \xrightarrow{e^1} L^0 \xrightarrow{e^0} \Lambda_0 \to 0$$

as a right Λ -module.

Before giving the precise results, we introduce notation and recall results from [5] which are used throughout the paper. For ease of notation, let R = kQ, let \mathcal{B} be the set of all paths in the quiver Q, and denote by \mathcal{B}_t all the paths of length t.

There exist integers $\{t_n\}_{n\geq 0}$ and elements $\{f_i^n\}_{i=0}^{t_n}$ in R such that a minimal right projective resolution (\mathbb{L}, e) of Λ_0 can be given in terms of a filtration of right ideals

$$\cdots \subseteq \coprod_{i=0}^{t_n} f_i^n R \subseteq \coprod_{i=0}^{t_{n-1}} f_i^{n-1} R \subseteq \cdots \subseteq \coprod_{i=0}^{t_1} f_i^1 R \subseteq \coprod_{i=0}^{t_0} f_i^0 R = R$$

in *R*. Then $L^n = \coprod_{i=0}^{t_n} f_i^n R / \coprod_{i=0}^{t_n} f_i^n I$ and the differential *e* is induced by the inclusion $\coprod_{i=0}^{t_n} f_i^n R \subseteq \coprod_{i=0}^{t_{n-1}} f_i^{n-1} R$. This inclusion gives elements $h_{ji}^{n-1,n}$ in *R* such that

$$f_i^n = \sum_{j=0}^{t_{n-1}} f_j^{n-1} h_{ji}^{n-1,n}$$

for all $i = 0, 1, \ldots, t_n$ and all $n \ge 1$, so that

$$e^{n}(\overline{f_{i}^{n}}) = (\overline{h_{0i}^{n-1,n}}, \overline{h_{1i}^{n-1,n}}, \dots, \overline{h_{t_{n-1}i}^{n-1,n}})$$

for all $n \geq 1$, where $\overline{*}$ denotes the natural residue class of * modulo I. It is shown in [5] that the f_i^n 's can be chosen so that (\mathbb{L}, e) is a minimal resolution of Λ_0 over Λ . We point out that an algorithmic construction of the elements f_i^n 's can be found in [4].

An important property of the elements $\{f_i^n\}_{i=0}^{t_n}$ is that there exist elements $f_j^{n+1'}$ in $\coprod_{i=0}^{t_{n-1}} f_i^n I$ such that

$$(\amalg_{i=0}^{t_n} f_i^n R) \cap (\amalg_{i=0}^{t_{n-1}} f_i^n I) = (\amalg_{i=0}^{t_{n+1}} f_i^{n+1} R) \amalg (\amalg_j f_j^{n+1'} R).$$

Recall that an element x in R is called *uniform* if x is non-zero and there exist vertices u and v in Q such that x = uxv. If x is a uniform element with x = uxv, then we write $\mathfrak{o}(x) = u$ and $\mathfrak{t}(x) = v$. The elements f_i^n can all be chosen uniform for $i = 0, 1, \ldots, t_n$ and all $n \ge 0$, and we assume that they are.

Note that $t_0 + 1$ is the number of non-isomorphic graded simple right Λ -modules, and that $\{f_i^0\}_{i=0}^{t_0}$ is the set of vertices of Q. Moreover, $t_1 + 1$ is the number of arrows of Q and $\{f_i^1\}_{i=0}^{t_1}$ is choosen to be the set of arrows of Q. The set $\{f_i^2\}_{i=0}^{t_2}$ is a set of uniform length homogeneous minimal generators for I.

In case Λ is a Koszul algebra, we have the following additional property of the elements f_i^n in R; namely each f_i^n is a linear combination of paths in \mathcal{B}_n for

 $i = 0, 1, \ldots, t_n$ and the length of each path occurring in $f_i^{n'}$ is at least n + 1. By length considerations, $h_{j_i}^{n-1,n}$ are all linear combinations of elements in \mathcal{B}_1 .

In section 1 we prove that the elements $\{f_i^n\}_{i=0,n\geq 0}^{t_n}$ have the following "comultiplicative structure", which is used in [2] to give the multiplicative structure of the Hochschild cohomology ring of a Koszul algebra and the structure constants for the basis associated to the elements $\{f_i^n\}$ for the Koszul dual.

Theorem. Let $\Lambda = kQ/I$ be a Koszul algebra. Then for each r, with $0 \le r \le n$, and i, with $0 \le i \le t_n$, there exist elements $c_{pq}(n, i, r)$ in k such that

$$f_i^n = \sum_{p=0}^{t_r} \sum_{q=0}^{t_{n-r}} c_{pq}(n, i, r) f_p^r f_q^{n-r}$$

for all $n \ge 1$, all *i* in $\{0, 1, ..., t_n\}$ and all *r* in $\{0, 1, ..., n\}$.

Viewing Λ_0 as a left module over Λ , it also has a minimal graded projective resolution given by $\{g_i^n\}_{i=0}^{s_n}$, where g_i^n 's are the left analogue of the right f_j^n 's in R. The above result is used to prove that one can choose the elements g_i^n 's to be the same as the elements f_j^n 's and then the formula $f_i^n = \sum_{p,q=0}^{t_1,t_{n-1}} c_{pq}(n,i,1) f_p^1 f_q^{n-1}$ gives the differential in the projective resolution of Λ_0 as a left Λ -module. Thus the knowledge of the minimal projective resolution (\mathbb{L}, e) via the elements $\{f_i^n\}$ contains all the information needed to construct a minimal projective resolution of Λ_0 as a left Λ -module.

In the final section of the paper, the elements f_i^n 's are shown to provide all the information needed to construct a minimal projective resolution of Λ as a right Λ^e -module. In particular, we prove the following.

Theorem. Let $\Lambda = kQ/I$ be a Koszul algebra, and let $\{f_i^n\}_{i=0}^{t_n}$ be defined as above for Λ_0 as a right Λ -module. A minimal projective resolution (\mathbb{P}, δ) of Λ over Λ^e is given by

$$P^n = \coprod_{i=0}^{t_n} \Lambda \mathfrak{o}(f_i^n) \otimes_k \mathfrak{t}(f_i^n) \Lambda$$

for $n \ge 0$, where *j*-th component of the differential $\delta^n \colon P^n \to P^{n-1}$ applied to the *i*-th generator $\mathfrak{o}(f_i^n) \otimes \mathfrak{t}(f_i^n)$ is given by

$$\sum_{p=0}^{t_1} c_{pj}(n,i,1) \overline{f_p^1} \mathfrak{o}(f_j^{n-1}) \otimes \mathfrak{t}(f_j^{n-1}) + (-1)^n \sum_{q=0}^{t_1} c_{jq}(n,i,n-1) \mathfrak{o}(f_j^{n-1}) \otimes \mathfrak{t}(f_j^{n-1}) \overline{f_q^1}$$

for $j = 0, 1, ..., t_{n-1}$ and $n \ge 1$, and $\delta^0 \colon \coprod_{i=0}^{t_0} \Lambda e_i \otimes_k e_i \Lambda \to \Lambda$ is the multiplication map.

As mentioned earlier, the final result of the paper is that Λ is a Koszul algebra if and only if Λ is a linear module as a right Λ^e -module.

1. A resolution with comultiplicative structure

In this section Theorem 1.1 provides a comultiplicative structure to a minimal projective resolution of Λ_0 as a right Λ -module. This result is then applied to show that the knowledge of a minimal projective resolution of Λ_0 as a right Λ -module is sufficient to construct a minimal projective resolution of Λ_0 as a left Λ -module.

Let $\Lambda = kQ/I$ be a graded algebra over a field k. Let $\{t_n\}_{n\geq 0}$ and $\{f_i^n\}_{i=0,n\geq 0}^{t_n}$ be as in the introduction. We say that $\{f_i^n\}_{i=0,n\geq 0}^{t_n}$ defines a minimal resolution if the resolution described in the introduction is minimal.

The next result shows that the elements $\{f_i^n\}$ have a comultiplicative structure for a Koszul algebra.

Theorem 1.1. Let $\Lambda = kQ/I$ be a Koszul algebra, and assume that $\{f_i^n\}_{i=0}^{t_n}$ defines a minimal resolution of Λ_0 as a right Λ -module. Then for each r, with $0 \le r \le n$, and i, with $0 \le i \le t_n$, there exist elements $c_{pq}(n, i, r)$ in k such that

$$f_i^n = \sum_{p=0}^{t_r} \sum_{q=0}^{t_{n-r}} c_{pq}(n, i, r) f_p^r f_q^{n-r}.$$

Proof. For any n, and r equal to 0 or n, the result follows from $f_i^n = f_i^n f_{\mathfrak{t}(f_i^n)}^0 = f_{\mathfrak{o}(f_i^n)}^0 f_i^n$ for $i = 0, 1, \ldots, t_n$. Also, this proves the result in the case n is equal to 1. Next we discuss the case n = 2. As we have remarked, each $f_i^2 = \sum_{j=0}^{t_1} f_j^1 h_{ji}^{1,2}$.

Next we discuss the case n = 2. As we have remarked, each $f_i^2 = \sum_{j=0}^{t_1} f_j^1 h_{ji}^{1,2}$. Since Λ is Koszul, each f_i^2 is a linear combination of paths in \mathcal{B}_2 , and hence $h_{ji}^{1,2}$ is a linear combination of elements in \mathcal{B}_1 . This gives the result for n = 2.

Now we proceed by induction on n and assume that the result is true for l < nand $n \ge 3$. We have that $f_i^n = \sum_{j=0}^{t_{n-1}} f_j^{n-1} h_{ji}^{n-1,n}$. As in our discussion for n = 2, we see that $h_{ji}^{n-1,n}$ is a linear combination of elements in \mathcal{B}_1 . There exist elements c_{ijs} in k such that

$$f_i^n = \sum_{j=0}^{t_{n-1}} \sum_{s=0}^{t_1} c_{ijs} f_j^{n-1} f_s^1$$

By induction, there exist elements c'_{juv} in k such that

$$f_j^{n-1} = \sum_{u=0}^{t_r} \sum_{v=0}^{t_{n-r-1}} c'_{juv} f_u^r f_v^{n-r-1}$$

for any r, with $0 \le r \le n-1$. Hence

$$f_i^n = \sum_j \sum_s \sum_u \sum_v c_{ijs} c'_{juv} f_u^r f_v^{n-r-1} f_s^1$$

for any r, with $0 \le r \le n-1$. The term after f_u^r is

(1)
$$A = \sum_{j} \sum_{s} \sum_{v} c_{ijs} c'_{juv} f_{v}^{n-r-1} f_{s}^{1}.$$

Theory tells us that

$$f_i^n = \sum_{w=0}^{t_{n-2}} f_w^{n-2} z_w,$$

where z_w is in *I*. Again by length considerations each z_w is a linear combination of f_l^2 's. Hence, there exist elements c''_{iwx} in k such that

$$f_i^n = \sum_{w=0}^{t_{n-2}} \sum_{x=0}^{t_2} c_{iwx}'' f_w^{n-2} f_x^2.$$

By induction each f_w^{n-2} is a linear combination of $f_u^r f_y^{n-r-2}$. We obtain

$$f_i^n = \sum_w \sum_x \sum_u \sum_y c_{iwx}'' c_{wuy}'' f_u^r f_y^{n-r-2} f_x^2$$

for some c''_{wuy} in k. So the term after f_u^r in this expression is

(2)
$$B = \sum_{w} \sum_{x} \sum_{u} c''_{wuy} c'''_{wuy} f_y^{n-r-2} f_x^2.$$

Since $\sum_{u} f_{u}^{r} R$ is a direct sum, we see that formulas (1) and (2) are equal. The equation (1) implies that A is in $\prod_{v=0}^{t_{n-r-1}} f_{v}^{n-r-1} R$, and the equation (2) implies that A is in $\prod_{y=0}^{t_{n-r-2}} f_{y}^{n-r-2} I$. It follows that A is contained in $(\prod_{t=0}^{t_{n-r}} f_{t}^{n-r} R) \prod$ $(\prod_{l} f_{l}^{n-r'} R)$. By length arguments we infer that A is in $\prod_{t=0}^{t_{n-r}} f_{t}^{n-r} R$ and that A is a k-linear combination of the f_{t}^{n-r} 's. Hence we conclude that f_{i}^{n} is a k-linear combination of $f_{s}^{r} f_{t}^{n-r}$, and this completes the proof of the result.

Since the maps in the minimal projective resolution of Λ_0 as a right Λ -module are given by the $h_{ji}^{n-1,n}$, we explicitly point out the following relationship.

Corollary 1.2. Keeping the notation of Theorem 1.1, we have

$$h_{ji}^{n-1,n} = \sum_{l=0}^{t_1} f_l^1 c_{jl}(n, i, n-1)$$

for $n \ge 1$, and i and j, with $0 \le i \le t_n$, and $0 \le j \le t_{n-1}$.

Before applying Theorem 1.1 we need the following lemma, where J denotes the ideal generated by the arrows in Q.

Lemma 1.3. Let $\{x_i\}_{i \in \mathcal{I}}$ be a set of elements in the linear span of \mathcal{B}_s . Suppose that $\{x_i\}_{i \in \mathcal{I}}$ is linearly independent viewed as vectors over k. Then $\sum_{i \in \mathcal{I}} Rx_i$ and $\sum_{i \in \mathcal{I}} x_i R$ are direct sums.

Proof. Suppose that $\sum_{i} \sum_{j \in \mathcal{I}} c_{ij} q_{ij} x_j = 0$ in R for some elements c_{ij} in k and some paths q_{ij} in R. Since all the paths occurring in any x_i have the same length, we can assume without loss of generality that the paths q_{ij} all have the same length, say t. Since $J^t = \prod_{q \in B_t} qR$, fixing q in \mathcal{B}_t , it follows that $\sum_{q_{ij}=q} q(c_{ij}x_j) = 0$, which implies that $\sum_{q_{ij}=q} c_{ij}x_j = 0$. By assumption, we have that $c_{ij} = 0$ for all $q_{ij} = q$. Hence we infer that $\sum_{i \in \mathcal{I}} Rx_i$ is a direct sum. Similarly, $\sum_{i \in \mathcal{I}} x_i R$ is a direct sum.

We now show that the $\{f_i^n\}$ obtained from a right minimal projective resolution of Λ_0 and the $\{g_j^n\}$ obtained from a left minimal projective resolution of Λ_0 can be chosen to be the same.

Proposition 1.4. Let $\Lambda = kQ/I$ be a Koszul algebra. Let $\{f_i^n\}_{i=0}^{t_n}$ and $\{g_i^n\}_{i=0}^{s_n}$ define a minimal resolution of Λ_0 as a right Λ -module and as a left Λ -module, respectively. Then $s_n = t_n$ for all $n \ge 0$ and the set $\{g_i^n\}_{i=0}^{t_n}$ can be chosen to be equal to the set $\{f_i^n\}_{i=0}^{t_n}$ for all $n \ge 0$.

Proof. For n equal to 0, 1, or 2 the result is clear. Let $n \ge 3$. We proceed by induction on n and assume that the result is true for all i < n. By Theorem 1.1, for each i with $0 \le i \le t_n$ the equalities

$$f_i^n = \sum_{p,q} c_{pq} f_p^1 f_q^{n-1} = \sum_{p',q'} c_{p',q'} f_{p'}^2 f_{q'}^{n-2}$$

hold for some c_{pq} and $c_{p',q'}$ in k. Hence f_i^n is in $(\coprod_q R f_q^{n-1}) \cap (\coprod_{q'} I f_{q'}^{n-2})$, which, by induction, is equal to $(\coprod_{i=0}^{s_n} R g_i^n) \amalg (\coprod_i R g_i^{n'})$. Since $\sum_i f_i^n R$ is direct, the set $\{f_i^n\}$ is linearly independent as vectors over k, and therefore the sum $\sum_i Rf_i^n$ is direct by Lemma 1.3. By length considerations, $\{f_i^n\}_{i=0}^{t_n}$ is contained in the k-linear span of $\{g_i^n\}_{i=0}^{s_n}$. Therefore $t_n \leq s_n$. By switching the roles of $\{f_i^n\}$ and $\{g_j^n\}$ and using the argument above, we conclude that $\{g_i^n\}_{i=0}^{s_n}$ is linearly independent and each g_i^n is in k-linear span of $\{f_i^n\}_{i=0}^{t_n}$. Hence $s_n = t_n$. By Lemma 1.3 it follows that $\coprod_{i=0}^{t_n} Rf_i^n = \coprod_{i=0}^{t_n} Rg_i^n$. This shows that we can choose the set $\{g_i^n\}_{i=0}^{t_n}$ equal to $\{f_i^n\}_{i=0}^{t_n}$.

Proposition 1.4 implies that, given a minimal projective resolution of Λ_0 as a right Λ -module in the from of $\{f_i^n\}$, we have all the information to construct a minimal projective resolution of Λ_0 as a left Λ -module. More precisely, take the $\{f_i^n\}$ as the $\{g_i^n\}$, and the maps in the left resolution are given by $g_i^n \mapsto \sum_{p=0}^{t_1} \sum_{q=0}^{t_{n-1}} c_{pq}(n, i, 1) g_p^1 g_q^{n-1}$.

2. A minimal projective bimodule resolution of Λ

In this section we turn our attention to the construction of a minimal projective Λ^{e} -resolution of Λ . This construction uses the comultiplicative structure of the minimal projective resolution of Λ_{0} as a right Λ -module found in Theorem 1.1. This is applied to show an unpublished result of E. L. Green and D. Zacharia that Λ is a Koszul algebra if and only if Λ is a (right) linear module over Λ^{e} .

The following result also shows that the knowledge of the $\{f_i^n\}$ from a minimal projective resolution of Λ_0 as a right Λ -module is sufficient to explicitly give the projective modules and the differentials in a minimal projective resolution of Λ as a right Λ^e -module. The structure of the projective modules in a minimal projective resolution of Λ as right Λ^e -module was first given in [6]. Recall that the notation $\overline{*}$ denotes the natural residue class of * modulo I. Let $\{c_{pq}(n, i, r)\}$ be as in Theorem 1.1.

Theorem 2.1. Let $\Lambda = kQ/I$ be a Koszul algebra, and let $\{f_i^n\}_{i=0}^{t_n}$ define a minimal resolution of Λ_0 as a right Λ -module. A minimal projective resolution (\mathbb{P}, δ) of Λ over Λ^e is given by

$$P^n = \coprod_{i=0}^{t_n} \Lambda \mathfrak{o}(f_i^n) \otimes_k \mathfrak{t}(f_i^n) \Lambda$$

for $n \ge 0$, where the *j*-th component of the differential $\delta^n \colon P^n \to P^{n-1}$ applied to the *i*-th generator $\mathfrak{o}(f_i^n) \otimes \mathfrak{t}(f_i^n)$ is given by

$$\sum_{p=0}^{t_1} c_{pj}(n,i,1) \overline{f_p^1} \mathfrak{o}(f_j^{n-1}) \otimes \mathfrak{t}(f_j^{n-1}) + (-1)^n \sum_{q=0}^{t_1} c_{jq}(n,i,n-1) \mathfrak{o}(f_j^{n-1}) \otimes \mathfrak{t}(f_j^{n-1}) \overline{f_q^1}$$

for $j = 0, 1, ..., t_{n-1}$ and $n \ge 1$, and $\delta^0 \colon \coprod_{i=0}^{t_0} \Lambda e_i \otimes_k e_i \Lambda \to \Lambda$ is the multiplication map.

In particular, Λ is a linear module over Λ^e .

Proof. Direct computations show that $(\delta)^2 = 0$, so that (\mathbb{P}, δ) is a linear complex. In addition, note that $(\Lambda_0 \otimes_{\Lambda} \mathbb{P}, 1_{\Lambda_0} \otimes \delta)$ is a minimal resolution of Λ_0 as a right Λ -module.

In our setting, we have that $\Lambda^e/\operatorname{rad} \Lambda^e \simeq \operatorname{Hom}_k(\Lambda_0, \Lambda_0)$. Let (\mathbb{F}, d) be a minimal projective resolution of Λ as a right Λ^e -module. Then by [3, Chap. IX, Proposition

[4.3] we have that

$$\operatorname{Hom}_{\Lambda^{e}/\operatorname{rad}\Lambda^{e}}(F^{n}/F^{n}\operatorname{rad}\Lambda^{e},\Lambda^{e}/\operatorname{rad}\Lambda^{e}) \simeq \operatorname{Ext}_{\Lambda^{e}}^{n}(\Lambda,\Lambda^{e}/\operatorname{rad}\Lambda^{e})$$
$$\simeq \operatorname{Ext}_{\Lambda}^{n}(\Lambda_{0},\Lambda_{0})$$
$$\simeq \operatorname{Hom}_{\Lambda_{0}}(\amalg_{i=0}^{t_{n}}f_{i}^{n}R/\amalg_{i=0}^{t_{n}}f_{i}^{n}J,\Lambda_{0})$$

for all $n \geq 0$. In particular, $P^n \simeq F^n$ as Λ^e -modules for all $n \geq 0$, and hence $P^n/P^n \operatorname{rad} \Lambda^e \simeq F^n/F^n \operatorname{rad} \Lambda^e$ as $\Lambda^e/\operatorname{rad} \Lambda^e$ -modules for all $n \geq 0$. Note that these need not be isomorphic as graded modules, but, we in fact show that this is the case.

Since (\mathbb{P}, δ) is a complex, we obtain the following commutative diagram

$$\begin{array}{c} \cdots \longrightarrow P^2 \xrightarrow{\delta^2} P^1 \xrightarrow{\delta^1} P^0 \xrightarrow{\delta^0} \Lambda \longrightarrow 0 \\ & \downarrow^{\alpha^2} & \downarrow^{\alpha^1} & \downarrow^{\alpha^0} \\ \cdots \longrightarrow F^2 \xrightarrow{d^2} F^1 \xrightarrow{d^1} F^0 \xrightarrow{d^0} \Lambda \longrightarrow 0 \end{array}$$

Clearly $\alpha^0 \colon P^0 \to F^0$ is an isomorphism, and we get an isomorphism $\alpha^0|_{\operatorname{Ker}\delta^0} \colon \operatorname{Ker}\delta^0 \to \operatorname{Ker}d^0$. Hence $\operatorname{Ker}\delta^0 \operatorname{rad}\Lambda^e \simeq F^1/F^1 \operatorname{rad}\Lambda^e$. Since $\operatorname{Im}\delta^1$ is contained in $\operatorname{Ker}\delta^0$, this induces a map $\beta^1 \colon P_1^1 \to \operatorname{Ker}\delta^0/\operatorname{Ker}\delta^0 \operatorname{rad}\Lambda^e$. If β^1 is an isomorphism, then α^1 is an isomorphism and we have exactness at P^0 . Suppose that β^1 is not an isomorphism. Since P^1 is generated in degree 1, there is some projective summand of P^1 which is mapped to zero by δ^1 . Using the observation that $(\Lambda_0 \otimes_{\Lambda} \mathbb{P}, \mathbb{1}_{\Lambda_0} \otimes \delta)$ is a minimal resolution of Λ_0 as a right Λ -module, we obtain a contradiction. Hence β^1 is an isomorphism.

Since α^1 is an isomorphism, we can use the above argument replacing α^0 by α^1 to show that α^2 is an isomorphism and exactness at P^1 . By induction we infer that (\mathbb{P}, δ) is exact. Since the terms $\overline{f_p^1}\mathfrak{o}(f_j^{n-1}) \otimes \mathfrak{t}(f_j^{n-1})$ and $\mathfrak{o}(f_j^{n-1}) \otimes \mathfrak{t}(f_j^{n-1}) \overline{f_q^1}$ are elements of degree one in Λ^e , we conclude that (\mathbb{P}, δ) is a minimal linear projective resolution of Λ over Λ^e . This also implies that Λ is a linear module over Λ^e . The proof is now complete.

As a consequence we obtain the next corollary which was first proved by E. L. Green and D. Zacharia.

Corollary 2.2. Let $\Lambda = kQ/I$ be a graded algebra. Then Λ is a Koszul algebra if and only if Λ as a (right) Λ^{e} -module is a linear module.

Proof. Suppose that Λ is a Koszul algebra. Then Theorem 2.1 implies that Λ has a linear projective Λ^e -resolution, and hence Λ is a linear module over Λ^e .

Suppose that Λ is a linear module over Λ^e as a right module. Let (\mathbb{P}, δ) be a linear projective resolution of Λ as a right Λ^e -module. Tensoring \mathbb{P} with Λ_0 , we obtain $\Lambda_0 \otimes_{\Lambda} \mathbb{P}$. But $\Lambda_0 \otimes_{\Lambda} \mathbb{P}$ is a linear projective resolution of Λ_0 as a right Λ -module. Hence Λ is a Koszul algebra, and we are done.

References

- Buchweitz, R.-O., Green, E. L., Madsen, D., Solberg, Ø., Finite Hochschild cohomology without finite global dimension, preprint.
- Buchweitz, R.-O., Green, E. L., Snashall, N., Solberg, Ø., Multiplicative structures for Koszul algebras, preprint.
- [3] Cartan, H., Eilenberg, S., Homology algebra, Princeton University Press, Princeton Mathematical series, 33 (1956).

- [4] Green, E. L., Solberg, Ø., An algorithmic resolution construction, preprint.
- [5] Green, E. L., Solberg, Ø., Zacharia, D., Minimal projective resolutions, Trans. Amer. Math. Soc., 353 (2001), 2915–2939.
- [6] Happel, D., Hochschild cohomology of finite-dimensional algebras, Séminaire d'Algèbre Paul Dubreil et Marie-Paul Malliavin, 39ème Année (Paris, 1987/1988), 108–126, Lecture Notes in Math., 1404, Springer, Berlin, 1989.

Edward L. Green, Department of Mathematics, Virginia Tech, Blacksburg, VA 24061, USA

E-mail address: green@math.vt.edu

Gregory Hartman, Department of Mathematics, The University of Arizona, 617 N. Santa Rita Ave., P.O. Box 210089, Tucson, AZ 85721-0089, USA

 $E\text{-}mail\ address: hartman@math.arizona.edu$

Eduardo N. Marcos, Instituto de Matemática e Estatística, Universidade São Paulo (IME-USP), Rua do Matão, 1010 - Cidade Universitária, CEP 05508-090, São Paulo - SP - Brazil

 $E\text{-}mail\ address:\ \texttt{enmarcos@ime.usp.br}$

Øvvind Solberg, Institutt for matematiske fag, NTNU, N-7491 Trondheim, Norway $E\text{-}mail\ address:\ oyvinso@math.ntnu.no$

8