

Classification of PM Quiver Hopf Algebras

Shouchuan Zhang ^{a, b}, Yao-Zhong Zhang ^b, Hui-xiang Chen ^c

a. Department of Mathematics, Human University

Changsha 410082, P.R. China

b. Department of Mathematics, University of Queensland

Brisbane 4072, Australia

c. Department of Mathematics, Yangzhou University

Yangzhou 225002, P.R. China

Abstract

We give the classification of quiver Hopf algebras with pointed module structures. This leads to the classification of multiple crown algebras and multiple Taft algebras as well as pointed Yetter-Drinfeld modules and their corresponding Nichols algebras. In particular, we give the classification of graded Hopf algebras on cotensor coalgebra $T_{kG}^c(M)$ of kG -bicomodule M over finite commutative group algebra kG and tensor algebra $T_{kG}(M)$ of dimensional finite kG - bimodule M over finite commutative group algebra kG with $|G|=n$, when k contains a primitive n -th root of 1 and the order of G is not divisible by $\text{char}(k)$. In this case, we also give the classification of Yetter-Drinfeld kG -modules and Nichols algebras of Yetter-Drinfeld kG -modules.

2000 Mathematics Subject Classification: 16W30, 16G10

keywords: Quiver, Hopf algebra, Hopf bimodule

Introduction

Classification of Hopf algebras would have applications to a number of other areas of mathematics, aside from its intrinsic algebraic interest. In mathematical physics, Drinfeld's and Jimbo's work was to provide solutions to quantum Yang-Baxter equation. In conformal field theory, I. Frenkel and Y. Zhu have shown how to assign a Hopf algebra to any conformal field theory model[13]. In topology, quasi-triangular and ribbon Hopf algebras provide many invariants of knots, links, tangles and 3-manifolds[14, 15, 23, 25]. In operator algebras, Hopf algebras can be assigned as an invariant for certain extensions[27].

It has been a long time since quivers were first used to study algebra structures. Quivers have played an important role in the representation theory of algebras (see e.g. [6]). It has been realized that quivers may be applied to the study of Hopf algebras and quantum groups. For instance, C. Cibils and M. Rosso obtained a necessary and sufficient condition for a path (co)algebra of a quiver to admit a graded Hopf algebra structure in [9] [10]. The classification of monomial Hopf algebras, which are a class of co-path Hopf algebras, and simple-pointed sub-Hopf algebras of co-path Hopf algebras were recently obtained in [7] and [21], respectively.

N. Andruskiewitsch and H. J. Schneider have obtained interesting result in classification of finite-dimensional pointed Hopf algebras with commutative coradical [2, 3, 4, 5]. More recently, they have also researched this problem in case of non-commutative coradical. Pavel Etingof and Shlomo Gelaki gave the complete and explicit classification of finite-dimensional triangular Hopf algebras over an algebraically closed field k of characteristic 0 [12],

Let G be a group and kG be the group algebra of G over a field k . It is well-known [10] that the kG -Hopf bimodule category ${}^{kG}_k\mathcal{M}_k^{kG}$ is equivalent to the direct product category $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, where $\mathcal{K}(G)$ is the set of conjugate classes in G , $u : \mathcal{K}(G) \rightarrow G$ is a map such that $u(C) \in C$ for any $C \in \mathcal{K}(G)$, $Z_{u(C)} = \{g \in G \mid gu(C) = u(C)g\}$ and $\mathcal{M}_{kZ_{u(C)}}$ denotes the category of right $kZ_{u(C)}$ -modules (see [10, Proposition 3.3] [9, Theorem 4.1] and Theorem 1.1). Thus for any Hopf quiver (Q, G, r) , the kG -Hopf bimodule structures on the arrow comodule kQ_1^c can be derived from the right $kZ_{u(C)}$ -module structures on ${}^{u(C)}(kQ_1^c)^1$ for all $C \in \mathcal{K}(G)$. If the arrow comodule kQ_1^c admits a kG -Hopf bimodule structure, then there exist six graded Hopf algebras: co-path Hopf algebra kQ^c , one-type-co-path Hopf algebras $kG[kQ_1^c]$, one-type-path Hopf algebras $(kG)^*[kQ_1^a]$, semi-path Hopf algebra kQ^s , semi-co-path Hopf algebra kQ^{sc} and path Hopf algebra kQ^a . We call these Hopf algebras quiver Hopf algebras. If the corresponding $kZ_{u(C)}$ -modules ${}^{u(C)}(kQ_1^c)^1$ is pointed (i.e. it is zero or a direct sum of one dimensional $kZ_{u(C)}$ -modules) for all $C \in \mathcal{K}(G)$, then kQ_1^c is called a PM kG -Hopf bimodule. The six graded Hopf algebras derived from the PM kG -Hopf bimodule kQ_1^c are called PM quiver Hopf algebras. A YD H -module V is *pointed* if $V = 0$ or V is a direct sum of one dimensional YD H -modules. If V is a pointed YD H -module, then the corresponding Nichols algebra $\mathcal{B}(V)$ is called a PM *Nichols algebra*. For example, when G is a finite abelian group with $|G| = n$ and k contains a primitive n -th root of 1 and the order of G is not divisible by $\text{char}(k)$, all quiver Hopf algebras are PM ones (see Lemma 1.2), all YD kG -modules are PM ones and all Nichols algebras of all YD kG -modules are PM ones (see Lemma 2.3).

The aim of this paper is to give the classification of PM quiver Hopf algebras and PM Nichols algebras. We use the ramification system with characters to classify PM

quiver Hopf algebras (see Theorem 4) and element system with characters to classify PM Nichols algebras (see Theorem 5). Moreover, we characterize quantum enveloping algebras in terms of semi-path Hopf algebras. We also point out that many finite dimensional Hopf algebras, such as Taft algebras, the bosonization of quantum linear spaces (see [3, Lemma 3.4]) and $H(C, n, c, c^*)$ (see [11, Definition 5.6.15]) are multiple Taft algebras. We also obtain an interesting quantum combinatoric formula:

$$\sum_{\sigma \in S_n} q^{\tau(\sigma)} = (n)_{\frac{1}{q}}! q^{\frac{n(n-1)}{2}}$$

(see Example 3.12).

This paper is organized as follows. In Section 0, we recall some basic definitions and results for later use. In Section 1, we restate a well-known theorem about the category of Hopf bimodules and classify the PM quiver Hopf algebras by ramification system with characters. In Section 2, we discuss the PM quiver Hopf algebras determined by central ramification system with characters. Moreover, we study the corresponding pointed Yetter-Drinfeld modules and PM Nichols algebra. In Section 3, we give the relation between quiver Hopf algebras and quotients of free algebras.

Throughout this paper, Hopf quivers are finite whenever arrow modules, arrow Hopf bimodules, path algebras, path Hopf algebras, semi-co-path Hopf algebras are considered. kG -Hopf bimodule M is finite dimensional and G is a finite group whenever tensor Hopf algebra $T_{(kG)^*}(M^*)$ over $(kG)^*$ are considered.

The main results in this paper are summarized in the following statement.

Theorem 1 (I) *Let $(G, r, \vec{\chi}, u)$ and $(G', r', \vec{\chi}', u')$ are two RSC's. Then the following statements are equivalent: (i) $RSC(G, r, \vec{\chi}, u) \cong RSC(G', r', \vec{\chi}', u')$. (ii) $(kG; (kQ_1^c, \alpha^X))$ and $(kG'; (kQ_1'^c, \alpha^{X'}))$ are pull-push Hopf bimodule isomorphism. (iii) $kQ^c(\alpha^X) \cong kQ'^c(\alpha^{X'})$. (iv) $kQ^s(\alpha^X) \cong kQ'^s(\alpha^{X'})$. (v) $kG[kQ_1, \alpha^X] \cong kG'[kQ_1', \alpha^{X'}]$. (vi) $kQ^a(\alpha^X) \cong kQ'^a(\alpha^{X'})$. (vii) $kQ^{sc}(\alpha^X) \cong kQ'^{sc}(\alpha^{X'})$. (viii) $(kG)^*[kQ_1^a, \alpha^X] \cong (kG')^*[kQ_1'^a, \alpha^{X'}]$. (ix) $(kG; \text{diag}(kG[kQ_1, \alpha^X])) \cong (kG'; \text{diag}(kG'[kQ_1', \alpha^{X'}]))$ as graded pull-push Yetter-Drinfeld Hopf algebras. (x) $((kG)^*; \text{diag}((kG)^*[kQ_1^a, \alpha^X])) \cong ((kG')^*; \text{diag}((kG')^*[kQ_1'^a, \alpha^{X'}]))$ as graded pull-push Yetter-Drinfeld Hopf algebras. (xi) $(kG; (M, [\alpha^X]))$ and $(kG'; (M', [\alpha^{X'}]))$ are pull-push Hopf bimodule isomorphism. (xii) $T_{kG}(M, [\alpha^X]) \cong T_{kG'}(M', [\alpha^{X'}])$. (xiii) $T_{kG}^c(M, [\alpha^X]) \cong T_{kG'}^c(M', [\alpha^{X'}])$. (xiv) $T_{(kG)^*}((M, [\alpha^X])^*) \cong T_{(kG')^*}((M', [\alpha^{X'}])^*)$. (xv) $T_{(kG)^*}^c((M, [\alpha^X])^*) \cong T_{(kG')^*}^c((M', [\alpha^{X'}])^*)$. (xvi) $kG[(M, [\alpha^X])] \cong kG'[(M', [\alpha^{X'}])]$. Note that all isomorphisms above are as graded Hopf algebras but in (i), (ii), (ix), (ix) and (xi).*

(II) *Let $(G, g_i, \chi_i; i \in J)$ and $(G', g'_j, \chi'_j; j \in J')$ be two ESC's. Then the following statements are equivalent: (i) $ESC(G, g_i, \chi_i; i \in J) \cong ESC(G', g'_j, \chi'_j; j \in J')$. (ii) $RSC(G, r(g_i, \chi_i; i \in J), \vec{\chi}(g_i, \chi_i; i \in J), u) \cong RSC(G', r'(g'_j, \chi'_j; j \in J'), \vec{\chi}'(g'_j, \chi'_j; j \in J'), u')$*

$J'), u')$. (iii) $(kG; (kQ_1^1, ad(g_i, \chi_i : i \in J)))$ and $(kG'; (kQ_1^1, ad(g'_i, \chi'_i : i \in J')))$ are pull-push Yetter-Drinfeld module isomorphism. (iv) $(kG; \mathcal{B}(kQ_1^1, ad(g_i, \chi_i : i \in J)))$ and $(kG'; \mathcal{B}(kQ_1^1, ad(g'_i, \chi'_i : i \in J')))$ are pull-push Yetter-Drinfeld Hopf algebra isomorphism. (v) $(kG; \mathcal{T}(G, g_i, \chi_i; i \in J)) \cong (kG'; \mathcal{T}(G', g'_j, \chi'_j, j \in J'))$ as graded pull-push Yetter-Drinfeld Hopf algebras.

(III) (i) If $(G, g_i, \chi_i; i \in J)$ is a quantum weakly commutative ESC, then $\mathcal{L}(G, g_i, \chi_i^{-1}; j \in J) \cong \text{diag}(kG[kQ_1, g_i, \chi_i; i \in J])$ as graded braided Hopf algebras in ${}_{kG}^{kG}\mathcal{YD}$ algebras. (ii) $\mathcal{T}(G, g_i, \chi_i^{-1}; j \in J) \cong \text{diag}(kQ^s(g_i, \chi_i; i \in J))$ as graded braided Hopf algebras in ${}_{kG}^{kG}\mathcal{YD}$ algebras. (iii) $\overline{kQ^s}(g_i, \chi_i; i \in J; A; d_i) \cong U(g_i, \chi_i; i \in J; A; d_i)$ as Hopf algebras for symmetrizable generalized Cartan matrix $A = (a_{ij})$. (iv) $U_q(L) \cong kQ^s(g_i, \chi_i; i \in J)/I$ as Hopf algebras, where $U_q(L)$ is the quantum enveloping algebra of semisimple Lie algebra L determined by Cartan matrix A .

Note that (I) follows from Theorem 4 and Lemma 24. (II) follows from Theorem 5, Theorem 7, Lemma 24 and Lemma 2.6, and (III) follows from Theorem 7 and Example 3.9.

0 Preliminaries

Throughout this paper, we work over a fixed field k . All algebras, coalgebras, Hopf algebras, and so on, are defined over k . Books [11, 19, 26] provide the necessary background for Hopf algebras and book [6] provides a nice description of the path algebra approach.

Let \mathbf{Z} denote the set of all integers. For sets X and Y , we denote by $|X|$ the cardinal number of X and by X^Y or $X^{\mathbb{M}}$ the Cartesian product $\prod_{y \in Y} X_y$ with $X_y = X$ for any $y \in Y$. If X is finite, then $|X|$ is the number of elements in X . For a vector space V , any two bases of V have the same cardinal number, which is called the dimension of V , written $\dim V$. If $X = \bigoplus_{i \in I} X_i$ as vector spaces, then we denote by ι_i the natural injection from X_i to X and by π_i the corresponding projection from X to X_i . We will use μ to denote the multiplication map of an algebra and use Δ to denote the comultiplication of a coalgebra. For a (left or right) module and a (left or right) comodule, denote by α^- , α^+ , δ^- and δ^+ the left module, right module, left comodule and right comodule structure maps, respectively. The Sweedler's sigma notations for coalgebras and comodules are $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$, $\delta^-(x) = \sum x_{(-1)} \otimes x_{(0)}$, $\delta^+(x) = \sum x_{(0)} \otimes x_{(1)}$. Let G be a group. We denote by $Z(G)$ the center of G . Let \widehat{G} denote the set of characters of all one-dimensional representations of G . It is clear that $\widehat{G} = \{\chi \mid \chi \text{ is a group homomorphism from } G \text{ to the multiplicative group of all non-zero elements in } k\}$.

A quiver $Q = (Q_0, Q_1, s, t)$ is an oriented graph, where Q_0 and Q_1 are the sets of vertices and arrows, respectively; s and t are two maps from Q_1 to Q_0 . For any arrow

$a \in Q_1$, $s(a)$ and $t(a)$ are called its start vertex and end vertex, respectively, and a is called an arrow from $s(a)$ to $t(a)$. For any $n \geq 0$, an n -path or a path of length n in the quiver Q is an ordered sequence of arrows $p = a_n a_{n-1} \cdots a_1$ with $t(a_i) = s(a_{i+1})$ for all $1 \leq i \leq n-1$. Note that a 0-path is exactly a vertex and a 1-path is exactly an arrow. In this case, we define $s(p) = s(a_1)$, the start vertex of p , and $t(p) = t(a_n)$, the end vertex of p . For a 0-path x , we have $s(x) = t(x) = x$. Let Q_n be the set of n -paths. Let ${}^y Q_n^x$ denote the set of all n -paths from x to y , $x, y \in Q_0$. That is, ${}^y Q_n^x = \{p \in Q_n \mid s(p) = x, t(p) = y\}$.

A quiver Q is *finite* if Q_0 and Q_1 are finite sets. A quiver Q is *locally finite* if ${}^y Q_1^x$ is a finite set for any $x, y \in Q_0$.

Let G be a group. Let $\mathcal{K}(G)$ denote the set of conjugate classes in G . A formal sum $r = \sum_{C \in \mathcal{K}(G)} r_C C$ of conjugate classes of G with cardinal number coefficients is called a *ramification* (or *ramification data*) of G , i.e. for any $C \in \mathcal{K}(G)$, r_C is a cardinal number. In particular, a formal sum $r = \sum_{C \in \mathcal{K}(G)} r_C C$ of conjugate classes of G with non-negative integer coefficients is a ramification of G . For any ramification r and a $C \in \mathcal{K}(G)$, since r_C is a cardinal number, we can choose a set $I_C(r)$ such that its cardinal number is r_C without lost generality. Let $\mathcal{K}_r(G) := \{C \in \mathcal{K}(G) \mid r_C \neq 0\} = \{C \in \mathcal{K}(G) \mid I_C(r) \neq \emptyset\}$.

Let G be a group. A quiver Q is called a *quiver of group G* if $Q_0 = G$ (i.e., $Q = (G, Q_1, s, t)$). In this paper we only consider quivers of group. If there exists a ramification r of G such that the cardinal number of ${}^y Q_1^x$ is equal to r_C for any $x, y \in G$ with $x^{-1}y \in C \in \mathcal{K}(G)$, then Q is called a *Hopf quiver with respect to the ramification data r* . In this case, there is a bijection from $I_C(r)$ to ${}^y Q_1^x$, and hence we may regard ${}^y Q_1^x = \{a_{y,x}^{(i)} \mid i \in I_C(r)\}$ for any $x, y \in G$ with $x^{-1}y \in C \in \mathcal{K}(G)$. With the notations above, one can see that a ramification r of G defines a unique Hopf quiver Q of G . This means that a Hopf quiver of G is uniquely determined by a ramification of G . Denote by (Q, G, r) the Hopf quiver of G with respect to r .

Let $Q = (Q_0, Q_1, s, t)$ and $Q' = (Q'_0, Q'_1, s', t')$ be two quivers with $Q_0 = G$ and $Q'_0 = G'$. Q and Q' are *isomorphic* if there exist a group isomorphism $\phi_0 : Q_0 \rightarrow Q'_0$ and a bijection $\phi_1 : Q_1 \rightarrow Q'_1$ such that $\phi_0(s(a)) = s'(\phi_1(a))$ and $\phi_0(t(a)) = t'(\phi_1(a))$ for any $a \in Q_1$. If r and r' are ramifications of G and G' , respectively, then the Hopf quivers (Q, G, r) and (Q', G', r') are isomorphic if and only if there exists a group isomorphism $\phi_0 : G \rightarrow G'$ such that $r_C = r'_{\phi_0(C)}$ for any $C \in \mathcal{K}(G)$, i.e., there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi_0(C)}(r')$. In this case, the path algebras (path coalgebras) of isomorphic quivers are isomorphic.

Throughout the following, let G be a group with unity 1 and let u be a fixed map from $\mathcal{K}(G)$ to G such that $u(C) \in C$ for any $C \in \mathcal{K}(G)$. Let $Z_{u(C)} = \{g \in G \mid gu(C) = u(C)g\}$. Then $Z_{u(C)}$ is a subgroup of G .

The coset decomposition of $Z_{u(C)}$ in G is

$$G = \bigcup_{\theta \in \Theta_C} Z_{u(C)}g_\theta, \quad (1)$$

where Θ_C is an index set. It is easy to check that $|\Theta_C| = |C|$. We always assume that the representative element of the coset $Z_{u(C)}$ is the identity 1 of G . We claim that $\theta = \eta$ if $g_\theta^{-1}u(C)g_\theta = g_\eta^{-1}u(C)g_\eta$. In fact, by the equation $g_\theta^{-1}u(C)g_\theta = g_\eta^{-1}u(C)g_\eta$ one gets $g_\eta g_\theta^{-1}u(C) = u(C)g_\eta g_\theta^{-1}$. Thus $g_\eta g_\theta^{-1} \in Z_{u(C)}$, and hence $\theta = \eta$. For any $x, y \in G$ with $x^{-1}y \in C \in \mathcal{K}(G)$, there exists a unique $\theta \in \Theta_C$ such that

$$x^{-1}y = g_\theta^{-1}u(C)g_\theta. \quad (2)$$

Without specification, we will always assume that x, y, θ and C satisfy the above relation (2). Note that θ is only determined by $x^{-1}y$. For any $h \in G$ and $\theta \in \Theta_C$, there exist unique $h' \in Z_{u(C)}$ and $\theta' \in \Theta_C$ such that $g_\theta h = h'g_{\theta'}$. Let $\zeta_\theta(h) = h'$. Then we have

$$g_\theta h = \zeta_\theta(h)g_{\theta'}. \quad (3)$$

If $u(C)$ lies in the center $Z(G)$ of G , we have $\zeta_\theta = id_G$. In particular, if G is abelian, $\zeta_\theta = id_G$ since $Z_{u(C)} = G$.

Let H be a bialgebra. Assume that M is simultaneously an H -bimodule (M, α^-, α^+) and an H -bicomodule (M, δ^-, δ^+) . Then $(M, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is called an H -Hopf bimodule, if the two module structure maps α^+ and α^- are H -bicomodule homomorphisms, or equivalently, the two comodule structure maps δ^- and δ^+ are H -bimodule homomorphisms. We denote by ${}^H_H\mathcal{M}_H^H$ the category of H -Hopf bimodules; the morphisms in this category are both H -bimodule and H -bicomodule homomorphisms.

Let H be a Hopf algebra. A (left-left) *Yetter-Drinfeld module* V over H (simply, YD H -module) is simultaneously a left H -module and a left H -comodule satisfying the following compatibility condition:

$$\sum (h \cdot v)_{(-1)} \otimes (h \cdot v)_{(0)} = \sum h_{(1)}v_{(-1)}S(h_{(3)}) \otimes h_{(2)} \cdot v_{(0)}, \quad v \in V, h \in H. \quad (4)$$

We denote by ${}^H_H\mathcal{YD}$ the category of YD H -modules; the morphisms in this category preserve both the action and the coaction of H . The category ${}^H_H\mathcal{YD}$ is a pre-braided monoidal category (see [24, 19]). Furthermore, if H be a Hopf algebra with invertible antipode, then ${}^H_H\mathcal{YD}$ is a braided monoidal category. A braided monoidal category is also called a braided tensor category.

For the definitions of an algebra, a coalgebra and a Hopf algebra in a braided monoidal category, the reader is directed to [18, 19].

The structure of a Nichols algebra appeared first in the paper [20] and was rediscovered later by several authors.

Let V be a YD H -module. A graded braided Hopf algebra $R = \bigoplus_{n \geq 0} R_n$ in ${}^H_H\mathcal{YD}$ is called a *Nichols algebra* of V if the following are satisfied:

- $k \cong R_0$,
- $V \cong R_1$ in ${}^H_H\mathcal{YD}$,
- $R_1 = P(R)$, the vector space of the primitive elements of R ,
- R is generated as an algebra by R_1 .

For any YD H -module V , it follows from [4, Proposition 2.2] that there exists a unique Nichols algebra $\mathcal{B}(V)$ of V up to graded braided Hopf algebra isomorphism in ${}^H_H\mathcal{YD}$.

If $\phi : A \rightarrow A'$ is an algebra homomorphism and (M, α^-) is a left A' -module, then M becomes a left A -module with the A -action given by $a \cdot x = \phi(a) \cdot x$ for any $a \in A$, $x \in M$, called a pullback A -module through ϕ , written as ${}_\phi M$. Similarly, if M is a right A' -module, then M becomes a right A -module with the A -action given by $x \cdot a = x \cdot \phi(a)$ for any $a \in A$ and $x \in M$, denoted by M_ϕ . In particular, if M is an A' -bimodule then ${}_\phi M_\phi$ is an A -bimodule. Dually, if $\phi : C \rightarrow C'$ be a coalgebra homomorphism and (M, δ^-) is a left C -comodule, then M is a left C' -comodule with the C' -comodule structure given by $\delta'^- := (\phi \otimes \text{id})\delta^-$, called a push-out C' -comodule through ϕ , written as ${}^\phi M$. Similarly, if (M, δ^+) be a right C -comodule, then M becomes a right C' -comodule with the C' -coaction given by $\delta'^+ := (\text{id} \otimes \phi)\delta^+$, denote it by M^ϕ . In particular, if M is a C -bicomodule then ${}^\phi M^\phi$ is a C' -bicomodule. For convenience, if B is a Hopf algebra and M is a B -Hopf bimodule, then we say that $(B; M)$ is a Hopf bimodules. For any two Hopf bimodules $(B; M)$ and $(B'; M')$, and ψ is a B -bimodule homomorphism from M to ${}_\phi M'_\phi$ and a B' -bicomodule homomorphism from ${}^\phi M^\phi$ to M' , then (ϕ, ψ) is called a pull-push Hopf bimodule homomorphism. If both ϕ and ψ are bijective, then (ϕ, ψ) is called a pull-push Hopf bimodule isomorphism. Similarly, we say that $(B; M)$ and $(B; X)$ are a Yetter-Drinfeld module and a Yetter-Drinfeld Hopf algebra if M is a Yetter-Drinfeld B -module and X is a braided Hopf algebra in Yetter-Drinfeld category ${}^B_B\mathcal{YD}$, respectively. For any two Yetter-Drinfeld modules $(B; M)$ and $(B'; M')$, if ϕ is a Hopf algebra homomorphism from B to B' , and ψ is simultaneously a left B -module homomorphism from M to ${}_\phi M'$ and a left B' -comodule homomorphism from ${}^\phi M$ to M' , then (ϕ, ψ) is called a pull-push Yetter-Drinfeld module homomorphism. For any two Yetter-Drinfeld Hopf algebras $(B; X)$ and $(B'; X')$, if ϕ is a Hopf algebra homomorphism from B to B' , ψ is simultaneously a left B -module homomorphism from X to ${}_\phi X'$ and a left B' -comodule homomorphism from ${}^\phi X$ to X' , and ψ is algebra and coalgebra homomorphism from X to X' , then (ϕ, ψ) is called a pull-push Yetter-Drinfeld Hopf algebra homomorphism.

Let A be an algebra and M be an A -bimodule. Then the tensor algebra $T_A(M)$ of M over A is a graded algebra with $T_A(M)_0 = A$, $T_A(M)_1 = M$ and $T_A(M)_n = \bigotimes_A^n M$ for $n > 1$. That is, $T_A(M) = A \oplus (\bigoplus_{n > 0} \bigotimes_A^n M)$ (see [20]). Let D be another algebra. If h is

an algebra map from A to D and f is an A -bimodule map from M to hD_h , then by the universal property of $T_A(M)$ (see [20, Proposition 1.4.1]) there is a unique algebra map $T_A(h, f) : T_A(M) \rightarrow D$ such that $T_A(h, f)\iota_0 = h$ and $T_A(h, f)\iota_1 = f$. One can easily see that $T_A(h, f) = h + \sum_{n>0} \mu^{n-1}T_n(f)$, where $T_n(f)$ is the map from $\otimes_A^n M$ to $\otimes_A^n D$ given by $T_n(f)(x_1 \otimes x_2 \otimes \cdots \otimes x_n) = f(x_1) \otimes f(x_2) \otimes \cdots \otimes f(x_n)$, i.e., $T_n(f) = f \otimes_A f \otimes_A \cdots \otimes_A f$. Note that μ can be viewed as a map from $D \otimes_A D$ to D . For the details, the reader is directed to [20, Section 1.4].

Dually, let C be a coalgebra and let M be a C -bicomodule. Then the cotensor coalgebra $T_C^c(M)$ of M over C is a graded coalgebra with $T_C^c(M)_0 = C$, $T_C^c(M)_1 = M$ and $T_C^c(M)_n = \square_C^n M$ for $n > 1$. That is, $T_C^c(M) = C \oplus (\bigoplus_{n>0} \square_C^n M)$ (see [20]). Let D be another coalgebra. If h is a coalgebra map from D to C and f is a C -bicomodule map from ${}^hD^h$ to M such that $f(\text{corad}(D)) = 0$, then by the universal property of $T_C^c(M)$ (see [20, Proposition 1.4.2]) there is a unique coalgebra map $T_C^c(h, f)$ from D to $T_C^c(M)$ such that $\pi_0 T_C^c(h, f) = h$ and $\pi_1 T_C^c(h, f) = f$. It is not difficult to see that $T_C^c(h, f) = h + \sum_{n>0} T_n^c(f)\Delta_{n-1}$, where $T_n^c(f)$ is the map from $\square_C^n D$ to $\square_C^n M$ induced by $T_n^c(f)(x_1 \otimes x_2 \otimes \cdots \otimes x_n) = f(x_1) \otimes f(x_2) \otimes \cdots \otimes f(x_n)$, i.e., $T_n^c(f) = f \otimes f \otimes \cdots \otimes f$.

1 Classification of PM quiver Hopf algebras

In this section we classify the PM quiver Hopf algebras.

We first describe the category of Hopf bimodules by categories of modules.

Let G be a group and let (B, δ^-, δ^+) be a kG -bicomodule. Then the (x, y) -isotypic component of B is

$${}^yB^x = \{b \in B \mid \delta^-(b) = y \otimes b, \delta^+(b) = b \otimes x\},$$

where $x, y \in G$. Let M be another kG -bicomodule and $f : B \rightarrow M$ be a kG -bicomodule homomorphism. Then $f({}^yB^x) \subseteq {}^yM^x$ for any $x, y \in G$. Denote by ${}^y f^x$ the restriction map $f|_{{}^yB^x} : {}^yB^x \rightarrow {}^yM^x$, $x, y \in G$.

The following theorem can be found in [10]. We list the result here for the completeness and give a slightly different proof.

Theorem 2 (See [10, Proposition 3.3] and [9, Theorem 4.1]) *The category ${}^{kG}_{kG}\mathcal{M}_{kG}^{kG}$ of kG -Hopf bimodules is equivalent to the Cartesian product category $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}^{kG}$ of categories $\mathcal{M}_{kZ_{u(C)}}$ of right $kZ_{u(C)}$ -modules for all $C \in \mathcal{K}(G)$.*

Proof. We define two functors W and V as follows.

$$\begin{aligned} W : {}^{kG}_{kG}\mathcal{M}_{kG}^{kG} &\rightarrow \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}^{kG} \\ B &\mapsto \{{}^{u(C)}B^1\}_{C \in \mathcal{K}(G)} \\ f &\mapsto W(f), \end{aligned}$$

where for any morphism $f : B \rightarrow B'$ in ${}_{kG}^k\mathcal{M}_{kG}^k$, $W(f) = \{u^{(C)}f^1\}_{C \in \mathcal{K}(G)}$.

$$\begin{aligned} V : \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}} &\rightarrow {}_{kG}^k\mathcal{M}_{kG}^k \\ \{M(C)\}_{C \in \mathcal{K}(G)} &\mapsto \bigoplus_{x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta, \\ f &\mapsto V(f), \end{aligned}$$

where for any morphism $f = \{f_C\}_{C \in \mathcal{K}(G)} : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$, $V(f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta) = x \otimes f_C(m) \otimes_{kZ_{u(C)}} g_\theta$ for any $m \in M_C$, $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$. That is, ${}^yV(f)^x = \text{id} \otimes f_C \otimes \text{id}$.

We will show that W and V are mutually inverse functors by three steps.

(i) Let B be a kG -Hopf bimodule. Then it is easy to see that ${}^{u(C)}B^1$ is a right $kZ_{u(C)}$ -module with the module action given by

$$b \triangleleft h = h^{-1} \cdot b \cdot h, \quad b \in {}^{u(C)}B^1, \quad h \in Z_{u(C)}. \quad (5)$$

Hence $W(B)$ is an object in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$. Moreover, one can see that $W(f)$ is a morphism from $W(B)$ to $W(B')$ in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ if $f : B \rightarrow B'$ is a morphism in ${}_{kG}^k\mathcal{M}_{kG}^k$. Now it is straightforward to check that W is a functor from ${}_{kG}^k\mathcal{M}_{kG}^k$ to $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$.

(ii) Let $M = \{M(C)\}_{C \in \mathcal{K}(G)} \in \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, we define

$$\begin{aligned} {}^yV(M)^x &= x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta, \\ h \cdot (x \otimes m \otimes_{kZ_{u(C)}} g_\theta) &= hx \otimes m \otimes_{kZ_{u(C)}} g_\theta, \\ (x \otimes m \otimes_{kZ_{u(C)}} g_\theta) \cdot h &= xh \otimes m \otimes_{kZ_{u(C)}} g_\theta h = xh \otimes (m \triangleleft \zeta_\theta(h)) \otimes_{kZ_{u(C)}} g_\theta, \end{aligned} \quad (6)$$

where $h, x, y \in G$ with $x^{-1}y \in C$ and the relation (2) and (3), $m \in M(C)$. It is easy to check that $V(M)$ is a kG -Hopf bimodule. If $f : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$ is a morphism in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, then one can see that $V(f) : V(M) \rightarrow V(N)$ is a morphism in ${}_{kG}^k\mathcal{M}_{kG}^k$. Now let $f : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$ and $f' : \{N(C)\} \rightarrow \{L(C)\}_{C \in \mathcal{K}(G)}$ be two morphisms in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$. Then for any $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$ and $m \in M(C)$, we have

$$\begin{aligned} V(f'f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta) &= x \otimes (f'f)_C(m) \otimes_{kZ_{u(C)}} g_\theta \\ &= x \otimes f'_C(f_C(m)) \otimes_{kZ_{u(C)}} g_\theta \\ &= V(f')(V(f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta)). \end{aligned}$$

Therefore, $V(f'f) = V(f')V(f)$. It follows that V is a functor from $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ to ${}_{kG}^k\mathcal{M}_{kG}^k$.

(iii) We now construct two natural isomorphisms φ and ψ as follows. Let $B \in {}_{kG}^k\mathcal{M}_{kG}^k$. Define a k -linear map φ_B from B to $VW(B) = \bigoplus_{x,y \in G} x \otimes {}^{u(C)}B^1 \otimes_{kZ_{u(C)}} g_\theta$ by

$$\varphi_B(b) = x \otimes (g_\theta x^{-1}) \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta, \quad (7)$$

for any $x, y \in G$ satisfying Eq.(2), $b \in {}^y B^x$. Now we have

$$\begin{aligned}
\delta^-(g_\theta x^{-1} \cdot b \cdot g_\theta^{-1}) &= (g_\theta x^{-1}) \cdot (\delta^-(b)) \cdot g_\theta^{-1} \quad (B \text{ is a } kG\text{-Hopf bimodule.}) \\
&= (g_\theta x^{-1}) \cdot (y \otimes b) \cdot g_\theta^{-1} \\
&= g_\theta x^{-1} y g_\theta^{-1} \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \\
&= u(C) \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \quad (\text{by Eq.(2)})
\end{aligned}$$

and

$$\delta^+(g_\theta x^{-1} \cdot b \cdot g_\theta^{-1}) = g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes 1.$$

It follows that $g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \in {}^{u(C)} B^1$, and hence $x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta \in {}^y VW(B)^x$.

This shows that φ_B is a kG -bicomodule map.

Next, let $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, $C \in \mathcal{K}(G)$, and let $b \in {}^y B^x$ and $h \in G$. Then $h \cdot b \in {}^{hy} B^{hx}$ and $(hx)^{-1}(hy) = x^{-1}y = g_\theta^{-1}u(C)g_\theta$. Hence

$$\begin{aligned}
\varphi_B(h \cdot b) &= hx \otimes g_\theta (hx)^{-1} \cdot (h \cdot b) \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta \\
&= hx \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta \\
&= h \cdot (x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta) \\
&= h \cdot \varphi_B(b).
\end{aligned}$$

This shows that φ_B is a left kG -module map. On the other hand, $b \cdot h \in {}^{yh} B^{xh}$ and $(xh)^{-1}(yh) = h^{-1}x^{-1}yh = h^{-1}g_\theta^{-1}u(C)g_\theta h$. By Eq.(3), one gets $g_\theta h = \zeta_\theta(h)g_{\theta'}$ with $\zeta_\theta(h) \in Z_{u(C)}$ and $\theta' \in \Theta_C$. Hence $(xh)^{-1}(yh) = g_{\theta'}^{-1}\zeta_\theta(h)^{-1}u(C)\zeta_\theta(h)g_{\theta'} = g_{\theta'}^{-1}u(C)g_{\theta'}$. Thus we have

$$\begin{aligned}
\varphi_B(b \cdot h) &= xh \otimes g_{\theta'} (xh)^{-1} \cdot (b \cdot h) \cdot g_{\theta'}^{-1} \otimes_{kZ_{u(C)}} g_{\theta'} \\
&= xh \otimes g_{\theta'} h^{-1} x^{-1} \cdot b \cdot h g_{\theta'}^{-1} \otimes_{kZ_{u(C)}} g_{\theta'} \\
&= xh \otimes \zeta_\theta(h)^{-1} g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \zeta_\theta(h) \otimes_{kZ_{u(C)}} g_{\theta'} \\
&= xh \otimes ((g_\theta x^{-1} \cdot b \cdot g_\theta^{-1}) \triangleleft \zeta_\theta(h)) \otimes_{kZ_{u(C)}} g_{\theta'} \quad (\text{by (5)}) \\
&= (x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta) \cdot h \\
&= \varphi_B(b) \cdot h.
\end{aligned}$$

It follows that φ_B is a right kG -module map, and hence φ_B is a kG -Hopf bimodule map.

Suppose that $\varphi_B(b) = 0$ with $b \in {}^y B^x$. Then $x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta = 0$, and hence $g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta = 0$. Since $M(C) \otimes_{kZ_{u(C)}} kZ_{u(C)} g_\theta \cong M(C) \otimes_{kZ_{u(C)}} kZ_{u(C)} \cong M(C)$ as k -vector spaces for any right $kZ_{u(C)}$ -module $M(C)$, we have $g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} = 0$. This implies $b = 0$. It follows that φ_B is injective.

Let $x \otimes b \otimes_{kZ_{u(C)}} g_\theta \in {}^y VW(B)^x$ with $b \in {}^{u(C)} B^1$ and $x^{-1}y = g_\theta^{-1}u(C)g_\theta$. Put $b' := xg_\theta^{-1} \cdot b \cdot g_\theta$. Then

$$\delta^-(b') = xg_\theta^{-1} \cdot \delta^-(b) \cdot g_\theta$$

$$\begin{aligned}
&= xg_\theta^{-1} \cdot (u(C) \otimes b) \cdot g_\theta \\
&= y \otimes b', \\
\delta^+(b') &= b' \otimes x.
\end{aligned}$$

Hence $b' \in {}^y B^x$ and $\varphi_B(b') = x \otimes b \otimes_{kZ_{u(C)}} g_\theta$. This shows that φ_B is surjective.

If $f : B \rightarrow B'$ is a kG -Hopf bimodule map, then it is easy to see that $\varphi_{B'} \circ f = (VW)(f) \circ \varphi_B$. Thus we have proven that φ is a natural equivalence from the identity functor $id_{\mathcal{B}(kG)}$ to the composition functor VW .

Now for any $M = \{M(C)\}_{C \in \mathcal{K}(G)} \in \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, it follows from Eq.(2) that ${}^{u(C)}V(M)^1 = 1 \otimes M(C) \otimes_{kZ_{u(C)}} 1$. Hence $WV(M) = \{1 \otimes M(C) \otimes_{kZ_{u(C)}} 1\}_{C \in \mathcal{K}(G)}$. Define a map ψ_M from M to $WV(M)$ by

$$(\psi_M)_C(m) = 1 \otimes m \otimes_{kZ_{u(C)}} 1,$$

where $C \in \mathcal{K}(G), m \in M(C)$. Clearly, $(\psi_M)_C$ is k -linear. Let $m \in M(C), h \in Z_{u(C)}$. Then

$$\begin{aligned}
((\psi_M)_C(m)) \triangleleft h &= (1 \otimes m \otimes_{kZ_{u(C)}} 1) \triangleleft h \\
&= h^{-1} \cdot (1 \otimes m \otimes_{kZ_{u(C)}} 1) \cdot h \\
&= 1 \otimes m \triangleleft h \otimes_{kZ_{u(C)}} 1 \\
&= (\psi_M)_C(m \triangleleft h).
\end{aligned}$$

That is, $(\psi_M)_C$ is a $kZ_{u(C)}$ -module map. Obviously, $(\psi_M)_C$ is bijective and ψ is a natural transformation. Thus ψ is a natural equivalence from the identity functor $id_{\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ to the composition functor WV . \square

An A -module M is called *pointed* if $M = 0$ or M is a direct sum of one dimensional A -modules.

Definition 1.1 *A kG -Hopf bimodule N is called a kG -Hopf bimodule with pointed module structure (or a PM kG -Hopf bimodule in short) if there exists an object $\prod_{C \in \mathcal{K}(G)} M(C)$ in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ such that $M(C)$ is a right pointed $kZ_{u(C)}$ -module for any $C \in \mathcal{K}(G)$ and $N \cong V(\{M(C)\}_{C \in \mathcal{K}(G)}) = \bigoplus_{x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ as kG -Hopf bimodules. Here $V(\{M(C)\}_{C \in \mathcal{K}(G)}) = \bigoplus_{x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ is defined in the proof of Theorem 2.*

Lemma 1.2 *Assume that G is a finite commutative group with $|G|=n$. If k contains a primitive n -th root of 1 and the order of G is not divisible by $\text{char}(k)$, then*

- (i) every kG -module is a pointed module;
- (ii) every kG -Hopf bimodule is a PM one.

Proof. (i) We first show that every irreducible representation of G is one dimensional. In fact, this is a well-known conclusion. For the completeness, we give a proof. Assume that ρ is an irreducible representation of G corresponding to V . For any $g \in G$, since $\rho(g^m) = id_V$, all eigenvalues of $\rho(g)$ is m -th roots of 1, which belong to k . Say that λ_g is an eigenvalue of $\rho(g)$. It is clear that $Ker(\rho(g) - \lambda_g id_V) := W$ is a kG -submodule of V and $W \neq 0$ since λ_g is an eigenvalue of $\rho(g)$. Therefore, $W = V$, which implies $\rho(g) = \lambda_g id_V$. Considering V is irreducible, we have that $dim V = 1$.

Using Maschke's Theorem, we have that every kG -module is a pointed module.

(ii) Assume M is a kG -Hopf bimodule. By Theorem 2, $M \cong \bigoplus_{x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g\theta$ as kG -Hopf bimodules with $M(C) = {}^{u(C)}M^1$ for any $C \in \mathcal{K}(G)$. It follows from (i) that $M(C)$ is a right pointed kG -module. Therefore, M is a PM kG -Hopf bimodule. \square

If ξ is a graded linear map from graded vector space $V = \bigoplus_{i \geq 0} V_i$ to graded vector space $V' = \bigoplus_{i \geq 0} V'_i$, then

$$\xi \iota_i \pi_i = \iota_i \pi_i \xi, \quad \xi \iota_i = \iota_i \pi_i \xi \iota_i, \quad \pi_i \xi = \pi_i \xi \iota_i \pi_i. \quad (8)$$

for $i = 1, 2, \dots$.

Lemma 1.3 *Let $H = \bigoplus_{i \geq 0} H_i$ be a graded Hopf algebra. Set $B := H_0$ and $M := H_1$. Then B is a Hopf algebra and M is a B -Hopf bimodule with the operations of B , the B -actions and B -coactions given by*

$$\mu_B = \pi_0 \mu(\iota_0 \otimes \iota_0), \quad \Delta_B = (\pi_0 \otimes \pi_0) \Delta \iota_0, \quad \epsilon_B = \epsilon \iota_0, \quad 1_B = 1_H. \quad (9)$$

$$\alpha^- = \pi_1 \mu(\iota_0 \otimes \iota_1), \quad \alpha^+ = \pi_1 \mu(\iota_1 \otimes \iota_0), \quad (10)$$

$$\delta^- = (\pi_0 \otimes \pi_1) \Delta \iota_1, \quad \delta^+ = (\pi_1 \otimes \pi_0) \Delta \iota_1. \quad (11)$$

Proof. Observe that (M, α^-, α^+) is a B -bimodule. Since Δ_H and μ_H are graded, we have the following equation by (8):

$$\Delta \iota_1 \pi_1 = (\iota_1 \pi_1 \otimes \iota_0 \pi_0) \Delta + (\iota_0 \pi_0 \otimes \iota_1 \pi_1) \Delta. \quad (12)$$

Since $\pi_1 \iota_1 = id$, it follows from Eq.(12) that

$$\Delta \iota_1 = (\iota_1 \pi_1 \otimes \iota_0 \pi_0) \Delta \iota_1 + (\iota_0 \pi_0 \otimes \iota_1 \pi_1) \Delta \iota_1. \quad (13)$$

By a straightforward computation, one can show the following two equations:

$$\delta^- \pi_1 = (\pi_0 \otimes \pi_1) \Delta, \quad \delta^+ \pi_1 = (\pi_1 \otimes \pi_0) \Delta. \quad (14)$$

Now we have

$$\begin{aligned} (\text{id} \otimes \delta^-)\delta^- &= (\text{id} \otimes \delta^-)(\pi_0 \otimes \pi_1)\Delta\iota_1 \\ &= (\pi_0 \otimes \pi_0 \otimes \pi_1)(\text{id} \otimes \Delta)\Delta\iota_1 \quad (\text{by Eq.(14)}), \end{aligned}$$

and

$$\begin{aligned} (\Delta_B \otimes \text{id})\delta^- &= (\Delta_B \otimes \text{id})(\pi_0 \otimes \pi_1)\Delta\iota_1 \\ &= (\pi_0 \otimes \pi_0 \otimes \pi_1)(\Delta \otimes \text{id})\Delta\iota_1. \end{aligned}$$

This shows that $(\text{id} \otimes \delta^-)\delta^- = (\Delta_B \otimes \text{id})\delta^-$. Clearly, $(\varepsilon \otimes \text{id})\delta^- = \text{id}$. Hence (M, δ^-) is a left B -comodule. Similarly, one can show that (M, δ^+) is a right B -comodule and (M, δ^-, δ^+) is a B -bicomodule. Next, we have

$$\begin{aligned} \delta^- \alpha^+ &= (\pi_0 \otimes \pi_1)\Delta\iota_1\pi_1\mu(\iota_1 \otimes \iota_0) \\ &= (\pi_0 \otimes \pi_1)\Delta\mu(\iota_1 \otimes \iota_0) \quad (\text{by Eq.(12)}) \\ &= (\pi_0\mu \otimes \pi_1\mu)(\text{id} \otimes \tau \otimes \text{id})(\Delta\iota_1 \otimes \Delta\iota_0) \\ &= (\mu_B \otimes \pi_1)(\pi_0 \otimes \pi_0 \otimes \mu)(\text{id} \otimes \tau \otimes \text{id})(\Delta \otimes \iota_0 \otimes \iota_0)(\iota_1 \otimes \Delta_B) \\ &= (\mu_B \otimes \pi_1)(\text{id} \otimes \text{id} \otimes \mu)(\text{id} \otimes \text{id} \otimes \iota_1 \otimes \iota_0)(\text{id} \otimes \tau \otimes \text{id})(\pi_0 \otimes \pi_1 \otimes \text{id} \otimes \text{id}) \\ &\quad (\Delta\iota_1 \otimes \Delta_B) \quad (\text{by Eq.(13)}) \\ &= (\mu \otimes \alpha^+)(\text{id} \otimes \tau \otimes \text{id})(\delta^- \otimes \Delta_B), \end{aligned}$$

where τ is the usual twist map. This shows that δ^- is a right B -module map. Similarly, one can show that δ^- is a left B -module map, and hence a B -bimodule map. A similar argument shows that δ^+ is also a B -bimodule map. Hence $(M, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a B -Hopf bimodule. \square

Lemma 1.4 *We have the following:*

(i) *Let B be an algebra and M be a B -bimodule. Then the tensor algebra $T_B(M)$ of M over B admits a graded Hopf algebra structure if and only if B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure.*

(ii) *Let B be a coalgebra and M be a B -bicomodule. Then the cotensor coalgebra $T_B^c(M)$ of M over B admits a graded Hopf algebra structure if and only if B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure.*

Proof. (i) Suppose that the tensor algebra $T_B(M)$ of M over B admits a graded Hopf algebra structure. Obviously, B is a Hopf algebra. It follows from Lemma 1.3 that M is a B -Hopf bimodule.

Conversely, suppose that B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure. By [20, Section 1.4], $T_B(M)$ is a bialgebra with the counit $\varepsilon = \varepsilon_B\pi_0$

and the comultiplication $\Delta = (\iota_0 \otimes \iota_0)\Delta_B + \sum_{n>0} \mu^{n-1}T_n(\Delta_M)$, where ε_B is the counit of B , $\Delta_M = (\iota_0 \otimes \iota_1)\delta_M^- + (\iota_1 \otimes \iota_0)\delta_M^+$. It is easy to see that $T_B(M)$ is a graded bialgebra. Then it follows from [20, Proposition 1.5.1] that $T_B(M)$ is a graded Hopf algebra.

(ii) Assume that the cotensor coalgebra $T_B^c(M)$ of M over B admits a graded Hopf algebra structure. Clearly, B is a Hopf algebra. By Lemma 1.3 one gets that M is a B -Hopf bimodule.

Conversely, assume that B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure. By [20, Section 1.4], $T_B^c(M)$ is a bialgebra containing B as a subbialgebra, and the multiplication of $T_B^c(M)$ is $\mu = \mu_B(\pi_0 \otimes \pi_0) + \sum_{n>0} T_n^c(\mu_M)\Delta_{n-1}$, where $\mu_M = \alpha_M^-(\pi_0 \otimes \pi_1) + \alpha_M^+(\pi_1 \otimes \pi_0)$. Since Δ preserves the grading, μ is a graded map. Thus $T_B^c(M)$ is a graded bialgebra. By [20, Proposition 1.5.1], $T_B^c(M)$ is a graded Hopf algebra. \square

Let B be a bialgebra (Hopf algebra) and M be a B -Hopf bimodule. Then $T_B^c(M)$ is a graded bialgebra (Hopf algebra) by Lemma 1.4. Let $B[M]$ denote the subalgebra of $T_B^c(M)$ generated by B and M . Then $B[M]$ is a bialgebra (Hopf algebra) of type one by [20, section 2.2, p.1533]. $B[M]$ is a graded subspace of $T_B^c(M)$.

Lemma 1.5 *Let B and B' be two Hopf algebras. Let M and M' be B -Hopf bimodule and B' -Hopf bimodule, respectively. Assume that $\phi : B \rightarrow B'$ be a Hopf algebra map. If ψ is a B -bimodule and B' -bicomodule map from ${}^\phi M^\phi$ to ${}_\phi M'_\phi$, then*

(i) $T_B(\iota_0\phi, \iota_1\psi) := \iota_0\phi + \sum_{n>0} \mu^{n-1}T_n(\iota_1\psi)$ is a graded Hopf algebra map from $T_B(M)$ to $T_{B'}(M')$.

(ii) $T_{B'}^c(\phi\pi_0, \psi\pi_1) := \phi\pi_0 + \sum_{n>0} T_n^c(\psi\pi_1)\Delta_{n-1}$ is a graded Hopf algebra map from $T_B^c(M)$ to $T_{B'}^c(M')$.

Proof. (i) Since ϕ is a Hopf algebra map, $\iota_0\phi : B \rightarrow T_{B'}(M')$ is an algebra map. Since ψ is a B -bimodule map from M to ${}_\phi M'_\phi$, $\iota_1\psi$ is a B -bimodule map from M to ${}_{\iota_0\phi}T_{B'}(M')_{\iota_0\phi}$. It follows that $T_B(\iota_0\phi, \iota_1\psi)$ is an algebra homomorphism from $T_B(M)$ to $T_{B'}(M')$. Clearly, $T_B(\iota_0\phi, \iota_1\psi)$ preserves the gradation of $T_B(M)$ and $T_{B'}(M')$.

Let $\Phi := T_B(\iota_0\phi, \iota_1\psi)$. Then both $\Delta_{T_{B'}(M')}\Phi$ and $(\Phi \otimes \Phi)\Delta_{T_B(M)}$ are graded algebra maps from $T_B(M)$ to $T_{B'}(M') \otimes T_{B'}(M')$. Now we show that $\Delta_{T_{B'}(M')}\Phi = (\Phi \otimes \Phi)\Delta_{T_B(M)}$. By the universal property of $T_B(M)$, we only have to prove that $\Delta_{T_{B'}(M')}\Phi\iota_n = (\Phi \otimes \Phi)\Delta_{T_B(M)}\iota_n$ for $n = 1, 2$. Since ϕ is a coalgebra map, one can see that $\Delta_{T_{B'}(M')}\Phi\iota_0 = (\Phi \otimes \Phi)\Delta_{T_B(M)}\iota_0$. Note that ψ is a B' -bicomodule map from ${}^\phi M^\phi$ to M' . Hence by the

proof of Lemma 1.4 we have

$$\begin{aligned}
\Delta_{T_{B'}(M')} \Phi \iota_1 &= \Delta_{T_{B'}(M')} \iota_1 \psi \\
&= (\iota_0 \otimes \iota_1) \delta_{M'}^- \psi + (\iota_1 \otimes \iota_0) \delta_{M'}^+ \psi \\
&= (\iota_0 \otimes \iota_1) (\phi \otimes \psi) \delta_M^- + (\iota_1 \otimes \iota_0) (\psi \otimes \phi) \delta_M^+ \\
&= (\iota_0 \phi \otimes \iota_1 \psi) \delta_M^- + (\iota_1 \psi \otimes \iota_0 \phi) \delta_M^+ \\
&= (\Phi \otimes \Phi) (\iota_0 \otimes \iota_1) \delta_M^- + (\Phi \otimes \Phi) (\iota_1 \otimes \iota_0) \delta_M^+ \\
&= (\Phi \otimes \Phi) [(\iota_0 \otimes \iota_1) \delta_M^- + (\iota_1 \otimes \iota_0) \delta_M^+] \\
&= (\Phi \otimes \Phi) \Delta_{T_B(M)} \iota_1.
\end{aligned}$$

Furthermore, it is easy to see that $\varepsilon_{T_{B'}(M')} \Phi = \varepsilon_{T_B(M)}$. It follows that Φ is a coalgebra map, and hence a Hopf algebra map since any bialgebra map between two Hopf algebras is a Hopf algebra map.

(ii) Since ϕ is a Hopf algebra map, $\phi \pi_0 : T_B^c(M) \rightarrow B'$ is a coalgebra map. Since ψ is a B' -bicomodule map from ${}^\phi M^\phi$ to M' , $\psi \pi_1$ is a B' -bicomodule map from ${}^{\phi \pi_0} T_B^c(M) {}^{\phi \pi_0}$ to M' . We also have $\text{corad}(T_B^c(M)) \subseteq \iota_0(B)$, and hence $(\psi \pi_1)(\text{corad}(T_B^c(M))) = 0$. It follows that $T_{B'}^c(\phi \pi_0, \psi \pi_1)$ is a coalgebra map from $T_B^c(M)$ to $T_{B'}^c(M')$. Clearly, $T_{B'}^c(\phi \pi_0, \psi \pi_1)$ is a graded map.

Put $\Psi := T_{B'}^c(\phi \pi_0, \psi \pi_1)$. Then both $\Psi \mu_{T_B^c(M)}$ and $\mu_{T_{B'}^c(M')}(\Psi \otimes \Psi)$ are graded coalgebra maps from $T_B^c(M) \otimes T_B^c(M)$ to $T_{B'}^c(M')$. Since $T_B^c(M) \otimes T_B^c(M)$ is a graded coalgebra, $\text{corad}(T_B^c(M) \otimes T_B^c(M)) \subseteq (T_B^c(M) \otimes T_B^c(M))_0 = \iota_0(B) \otimes \iota_0(B)$. It follows that $(\pi_1 \Psi \mu_{T_B^c(M)})(\text{corad}(T_B^c(M) \otimes T_B^c(M))) = 0$ and $(\pi_1 \mu_{T_{B'}^c(M')}(\Psi \otimes \Psi))(\text{corad}(T_B^c(M) \otimes T_B^c(M))) = 0$. Thus by the universal property of $T_{B'}^c(M')$, in order to prove $\Psi \mu_{T_B^c(M)} = \mu_{T_{B'}^c(M')}(\Psi \otimes \Psi)$, we only need to show $\pi_n \Psi \mu_{T_B^c(M)} = \pi_n \mu_{T_{B'}^c(M')}(\Psi \otimes \Psi)$ for $n = 1, 2$. However, this follows from a straightforward computation dual to Part (i). Furthermore, one can see $\Psi(1) = 1$. Hence Ψ is an algebra map, and so a Hopf algebra map. \square

Lemma 1.6 *Let B and B' be two Hopf algebras. Let M and M' be B -Hopf bimodule and B' -Hopf bimodule, respectively. Then the following statements are equivalent:*

- (i) *There exists a Hopf algebra isomorphism $\phi : B \rightarrow B'$ such that $M \cong {}_\phi^{\phi^{-1}} M' {}_\phi^{\phi^{-1}}$ as B -Hopf bimodules.*
- (ii) *$T_B(M)$ and $T_{B'}(M')$ are isomorphic as graded Hopf algebras.*
- (iii) *$T_B^c(M)$ and $T_{B'}^c(M')$ are isomorphic as graded Hopf algebras.*
- (iv) *$B[M]$ and $B'[M']$ are isomorphic as graded Hopf algebras.*

Proof. (i) \Rightarrow (ii) Assume that $\phi : B \rightarrow B'$ is a Hopf algebra isomorphism and $\psi : M \rightarrow {}_\phi^{\phi^{-1}} M' {}_\phi^{\phi^{-1}}$ is a B -Hopf bimodule isomorphism. Then $\phi^{-1} : B' \rightarrow B$ is also a Hopf algebra isomorphism and $\psi^{-1} : M' \rightarrow {}_{\phi^{-1}} M {}_{\phi^{-1}}$ is a B' -Hopf bimodule isomorphism. By Lemma 1.5(i), we have two graded Hopf algebra homomorphisms $T_B(\iota_0 \phi, \iota_1 \psi) : T_B(M) \rightarrow T_{B'}(M')$

and $T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1}) : T_{B'}(M') \rightarrow T_B(M)$. Then we have

$$\begin{aligned} T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})T_B(\iota_0\phi, \iota_1\psi)\iota_0 &= T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})\iota_0\phi \\ &= \iota_0\phi^{-1}\phi = \iota_0 \end{aligned}$$

and

$$\begin{aligned} T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})T_B(\iota_0\phi, \iota_1\psi)\iota_1 &= T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})\iota_1\psi \\ &= \iota_1\psi^{-1}\psi = \iota_1. \end{aligned}$$

It follows from the universal property of $T_B(M)$ that $T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})T_B(\iota_0\phi, \iota_1\psi) = \text{id}_{T_B(M)}$. Similarly, one can show that $T_B(\iota_0\phi, \iota_1\psi)T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1}) = \text{id}_{T_{B'}(M')}$. Hence $T_B(M) \cong T_{B'}(M')$ as graded Hopf algebras.

(i) \Rightarrow (iii) It is similar to the proof above.

(ii) \Rightarrow (i) Assume that $f : T_B(M) \rightarrow T_{B'}(M')$ is a graded Hopf algebra isomorphism. Put $\phi := \pi_0 f \iota_0$ and $\psi := \pi_1 f \iota_1$. Then it is easy to check that ϕ is a Hopf algebra isomorphism from B to B' and ψ is a B -Hopf bimodule isomorphism from M to ${}_{\phi}^{\phi^{-1}}M'_{\phi}^{\phi^{-1}}$.

(iv) \Rightarrow (i) Assume that φ is a graded Hopf algebra isomorphism from $B[M]$ to $B'[M']$. Let $\phi := \pi_0 \varphi \iota_0$ and $\psi := \pi_1 \varphi \iota_1$. Obviously, ϕ is a Hopf algebra isomorphism from B to B' and ψ is a bijection from M to M' . Now we show that ψ is a B -Hopf bimodule isomorphism from M to ${}_{\phi}^{\phi^{-1}}M'_{\phi}^{\phi^{-1}}$. Assume that $\alpha^-, \alpha^+, \delta^-$ and δ^+ are the module operations and comodule operations of M ; $\alpha'^-, \alpha'^+, \delta'^-$ and δ'^+ are the module operations and comodule operations of M' .

See that

$$\begin{aligned} \alpha'^-(\phi \otimes \psi) &= \pi_1 \mu(\iota_0 \pi_0 \varphi \iota_0 \otimes \iota_1 \pi_1 \varphi \iota_1) \\ &= \pi_1 \mu(\varphi \iota_0 \otimes \varphi \iota_1) \quad \text{and} \\ \psi \alpha^- &= \pi_1 \varphi \iota_1 \pi_1 \mu(\iota_0 \otimes \iota_1) \\ &= \pi_1 \varphi \mu(\iota_0 \otimes \iota_1) \\ &= \pi_1 \mu(\varphi \iota_0 \otimes \varphi \iota_1). \end{aligned}$$

Thus ψ is a left B -module homomorphism from M to ${}_{\phi}M'$. Similarly, we have that ψ is a right B -module homomorphism from M to M'_{ϕ} . See that

$$\begin{aligned} (\phi^{-1} \otimes \text{id}_{M'})\delta'^-\psi &= (\phi^{-1} \otimes \text{id}_{M'})(\pi_0 \otimes \pi_1)\Delta\iota_1\pi_1\varphi\iota_1 \\ &= (\phi^{-1}\pi_0 \otimes \pi_1)\Delta\varphi\iota_1 \quad \text{and} \\ (\text{id}_B \otimes \psi)\delta^- &= (\phi^{-1}\pi_0\varphi\iota_0\pi_0 \otimes \pi_1\varphi\iota_0\pi_0)\Delta\iota_1 \\ &= (\phi^{-1}\pi_0 \otimes \pi_1)\Delta\varphi\iota_1. \end{aligned}$$

Thus ψ is a left B -comodule homomorphism from M to ${}^{\phi^{-1}}M'$. Similarly, we have that ψ is a right B -comodule homomorphism from M to $M'^{\phi^{-1}}$.

(iii) \Rightarrow (iv) Obvious. \square

Let $Q = (G, Q_1, s, t)$ be a quiver of a group G . Then kQ_1 becomes a kG -bicomodule under the natural comodule structures:

$$\delta^-(a) = t(a) \otimes a, \quad \delta^+(a) = a \otimes s(a), \quad a \in Q_1, \quad (15)$$

called an *arrow comodule*, written as kQ_1^c or kQ_1 . In this case, the path coalgebra kQ^c is exactly isomorphic to the cotensor coalgebra $T_{kG}^c(kQ_1^c)$ over kG in a natural way (see [8] and [9]). We will regard $kQ^c = T_{kG}^c(kQ_1^c)$ in the following. Moreover, kQ_1 becomes a $(kG)^*$ -bimodule with the module structures defined by

$$p \cdot a := \langle p, t(a) \rangle a, \quad a \cdot p := \langle p, s(a) \rangle a, \quad p \in (kG)^*, a \in Q_1, \quad (16)$$

written as kQ_1^a , called an *arrow module*. Therefore, we have a tensor algebra $T_{(kG)^*}(kQ_1^a)$. Note that the tensor algebra $T_{(kG)^*}(kQ_1^a)$ of kQ_1^a over $(kG)^*$ is exactly isomorphic to the path algebra kQ^a . We will regard $kQ^a = T_{(kG)^*}(kQ_1^a)$ in the following.

Assume that Q is a finite quiver on finite group G . Let $\xi_{kQ_1^a}$ denote the linear map from kQ_1^a to $(kQ_1^a)^*$ by sending a to a^* for any $a \in Q_1$ and $\xi_{kQ_1^c}$ denote the linear map from kQ_1^c to $(kQ_1^a)^*$ by sending a to a^* for any $a \in Q_1$. Here $\{a^* \mid a^* \in (kQ_1^a)^*\}$ is the dual basis of $\{a \mid a \in Q_1\}$.

Lemma 1.7 *Assume that Q is a finite Hopf quiver on finite group G . Then*

(i) *If $(M, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a finite dimensional B -Hopf bimodule and B is a finite dimensional Hopf algebra, then $(M^*, \delta^{*-}, \delta^{*+}, \alpha^{*-}, \alpha^{*+})$ is a B^* -Hopf bimodule.*

(ii) *If $(kQ_1^c, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a kG -Hopf bimodule, then there exist unique left $(kG)^*$ -comodule operation $\delta_{kQ_1^a}^-$ and right $(kG)^*$ -comodule $\delta_{kQ_1^a}^+$ such that $(kQ_1^a, \alpha_{kQ_1^a}^-, \alpha_{kQ_1^a}^+, \delta_{kQ_1^a}^-, \delta_{kQ_1^a}^+)$ becomes a $(kG)^*$ -Hopf bimodule and $\xi_{kQ_1^a}$ becomes a $(kG)^*$ -Hopf bimodule isomorphism from $(kQ_1^a, \alpha_{kQ_1^a}^-, \alpha_{kQ_1^a}^+, \delta_{kQ_1^a}^-, \delta_{kQ_1^a}^+)$ to $((kQ_1^a)^*, \delta^{*-}, \delta^{*+}, \alpha^{*-}, \alpha^{*+})$.*

(iii) *If $(kQ_1^a, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a kG^* -Hopf bimodule, then there exist unique left kG -module operation $\alpha_{kQ_1^c}^-$ and right kG -module $\alpha_{kQ_1^c}^+$ such that $(kQ_1^c, \alpha_{kQ_1^c}^-, \alpha_{kQ_1^c}^+, \delta_{kQ_1^c}^-, \delta_{kQ_1^c}^+)$ become a kG -Hopf bimodule and $\xi_{kQ_1^c}$ becomes a kG -Hopf bimodule isomorphism from $(kQ_1^c, \alpha_{kQ_1^c}^-, \alpha_{kQ_1^c}^+, \delta_{kQ_1^c}^-, \delta_{kQ_1^c}^+)$ to $((kQ_1^a)^*, \delta^{*-}, \delta^{*+}, \alpha^{*-}, \alpha^{*+})$.*

(iv) *$\xi_{kQ_1^a}$ is a $(kG)^*$ -Hopf bimodule isomorphism from $(kQ_1^a, \alpha_{kQ_1^a}^-, \alpha_{kQ_1^a}^+, \delta_{kQ_1^a}^-, \delta_{kQ_1^a}^+)$ to $((kQ_1^a)^*, \delta_{kQ_1^a}^{*-}, \delta_{kQ_1^a}^{*+}, \alpha_{kQ_1^a}^{*-}, \alpha_{kQ_1^a}^{*+})$ if and only if $\xi_{kQ_1^c}$ becomes a kG -Hopf bimodule isomorphism from $(kQ_1^c, \alpha_{kQ_1^c}^-, \alpha_{kQ_1^c}^+, \delta_{kQ_1^c}^-, \delta_{kQ_1^c}^+)$ to $((kQ_1^a)^*, \delta_{kQ_1^a}^{*-}, \delta_{kQ_1^a}^{*+}, \alpha_{kQ_1^a}^{*-}, \alpha_{kQ_1^a}^{*+})$.*

Proof. It is easy to check (i)–(iii). Now we show (iv). Let $A := kQ_1^a$, $B := kQ_1^c$. Let σ_B denote the canonical linear isomorphism from B to B^{**} by sending b to b^{**} for any $b \in B$, where $\langle b^{**}, f \rangle = \langle f, b \rangle$ for any $f \in B^*$. If $A \stackrel{\xi_A}{\cong} B^*$ as $(kG)^*$ -Hopf bimodules, then

$B \xrightarrow{\sigma_B} B^{**} \xrightarrow{(\xi_A)^*} A^*$ as kG -Hopf bimodules. It is easy to check $\xi_B = (\xi_A)^* \sigma_B$. Therefore ξ_B is a kG -Hopf bimodule isomorphism. Conversely, if $B \xrightarrow{\xi_B} A^*$ as kG -Hopf bimodules, we can similarly show that $A \cong B^*$ as $(kG)^*$ -Hopf bimodules. \square

The following theorem can be found in [10, 9]. We list the result here for the completeness and give a slightly different proof.

Theorem 3 (see [9, Theorem 3.3] and [10, Theorem 3.1]) *Let Q be a quiver over group G . Then the following statements are equivalent:*

- (i) Q is a Hopf quiver.
- (ii) Arrow comodule kQ_1^c admits a kG -Hopf bimodule structure.
- (iii) Arrow module kQ_1^a admits a $(kG)^*$ -Hopf bimodule structure.

Proof. (i) \Rightarrow (ii). Assume that (Q, G, r) is a Hopf quiver, where r is a ramification of G . For any $C \in \mathcal{K}(G)$, let $M(C)$ be a trivial right $kZ_{u(C)}$ -module with $\dim M(C) = r_C$. That is, $m \cdot h = m$ for any $m \in M(C)$ and $h \in Z_{u(C)}$. From the proof of Theorem 2, $M := \bigoplus_{x,y \in G} (x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta)$ is a kG -Hopf bimodule. Let $\{\xi_C^{(i)} \mid i \in I_C(r)\}$ be a basis of $M(C)$ over k . Let $\phi : kQ_1 \rightarrow M$ be a k -linear map defined by

$$\phi(a_{x,y}^{(i)}) = x \otimes \xi_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta,$$

where $x, y \in G$ with $x^{-1}y = g_\theta u(C)g_\theta^{-1}$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. It is easy to see that ϕ is a k -linear isomorphism and a kG -bicomodule homomorphism from kQ_1^c to M . It follows that the arrow comodule kQ_1^c admits a kG -Hopf bimodule structure.

(ii) \Rightarrow (i). Assume that the arrow comodule kQ_1^c admits a kG -Hopf bimodule structure. Then by Theorem 2 and its proof, there is a vector space isomorphism ${}^y(kQ_1^c)^x \cong x \otimes {}^{u(C)}(kQ_1^c)^1 \otimes_{kZ_{u(C)}} g_\theta$ for any $x, y \in G$ with $y = xg_\theta u(C)g_\theta^{-1}$. Hence for any $x, y \in G$ with $y = xg_\theta u(C)g_\theta^{-1}$, we have $\dim {}^y(kQ_1^c)^x = \dim {}^{u(C)}(kQ_1^c)^1$. This shows that Q is a Hopf quiver.

(ii) \Leftrightarrow (iii) It follows from Lemma 1.7. \square

Assume that Q is a Hopf quiver. It follows from Theorem 3 that there exist a left kG -module structure α^- and a right kG -module structure α^+ on arrow comodule $(kQ_1^c, \delta^-, \delta^+)$ such that $(kQ_1^c, \alpha^-, \alpha^+, \delta^-, \delta^+)$ becomes a kG -Hopf bimodule, called a co-arrow Hopf bimodule, written $(kQ_1^c, \alpha^-, \alpha^+)$ in short. We obtain two graded Hopf algebras $T_{kG}(kQ_1^c)$ and $T_{kG}^c(kQ_1^c)$, called semi-path Hopf algebra and co-path Hopf algebra, written $kQ^s(\alpha^-, \alpha^+)$ and $kQ^c(\alpha^-, \alpha^+)$, respectively. Let $kG[kQ_1^c, \alpha_1, \alpha^+]$ be the subalgebra of kQ^c generated by kG and kQ_1^c , which is a Hopf algebra of type one (see [20, section 2.2]), called a *one-type-co-path Hopf algebra*. Dually, it follows from Theorem 3 that there exist a left $(kG)^*$ -comodule structure δ^- and a right $(kG)^*$ -comodule structure δ^+ on arrow module $(kQ_1^a, \alpha^-, \alpha^+)$ such that $(kQ_1^a, \alpha^-, \alpha^+, \delta^-, \delta^+)$ becomes a

$(kG)^*$ -Hopf bimodule, called a arrow Hopf bimodule, written $(kQ_1^a, \delta^-, \delta^+)$ in short. We obtain two graded Hopf algebras $T_{(kG)^*}(kQ_1^a)$ and $T_{(kG)^*}^c(kQ_1^a)$, called path Hopf algebra and semi-co-path Hopf algebra, written $kQ^a(\delta^-, \delta^+)$ and $kQ^{sc}(\delta^-, \delta^+)$, respectively. Let $(kG)^*[kQ_1^a, \alpha_1, \alpha^+]$ be the subalgebra of kQ^{sc} generated by $(kG)^*$ and kQ_1^a , called *one-type-path Hopf algebra*. We call the six graded Hopf algebras the quiver Hopf algebras. We usually omit the (co)module operations when we write these quiver Hopf algebras.

If $\xi_{kQ_1^a}$ or $\xi_{kQ_1^c}$ is a Hopf bimodule isomorphism, then, by Lemma 1.6 and Lemma 1.7, $T_{(kG)^*}(\iota_0, \iota_1 \xi_{kQ_1^a})$ and $T_{kG}^c(\pi_0, \xi_{kQ_1^c} \pi_1)$ are graded Hopf algebra isomorphisms from $T_{(kG)^*}(kQ_1^a)$ to $T_{(kG)^*}((kQ_1^c)^*)$ and from $T_{kG}^c(kQ_1^c)$ to $T_{kG}^c((kQ_1^a)^*)$, respectively; $T_{(kG)^*}^c(\pi_0, \xi_{kQ_1^a} \pi_1)$ and $T_{kG}(\iota_0, \iota_1 \xi_{kQ_1^c})$ are graded Hopf algebra isomorphisms from $T_{(kG)^*}^c(kQ_1^a)$ to $T_{(kG)^*}^c((kQ_1^c)^*)$ and from $T_{kG}(kQ_1^c)$ to $T_{kG}((kQ_1^a)^*)$, respectively. In this case, (kQ_1^a, kQ_1^c) , (kQ^a, kQ^c) and (kQ^s, kQ^{sc}) are said to be arrow dual pairings.

If $(kQ_1^c, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a PM kG -Hopf bimodule, and (kQ_1^c, kQ_1^a) is an arrow pairing, then $(kQ_1^a, \alpha^{-*}, \alpha^{+*})$ is called a PM $(kG)^*$ -Hopf bimodule and six quiver Hopf algebras induced by kQ_1^c and kQ_1^a are called PM quiver Hopf algebras.

Now we are going to describe the structure of all PM kG -Hopf bimodules and the corresponding graded Hopf algebras.

Definition 1.8 $(G, r, \vec{\chi}, u)$ is called a *ramification system with characters* (or RSC in short), if r is a ramification of G , u is a map from $\mathcal{K}(G)$ to G with $u(C) \in C$ for any $C \in \mathcal{K}(G)$, and $\vec{\chi} = \{\chi_C^{(i)}\}_{i \in I_C(r), C \in \mathcal{K}_r(G)} \in \prod_{C \in \mathcal{K}_r(G)} (\widehat{Z_{u(C)}})^{r_C}$ with $\chi_C^{(i)} \in \widehat{Z_{u(C)}}$ for any $i \in I_C(r), C \in \mathcal{K}_r(G)$.

$RSC(G, r, \vec{\chi}, u)$ and $RSC(G', r', \vec{\chi}', u')$ are said to be *isomorphic* if the following conditions are satisfied:

- There exists a group isomorphism $\phi : G \rightarrow G'$.
- For any $C \in \mathcal{K}(G)$, there exists an element $h_C \in G$ such that $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$.
- For any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}(\phi_C^{(i)}) (\phi(h_C^{-1}hh_C)) = \chi_C^{(i)}(h)$ for all $h \in Z_{u(C)}$ and $i \in I_C(r)$.

Obviously, if $u(C) = u'(C)$ for any $C \in \mathcal{K}_r(G)$, then $RSC(G, r, \vec{\chi}, u) \cong RSC(G, r, \vec{\chi}, u')$. Indeed, for any $C \in \mathcal{K}(G)$, there exists $h_C \in G$ such that $u'(C) = h_C^{-1}u(C)h_C$ and $h_C = 1$ when $C \in \mathcal{K}_r(G)$. Let $\phi = id_G$ and $\phi_C = id_{I_C(r)}$. It is straightforward to check the conditions in Definition 1.8 hold.

Proposition 1.9 If N is a PM kG -Hopf bimodule, then there exist an RSC $(G, \vec{\chi}, r, u)$, a Hopf quiver (Q, G, r) and a kG -Hopf bimodule $(kQ_1, \alpha^-, \alpha^+)$ with

$$\alpha^-(h \otimes a_{y,x}^{(i)}) = h \cdot a_{y,x}^{(i)} = a_{hy, hx}^{(i)}, \quad \alpha^+(a_{y,x}^{(i)} \otimes h) = a_{y,x}^{(i)} \cdot h = \chi_C^{(i)}(\zeta_\theta(h)) a_{yh, xh}^{(i)}$$

where $x, y, h \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, ζ_θ is given by (3), $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$, such that $N \cong (kQ_1, \alpha^-, \alpha^+)$ as kG -Hopf bimodule.

Proof. Since N is a PM kG -Hopf bimodule, there exists an object $\prod_{C \in \mathcal{K}(G)} M(C)$ in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ such that $M(C)$ is a pointed $kZ_{u(C)}$ -module for any $C \in \mathcal{K}(G)$ and $N \cong V(\{M(C)\}_{C \in \mathcal{K}(G)}) = \bigoplus_{x, y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ as kG -Hopf bimodules. Let $r = \sum_{C \in \mathcal{K}(G)} r_C C$ with $r_C = \dim M(C)$ for any $C \in \mathcal{K}(G)$. Since $(M(C), \alpha_C)$ is a pointed $kZ_{u(C)}$ -module, there exist a k -basis $\{x_C^{(i)} \mid i \in I_C(r)\}$ in $M(C)$ and a family of characters $\{\chi_C^{(i)} \in \widehat{Z_{u(C)}} \mid i \in I_C(r)\}$ such that $\alpha_C(x_C^{(i)} \otimes h) = x_C^{(i)} \triangleleft h = \chi_C^{(i)}(h)x_C^{(i)}$ for any $i \in I_C(r)$ and $h \in Z_{u(C)}$.

We have to show that $(kQ_1, \alpha^-, \alpha^+)$ is isomorphic to $\bigoplus_{x, y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ as kG -Hopf bimodules. Observe that there is a canonical kG -bicomodule isomorphism $\varphi : kQ_1 \rightarrow \bigoplus_{x, y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ given by

$$\varphi(a_{y,x}^{(i)}) = x \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta \quad (17)$$

where $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. See that

$$\begin{aligned} \varphi(\alpha^-(h \otimes a_{y,x}^{(i)})) &= \varphi(a_{hy, hx}^{(i)}) \\ &= hx \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta \\ &= h \cdot (x \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta) \quad (\text{see (6)}) \\ &= h \cdot \varphi(a_{y,x}^{(i)}) \end{aligned}$$

and

$$\begin{aligned} \varphi(\alpha^+(a_{y,x}^{(i)} \otimes h)) &= \chi_C^{(i)}(\zeta_\theta(h))\varphi(a_{yh, xh}^{(i)}) \\ &= \chi_C^{(i)}(\zeta_\theta(h))xh \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta \\ &\quad (\text{since } hg_\theta^{-1} = g_\theta^{-1}\zeta_\theta(h) \text{ and } yh = xhg_\theta^{-1}u(C)g_\theta) \\ &= (x \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta) \cdot h \quad (\text{see (6)}) \\ &= \varphi(a_{y,x}^{(i)}) \cdot h, \end{aligned}$$

where $x, y, h \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, ζ_θ is given by (3), $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. Consequently, φ is a kG -Hopf bimodule isomorphism. \square

Let (kQ_1, α^χ) denote $(kQ_1, \alpha^-, \alpha^+)$ in Lemma 1.9. Furthermore, if (kQ_1, kQ_1^a) is an arrow dual pairing, then we denote kQ_1^a by (kQ_1^a, α^χ) . We obtain six quiver Hopf algebras $kQ^c(\alpha^\chi)$, $kQ^s(\alpha^\chi)$, $kG[kQ_1, \alpha^\chi]$, $kQ^a(\alpha^\chi)$, $kQ^{sc}(\alpha^\chi)$, $(kG)^*[kQ_1^a, \alpha^\chi]$.

Definition 1.10 $(M, [\alpha^\chi])$ denotes a kG -Hopf bimodule, which is isomorphic to (kQ_1^c, α^χ) as kG -Hopf bimodule.

By Definition 1.10, every $(M, [\alpha^\lambda])$ is a PM kG -Hopf bimodule. Conversely, it follows from Proposition 1.9 that, for every PM kG -Hopf bimodule N , there exists a $RSC(G, r, \vec{\chi}, u)$ such that $N \cong (kQ_1^c, \alpha^\lambda)$, i.e. N can be written as $(N, [\alpha^\lambda])$.

Proposition 1.11 $(kG)^*$ -coactions on the $(kG)^*$ -Hopf bimodule (kQ_1^a, α^λ) are given by

$$\delta^-(a_{y,x}^{(i)}) = \sum_{h \in G} p_h \otimes a_{h^{-1}y, h^{-1}x}^{(i)}, \quad \delta^+(a_{y,x}^{(i)}) = \sum_{h \in G} \chi_C^{(i)}(\zeta_\theta(h^{-1})^{-1}) a_{yh^{-1}, xh^{-1}}^{(i)} \otimes p_h$$

where $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, ζ_θ is given by (3), $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$ and $p_h \in (kG)^*$ is defined by $p_h(g) = \delta_{h,g}$ for all $g, h \in G$.

Proof. Let $(kQ_1^c, \alpha^\lambda) = (kQ_1, \alpha^-, \alpha^+)$ and $(kQ_1^a, \alpha^\lambda) = (kQ_1^a, \delta^-, \delta^+)$. Since $((kQ_1, \alpha^\lambda), (kQ_1^a, \alpha^\lambda))$ is an arrow dual pairing, $\xi_{kQ_1^a}$ is a $(kG)^*$ -Hopf bimodule isomorphism from $(kQ_1^a, \delta^-, \delta^+)$ to $((kQ_1^c)^*, \alpha^{-*}, \alpha^{+*})$ (see Lemma 1.7). Now let $h, x, y, v, w \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, where $C \in \mathcal{K}_r(G)$, $\theta \in \Theta_C$. By Proposition 1.9, we have

$$\begin{aligned} \langle \delta^-((a_{y,x}^{(i)})^*), h \otimes a_{w,v}^{(j)} \rangle &= \langle (a_{y,x}^{(i)})^*, h \cdot a_{w,v}^{(j)} \rangle \\ &= \langle (a_{y,x}^{(i)})^*, a_{hw, hv}^{(j)} \rangle \\ &= \delta_{x, hv} \delta_{y, hw} \delta_{i,j} \\ &= \delta_{h^{-1}x, v} \delta_{h^{-1}y, w} \delta_{i,j}. \end{aligned}$$

This shows that $\delta^-((a_{y,x}^{(i)})^*) = \sum_{h \in G} p_h \otimes (a_{h^{-1}y, h^{-1}x}^{(i)})^*$. Hence the left $(kG)^*$ -coaction of the $(kG)^*$ -Hopf bimodule $(kQ_1^a; \alpha^\lambda)$ is given by

$$\delta^-(a_{y,x}^{(i)}) = \sum_{h \in G} p_h \otimes a_{h^{-1}y, h^{-1}x}^{(i)}.$$

In order to show the second equation, we only need to show that

$$\langle \delta^+((a_{y,x}^{(i)})^*), a_{wv}^{(j)} \otimes h \rangle = \sum_{g \in G} \langle \chi_C^{(i)}(\zeta_\theta(g^{-1})^{-1})(a_{yg^{-1}, xg^{-1}}^{(i)})^* \otimes p_g, a_{wv}^{(j)} \otimes h \rangle \quad (18)$$

for any $h, v, w \in G, D \in \mathcal{K}(G), j \in I_D(r)$ and $v^{-1}w \in D$, since $\xi_{kQ_1^a}$ is a $(kG)^*$ -Hopf bimodule isomorphism. If $v \neq xh^{-1}$ or $w \neq yh^{-1}$, then there exists $\beta \in k$ such that $a_{w,v}^{(j)} \cdot h = \beta a_{wh, vh}^{(j)}$ by Proposition 1.9. By simple computation, both the right hand side and the left hand side of (18) are zero.

Now assume $v = xh^{-1}$ and $w = yh^{-1}$. Assume $g_\theta h^{-1} = \zeta_\theta(h^{-1})g_{\theta'}$. Thus $v^{-1}w = hg_\theta^{-1}u(C)g_\theta h^{-1} = g_{\theta'}^{-1}\zeta_\theta(h^{-1})^{-1}u(C)\zeta_\theta(h^{-1})g_{\theta'} = g_{\theta'}^{-1}u(C)g_{\theta'}$. Since $g_{\theta'}h = \zeta_{\theta'}(h^{-1})^{-1}g_\theta$,

$\zeta_{\theta'}(h) = \zeta_{\theta}(h^{-1})^{-1}$. Now we have

$$\begin{aligned}
\text{the left hand side of (18)} &= \langle \delta^+((a_{y,x}^{(i)})^*), a_{w,v}^{(j)} \otimes h \rangle \\
&= \langle (a_{y,x}^{(i)})^*, a_{w,v}^{(j)} \cdot h \rangle \\
&= \chi_C^{(j)}(\zeta_{\theta'}(h)) \langle (a_{y,x}^{(i)})^*, a_{wh,vh}^{(j)} \rangle \\
&= \chi_C^{(j)}(\zeta_{\theta'}(h)) \delta_{i,j} \\
&= \chi_C^{(i)}(\zeta_{\theta}(h^{-1})^{-1}) \delta_{i,j} \\
&= \text{the right hand side of (18)}. \quad \square
\end{aligned}$$

From Proposition 1.9, it seems that the right kG -action on (kQ_1, α^x) depends on the choice of the set $\{g_{\theta} \mid \theta \in \Theta_C\}$ of coset representatives of $Z_{u(C)}$ in G (see, Eq.(1)). The following lemma shows that (kQ_1, α^x) is, in fact, independent of the choice of the coset representative set $\{g_{\theta} \mid \theta \in \Theta_C\}$, up to kG -Hopf bimodule isomorphisms. For a while, we write $(kQ_1, \alpha^x) = (kQ_1, \alpha^x, \{g_{\theta}\})$ given before. Now let $\{h_{\theta} \in G \mid \theta \in \Theta_C\}$ be another coset representative set of $Z_{u(C)}$ in G for any $C \in \mathcal{K}(G)$. That is,

$$G = \bigcup_{\theta \in \Theta_C} Z_{u(C)} h_{\theta}. \quad (19)$$

Lemma 1.12 *With the above notations, $(kQ_1, \alpha^x, \{g_{\theta}\})$ and $(kQ_1, \alpha^x, \{h_{\theta}\})$ are isomorphic kG -Hopf bimodules.*

Proof. We may assume $Z_{u(C)} h_{\theta} = Z_{u(C)} g_{\theta}$ for any $C \in \mathcal{K}(G)$ and $\theta \in \Theta_C$. Then $g_{\theta} h_{\theta}^{-1} \in Z_{u(C)}$. Now let $x, y, h \in G$ with $x^{-1}y = g_{\theta}^{-1}u(C)g_{\theta}$. Then $x^{-1}y = h_{\theta}^{-1}(g_{\theta} h_{\theta}^{-1})^{-1}u(C)(g_{\theta} h_{\theta}^{-1})h_{\theta} = h_{\theta}^{-1}u(C)h_{\theta}$ and $h_{\theta}h = (h_{\theta}g_{\theta}^{-1})g_{\theta}h = (h_{\theta}g_{\theta}^{-1})\zeta_{\theta}(h)g_{\theta'} = (h_{\theta}g_{\theta}^{-1})\zeta_{\theta}(h)(g_{\theta'}h_{\theta'}^{-1})h_{\theta'}$, where $g_{\theta}h = \zeta_{\theta}(h)g_{\theta'}$. Hence from Proposition 1.9 we know that the right kG -action on $(kQ_1, \alpha^x, \{h_{\theta}\})$ is given by

$$\begin{aligned}
a_{y,x}^{(i)} \cdot h &= \chi_C^{(i)}((h_{\theta}g_{\theta}^{-1})\zeta_{\theta}(h)(g_{\theta'}h_{\theta'}^{-1}))a_{y_{h,x}h}^{(i)} \\
&= \chi_C^{(i)}((g_{\theta}h_{\theta}^{-1})^{-1})\chi_C^{(i)}(\zeta_{\theta}(h))\chi_C^{(i)}(g_{\theta'}h_{\theta'}^{-1})a_{y_{h,x}h}^{(i)}, \quad i \in I_C(r).
\end{aligned}$$

However, we also have

$$(xh)^{-1}(yh) = h^{-1}(x^{-1}y)h = h^{-1}g_{\theta}^{-1}u(C)g_{\theta}h = g_{\theta'}^{-1}u(C)g_{\theta'}.$$

It follows that the k -linear isomorphism $f : kQ_1 \rightarrow kQ_1$ given by

$$f(a_{y,x}^{(i)}) = \chi_C^{(i)}(g_{\theta}h_{\theta}^{-1})a_{y,x}^{(i)}$$

for any $x, y \in G$ with $x^{-1}y = g_{\theta}^{-1}u(C)g_{\theta}$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$, is a kG -Hopf bimodule isomorphism from $(kQ_1, \alpha^x, \{g_{\theta}\})$ to $(kQ_1, \alpha^x, \{h_{\theta}\})$. \square

Now we state one of our main results, which classifies the PM (co-)path Hopf algebras, PM semi-(co-)path Hopf algebras and PM one-type-path Hopf algebras.

Theorem 4 Let $(G, r, \vec{\chi}, u)$ and $(G', r', \vec{\chi}', u')$ are two RSC's. Then the following statements are equivalent:

(i) $RSC(G, r, \vec{\chi}, u) \cong RSC(G', r', \vec{\chi}', u')$. (ii) $(kG; (kQ_1^c, \alpha^x))$ and $(kG'; (kQ_1'^c, \alpha^{x'}))$ are pull-push Hopf bimodule isomorphism. (iii) $kQ^c(\alpha^x) \cong kQ'^c(\alpha^{x'})$. (iv) $kQ^s(\alpha^x) \cong kQ'^s(\alpha^{x'})$. (v) $kG[kQ_1, \alpha^x] \cong kG'[kQ_1', \alpha^{x'}]$. (vi) $kQ^a(\alpha^x) \cong kQ'^a(\alpha^{x'})$. (vii) $kQ^{sc}(\alpha^x) \cong kQ'^{sc}(\alpha^{x'})$. (viii) $(kG)^*[kQ_1^a, \alpha^x] \cong (kG')^*[kQ_1'^a, \alpha^{x'}]$. (ix) $(kG; (M, [\alpha^x]))$ and $(kG'; (M', [\alpha^{x'}]))$ are pull-push Hopf bimodule isomorphism. (x) $T_{kG}(M, [\alpha^x]) \cong T_{kG'}(M', [\alpha^{x'}])$. (xi) $T_{kG}^c(M, [\alpha^x]) \cong T_{kG'}^c(M', [\alpha^{x'}])$. (xii) $T_{(kG)^*}((M, [\alpha^x])^*) \cong T_{(kG')^*}((M', [\alpha^{x'}])^*)$. (xiii) $T_{(kG)^*}^c((M, [\alpha^x])^*) \cong T_{(kG')^*}^c((M', [\alpha^{x'}])^*)$. (xiv) $kG[(M, [\alpha^x])] \cong kG'[(M', [\alpha^{x'}])]$. (xv) $(kG)^*[(M^*, [\alpha^x])] \cong (kG')^*[(M'^*, [\alpha^{x'}])]$. Note that all isomorphisms above are as graded Hopf algebras but in (i), (ii) and (ix).

Proof. By Lemma 1.6 and Lemma 1.7, we only have to prove (i) \Leftrightarrow (ii) and (ii) \Leftrightarrow (ix).

(ii) \Rightarrow (ix). By Definition 1.10, $(M, [\alpha^x]) \cong (Q_1^c, \alpha^x)$ as kG -Hopf bimodules and $(M', [\alpha^{x'}]) \cong (Q_1'^c, \alpha^{x'})$ as kG' -Hopf bimodules. Thus $(kG; (M, [\alpha^x])) \cong (kG'; (M', [\alpha^{x'}]))$ as pull-push Hopf bimodules. Similarly we can show (ix) \Rightarrow (ii).

(i) \Rightarrow (ii). Assume that $RSC(G, r, \vec{\chi}, u) \cong RSC(G', r', \vec{\chi}', u')$. Then there exist a group isomorphism $\phi : G \rightarrow G'$, an element $h_C \in G$ such that $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$ for any $C \in \mathcal{K}(G)$ and a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}(\phi(h_C^{-1}hh_C)) = \chi_C^{(i)}(h)$ for any $h \in Z_{u(C)}$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. Then $\phi(h_C^{-1}Z_{u(C)}h_C) = Z_{u'(\phi(C))}$ and $\phi : kG \rightarrow kG'$ is a Hopf algebra isomorphism. Now let $G = \bigcup_{\theta \in \Theta_C} Z_{u(C)}g_\theta$ be given as in (2) for any $C \in \mathcal{K}(G)$, and assume that the kG -Hopf bimodule (kQ_1, α^x) is defined by using these coset decompositions. Then

$$G' = \bigcup_{\theta \in \Theta_C} Z_{u'(\phi(C))}(\phi(h_C^{-1}g_\theta h_C)) \quad (20)$$

is a coset decomposition of $Z_{u'(\phi(C))}$ in G' for any $\phi(C) \in \mathcal{K}(G')$. By Lemma 1.12, we may assume that the structure of the kG' -Hopf bimodule $(kQ_1', \alpha^{x'})$ is obtained by using these coset decompositions (20). Define a k -linear isomorphism $\psi : (kQ_1, \alpha^x) \rightarrow (kQ_1', \alpha^{x'})$ by

$$\psi(a_{y,x}^{(i)}) = \chi_C^{(i)}(\zeta_\theta(h_C^{-1}))a_{\phi(y), \phi(x)}^{(\phi_C(i))}$$

for any $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, and $i \in I_C(r)$, where $C \in \mathcal{K}_r(G)$ and $g_\theta h_C^{-1} = \zeta_\theta(h_C^{-1})g_\eta$ with $\zeta_\theta(h_C^{-1}) \in Z_{u(C)}$ and $\theta, \eta \in \Theta_C$. It is easy to see that ψ is a kG -bicomodule homomorphism from (kQ_1, α^x) to $\phi^{-1}(kQ_1', \alpha^{x'})\phi^{-1}$. Since $(hx)^{-1}(hy) = x^{-1}y$ for any $x, y, h \in G$, it follows from Proposition 1.9 that ψ is also a left kG -module homomorphism from (kQ_1, α^x) to $\phi^{-1}(kQ_1', \alpha^{x'})\phi^{-1}$.

Now let $x, y, h \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, $C \in \mathcal{K}_r(G)$ and $\theta \in \Theta_C$. Assume that $g_\theta h_C^{-1} = \zeta_\theta(h_C^{-1})g_\eta$, $g_\theta h = \zeta_\theta(h)g_{\theta'}$, $g_\eta(h_C h h_C^{-1}) = \zeta_\eta(h_C h h_C^{-1})g_{\eta'}$ and $g_{\theta'} h_C^{-1} = \zeta_{\theta'}(h_C^{-1})g_{\theta''}$

with $\zeta_\theta(h_C^{-1}), \zeta_\theta(h), \zeta_\eta(h_C h h_C^{-1}), \zeta_{\theta'}(h_C^{-1}) \in Z_{u(C)}$ and $\eta, \theta', \eta', \theta'' \in \Theta_C$. Then we have

$$\begin{aligned} g_\theta h h_C^{-1} &= \zeta_\theta(h) g_{\theta'} h_C^{-1} \\ &= \zeta_\theta(h) \zeta_{\theta'}(h_C^{-1}) g_{\theta''} \end{aligned}$$

and

$$\begin{aligned} g_\theta h h_C^{-1} &= (g_\theta h_C^{-1})(h_C h h_C^{-1}) \\ &= \zeta_\theta(h_C^{-1}) g_\eta(h_C h h_C^{-1}) \\ &= \zeta_\theta(h_C^{-1}) \zeta_\eta(h_C h h_C^{-1}) g_{\eta'}. \end{aligned}$$

It follows that $\theta'' = \eta'$ and

$$\zeta_\theta(h) \zeta_{\theta'}(h_C^{-1}) = \zeta_\theta(h_C^{-1}) \zeta_\eta(h_C h h_C^{-1}). \quad (21)$$

By Proposition 1.9, $a_{y,x}^{(i)} \cdot h = \chi_C^{(i)}(\zeta_\theta(h)) a_{yh,xh}^{(i)}$ for any $i \in I_C(r)$. Moreover, we have $(xh)^{-1}(yh) = h^{-1}g_\theta^{-1}u(C)g_\theta h = g_{\theta'}^{-1}u(C)g_{\theta'}$. This implies that

$$\psi(a_{y,x}^{(i)} \cdot h) = \chi_C^{(i)}(\zeta_\theta(h)) \psi(a_{yh,xh}^{(i)}) = \chi_C^{(i)}(\zeta_\theta(h)) \chi_C^{(i)}(\zeta_{\theta'}(h_C^{-1})) a_{\phi(yh), \phi(xh)}^{(\phi_C(i))}.$$

On the other hand, we have $g_\theta = g_\theta h_C^{-1} h_C = \zeta_\theta(h_C^{-1}) g_\eta h_C$, and hence

$$\begin{aligned} \phi(x)^{-1} \phi(y) &= \phi(x^{-1}y) = \phi(g_\theta^{-1}u(C)g_\theta) = \phi(h_C^{-1}g_\eta^{-1}u(C)g_\eta h_C) \\ &= \phi(h_C^{-1}g_\eta^{-1}h_C) \phi(h_C^{-1}u(C)h_C) \phi(h_C^{-1}g_\eta h_C) \\ &= \phi(h_C^{-1}g_\eta h_C)^{-1} u'(\phi(C)) \phi(h_C^{-1}g_\eta h_C). \end{aligned}$$

We also have

$$\begin{aligned} \phi(h_C^{-1}g_\eta h_C) \phi(h) &= \phi(h_C^{-1}g_\eta h_C h h_C^{-1} h_C) \\ &= \phi(h_C^{-1} \zeta_\eta(h_C h h_C^{-1}) g_{\eta'} h_C) \\ &= \phi(h_C^{-1} \zeta_\eta(h_C h h_C^{-1}) h_C) \phi(h_C^{-1} g_{\eta'} h_C). \end{aligned}$$

Thus by Proposition 1.9 one gets

$$\begin{aligned} a_{\phi(y), \phi(x)}^{(\phi_C(i))} \cdot \phi(h) &= \chi_{\phi(C)}^{(\phi_C(i))} (\phi(h_C^{-1} \zeta_\eta(h_C h h_C^{-1}) h_C)) a_{\phi(yh), \phi(xh)}^{(\phi_C(i))} \\ &= \chi_C^{(i)} (\zeta_\eta(h_C h h_C^{-1})) a_{\phi(yh), \phi(xh)}^{(\phi_C(i))}. \end{aligned}$$

Now it follows from Eq.(21) that

$$\begin{aligned} \psi(a_{y,x}^{(i)} \cdot \phi(h)) &= \chi_C^{(i)}(\zeta_\theta(h_C^{-1})) a_{\phi(y), \phi(x)}^{(\phi_C(i))} \cdot \phi(h) \\ &= \chi_C^{(i)}(\zeta_\theta(h_C^{-1})) \chi_C^{(i)}(\zeta_\eta(h_C h h_C^{-1})) a_{\phi(yh), \phi(xh)}^{(\phi_C(i))} \\ &= \psi(a_{y,x}^{(i)} \cdot h). \end{aligned}$$

This shows that ϕ is a right kG -module homomorphism, and hence a kG -Hopf bimodule isomorphism from (kQ_1, α^X) to $\phi^{-1}(kQ'_1, \alpha^{X'})^{\phi^{-1}}$.

(ii) \Rightarrow (i). Assume that there exist a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ and a kG -Hopf bimodule isomorphism $\psi : (kQ_1, \alpha^X) \rightarrow \phi^{-1}(kQ'_1, \alpha^{X'})\phi^{-1}$. Then $\phi : G \rightarrow G'$ is a group isomorphism. Let $C \in \mathcal{K}(G)$. Then $\phi(u(C)), u'(\phi(C)) \in \phi(C) \in \mathcal{K}(G')$, and hence $u'(\phi(C)) = \phi(h_C)^{-1}\phi(u(C))\phi(h_C) = \phi(h_C^{-1}u(C)h_C)$ for some $h_C \in G$. Since ψ is a kG' -bicomodule isomorphism from $\phi(kQ_1, \alpha^X)\phi$ to $(kQ'_1, \alpha^{X'})$ and $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$, by restriction one gets a k -linear isomorphism

$$\psi_C : h_C^{-1}u(C)h_C(kQ_1)^1 \rightarrow u'(\phi(C))(kQ'_1)^1, x \mapsto \psi(x).$$

We also have a k -linear isomorphism

$$f_C : u(C)(kQ_1)^1 \rightarrow h_C^{-1}u(C)h_C(kQ_1)^1, x \mapsto h_C^{-1} \cdot x \cdot h_C.$$

Since $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$ and $h_C^{-1}Z_{u(C)}h_C = Z_{h_C^{-1}u(C)h_C}$, one gets $\phi(h_C^{-1}Z_{u(C)}h_C) = Z_{u'(\phi(C))}$. Hence ϕ and h_C induce an algebra isomorphism

$$\sigma_C : kZ_{u(C)} \rightarrow kZ_{u'(\phi(C))}, h \mapsto \phi(h_C^{-1}hh_C).$$

Using the hypothesis that ψ is a kG -bimodules homomorphism from (kQ_1, α^X) to $\phi(kQ'_1, \alpha^{X'})\phi$, one can easily check that the composition $\psi_C f_C$ is a right $kZ_{u(C)}$ -module isomorphism from $u(C)(kQ_1)^1$ to $(u'(\phi(C))(kQ'_1)^1)_{\sigma_C}$. Since both $u(C)(kQ_1)^1$ and $(u'(\phi(C))(kQ'_1)^1)_{\sigma_C}$ are pointed right $kZ_{u(C)}$ -modules, they are semisimple $kZ_{u(C)}$ -modules for any $C \in \mathcal{K}_r(G)$. Moreover, $ka_{u(C),1}^{(i)}$ and $ka_{u'(\phi(C)),1}^{(j)}$ are simple submodules of $u(C)(kQ_1)^1$ and $(u'(\phi(C))(kQ'_1)^1)_{\sigma_C}$, respectively, for any $i \in I_C(r)$ and $j \in I_{\phi(C)}(r')$, where $C \in \mathcal{K}_r(G)$. Thus for any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $ka_{u(C),1}^{(i)}$ and $(ka_{u'(\phi(C)),1}^{(\phi_C(i))})_{\sigma_C}$ are isomorphic right $kZ_{u(C)}$ -modules for any $i \in I_C(r)$, which implies $\chi_C^{(i)}(h) = \chi'_{\phi(C)}^{(\phi_C(i))}(\phi(h_C^{-1}hh_C))$ for any $h \in Z_{u(C)}$ and $i \in I_C(r)$. It follows that $RSC(G, r, \vec{\chi}, u) \cong RSC(G', r', \vec{\chi}', u')$. \square

By Theorem 4 we have

Corollary 1.13 *Let (Q, G, r) be a Hopf quiver. Then there are one-to-one correspondences between the following sets:*

(i) $\{[(G, r, \vec{\chi}, u)] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (ii) $\{[(kG; (kQ_1, \alpha^X))] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (iii) $\{[kQ^c(\alpha^X)] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (iv) $\{[kQ^s(\alpha^X)] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (v) $\{[kG[kQ_1, \alpha^X]] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (vi) $\{[kQ^a(\alpha^X)] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (vii) $\{[kQ^{sc}(\alpha^X)] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (viii) $\{[(kG)^*[kQ_1^a, \alpha^X]] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. (ix) $\{[(kG)^*; (kQ_1^a, \alpha^X)] \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. Note that $[x]$ in (i) denotes the isomorphic class of RSC containing x ; $[x]$ in (ii) denotes the isomorphic class of pull-push Hopf bimodules containing x ; $[x]$ in (iii) – (viii) denote the isomorphic class of graded Hopf algebras containing x . $[x]$ in (ix) denotes the isomorphic class of pull-push Hopf bimodules containing x .

Let $\mathcal{N}(G, r)$ denote the number of elements in the set in Corollary 1.13 (i) when the Hopf quiver (Q, G, r) is finite. Thus all the sets in Corollary 1.13 have the same cardinal number $\mathcal{N}(G, r)$. Then we have

Corollary 1.14 *Let (Q, G, r) be a finite Hopf quiver. Then*

$$\mathcal{N}(G, r) \leq |G|^{|G|} \prod_{C \in \mathcal{K}_r(G)} |\widehat{Z}_{u(C)}|^{r_C} .$$

Example 1.15 *Assume that k is a field with $\text{char}(k) \neq 2$. Let $G = \{1, g\} \cong \mathbf{Z}_2$ be the cyclic group of order 2 with the generator g . Let r be a ramification data of G with $r_1 = m$ and $r_g = 0$ and (Q, G, r) be the corresponding Hopf quiver, where m is a positive integer. Then ${}^1(Q_1)^1 = \{a_{1,1}^{(i)} \mid i = 1, 2, \dots, m\}$, ${}^g(Q_1)^g = \{a_{g,g}^{(i)} \mid i = 1, 2, \dots, m\}$, ${}^1(Q_1)^g$ and ${}^g(Q_1)^1$ are two empty sets. For simplification, we write $x_i = a_{1,1}^{(i)}$ and $y_i = a_{g,g}^{(i)}$ for any $1 \leq i \leq m$. Clearly, $Z_{u(\{1\})} = G$ and $\widehat{G} = \{\chi_+, \chi_-\}$, where $\chi_{\pm}(g) = \pm 1$. For any $0 \leq n \leq m$, put $\overrightarrow{\chi}_n \in (\widehat{G})^m$ with*

$$\chi_{n\{1\}}^{(i)} = \begin{cases} \chi_-, & \text{if } i > n; \\ \chi_+, & \text{otherwise.} \end{cases}$$

Then $\{RSC(G, r, \overrightarrow{\chi}_n, u_n) \mid n = 0, 1, 2, \dots, m\}$ are all non-isomorphic RSC's. Thus by Theorem 4 we know that the path coalgebra kQ^c exactly admits $m+1$ distinct PM co-path Hopf algebra structures $kQ^c(\alpha^{\chi_n})$, $0 \leq n \leq m$, up to graded Hopf algebra isomorphism. Now let $0 \leq n \leq m$. Then by Proposition 1.9, the kG -actions on (kQ_1, α^{χ_n}) are given by

$$g \cdot x_i = y_i, \quad g \cdot y_i = x_i, \quad 1 \leq i \leq m,$$

$$x_i \cdot g = \begin{cases} -y_i, & \text{if } i > n, \\ y_i, & \text{otherwise,} \end{cases} \quad y_i \cdot g = \begin{cases} -x_i, & \text{if } i > n, \\ x_i, & \text{otherwise.} \end{cases}$$

Thus by [9, p.245 or Theorem 3.8], the products of these arrows x_i, y_j in $kQ^c(\alpha^{\chi_n})$ can be described as follow. For any $i, j = 1, 2, \dots, m$,

$$x_i \cdot x_j = x_i x_j + x_j x_i, \quad y_i \cdot x_j = y_i y_j + y_j y_i,$$

$$x_i \cdot y_j = \begin{cases} -(y_i y_j + y_j y_i) & , \text{ if } i > n, \\ y_i y_j + y_j y_i & , \text{ otherwise,} \end{cases} \quad y_i \cdot y_j = \begin{cases} -(x_i x_j + x_j x_i) & , \text{ if } i > n, \\ x_i x_j + x_j x_i & , \text{ otherwise,} \end{cases}$$

where $x \cdot y$ denotes the product of x and y in $kQ^c(\alpha^{\chi_n})$ for any $x, y \in kQ^c(\alpha^{\chi_n})$, $x_i x_j$ and $y_i y_j$ denote the 2-paths in the quiver Q as usual for any $1 \leq i, j \leq m$.

2 Multiple crown algebras and multiple Taft algebras

In this section we discuss the PM quiver Hopf algebras defined by the RSC's with $\mathcal{K}_r(G) \subseteq Z(G)$.

Let r be a ramification data of G and (Q, G, r) be the corresponding Hopf quiver. If C contains only one element of G for any $C \in \mathcal{K}_r(G)$, then $C = \{g\}$ for some $g \in Z(G)$, the center of G . In this case, we say that the ramification r is *central*, and that $RSC(G, r, \vec{\chi}, u)$ a *central ramification system with characters*, or a CRSC in short. If r is a central ramification data of G , then we may write g for $\{g\}$ when $\{g\} \in \mathcal{K}_r(G)$. If $RSC(G, r, \vec{\chi}, u)$ is CRSC, then the PM co-path Hopf algebra $kQ^c(\alpha^x)$ is called a *multiple crown algebra* and $kG[kQ_1, \alpha^x]$ is called a *multiple Taft algebra*.

Definition 2.1 $(G, \vec{g}, \vec{\chi}, J)$ is called an *element system with characters* (simply, ESC) if G is a group, J is a set, $\vec{g} = \{g_i\}_{i \in J} \in Z(G)^J$ and $\vec{\chi} = \{\chi_i\}_{i \in J} \in \widehat{G}^J$ with $g_i \in Z(G)$ and $\chi_i \in \widehat{G}$. $ESC(G, \vec{g}, \vec{\chi}, J)$ and $ESC(G', \vec{g}', \vec{\chi}', J')$ are said to be *isomorphic* if there exist a group isomorphism $\phi : G \rightarrow G'$ and a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_i) = g'_{\sigma(i)}$ and $\chi'_{\sigma(i)}\phi = \chi_i$ for any $i \in J$.

$ESC(G, \vec{g}, \vec{\chi}, J)$ can be written as $ESC(G, g_i, \chi_i; i \in J)$ for convenience. Let $q_{ij} = \chi_i(g_j)$, $q_i = q_{ii}$ and N_i be the order of q_i ($N_i = \infty$ when q_i is not a root of unit, or $q_i = 1$) for $i, j \in J$.

Let H be a Hopf algebra with a bijective antipode S . A YD H -module V is *pointed* if $V = 0$ or V is a direct sum of one dimensional YD H -modules. If V is a pointed YD H -module, then the corresponding Nichols algebra $\mathcal{B}(V)$ is called a PM *Nichols algebra*.

Lemma 2.2 Let (V, α^-, δ^-) be a YD kG -module. Then (V, α^-, δ^-) is a pointed YD kG -module if and only if (V, α^-) is a pointed (left) kG -module with $\delta^-(V) \subseteq kZ(G) \otimes V$.

Proof. We may assume $V \neq 0$. Assume that V is a pointed YD kG -module. Then V is obviously a pointed kG -module. Now let U be a one dimensional YD kG -submodule of V and let $0 \neq x \in U$. Then $\delta^-(x) = g \otimes x$ for some $g \in G$. Let $h \in G$. Then $h \cdot x = \beta x$ for some $0 \neq \beta \in k$. By Eq.(4), one gets that $\beta g \otimes x = \delta^-(\beta x) = \delta^-(h \cdot x) = hgh^{-1} \otimes h \cdot x = \beta hgh^{-1} \otimes x$. This implies that $hgh^{-1} = g$ for any $h \in G$, and hence $g \in Z(G)$. It follows that $\delta^-(V) \subseteq kZ(G) \otimes V$. Conversely, assume that V is a pointed kG -module and $\delta^-(V) \subseteq kZ(G) \otimes V$. For any $g \in G$, let

$$V_g = \{v \in V \mid \delta^-(v) = g \otimes v\}.$$

Then V_g is a kG -subcomodule of V and any subspace of V_g is a kG -subcomodule for all $g \in G$. By $\delta^-(V) \subseteq kZ(G) \otimes V$, we know that $V = \bigoplus_{g \in Z(G)} V_g$. Now let $g \in Z(G)$.

Then it follows from Eq.(4) that V_g is a kG -submodule of V . Since V is a pointed kG -module, any submodule of V is pointed, and hence V_g is a pointed kG -module. Thus if $V_g \neq 0$ then V_g is a direct sum of some one dimensional kG -submodules of V_g . However, any one dimensional kG -submodule of V_g is also a kG -subcomodule, and hence a YD kG -submodule. Thus V is a pointed YD kG -module. \square

Lemma 2.3 *If G is a finite commutative group with $|G|=n$, k contains all roots of all irreducible polynomials over k with degree $\leq n$ and the order of G is not divisible by $\text{char}(k)$, then every YD kG -module is a pointed one and Nichols algebra of every YD kG -module is a PM one.*

Proof. By Lemma 1.2 and Lemma 2.2, every YD kG -module is a pointed one. It follows from the definition of PM Nichols algebra that Nichols algebra of every YD kG -module is a PM one. \square

Let $(G, g_i, \chi_i; i \in J)$ be an ESC. Let V be a k -vector space with $\dim(V) = |J|$. Let $\{x_i \mid i \in J\}$ be a basis of V over k . Define a left kG -action and a left kG -coaction on V by

$$g \cdot x_i = \chi_i(g)x_i, \quad \delta^-(x_i) = g_i \otimes x_i, \quad i \in J, \quad g \in G.$$

Then it is easy to see that V is a pointed YD kG -module and kx_i is a one dimensional YD kG -submodule of V for any $i \in J$. Denote by $V(g_i, \chi_i; i \in J)$ the pointed YD kG -module V . Note that $V(g_i, \chi_i; i \in J) = 0$ if J is empty.

Proposition 2.4 *V is pointed YD kG -module if and only if V is isomorphic to $V(g_i, \chi_i; i \in J)$ for some ESC $(G, g_i, \chi_i; i \in J)$.*

Proof. If $V \cong V(g_i, \chi_i; i \in J)$ for some ESC $(g_i, \chi_i; i \in J)$ of G , then V is obviously a pointed YD kG -module. Conversely, assume that V is a nonzero pointed YD kG -module. By Lemma 2.2 and its proof, $V = \bigoplus_{g \in Z(G)} V_g$ and V_g is a pointed YD kG -submodule of V for any $g \in Z(G)$. Let $g \in Z(G)$ with $V_g \neq 0$. Then V_g is a nonzero pointed kG -module. Hence there is a k -basis $\{x_i \mid i \in J_g\}$ such that kx_i is a kG -submodule of V_g for any $i \in J_g$. It follows that there is a character $\chi_i \in \widehat{G}$ for any $i \in J_g$ such that $h \cdot x_i = \chi_i(h)x_i$ for all $h \in G$. For any $i \in J_g$, put $g_i = g$. We may assume that these index sets J_g are disjoint, that is, $J_g \cap J_h = \emptyset$ for any $g \neq h$ in $Z(G)$ with $V_g \neq 0$ and $V_h \neq 0$. Now let J be the union of all the J_g with $g \in Z(G)$ and $V_g \neq 0$. Then one can see that $(g_i, \chi_i; i \in J)$ is an ESC of G , and that V is isomorphic to $V(g_i, \chi_i; i \in J)$ as a YD kG -module. \square

Assume that $(g_i, \chi_i; i \in J)$ is an ESC of G . We define a binary relation \sim on J by

$$i \sim j \Leftrightarrow g_i = g_j,$$

where $i, j \in J$. Clearly, this is an equivalence relation. Denote by J/\sim the quotient set of J modulo \sim . For any $i \in J$, let $[i]$ denote the equivalence class containing i . That is, $[i] := \{j \in J \mid j \sim i\}$. Choose a subset $\bar{J} \subseteq J$ such that the assignment $i \mapsto [i]$ is a bijective map from \bar{J} to J/\sim . Then $J = \bigcup_{i \in \bar{J}} [i]$ is a disjoint union. Let $r_{g_i} = |[i]|$ for any $i \in \bar{J}$. Then $r = \sum_{i \in \bar{J}} r_{g_i} g_i$ is a central ramification of G with $I_{g_i}(r) = [i]$ for $i \in \bar{J}$. Moreover, $\mathcal{K}_r(G) = \{\{g_i\} \mid i \in \bar{J}\}$. Put $\chi_{g_i}^{(j)} := \chi_j$ for any $i \in \bar{J}$ and $j \in [i]$. We obtain an *CRSC*, written $CRSC(G, r(g_i, \chi_i; i \in J), \vec{\chi}(g_i, \chi_i; i \in J), u)$. Let (Q, G, r) be the corresponding Hopf quiver with $r = r(g_i, \chi_i; i \in J)$ and denote by $(kQ_1; g_i, \chi_i; i \in J)$ the kG -Hopf bimodule (kQ_1, α^χ) . Denote by $kQ^c(g_i, \chi_i; i \in J)$ and $kG[kQ_1, g_i, \chi_i; i \in J]$ the corresponding multiple crown algebra $kQ^c(\alpha^\chi)$ and multiple Taft algebra $kG[kQ_1, \alpha^\chi]$, respectively. We also denote by $kQ^a(g_i, \chi_i; i \in J)$, $kQ^{sc}(g_i, \chi_i; i \in J)$ and $kQ^s(g_i, \chi_i; i \in J)$ the corresponding path Hopf algebra $kQ^a(\alpha^\chi)$, semi-co-path Hopf algebra $kQ^{sc}(\alpha^\chi)$, semi-path Hopf algebra $kQ^s(\alpha^\chi)$ and one-type- path Hopf algebra $(kG)^*[kQ_1^a, \alpha^\chi]$, respectively.

Conversely, assume that $(G, r, \vec{\chi}, u)$ is a *CRSC*. We may assume $I_g(r) \cap I_h(r) = \emptyset$ for any $g \neq h$ in $\mathcal{K}_r(G)$. Let $J := \bigcup_{g \in \mathcal{K}_r(G)} I_g(r)$. For any $i \in J$, put $g_i := g$ and $\chi_i := \chi_g^{(i)}$ if $i \in I_g(r)$ with $g \in \mathcal{K}_r(G)$. We obtain an *ESC*, written $ESC(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J)$.

From now on, assume $I_g(r) \cap I_h(r) = \emptyset$ for any $g, h \in \mathcal{K}_r(G)$ with $g \neq h$. Let E_j denote $a_{g_j, 1}^{(j)}$ for $j \in I_{\{g_j\}}(r)$. In fact, in two cases above, for every arrow $b \in Q_1^1$, there exists $j \in J$ such that $b = a_{g_j, 1}^{(j)}$.

Proposition 2.5 $CRSC(G, r, \vec{\chi}, u) \cong CRSC(G', r', \vec{\chi}', u')$ if and only if $ESC(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J) \cong ESC(G', \vec{g}'(r', \vec{\chi}', u'), \vec{\chi}'(r', \vec{\chi}', u'), J')$.

Proof. We use notations above. If $CRSC(G, r, \vec{\chi}, u) \cong CRSC(G', r', \vec{\chi}', u')$, then there exists a group isomorphism $\phi : G \rightarrow G'$; for any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}^{(\phi_C(i))} \phi = \chi_C^{(i)}$. Let σ be the bijection from J to J' such that $\sigma(i) = \phi_C(i)$ for any $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$. It is clear that $\phi(g_j) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)} \phi = \chi_j$ for any $j \in J$. Thus $ESC(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J) \cong ESC(G', \vec{g}'(r', \vec{\chi}', u'), \vec{\chi}'(r', \vec{\chi}', u'), J')$.

Conversely, if $ESC(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J) \cong ESC(G', \vec{g}'(r', \vec{\chi}', u'), \vec{\chi}'(r', \vec{\chi}', u'), J')$, then there exist a group isomorphism $\phi : G \rightarrow G'$, a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_j) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)} \phi = \chi_j$ for any $j \in J$. For any $C \in \mathcal{K}_r(G)$, $j \in I_C(r)$, there exists $i \in \bar{J}$ such that $\{g_i\} = C$ and we have

$$\begin{aligned} \chi'_{\phi(C)}^{(\phi_C(j))} \phi &= \chi'_{g'_{\sigma(i)}}^{(\sigma(j))} \phi \\ &= \chi'_{\sigma(j)} \phi = \chi_j \\ &= \chi_C^{(j)}. \quad \square \end{aligned}$$

Let (Q, G, r) be a Hopf quiver. Let $Q_1^1 := \{a \in Q_1 \mid s(a) = 1\}$. Clearly, kQ_1^1 is a left kG -subcomodule of kQ_1 . Now assume that $(kQ_1, \alpha^-, \alpha^+)$ is a kG -Hopf bimodule for some α^- and α^+ . Define a new left kG -action on kQ_1 by

$$g \triangleright x := g \cdot x \cdot g^{-1}, \quad g \in G, x \in kQ_1,$$

where $g \cdot x = \alpha^-(g \otimes x)$ and $x \cdot g = \alpha^+(x \otimes g)$ for any $g \in G$ and $x \in kQ_1$. With this left kG -action and the original left (arrow) kG -coaction δ^- , kQ_1 is a YD kG -module and kQ_1^1 is a YD kG -submodule of kQ_1 . Denote them by $(kQ_1, ad(\alpha^-, \alpha^+))$ and $(kQ_1^1, ad(\alpha^-, \alpha^+))$, respectively. We denote by $(kQ_1, ad(\alpha^x))$ and $(kQ_1^1, ad(\alpha^x))$ the two corresponding YD kG -modules, respectively, when $(kQ_1, \alpha^-, \alpha^+) = (kQ_1, \alpha^x)$ for some $RSC(G, r, \vec{\chi}, u)$. In particular, when $RSC(G, r, \vec{\chi}, u)$ determined by a $(G, g_i, \chi_i; i \in J)$, i.e. $RSC(G, r, \vec{\chi}, u) = CRSC(G, r(G, g_i, \chi_i; i \in J), \vec{\chi}(G, g_i, \chi_i; i \in J), u)$ we write $(kQ_1, ad(g_i, \chi_i; i \in J)) := (kQ_1, ad(\alpha^x))$ and $(kQ_1^1, ad(g_i, \chi_i; i \in J)) := (kQ_1^1, ad(\alpha^x))$, respectively.

Lemma 2.6 $(kQ_1^1, ad(g_i, \chi_i; i \in J))$ and $V(g_i, \chi_i^{-1}; i \in J)$ are isomorphic YD kG -modules.

Proof. By definition, $V(g_i, \chi_i^{-1}; i \in J)$ has a k -basis $\{x_i \mid i \in J\}$ such that $\delta^-(x_i) = g_i \otimes x_i$ and $g \cdot x_i = \chi_i^{-1}(g)x_i$ for all $i \in J$ and $g \in G$. By Proposition 1.9, for any $j \in J$, we have that $g \triangleright a_{g_j, 1}^{(j)} = \chi_j(g^{-1})a_{g_j, 1}^{(j)}$ and $\delta^-(a_{g_j, 1}^{(j)}) = g_j \otimes a_{g_j, 1}^{(j)}$. It follows that there is a YD kG -module isomorphism from $(kQ_1^1, ad(g_i, \chi_i; i \in J))$ to $V(g_i, \chi_i^{-1}; i \in J)$ given by $a_{g_j, 1}^{(j)} \mapsto x_j$ for any $j \in J$. \square

Lemma 2.7 Let B and B' be two Hopf algebras with bijective antipodes. Let V be a YD B -module. Assume that there is a Hopf algebra isomorphism $\phi : B' \rightarrow B$. Then $\phi^{-1} \mathcal{B}(V) \cong \mathcal{B}(\phi^{-1} V)$ as graded braided Hopf algebras in ${}_{B'}^B \mathcal{YD}$.

Proof. We first show that $\phi^{-1} \mathcal{B}(V)$ is a Nichols algebra of $\phi^{-1} V$ by following steps. Let $R := \phi^{-1} \mathcal{B}(V)$.

(i) Obviously, $R_0 = k$, $R_1 = P(R)$ and R is generated by R_1 as algebras. Since $C(x \otimes y) = C'(x' \otimes y')$ for any $x', y' \in R$, $x, y \in \mathcal{B}(V)$ with $x' = x, y' = y$, where C and C' denote the braidings in ${}_{B'}^B \mathcal{YD}$ and ${}_{B'}^{B'} \mathcal{YD}$, respectively, we have that R is a graded braided Hopf algebra in ${}_{B'}^{B'} \mathcal{YD}$.

(ii) $R_1 = \phi^{-1} \mathcal{B}(V)_1 \cong \phi^{-1} V$ as YD B' -modules, since $\mathcal{B}(V)_1 \cong V$ as YD B -modules.

Consequently, $\phi^{-1} \mathcal{B}(V)$ is a Nichols algebra of $\phi^{-1} V$. By [4, Proposition 2.2 (iv)], $R \cong \mathcal{B}(\phi^{-1} V)$ as graded braided Hopf algebra in ${}_{B'}^{B'} \mathcal{YD}$. \square

Theorem 5 Assume that $(G, g_i, \chi_i; i \in J)$ and $(G', g'_i, \chi'_i; i \in J')$ are two ESC's. Then the following statements are equivalent:

- (i) $ESC(G, g_i, \chi_i; i \in J) \cong ESC(G', g'_i, \chi'_i; i \in J')$
- (ii) $CRSC(G, r(G, g_i, \chi_i; i \in J), \vec{\chi}(G, g_i, \chi_i; i \in J), u) \cong CRSC(G', r'(G', g'_i, \chi'_i; i \in J'), \vec{\chi}'(G', g'_i, \chi'_i; i \in J'), u')$.
- (iii) *There is a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $V(G, g_i, \chi_i; i \in J) \cong \phi^{-1} V'(G', g'_i, \chi'_i; i \in J')$ as YD kG -modules.*
- (iv) *There is a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $\mathcal{B}(V(G, g_i, \chi_i; i \in J)) \cong \phi^{-1} \mathcal{B}(V'(G', g'_i, \chi'_i; i \in J'))$ as graded braided Hopf algebras in ${}^{kG}_{kG} \mathcal{YD}$.*
- (v) *There is a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $(kQ_1^1, ad(g_i, \chi_i; i \in J)) \cong \phi^{-1} (kQ_1^1, ad(g'_i, \chi'_i; i \in J'))$ as YD kG -modules.*

Proof. We use the notations before Proposition 2.5.

(i) \Rightarrow (ii). There exist a group isomorphism $\phi : G \rightarrow G'$ and a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_j) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)}\phi = \chi_j$ for any $j \in J$. For any $C = \{g_i\} \in \mathcal{K}_r(G)$ and $j \in I_C(r)$, we have $g_i = g_j$ and

$$\begin{aligned} \chi_C^{(j)} = \chi_{g_i}^{(j)} &= \chi_j = \chi'_{\sigma(j)}\phi \\ &= \chi'_{g'_{\sigma(i)}}(\sigma(j))\phi = \chi'_{\phi(g_i)}(\phi_{g_i}(j))\phi = \chi'_{\phi(C)}(\phi_C(j))\phi. \end{aligned}$$

(ii) \Rightarrow (i). There is a group isomorphism $\phi : G \rightarrow G'$ and a bijection $\tau : \bar{J} \rightarrow \bar{J}'$ such that $\phi(g_i) = g'_{\tau(i)}$ for any $i \in \bar{J}$, and there is a bijection $\phi_{g_i} : [i] \rightarrow [\tau(i)]$ for any $i \in \bar{J}$ such that $\chi'_{g'_{\tau(i)}}(\phi_{g_i}(j))\phi = \chi_{g_i}^{(j)}$ for any $i \in \bar{J}$ and $j \in [i]$. Since $J = \bigcup_{i \in \bar{J}} [i]$ and $J' = \bigcup_{i \in \bar{J}'} [\tau(i)]$, there is a bijective map $\sigma : J \rightarrow J'$ given by

$$\sigma(j) = \phi_{g_i}(j), \quad \text{for any } j \in [i], i \in \bar{J}.$$

Now let $j \in J$. Then there is a unique $i \in \bar{J}$ such that $j \in [i]$. Hence $\tau(i) \in \bar{J}'$ and $\sigma(j) = \phi_{g_i}(j) \in [\tau(i)]$. Thus we have that $\phi(g_j) = \phi(g_i) = g'_{\tau(i)} = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)}\phi = \chi'_{g'_{\tau(i)}}(\phi_{g_i}(j))\phi = \chi_{g_i}^{(j)} = \chi_j$. This shows that $ESC(G, \vec{g}, \vec{\chi}, J) \cong ESC(G', \vec{g}', \vec{\chi}', J')$.

(i) \Leftrightarrow (iii). Let $V := V(G, g_i, \chi_i; i \in J)$ and $V' := V'(G', g'_i, \chi'_i; i \in J')$. By definition V has a k -basis $\{x_i \mid i \in J\}$ such that $g \cdot x_i = \chi_i(g)x_i$ and $\delta^-(x_i) = g_i \otimes x_i$ for any $i \in J$ and $g \in G$. Similarly, V' has a k -basis $\{y_j \mid j \in J'\}$ such that $h \cdot y_j = \chi'_j(h)y_j$ and $\delta^-(y_j) = g'_j \otimes y_j$ for all $j \in J'$ and $h \in G'$.

Assume $ESC(G, \vec{g}, \vec{\chi}, J) \cong ESC(G', \vec{g}', \vec{\chi}', J')$. Then there is a group isomorphism $\phi : G \rightarrow G'$ and a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_i) = g'_{\sigma(i)}$ and $\chi'_{\sigma(i)}\phi = \chi_i$ for any $i \in J$. Hence $\phi : kG \rightarrow kG'$ is a Hopf algebra isomorphism. Define a k -linear isomorphism $\psi : V \rightarrow V'$ by $\psi(x_i) = y_{\sigma(i)}$ for any $i \in J$. Then it is straightforward to check that ψ is a YD kG -module homomorphism from V to $\phi^{-1} V'$.

Conversely, assume that $\phi : kG \rightarrow kG'$ is a Hopf algebra isomorphism and $\psi : V \rightarrow \phi^{-1}V'$ is a YD kG -module isomorphism. Then $\phi : G \rightarrow G'$ is a group isomorphism. We use the notations in the proof of Lemma 2.2 and the notations above. Then $\psi(V_g) = V'_{\phi(g)}$ for any $g \in G$. Since $V = \bigoplus_{i \in \bar{J}} V_{g_i}$ and $V' = \bigoplus_{i \in \bar{J}'} V'_{g'_i}$, there is a bijection $\tau : \bar{J} \rightarrow \bar{J}'$ such that $\psi(V_{g_i}) = V'_{g'_{\tau(i)}}$ for any $i \in \bar{J}$. This shows that $\phi(g_i) = g'_{\tau(i)}$ and $V_{g_i} \cong \phi(V'_{g'_{\tau(i)}})$ as left kG -modules for any $i \in \bar{J}$. However, V_{g_i} is a pointed kG -module and kx_j is its one dimensional submodule for any $j \in [i]$. Similarly, $V'_{g'_{\tau(i)}}$ is a pointed kG' -module and ky_j is its one dimensional submodule for any $j \in [\tau(i)]$. Hence there is a bijection $\phi_{g_i} : [i] \rightarrow [\tau(i)]$ for any $i \in \bar{J}$ such that kx_j and $\phi(ky_{\phi_{g_i}(j)})$ are isomorphic kG -modules for all $j \in [i]$. This implies that $\chi_j = \chi'_{\phi_{g_i}(j)}\phi$ for all $j \in [i]$ and $i \in \bar{J}$. Then the same argument as in the proof of (ii) \Rightarrow (i) shows that $ESC(G, \vec{g}, \vec{\chi}, J) \cong ESC(G', \vec{g}', \vec{\chi}', J')$.

(iii) \Leftrightarrow (iv) It follows from Lemma 2.7.

(iii) \Leftrightarrow (v) It follows from Lemma 2.6. \square

3 The relation between quiver Hopf algebras and quotients of free algebras

$H = \bigoplus_{i \geq 0} H_i$ be a graded Hopf algebra with antipode S . Let $B = H_0$, and let $\pi_0 : H \rightarrow H_0 = B$ and $\iota_0 : B = H_0 \rightarrow H$ denote the canonical projection and injection. Set $\omega := id_H * (\iota_0 \pi_0 S) : H \rightarrow H$. Then it is clear that (H, δ^+, α^+) is a right B -Hopf module with $\delta^+ := (id \otimes \pi_0)\Delta$ and $\alpha^+ := \mu(id \otimes \iota_0)$. Let $R := H^{coB} := \{h \in H \mid \delta^+(h) = h \otimes 1\}$, which is a graded subspace of H . Then it is known that $R = \text{Im}(\omega)$ and $\Delta(R) \subseteq H \otimes R$. Hence R is a left coideal subalgebra of H , and so R is a left H -comodule algebra. It is well known that R is a graded braided Hopf algebra in ${}^B_B\mathcal{YD}$ with the same multiplication, unit and counit as in H , the comultiplication $\Delta_R = (\omega \otimes id)\Delta$, where the left B -action α_R and left B -coaction δ_R on R are given by

$$b \rightharpoonup_{ad} x = \sum b_{(1)}xS(b_{(2)}), \quad \delta_R^-(x) = \sum \pi_0(x_{(1)}) \otimes x_{(2)}, \quad b \in B, \quad x \in R$$

(see the proof of [22, Theorem 3]). R is called the diagram of H , written $diag(H)$. By [22, Theorem 1], the biproduct of R and B is a Hopf algebra, written $R_{\alpha_R}^{\delta_R} \# B$, or $R \# B$ in short. The biproduct $R_{\alpha_R}^{\delta_R} \# B$ is also called the bosonization of R . Furthermore, we have the following well known result.

Theorem 6 (see [20, p.1530], [2] and [22]) *Under notations above, if $H = \bigoplus_{i \geq 0} H_i$ is a graded Hopf algebra, then R is a graded braided Hopf algebra in ${}^B_B\mathcal{YD}$ and $R \# B \cong H$ as graded Hopf algebras, where the isomorphism is $\alpha^+ := \mu_H(id_H \otimes \iota_0)$.*

Lemma 3.1 (i) Let H and H' be two graded Hopf algebras with $B = H_0$ and $B' = H'_0$. Then $H \cong H'$ as graded Hopf algebras if and only if $(B; \text{diag}(H)) \cong (B'; \text{diag}(H'))$ as graded pull-push YD Hopf algebras.

(ii) Let B and B' be two Hopf algebras. Let M and M' be B -Hopf bimodule and B' -Hopf bimodule, respectively. Then $B[M] \cong B'[M']$ as graded Hopf algebras if and only if $(B; \text{diag}(B[M])) \cong (B'; \text{diag}(B'[M']))$ as graded pull-push YD Hopf algebras.

Proof. (i) Assume that ξ is a graded Hopf algebra isomorphism from H to H' . Let $R := \text{diag}(H)$, $R' := \text{diag}(H')$, $\phi := \xi|_B$ and $\psi := \xi|_{R'}$. Obviously, ψ is a algebra homomorphism from R to R' . We show that (ϕ, ψ) is graded pull-push YD Hopf algebra isomorphism from $(B; R)$ to $(B'; R')$ by following several steps.

(1). ψ is a B -module homomorphism from R to $\phi^{-1}R'$. For any $b \in B, r \in R$, see

$$\begin{aligned} \psi(b \cdot r) &= \xi(\mu_H(\iota_0 \otimes id)(b \otimes r)) \\ &= \mu_{H'}(\xi \otimes \xi)(\iota_0 \otimes id)(b \otimes r) \\ &= \mu_{H'}(\iota_0 \phi \otimes \xi)(b \otimes r) \quad (\text{since } \xi \text{ is graded}) \\ &= \phi(b) \cdot \psi(r). \end{aligned}$$

(2). Similarly, ψ is a B -comodule homomorphism from R to $\phi^{-1}R'$.

(3). ψ is a coalgebra homomorphism from R to R' . Indeed, for any $r \in R$, see

$$\begin{aligned} (\psi \otimes \psi)\Delta_R(r) &= (\psi \otimes \psi)(\omega_H \otimes id)\Delta_H(r) \\ &= (\psi \otimes \psi)(\mu_H(id \otimes \iota_0 \pi_0 S_H) \otimes id)(\Delta_H \otimes id)\Delta_H(r) \\ &= \Delta_{R'}\psi(r). \end{aligned}$$

Conversely, assume that (ϕ, ψ) is graded pull-push YD Hopf algebra isomorphism from $(B; R)$ to $(B'; R')$ with $R := \text{diag}(H)$ and $R' := \text{diag}(H')$. By Theorem 6, $R\#B \cong H$ and $R'\#B' \cong H'$ as graded Hopf algebras. Let ξ be a linear map from $R\#B$ to $R'\#B'$ by sending $r\#b$ to $\psi(r)\#\phi(b)$ for any $r \in R, b \in B$. Let ν be a linear map from $R'\#B'$ to $R\#B$ by sending $r'\#b'$ to $\psi^{-1}(r')\#\phi^{-1}(b')$ for any $r' \in R', b' \in B'$. Obviously, ν is the inverse of ξ . Since ψ is graded, so is ξ .

Now we show that ξ is an algebra homomorphism. For any $r, r' \in R, b, b' \in B$, see

$$\begin{aligned} \xi\mu_{R\#B}((r\#b) \otimes (r'\#b')) &= \psi(r(b_{(1)} \cdot r'))\#\phi(b_{(2)}b') \\ &= \psi(r)(\phi(b_{(1)}) \cdot \psi(r'))\#\phi(b_{(2)})\phi(b) \\ &\quad (\text{since } \psi \text{ is a pullback module homomorphism} \\ &\quad \text{and an algebra homomorphism.}) \\ &= \mu_{R'\#B'}(\xi(r\#b) \otimes \xi(r'\#b')). \end{aligned}$$

Similarly, we can show that ξ is a coalgebra homomorphism.

(ii) It follows from (i). \square

Lemma 3.2 (i) $(kQ^c(\alpha^X))^{co\ kG} = \text{span}\{\beta \mid \beta \text{ is a path with } s(\beta) = 1\}$.

(ii) $(kG[kQ_1, \alpha^X])^{co\ kG}$ is the subalgebra of $kG[kQ_1^c, \alpha^X]$ generated by Q_1^1 as algebras.

(iii) $(kG[kQ_1, \alpha^X])^{co\ kG} \# kG \cong kG[kQ_1, \alpha^X]$ as graded Hopf algebra isomorphism

(iv) $(kQ^s(\alpha^X))^{co\ kG}$ is the subalgebra of $kG^s(\alpha^X)$ generated by Q_1^1 as algebras.

Proof. (i) For a path β , see that

$$\delta^+(\beta) = (id \otimes \pi_0)\Delta(\beta) = \beta \otimes s(\beta).$$

This implies $(kQ^c)^{co\ kG} = \text{span}\{\beta \mid \beta \text{ is a path with } s(\beta) = 1\}$.

(ii) Since every path generated by arrows in Q_1^1 is of start vertex 1, this path belongs to $(kG[kQ_1, \alpha^X])^{co\ kG}$. Let $R := (kG[kQ_1, \alpha^X])^{co\ kG}$ and $A :=$ the subalgebra of $kG[kQ_1^c, \alpha^X]$ generated by Q_1^1 as algebras. Obviously, $A \subseteq R$. It is clear that $\alpha^+(R \# kG) = \alpha^+(A \# kG) = kG[kQ_1, \alpha^X]$ and α^+ is injective. Thus $R \# kG = A \# kG$ and $R = A$.

(iii) It follows from Theorem 6.

(iv) We first show $(kQ^s)^{co\ kG} = \text{span}\{\beta \mid \beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_1 \text{ with } \prod_{i=1}^n s(\beta_i) = 1 \text{ and } \beta_i \in Q_1, i = 1, 2, \dots, n\}$. Indeed, obviously right hand side \subseteq the left hand side. For any $\beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_1$ with $\beta_i \in Q_1$, called a monomial, define $s(\beta) = \prod_{i=1}^n s(\beta_i)$. For any $0 \neq u \in (kQ^s)^{co\ kG}$ with $u \notin kG$, assume that u_1, u_2, \dots, u_n is linear independent, $u = \sum_{i=1}^n b_i u_i$ with $0 \neq b_i \in k$ and u_i is a monomial for $i = 1, 2, \dots, n$. See $\delta^+(u) = \sum_{i=1}^n b_i u_i \otimes s(u_i) = u \otimes 1$. Consequently, $s(u_i) = 1$ for $i = 1, 2, \dots, n$. This implies that u belongs to the right hand side.

For any $\beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_1$ with $\prod_{i=1}^n s(\beta_i) = 1$ and $\beta_i \in Q_1, i = 1, 2, \dots, n$, we show that β can be written as multiplication of arrows in Q_1^1 by induction. When $n = 1$, it is clear. For $n > 1$, see $\beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_2 \cdot s(\beta_1) \otimes_{kG} (s(\beta_1))^{-1} \cdot \beta_1$. Thus β can be written as multiplication of arrows in Q_1^1 . Consequently, we complete the proof of (iv). \square

Recall that a braided algebra A in braided tensor category (\mathcal{C}, C) with braiding C is said to be braided commutative or quantum commutative, if $ab = \mu C(a \otimes b)$ for any $a, b \in A$. An $ESC(G, g_i, \chi_i; i \in J)$ is said to be quantum commutative if

$$\chi_i(g_j)\chi_j(g_i) = 1$$

for any $i, j \in J$. An $ESC(G, g_i, \chi_i; i \in J)$ is said to be quantum weakly commutative if

$$\chi_i(g_j)\chi_j(g_i) = 1$$

for any $i, j \in J$ with $i \neq j$.

Lemma 3.3 (i) $ESC(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative if and only if in $diag(kG[kQ_1, g_i, \chi_i; i \in J])$,

$$E_i \cdot E_j = \chi_j(g_i^{-1})E_j \cdot E_i \quad (22)$$

for any $i, j \in J$ with $i \neq j$.

(ii) $diag(kG[kQ_1, g_i, \chi_i; i \in J])$ is quantum commutative in ${}^{kG}_{kG}\mathcal{YD}$ if and only if $ESC(G, g_i, \chi_i; i \in J)$ is quantum commutative.

Proof. (ii) By Theorem 6, $R := diag(kG[kQ_1, g_i, \chi_i; i \in J])$ is a braided Hopf algebra in ${}^{kG}_{kG}\mathcal{YD}$. If $diag(kG[kQ_1, g_i, \chi_i; i \in J])$ is quantum commutative in ${}^{kG}_{kG}\mathcal{YD}$, then

$$E_i \cdot E_j = \chi_j(g_i^{-1})E_j \cdot E_i \quad (23)$$

for any $i, j \in J$. By [9, Theorem 3.8] and Proposition 1.9, we have

$$\begin{aligned} E_i \cdot E_j &= a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + q_{ij} a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)}, \\ E_j \cdot E_i &= q_{ji} a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)}. \end{aligned}$$

Thus $\chi_i(g_j^{-1}) = \chi_j(g_i)$.

Conversely, if $\chi_i(g_j^{-1}) = \chi_j(g_i)$ for any $i, j \in J$, see

$$\begin{aligned} E_i \cdot E_j &= a_{g_i, 1}^{(i)} \cdot a_{g_j, 1}^{(j)} \\ &= a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + q_{ij} a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)} \\ &= q_{ij} (q_{ji} a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)}) \\ &= q_{ij} a_{g_j, 1}^{(j)} \cdot a_{g_i, 1}^{(i)} \\ &= q_{ij} E_j \cdot E_i. \end{aligned}$$

Since E'_i 's generate R as algebras, R is quantum commutative.

(i) It is similar to the proof of (ii). \square

For any positive integers m and n , let

$$D_n^{n+m} = \{d = (d_{n+m}, d_{n+m-1}, \dots, d_1) \mid d_i = 0 \text{ or } 1, \sum_{i=1}^{n+m} d_i = n\}.$$

Let $d \in D_n^{n+m}$ and let $A = a_n a_{n-1} \cdots a_1 \in Q_n$ be an n -path, i.e., $a_1, a_2, \dots, a_n \in Q_1$ with $t(a_i) = s(a_{i+1})$ for all $1 \leq i \leq n-1$. Then we define a sequence $dA = ((dA)_{n+m}, \dots, (dA)_1)$ by

$$(dA)_i = \begin{cases} t(a_{d(i)}) & , \text{ if } d_i = 0; \\ a_{d(i)} & , \text{ if } d_i = 1, \end{cases}$$

where $1 \leq i \leq n+m$ and $d(i) = \sum_{j=1}^i d_j$. Such a sequence dA is called an $n+m$ -thin splits of the n -path A . Note that if $d(i) = 0$ then we regard $t(a_{d(i)}) = s(a_1)$, since $s(a_{d(i)+1}) = s(a_1)$ in this case.

If $q \in k$ and $0 \leq i \leq n < \text{ord}(q)$ (the order of q), we set $(0)_q! = 1$,

$$\binom{n}{i}_q = \frac{(n)_q!}{(i)_q!(n-i)_q!}, \quad \text{where } (n)_q! = \prod_{1 \leq i \leq n} (i)_q, \quad (n)_q = \frac{q^n - 1}{q - 1}.$$

In particular, $(n)_q = n$ when $q = 1$.

Lemma 3.4 *In $kQ^c(\alpha^X)$, assume $\{g\} \in \mathcal{K}_r(G)$ and $j \in I_{\{g\}}(r)$. Let $q := \chi_g^{(j)}(g)$. If i_1, i_2, \dots, i_m be non-negative integers, then*

$$a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)} = q^{\beta_m}(m)_q! P_{g^{\alpha_m}}^{(j)}(g, m)$$

where $\alpha_m = i_1 + i_2 + \dots + i_m$, $P_h^{(j)}(g, m) = a_{g^m h, g^{m-1} h}^{(j)} a_{g^{m-1} h, g^{m-2} h}^{(j)} \cdots a_{g h, h}^{(j)}$, $\beta_1 = 0$ and $\beta_m = \sum_{j=1}^{m-1} (i_1 + i_2 + \dots + i_j)$ if $m > 1$.

Proof. We prove the equality by induction on m . For $m = 1$, it is easy to see that the equality holds. Now suppose $m > 1$. We have

$$\begin{aligned} & a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)} \\ = & a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot (a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)}) \\ = & q^{\beta_{m-1}}(m-1)_q! a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot P_{g^{\alpha_{m-1}}}^{(j)}(g, m-1) \quad (\text{by inductive assumption}) \\ = & q^{\beta_{m-1}}(m-1)_q! a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot (a_{g^{\alpha_{m-1}+m-1}, g^{\alpha_{m-1}+m-2}}^{(j)} \cdots a_{g^{\alpha_{m-1}+1}, g^{\alpha_{m-1}}}^{(j)}) \\ = & q^{\beta_{m-1}}(m-1)_q! \sum_{l=1}^m [(g^{i_m+1} \cdot a_{g^{\alpha_{m-1}+m-1}, g^{\alpha_{m-1}+m-2}}^{(j)}) \cdots (g^{i_m+1} \cdot a_{g^{\alpha_{m-1}+l}, g^{\alpha_{m-1}+l-1}}^{(j)}) \\ & (a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot g^{\alpha_{m-1}+l-1}) (g^{i_m} \cdot a_{g^{\alpha_{m-1}+l-1}, g^{\alpha_{m-1}+l-2}}^{(j)}) \cdots (g^{i_m} \cdot a_{g^{\alpha_{m-1}+1}, g^{\alpha_{m-1}}}^{(j)})] \\ & (\text{by [9, Theorem 3.8]}) \\ = & q^{\beta_{m-1}}(m-1)_q! \sum_{l=1}^m [a_{g^{\alpha_m+m}, g^{\alpha_m+m-1}}^{(j)} \cdots a_{g^{\alpha_m+l+1}, g^{\alpha_m+l}}^{(j)} \\ & (\chi(g^{\alpha_{m-1}+l-1}) a_{g^{\alpha_m+l}, g^{\alpha_m+l-1}}^{(j)} a_{g^{\alpha_m+l-1}, g^{\alpha_m+l-2}}^{(j)} \cdots a_{g^{\alpha_m+1}, g^{\alpha_m}}^{(j)})] \quad (\text{by Proposition 1.9}) \\ = & q^{\beta_{m-1}}(m-1)_q! \sum_{l=1}^m q^{\alpha_{m-1}+l-1} P_{g^{\alpha_m}}^{(j)}(g, m) \\ = & q^{\beta_{m-1}+\alpha_{m-1}}(m)_q! P_{g^{\alpha_m}}^{(j)}(g, m) \\ = & q^{\beta_m}(m)_q! P_{g^{\alpha_m}}^{(j)}(g, m). \quad \square \end{aligned}$$

Lemma 3.5 *(See [3, Lemma 3.3]) Let B be a Hopf algebra and R a braided Hopf algebra in ${}^B_B\mathcal{YD}$ with a linear independent set $\{x_1, \dots, x_t\} \subseteq P(R)$. Assume that there exist $g_j \in G(B)$ and $\chi_j \in \text{Alg}(B, k)$ such that*

$$\delta(x_j) = g_j \otimes x_j, \quad h \cdot x_j = \chi_j(h)x_j, \quad \text{for all } h \in B, j = 1, 2, \dots, t.$$

Then

$$\{x_1^{m_1} x_2^{m_2} \cdots x_t^{m_t} \mid 0 \leq m_j < N_j, 1 \leq j \leq t\}.$$

is linearly independent, where N_i is the order of $q_i := \chi_i(g_i)$ ($N_i = \infty$ when q_i is not a root of unit, or $q_i = 1$) for $1 \leq i \leq t$.

Proof. By the quantum binomial formula, if $1 \leq n_j < N_j$, then

$$\Delta(x_j^{n_j}) = \sum_{0 \leq i_j \leq n_j} \binom{n_j}{i_j}_{q_j} x_j^{i_j} \otimes x_j^{n_j - i_j}.$$

We use the following notation:

$$\mathbf{n} = (n_1, \dots, n_j, \dots, n_t), \quad x^{\mathbf{n}} = x_1^{n_1} \dots x_j^{n_j} \dots x_t^{n_t}, \quad |\mathbf{n}| = n_1 + \dots + n_j + \dots + n_t;$$

accordingly, $\mathbf{N} = (N_1, \dots, N_t)$, $\mathbf{1} = (1, \dots, 1)$. Also, we set

$$\mathbf{i} \leq \mathbf{n} \quad \text{if } i_j \leq n_j, \quad j = 1, \dots, t.$$

Then, for $\mathbf{n} < \mathbf{N}$, we deduce from the quantum binomial formula that

$$\Delta(x^{\mathbf{n}}) = x^{\mathbf{n}} \otimes 1 + 1 \otimes x^{\mathbf{n}} + \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}},$$

where $c_{\mathbf{n}, \mathbf{i}} \neq 0$ for all \mathbf{i} . We shall need

$$\begin{aligned} \Delta(x_j x_i) &= (x_j \otimes 1 + 1 \otimes x_j)(x_i \otimes 1 + 1 \otimes x_i) \\ &= x_j x_i \otimes 1 + x_j \otimes x_i + \chi_i(g_j) x_i \otimes x_j + 1 \otimes x_j x_i, \end{aligned} \tag{24}$$

for $1 \leq j, i \leq t$.

We shall prove by induction on r that the set

$$\{x^{\mathbf{n}} \mid |\mathbf{n}| \leq r, \quad \mathbf{n} < \mathbf{N}\}$$

is linearly independent.

Let $r = 1$ and let $a_0 + \sum_{i=1}^t a_i x_i = 0$, with $a_j \in k$, $0 \leq j \leq t$. Applying ϵ , we see that $a_0 = 0$; by hypothesis we conclude that the other a_j 's are also 0.

Now let $r > 1$ and suppose that $z = \sum_{|\mathbf{n}| \leq r} a_{\mathbf{n}} x^{\mathbf{n}} = 0$. Applying ϵ , we see that $a_0 = 0$. Then

$$\begin{aligned} 0 &= \Delta(z) = z \otimes 1 + 1 \otimes z + \sum_{1 \leq |\mathbf{n}| \leq r} a_{\mathbf{n}} \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}} \\ &= \sum_{1 \leq |\mathbf{n}| \leq r} \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} a_{\mathbf{n}} c_{\mathbf{n}, \mathbf{i}} x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}}. \end{aligned}$$

Now, if $|\mathbf{n}| \leq r$, $0 \leq \mathbf{i} \leq \mathbf{n}$, and $0 \neq \mathbf{i} \neq \mathbf{n}$, then $|\mathbf{i}| < r$ and $|\mathbf{n} - \mathbf{i}| < r$. By inductive hypothesis, the elements $x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}}$ are linearly independent. Hence $a_{\mathbf{n}} c_{\mathbf{n}, \mathbf{i}} = 0$ and $a_{\mathbf{n}} = 0$ for all \mathbf{n} , $|\mathbf{n}| \geq 1$. Thus $a_{\mathbf{n}} = 0$ for all \mathbf{n} . \square

Assume that $(G, g_i, \chi_i; j \in J)$ is an *ESC*. Let $\mathcal{T}(G, g_i, \chi_i; j \in J)$ be the free algebra generated by set $\{x_j \mid j \in J\}$. Let $\mathcal{S}(G, g_i, \chi_i; j \in J)$ be the algebra generated by set $\{x_j \mid j \in J\}$ with relations

$$x_i x_j = \chi_j(g_i) x_j x_i \quad \text{for any } i, j \in J \text{ with } i \neq j. \tag{25}$$

Let $\mathcal{L}(G, g_i, \chi_i; j \in J)$ be the algebra generated by set $\{x_j \mid j \in J\}$ with relations

$$x_l^{N_l} = 0, \quad x_i x_j = \chi_j(g_i) x_j x_i \quad \text{for any } i, j, l \in J \text{ with } N_l < \infty, i \neq j. \quad (26)$$

Define coalgebra operations and kG -(co-)module operations in $\mathcal{T}(G, g_i, \chi_i; j \in J)$, $\mathcal{S}(G, g_i, \chi_i; j \in J)$ and $\mathcal{L}(G, g_i, \chi_i; j \in J)$ as follows:

$$\Delta x_i = x_i \otimes 1 + 1 \otimes x_i, \quad \epsilon(x_i) = 0,$$

$$\delta^-(x_i) = g_i \otimes x_i, \quad h \cdot x_i = \chi_i(h) x_i.$$

$\mathcal{T}(G, g_i, \chi_i; j \in J)$ is called a quantum tensor algebra in ${}^{kG}\mathcal{YD}$, $\mathcal{S}(G, g_i, \chi_i; j \in J)$ is called a quantum symmetric algebra in ${}^{kG}\mathcal{YD}$ and $\mathcal{L}(G, g_i, \chi_i; j \in J)$ is called a quantum linear space in ${}^{kG}\mathcal{YD}$. Note that when $ESC(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative with finite J and finite N_j for any $j \in J$, the definition of quantum linear space is the same as in [3, Lemma 3.4]. Obviously, if N_i is infinite for all $i \in J$, then $\mathcal{S}(G, g_i, \chi_i; j \in J) = \mathcal{L}(G, g_i, \chi_i; j \in J)$.

Let \mathbf{Z} and \mathbf{Z}^+ denote sets of integers and positive integers, respectively.

Theorem 7 *Assume that $ESC(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative. Let \prec be a total order of J . Then*

(i) *The multiple Taft algebra $kG[kQ_1, g_i, \chi_i; i \in J]$ has a k -basis*

$$\{g \cdot E_{\nu_1}^{m_1} \cdot E_{\nu_2}^{m_2} \cdots E_{\nu_t}^{m_t} \mid 0 \leq m_j < N_j; \nu_j \prec \nu_{j+1}, \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbf{Z}^+, g \in G\}.$$

Moreover, $kG[kQ_1, g_i, \chi_i; i \in J]$ is finite dimensional if and only if $|G|$, $|J|$ and N_j are finite for any $j \in J$. In this case, $\dim_k(kG[kQ_1, g_i, \chi_i; i \in J]) = |G|N_1N_2 \cdots N_t$ with $J = \{1, 2, \dots, t\}$.

(ii) *diag($kG[kQ_1, g_i, \chi_i; i \in J]$) has a k -basis*

$$\{E_{\nu_1}^{m_1} \cdot E_{\nu_2}^{m_2} \cdots E_{\nu_t}^{m_t} \mid 0 \leq m_j < N_j; \nu_j \prec \nu_{j+1}, \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbf{Z}^+\}. \quad (27)$$

(iii) $\mathcal{L}(G, g_i, \chi_i^{-1}; j \in J) \cong \text{diag}(kG[kQ_1, g_i, \chi_i; i \in J])$ as graded braided Hopf algebras in ${}^{kG}\mathcal{YD}$, by sending x_j to $a_{g_j, 1}^{(j)}$ for any $j \in J$.

(iv) $\mathcal{T}(G, g_i, \chi_i^{-1}; j \in J) \cong \text{diag}(kQ^s(g_i, \chi_i; i \in J))$ as graded braided Hopf algebras in ${}^{kG}\mathcal{YD}$ algebras, by sending x_j to $a_{g_j, 1}^{(j)}$ for any $j \in J$.

(v) $kQ^s(g_i, \chi_i; i \in J) \cong \mathcal{T}(G, g_i, \chi_i^{-1}; j \in J) \# kG$ as graded Hopf algebras and $kQ^s(g_i, \chi_i; i \in J)$ has a k -basis

$$\{g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} \mid \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbf{Z}^+ \cup \{0\}, g \in G\},$$

where $g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} = g$ when $t = 0$.

Note that (iv) and (v) hold without quantum weakly commutative condition.

Proof. (ii) Since $(N_j)!_{q_j} = 0$, it follows from Lemma 3.4 that $E_j^{N_j} = 0$ when $N_j < \infty$. By Lemma 3.3, $E_i \cdot E_j = \chi_j(g_i^{-1})E_j \cdot E_i$ for any $i, j \in J$ with $i \neq j$. Considering Lemma 3.5, we complete the proof.

(iii) By Lemma 3.4 and Eq.(23), there exists an algebra homomorphism ψ from $\mathcal{L}(G, g_i, \chi_i; j \in J)$ to $\text{diag}(kG[kQ_1, g_i, \chi_i; i \in J])$ by sending x_j to $a_{g_j,1}^{(j)}$ for any $j \in J$. By (ii), ψ is bijective.

(iv) and (v). Let $A := \mathcal{T}(G, g_i, \chi_i^{-1}; j \in J)$ and $R = \text{diag}(kQ^s(g_i, \chi_i; i \in J))$. Let ψ be an algebra homomorphism from $\mathcal{T}(G, g_i, \chi_i; j \in J)$ to $\text{diag}(kQ^s, g_i, \chi_i; i \in J)$ by sending x_j to $a_{g_j,1}^{(j)}$.

It is clear that $\mathcal{T}(G, g_i, \chi_i; j \in J)$ is a kG -module algebra. Define a linear map ν from $A \# kG$ to kQ^s by sending $x_j \# g$ to $a_{g_j,1}^{(j)} \cdot g = \alpha^+(a_{g_j,1}^{(j)} \otimes g)$ for any $g \in G, j \in J$. That is, ν is the composition of

$$A \# kG \xrightarrow{\psi \otimes id} R \# kG \xrightarrow{\alpha^+} kQ^s,$$

where $\alpha^+ = \mu_{kQ^s}(id \otimes \iota_0)$ (see Theorem 6). Define a linear map λ from kG to $A \# kG$ by sending g to $1 \# g$ for any $g \in G$ and another linear map γ from kQ_1 to $A \# kG$ by sending $a_{g_i h, h}^{(i)}$ to $\chi_i^{-1}(h)x_i \# h$ for any $h \in G, i \in J$. It is clear that γ is a kG -bimodule homomorphism from kQ_1 to ${}_{\lambda}(A \# kG)_{\lambda}$. Considering $kQ^s = T_{kG}(kQ_1)$ and universal property of tensor algebra over kG , we have that there exists an algebra homomorphism $\phi = T_{kG}(\lambda, \gamma)$ from kQ^s to $A \# kG$. Obviously, ϕ is the inverse of ν . Thus ϕ is bijective. It is easy to check that ϕ is graded Hopf algebra isomorphism. Obviously, $\{x_{\nu_1} x_{\nu_2} \cdots x_{\nu_t} \# g \mid \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbf{Z}^+ \cup \{0\}, g \in G\}$ is a basis of $\mathcal{T}(G, g_i, \chi_i; j \in J) \# kG$. See

$$\begin{aligned} & \nu(x_{\nu_1} x_{\nu_2} \cdots x_{\nu_t} \# g) \\ &= E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} \cdot g \\ &= \chi_{\nu_1}(g) \chi_{\nu_2}(g) \cdots \chi_{\nu_t}(g) g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t}. \end{aligned}$$

Thus $\{g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} \mid \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbf{Z}^+ \cup \{0\}, g \in G\}$ is a basis of kQ^s . It is easy to check that ψ is graded braided Hopf algebra isomorphism.

(i) Considering (ii), Lemma 3.2 and Theorem 6, we complete the proof. \square

We now recall the Hopf algebra defined in [11, Definition 5.6.8 and 5.6.15].

Definition 3.6 For a quantum weakly commutative ESC $(G, g_i, \chi_i; i \in J)$ with $J = \{1, 2, \dots, t\}$ and commutative group G , Let A_t be the Hopf algebra, which is generated by $\{g, X_j \mid g \in G, j = 1, 2, \dots, t\}$ as algebras, where

- (i) elements of group G are abelian group-like elements of A_t ;
- (ii) X_j is $(1, g_j)$ -primitive, i.e., $\Delta(X_j) = X_j \otimes 1 + g_j \otimes X_j, 1 \leq j \leq t$;
- (iii) $X_j g = \chi_j(g) g X_j, 1 \leq j \leq t, g \in G$;
- (iv) $X_j X_i = \chi_j(g_i) X_i X_j$, where $1 \leq i, j \leq t$ with $i \neq j$.

Furthermore, assume that $J(a)$ is the ideal of A_t generated by

$$\{X_1^{N_1}, X_2^{N_2}, \dots, X_t^{N_t}\}.$$

Then we denote by $H(G, g_i, \chi_i; i \in J)$ the quotient algebra $A_t/J(a)$.

Remark: $H(G, g_i, \chi_i; i \in J)$ just is $H(C, n, c, c^*, 0, 0)$ in [11, Definition 5.6.8 and Definition 5.6.15] with $G = C$, $J = \{1, 2, \dots, t\}$, $n_i = N_i < \infty$, $g_i = c_i, c_i^* = \chi_i$ for $i = 1, 2, \dots, t$.

Proposition 3.7 For a quantum weakly commutative ESC($G, g_i, \chi_i; i \in J$) with $J = \{1, 2, \dots, t\}$ and commutative group G , we have that $H(G; g_i, \chi_i; i \in J)$ and $kG[kQ_1, g_i, \chi_i; i \in J]$ are isomorphic as graded Hopf algebras.

Proof. Let $H := H(G; g_i, \chi_i; i \in J)$. H is a graded Hopf algebra under ordinary gradation similar to polynomial algebras. By the method similar to the proof of Lemma 3.2(ii), we can obtain that $H^{co\ kG}$ is a subalgebra of H generated by X_i 's. By Theorem 6, $H^{co\ kG}$ is a braided Hopf algebra in ${}^{kG}\mathcal{YD}$ and H is the bosonization of $H^{co\ kG}$. Considering Theorem 7 (iii), we only need to show that $\mathcal{L}(G, g_i, \chi_i^{-1}; j \in J) \cong H^{co\ kG}$ as graded braided Hopf algebras. It follows from [11, Proposition 5.6.14] that $\{X_1^{m_1} X_2^{m_2} \dots X_t^{m_t} \mid 0 \leq m_j < N_j, j = 1, 2, \dots, t\}$ is a basis of $H^{co\ kG}$. Consequently, we have an algebra isomorphism ψ from $\mathcal{L}(G, g_i, \chi_i; i \in J)$ to $H^{co\ kG}$ by sending x_i to X_i . It is easy to check that ψ is also a graded braided Hopf algebra isomorphism. \square

Let B be a Hopf algebra and R a braided Hopf algebra in ${}^B\mathcal{YD}$. For convenience, we denote $r\#1$ by r and $1\#b$ by b in biproduct $R\#B$ for any $r \in R$, $b \in B$.

Lemma 3.8 Let B be a Hopf algebra and R a braided Hopf algebra in ${}^B\mathcal{YD}$ with $x_1, x_2 \in P(R)$ and $g_1, g_2 \in Z(G(B))$. Assume that there exist $\chi_1, \chi_2 \in \text{Alg}(B, k)$ with $\sqrt{\chi_i(g_j)} \in k$ such that

$$\delta_R(x_j) = g_j \otimes x_j, \quad h \cdot x_j = \chi_j(h^{-1})x_j,$$

for all $h \in B, j = 1, 2$.

(i) If r is a positive integer and

$$\chi_2(g_1)\chi_1(g_2)\chi_1(g_1)^{r-1} = 1, \quad \chi_1(g_1)^{\frac{1}{2}(r-1)}\chi_2(g_1) = 1, \quad r - 1 < \text{ord}(\chi_1(g_1)), \quad (28)$$

then

$$\sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix} \sqrt{\chi_1(g_1^{-1})} x_1^{r-m} x_2 x_1^m \quad (29)$$

$= (ad_c x_1)^r x_2$ is a primitive element of R and a $(1, g_1^r g_2)$ -primitive element of biproduct $R \# B$, where $(ad_c x_1)x_2 = x_1 x_2 - \chi_2(g_1^{-1})x_2 x_1$.

(ii) If $\sqrt{\chi_1(g_2)}\sqrt{\chi_2(g_1)} = 1$ and $x_i g_j = \chi_i(g_j)g_j x_i$ for $i, j = 1, 2$, then

$$\sqrt{\chi_2(g_1)}x_1 x_2 - \sqrt{\chi_1(g_2)}x_2 x_1$$

is a primitive element of R and

$$\sqrt{\chi_2(g_1)}x_1 x_2 - \sqrt{\chi_1(g_2)}x_2 x_1 - \beta(g_1 g_2 - 1)$$

is a $(1, g_1 g_2)$ -primitive element of biproduct $R \# B$ for any $\beta \in k$.

Proof. (i) Let $z := \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} x_1^{r-m} x_2 x_1^m$. By [5, Lemma A.1], $(ad_c x_1)^r x_2$ is primitive. However,

$$\begin{aligned} & (ad_c x_1)^r x_2 \\ &= \sum_{m=0}^r (-1)^m \begin{pmatrix} r \\ m \end{pmatrix}_{\chi_1(g_1^{-1})} \chi_1(g_1^{-1})^{\frac{1}{2}m(m-1)} \chi_2(g_1^{-1})^m x_1^{r-m} x_2 x_1^m \quad (\text{by [5, (A.8)]}) \\ &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} \chi_1(g_1^{-1})^{\frac{1}{2}m(r-1)} \chi_2(g_1^{-1})^m x_1^{r-m} x_2 x_1^m \quad (\text{by [5, P40]}) \\ &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} x_1^{r-m} x_2 x_1^m \quad (\text{by assumption (28)}). \end{aligned}$$

Thus z is primitive in R , i.e.

$$\Delta_R(z) = z \otimes 1 + 1 \otimes z. \quad (30)$$

Since

$$\begin{aligned} \delta_R(z) &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} \delta_R(x_1^{r-m} x_2 x_1^m) \\ &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} g_1^r g_2 \otimes x_1^{r-m} x_2 x_1^m \\ &= g_1^r g_2 \otimes z, \end{aligned}$$

we have

$$\begin{aligned}
\Delta_{R\#B}(z) &= \Delta_{R\#B}(z\#1) \quad (\text{by Theorem 6}) \\
&= (id_R \otimes \mu_B \otimes id_R \otimes id_B)(id_R \otimes id_B \otimes C_{R,B} \otimes id_B) \\
&\quad (id_R \otimes \delta_R \otimes \Delta_B)(1\#z \otimes 1) + z \otimes 1 \\
&= g_1^r g_2 \otimes z + z \otimes 1,
\end{aligned}$$

where $C_{R,B}$ denotes the ordinary twist map from $R \otimes B$ to $B \otimes R$ by sending $r \otimes b$ to $b \otimes r$ for any $r \in R, b \in B$.

(ii) See

$$\begin{aligned}
&\Delta_{R\#B}(\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \\
&= \sqrt{\chi_2(g_1)}(g_1g_2 \otimes x_1x_2 + x_1g_2 \otimes x_2 + g_1x_2 \otimes x_1 + x_1x_2 \otimes 1) \\
&\quad - \sqrt{\chi_1(g_2)}(g_1g_2 \otimes x_2x_1 + x_2g_1 \otimes x_1 + g_2x_1 \otimes x_2 + x_2Ei \otimes 1) \\
&= g_1g_2 \otimes (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) + (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \otimes 1 \\
&\quad (\text{by assumption of (ii)})
\end{aligned}$$

and

$$\begin{aligned}
&\Delta_{R\#B}(\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) - \Delta_{R\#B}(\beta(g_1g_2 - 1)) \\
&= g_1g_2 \otimes (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1 - \beta(g_1g_2 - 1)) \\
&\quad + (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1 - \beta(g_1g_2 - 1)) \otimes 1.
\end{aligned}$$

Thus $\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1 - \beta(g_1g_2 - 1)$ is a $(1, g_1g_2)$ -primitive element of $R\#B$. See

$$\begin{aligned}
&\Delta_R(\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \\
&= \sqrt{\chi_2(g_1)}(1 \otimes x_1x_2 + x_1 \otimes x_2 + \chi_2(g_1)^{-1}x_2 \otimes x_1 + x_1x_2 \otimes 1) \\
&\quad - \sqrt{\chi_1(g_2)}(1 \otimes x_2x_1 + x_2 \otimes x_1 + \chi_1(g_2)^{-1}x_1 \otimes x_2 + x_2 \otimes 1) \\
&= 1 \otimes (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) + (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \otimes 1 \\
&\quad (\text{by assumption of (ii)}).
\end{aligned}$$

Thus $\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1$ is a primitive element of R . \square

Let N be a set and $J = \cup_{s \in N} (J_s \cup J'_s)$ a disjoint union. Let $J^{(1)} := \cup_{u \in N} J_u$ and $J^{(2)} := \cup_{u \in N} J'_u$. There exists a bijection $\sigma : J^{(1)} \rightarrow J^{(2)}$ such that $\sigma|_{J_u}$ is a bijection from J_u to J'_u for $u \in N$. Let G be a free commutative group with generators $\{\xi_i \mid i \in J^{(1)}\}$. Put $\chi_{\sigma(i)}(\xi_j) = \chi_i^{-1}(\xi_j)$, $\xi_{\sigma(i)} := \xi_i^{-1}$ and $g_i = g_{\sigma(i)} := \xi_i^2$ for any $i, j \in J_u$, $u \in N$.

$(G, g_i, \chi_i; i \in J)$ is an *ESC* and r_{ij} is a positive integer with $r_{\sigma(i), \sigma(j)} = r_{ij}$ for any $i, j \in J^{(1)}$ with $i \neq j$. Let $\text{ord}(\chi_i(g_i))$ denote the order of $\chi_i(g_i)$.

Let I be the ideal of $kQ^s(g_i, \chi_i; i \in J)$ generated by the following elements:

$$\chi_j(\xi_i)E_iE_j - \chi_i^{-1}(\xi_j)E_jE_i - \delta_{\sigma(i), j} \frac{g_i^2 - 1}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}, \text{ for } i \in J_u, j \in J'_u, u \in N; \quad (31)$$

$$\sum_{m=0}^{r_{ij}} (-1)^m \begin{bmatrix} r_{ij} \\ m \end{bmatrix}_{\chi_i(\xi_i^{-1})} E_i^{r_{ij}-m} E_j E_i^m, \quad (32)$$

for any $i, j \in J_u$ or $i, j \in J'_u, i \neq j$ and $r_{ij} - 1 < \text{ord}(\chi_i(g_i)); u \in N$.

Let U be an algebra generated by $\{K_i, X_i \mid i \in J\}$ with relations

$$\begin{aligned} X_i X_j - X_j X_i &= \delta_{\sigma(i), j} \frac{K_i^2 - K_i^{-2}}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}, \text{ for any } i \in J_u, j \in J'_u; u \in N; \\ K_i K_{\sigma(i)} &= K_{\sigma(i)} K_i = 1 \text{ for any } i \in J^{(1)}; \\ \chi_j(\xi_i) K_i X_j &= X_j K_i, \quad K_i K_j - K_j K_i = 0, \text{ for any } i, j \in J; \\ \sum_{m=0}^{r_{ij}} (-1)^m \begin{bmatrix} r_{ij} \\ m \end{bmatrix}_{\chi_i(\xi_i^{-1})} X_i^{r_{ij}-m} X_j X_i^m &= 0, \end{aligned} \quad (33)$$

for $i, j \in J_u$ or $i, j \in J'_u, i \neq j$ and $r_{ij} - 1 < \text{ord}(\chi_i(g_i)); u \in N$.

The comultiplication, counit and antipode of U are defined by

$$\begin{aligned} \Delta(X_j) &= X_j \otimes K_{\sigma(j)} + K_j \otimes X_j, & S(X_i) &= -\chi_i(\xi_i) X_i, & \epsilon(X_i) &= 0, \\ \Delta(K_i) &= K_i \otimes K_i, & S(K_j) &= K_{\sigma(j)}, & \epsilon(K_i) &= 1, \\ \Delta(X_{\sigma(j)}) &= X_{\sigma(j)} \otimes K_{\sigma(j)} + K_j \otimes X_{\sigma(j)}, \end{aligned}$$

for any $j \in J^{(1)}, i \in J$.

In fact, $K_{\sigma(j)} = K_j^{-1}$ in U for any $j \in J^{(1)}$.

Theorem 8 *Under notation above, Assume*

$$\chi_j(g_i) \chi_i(g_j) \chi_i(g_i)^{r_{ij}-1} = 1, \quad \chi_i(\xi_j) = \chi_j(\xi_i), \quad \chi_i(g_i)^{\frac{1}{2}(r_{ij}-1)} \chi_j(g_i) = 1, \quad (34)$$

for any $i, j \in J_u, i \neq j, u \in N$. Then $kQ^s(g_i, \chi_i; i \in J)/I \cong U$ as Hopf algebras.

Proof. For any $i, j \in J_u, u \in N$, see $\chi_{\sigma(i)}(\xi_{\sigma(j)}) = \chi_i(\xi_j) = \chi_j(\xi_i) = \chi_{\sigma(j)}(\xi_{\sigma(i)})$, $\chi_{\sigma(i)}(\xi_j) = \chi_i^{-1}(\xi_j) = \chi_j^{-1}(\xi_i) = \chi_j(\xi_{\sigma(i)})$, $\chi_i(\xi_{\sigma(j)}) = \chi_i(\xi_j)^{-1} = \chi_j(\xi_i)^{-1} = \chi_{\sigma(j)}(\xi_i)$. Therefore,

$$\chi_i(\xi_j) = \chi_j(\xi_i) \quad (35)$$

For any $i, j \in J_u \cup J'_u, u \in N$. Obviously, for $i, j \in J'_u, i \neq j, u \in N$, (34) holds.

We show this theorem by following several steps.

(i) There is a algebra homomorphism Φ from kQ^s to U such that $\Phi(\xi_i) = K_i$ and $\Phi(a_{hg_i,h}^{(i)}) = \Phi(h)K_iX_i$ and $\Phi(a_{hg_{\sigma(i)},h}^{(\sigma(i))}) = \Phi(h)K_iX_{\sigma(i)}$ for all $h \in G$ and $i \in J^{(1)}$. Indeed, define algebra homomorphism $\phi : kG \rightarrow U$ given by $\phi(\xi_i) = K_i$ for $i \in J$ and a k -linear map $\psi : kQ_1 \rightarrow U$ by $\psi(a_{gg_j,g}^{(j)}) = \phi(g)K_jX_j$ and $\psi(a_{gg_{\sigma(j)},g}^{(\sigma(j))}) = \phi(g)K_jX_{\sigma(j)}$ for any $j \in J^{(1)}$, $g \in G$. For any $g, h \in G, j \in J^{(1)}$, see

$$\begin{aligned}\psi(h \cdot a_{gg_j,g}^{(j)}) &= \psi(a_{hgg_j,hg}^{(j)}) \\ &= \phi(hg)K_jX_j \\ &= \phi(h)\phi(g)K_jX_j \\ &= \phi(h)\psi(a_{gg_j,g}^{(j)}),\end{aligned}$$

and

$$\begin{aligned}\psi(a_{gg_j,g}^{(j)} \cdot h) &= \chi_j(h)\psi(a_{hgg_j,hg}^{(j)}) \\ &= \chi_j(h)\phi(hg)K_jX_j \\ &= \phi(g)K_jX_j\phi(h) \quad (\text{since } X_j\phi(h) = \phi(h)\chi_j(h)X_j) \\ &= \psi(a_{gg_j,g}^{(j)})\phi(h).\end{aligned}$$

Similarly, $\psi(h \cdot a_{gg_{\sigma(j)},g}^{(\sigma(j))}) = \phi(h)\psi(a_{gg_{\sigma(j)},g}^{(\sigma(j))})$ and $\psi(a_{gg_{\sigma(j)},g}^{(\sigma(j))} \cdot h) = \psi(a_{gg_{\sigma(j)},g}^{(\sigma(j))})\phi(h)$. This implies that ψ is a kG -bimodule map from $(kQ_1; g_i, \chi_i; i \in J)$ to ${}_{\phi}U_{\phi}$. Using the universal property of tensor algebra over kG , we complete the proof.

(ii) $\Phi(I) = 0$. For any $i \in J_u, j \in J'_u$, see that

$$\begin{aligned}&\Phi(\chi_j(\xi_i)E_iE_j - \chi_i(\xi_j)^{-1}E_jE_i - \delta_{\sigma(i),j} \frac{g_i^2-1}{\chi_i(\xi_i)-\chi_i(\xi_i)^{-1}}) \\ &= \chi_j(\xi_i)K_iX_iK_j^{-1}X_j - \chi_i(\xi_j)^{-1}K_j^{-1}X_jK_iX_i - \delta_{\sigma(i),j} \frac{K_i^4-1}{\chi_i(\xi_i)-\chi_i(\xi_i)^{-1}} \\ &= K_iK_j^{-1}X_iX_j - K_j^{-1}K_iX_jX_i - \delta_{\sigma(i),j} \frac{K_i^4-1}{\chi_i(\xi_i)-\chi_i(\xi_i)^{-1}} \\ &= K_iK_j^{-1}(X_iX_j - X_jX_i - \delta_{\sigma(i),j} \frac{K_i^3K_j - K_i^{-1}K_j}{\chi_i(\xi_i)-\chi_i(\xi_i)^{-1}}) \\ &= K_iK_j^{-1}(X_iX_j - X_jX_i - \delta_{\sigma(i),j} \frac{K_i^2 - K_i^{-2}}{\chi_i(\xi_i)-\chi_i(\xi_i)^{-1}}) \\ &= 0.\end{aligned}$$

For $i, j \in J_u, i \neq j$, see that

$$\begin{aligned}&\Phi(E_i^{r_{ij}-m}E_jE_i^m) \\ &= (K_iX_i)^{r_{ij}-m}K_jX_j(K_iX_i)^m \\ &= \chi_i(\xi_i)^{\frac{1}{2}((r_{ij}-m)(r_{ij}-m-1)+m(m-1)+2(r_{ij}-m)m)}\chi_i(\xi_j)^{r_{ij}-m}\chi_j(\xi_i)^mX_i^{r_{ij}-m}X_jX_i^m \\ &= \chi_i(\xi_i)^{\frac{1}{2}(r_{ij}-1)r_{ij}}\chi_i(\xi_j)^{r_{ij}}X_i^{r_{ij}-m}X_jX_i^m \quad (\text{by assumption (34)})\end{aligned}$$

and

$$\Phi\left(\sum_{m=0}^{r_{ij}} (-1)^m \begin{bmatrix} r_{ij} \\ m \end{bmatrix}_{\chi_i(\xi_i^{-1})} E_i^{r_{ij}-m}E_jE_i^m\right) = 0.$$

Similarly, the equation above holds for $i, j \in J'_u, i \neq j, u \in N$.

By (ii), there exists an algebra homomorphism $\bar{\Phi}$ from kQ^s/I to U such that $\bar{\Phi}(x+I) = \Phi(x)$ for any $x \in kQ^s$. For convenience, we will still use x to denote $x + I$.

(iii) It follows from the definition of U that there exists a unique algebra map $\Psi : U \rightarrow kQ^s/I$ such that $\Psi(K_i) = \xi_i$, $\Psi(X_i) = \xi_i^{-1}E_i$ and $\Psi(X_{\sigma(i)}) = \xi_i^{-1}E_{\sigma(i)}$ for all $i \in J^{(1)}$. It is easy to see that $\Psi\bar{\Phi} = id$ and $\bar{\Phi}\Psi = id$.

(iv) We show that I is a Hopf ideal of kQ^s . It follows from Theorem 7 that $R := \text{diag}(kQ^s)$ is a braided Hopf algebra with $\delta^-(E_i) = g_i \otimes E_i$, $h \triangleright E_i = \chi_i^{-1}(h)E_i$ for any $h \in G$, $i \in J$. By Theorem 6, $R\#B \cong kQ^s$ as Hopf algebras. It follows from Lemma 3.8 that I is a Hopf ideal of kQ^s .

Obviously, Ψ preserve the comultiplication and counit. Since kQ^s/I is a Hopf algebra, then Ψ is a Hopf algebra isomorphism. \square

Denote by $kQ^s(g_i, \chi_i; i \in J; r_{ij}; J_u, J'_u; u \in N)$ the semi-path Hopf algebra kQ^s above. Let $\overline{kQ^s}(g_i, \chi_i; i \in J; r_{ij}; J_u, J'_u; u \in N) = kQ^s(g_i, \chi_i; i \in J; r_{ij}; J_u, J'_u; u \in N)/I$ and $U(g_i, \chi_i; i \in J; r_{ij}; J_u, J'_u; u \in N) = U$.

Example 3.9 Under the notation above, let $A = (a_{ij})$ be a symmetrizable $J^{(1)} \times J^{(1)}$ -matrix with $a_{ii} = 2$ and non-positive integer a_{ij} for any $i, j \in J^{(1)}$ and $i \neq j$. There exists integer d_i such that $d_i a_{ij} = d_j a_{ji}$ for any $i, j \in J^{(1)}$. Put $\chi_{\sigma(i)}(\xi_j) = \chi_i^{-1}(\xi_j)$, $\xi_{\sigma(i)} := \xi_i^{-1}$ and $g_i = g_{\sigma(i)} := \xi_i^2$ for any $i, j \in J_u$, $u \in N$. Define $r_{ij} = 1 - a_{ij}$, $\chi_i(\xi_j) = q^{-d_i a_{ij}}$ for $i, j \in J_u$, $u \in N$. It is easy to check the conditions (34) are satisfied, and hence Theorem 8 holds. In this case, we write the three Hopf algebras as $kQ^s(g_i, \chi_i; i \in J; A; d_i; J_u, J'_u; u \in N)$, $\overline{kQ^s}(g_i, \chi_i; i \in J; A; d_i; J_u, J'_u; u \in N)$, and $U(g_i, \chi_i; i \in J; A; d_i; J_u, J'_u; u \in N)$, respectively. If $A = (a_{ij})$ is a symmetrizable generalized Cartan matrix with $|N| = 1$ and $J^{(1)} = \{1, 2, \dots, n\}$, then the three Hopf algebras above is denoted by $kQ^s(g_i, \chi_i; i \in J; A; d_i)$, $\overline{kQ^s}(g_i, \chi_i; i \in J; A; d_i)$ and $U(g_i, \chi_i; i \in J; A; d_i)$, respectively. In particular, if k is the complex field, q is not a root of unit and A is a symmetrizable Cartan matrix with $d_i \in \{1, 2, 3\}$, then U is the quantum enveloping algebra $U_q(L)$ of L , where L is the semisimple Lie algebra determined by A (see [19, p.218] or [16, P. 90]). Furthermore, $U_q(L)$ is isomorphic, as a Hopf algebra, to $kQ^s(g_i, \chi_i; i \in J)/I = \overline{kQ^s}(g_i, \chi_i; i \in J; A; d_i)$.

An $ESC(G, g_i, \chi_i; i \in J)$ is said to be a local FL-type (see [5, P.4]) if the following conditions are satisfied:

- N is a set and $J = \cup_{s \in N} (J_s \cup J'_s)$ is a disjoint union.
- There exists a bijection $\sigma : J^{(1)} \rightarrow J^{(2)}$ such that $\sigma|_{J_u}$ is a bijection from J_u to J'_u for $u \in N$, where $J^{(1)} := \cup_{u \in N} J_u$ and $J^{(2)} := \cup_{u \in N} J'_u$.
- There exists a $J^{(1)} \times J^{(1)}$ -matrix $A = (a_{ij})$ with $a_{ii} = 2$ and non-positive integer a_{ij} for any $i, j \in J^{(1)}$ and $i \neq j$. For any $u \in N$, there exists an integer $d_i^{(u)}$ such that $d_i^{(u)} a_{ij} = d_j^{(u)} a_{ji}$ for any $i, j \in J_u$.

• For any $u \in N$, there exists $0 \neq q_u \in k$ such that $\chi_i(g_j) = q_u^{-2d_i^{(u)} a_{ij}}$, $\chi_{\sigma(i)}(g_j) = \chi_i^{-1}(g_j)$ and $g_{\sigma(j)} = g_j$ for $i, j \in J_u$.

In addition. if A is a generalized Cartan matrix, then $ESC(G, g_i, \chi_i; i \in J)$ is said to be a local FL Cartan-type. If N only contains one element and $J^{(1)} = \{1, 2, \dots, n\}$, then $ESC(G, g_i, \chi_i; i \in J)$ is said to be a FL -type.

Obviously, $ESC(G, g_i, \chi_i; i \in J)$ in Example 3.9 is a local FL-type. By Lemma 3.8 and Theorem 7, we have

Example 3.10 *Under the notations above, assume that $ESC(G, g_i, \chi_i; i \in J)$ is a local FL-type.*

(i) *Let I be the ideal of $kQ^s(g_i, \chi_i; i \in J)$ generated by the following elements:*

$$\sqrt{\chi_j(g_i)} E_i E_j - \sqrt{\chi_i(g_j)} E_j E_i - \delta_{\sigma(i), j} \frac{g_i^2 - 1}{\sqrt{\chi_i(g_i)} - \sqrt{\chi_i(g_i)^{-1}}}, \text{ for } i \in J_u, j \in J'_u, u \in N;$$

$$\sum_{m=0}^{1-a_{ij}} (-1)^m \begin{bmatrix} 1 - a_{ij} \\ m \end{bmatrix}_{\sqrt{\chi_i(g_i)^{-1}}} E_i^{1-a_{ij}-m} E_j E_i^m,$$

for any $i, j \in J_u$ or $i, j \in J'_u, i \neq j$, and $-a_{ij} < \text{ord}(\chi_i(g_i)); u \in N$. Then I is a Hopf ideal of $kQ^s(g_i, \chi_i; i \in J)$.

(ii) *Let I be the ideal of $kQ^c(g_i, \chi_i; i \in J)$ generated by the following elements:*

$$\sqrt{\chi_j(g_i)} E_i \cdot E_j - \sqrt{\chi_i(g_j)} E_j \cdot E_i - \delta_{\sigma(i), j} \frac{g_i^2 - 1}{\sqrt{\chi_i(g_i)} - \sqrt{\chi_i(g_i)^{-1}}}, \quad (36)$$

for $i \in J_u, j \in J'_u, u \in N$;

$$\sum_{m=0}^{1-a_{ij}} (-1)^m \begin{bmatrix} 1 - a_{ij} \\ m \end{bmatrix}_{\sqrt{\chi_i(g_i)^{-1}}} E_i^{1-a_{ij}-m} \cdot E_j \cdot E_i^m, \quad (37)$$

for any $i, j \in J_u$ or $i, j \in J'_u, i \neq j$, and $a_{ij} < \text{ord}(\chi_i(g_i)); u \in N$. Then I is a Hopf ideal of $kQ^c(g_i, \chi_i; i \in J)$.

(iii) *Let I be the ideal of $kG[kQ_1, g_i, \chi_i; i \in J]$ generated by elements (36) and (37). Then I is a Hopf ideal of $kG[kQ_1, g_i, \chi_i; i \in J]$.*

Finally, we give an interesting quantum combinatoric formula by means of multiple Taft algebras. Let n be a positive integer and S_n be the symmetric group on the set $\{1, 2, \dots, n\}$. For any permutation $\sigma \in S_n$, let $\tau(\sigma)$ denote the number of reverse order of σ , i.e., $\tau(\sigma) = |\{(i, j) \mid 1 \leq i < j \leq n, \sigma(i) > \sigma(j)\}|$. For any $0 \neq q \in k$, let $S_n(q) := \sum_{\sigma \in S_n} q^{\tau(\sigma)}$.

Lemma 3.11 In $kQ^c(\alpha^\chi)$, assume $g \in \mathcal{K}_r(G)$ and $j \in I_{\{g\}}(r)$. Let $q := \chi_g^{(j)}(g)$. If i_1, i_2, \dots, i_m be non-negative integers, then

$$a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)} = q^{\beta_m + \frac{m(m-1)}{2}} S_m(q^{-1}) P_{g^{\alpha_m}}^{(j)}(g, m)$$

where $\alpha_m = i_1 + i_2 + \dots + i_m$, $P_h^{(j)}(g, m) = a_{g^m h, g^{m-1} h}^{(j)} a_{g^{m-1} h, g^{m-2} h}^{(j)} \cdots a_{g h, h}^{(j)}$, $\beta_1 = 0$ and $\beta_m = \sum_{j=1}^{m-1} (i_1 + i_2 + \dots + i_j)$ if $m > 1$.

Proof. Now let $a_l = a_{g^{i_l+1}, g^{i_l}}^{(j)}$ for $l = 1, 2, \dots, m$ and consider the sequence of m arrows $A = (a_m, \dots, a_1)$. We will use the notations of [9, p.247]. For any $\sigma \in S_m$ and $1 \leq l \leq m$, let $m(\sigma, l) = |\{i \mid i < l, \sigma(i) < \sigma(l)\}|$. Then

$$m(\sigma, l) = (l-1) - |\{i \mid i < l, \sigma(i) > \sigma(l)\}|,$$

and hence

$$\begin{aligned} \sum_{l=1}^m m(\sigma, l) &= \frac{m(m-1)}{2} - \sum_{l=1}^m |\{i \mid i < l, \sigma(i) > \sigma(l)\}| \\ &= \frac{m(m-1)}{2} - \tau(\sigma). \end{aligned}$$

Now it follows from [9, Proposition 3.13] that

$$\begin{aligned} a_m \cdot a_{m-1} \cdots \cdots a_1 &= \sum_{\sigma \in S_m} A_m^\sigma \cdots A_2^\sigma A_1^\sigma \\ &= \sum_{\sigma \in S_m} \left(\prod_{l=1}^m q^{i_1 + i_2 + \dots + i_{\sigma(l)-1} + m(\sigma, l)} \right) P_{g^{\alpha_m}}^{(j)}(g, m) \\ &= q^{\beta_m + \frac{m(m-1)}{2}} S_m(q^{-1}) P_{g^{\alpha_m}}^{(j)}(g, m). \quad \square \end{aligned}$$

Let $G \cong \mathbf{Z}$ be the infinite cyclic group with generator g and let r be a ramification of G given by $r_g = 1$ and $r_{g^n} = 0$ if $n \neq 1$. Let $0 \neq q \in k$. Define $\chi_g^{(1)} \in \widehat{G}$ by $\chi_g^{(1)}(g) = q$. Then $(G, r, \overrightarrow{\chi}, u)$ is an RSC. In Lemma 3.4 and Lemma 3.11, one gets the following result.

Example 3.12 (i) For any $0 \neq q \in k$, $(m)_q! = q^{\frac{m(m-1)}{2}} S_m(q^{-1})$, and $S_m(q) = (m)_\perp q^{\frac{m(m-1)}{2}}$, where m is a positive integer.

(ii) Assume q is a primitive n -th root of unity with $n > 1$. Then $S_m(q) = 0$ if $m \geq n$, and $S_m(q) \neq 0$ if $0 < m < n$.

Acknowledgement: We would like to thank Prof. N. Andruskiewitsch for comments and suggestions. The first two authors were financially supported by the Australian Research Council, and the 3rd author was supported by NSF of China (10471121). S.C.Z thanks the Department of Mathematics, University of Queensland for hospitality.

References

- [1] F. W. Anderson and K. R. Fuller, Rings and categories of modules, GTM **13**, Springer-Verlag, New York, 1992.
- [2] N. Andruskiewitsch and H. J. Schneider, Hopf algebras of order p^2 and braided Hopf algebras, J. Alg. **199** (1998), 430–454.
- [3] N. Andruskiewitsch and H. J. Schneider, Lifting of quantum linear spaces and pointed Hopf algebras of order p^3 , J. Alg. **209** (1998), 645–691.
- [4] N. Andruskiewitsch and H. J. Schneider, Pointed Hopf algebras, new directions in Hopf algebras, edited by S. Montgomery and H.J. Schneider, Cambridge University Press, 2002.
- [5] N. Andruskiewitsch and H. J. Schneider, Finite quantum groups and Cartan matrices, Adv. Math. **154** (2000), 1–45.
- [6] M. Auslander, I. Reiten and S. O. Smalø, Representation theory of Artin algebras, Cambridge University Press, 1995.
- [7] X. W. Chen, H. L. Huang, Y. Ye and P. Zhang, Monomial Hopf algebras, J. Alg. **275** (2004), 212–232.
- [8] W. Chin and S. Montgomery, Basic coalgebras, modular interfaces, AMS/IP Stud. Adv. Math. **4**, Amer. Math. Soc., Providence, RI, 1997, pp.41–47.
- [9] C. Cibils and M. Rosso, Hopf quivers, J. Alg. **254** (2002), 241–251.
- [10] C. Cibils and M. Rosso, Algebres des chemins quantiques, Adv. Math. **125** (1997), 171–199.
- [11] S. Dascalescu, C. Nastasecu and S. Raianu, Hopf algebras: an introduction, Marcel Dekker Inc., 2001.
- [12] Pavel Etingof, Shlomo Gelaki, The classification of finite-dimensional triangular Hopf algebras over an algebraically closed field of characteristic 0, Internat. Math. Res. Notices, 5(2000), 223–234.
- [13] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Math. J., **66** (1992), 123–168.
- [14] M. A. Hennings, Hopf algebras and regular isotopy invariants for link diagrams, Pro. Cambridge Phil. Soc **109** (1991), 59–77.

- [15] L. Kauffman, invariants of links and 3-manifolds via Hopf algebras, in *Geometry and Physics*, Marcel Dekker Lecture Notes in Pure and Appl. Math. vol 184, 1997, 471-479.
- [16] G. Lusztig, Quantum groups at roots of 1, *Geom. Dedicata*, **35** (1990), 89-114.
- [17] S. Majid, Physics for algebraists: non-commutative and non-cocommutative Hopf algebras by a bicross product construction, *J. Alg.* **130** (1990), 17–64.
- [18] S. Majid. Algebras and Hopf algebras in braided categories. Lecture notes in pure and applied mathematics advances in Hopf algebras, Vol. 158, edited by J. Bergen and S. Montgomery, 1995.
- [19] S. Montgomery, Hopf algebras and their actions on rings, CBMS no.82, AMS, Providence, RI, 1993.
- [20] W. Nichols, Bialgebras of type one, *Commun. Alg.* **6** (1978), 1521–1552.
- [21] F. Van Oystaeyen and P. Zhang, Quiver Hopf algebras, *J. Alg.* **280** (2004), 577–589.
- [22] D. E. Radford, The structure of Hopf algebras with a projection, *J. Alg.* **92** (1985), 322–347.
- [23] D. E. Radford, On Kauffman’s knot invariants arising from finite-dimensional Hopf algebras, in *Advances in Hopf Algebras*, Marcel Dekker Lecture Notes in Pure and Appl. Math vol 158 (1994), 205-266.
- [24] D.E.Radford, J.Towber, Yetter-Drinfeld categories associated to an arbitrary bialgebra, *J. Pure and Applied Algebra*, 87(1993) 259-279.
- [25] N. Yu. Reshetikhin and V.G. Turaev, Ribbon graphs and their invariants derived from quantum groups, *Commun. Math. Phys.* **127** (1990), 1–26.
- [26] M. E. Sweedler, *Hopf algebras*, Benjamin, New York, 1969.
- [27] W. Szymanski, Finite index subfactors and Hopf algebra crossed products, *Proc. AMS*, 120 (1994), 519-528.
- [28] S. L. Woronowicz, Differential calculus on compactmatrix pseudogroups (quantum groups), *Commun. Math. Phys.* 122 (1989), 125-170.