## The smallest eigenvalue of $K_r$ -free graphs

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis, Memphis TN 38152, USA

Submitted: 2004; Accepted: 200X; Published: 200X MR Subject Classifications: 05C35

## Abstract

Let G be a  $K_{r+1}$ -free graph with n vertices and m edges, and let  $\mu_n(G)$  be the smallest eigenvalue of its adjacency matrix. We show that

$$\mu_n(G) < -\frac{2^{r+1}m^r}{rn^{2r-1}}.$$

Our notation and terminology are standard (see, e.g. [1]). If G is a graph of order n, we write e(X) for the number of edges induced by a set  $X \subset V(G)$ ,  $\Gamma(u)$  for the set of neighbors of a vertex  $u \in V(G)$ , and  $\mu_n(G)$  for the smallest eigenvalue of the adjacency matrix of G.

Bollobás and Nikiforov [2] observed that if G is a dense  $K_r$ -free graph then  $\mu_n(G) < -\alpha n$  for some  $\alpha > 0$ , independent of n. In this note we prove the following more precise statement.

**Theorem 1** Suppose  $r \geq 2$  and G is a  $K_{r+1}$ -free graph with n vertices and m edges. Then

$$\mu_n(G) < -\frac{2^{r+1}m^r}{rn^{2r-1}}.$$

To prove this theorem we need two preliminary results. Our main tool will be the following inequality proved in [2]: If  $V(G) = V_1 \cup V_2$  is a bipartition of the vertices of a graph G then

$$\mu_n(G) \le \frac{2e(V_1)}{|V_1|} + \frac{2e(V_2)}{|V_2|} - \frac{2e(G)}{|V(G)|}.$$
(1)

Write t(G) for the number of triangles of a graph G and t''(G) for the number of its induced subgraphs of order 3 and size 1. For every vertex u, let  $t(u) = e(\Gamma(u))$  and  $t''(u) = e(V(G) \setminus \Gamma(u))$ . Observe the following simple equalities

$$3t(G) = \sum_{u \in V(G)} t(u)$$
 and  $t''(G) = \sum_{u \in V(G)} t''(u)$ .

Inequality (1) implies the following lemma.

**Lemma 2** Let G be a graph with n vertices and m edges with no isolated vertices. Then

$$\mu_n(G) \le \frac{2n}{(n^2 - 2m)} \sum_{u \in V(G)} \frac{t(u)}{d(u)} - \frac{4m^2}{n(n^2 - 2m)}.$$
(2)

**Proof** We start by recalling the equality

$$3t(G) = \sum_{u \in V(G)} d^2(u) - nm + t''(G)$$
(3)

THE ELECTRONIC JOURNAL OF COMBINATORICS XX (200X), #R00

1

whose proof we shall outline for convenience. For every edge  $uv \in E(G)$  we have

$$\left|\Gamma\left(u\right)\cap\Gamma\left(v\right)\right| = \left|\Gamma\left(u\right)\right| + \left|\Gamma\left(v\right)\right| - \left|\Gamma\left(u\right)\cup\Gamma\left(v\right)\right| = d\left(u\right) + d\left(v\right) - n + \left|\overline{\Gamma\left(u\right)}\cap\overline{\Gamma\left(v\right)}\right|.$$

Summing this equality over all  $uv \in E(G)$  we obtain

$$3t(G) = \sum_{uv \in E(G)} \left( d(u) + d(v) - n + \left| \overline{\Gamma(u)} \cap \overline{\Gamma(v)} \right| \right) = \sum_{u \in V(G)} d^2(u) - nm + t''(G),$$

as claimed.

For every  $u \in V(G)$  and bipartition  $V_1 = \Gamma(u), V_2 = V(G) \setminus \Gamma(u)$ , inequality (1) implies

$$\mu_n(G) \le \frac{2e(V_1)}{d(u)} + \frac{2e(V_2)}{n-d(u)} - \frac{2m}{n} = \frac{2t(u)}{d(u)} + \frac{2t''(u)}{n-d(u)} - \frac{2m}{n},$$

and therefore,

$$\mu_n(G)(n-d(u)) \le \frac{2t(u)}{d(u)}(n-d(u)) + 2t''(u) - \frac{2m}{n}(n-d(u)).$$

Summing this inequality for all  $u \in V(G)$ , in view of (3), we obtain

$$\mu_n(G)(n^2 - 2m) \le n \sum_{u \in V(G)} \frac{2t(u)}{d(u)} - 6t(G) + 2t''(G) - \frac{2m}{n}(n^2 - 2m)$$
$$= n \sum_{u \in V(G)} \frac{2t(u)}{d(u)} + 2nm - 2 \sum_{u \in V(G)} d^2(i) - \frac{2m}{n}(n^2 - 2m)$$
$$\le 2n \sum_{u \in V(G)} \frac{t(u)}{d(u)} - \sum_{u \in V(G)} d^2(u) \le 2n \sum_{u \in V(G)} \frac{t(u)}{d(u)} - \frac{4m^2}{n},$$

completing the proof.

**Proof of Theorem 1** Assume that G has no isolated vertices; the general case follows immediately. Our proof is by induction on r. If G is triangle-free, Lemma 2 implies

$$\mu_n(G) \le \frac{-4m^2}{n(n^2 - 2m)} < -\frac{4m^2}{n^3},$$

so the assertion holds for r = 2; assume it holds for  $r - 1 \ge 2$ . Since, for every  $u \in V(G)$ , the graph induced by  $\Gamma(u)$  is  $K_r$ -free, the induction hypothesis implies

$$\mu_n(G) < -\frac{2^r t^{r-1}(u)}{(r-1) d^{2r-3}(u)}.$$

Summing this inequality for all  $u \in V(G)$ , we obtain

$$n\mu_n(G) < -\frac{2^r}{r-1} \sum_{u \in V(G)} \frac{t^{r-1}(u)}{d^{2r-3}(u)}.$$
(4)

By Hölder's inequality, we find that

$$\left(\sum_{u\in V(G)} d\left(u\right)\right)^{\frac{r-2}{r-1}} \left(\sum_{u\in V(G)} \frac{t^{r-1}\left(u\right)}{d^{2r-3}\left(u\right)}\right)^{\frac{1}{r-1}} \ge \sum_{u\in V(G)} \left(\frac{t^{r-1}\left(u\right)}{d^{2r-3}\left(u\right)}\right)^{\frac{1}{r-1}} d^{\frac{r-2}{r-1}}\left(u\right) = \sum_{u\in V(G)} \frac{t\left(u\right)}{d\left(u\right)},$$

THE ELECTRONIC JOURNAL OF COMBINATORICS XX (200X), #R00

and so,

$$\sum_{u \in V(G)} \frac{t^{r-1}(u)}{d^{2r-3}(u)} \ge \frac{1}{(2m)^{r-2}} \left(\sum_{u \in V(G)} \frac{t(u)}{d(u)}\right)^{r-1}.$$

Hence, from (4), we find that

$$n\mu_n(G) \le -\frac{4}{(r-1)m^{r-2}} \left(\sum_{u \in V(G)} \frac{t(u)}{d(u)}\right)^{r-1}.$$

Assume the assertion of the theorem is false, that is to say

$$\mu_n(G) \ge -\frac{2^{r+1}m^r}{rn^{2r-1}},\tag{5}$$

and so,

$$\frac{4}{(r-1)m^{r-2}} \left( \sum_{u \in V(G)} \frac{t(u)}{d(u)} \right)^{r-1} \le -n\mu_n(G) \le \frac{2^{r+1}m^r}{rn^{2r-2}},$$

implying

$$\sum_{u \in V(G)} \frac{t(u)}{d(u)} \le 2\left(\frac{r-1}{r}\right)^{\frac{1}{r-1}} \frac{m^2}{n^2}.$$

Hence, in view of (2), we obtain,

$$\mu_n(G) \le \left(4\left(\frac{r-1}{r}\right)^{\frac{1}{r-1}} - 4\right) \frac{m^2}{n(n^2 - 2m)^2}$$

and from (5) it follows that

$$-\frac{2^{r+1}m^r}{rn^{2r-1}} \le \mu_n(G) \le 4\left(\left(\frac{r-1}{r}\right)^{\frac{1}{r-1}} - 1\right)\frac{m^2}{n(n^2 - 2m)}.$$

Then,

$$\left(1 - \left(\frac{r-1}{r}\right)^{\frac{1}{r-1}}\right) \frac{m^2}{n\left(n^2 - 2m\right)} \le \frac{2^{r-1}m^r}{rn^{2r-1}}$$

and so,

$$1 - \left(\frac{r-1}{r}\right)^{\frac{1}{r-1}} \le \frac{2^{r-1}m^{r-2}\left(n^2 - 2m\right)}{rn^{2r-2}}.$$

Since  $0 < m < n^2$ , the expression  $m^{r-2}n^2 - 2m^{r-1}$  attains its maximum at  $m = \frac{r-2}{2(r-1)}n^2$ ; thus,

$$1 - \left(\frac{r-1}{r}\right)^{\frac{1}{r-1}} < \frac{2}{r} \left(\frac{r-2}{r-1}\right)^{r-2} \frac{1}{r-1}.$$

Hence, by Bernoulli's inequality, we see that

$$\frac{1}{r(r-1)} \le 1 - \left(\frac{r-1}{r}\right)^{\frac{1}{r-1}} < \frac{2}{r} \left(\frac{r-2}{r-1}\right)^{r-2} \frac{1}{r-1},$$

a contradiction for  $r \ge 3$ . Therefore, assumption (5) is false and the proof is completed.

We conclude with the following problem.

**Problem 3** Find a simple lower bound on the second eigenvalue of graphs of order n, size m, and independence number r.

## References

- B. Bollobás, Modern Graph Theory, Graduate Texts in Mathematics, 184, Springer-Verlag, New York (1998), xiv+394 pp.
- [2] B. Bollobás and V. Nikiforov, Graphs and Hermitian matrices: eigenvalue interlacing, to appear in *Discr. Math.*