New approach to Hermitian q-differential operators on \mathbb{R}_q^N

Gaetano Fiore *

Dip. di Matematica e Applicazioni, Fac. di Ingegneria Università di Napoli, V. Claudio 21, 80125 Napoli and I.N.F.N., Sezione di Napoli, Complesso MSA, V. Cintia, 80126 Napoli

Abstract

We report on our recent breakthrough [9] in the costruction for q > 0 of Hermitean and "tractable" differential operators out of the $U_q so(N)$ -covariant differential calculus on the noncommutative manifolds \mathbb{R}_q^N (the socalled "quantum Euclidean spaces").

1 Introduction

Noncommutative geometry for non-compact noncommutative manifolds is usually much more difficult than for compact ones, especially when trying to proceed from an algebraic to a functional-analytical treatment (see e.g. [12]). Major sources of complications are \star -structures and \star -representations of the involved algebras. As a noncommutative space here we adopt the socalled N-dimensional quantum Euclidean space \mathbb{R}_q^N ($N \geq 3$), a deformation of \mathbb{R}^N characterized by its covariance under the quantum group $U_q so(N)$ (with q real so that the latter is compact; for \star -representation purposes we shall later need and choose q > 0) and shall denote by F the \star -algebra "of functions on \mathbb{R}_q^N ". The \star -structure [13] associated to the corresponding $U_q so(N)$ -covariant differential calculus [1]

^{*}Talk given at the Conference "Noncommutative Geometry and Representation Theory in Mathematical Physics", Karlstad (Sweden), July 2004. Preprint 04-45 Dip. Matematica e Applicazioni, Università di Napoli; DSF/04-35; ESI 1529. Work partially supported by the European Commission RTN Programme HPRN-CT-2000-00131 and by MIUR

on \mathbb{R}_q^N [6] is however characterized by an unpleasant nonlinear action on the differentials, the partial derivatives and the exterior derivative. This at the origin of a host of formal and substantial complications: it seems hardly possible to construct physically relevant differential operators (e.g. momentum components, Laplacian, kinetic term in a would-be field theory on \mathbb{R}_q^N) which are at the same time Hermitean and "tractable".

Here we report on a recent breakthrough [9] to the approach to these problems. This is based on our result (Thm 1 in [9]) that the transformation laws [13] of the N partial derivatives ∂^{α} and of the exterior derivative d of the differential calculus [1] on \mathbb{R}_q^N under the \star -structure can be expressed as the similarity transformations

$$\partial^{\alpha \star} = -\tilde{\nu}'^{-2} \tilde{\partial}^{\alpha} \tilde{\nu}'^{2}, \qquad d^{\star} = -\tilde{\nu}'^{2} d\tilde{\nu}'^{-2}, \qquad (1)$$

where $\tilde{\nu}'$ is a $U_q so(N)$ -invariant positive-definite pseudodifferential operator, more precisely the realization of the fourth root of the ribbon element of the extension of $U_q so(N)$ with a central element generating dilatations of \mathbb{R}_q^N .

The situation reminds us of what one encounters in functional analysis on the real line when taking the Hermitean conjugate of a differential operator like

$$\partial := \tilde{\nu}'(x) \frac{d}{dx} \frac{1}{\tilde{\nu}'(x)},$$

where $\tilde{\nu}'(x)$ is a positive smooth function; as an element of the Heisenberg algebra ∂ is not imaginary (excluding the trivial case $\tilde{\nu}' \equiv 1$) w.r.t. the \star -structure

$$x^* = x, \qquad \left(\frac{d}{dx}\right)^* = -\frac{d}{dx},$$

but fulfills the similarity transformation (1). This corresponds to the fact that it is not antihermitean as an operator on $L^2(\mathbb{R})$. ∂ is however (formally) antihermitean on $L^2(\mathbb{R}, \tilde{\nu}'^{-2}dx)$, the Hilbert space of square-integrable functions over \mathbb{R} endowed with measure $\tilde{\nu}'^{-2}dx$. In other words, if we introduce the scalar product

$$(\phi, \psi) = \int \phi^{\star}(x) \,\tilde{\nu}'^{-2}(x) \,\psi(x) dx, \tag{2}$$

[as one does when setting the Sturm-Liouville problem for ∂^2], ∂ becomes antihermitean under the corresponding Hermitean conjugation \dagger^1 :

$$(A^{\dagger}\phi, \psi) := (\phi, A\psi) \qquad \Rightarrow \qquad \partial^{\dagger} = -\partial.$$

The idea is to adopt the analog of (2), where now ν' is a *pseudodifferential* operator rather than a function, as a scalar product also in our noncommutative space.

¹The Hermitean conjugation † is the representation of the following modified \star -structure \star' of the Heisenberg algebra: $a^{\star'} = (\tilde{\nu}'^{-2} a \tilde{\nu}'^{2})^{\star} = \tilde{\nu}'^{2} a^{*} \tilde{\nu}'^{-2}$.

Preliminaries and notation 2

The algebras.

F is essentially the unital associative algebra over $\mathbb{C}[[h]]$ generated by N elements x^{α} (the cartesian "coordinates") modulo the relations (3) given below, and is extended to include formal power series in the generators; out of F one can extract subspaces consisting of elements that can be considered integrable or square-integrable functions. The $U_qso(N)$ -covariant differential calculus on \mathbb{R}_q^N [1] is defined introducing the invariant exterior derivative d, satisfying nilpotency and the Leibniz rule d(fg) = dfg + fdg, and imposing the covariant commutation relations (4) between the x^{α} and the differentials $\xi^{\alpha} := dx^{\alpha}$. Partial derivatives are introduced through the decomposition $d =: \xi^{\alpha} \partial_{\alpha}$. All the other commutation relations are derived by consistency. The complete list is given by

$$\mathcal{P}_{a\gamma\delta}^{\ \alpha\beta}x^{\gamma}x^{\delta} = 0, \tag{3}$$

$$x^{\gamma} \xi^{\alpha} = q \hat{R}^{\gamma \alpha}_{\beta \delta} \xi^{\beta} x^{\delta}, \tag{4}$$

$$(\mathcal{P}_s + \mathcal{P}_t)^{\alpha\beta}_{\gamma\delta} \xi^{\gamma} \xi^{\delta} = 0, \tag{5}$$

$$\mathcal{P}_{a\gamma\delta}^{\ \alpha\beta}\partial_{\beta}\partial_{\alpha} = 0, \tag{6}$$

$$\partial_{\alpha} x^{\beta} = \delta_{\alpha}^{\beta} + q \hat{R}_{\alpha\delta}^{\beta\gamma} x^{\delta} \partial_{\gamma}, \qquad (7)$$

$$\partial^{\gamma} \xi^{\alpha} = q^{-1} \hat{R}_{\beta\delta}^{\gamma\alpha} \xi^{\beta} \partial^{\delta}. \qquad (8)$$

$$\partial^{\gamma} \xi^{\alpha} = q^{-1} \hat{R}^{\gamma \alpha}_{\beta \delta} \xi^{\beta} \partial^{\delta}. \tag{8}$$

Here the $N^2 \times N^2$ matrix \hat{R} is the braid matrix of $SO_q(N)$ [6]. The matrices \mathcal{P}_s , \mathcal{P}_a , \mathcal{P}_t are $SO_q(N)$ -covariant deformations of the symmetric trace-free, antisymmetric and trace projectors respectively, which appear in the orthogonal projector decomposition of R

$$\hat{R} = q\mathcal{P}_s - q^{-1}\mathcal{P}_a + q^{1-N}\mathcal{P}_t. \tag{9}$$

Thus they satisfy the equations (with A, B = s, a, t)

$$\mathcal{P}_A \mathcal{P}_B = \mathcal{P}_A \delta_{AB}, \qquad \qquad \mathcal{P}_s + \mathcal{P}_a + \mathcal{P}_t = \mathbf{1}_{N^2}. \tag{10}$$

The \mathcal{P}_t projects on a one-dimensional sub-space and can be written in the

$$\mathcal{P}_{t\gamma\delta}^{\alpha\beta} \sim g^{\alpha\beta}g_{\gamma\delta};\tag{11}$$

here the $N \times N$ matrix $g_{\alpha\beta}$ is a $SO_q(N)$ -isotropic tensor, deformation of the ordinary Euclidean metric. The elements

$$r^2 \equiv x \cdot x := x^{\alpha} g_{\alpha\beta} x^{\beta}, \qquad \partial \cdot \partial := g^{\alpha\beta} \partial_{\beta} \partial_{\alpha}$$

are $U_q so(N)$ -invariant and respectively generate the centers of F, F'. r^2 is real and positive-definite w.r.t. the \star -structure in F for real q introduced in [6]. $\partial \cdot \partial$ is a deformation of the Laplacian on \mathbb{R}^N . We shall adopt here mainly a set $\{x^{\alpha}\}$ of real generators $x^{\alpha} = V_i^{\alpha} x^i$, whereas by $\{x^i\}$ we mean the standard non-real basis adopted since the works [6, 1], in which r^2 takes the form $r^2 = \sum_i x^{i\star} x^i$ and the relations (3-8) take the simplest explicit form, and V is the (complex) linear transformation relating the two bases.²

We shall slightly extend F by introducing the square root r of r^2 and its inverse r^{-1} as new (central) generators; r is "the deformed Euclidean distance of the generic point of coordinates (x^i) of \mathbb{R}_q^N from the origin". The elements

$$t^{\alpha} := r^{-1}x^{\alpha}$$

fulfill $t \cdot t = 1$ and thus generate the subalgebra $F(S_q^{N-1})$ of "functions on the unit quantum Euclidean sphere". $F(S_q^{N-1})$ can be completely decomposed into eigenspaces V_l of the quadratic Casimir of $U_q so(N)$, or equivalently of the Casimir w defined in (2) with eigenvalues $w_l := q^{-l(l+N-2)}$, implying a corresponding decomposition for F:

$$F(S_q^{N-1}) = \bigoplus_{l=0}^{\infty} V_l, \qquad F = \bigoplus_{l=0}^{\infty} \left(V_l \otimes \mathbb{C}[[r, r^{-1}]] \right). \tag{12}$$

An orthonormal basis $\{S_l^I\}$ (consisting of 'spherical harmonics') of V_l can be extracted out of the set of homogeneous, completely symmetric and trace-free polynomials of degree l, suitably normalized. Therefore for the generic $f \in F$

$$f = \sum_{l=0}^{\infty} f_l = \sum_{l=0}^{\infty} \sum_{I} S_l^I f_{l,I}(r).$$
 (13)

We shall call \mathcal{DC}^* (differential calculus algebra on \mathbb{R}_q^N) the unital associative algebra over $\mathbb{C}[[h]]$ generated by $x^{\alpha}, \xi^{\alpha}, \partial_{\alpha}$ modulo relations (3-8). We shall denote by \bigwedge^* (exterior algebra, or algebra of exterior forms) the graded unital subalgebra generated by the ξ^{α} alone, with grading $\natural \equiv$ the degree in ξ^{α} , and by \bigwedge^p (vector space of exterior p-forms) the component with grading $\natural = p, \ p = 0, 1, 2...$ Each \bigwedge^p carries an irreducible representation of $U_q so(N)$, and its dimension is the binomial coefficient $\binom{N}{p}$ [10], exactly as in the q = 1 (i.e. undeformed) case. We shall endow \mathcal{DC}^* with the same grading \natural , and call \mathcal{DC}^p its component with grading $\natural = p$. The elements of \mathcal{DC}^p can be considered differential-operator-valued p-forms. We

$$x^-x^0 = qx^0x^- \qquad x^0x^+ = qx^+x^0, \qquad [x^-, x^+] = (q^{\frac{1}{2}} - q^{-\frac{1}{2}})x^0x^0,$$

$$q\xi^-\xi^0 + \xi^0\xi^- = q\xi^0\xi^+ + \xi^+\xi^0 = \xi^-\xi^+ + \xi^+\xi^- = (\xi^\pm)^2 = (\xi^0)^2 - (q^{\frac{1}{2}} - q^{-\frac{1}{2}})\xi^-\xi^+ = 0.$$

The *-structure acts on the coordinates as follows:

$$(x^{-})^{*} = q^{\frac{1}{2}}x^{+}, \qquad (x^{0})^{*} = x^{0}, \qquad (x^{+})^{*} = q^{-\frac{1}{2}}x^{-}.$$

For instance, relations (3), (5) for \mathbb{R}_q^3 in the basis $\{x^i\} = \{x^-, x^0, x^+\}$ [6, 1] amount to:

shall denote by \mathcal{H} (Heisenberg algebra on \mathbb{R}_q^N) the unital subalgebra generated by the $x^{\alpha}, \partial_{\alpha}$. Note that by definition $\mathcal{DC}^0 = \mathcal{H}$, and that both \mathcal{DC}^* and \mathcal{DC}^p are \mathcal{H} -bimodules. We shall denote by Ω^* (algebra of differential forms) the graded unital subalgebra generated by the ξ^{α}, x^{α} , with grading ξ , and by Ω^p (space of differential p-forms) its component with grading p; by definition $\Omega^0 = F$ itself. Clearly both Ω^* and Ω^p are F-bimodules. Finally, we recall that by replacing $q \to q^{-1}, \hat{R} \to \hat{R}^{-1}$ in (3-8) one gets an alternative $U_q so(N)$ -covariant calculus, whose exterior/partial derivative and differentials $\hat{d}, i\hat{\partial}^{\alpha}, \hat{\xi}^{\alpha}$ are the \star -conjugates of $d, i\partial^{\alpha}, \xi^{\alpha}$ and are related to the latter by a rather complicated rational transformation [14].

Hodge duality.

The "Hodge map" [10] is a $U_q so(N)$ -covariant, \mathcal{H} -bilinear map

$$*: \mathcal{DC}^p \to \mathcal{DC}^{N-p}$$
 (14)

(p=0,1,...,N) such that * $\mathbf{1}=dV$ and on each \mathcal{DC}^p (hence on the whole \mathcal{DC}^*)

$$*^2 \equiv * \circ * = \mathrm{id} \,, \tag{15}$$

defined by setting on the monomials in the ξ^{α}

$$^{*}(\xi^{\alpha_{1}}...\xi^{\alpha_{p}}) = q^{-N(p-N/2)}c_{p}\,\xi^{\alpha_{p+1}}...\xi^{\alpha_{N}}\varepsilon_{\alpha_{N}...\alpha_{p+1}}{}^{\alpha_{1}...\alpha_{p}}\Lambda^{2p-N}. \tag{16}$$

Here c_p is a suitable normalization factor [going to 1/(N-p)! in the commutative limit $q \to 1$], $\varepsilon^{\alpha_1 \dots \alpha_N}$ is the q-deformed ε -tensor and $\Lambda^{\mp 1}$ are the square root and the inverse square root of

$$\Lambda^{-2} := 1 + (q^2 - 1)x^{\alpha}\partial_{\alpha} + \frac{q^{N-2}(q^2 - 1)^2}{(1 + q^{N-2})^2}r^2\partial \cdot \partial \equiv 1 + O(h); \tag{17}$$

as a pseudodifferential operator Λ acts as a dilatation:

$$\Lambda x^{\alpha} = q^{-1} x^{\alpha} \Lambda, \qquad \Lambda \partial^{\alpha} = q \partial^{\alpha} \Lambda, \qquad \Lambda \xi^{\alpha} = \xi^{\alpha} \Lambda, \qquad \Lambda 1 = 1.$$
(18)

In \mathcal{DC}^* one can introduce [9, 2] as an alternative basis of 1-forms a "frame" [4] $\{\theta^{\alpha}\}$, in the sense that $[\theta^{\alpha}, \mathcal{H}] = 0$ (implying in particular $[\theta^{\alpha}, F] = 0$). In the latter (16) takes the form

$$^*(\theta^{\alpha_1}\theta^{\alpha_2}...\theta^{\alpha_p}) = c_p \,\theta^{\alpha_{p+1}}...\theta^{\alpha_N} \varepsilon_{\alpha_N...\alpha_{p+1}}{}^{\alpha_1...\alpha_p}, \tag{19}$$

Restricting the domain of * to the unital subalgebra $\widetilde{\Omega}^* \subset \mathcal{DC}^*$ generated by $x^{\alpha}, \xi^{\alpha}, \Lambda^{\pm 1}$ one obtains a $U_q so(N)$ -covariant, \widetilde{F} -bilinear map (with $\widetilde{F} \equiv \widetilde{\Omega}^0$)

$$*: \widetilde{\Omega}^p \to \widetilde{\Omega}^{N-p} \tag{20}$$

fulfilling again $^*\mathbf{1}=dV$ and (15). Finally, introducing the exterior coderivatives

$$\delta := -^* d^* \qquad \qquad \hat{\delta} := -^* \hat{d}^* \qquad (21)$$

one finds that on all of \mathcal{DC}^* , and in particular on all of Ω^* , the Laplacians

$$\hat{\Delta} := d\,\delta + \delta\,d = -q^2\,\partial \cdot \partial\Lambda^2 = -q^{-N}\hat{\partial}\cdot\hat{\partial},\tag{22}$$

$$\Delta := \hat{d}\,\hat{\delta} + \hat{\delta}\,\hat{d} = -q^{-2}\,\hat{\partial}\cdot\hat{\partial}\Lambda^{-2} = -q^N\,\partial\cdot\partial\tag{23}$$

are respectively quadratic in the $\hat{\partial}_{\alpha}$, ∂_{α} (note: not in the ∂_{α} , $\hat{\partial}_{\alpha}$!).

The restriction (20) is the notion closest to the conventional notion of a Hodge map on \mathbb{R}_q^N : as a matter of fact, there is no F-bilinear restriction of * to Ω^* . However $\widetilde{\Omega}^*$ is not closed under the \star -structure which we shall recall below.

$U_qso(N)$ and its realization by (pseudo)differential operators.

We extend [11] the Hopf algebra $U_q so(N)$ by adding a central, primitive generator η

$$\phi(\eta) = \mathbf{1} \otimes \eta + \eta \otimes \mathbf{1}, \qquad \epsilon(\eta) = 0, \qquad S\eta = -\eta$$

(here ϕ, ϵ, S respectively denote the coproduct, counit, antipode), and we endow the resulting Hopf algebra $\widetilde{U_qso}(N)$ by the quasitriangular structure

$$\tilde{\mathcal{R}} := \mathcal{R} \ q^{\eta \otimes \eta}, \tag{24}$$

where $\mathcal{R} \equiv \mathcal{R}^{(1)} \otimes \mathcal{R}^{(2)}$ (in a Sweedler notation with upper indices and suppressed summation index) denotes the quasitriangular structure of $U_q so(N)$. The action (which here and in [9] we choose to be right) on \mathcal{DC}^* is completely specified by the transformation laws of the generators $\sigma^{\alpha} = x^{\alpha}, \xi^{\alpha}, \partial^{\alpha}$, which read

$$\sigma^{\alpha} \triangleleft g = \rho^{\alpha}_{\beta}(g)\sigma^{\beta} \qquad \qquad \sigma^{\alpha} = x^{\alpha}, \xi^{\alpha}, \partial^{\alpha}, \qquad g \in U_{q}so(N),$$
$$x^{\alpha} \triangleleft \eta = x^{\alpha}, \qquad \xi^{\alpha} \triangleleft \eta = \xi^{\alpha}, \qquad \partial^{\alpha} \triangleleft \eta = -\partial^{\alpha};$$

here ρ denotes the N-dimensional representation of $U_q so(N)$. The elements

$$Z^{\alpha}_{\beta} := T^{(1)} \rho^{\alpha}_{\beta}(T^{(2)}), \qquad T := \mathcal{R}_{21} \mathcal{R} \equiv T^{(1)} \otimes T^{(2)}, \qquad \mathcal{R}_{21} \equiv \mathcal{R}^{(2)} \otimes \mathcal{R}^{(1)}$$

are generators of $U_q so(N)$, and make up the " $SO_q(N)$ vector field matrix" Z [17, 15]. We recall that the ribbon element $w \in U_q so(N)$ is the central element such that

$$w^2 = u_1 S(u_1), u_1 := (S\mathcal{R}^{(2)})\mathcal{R}^{(1)}.$$

It is well-known [5] that there exist isomorphisms $U_h so(N)[[h]] \simeq U so(N))[[h]]$ of \star -algebras over $\mathbb{C}[[h]]$. This essentially means that it is possible to express the elements of either algebra as power series in $h = \ln q$ with coefficients in the other. In particular w has an extremely simple expression in terms of the quadratic Casimir C of so(N):

$$w = q^{-C} = e^{-hC} = \mathbf{1} + O(h),$$
 $C := X^a X_a =: L(L + N - 2).$ (25)

($\{X^a\}$ is a basis of so(N)). We denote by $\nu:=w^{1/4}$ and by $\tilde{w},\tilde{\nu},\tilde{T},\tilde{Z}$ the analogs of w,ν,T,Z obtained by replacing $\mathcal R$ by $\tilde{\mathcal R}$. As an immediate consequence

$$\tilde{w} = q^{-C}q^{-\eta^2}, \qquad \tilde{\nu} = q^{-C/4}q^{-\eta^2/4}, \qquad \tilde{T} = Tq^{2\eta \otimes \eta}.$$

As shown in [7, 3, 10], for real q there exists a \star -algebra homomorphism³

$$\varphi: \mathcal{H} \rtimes \widetilde{U_q so(N)} \to \mathcal{H},$$
 (26)

acting as the identity on \mathcal{H} itself,

$$\varphi(a) = a \qquad a \in \mathcal{H}. \tag{27}$$

(This requires introducing an additional generator $\eta' = \varphi(\eta) \in \mathcal{DC}^*$ subject to the condition $\varphi(q^{\eta}) = q^{\eta'} = q^{-N/2}\Lambda$, so that $[\eta', \xi^{\alpha}] = 0$, $[\eta', x^{\alpha}] = -x^{\alpha}$, $[\eta', \partial^{\alpha}] = \partial^{\alpha}$. In the sequel we shall often use the shorthand notation

$$\varphi(g) =: g', \qquad g \in \widetilde{U_q so(N)}.$$

One finds [9] on the spherical harmonics of level l (with l = 0, 1, 2, ...)

$$\nu' S_I^I = q^{-l(l+N-2)/4} S_I^I = S_I^I \triangleleft \nu. \tag{28}$$

*-Structure and Hermiticity in configuration space representation

Theorem 2.1 [9] For q > 0 the \star -structure of \mathcal{DC}^* given in [6, 14] can be expressed in the form

$$x^{\alpha \star} = x^{\alpha}, \tag{29}$$

$$\xi^{\alpha\star} = q^N \, \xi^\beta \, Z'^\alpha_\beta \Lambda^{-2},\tag{30}$$

$$\partial^{\alpha \star} = -\tilde{\nu}'^{-2} \partial^{\alpha} \tilde{\nu}'^{2} = -q^{\frac{1-N}{2}} \tilde{\nu}'^{-2} \partial_{\alpha} \tilde{\nu}'^{2} \Lambda, \tag{31}$$

$$d^{\star} = -\tilde{\nu}^{\prime 2} d\tilde{\nu}^{\prime - 2}.\tag{32}$$

$$\varphi(L^{\alpha\beta}) = x^{\alpha}\partial^{\beta} - x^{\beta}\partial^{\alpha}, \qquad \varphi(x^{\alpha}) = x^{\alpha} \qquad \varphi(\partial_{\alpha}) = \partial_{\alpha}, \qquad \varphi(\eta) = -x^{\alpha}\partial_{\alpha}.$$

³This is the q-deformed analog of the realization in the q=1 case of the Cartan-Weyl $L^{\alpha\beta}$ generators of Uso(N) through vector fields (i.e. 1^{st} -order differential operators):

Integration over \mathbb{R}_q^N .

Up to normalization this is defined by [16]

$$\int_{q} f(x) d^{N}x = \int_{0}^{\infty} dr \, m(r) \, r^{N-1} \int_{S_{q}^{N-1}} d^{N-1}t \, f(t,r) = \int_{0}^{\infty} dr \, m(r) \, r^{N-1} f_{0}(r).$$
(33)

where f_0 denotes the l=0 component in (13). The weight m along the radial direction r has to fulfill, beside m(r) > 0, the q-scaling condition

$$m(r) = m(qr). (34)$$

Among the weights fulfilling the latter we can single out two "extreme" cases:

1. $m(r) \equiv 1$;

2.
$$m(r) \equiv m_{J,r_0}(r) := |q-1| \sum_{n=-\infty}^{\infty} r\delta(r-r_0q^n)$$
 ('Jackson' integral).

Integration over \mathbb{R}_q^N fulfills the standard properties of reality, positivity, $\widetilde{U_qso(N)}$ -invariance, and a slightly modified cyclicity property [16]. Moreover, if f is a regular function decreasing faster than $1/r^{N-1}$ as $r \to \infty$ the Stokes theorem holds

$$\int_{q} \partial_{\alpha} f(x) d^{N}x = 0, \qquad \int_{q} \hat{\partial}_{\alpha} f(x) d^{N}x = 0.$$
 (35)

Integration of functions immediately leads to integration of N-forms ω_N : upon moving all the ξ 's to the right of the x's we can express ω_N in the form $\omega_N = f d^N x$ ($d^N x$ denotes the unique independent exterior N-form), and just have to set

$$\int_{q} \omega_{N} = \int_{q} f \, d^{N}x. \tag{36}$$

We introduce the scalar product of two "wave-functions" $\phi, \psi \in F$ and more generally of two "wave-forms" $\alpha_p, \beta_p \in \Omega^p$ by

$$(\boldsymbol{\phi}, \boldsymbol{\psi}) := \int_{q} \boldsymbol{\phi}^{*} \tilde{\nu}'^{-2} \boldsymbol{\psi} \, d^{N} x, \qquad (\alpha_{p}, \beta_{p}) := \int_{q} \alpha_{p}^{*} \tilde{\nu}'^{-2} \beta_{p}. \tag{37}$$

Setting $U^{-1}_{\gamma}^{\delta} := g^{\lambda \delta} g_{\lambda \gamma}$, the second can be easily expressed in terms of integrals

$$(\alpha_p, \beta_p) = \frac{1}{c_{N-p}} \int_q \alpha_{\gamma_p \dots \gamma_1}^{\star} U^{-1}_{\gamma_p}^{\delta_p} \dots U^{-1}_{\gamma_1}^{\delta_1} \tilde{\nu}'^{-2} \beta_{\delta_p \dots \delta_1} d^N x$$
 (38)

involving the form components of the generic $\omega_p \in \Omega^p$ in the frame basis:

$$\omega_p = \theta^{\gamma_1} ... \theta^{\gamma_p} \omega_{\gamma_p ... \gamma_1}.$$

[Note that for $q \neq 1$ $\omega_{\gamma_p...\gamma_1} \in \mathcal{H} \setminus F$, because $\theta^a \in \mathcal{DC}^* \setminus \Omega^*$. As an integrand funtion at the rhs(38) we mean what one obtains after letting all ∂_{α} present under the integral sign act on functions on their right, using the derivation rule (7)].

As a consequence of Theorem 2.1 we obtain

$$(\alpha_p, \hat{d}\beta_{p-1}) = (\hat{\delta}\alpha_p, \beta_{p-1}), \qquad (\hat{d}\beta_{p-1}, \alpha_p) = (\beta_{p-1}, \hat{\delta}\alpha_p). \tag{39}$$

and the (formal) hermiticity of both the momenta $p^{\alpha} = i\partial^{\alpha}$ and the Laplacian Δ :

$$(\boldsymbol{\phi}, p^{\alpha} \boldsymbol{\psi}) = (p^{\alpha} \boldsymbol{\phi}, \boldsymbol{\psi}), \qquad (\alpha_p, \Delta \beta_p) = (\Delta \alpha_p, \beta_p). \tag{40}$$

The kinetic term in the action for a (real) p-form (i.e. an antisymmetric tensor with p-indices) field theory of mass M can be now introduced as

$$S_k = (\Delta \alpha_p, \alpha_p) + M^2(\alpha_p, \alpha_p). \tag{41}$$

Spectral analysis for p^{α} , Δ is affordable (see e.g. [8]), because of the relatively simple derivation rules (7), compared e.g. to those of $p_R^{\alpha} := p^{\alpha} + p^{\alpha \star}$.

The above considerations are still formal unless we make sense out of

$$\tilde{\nu}' = q^{-C'/4} q^{-\eta'^2/4}$$

and its inverse as positive-definite (pseudodifferential) operators on suitable Hilbert subspaces of F, Ω^* . Now $q^{-C'/4}$ is well-defined and positive-definite by (12), (28). To define the action of $q^{-\eta'^2/4}$ on functions $\phi(r)$ we change variables $r \to y = \ln r$, whereby $\eta' = -\partial_y - N/2$ and $r^{N-1}dr = e^{Ny}dy$, for any function $\phi(r)$ denote $\tilde{\phi}(y) := \phi(r)$, and express $e^{yN/2}\tilde{\phi}(y) = r^{N/2}\phi(r)$ in terms of its Fourier transform,

$$e^{\frac{N}{2}y}\tilde{\phi}(y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{\phi}(\omega)e^{i\omega y}d\omega. \tag{42}$$

One finds that $q^{-\eta'^2/4}$ acts as multiplication by $q^{\omega^2/4}$ on the Fourier transform:

$$e^{\frac{N}{2}y}q^{-\eta'^2/4}\tilde{\phi}(y) = q^{-\partial_y^2/4}e^{\frac{N}{2}y}\tilde{\phi}(y)| = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\infty}d\omega\hat{\phi}(\omega)e^{i\omega y}q^{\omega^2/4}.$$

For q < 1 the UV convergence of scalar products is drastically improved by $q^{\omega^2/4}$. Introducing tentatively the 'Hilbert space of square integrable functions on \mathbb{R}_q^{N} ,

$$L_2^m := \left\{ \mathbf{f}(x) \equiv \sum_{l=0}^{\infty} \sum_{I} S_l^I f_{l,I}(r) \in F \mid (\mathbf{f}, \mathbf{f}) < \infty \right\}$$
 (43)

one finds [9] that all works automatically with $L_2^{m\equiv 1}$ (i.e. if we choose $m(r)\equiv 1$). Otherwise, we have to

- 1. further require that m is invariant under inversion, $m(r^{-1}) = m(r)$, and fulfills condition (A.26) in [9] in order that the scalar product (,) be positive-definite;
- 2. at rhs(43) replace F with the subspace $E^{m,\beta,n}$ whose elements $f_{l,I}(r)$'s admit each an analytic continuation in the complex r-plane with poles possibly only in

$$r_{j,k} := q^{j+\beta} e^{i\frac{\pi(2k+1)}{n}},$$
 (44)

where $\beta \in \{0, 1/2\}$, n is an integer submultiple of N, k = 0, 1, ..., n-1, and $j \in \mathbb{Z}$, in order that the hermiticity (40) is actually implemented.

Condition 1. is fulfilled by a large class of weights, including the 'Jackson' ones $m_{J,q^{\beta}}$. Condition 2. selects interesting subclasses of functions, including q-special functions with "quantized parameters", among the simplest one e.g. $1/[1+(q^jr)^n]$.

References

- [1] U. Carow-Watamura, M. Schlieker, S. Watamura, $SO_q(N)$ covariant differential calculus on quantum space and quantum deformation of Schrödinger equation, Z. Phys. C **49** (1991), 439.
- [2] B. L. Cerchiai, G. Fiore, J. Madore, Geometrical Tools for Quantum Euclidean Spaces, Commun. Math. Phys. 217 (2001), 521-554. math.QA/0002007
- [3] C.-S. Chu, B. Zumino, Realization of vector fields for quantum groups as pseudodifferential operators on quantum spaces, Proc. XX ICGTMP, Toyonaka 1995; q-alg/9502005.
- [4] A. Dimakis and J. Madore, Differential calculi and linear connections,
 J. Math. Phys. 37 (1996), 4647-4661.
- [5] V. G. Drinfeld, Quasi Hopf Algebras, Leningrad Math. J. 1 (1990), 1419.
- [6] L. D. Faddeev, N. Y. Reshetikhin, L. Takhtadjan, Quantization of Lie groups and Lie algebras, Leningrad Math. J. 1 (1990), 193-225.
- [7] G. Fiore, Realization of $U_q(so(N))$ within the Differential Algebra on \mathbb{R}_q^N , Commun. Math. Phys. **169** (1995), 475-500.
- [8] G. Fiore, The Euclidean Hopf algebra $U_q(e^N)$ and its fundamental Hilbert space representations, J. Math. Phys. **36** (1995), 4363-4405.
- [9] G. Fiore, On the hermiticity of q-differential operators and forms on the quantum Euclidean spaces \mathbb{R}_q^N , math.QA/0403463
- [10] G. Fiore, Quantum group covariant (anti)symmetrizers, ε-tensors, vielbein, Hodge map and Laplacian, J. Phys. A: Math Gen. 37 (2004), 9175-9193; and references therein.

- [11] S. Majid, Braided Momentum Structure of the q-Poincare Group, J. Math. Phys. **36** (1995), 1991-2007.
- [12] T. Masuda, Y. Nakagami, S. L. Woronowicz, A C* algebraic framework for quantum groups, math.QA/0309338.
- [13] O. Ogievetsky Differential operators on quantum spaces for $GL_q(n)$ and $SO_q(n)$, Lett. Math. Phys. **24** (1992), 245.
- [14] O. Ogievetsky, B. Zumino Reality in the Differential calculus on the q-Euclidean Spaces, Lett. Math. Phys. 25 (1992), 121-130.
- [15] P. Schupp, P. Watts, and B. Zumino, Bicovariant Quantum Algebras and Quantum Lie Algebras, Commun. Math. Phys. 157 (1993), 305; and references therein.
- [16] H. Steinacker, Integration on quantum Euclidean space and sphere in N dimensions, J. Math Phys. 37 (1996), 4738.
- [17] B. Zumino, Introduction to the Differential Geometry of Quantum Groups, In: K. Schmüdgen (ed.), Math. Phys. X, Proc. X-th IAMP Conf. Leipzig (1991), Springer-Verlag (1992).