Van der Waerden's Theorem on Homothetic Copies of $\{1, 1+s, 1+s+t\}$ Byeong Moon Kim 1 and Yoomi Rho 2

Abstract

For all positive integers s and t, Brown et. al [1] defined f(s,t) to be the smallest positive integer N such that every 2-coloring of [1, N] has a monochromatic homothetic copy of $\{1, 1+s, 1+s+t\}$. They proved that $f(s,t) \leq 4(s+t) + 1$ for all s, t and that the equality holds in the case where both $s/g \not\equiv 0 \pmod{4}$ and $t/g \not\equiv 0 \pmod{4}$ with g = gcd(s,t) and in many other cases. Also they proved that for all positive integer m, f(4mt,t) = f(t,4mt) = 4(4mt+t) - t + 1 or 4(4mt+t) + 1. In this paper, we show that f(4mt,t) = f(t,4mt) = 4(4mt+t) - t + 1 and that for all the other (s,t), f(s,t) = 4(s+t) + 1.

keywords: van der Waerden's theorem, arithmetic progression, homothetic copy, 2-coloring, monochromatic triple

1 Introduction

Van der Waerden's theorem on arithmetic progressions [2] states that for every positive integer k there is a smallest positive integer w(k) such that every k-coloring of $[1, w(k)] = \{1, 2, \dots, w(k)\}$ has a monochromatic k-term arithmetic progression. There are lots of results on the estimation of w(k) for large k.(See [3]].)

Let s, t, m be positive integers. A homothetic copy of $\{1, 1+s, 1+s+t\}$ is any set of the form $\{x, x + ys, x + ys + yt\}$ where x and y are positive integers. Regarding 3-term arithmetic progressions as homothetic copies of $\{1, 1+1, 1+1+1\}$, Brown et. al [1] considered van der Waerden's theorem on homothetic copies of $\{1, 1+s, 1+s+t\}$. They defined f(s,t) to be the smallest positive integer N such that every 2-coloring of [1, N] has a monochromatic homothetic copy of $\{1, 1 + s, 1 + s + t\}$. As they have noted, f(s,t) = f(t,s) and hence we assume that $s \ge t$. They proved that $f(s,t) \le 4(s+t) + 1$ for all s, t and that the equality holds in the case where both $s/g \neq 0 \pmod{4}$ and $t/g \neq 0 \pmod{4}$ with g = gcd(s,t) and in many other cases. Also they proved that f(4mt,t) = 4(4mt+t) - t + 1 or 4(4mt+t)+1. In this paper, we show that f(4mt,t) = 4(4mt+t)-t+1 and that for all the other (s,t), f(s,t) = 4(s+t) + 1.

The following theorem is stated as THEOREM 1 in Brown et. al [1].

Theorem 1 Let s, t and c be positive integers. Then f(cs, ct) = c(f(s, t) - 1) + 1.

¹Department of Mathematics, Kangnung National University, Kangnung, Kangweondo, KOREA E-mail: bmk@knusun.kangnung.ac.kr

²Department of Mathematics, Incheon University, Incheon 402-749, Korea E-mail: rho@incheon.ac.kr

2 f(4mt, t)=4(4mt+t)-t+1

Lemma 2 Let N, s, $t \in Z^+$ and C: $[1, N] \to \{0.1\}$ be a coloring without monochromatic triple $\{x, x + sy, x + (s + t)y\}$ for y = 1, 2 or 3. Note that if $u \in \{0, 1\}$ denotes one color of C, then 1 - u denotes the other color. Define V(a) = (C(a), C(a + s), C(a + 2s), C(a + 3s)) for $a \in [1, N - 3s]$. Then the followings are true for each $u \in \{0, 1\}$.

$$\begin{array}{l} 1) \ For \ 1 \leq a \leq N-2s-2t, \ if \ C(a) = C(a+s) = u, \ then \ C(a+2s) = 1-u. \\ For \ s+1 \leq a \leq N-s-2t, \ if \ C(a) = C(a+s) = u, \ then \ C(a-s) = 1-u. \\ For \ 2s+1 \leq a \leq N-2t, \ if \ C(a) = C(a+t) = u, \ then \ C(a+2t) = 1-u. \\ For \ 2s+t+1 \leq a \leq N-t, \ if \ C(a) = C(a+t) = u, \ then \ C(a-t) = 1-u. \\ 2) \ For \ 1 \leq a \leq N-3s-4t, \ if \ V(a) = (u, 1-u, 1-u, u), \ then \ V(a+t) = (1-u, 1-u, u, u) \ or \\ (1-u, u, u, 1-u). \\ 3) \ For \ 1 \leq a \leq N-3s-6t, \ if \ V(a) = (u, 1-u, 1-u, u), \ then \ V(a+t) = (1-u, 1-u, u, u), \ V(a+2t) = \\ (1-u, u, u, 1-u) \ and \ V(a+3t) = (u, u, 1-u, 1-u). \\ 4) \ For \ 1 \leq a \leq N-3s-5t, \ if \ V(a) = (u, 1-u, 1-u, u), \ then \ V(a+2t) = (u, 1-u, 1-u, u, u), \ and \ V(a+3t) = (1-u, u, u, 1-u). \\ 5) \ For \ 1 \leq a \leq N-3s-4t, \ if \ V(a) = (u, 1-u, 1-u, u) \ and \ V(a+t) = (1-u, 1-u, u, u), \ then \ V(a+2t) = (1-u, u, u, 1-u). \\ For \ 1 \leq a \leq N-3s-4t, \ if \ V(a) = (u, 1-u, 1-u, u) \ and \ V(a+t) = (1-u, 1-u, u, u), \ then \ V(a+2t) = (1-u, u, u, 1-u). \\ For \ 1 \leq a \leq N-3s-4t, \ if \ V(a) = (u, 1-u, 1-u, u) \ and \ V(a+t) = (1-u, u, u, 1-u), \ then \ V(a+2t) = (1-u, u, u, 1-u). \\ For \ 1 \leq a \leq N-3s-4t, \ if \ V(a) = (u, 1-u, 1-u, u) \ and \ V(a+t) = (1-u, u, u, 1-u), \ then \ V(a+2t) = (u, 1-u, u, u, 1-u). \\ \end{array}$$

Proof. Throughout the proof, we may assume that u = 0.

1) For $1 \le a \le N - 2s - 2t$, if C(a) = C(a+s) = 0, then C(a+s+t) = 1. Suppose C(a+2s) = 0. Then C(a+2s+t) = C(a+2s+2t) = 1. This gives a monochromatic triple $\{a+s+t, a+2s+t, a+2s+2t\} \subset [1, N]$, a contradiction. Hence C(a+2s) = 1. The others are true by similar methods.

2) For $1 \le a \le N - 3s - 4t$, if V(a) = (0, 1, 1, 0), then C(a + 2s + t) = 0 and C(a + 3s + 3t) = 1. If C(a + s + t) = 0, then V(a + t) = (1, 0, 0, 1) by 1). If C(a + s + t) = 1, then C(a + 3s + t) = 0 and hence C(a + 3s + 2t) = 1. This implies C(a + 3s + 4t) = 0 and then C(a + t) = 1. Thus V(a + t) = (1, 1, 0, 0).

3) For $1 \le a \le N - 3s - 6t$, if V(a) = (0, 0, 1, 0), then C(a + s + t) = C(a + 3s + 2t) = C(a + 3s + 3t) = 1. Thus by 1), C(a + 3s + t) = C(a + 3s + 4t) = 0. This implies C(a + t) = 1. Again by 1) V(a + t) = (1, 1, 0, 0). Now C(a + s + 2t) = 0. Also C(a + 3s + 2t) = C(a + 3s + 3t) = 1 leads to C(a + 2s + 2t) = 0. Again by 1) V(a + 2t) = (1, 0, 0, 1).

4) For $1 \le a \le N-3s-5t$, if V(a) = (0, 1, 0, 0), then C(a+2s+2t) = C(a+3s+t) = C(a+3s+3t) = 1. So C(a+3s+2t) = C(a+s+t) = 0. If C(a+s+2t) = 0, then C(a+t) = 1. And hence either $\{a+t, a+3s+t, a+3s+4t\}$ or $\{a+s+2t, a+3s+2t, a+3s+4t\}$ is monochromatic, a contradiction. Therefore C(a+s+2t) = 1 and hence V(a+2t) = (0, 1, 1, 0) by 1). Thus C(a+2s+3t) = 0. C(a+2t) = C(a+3s+2t) = 0 leads to C(a+3s+5t) = 1 and hence it leads to C(a+s+3t) = 0. By 1), V(a+3t) = (1, 0, 0, 1).

5) Let $1 \le a \le N - 3s - 4t$. Assume V(a) = (0, 1, 1, 0) and V(a+t) = (1, 1, 0, 0). Then C(a+s+2t) = 0and C(a+3s+2t) = C(a+3s+3t) = 1. So C(a+2s+2t) = 0 and hence V(a+2t) = (1, 0, 0, 1). If $1 \le a \le N - 3s - 6t$, then C(a+2s+3t) = 1 and C(a+3s+4t) = C(a+3s+5t) = 0. So by 1) C(a+3s+6t) = 1 and therefore C(a+3t) = (0, 0, 1, 1). Assume V(a) = (0, 1, 1, 0) and V(a+t) = (1, 0, 0, 1). Then C(a+2s+2t) = C(a+3s+3t) = 1 and C(a+3s+4t) = 0. So C(a+3s+2t) = 0 and hence C(a+s+2t) = 1. Thus V(a+2t) = (1, 0, 0, 1).

Lemma 3 Let N, s, t, C and V be the same as in Lemma 2 and $a \in [1, N - 3s]$. Then the followings are true for each $u \in \{0, 1\}$.

1) If V(a) = (u, 1 - u, 1 - u, u) and V(a + 1) = (1 - u, 1 - u, u, u), then $V(a + jt) = \begin{cases} (u, 1 - u, 1 - u, u, u), & j \equiv 0 \pmod{4} \\ (1 - u, 1 - u, u, u), & j \equiv 1 \pmod{4} \\ (1 - u, u, u, 1 - u), & j \equiv 2 \pmod{4} \\ (u, u, 1 - u, 1 - u), & j \equiv 3 \pmod{4} \end{cases}$ (1)

for all j such that $0 \le j \le 2\lfloor \frac{N-a-3s-2t}{2t} \rfloor$.

2) If V(a) = (u, 1 - u, 1 - u, u) and V(a + 1) = (1 - u, u, u, 1 - u), then

$$V(a+jt) = \begin{cases} (u, 1-u, 1-u, u), & j \equiv 0 \pmod{2} \\ (1-u, u, u, 1-u), & j \equiv 1 \pmod{2} \end{cases}$$
(2)

for all j such that $0 \leq j \leq \lfloor \frac{N-a-3s-2t}{t} \rfloor.$

Proof. Throughout the proof, we may assume that u = 0.

1) Let k be the largest integer such that (1) holds for all $0 \le j \le k$. Assume $k \equiv 0 \pmod{4}$. Then V(k-2) = (1,0,0,1) and V(k-1) = (0,0,1,1). Suppose $k < 2\lfloor \frac{N-a-3s-2t}{2t} \rfloor$. Then $a + (k-2)t \le 1$.

N-3s-6t and hence by 5) of Lemma 2 V(k+1) = (1,1,0,0), a contradiction. So $k \ge 2\lfloor \frac{N-a-3s-2t}{2t} \rfloor$. Similarly, $k \ge 2\lfloor \frac{N-a-3s-2t}{2t} \rfloor$ if $k \equiv 1,2,3 \pmod{4}$. Thus (1) holds for all $0 \le j \le 2\lfloor \frac{N-a-3s-2t}{2t} \rfloor$.

2) Let k be the largest integer such that (2) holds for all $0 \le j \le k$. Assume $k \equiv 0 \pmod{2}$. Then V(k-1) = (1,0,0,1) and V(k) = (0,1,1,0). Suppose $k < \lfloor \frac{N-a-3s-2t}{t} \rfloor$. Then $a + (k-1)t \le N-3s-4t$ and hence by 5) of Lemma 2 V(k+1) = (1,0,0,1), a contradiction. So $k \ge \lfloor \frac{N-a-3s-2t}{t} \rfloor$. Similarly, $k \ge \lfloor \frac{N-a-3s-2t}{t} \rfloor$ if $k \equiv 1 \pmod{2}$. Thus (2) holds for all $0 \le j \le \lfloor \frac{N-a-3s-2t}{t} \rfloor$.

Theorem 4 For all $m \in Z^+$, f(4m, 1) = 16m + 4.

Proof. THEOREM 4 of Brown et. al [1] states that f(4mt, t) = 4(4mt + t) - t + 1 or 4(4mt + t) + 1for all positive integers m and t. So it suffices to show that f(4m, 1) < 16m + 5. Suppose not. Then there is a coloring of [1, 16m+4] which has no monochromatic homothetic copy of $\{1, 4m + 1, 4m + 2\}$. Let C be such a coloring and V(a) = (C(a), C(a + 4m), C(a + 8m), C(a + 12m)) for each $a \in [1, 4m + 4]$. We may assume that C(1) = 0. There are five cases to consider. In each case we show that we have contradictions.

Case 1: V(1) = (0, 0, 1, 0). Then by 3) of Lemma 2 V(2) = (1, 1, 0, 0), V(3) = (1, 0, 0, 1) and V(4) = (0, 0, 1, 1). So by 1) of Lemma 3

$$V(3+j) = \begin{cases} (1,0,0,1), & j \equiv 0 \pmod{4} \\ (0,0,1,1), & j \equiv 1 \pmod{4} \\ (0,1,1,0), & j \equiv 2 \pmod{4} \\ (1,1,0,0), & j \equiv 3 \pmod{4} \end{cases}$$
(3)

for all $j \le 4m - 2$. In particular, V(4m + 1) = (0, 1, 1, 0), which contradicts to C(12m + 1) = 0. Case 2: V(1) = (0, 1, 0, 0). Then by 4) of Lemma 2 V(3) = (0, 1, 1, 0) and V(4) = (1, 0, 0, 1). So by 2) of Lemma 3

$$V(3+j) = \begin{cases} (0,1,1,0), & j \equiv 0 \pmod{2} \\ (1,0,0,1), & j \equiv 1 \pmod{2} \end{cases}$$
(4)

for all $j \leq 4m - 1$. In particular, V(4m + 1) = (0, 1, 1, 0), which contradicts to C(12m + 1) = 0.

Case 3: V(1) = (0, 1, 1, 0). Then by 2) of Lemma 2 V(2) is (1, 1, 0, 0) or (1, 0, 0, 1). If V(2) = (1, 1, 0, 0), then by 1) of LEMMA 3

$$V(1+j) = \begin{cases} (0,1,1,0), & j \equiv 0 \pmod{4} \\ (1,1,0,0), & j \equiv 1 \pmod{4} \\ (1,0,0,1), & j \equiv 2 \pmod{4} \\ (0,0,1,1), & j \equiv 3 \pmod{4} \end{cases}$$
(5)

for all $j \le 4m$. In particular, V(4m + 1) = (0, 1, 1, 0), which contradicts to C(12m + 1) = 0. If V(2) = (1, 0, 0, 1), then by 2) of Lemma 3

$$V(1+j) = \begin{cases} (0,1,1,0), & j \equiv 0 \pmod{2} \\ (1,0,0,1), & j \equiv 1 \pmod{2} \end{cases}$$
(6)

for all $j \leq 4m + 1$. In particular, V(4m + 1) = (0, 1, 1, 0), which contradicts to C(12m + 1) = 0.

Case 4: V(1) = (0, 0, 1, 1). Then $V(4m + 1) \neq (0, 0, 1, 1)$ since C(8m + 1) = 1. So there is a minimal k such that $V(k) \neq (u(k), u(k), 1 - u(k), 1 - u(k))$ where $u(k) = \begin{cases} 0, & k \text{ is odd} \\ 1, & k \text{ is even} \end{cases}$. Consider the case when k is even. As V(k - 1) = (0, 0, 1, 1), V(k) is (0, 1, 0, 0) or (0, 1, 1, 0). If

Consider the case when k is even. As V(k-1) = (0, 0, 1, 1), V(k) is (0, 1, 0, 0) or (0, 1, 1, 0). If k = 4m, then it contradicts to C(12m+2) = 0 and C(12m+1) = 1. Assume $k \le 4m - 2$.

Firstly if V(k) = (0, 1, 0, 0), then by 4) of Lemma 2 V(k+2) = (0, 1, 1, 0) and V(k+3) = (1, 0, 0, 1). So by 2) of Lemma 3

$$V(k+2+j) = \begin{cases} (0,1,1,0), & j \equiv 0 \pmod{2} \\ (1,0,0,1), & j \equiv 1 \pmod{2} \end{cases}$$
(7)

for all $j \leq 4m - k$. In particular, V(4m + 1) = (1, 0, 0, 1), which contradicts to C(12m + 1) = 1.

Secondly if V(k) = (0, 1, 1, 0), then by 2) of Lemma 2 V(k + 1) is (1, 1, 0, 0) or (1, 0, 0, 1). If V(k + 1) = (1, 1, 0, 0), then by 1) of Lemma 3

$$V(k+j) = \begin{cases} (0,1,1,0), & j \equiv 0 \pmod{4} \\ (1,1,0,0), & j \equiv 1 \pmod{4} \\ (1,0,0,1), & j \equiv 2 \pmod{4} \\ (0,0,1,1), & j \equiv 3 \pmod{4} \end{cases}$$
(8)

for all $j \le 4m - k + 2$. So V(4m + 1) is (1, 1, 0, 0) or (0, 0, 1, 1), which contradicts to C(12m + 1) = C(8m + 1) = 1. If V(k + 1) = (1, 0, 0, 1), then by 2) of Lemma 3

$$V(k+j) = \begin{cases} (0,1,1,0), & j \equiv 0 \pmod{2} \\ (1,0,0,1), & j \equiv 1 \pmod{2} \end{cases}$$
(9)

for all $j \leq 4m - k + 2$. In particular, V(4m + 1) is (1, 0, 0, 1), which contradicts to C(8m + 1) = 1.

When k is odd, then we get contradictions similarly.

Case 5: V(1) = (0, 1, 0, 1). Then C(8m+3) = 1 and C(12m+3) = 0. Thus V(2) is (0, 0, 1, 0), (0, 0, 1, 1), (0, 1, 0, 1), (1, 0, 1, 1), (1, 1, 0, 1), (1, 0, 1, 0) or (1, 0, 0, 1).

If V(2) = (0, 0, 1, 0), then by 3) of Lemma 2 V(3) = (1, 1, 0, 0), which contradicts to C(8m + 3) = 1. If V(2) = (1, 1, 0, 1), then similarly V(3) = (0, 0, 1, 1), which contradicts to C(12m + 3) = 0.

If V(2) = (0, 0, 1, 1), then C(4m + 3) = 1 and hence by 1) of Lemma 2 V(3) = (0, 1, 1, 0). By the method used in Case 4, we can show that V(4m + 1) is (0, 1, 1, 0) or (1, 0, 0, 1), which contradicts to C(4m + 1) = C(12m + 1) = 1.

If V(2) = (1, 0, 1, 1), then by 4) of Lemma 2 V(4) = (1, 0, 0, 1) and V(5) = (0, 1, 1, 0). So by 2) of

Lemma 3

$$V(4+j) = \begin{cases} (1,0,0,1), & j \equiv 0 \pmod{2} \\ (0,1,1,0), & j \equiv 1 \pmod{2} \end{cases}$$
(10)

for all $j \leq 4m-2$. So V(4m+1) = (0, 1, 1, 0), which contradicts to C(4m+1) = 1.

If V(2) = (1, 0, 0, 1), then by 1) and 2) of Lemma 2 V(3) = (0, 1, 1, 0). So by 2) of Lemma 3

$$V(2+j) = \begin{cases} (1,0,0,1), & j \equiv 0 \pmod{2} \\ (0,1,1,0), & j \equiv 1 \pmod{2} \end{cases}$$
(11)

for all $j \leq 4m$. In particular, V(4m+1) = (0, 1, 1, 0), which contradicts to C(4m+1) = 1.

If V(2) = (0, 1, 0, 1), then C(8m + 3) = C(8m + 4) = 1 and C(12m + 3) = C(12m + 4) = 0 and hence C(4m + 3) = 0. C(3) = 1 as otherwise V(3) = (0, 0, 1, 0) and hence by 3) of Lemma 2 V(4) = (1, 1, 0, 0), a contradiction. Thus V(3) = (1, 0, 1, 0). As C(4m + 1) = 1, $V(4m + 1) \neq (0, 1, 0, 1)$. So there is a minimal k such that $V(k) \neq (u(k), 1 - u(k), u(k), 1 - u(k))$ where $u(k) = \begin{cases} 0, & k \equiv 1, 2 \pmod{4} \\ 1, & k \equiv 0, 3 \pmod{4} \end{cases}$. Firstly assume $k \equiv 1 \pmod{4}$. Then $k \leq 4m + 1$. When $k \leq 4m - 3$, V(k - 2) = V(k - 1) = (1, 0, 1, 0) and hence as above V(k) = (1, 1, 0, 1), a contradiction. When k = 4m + 1, C(8m + k = 12m + 1) = 0 as V(k - 2) = (1, 0, 1, 0), a contradiction. Secondly assume $k \equiv 2 \pmod{4}$. Then $k \leq 4m - 2$. From V(k - 2) = (1, 0, 1, 0) and V(k - 1) = (0, 1, 0, 1), we get C(8m + k) = C(12m + k + 1) = 0 and C(8m + k + 1) = C(12m + k) = 1. Suppose C(4m + k) = 1. Then C(12m + k + 2) = 0 and hence by 1) of Lemma 2 C(12m + k + 3) = 1. Thus C(k) = 0 i.e. V(k) = (0, 1, 0, 1), a contradiction. Therefore C(4m + k) = 0 and hence V(k) = (1, 0, 0, 1) by 1) of lemma 2. By 2) of Lemma 2 V(k + 1) = (0, 1, 1, 0).

$$V(k+j) = \begin{cases} (1,0,0,1), & j \equiv 0 \pmod{2} \\ (0,1,1,0), & j \equiv 1 \pmod{2} \end{cases}$$
(12)

for all $j \leq 4m-k+2$. In particular, V(4m+1) = (0, 1, 1, 0), which contradicts to C(4m+1) = 1. Thirdly assume $k \equiv 3 \pmod{4}$. Then $k \leq 4m-1$. When $k \leq 4m-5$, as in the case when $k \equiv 1 \pmod{4}$, we get a contradiction. When k = 4m-1, V(k+1) = (0, 1, 1, 0) and hence C(8m+k+2 = 12m+1) = 0, a contradiction. Lastly assume $k \equiv 0 \pmod{4}$. Then $k \leq 4m-4$. As in the case when $k \equiv 2 \pmod{4}$, we get contradictions.

If V(2) = (1, 0, 1, 0), then $V(4m+1) \neq (0, 1, 0, 1)$ since C(4m+1) = 1. So there is a minimal k such that $V(k) \neq (u(k), 1 - u(k), u(k), 1 - u(k))$ where $u(k) = \begin{cases} 0, & k \equiv 0, 1 \pmod{4} \\ 1, & k \equiv 2, 3 \pmod{4} \end{cases}$. For each k, the case reduces to the previous case where V(2) = (0, 1, 0, 1) for k+1 and we get contradictions.

Remark 5 Investigating the proof of Theorem 4, we see that we may have a coloring C of [1, 16m + 3]such that C(1) = 0 which has no monochromatic homothetic copy of $\{1, 4m + 1, 4m + 2\}$ only in Case 4, where either k is even, $k \le 4m - 2$, V(k) = (0, 1, 1, 0) and V(k+1) = (1, 1, 0, 0) or k is odd, $k \le 4m - 1$, V(k) = (1, 0, 0, 1) and V(k+1) = (0, 0, 1, 1). In fact only in the latter, if $k \equiv 3 \pmod{4}$, we have such a coloring which satisfies

$$V(j) = \begin{cases} (0,0,1,1), & j \equiv 1 \pmod{2}, j < k\\ (1,1,0,0), & j \equiv 0 \pmod{2}, j < k\\ (1,0,0,1), & j \equiv 3 \pmod{4}, k \le j \le 4m\\ (0,0,1,1), & j \equiv 0 \pmod{4}, k \le j \le 4m\\ (0,1,1,0), & j \equiv 1 \pmod{4}, k \le j \le 4m\\ (1,1,0,0), & j \equiv 2 \pmod{4}, k \le j \le 4m \end{cases}$$
(13)

, C(16m + 1) = C(16m + 2) = 0 and C(16m + 3) = 1. We can dispose it as 0 0 1 1 0 1 1 0 0 0 0 0 1 1 1

0	0	1	1
1	1	0	0
	÷	÷	
0	0	1	1
1	1	0	0
1	0	0	1
0	0	1	1
0	1	1	0
1	1	0	0
	÷	÷	
1	0	0	1
0	0	1	1

with C(4m(j-1)+i) in the *i*-th row and the *j*-th column. For each C, 1-C has no monochromatic homothetic copy of $\{1, 4m+1, 4m+2\}$. Thus we have altogether 2m colorings of [1, 16m+3] which has no monochromatic homothetic copy of $\{1, 4m+1, 4m+2\}$.

Thus together with Theorem 1, the following theorem is true.

Theorem 6 For all $m, t \in Z^+$, f(4mt, t) = 4(4mt + t) - t + 1.

3 Remaining cases

The following theorem which is stated as THEOREM 3 in Brown et. al[1] implies that f(s,t) = 4(s+t) + 1 if t divides s and $s/t \neq 0 \pmod{4}$.

Theorem 7 Let s, t be positive integers with g = gcd(s,t). If $s/g \not\equiv 0 \pmod{4}$ and $t/g \not\equiv 0 \pmod{4}$, then f(s,t) = 4(s+t) + 1. **Theorem 8** Let s, t be positive integers such that s > t > 1 and t does not divide s. If $\lfloor s/t \rfloor$ is even or $\lfloor 2s/t \rfloor$ is even, then f(s,t) = 4(s+t) + 1. If $\lfloor s/t \rfloor$ and $\lfloor 2s/t \rfloor$ are both odd, then f(s,t) = 4(s+t) + 1 provided s, t satisfy the additional condition $s/t \notin (1.5, 2)$.

This theorem is stated as THEOREM 5 in Brown et. al [1]. Together with the following theorem it implies that f(s,t) = 4(s+t) + 1 if t does not divide s.

Theorem 9 Let s, t be positive integers such that s > t > 1 and t does not divide s. If $\lfloor s/t \rfloor$ and $\lfloor 2s/t \rfloor$ are both odd and $s/t \in (1.5, 2)$, then f(s, t) = 4(s + t) + 1.

The following lemma is directly from the Chinese Remainder Theorem.

Lemma 10 Let s, t be positive integers such that gcd(s,t) = 1. Then any integer in [1, 4s + 4t] is equal to 1 + is + jt for some unique $0 \le j < s$ and $-jt/s \le i < 4 + (4-j)t/s$.

Proof of Theorem 9. By Theorem 1, it is enough to consider s, t such that gcd(s,t) = 1. By Theorem 7, we need to consider only the cases where $s \equiv 0 \pmod{4}$ or $t \equiv 0 \pmod{4}$. By THEOREM 2 of Brown et. al [1], $f(s,t) \leq 4(s+t) + 1$ for all positive integers s and t. In each of the following exhaustive cases, we show that the equality holds by choosing a 2-coloring C of [1, 4s + 4t] which contains no homothetic copy of $\{1, 1 + s, 1 + s + t\}$. By Lemma 10, for each $1 + is + jt \in [1, 4s + 4t]$, $0 \leq j < s$ and $-jt/s \leq i < 4 + (4 - j)t/s$. Figure 1-Figure 5 show C(1 + is + jt) for those i and j.

Case 1: $s \equiv 0 \pmod{4}$ and $t \equiv 3 \pmod{4}$. Consider the following two subcases.

Case 1a: $s/t \ge 5/3$. Note that $1 + (-t+7)s + (s-1)t \ge 1 + 4s + 4t$ and hence $1 + (-t+7)s + (s-1)t \notin [1, 4s + 4t]$. We marked @ on its position in Figure 1. Let $A = \{1 + is + jt \mid i \ge -t+7 \text{ or } j \le s-4\}$ and $B = \{1 + is + jt \mid i \le -t+6, j \ge s-3\}$. For $1 + is + jt \in A$ let $C(1 + is + jt) = \begin{cases} 0, & j \text{ even}, i \equiv 0, 1 \pmod{4}; j \text{ odd}, i \equiv 2, 3 \pmod{4} \\ 1, & \text{otherwise} \end{cases}$ and for $1 + is + jt \in B$ let

$$C(1+is+jt) = \begin{cases} 0, & j \text{ even } i \equiv 1,2 \pmod{4}; j \text{ odd } i \equiv 0,3 \pmod{4} \\ 1, & \text{otherwise} \end{cases}$$

To show that C has no monochromatic homothetic copy of $\{1, 1 + s, 1 + s + t\}$, it is enough to show that C has no monochromatic triple $\{1 + is + jt, 1 + (i + y)s + jt, 1 + (i + y)s + (j + y)t\}$ where y = 1, 2 or 3. Suppose C has such a triple. Then the triple is not in A as $C(1 + (i + y)s + jt) \neq$ C(1 + (i + y)s + (j + y)t) for y = 1, 3 and $C(1 + (i + y)s + jt) \neq C(1 + is + jt)$ for y = 2. Similarly the




Figure 1: Case 1a

triple is not in B either. Note that $1 + (i+y)s + jt \in A$ from the definitions of A and B. If y = 1, then $1 + (i+y)s + (j+y)t \in B$ and hence $i \leq -t+5$ and j = s-4. As $i \geq -jt/s$, $i \geq -t+3$. For each of such an (i, j), $\{1+is+jt, 1+(i+y)s+jt, 1+(i+y)s+(j+y)t\}$ is not monochromatic by the definitions of A and B, a contradiction. By a similar argument, the triple is not monochromatic if y = 3. If y = 2, then $1 + is + jt \in B$. Thus $i \geq -t+5$ and $j \geq s-3$ and therefore $1 + (i+2)s + (j+2)t \geq 1 + 4s + 4t$, a contradiction. Therefore C avoids monochromatic homothetic copy of $\{1, 1+s, 1+s+t\}$.

The following 2-coloring C of [1, 4s + 4t] in each case has no monochromatic homothetic copy of $\{1, 1+s, 1+s+t\}$ by similar arguments.

 $\begin{aligned} \text{Case 1b: } s/t < 5/3. \text{ Note that } 1 + (-t+3)s + (s-5)t < 1 \text{ and hence } 1 + (-t+3)s + (s-5)t \notin [1, 4s+4t]. \end{aligned}$ We marked @ on its position in Figure 2. Let C(1+4s+2t) = 1 and C(1+5s+2t) = C(1+4s+3t) = 0.For the other (i, j) if $i \ge -t+7$ or $2 \le j \le s-5$, then let $C(1+is+jt) = \begin{cases} 0, \ j \text{ even } i \equiv 0, 3 \pmod{4}; \ j \text{ odd } i \equiv 1, 2 \pmod{4} \\ 1, \ \text{ otherwise} \end{cases}$ For the other (i, j), let $C(1+is+jt) = \begin{cases} 0, \ (i, j) = (0, 0), (2, 0), (5, 0), (0, 1), (1, 1), (4, 1), (-t+4, s-4), \\ (-t+3, s-3), (-t+5, s-3), (-t+6, s-3), (-t+2, s-2), \\ (-t+4, s-2), (-t+5, s-2), (-t+1, s-1), (-t+3, s-1), \\ 1, \ \text{ otherwise} \end{cases}$

Case 2: $s \equiv 0 \pmod{4}$ and $t \equiv 1 \pmod{4}$. Consider the following two subcases.




Figure 2: Case 1b

1	i=-t+1		i=-	t+12	i	i=0	
j=0						1 0 0 1	1 0 1
						0 1 1 0	0 1
					1	1 0 0 1	1 0
					0	0 1 1 0	
					0 0	1 1 0 1	
					1 1	0 0 1 0	
					1 0 0	1 1 0	
					0 1 1	0 0 1	
		•	•		•	•	
		•	•		•	•	
j=s-8		1 1 0 0	1 1 0				
	1	0 0 1 1	0 0				
	0	1 1 0 0	1 1	• • •			
	1 1	0 0 1 1	0				
	0 0	1 1 0 0					
	0 0 1	1 0 1 1					
	1 1 0	0 1 0					
j=s-1	1 1 0 0	1 0 @					

Figure 3: Case 2a

Case 2a: $s/t \ge 5/3$. As in Case 1a, $1 + (-t+7)s + (s-1)t \notin [1, 4s + 4t]$. We marked @ on its position in Figure 3. Let C(1 + 3s + 4t) = 1, C(1 + 3s + 5t) = 0 and C(1 + 6s) = 0. For the other (i, j), if $4 \le j \le s - 4$, then let $C(1 + is + jt) = \begin{cases} 0, \ j \text{ even } i \equiv 2, 3 \pmod{4}; \ j \text{ odd } i \equiv 0, 1 \pmod{4} \\ 1, \ \text{otherwise} \end{cases}$. For the other (i, j), if $j \le 3$, then let $C(1 + is + jt) = \begin{cases} 0, \ j \text{ even } i \equiv 1, 2 \pmod{4}; \ j \text{ odd } i \equiv 0, 3 \pmod{4} \\ 1, \ \text{otherwise} \end{cases}$. For the other (i, j), let $C(1 + is + jt) = \begin{cases} 0, \ (i, j) = (-t + 2, s - 3), (-t + 3, s - 3), (-t + 6, s - 3), \\ (-t + 4, s - 2), (-t + 5, s - 2), (-t + 7, s - 2), (-t + 3, s - 1), \\ (-t + 4, s - 1), (-t + 6, s - 1), \\ 1, \ \text{otherwise} \end{cases}$. Case 2b: s/t < 5/3. As in Case 1b, $1 + (-t+3)s + (s-5)t \notin [1, 4s + 4t]$. We marked @ on its position in

Figure 4. Let C(1 + (-t+4)s + (s-6)t) = 1, C(1 + (-t+4)s + (s-5)t) = 0, C(1 + (-t+8)s + (s-4)t) = 0and C(1 + (-t+8)s + (s-3)t) = 1. For the other (i, j), if $i \le 3$ and $2 \le j \le s-5$, then let $C(1 + is + jt) = \begin{cases} 0, & j \text{ even } i \equiv 0, 3 \pmod{4}; \ j \text{ odd } i \equiv 1, 2 \pmod{4} \\ 1, & \text{otherwise} \end{cases}$. For the other (i, j), if $j \ge s-4$, then let

i	=-t+1		i=-	t+12	i	=0	
j=0						0 1 0 1	1 0 0
						1 0 1 0	0 1
					0	0 1 1 0	1 0
					1	1 0 0 1	0
					0 1 1 0	0 1 1 0	
					1 0 0 1	1 0 0 1	
				• • •	0 1 1 0	0 1 1 0	
					1 0 0 1	1 0	
		•	•				
		•	•	• • •		•	
j=s-8		0 1 1 0	0 1 1 0				
		1 0 0 1	1 0 0 1				
	1	0 1 1 0	0 1 1	• • •			
	@ 0	1 0 0 1	1 0				
	1 1	0 0 1 0	0				
	1 0 0	1 1 0 1					
	0 1 1	0 0 1					
j=s-1	1 1 0 0	1 1 0					

Figure 4: Case 2b

$$\begin{split} C(1+is+jt) &= \begin{cases} 0, & j \text{ even } i \equiv 0,1 \pmod{4}; j \text{ odd } i \equiv 2,3 \pmod{4} \\ 1, & \text{otherwise} \end{cases} \\ \text{For the other } (i,j), \text{ let} \\ C(1+is+jt) &= \begin{cases} 0, & (i,j) = (0,0), (2,0), (5,0), (6,0), (1,1), (3,1), (4,1), (5,2), (4,3), \\ 1, & \text{otherwise} \end{cases} . \end{split}$$

Case 3: $s \equiv 1,3 \pmod{4}$ and $t \equiv 0 \pmod{4}$. Notice that either $1 + (-t+7)s + (s-1)t \notin [1, 4s + 4t]$ or $1 + (-t+3)s + (s-5)t \notin [1, 4s + 4t]$ depends on $s/t \ge 5/3$ or not. We marked @' and @ on their positions in Figure 5. Let C(1 + 3s + 4t) = 1, C(1 + 3s + 5t) = 0, C(1 + 6s) = 1, C(1 + s - t) = 1, C(1 + 2s - t) = 1, C(1 + 3s - t) = 0, C(1 + 4s - t) = 0, C(1 + 5s - t) = 1, C(1 + 6s - t) = 0, C(1 + 7s - t) = 0. For the other (i, j), if $4 \le j \le s - 2$, then let $C(1 + is + jt) = \begin{cases} 0, \ j \text{ even } i \equiv 2, 3 \pmod{4}; j \text{ odd } i \equiv 0, 1 \pmod{4} \\ 1, \ \text{ otherwise} \end{cases}$. For the other (i, j), if $j \le 3$, then let $C(1 + is + jt) = \begin{cases} 0, \ j \text{ even } i \equiv 1, 2 \pmod{4}; j \text{ odd } i \equiv 0, 3 \pmod{4} \\ 1, \ \text{ otherwise} \end{cases}$.

By Theorem 7, Theorem 8 and Theorem 9, for all pair of positive integers $(s,t) \neq (4mt,t)$ for a positive integer m, f(s,t) = 4(s+t) + 1.




Figure 5: Case 3

Acknowledgement: We thank Seoul National University for providing us with a place to research during the winter break of 2004.

References

- [1]; T. C. Brown, B. M. Landman and M. Mishna, Monochromatic homothetic copies of $\{1, 1 + s, 1 + s + t\}$, Canad. Math. Bull., 40. No.2 (1997); 149-157.
- [2]; B. L. Van der Waerden, Beweis einer Baudetschen Vermutung, Nieuw Arch. Wisk, 15 (1927);
 212-216.
- [3]; R. L. Graham, B. L. Rothschild and J. H. Spencer, *Ramsey Theory*, Wiley-Interscience, New York, 1990.