

# HOLOMORPHIC DISCS WITH DENSE IMAGES

FRANC FORSTNERIČ AND JÖRG WINKELMANN

ABSTRACT. Let  $\Delta$  be the open unit disc in  $\mathbb{C}$ ,  $X$  a connected complex manifold and  $\mathcal{D}$  the set of all holomorphic maps  $f: \Delta \rightarrow X$  with  $\overline{f(\Delta)} = X$ . We prove that  $\mathcal{D}$  is dense in  $Hol(\Delta, X)$ .

## 1. INTRODUCTION

Let  $\Delta_r = \{z \in \mathbb{C}: |z| < r\}$  and  $\Delta = \Delta_1$ . In [7] the second author proved that *for any irreducible complex space  $X$  there exists a holomorphic map  $\Delta \rightarrow X$  with dense image*, and he raised the question whether the set of all holomorphic maps  $\Delta \rightarrow X$  with dense image forms a dense subset of the set  $Hol(\Delta, X)$  of all holomorphic maps  $\Delta \rightarrow X$  with respect to the topology of locally uniform convergence.

In this paper we show that the answer to this question is positive if  $X$  is smooth, but negative for some singular space.

**Theorem 1.** *For any connected complex manifold  $X$  the set of holomorphic maps  $\Delta \rightarrow X$  with dense images forms a dense subset in  $Hol(\Delta, X)$ . The conclusion fails for some singular complex surface  $X$ .*

The situation is quite different for *proper discs*, i.e., proper holomorphic maps  $\Delta \rightarrow X$ . The paper [3] contains an example of a non-pseudoconvex bounded domain  $X \subset \mathbb{C}^2$  such that a certain nonempty open subset  $U \subset X$  is not intersected by the image of any proper holomorphic disc  $\Delta \rightarrow X$ . On the other hand, proper holomorphic discs exist in great abundance in *Stein manifolds* [5], [1], [2].

## 2. PREPARATIONS

**Lemma 1.** *Let  $W_n$  be a decreasing sequence (i.e.,  $W_{n+1} \subset W_n$ ) of open sets with  $\Delta \subset W_n \subset \Delta_2$  for every  $n$ . Let  $K = \bigcap_n \overline{W_n}$  and assume that the interior of  $K$  coincides with  $\Delta$ . Furthermore assume that there are biholomorphic maps  $\phi_n: \Delta \rightarrow W_n$  with  $\phi_n(0) = 0$  for  $n = 1, 2, \dots$*

---

*Date:* August 7, 2018.

*2000 Mathematics Subject Classification.* 32E30, 32H02.

*Key words and phrases.* Holomorphic discs, complex spaces.

Research of the first author was supported by grants P1-0291 and J1-6173, Republic of Slovenia.

Then there exists an automorphism  $\alpha \in \text{Aut}(\Delta)$  and a subsequence  $(\phi_{n_k})$  of the sequence  $(\phi_n)$  such that  $\phi_{n_k} \circ \alpha^{-1}$  converges locally uniformly to the identity map  $\text{id}_\Delta$  on  $\Delta$ .

*Proof.* Montel's theorem shows that, after passing to a suitable subsequence, we have  $\lim_{n \rightarrow \infty} \phi_n = \alpha: \Delta \rightarrow K$  and  $\lim_{n \rightarrow \infty} (\phi_n^{-1}|_\Delta) = \beta: \Delta \rightarrow \overline{\Delta}$ . Since the limit maps are holomorphic and satisfy  $\alpha(0) = 0$  and  $\beta(0) = 0$ , we conclude that  $\alpha(\Delta) \subset \text{Int}K = \Delta$  and  $\beta(\Delta) \subset \Delta$ . Moreover  $\alpha \circ \beta = \text{id}_\Delta = \beta \circ \alpha$ , and hence both  $\alpha$  and  $\beta$  are automorphisms of  $\Delta$  (indeed, rotations  $z \rightarrow ze^{it}$ ).  $\square$

We also need the following special case of a result of the first author (theorem 3.2 in [4]):

**Proposition 1.** *Let  $X$  be a complex manifold,  $0 < r < 1$ ,  $E$  the real line segment  $[1, 2] \subset \mathbb{C}$ ,  $K = \overline{\Delta} \cup E$ ,  $U$  an open neighbourhood of  $\overline{\Delta}$  in  $\mathbb{C}$ ,  $S$  a finite subset of  $K$  and  $f: U \cup E \rightarrow X$  a continuous map which is holomorphic on  $U$ .*

*Then there is a sequence of pair of open neighbourhoods  $W_n \subset \mathbb{C}$  of  $K$  and holomorphic maps  $g_n: W_n \rightarrow X$  such that:*

- (1)  $g_n|_K$  converges uniformly to  $f|_K$  as  $n \rightarrow \infty$ , and
- (2)  $g_n(a) = f(a)$  for all  $a \in S$  and  $n \in \mathbb{N}$ .

### 3. TOWARDS THE MAIN RESULT

In this section we prove the following proposition which is the main technical result in the paper. The first statement in theorem 1 (§1) is an immediate corollary.

**Proposition 2.** *Let  $X$  be a connected complex manifold endowed with a complete Riemannian metric and induced distance  $d$ ,  $S$  a countable subset of  $X$ ,  $f: \Delta \rightarrow X$  a holomorphic map,  $\epsilon > 0$  and  $0 < r < 1$ .*

*Then there exists a holomorphic map  $F: \Delta \rightarrow X$  such that*

- (a)  $S \subset F(\Delta)$ , and
- (b)  $d(f(z), F(z)) \leq \epsilon$  for all  $z \in \Delta_r$ .

*Proof.* Let  $s_1, s_2, s_3, \dots$  be an enumeration of the elements of  $S$ . We shall inductively construct a sequence of holomorphic maps  $f_n: \Delta \rightarrow X$ , numbers  $r_n \in (0, 1)$  and points  $a_{1,n}, \dots, a_{n,n} \in \Delta$  satisfying the following properties for  $n = 0, 1, 2, \dots$ :

- (1)  $f_0 = f$  and  $r_0 = r$ ,
- (2)  $(r_n + 1)/2 < r_{n+1} < 1$ ,
- (3)  $f_n(a_{j,n}) = s_j$  for  $n \geq 1$  and  $j = 1, 2, \dots, n$ ,
- (4)  $d(f_n(z), f_{n+1}(z)) < 2^{-(n+1)}\epsilon$  for all  $z \in \Delta_{r_n}$ , and

- (5)  $d_\Delta(a_{j,n}, a_{j,n+1}) < 2^{-n}$  for  $j = 1, 2, \dots, n$  where  $d_\Delta$  denotes the Poincaré distance on the unit disc.

Assume inductively that the data for level  $n$  (i.e.,  $f_n, r_n, a_{j,n}$ ) have been chosen. (For  $n = 0$  we do not have any points  $a_{j,0}$ .) With  $n$  fixed we choose an increasing sequence of real numbers  $\lambda_k$  with  $\lambda_k > r_n$  and  $\lim_{k \rightarrow \infty} \lambda_k = 1$ . For every  $k \in \mathbb{N}$  the map  $\tilde{g}_k(z) \stackrel{\text{def}}{=} f_n(\lambda_k z) \in X$  is defined and holomorphic on the disc  $\Delta_{1/\lambda_k} \supset \overline{\Delta}$ . After a slight shrinking of its domain we can extend it continuously to the segment  $E = [1, 2] \subset \mathbb{C}$  such that the right end point 2 of  $E$  is mapped to the next point  $s_{n+1} \in S$  (this is possible since  $X$  is connected).

Applying proposition 1 to the extended map  $\tilde{g}_k$  we obtain for every  $k \in \mathbb{N}$  an open neighbourhood  $V_k \subset \mathbb{C}$  of  $K = \overline{\Delta} \cup E$  and a holomorphic map  $g_k: V_k \rightarrow X$  such that

- (i)  $|g_k(z) - f_n(\lambda_k z)| < 2^{-k}$  for all  $z \in \overline{\Delta}$ ,
- (ii)  $g_k(2) = s_{n+1}$ , and
- (iii)  $g_k(a_{j,n}/\lambda_k) = f_n(a_{j,n}) = s_j$  for  $j = 1, \dots, n$ .

Next we choose a decreasing sequence of simply connected open sets  $W_k \subset \mathbb{C}$  ( $k \in \mathbb{N}$ ) with  $K \subset W_k \subset V_k$  and  $K = \cap_k \overline{W}_k$ . Notice that  $\text{Int} K = \Delta$ . By lemma 1 there is a sequence of biholomorphic maps  $\phi_k: \Delta \rightarrow W_k$  with  $\lim_{k \rightarrow \infty} \phi_k = id_\Delta$ .

Consider the holomorphic maps  $h_k = g_k \circ \phi_k: \Delta \rightarrow X$ . By our construction we know that  $\lim_{k \rightarrow \infty} h_k = f_n$  locally uniformly on  $\Delta$ .

To fulfill the inductive step it thus suffices to choose  $f_{n+1} = h_k$  for a sufficiently large  $k$ ,  $a_{j,n+1} = a_{j,n}/\lambda_k$  ( $j = 1, \dots, n$ ),  $a_{n+1,n+1} = \phi_k^{-1}(2)$ . Finally we choose a number  $r_{n+1}$  satisfying

$$\max\{|a_{n+1,n+1}|, \frac{r_n + 1}{2}\} < r_{n+1} < 1.$$

This completes the inductive step.

By properties (2) and (4) the sequence  $f_n$  converges locally uniformly in  $\Delta$  to a holomorphic map  $F: \Delta \rightarrow X$ . Aided by property (1) we also control  $d(f(z), F(z))$  for  $z \in \Delta_r$ . Since the Poincaré metric is complete, property (5) insures that for every fixed  $j \in \mathbb{N}$  the sequence  $a_{j,n} \in \Delta$  ( $n = j, j+1, \dots$ ) has an accumulation point  $b_j$  inside of  $\Delta$ , and (3) implies  $F(b_j) = s_j$  for  $j = 1, 2, \dots$ . Hence  $S \subset F(\Delta)$ .  $\square$

#### 4. SINGULAR SPACES

We use an example of Kaliman and Zaidenberg [6] to show that for a complex spaces  $X$  with singularities the set of maps  $\Delta \rightarrow X$  with dense image need not be dense in  $Hol(\Delta, X)$ . We denote by  $Sing(X)$  the singular locus of  $X$ .

**Proposition 3.** *There is a singular compact complex surface  $S$ , a non-constant holomorphic map  $f: \Delta \rightarrow S$  and an open neighbourhood  $\Omega$  of  $f$  in  $\text{Hol}(\Delta, S)$  such that  $g(\Delta) \subset \text{Sing}(S)$  for every  $g \in \Omega$ .*

*Proof.* In [6] Kaliman and Zaidenberg constructed an example of a singular surface  $S$  with normalization  $\pi: Z \rightarrow S$  such that  $S$  contains a rational curve  $C \simeq \mathbb{P}^1$  while  $Z$  is smooth and hyperbolic. Denote by  $d_Z$  the Kobayashi distance function on  $Z$ . We choose two distinct points  $p, q \in C$  and open relatively compact neighbourhoods  $V$  of  $p$  and  $W$  of  $q$  in  $S$  such that  $\overline{V} \cap \overline{W} = \emptyset$ . The preimages  $\pi^{-1}(\overline{V})$  and  $\pi^{-1}(\overline{W})$  in  $Z$  are also compact, and since  $Z$  is hyperbolic we have

$$r = \min\{d_Z(x, y) : x \in \pi^{-1}(\overline{V}), y \in \pi^{-1}(\overline{W})\} > 0.$$

Fix a point  $a \in \Delta$  with  $0 < d_\Delta(0, a) < r$  and let  $\Omega$  consist of all holomorphic maps  $g: \Delta \rightarrow S$  satisfying  $g(0) \in V$  and  $g(a) \in W$ . Since both  $p$  and  $q$  are lying on the rational curve  $C$ , there is a holomorphic map  $g: \Delta \rightarrow C$  with  $g(0) = p \in V$  and  $g(a) = q \in W$ ; hence the set  $\Omega$  is not empty. Clearly  $\Omega$  is open in  $\text{Hol}(\Delta, S)$ .

To conclude the proof it remains to show that  $g(\Delta) \subset \text{Sing}(S)$  for all  $g \in \Omega$ . Indeed, a holomorphic map  $g: \Delta \rightarrow S$  with  $g(\Delta) \not\subset \text{Sing}(S)$  admits a holomorphic lifting  $\tilde{g}: \Delta \rightarrow Z$  with  $\pi \circ \tilde{g} = g$ . If  $g \in \Omega$  then by construction

$$d_Z(\tilde{g}(0), \tilde{g}(a)) \geq r > d_\Delta(0, a)$$

which violates the distance decreasing property for the Kobayashi pseudometric. This contradiction establishes the claim.  $\square$

In particular, we see that in this example the set of all holomorphic maps  $f: \Delta \rightarrow S$  with dense image *does not* constitute a dense subset of  $\text{Hol}(\Omega, S)$ .

## REFERENCES

- [1] Drinovec-Drnovšek, B.: Discs in Stein manifolds containing given discrete sets. *Math. Z.*, **239** (2002), no. 4, 683–702.
- [2] ——— Proper discs in Stein manifolds avoiding complete pluripolar sets. *Math. Res. Lett.*, **11** (2004), no. 5, 575–581.
- [3] Forstnerič, F., Globevnik, J.: Discs in pseudoconvex domains. *Comment. Math. Helv.*, **67** (1992), no. 1, 129–145.
- [4] Forstnerič, F.: Holomorphic Submersions from Stein Manifolds. *Ann. Inst. Fourier*, to appear. [arXiv: math.CV/0309093]
- [5] Globevnik, J.: Discs in Stein manifolds. *Indiana Univ. Math. J.*, **49** (2000), no. 2, 553–574.
- [6] Kaliman, S., Zaidenberg, M.: Non-hyperbolic complex space with a hyperbolic normalization. *Proc. Amer. Math. Soc.*, **129** (2001), no. 5, 1391–1393.
- [7] Winkelmann, J.: Non-degenerate Maps and Sets. *Math. Z.*, to appear.

INSTITUT OF MATHEMATICS, PHYSICS AND MECHANICS, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

*E-mail address:* `franc.forstneric@fmf.uni-lj.si`

INSTITUT ELIE CARTAN (MATHÉMATIQUES), UNIVERSITÉ HENRI POINCARÉ NANCY 1, B.P. 239, F-54506 VANDŒUVRE-LES-NANCY CEDEX, FRANCE

*E-mail address:* `jwinkel@member.ams.org`

*Webpage:* <http://www.math.unibas.ch/~winkel/>