QUANTUM GROUPS AND NON-COMMUTATIVE COMPLEX ANALYSIS

S. Sinel'shchikov and L. Vaksman

The problem of uniform approximation by analytic polynomials on a compact $K \subset \mathbb{C}$ made an essential impact to the function theory and the theory of commutative Banach algebras. This problem was solved by S. Mergelyan. Much later a theory of uniform algebras was developed and an abstract proof of Mergelyan's theorem was obtained.

The work by W. Arveson [1] starts an investigation of non-commutative analogs for uniform algebras. In particular, a notion of the Shilov boundary for a subalgebra of a C^* -algebra have been introduced therein. So, the initial results of non-commutative complex analysis were obtained. We assume the basic concepts of that work known to the reader.

In mid'90-s an investigation of quantum analogs for bounded symmetric domains has been started within the framework of the quantum group theory [2]. The simplest of those is a unit ball in \mathbb{C}^n . Our goal is to explain that the quantum sphere is the Shilov boundary for this quantum domain. The subsequent results of the authors on non-commutative function theory and quantum groups are available at www.arxiv.org. Specifically, we obtained some results on weighted Bergman spaces, the Berezin transform, and the Cauchy-Szegö kernels for quantum bounded symmetric domains introduced in [7].

In what follows the complex numbers are assumed as a ground field and all the algebras are assumed to be unital. In what follows $q \in (0, 1)$.

To introduce a quantum unit ball, consider a *-algebra $\operatorname{Pol}(\mathbb{C}^n)_q$ given by the generators z_1, z_2, \ldots, z_n and the defining relations $z_j z_k = q z_k z_j$ for j < k,

$$z_j^* z_k = q z_k z_j^*, \quad j \neq k, \qquad z_j^* z_j = q^2 z_j z_j^* + (1 - q^2) \left(1 - \sum_{k > j} z_k z_k^* \right).$$

This *-algebra has been introduced by W. Pusz and S. Woronowicz [6] where one can find a description (up to unitary equivalence) of its irreducible *-representations T. One can demonstrate that $0 < \|f\| \stackrel{\text{def}}{=} \sup_{T} \|T(f)\| < \infty$ for all non-zero $f \in \text{Pol}(\mathbb{C}^n)_q$ and that its C^* -enveloping algebra $C(\mathbb{B})_q$ is a q-analogue for the C^* -algebra of continuous functions in the closed unit ball in \mathbb{C}^n . A plausible description of this C^* -algebra has been obtained by D. Proskurin and Yu. Samoilenko [5].

To introduce a quantum unit sphere, consider a closed two-sided ideal J of the C^* -algebra $C(\mathbb{B})_q$ generated by $1 - \sum_{j=1}^n z_j z_j^*$. Obviously, the C^* -algebra $C(\partial \mathbb{B})_q \stackrel{\text{def}}{=} C(\mathbb{B})_q/J$ is a q-analogue for the algebra we need. Thus the canonical onto morphism

$$j_q: C(\mathbb{B})_q \to C(\partial \mathbb{B})_q$$

is a q-analogue for the restriction operator of a continuous function onto the boundary of the ball.

The closed subalgebra $A(\mathbb{B})_q \subset C(\partial \mathbb{B})_q$ generated by z_1, z_2, \ldots, z_n is a q-analogue for the algebra of continuous functions in the closed ball which are holomorphic in its interior.

Let $j_{A(\mathbb{B})_q}$ be the restriction of the homomorphism j_q onto the subalgebra $A(\mathbb{B})_q$.

Theorem 1 The homomorphism $j_{A(\mathbb{B})_q}$ is completely isometric.

This result is a q-analogue of the well known maximum principle for holomorphic functions. By an Arveson's definition [1] this means that J is a boundary ideal for the subalgebra $A(\mathbb{B})_q$. A proof of this theorem elaborates the methods of quantum group theory and theory of unitary dilations [4].

One can use the deep result of M. Hamana [3] (on existence of the Shilov boundary) to prove the following simple

Proposition 2 *J* is the largest boundary ideal for $A(\mathbb{B})_q$.

Thus the quantum sphere is the Shilov boundary of the quantum ball. A more detailed exposition of this talk is available in [8].

The second named author would like to express his gratitude to M. Livšitz and V. Drinfeld who taught him the theory of non-selfadjoint linear operators and quantum groups, respectively.

References

- [1] W. B. Arveson. Subalgebras of C^* -algebras. Acta Math., 123:141–122, 1969.
- [2] V. Drinfeld. Quantum groups. In A. M. Gleason, editor, *Proceedings of the International Congress of Mathematicians (Berkeley, 1986)*, pages 798–820. Amer. Math. Soc., Providence, RI, 1987.
- [3] M. Hamana. Injective envelopes of operator systems. *Publ. RIMS Kyoto Univ*, 15:773 785, 1979.
- [4] B. Sz.-Nagy and C. Foiaş. Analyse Harmonique des Opérateurs de l'Espace de Hilbert. Masson, Académiai Kiado, 1967.
- [5] D. Proskurin and Yu. Samoilenko. Stability of the C^* -algebra associated with the twisted CCR. Algebra and Representation Theory, 5(4):456–460, 2002.
- [6] W. Pusz and S. Woronowich. Twisted second quantization. Reports Math. Phys., 27:231 257, 1989.
- [7] S. Sinel'shchikov and L. Vaksman. On q-analogues of bounded symmetric domains and Dolbeault complexes. *Mathematical Physics, Analysis, and Geometry*, 1:75–100, 1998.
- [8] L. Vaksman. Maximum principle for holomorphic functions in the quantum ball. *Matematicheskaya fizika, analiz, geometriya (in Russian)*, 10(1):12 28, 2003.