

FINE REGULARITY FOR ELLIPTIC SYSTEMS WITH DISCONTINUOUS INGREDIENTS

DIAN PALAGACHEV AND LUBOMIRA SOFTOVA

ABSTRACT. We propose results on interior Morrey, *BMO* and Hölder regularity for the strong solutions to linear elliptic systems of order $2b$ with discontinuous coefficients and right-hand sides belonging to the Morrey space $L^{p,\lambda}$.

1. MAIN RESULTS

It is well known that, when dealing with elliptic systems, in contrast to the case of a single second-order elliptic equation, the solely essential boundedness of the principal coefficients is not sufficient to ensure Hölder continuity even of the solution (see [8, Chapter 1]). On the other hand, precise estimates on Hölder's seminorms of the solution and its lower order derivatives is a matter of great concern in the study of nonlinear elliptic systems. In fact, these bounds imply good mapping properties of certain Carathéodory operators ensuring this way the possibility to apply the powerful tools of the nonlinear analysis and differential calculus. It turns out that "suitable continuity" of the principal coefficients of the system under consideration is sufficient to guarantee good regularity (e.g. Sobolev) of the solutions (see [5, 12]). We deal here with discontinuous coefficients systems for which the discontinuity is expressed in terms of appurtenance to the class of functions with vanishing mean oscillation. Although such systems have been already studied in Sobolev spaces $W^{2b,p}$ (cf. [3]), our functional framework is that of the Sobolev–Morrey classes $W^{2b,p,\lambda}$. These possess better embedding properties into Hölder spaces than $W^{2b,p}$ and as outgrowth of suitable Caccioppoli-type estimates, we give precise characterization of the Morrey, *BMO* or Hölder regularity of the solution and its derivatives up to order $2b - 1$.

Let Ω be a domain in $\mathbb{R}^n$, $n \geq 2$, and consider the linear system

$$(1.1) \quad \mathfrak{L}(x, D)\mathbf{u} := \sum_{|\alpha|=2b} \mathbf{A}_\alpha(x) D^\alpha \mathbf{u}(x) = \mathbf{f}(x)$$

for the unknown vector-valued function $\mathbf{u}: \Omega \rightarrow \mathbb{R}^m$ given by the transpose $\mathbf{u}(x) = (u_1(x), \dots, u_m(x))^T$, $m \geq 1$, $\mathbf{f}(x) = (f_1(x), \dots, f_m(x))^T$, where $\mathbf{A}_\alpha(x)$ is the $m \times m$ -matrix $\{a_\alpha^{jk}(x)\}_{j,k=1}^m$ and $a_\alpha^{jk}: \Omega \rightarrow \mathbb{R}$ are measurable functions. Hereafter, $b \geq 1$ is

1991 *Mathematics Subject Classification.* Primary 35J45; Secondary 35R05, 35B45, 35B65, 46E35.

Key words and phrases. Elliptic systems, a priori estimates, Hölder regularity, singular integrals.

a fixed integer, $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multiindex of length $|\alpha| = \alpha_1 + \dots + \alpha_n$ and $D^\alpha := D_1^{\alpha_1} \dots D_n^{\alpha_n}$ with $D_i := \partial/\partial x_i$. This way, the matrix differential operator $\mathfrak{L}(x, D)$ has entries

$$l^{jk}(x, D) := \sum_{|\alpha|=2b} a_\alpha^{jk}(x) D^\alpha$$

and for fixed j and k the polynomial

$$l^{jk}(x, \xi) := \sum_{|\alpha|=2b} a_\alpha^{jk}(x) \xi^\alpha, \quad \xi \in \mathbb{R}^n, \quad \xi^\alpha := \xi_1^{\alpha_1} \xi_2^{\alpha_2} \dots \xi_n^{\alpha_n},$$

is homogeneous of degree $2b$.

We suppose (1.1) to be an *elliptic system*, that is, the characteristic determinant of $\mathfrak{L}(x, \xi)$ is non-vanishing for a.a. $x \in \Omega$ and all $\xi \neq 0$. In view of the homogeneity of l^{jk} 's, this rewrites as (see [3, 5])

$$(1.2) \quad \exists \delta > 0: \quad \det \left\{ \sum_{|\alpha|=2b} \mathbf{A}_\alpha(x) \xi^\alpha \right\} \geq \delta |\xi|^{2bm} \quad \text{a.a. } x \in \Omega, \quad \forall \xi \in \mathbb{R}^n.$$

Our goal is to obtain interior Hölder regularity of the solutions to (1.1) as a byproduct of a priori estimates in Sobolev and Sobolev-Morrey spaces. Let us recall the definitions of these functional classes.

Definition 1.1. *The Sobolev space $W^{2b,p}(\Omega)$, $p \in (1, +\infty)$, is the collection of $L^p(\Omega)$ functions $u: \Omega \rightarrow \mathbb{R}$ all of which distribution derivatives $D^\alpha u$ with $|\alpha| \leq 2b$, belong to $L^p(\Omega)$. The norm in $W^{2b,p}(\Omega)$ is*

$$\|u\|_{W^{2b,p}(\Omega)} := \sum_{s=0}^{2b} \sum_{|\alpha|=s} \|D^\alpha u\|_{p;\Omega}, \quad \|\cdot\|_{p;\Omega} := \left(\int_\Omega |\cdot|^p dx \right)^{1/p}.$$

For the sake of brevity, the cross-product of m copies of $L^p(\Omega)$ is denoted by the same symbol. Thus, if $\mathbf{u} = (u_1, \dots, u_m)$ is a vector-valued function, $\mathbf{u} \in L^p(\Omega)$ means that $u_k \in L^p(\Omega)$ for all $k = 1, \dots, m$, and $\|\mathbf{u}\|_{p;\Omega} := \sum_{k=1}^m \|u_k\|_{p;\Omega}$.

Definition 1.2. *Let $p \in (1, +\infty)$ and $\lambda \in (0, n)$. The function $u \in L^p(\Omega)$ belongs to the Morrey space $L^{p,\lambda}(\Omega)$ if*

$$\|u\|_{p,\lambda;\Omega} := \left(\sup_{r>0} \frac{1}{r^\lambda} \int_{B_r \cap \Omega} |u(x)|^p dx \right)^{1/p} < \infty$$

where B_r ranges in the set of balls with radius r in $\mathbb{R}^n$. The Sobolev-Morrey space $W^{2b,p,\lambda}(\Omega)$ consists of all functions $u \in W^{2b,p}(\Omega)$ with generalized derivatives $D^\alpha u$, $|\alpha| \leq 2b$, belonging to $L^{p,\lambda}(\Omega)$. The norm in $W^{2b,p,\lambda}(\Omega)$ is given by

$$\|u\|_{W^{2b,p,\lambda}(\Omega)} := \sum_{s=0}^{2b} \sum_{|\alpha|=s} \|D^\alpha u\|_{p,\lambda;\Omega}.$$

We refer the reader to [1, 2, 9] for various properties of the Morrey and Sobolev–Morrey spaces.

Definition 1.3. For a locally integrable function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ define

$$\eta_f(R) := \sup_{r \leq R} \frac{1}{|B_r|} \int_{B_r} |f(y) - f_{B_r}| dy \quad \text{for every } R > 0,$$

where B_r ranges over the balls in $\mathbb{R}^n$ and $f_{B_r} = \frac{1}{|B_r|} \int_{B_r} f(y) dy$. Then:

- $f \in BMO$ (*bounded mean oscillation*, see John–Nirenberg [7]) if $\|f\|_* := \sup_R \eta_f(R) < +\infty$. $\|f\|_*$ is a norm in BMO modulo constant functions under which BMO is a Banach space.
- $f \in VMO$ (*vanishing mean oscillation*, see Sarason [11]) if $f \in BMO$ and $\lim_{R \downarrow 0} \eta_f(R) = 0$. The quantity $\eta_f(R)$ is referred to as VMO -modulus of f .

The spaces $BMO(\Omega)$ and $VMO(\Omega)$, and $\|\cdot\|_{*,\Omega}$ are defined in a similar manner taking $B_r \cap \Omega$ instead of B_r above.

As already mentioned, the desired Hölder bounds will be derived on the base of the following a priori estimate in Sobolev–Morrey classes.

Theorem 1.4. Suppose (1.2), $a_\alpha^{jk} \in VMO(\Omega) \cap L^\infty(\Omega)$, $\mathbf{f} \in L_{\text{loc}}^{p,\lambda}(\Omega)$, $1 < p < \infty$, $0 < \lambda < n$, and let $\mathbf{u} \in W_{\text{loc}}^{2b,p,\lambda}(\Omega)$ be a strong solution of (1.1). Then, for any $\Omega' \Subset \Omega'' \Subset \Omega$ there is a constant C depending on $n, m, b, p, \lambda, \delta, \|a_\alpha^{jk}\|_{\infty;\Omega}$, the VMO -moduli $\eta_{a_\alpha^{jk}}$ of the coefficients (cf. [3, 4]) and $\text{dist}(\Omega', \partial\Omega'')$, such that

$$(1.3) \quad \|\mathbf{u}\|_{W^{2b,p,\lambda}(\Omega')} \leq C (\|\mathbf{f}\|_{p,\lambda;\Omega''} + \|\mathbf{u}\|_{p,\lambda;\Omega''}).$$

It turns out, moreover, that the operator $\mathfrak{L}$ improves the integrability of solutions to (1.1). In fact, by means of standard homotopy arguments and making use of formula (2.1) (cf. [3], [10, Section 3]), it is easy to get

Corollary 1.5. Under the hypotheses of Theorem 1.4, suppose $\mathbf{u} \in W_{\text{loc}}^{2b,q}(\Omega)$ with $q \in (1, p]$. Then $\mathbf{u} \in W_{\text{loc}}^{2b,p,\lambda}(\Omega)$.

A combination of (1.3) with the embedding properties of Sobolev–Morrey spaces leads to a precise characterization of the Morrey, BMO and Hölder regularity of the solutions to (1.1).

Corollary 1.6. Under the hypotheses of Theorem 1.4 define s_0 as the least non-negative integer such that $\frac{n}{2b-s_0} > 1$ and fix an $s \in \{s_0, \dots, 2b-1\}$. Then there is a constant C such that:

- a) if $p \in (1, \frac{n-\lambda}{2b-s})$ then $D^s \mathbf{u} \in L^{p,(2b-s)p+\lambda}(\Omega')$ and

$$\|D^s \mathbf{u}\|_{p,(2b-s)p+\lambda;\Omega'} \leq C (\|\mathbf{f}\|_{p,\lambda;\Omega''} + \|\mathbf{u}\|_{p,\lambda;\Omega''});$$

b) if $p = \frac{n-\lambda}{2b-s}$ then $D^s \mathbf{u} \in BMO(\Omega')$ and


$$\|D^s \mathbf{u}\|_{BMO(\Omega')} \leq C (\|\mathbf{f}\|_{p,\lambda;\Omega''} + \|\mathbf{u}\|_{p,\lambda;\Omega''});$$

c) if $p \in (\frac{n-\lambda}{2b-s}, \frac{n-\lambda}{2b-s-1})^*$ then $D^s \mathbf{u} \in C^{0,\sigma_s}(\Omega')$ with $\sigma_s = 2b - s - \frac{n-\lambda}{p}$ and

$$\sup_{\substack{x \neq x' \\ x, x' \in \Omega'}} \frac{|D^s \mathbf{u}(x) - D^s \mathbf{u}(x')|}{|x - x'|^{\sigma_s}} \leq C (\|\mathbf{f}\|_{p,\lambda;\Omega''} + \|\mathbf{u}\|_{p,\lambda;\Omega''}).$$

If $s_0 \geq 1$ (i.e., $2b \geq n$) and $p \in (1, \frac{n-\lambda}{2b-s_0})$ then $\mathbf{u} \in C^{s_0-1, 2b-s_0+1-\frac{n-\lambda}{p}}(\Omega')$ and

$$\sup_{\substack{x \neq x' \\ x, x' \in \Omega'}} \frac{|D^{s_0-1} \mathbf{u}(x) - D^{s_0-1} \mathbf{u}(x')|}{|x - x'|^{2b-s_0+1-\frac{n-\lambda}{p}}} \leq C (\|\mathbf{f}\|_{p,\lambda;\Omega''} + \|\mathbf{u}\|_{p,\lambda;\Omega''}).$$



The picture illustrates geometrically the results of Corollary 1.6 with the couple (p, λ) lying in the semistrip $\{(p, \lambda): p > 1, 0 < \lambda < n\}$ and $s \in \{s_0, \dots, 2b-1\}$. The points B_s on the p -axis are $B_s = (\frac{n}{2b-s}, 0)$, $B = (1, 0)$, and $A_s = (1, n - 2b + s)$ is the intersection of the line through $(0, n)$ and B_s with the vertical line $\{p = 1\}$.

When (p, λ) belongs to the open right triangle BB_sA_s then we have case a), i.e. $D^s \mathbf{u} \in L^{p, (2b-s)p+\lambda}(\Omega')$. If (p, λ) lies on the open segment A_sB_s we have $D^s \mathbf{u} \in BMO(\Omega')$

*This rewrites as $p \in (n - \lambda, \infty)$ when $s = 2b - 1$.

(case b)). In particular, $(p, \lambda) \in \triangle BB_{s_0}A_{s_0}$ yields $D^{s_0}\mathbf{u} \in L^{p, (2b-s_0)p+\lambda}(\Omega')$, while $\mathbf{u} \in C^{s_0-1, 2b-s_0+1-\frac{n-\lambda}{p}}(\Omega')$ if $s_0 \geq 1$. Further on, $(p, \lambda) \in A_{s_0}B_{s_0}$ gives $D^{s_0}\mathbf{u} \in BMO(\Omega')$.

Let $s \in \{s_0, \dots, 2b-2\}$ and suppose (p, λ) lies in the interior of the quadrilateral $Q_s := B_s B_{s+1} A_{s+1} A_s$. Then $D^s \mathbf{u} \in C^{0, \sigma_s}(\Omega')$ (case c)) whereas $D^{s+1} \mathbf{u} \in L^{p, (2b-s-1)p+\lambda}(\Omega')$. Moreover, the exponent σ_s is the length of the segment CA_s where $C = C(p, \lambda)$ is the intersection of the line $\{p = 1\}$ with the line passing through the points (p, λ) and $(0, n)$. In particular, $(p, \lambda) \in A_{s+1}B_{s+1}$ implies $D^{s+1}\mathbf{u} \in BMO(\Omega')$.

Similarly, set Q_{2b-1} for the shadowed *unbounded region* on the picture. Then $(p', \lambda') \in Q_{2b-1}$ gives $D^{2b-1}\mathbf{u} \in C^{0, \sigma_{2b-1}}(\Omega')$ with σ_{2b-1} equals to the length of $C'A_{2b-1}$, $C' = C'(p', \lambda')$, while $D^{2b-1}\mathbf{u} \in BMO(\Omega')$ if $(p', \lambda') \in A_{2b-1}B_{2b-1}$.

2. NEWTONIAN-TYPE POTENTIALS

Fix the coefficients of (1.1) at a point $x_0 \in \Omega$ and consider the constant coefficients operator $\mathfrak{L}(x_0, D) := \sum_{|\alpha|=2b} \mathbf{A}_\alpha(x_0) D^\alpha$. The $2bm$ -order differential operator

$$L(x_0, D) := \det \mathfrak{L}(x_0, D) = \det \left\{ \sum_{|\alpha|=2b} a_\alpha^{jk}(x_0) D^\alpha \right\}$$

is elliptic in view of (1.2), and therefore there exists its fundamental solution $\tilde{\Gamma}(x_0; x - y)$ (see [5, 6]). If the space dimension n is an *odd* number, then

$$\tilde{\Gamma}(x_0; x - y) = |x - y|^{2bm-n} P \left(x_0; \frac{x - y}{|x - y|} \right)$$

where $P(x_0; \xi)$ is a real analytic function of $\xi \in \mathbb{S}^{n-1} := \{\xi \in \mathbb{R}^n : |\xi| = 1\}$. In case of *even* dimension n the procedure is standard and is based on introduction of a fictitious new variable x_{n+1} and extension of all functions as constants with respect to it, see [5] for details. Set $\{L_{jk}(x_0, \xi)\}_{j,k=1}^m$ for the *cofactor matrix* of $\{l^{jk}(x_0, \xi)\}_{j,k=1}^m$ and note that for all fixed $j, k = 1, \dots, m$, $L_{jk}(x_0, D)$ is either a differential operator of order $2b(m-1)$, or the operator of multiplication by 0. Making use of the identities

$$\sum_{k=1}^m l^{ik}(x_0, \xi) L_{jk}(x_0, \xi) = \delta_{ij} L(x_0, \xi)$$

with Kronecker's δ_{ij} , it is not hard to check (cf. [5, 12]) that the fundamental matrix $\Gamma(x_0; x) = \{\Gamma^{jk}(x_0; x)\}_{j,k=1}^m$ of $\mathfrak{L}(x_0, D)$ has entries $\Gamma^{jk}(x_0; x) = L_{kj}(x_0, D) \tilde{\Gamma}(x_0; x)$.

Let $r > 0$ be so small that $B_r = \{x \in \mathbb{R}^n : |x - x_0| < r\} \Subset \Omega$, and let $\mathbf{v} \in C_0^\infty(B_r)$. Then, employing

$$\mathfrak{L}(x_0, D)\mathbf{v}(x) = (\mathfrak{L}(x_0, D) - \mathfrak{L}(x, D))\mathbf{v}(x) + \mathfrak{L}(x, D)\mathbf{v}(x)$$

and using standard approach (cf. [3, 5]), we obtain a representation of $\mathbf{v}$ in terms of the Newtonian-type potentials

$$\mathbf{v}(x) = \int_{B_r} \mathbf{\Gamma}(x_0; x - y) \mathfrak{L}\mathbf{v}(y) dy + \int_{B_r} \mathbf{\Gamma}(x_0; x - y) (\mathfrak{L}(x_0, D) - \mathfrak{L}(y, D)) \mathbf{v}(y) dy.$$

Taking the $2b$ -order derivatives and then unfreezing the coefficients by putting $x_0 = x$, we get

$$\begin{aligned} (2.1) \quad D^\alpha \mathbf{v}(x) &= \underbrace{p.v. \int_{B_r} D^\alpha \mathbf{\Gamma}(x; x - y) \mathfrak{L}\mathbf{v}(y) dy}_{=: \mathfrak{K}_\alpha(\mathfrak{L}\mathbf{v})} \\ &+ \sum_{|\alpha'|=2b} p.v. \int_{B_r} D^\alpha \mathbf{\Gamma}(x; x - y) (\mathbf{A}_{\alpha'}(x) - \mathbf{A}_{\alpha'}(y)) D^{\alpha'} \mathbf{v}(y) dy \\ &\quad \underbrace{\hspace{10em}}_{=: \mathfrak{C}_\alpha[\mathbf{A}_{\alpha'}, D^{\alpha'} \mathbf{v}]} \\ &+ \int_{\mathbb{S}^{n-1}} D^{\beta^s} \mathbf{\Gamma}(x; y) \nu_s d\sigma_y \mathfrak{L}\mathbf{v}(x) \quad \forall \alpha: |\alpha| = 2b \end{aligned}$$

where the derivatives $D^\alpha \mathbf{\Gamma}(\cdot; \cdot)$ are taken with respect to the second variable, the multi-indices β^s are such that $\beta^s := (\alpha_1, \dots, \alpha_{s-1}, \alpha_s - 1, \alpha_{s+1}, \dots, \alpha_n)$, $|\beta^s| = 2b - 1$ and ν_s is the s -th component of the outer normal to $\mathbb{S}^{n-1}$.

Noting that each entry of the matrix $D^\alpha \mathbf{\Gamma}(x; y)$, $|\alpha| = 2b$, is a *Calderón-Zygmund kernel* (cf. [3, 4]), we have

Lemma 2.1. *Let $|\alpha| = |\alpha'| = 2b$ and $\mathbf{A}_\alpha \in L^\infty(\Omega)$. For each $p \in (1, \infty)$ and each $\lambda \in (0, n)$ there is a constant C depending on $n, m, b, \delta, \|\mathbf{A}_\alpha\|_{\infty; \Omega}, p$ and λ such that*

$$(2.2) \quad \|\mathfrak{K}_\alpha \mathbf{f}\|_{p, \lambda; \Omega} \leq C \|\mathbf{f}\|_{p, \lambda; \Omega}, \quad \|\mathfrak{C}_\alpha[\mathbf{A}_{\alpha'}, \mathbf{f}]\|_{p, \lambda; \Omega} \leq C \|\mathbf{A}_{\alpha'}\|_{*, \Omega} \|\mathbf{f}\|_{p, \lambda; \Omega}$$

for all $\mathbf{f} \in L^{p, \lambda}(\Omega)$. Moreover, let $\mathbf{A}_\alpha \in VMO(\Omega) \cap L^\infty(\Omega)$ with VMO -modulus $\eta_{\mathbf{A}_\alpha}$. Then for each $\varepsilon > 0$ there exists $r_0 = r_0(\varepsilon, \eta_{\mathbf{A}_\alpha})$ such that if $r < r_0$ we have

$$(2.3) \quad \|\mathfrak{C}_\alpha[\mathbf{A}_{\alpha'}, \mathbf{f}]\|_{p, \lambda; B_r} \leq C \varepsilon \|\mathbf{f}\|_{p, \lambda; B_r}$$

for all $B_r \Subset \Omega$ and all $\mathbf{f} \in L^{p, \lambda}(B_r)$.

Lemma 2.1 is proved in [10, Theorem 2.1, Corollary 2.7] in the general case of Calderón-Zygmund's kernels of mixed homogeneity.

3. PROOF OF THEOREM 1.4

Fix an arbitrary $x_0 \in \text{supp } \mathbf{u}$ and let $B_r := \{x \in \mathbb{R}^n : |x - x_0| < r\}$. The operators $\mathfrak{K}_\alpha$ and $\mathfrak{C}_\alpha[\mathbf{A}_{\alpha'}, \cdot]$ are bounded from L^p into itself (cf. [4]) and therefore the representation (2.1) still holds true (a.e. in Ω) for any function $\mathbf{v} \in W_0^{2b, p}(\Omega) := \text{closure}_{W^{2b, p}(\Omega)} C_0^\infty(\Omega)$

and moreover for $\mathbf{v} \in W_0^{2b,p}(\Omega) \cap W_{\text{loc}}^{2b,p,\lambda}(\Omega)$ as well. Let $\text{supp } \mathbf{v} \subset B_r$. In view of (2.1), (2.2) and (2.3), for each $\varepsilon > 0$ there exists $r_0(\varepsilon, \eta_{\mathbf{A}_\alpha})$ such that for $r < r_0$ one has

$$\|D^{2b}\mathbf{v}\|_{p,\lambda;B_r} \leq C(\|\mathfrak{L}\mathbf{v}\|_{p,\lambda;B_r} + \varepsilon\|D^{2b}\mathbf{v}\|_{p,\lambda;B_r})$$

whence, choosing ε small enough we obtain

$$(3.1) \quad \|D^{2b}\mathbf{v}\|_{p,\lambda;B_r} \leq C\|\mathfrak{L}\mathbf{v}\|_{p,\lambda;B_r}.$$

Let $\theta \in (0, 1)$, $\theta' = \theta(3 - \theta)/2 > \theta$ and define a cut-off function $\varphi(x) \in C_0^\infty(B_r)$ such that $\varphi(x) = 1$ for $x \in B_{\theta r}$ whereas $\varphi(x) = 0$ for $x \notin B_{\theta' r}$. It is clear that $|D^s \varphi| \leq C(s)[\theta(1 - \theta)r]^{-s}$ for any $1 \leq s \leq 2b$ because of $\theta' - \theta = \theta(1 - \theta)/2$. Applying (3.1) to $\mathbf{v}(x) := \varphi(x)\mathbf{u}(x) \in W_0^{2b,p}(B_r) \cap W^{2b,p,\lambda}(B_r)$, we get

$$\begin{aligned} \|D^{2b}\mathbf{u}\|_{p,\lambda;B_{\theta r}} &\leq \|D^{2b}\mathbf{v}\|_{p,\lambda;B_{\theta' r}} \leq C\|\mathfrak{L}\mathbf{v}\|_{p,\lambda;B_{\theta' r}} \\ &\leq C \left(\|\mathbf{f}\|_{p,\lambda;B_{\theta' r}} + \sum_{s=1}^{2b-1} \frac{\|D^{2b-s}\mathbf{u}\|_{p,\lambda;B_{\theta' r}}}{[\theta(1 - \theta)r]^s} + \frac{\|\mathbf{u}\|_{p,\lambda;B_{\theta' r}}}{[\theta(1 - \theta)r]^{2b}} \right). \end{aligned}$$

Hence, the choice of θ' and $\theta(1 - \theta) \leq 2\theta'(1 - \theta')$ imply

$$\begin{aligned} [\theta(1 - \theta)r]^{2b} \|D^{2b}\mathbf{u}\|_{p,\lambda;B_{\theta r}} &\leq C \left([\theta'(1 - \theta')r]^{2b} \|\mathbf{f}\|_{p,\lambda;B_{\theta' r}} \right. \\ &\quad \left. + \sum_{s=1}^{2b-1} [\theta'(1 - \theta')r]^s \|D^s \mathbf{u}\|_{p,\lambda;B_{\theta' r}} + \|\mathbf{u}\|_{p,\lambda;B_{\theta' r}} \right) \end{aligned}$$

Setting Θ_s for the weighted Morrey seminorms $\sup_{0 < \theta < 1} [\theta(1 - \theta)r]^s \|D^s \mathbf{u}\|_{p,\lambda;B_{\theta r}}$, $s \in \{0, \dots, 2b\}$, the last inequality rewrites as

$$(3.2) \quad \Theta_{2b} \leq C \left(r^{2b} \|\mathbf{f}\|_{p,\lambda;B_r} + \sum_{s=1}^{2b-1} \Theta_s + \Theta_0 \right).$$

To manage the seminorms Θ_s with $1 \leq s \leq 2b - 1$, we use the next interpolation inequality which follows from [12, (5.6)] as in [10, Proposition 3.2]:

Proposition 3.1. *There exists a constant $C = C(n, m, b, p, \lambda, s)$ independent of r and such that*

$$\Theta_s \leq \varepsilon \Theta_{2b} + \frac{C}{\varepsilon^{s/(2b-s)}} \Theta_0 \quad \forall \varepsilon \in (0, 2).$$

Therefore, interpolating the intermediate seminorms in (3.2) and fixing $\theta = 1/2$, we obtain the following *Caccioppoli-type* inequality

$$(3.3) \quad \|D^{2b}\mathbf{u}\|_{p,\lambda;B_{r/2}} \leq C(\|\mathbf{f}\|_{p,\lambda;B_r} + r^{-2b}\|\mathbf{u}\|_{p,\lambda;B_r}).$$

The estimate (1.3) follows from (3.3) by a finite covering of Ω' with balls $B_{r/2}$, $r < \text{dist}(\Omega', \partial\Omega'')$.

4. PROOF OF COROLLARY 1.6

To begin with, let $s = 2b - 1$ and $\Omega'_r = B_r \cap \Omega'$ with $2r < \text{dist}(\Omega', \partial\Omega'')$. Direct calculations based on the classical Poincaré inequality lead to

$$(4.1) \quad \frac{1}{r^{p+\lambda}} \int_{\Omega'_r} |D^{2b-1}\mathbf{u}(x) - (D^{2b-1}\mathbf{u})_{\Omega'_r}|^p dx \leq C(n, p, m) \|D^{2b}\mathbf{u}\|_{p, \lambda; \tilde{\Omega}'}^p$$

with suitable $\tilde{\Omega}'$ such that $\Omega' \Subset \tilde{\Omega}' \Subset \Omega''$ and $(\mathbf{f})_{\Omega'_r}$ standing for the integral average of $\mathbf{f}$ over Ω'_r . Taking the supremum with respect to r we get that $D^{2b-1}\mathbf{u}$ belongs to the Campanato space $\mathfrak{L}^{p, p+\lambda}(\Omega')$. Now, employing the embedding properties of Campanato spaces into Morrey, BMO and Hölder ones (cf. [1, Theorem 2.1]) and (1.3), we obtain as follows. If $p + \lambda < n$ then $D^{2b-1}\mathbf{u} \in L^{p, p+\lambda}(\Omega')$, $p + \lambda = n$ implies $D^{2b-1}\mathbf{u} \in BMO(\Omega')$, while $D^{2b-1}\mathbf{u} \in C^{0, \sigma_{2b-1}}(\Omega')$ with $\sigma_{2b-1} = 1 - (n - \lambda)/p$ when $p + \lambda > n$.

To prove the statement for any s , we run induction for decreasing values of s until s_0 . Thus, suppose $D^{s+1}\mathbf{u}$ satisfies the statement of Corollary 1.6. Then

$$\begin{aligned} \frac{1}{r^{(2b-s)p+\lambda}} \int_{\Omega'_r} |D^s\mathbf{u}(x) - (D^s\mathbf{u})_{\Omega'_r}|^p dx &\leq \|D^{s+1}\mathbf{u}\|_{p, (2b-s-1)p+\lambda; \tilde{\Omega}'}^p \\ &\leq C (\|\mathbf{f}\|_{p, \lambda; \Omega''}^p + \|\mathbf{u}\|_{p, \lambda; \Omega''}^p) \end{aligned}$$

whence $D^s\mathbf{u} \in \mathfrak{L}^{p, (2b-s)p+\lambda}(\Omega')$ and the conclusions follow as above.

Similarly, if $s_0 \geq 1$ and $p \in \left(1, \frac{n-\lambda}{2b-s_0}\right)$ we have $D^{s_0-1}\mathbf{u} \in \mathfrak{L}^{p, (2b-s_0+1)p+\lambda}(\Omega')$. It follows from the definition of s_0 that $n < (2b - s_0 + 1)p + \lambda < n + p$, whence $D^{s_0-1}\mathbf{u} \in C^{0, 2b-s_0+1-\frac{n-\lambda}{p}}(\Omega')$.

REFERENCES

- [1] S. Campanato, *Sistemi ellittici in forma divergenza. Regolarità all'interno*, Pubblicazioni della Classe di Scienze: Quaderni, Scuola Norm. Sup., Pisa, 1980.
- [2] S. Campanato, Proprietà di Hölderianità di alcune classi di funzioni. *Ann. Scuola Norm. Sup. Pisa* **17**, (1963) 175-188.
- [3] F. Chiarenza, M. Franciosi and M. Frasca, L^p -estimates for linear elliptic systems with discontinuous coefficients, *Rend. Accad. Naz. Lincei, Mat. Appl.*, **5** (1994) 27-32.
- [4] F. Chiarenza, M. Frasca and P. Longo, Interior $W^{2,p}$ estimates for non divergence elliptic equations with discontinuous coefficients, *Ric. Mat.*, **60** (1991) 149-168.
- [5] A. Douglis and L. Nirenberg, Interior estimates for elliptic systems of partial differential equations, *Commun. Pure Appl. Math.*, **8** (1955) 503-538.
- [6] F. John, *Partial Differential Equations*, Appl. Math. Sci., Vol. 1, Springer-Verlag, Berlin, 1991.
- [7] F. John and L. Nirenberg, On functions of bounded mean oscillation, *Commun. Pure Appl. Math.*, **14** (1961) 415-426.
- [8] A. Maugeri, D.K. Palagachev and L.G. Softova, *Elliptic and Parabolic Equations with Discontinuous Coefficients*, Wiley-VCH, Berlin, 2000.
- [9] C.B. Morrey, Jr., *Multiple Integrals in the Calculus of Variations*. Springer-Verlag, Berlin, 1966.
- [10] D. Palagachev and L. Softova, Singular integral operators, Morrey spaces and fine regularity of solutions to PDE's, *Potential Anal.*, **20** (2004) 237-263.

- [11] D. Sarason, Functions of vanishing mean oscillation, *Trans. Amer. Math. Soc.*, **207** (1975) 391–405.
- [12] V. A. Solonnikov, On the boundary value problems for linear parabolic systems of differential equations of general form, *Proc. Steklov Inst. Math.*, **83** (1965); English translation: O. A. Ladyzhenskaya (Ed.) *Boundary Value Problems of Mathematical Physics III*, Amer. Math. Soc., Providence, R.I., 1967.

POLITECNICO DI BARI, DIPARTIMENTO DI MATEMATICA, VIA E. ORABONA, 4, 70 125 BARI, ITALY
E-mail address: dian@dm.uniba.it

BULGARIAN ACADEMY OF SCIENCES, INSTITUTE OF MATHEMATICS, SOFIA, BULGARIA
E-mail address: luba@dm.uniba.it