# arXiv:math/0410463v4 [math.QA] 24 Jul 2006

# A QUANTUM ANALOGUE OF THE BERNSTEIN FUNCTOR

S. Sinel'shchikov<sup> $\dagger$ </sup> A. Stolin<sup> $\ddagger$ </sup> L. Vaksman<sup> $\dagger$ </sup>

<sup>†</sup>Mathematics Division, Institute for Low Temperature Physics & Engineering 47 Lenin Ave, 61103 Kharkov, Ukraine

<sup>‡</sup>Chalmers Tekniska Högskola, Mathematik, 412 96, Göteborg, Sweden

### Abstract

We consider Knapp-Vogan Hecke algebras in the quantum group setting. This allows us to produce a quantum analogue of the Bernstein functor as a first step towards the cohomological induction for quantum groups.

The cohomological induction is one of the most important tools for producing unitarizable Harish-Chandra modules [7]. The present paper demonstrates that this method can be applied to modules over quantum universal enveloping algebras [6] as well. Our way to this circle of problems was as follows.

The papers [11, 9] introduce quantum analogues of the bounded symmetric domains and the associated Harish-Chandra modules. One special case considered in [10] leads to two geometric realizations for the ladder representation of the quantum universal enveloping algebra  $U_q \mathfrak{su}_{2,2}$  and to the quantum Penrose transform [1]. One of the main tools used in [10] was studying q-analogues of the Čech cohomology. Unfortunately this method can be hardly generalized because it relates to non-trivial problems in non-commutative algebraic geometry. We hope to overcome these obstacles via replacing the q-analogues of the Čech cohomology by q-analogues of the Dolbeault cohomology. It is well-known that in the classical situation the Dolbeault cohomology can be constructed algebraically using the so-called cohomological induction (see [7]).

It is convenient to work with the Bernstein functor (the projective Zuckerman functor) for producing unitarizable Harish-Chandra modules. The definition of the Bernstein functor uses essentially the algebra of distributions on a real reductive group G with support in a maximal compact subgroup K. The principal obstacle to cope with in this work is to construct a q-analogue for this algebra.

In the present paper we construct a q-analogue of the algebra of distributions as above in certain cases. This allows us to construct a q-analogue of the Bernstein functor.

The authors are grateful to V. Kruglov for assistance in preparation of a preliminary version of this work and for helpful discussions.

# 1 The category $C(\mathfrak{l},\mathfrak{l})_q$ and the algebra $R(\mathfrak{l})_q$

We will use  $\mathbb{C}$  as the ground field.

Let  $\mathbf{a} = (a_{ij})_{i,j=1,2,\dots,l}$ , be the Cartan matrix for a simple complex Lie algebra of rank *l*. Its universal universal algebra is a unital algebra *U* determined by the generators  $\{H_i, E_i, F_i\}_{i,j=1,2,\dots,l}$  and the relations [6, p. 51]:

$$H_iH_j - H_jH_i = 0, \qquad E_iF_j - F_jE_i = \delta_{ij}H_i,$$

$$H_i E_j - E_j H_i = a_{ij} E_j, \qquad H_i F_j - F_j H_i = -a_{ij} F_j, \qquad i, j = 1, 2, \dots, l,$$

and

$$\sum_{m=0}^{1-a_{ij}} (-1)^m \binom{1-a_{ij}}{m} E_i^{1-a_{ij}-m} \cdot E_j \cdot E_i^m = 0,$$
$$\sum_{m=0}^{1-a_{ij}} (-1)^m \binom{1-a_{ij}}{m} F_i^{1-a_{ij}-m} \cdot F_j \cdot F_i^m = 0,$$

for all  $i \neq j$ .

The Cartan subalgebra spanned by  $H_1, H_2, \ldots, H_l$  is denoted by  $\mathfrak{h}$ , and  $\alpha_1, \alpha_2, \ldots, \alpha_l$ ,

$$\alpha_j(H_i) = a_{ij}, \qquad i, j = 1, 2, \dots, l,$$

are simple roots.

There exists a unique collection of coprime positive integers  $d_1, d_2, \ldots, d_l$  such that

$$d_i a_{ij} = d_j a_{ji}, \qquad i, j = 1, 2, \dots, l.$$

The bilinear form in  $\mathfrak{h}^*$  given by

$$(\alpha_i, \alpha_j) = d_i a_{ij}, \qquad i, j = 1, 2, \dots, l,$$

is positive definite.

Recall the definition of the quantum universal enveloping algebra  $U_q$  introduced by V. Drinfeld and M. Jimbo. We assume that  $q \in (0, 1)$ .<sup>1</sup>

The unital algebra  $U_q$  is determined by its generators  $K_i$ ,  $K_i^{-1}$ ,  $E_i$ ,  $F_i$ , i = 1, 2, ..., l, and the relations

$$K_{i}K_{j} = K_{j}K_{i}, \qquad K_{i}K_{i}^{-1} = K_{i}^{-1}K_{i} = 1,$$

$$K_{i}E_{j} = q_{i}^{a_{ij}}E_{j}K_{i}, \qquad K_{i}F_{j} = q_{i}^{-a_{ij}}F_{j}K_{i}, \qquad E_{i}F_{j} - F_{j}E_{i} = \delta_{ij}(K_{i} - K_{i}^{-1})/(q_{i} - q_{i}^{-1}),$$

$$\sum_{s=0}^{1-a_{ij}} (-1)^{s} \begin{bmatrix} 1 - a_{ij} \\ s \end{bmatrix}_{q_{i}} E_{i}^{1-a_{ij}-s}E_{j}E_{i}^{s} = 0,$$

$$\sum_{s=0}^{1-a_{ij}} (-1)^{s} \begin{bmatrix} 1 - a_{ij} \\ s \end{bmatrix}_{q_{i}} F_{i}^{1-a_{ij}-s}F_{j}F_{i}^{s} = 0,$$

with

$$q_i = q^{d_i}, \qquad 1 \le i \le l,$$

$$\begin{bmatrix} m \\ n \end{bmatrix}_q = \frac{[m]_q!}{[n]_q![m-n]_q!}, \qquad [n]_q! = [n]_q \dots [2]_q[1]_q, \qquad [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

<sup>1</sup>The purpose of so strong assumptions on q will become evident while investigating the unitarizability of the Harish-Chandra modules in question over  $U_q$ .

The Hopf algebra structure on  $U_q$  is given by the following formulas:

$$\Delta(E_i) = E_i \otimes 1 + K_i \otimes E_i, \qquad \Delta(F_i) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \qquad \Delta(K_i) = K_i \otimes K_i,$$
$$S(E_i) = -K_i^{-1}E_i, \qquad S(F_i) = -F_iK_i, \qquad S(K_i) = K_i^{-1},$$
$$\varepsilon(E_i) = \varepsilon(F_i) = 0, \qquad \varepsilon(K_i) = 1.$$

Any subset  $\mathbb{L} \subset \{1, 2, \dots, l\}$  determines a Hopf subalgebra  $U_q \mathfrak{l} \subset U_q$  generated by

 $K_i^{\pm 1}, \quad i=1,2,\ldots,l; \qquad E_j, F_j, \quad j \in \mathbb{L}.$ 

We use the following notation:

$$P = \mathbb{Z}^l, \qquad P_+ = \mathbb{Z}^l_+, \qquad P_+^{\mathbb{L}} = \{ \boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_l) \in P \mid \lambda_i \ge 0 \text{ for } i \in \mathbb{L} \}.$$

Let V be a  $U_q \mathfrak{l}$ -module and

$$V_{\mu} = \{ v \in V | K_i v = q_i^{\mu_i} v, \quad i = 1, 2, \dots, l \}$$

the weight space, where  $\mu = (\mu_1, \mu_2, \dots, \mu_l) \in P$ .

**Definition 1.1** We say V is a weight  $U_q \mathfrak{l}$ -module if  $V = \bigoplus_{\mu \in P} V_{\mu}$ .

**Example 1.2** Consider the Verma module  $U_q\mathfrak{l}$ -module  $M(\mathfrak{l}, \lambda)$  with the highest weight  $\lambda \in P_+^{\mathbb{L}}$ . This is a module determined by its generator  $v(\mathfrak{l}, \lambda)$  and the relations

$$E_j v(\mathfrak{l}, \lambda) = 0, \quad j \in \mathbb{L}; \qquad K_i^{\pm 1} v(\mathfrak{l}, \lambda) = q_i^{\pm \lambda_i} v(\mathfrak{l}, \lambda), \quad i = 1, 2, \dots, l.$$

It is easy to prove that its submodule  $K(\mathfrak{l},\lambda)$  generated by  $F_j^{\lambda_j+1}v(\mathfrak{l},\lambda)$ ,  $j \in \mathbb{L}$ , is the unique submodule of finite codimension. In particular, the factor-module  $L(\mathfrak{l},\lambda) = M(\mathfrak{l},\lambda)/K(\mathfrak{l},\lambda)$  is simple (see [6]).

The following claims are well-known (see for instance [6]). The dimensions of the weight subspaces of the  $U_q\mathfrak{l}$ -modules  $L(\mathfrak{l}, \lambda)$  are the same as in the classical case q = 1. The weight finite dimensional  $U_q\mathfrak{l}$ -modules are completely reducible. Irreducible weight finite dimensional  $U_q\mathfrak{l}$ -modules are isomorphic to  $L(\mathfrak{l}, \lambda)$  with  $\lambda \in P_+^{\mathbb{L}}$ . The center of  $U_q\mathfrak{l}$  (denoted by  $Z(U_q\mathfrak{l})$ ) admits an easy description via the Harish-Chandra isomorphism [6, p. 109] and separates the simple weight finite dimensional  $U_q\mathfrak{l}$ -modules [6, p. 125]. In other words, for any  $\lambda_1, \lambda_2 \in P_+^{\mathbb{L}}$  there exists  $z \in Z(U_q\mathfrak{l})$  such that  $z|_{L(\mathfrak{l},\lambda_1)} \neq z|_{L(\mathfrak{l},\lambda_2)}$ .

A module V over  $U_q \mathfrak{l}$  is called locally finite dimensional if  $\dim(U_q \mathfrak{l} v) < \infty$  for all  $v \in V$ . Introduce the notation  $C(\mathfrak{l}, \mathfrak{l})_q$  for the full subcategory of weight locally finite dimensional  $U_q \mathfrak{l}$ -modules.

It is easy to prove that any  $U_q\mathfrak{l}$ -module  $V \in C(\mathfrak{l}, \mathfrak{l})_q$  is uniquely decomposable as a direct sum  $V = \bigoplus_{\lambda \in P_+^{\mathbb{L}}} V_\lambda$  of its submodules. Each  $V_\lambda$  here is a multiple of  $L(\mathfrak{l}, \lambda), \lambda \in P_+^{\mathbb{L}}$ .

The sum is direct since  $Z(U_q\mathfrak{l})$  separates simple finite dimensional weight  $U_q\mathfrak{l}$ -modules.

Let End V be the algebra of all endomorphisms of V considered as a vector space. Now consider the  $U_q\mathfrak{l}$ -module  $L^{\text{univ}} = \bigoplus_{\lambda \in P_+^{\mathbb{L}}} L(\mathfrak{l}, \lambda)$  and the projections  $P_{\lambda}$  in  $L^{\text{univ}}$  onto

the isotypic component  $L(\mathfrak{l}, \lambda)$  parallel to the sum of all other isotypic components. The

algebra End  $L^{\text{univ}}$  is a  $U_q\mathfrak{l}$ -module algebra.<sup>2</sup> The natural homomorphism  $U_q\mathfrak{l} \to \text{End } L^{\text{univ}}$  is injective [6, p. 76 – 77]; this allows one to identify  $U_q\mathfrak{l}$  with its image in End  $L^{\text{univ}}$ . We are interested in the following  $U_q\mathfrak{l}$ -module subalgebras of End  $L^{\text{univ}}$ :

$$R(\mathfrak{l})_q \stackrel{\text{def}}{=} \bigoplus_{\lambda \in P_+^{\mathbb{L}}} \operatorname{End} L(\mathfrak{l}, \lambda), \qquad F(\mathfrak{l})_q \stackrel{\text{def}}{=} R(\mathfrak{l})_q \oplus U_q \mathfrak{l}.^3$$

Obviously,  $R(\mathfrak{l})_q$  is a two-sided ideal of  $F(\mathfrak{l})_q$  generated by the projections  $P_{\lambda}$ .

 $R(\mathfrak{l})_q$  is not a unital algebra, but it admits a distinguished approximate identity. Specifically, any finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  determines an idempotent  $\chi_{\Lambda} = \sum_{\lambda \in \Lambda} P_{\lambda}$  in  $R(\mathfrak{l})_q$ . Obviously,  $\chi_{\Lambda_1}\chi_{\Lambda_2} = \chi_{\Lambda_2}\chi_{\Lambda_1} = \chi_{\Lambda_1 \cap \Lambda_2}$ , and for any  $r \in R(\mathfrak{l})_q$  there exists a finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  such that  $\chi_{\Lambda}r = r\chi_{\Lambda} = r$ . With some abuse of terminology, one can say that in  $F(\mathfrak{l})_q$  we have

$$\lim_{\Lambda\uparrow P_+^{\mathbb{L}}}\chi_{\Lambda}=1.$$

A module V over  $R(\mathfrak{l})_q$  is called approximately unital if for any  $v \in V$  there exists a finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  such that  $\chi_{\Lambda} v = v$ .

We are going to demonstrate that the full subcategory of approximately unital  $R(\mathfrak{l})_q$ -modules is canonically isomorphic to  $C(\mathfrak{l},\mathfrak{l})_q$ .

**Proposition 1.3** (\*). For any approximately unital  $R(\mathfrak{l})_q$ -module V and any  $\xi \in F(\mathfrak{l})_q$ ,  $v \in V$  there exists a limit

$$\xi v \stackrel{\text{def}}{=} \lim_{\Lambda \uparrow P_+^{\mathbb{L}}} (\xi \chi_{\Lambda}) v \tag{1.1}$$

in the sense that  $(\xi \chi_{\Lambda}) v$  does not depend on  $\Lambda$  for large  $\Lambda$ .

(\*\*). The relation (1.1) equips V with a structure of  $F(\mathfrak{l})_q$ -module and defines a functor from the category of approximately unital  $R(\mathfrak{l})_q$ -modules into the category  $C(\mathfrak{l},\mathfrak{l})_q$ . (\*\*\*). This functor is an isomorphism of categories.

**Proof.** Any approximately unital  $R(\mathfrak{l})_q$ -module V admits an embedding into an approximately unital  $R(\mathfrak{l})_q$ -module that is a multiple of  $L^{\text{univ}}$ . This observation implies the first two statements, since they can be easily verified for  $L^{\text{univ}}$ .

Any  $U_q\mathfrak{l}$ -module that is a multiple of  $L^{\text{univ}}$ , is a  $F(\mathfrak{l})_q$ -module. This allows one to construct an inverse functor by embedding any  $U_q\mathfrak{l}$ -module  $V \in C(\mathfrak{l}, \mathfrak{l})_q$  into a  $U_q\mathfrak{l}$ -module that is a multiple of  $L^{\text{univ}}$ . This implies (\*\*\*).

Note that the action of  $P_{\lambda} \in R(\mathfrak{l})_q$  on vectors  $v \in V \in C(\mathfrak{l}, \mathfrak{l})_q$  can be described without referring to an embedding into a  $U_q\mathfrak{l}$ -module multiple to  $L^{\mathrm{univ}}$ . More exactly, for any  $\lambda \in P_+^{\mathbb{L}}$  there exists a sequence  $z_1(\lambda), z_2(\lambda), \ldots$  of elements of the center of  $U_q\mathfrak{l}$  such that  $P_{\lambda}v = \lim_{n \to \infty} z_n(\lambda)v$  for all  $v \in V \in C(\mathfrak{l}, \mathfrak{l})_q$ . To see this, note that for any finite subset  $\lambda \in \Lambda \subset P_+^{\mathbb{L}}$  there exists such  $z \in Z(U_q\mathfrak{l})$  that  $z|_{L(\mathfrak{l},\mu)} = 0$  for  $\lambda \in \Lambda \setminus \{\lambda\}$  and  $z|_{L(\mathfrak{l},\lambda)} = 1$ . It remains to choose a sequence  $\Lambda_n \uparrow P_+^{\mathbb{L}}$ .

<sup>2</sup>If  $\xi \in U_q \mathfrak{l}$  and  $\Delta \xi = \sum_i \xi'_i \otimes \xi''_i$ , then  $(\xi A)v = \sum_i \xi'_i AS(\xi''_i)v$ ,  $v \in L^{\text{univ}}$ ,  $A \in \text{End } L^{\text{univ}}$ , with  $\Delta$  and S being the comultiplication and the antipode of  $U_q \mathfrak{l}$ .

<sup>&</sup>lt;sup>3</sup>It is easy to prove that  $R(\mathfrak{l})_q \cap U_q \mathfrak{l} = 0$ . In fact,  $U_q \mathfrak{l}$  has no zero divisors [3], while the Harish-Chandra isomorphism allows one to produce for any  $a \in R(\mathfrak{l})_q$  a non-zero element  $z \in Z(U_q \mathfrak{l})$  such that za = 0.

## 2 The category $C(\mathfrak{g},\mathfrak{l})_q$ and the algebra $R(\mathfrak{g},\mathfrak{l})_q$

Let  $\mathbb{G} \supset \mathbb{L}$  be a pair of subsets of  $\{1, 2, \ldots, l\}$  and let  $U_q \mathfrak{g} \supset U_q \mathfrak{l}$  be the associated pair of Hopf subalgebras of  $U_q$ .

Consider the category of all  $U_q \mathfrak{g}$ -modules and its full subcategory  $C(\mathfrak{g}, \mathfrak{l})_q$  that consists of weight  $U_q \mathfrak{l}$ -locally finite dimensional  $U_q \mathfrak{g}$ -modules.

Let us turn to a description of the algebra  $F(\mathfrak{g}, \mathfrak{l})_q$ . This algebra is an important tool in studying  $U_q\mathfrak{g}$ -modules from  $C(\mathfrak{g}, \mathfrak{l})_q$ . First, we consider a simple special case, namely  $\mathbb{G} = \mathbb{L}$ . In this situation we define  $F(\mathfrak{g}, \mathfrak{l})_q$  as  $F(\mathfrak{l})_q$ . We intend to construct  $F(\mathfrak{g}, \mathfrak{l})_q$ along with embeddings  $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ ,  $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ , which possess the following properties.

(F1) The diagram



commutes<sup>4</sup>, and the subalgebras  $U_q \mathfrak{g}$  and  $F(\mathfrak{l})_q$  generate the algebra  $F(\mathfrak{g}, \mathfrak{l})_q$ .

- (F2) The approximate identity  $\{\chi_{\Lambda}\}$  of  $R(\mathfrak{l})_q$  is also an approximate identity of the twosided ideal  $R(\mathfrak{g}, \mathfrak{l})_q$  of  $F(\mathfrak{g}, \mathfrak{l})_q$ , generated by  $R(\mathfrak{l})_q$ .
- (F3) For any approximately unital  $R(\mathfrak{g}, \mathfrak{l})_q$ -module V and any  $\xi \in F(\mathfrak{g}, \mathfrak{l})_q$ ,  $v \in V$  there exists a limit

$$\xi v \stackrel{\text{def}}{=} \lim_{\Lambda \uparrow P_{+}^{\mathbb{L}}} (\xi \chi_{\Lambda}) v, \qquad (2.1)$$

and the equation (2.1) equips V with a structure of  $F(\mathfrak{g}, \mathfrak{l})_q$ -module.

The intersection of the kernels for all representations of  $F(\mathfrak{g}, \mathfrak{l})_q$  obtained in this way is 0.

(F4) The functor from the category of approximately unital  $R(\mathfrak{g}, \mathfrak{l})_q$ -modules into  $C(\mathfrak{g}, \mathfrak{l})_q$  given by (2.1) is an isomorphism of categories.

We are going to show that the algebra  $F(\mathfrak{g},\mathfrak{l})_q$  and the embeddings  $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g},\mathfrak{l})_q$ ,  $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g},\mathfrak{l})_q$  are essentially uniquely determined by (F1) – (F4), i.e. there exists a unique isomorphism of algebras that respects the embeddings.

Assign to any  $U_q\mathfrak{l}$ -module V a  $U_q\mathfrak{g}$ -module  $P(V) = U_q\mathfrak{g} \otimes_{U_q\mathfrak{l}} V$ . The following result is well-known in the classical case q = 1 (see [7]). It is proved in the Appendix (see Corollary A.3).

**Lemma 2.1** 1. If  $V \in C(\mathfrak{l}, \mathfrak{l})_q$  then P(V) is a projective object in  $C(\mathfrak{g}, \mathfrak{l})_q$ . 2. For any  $V \in C(\mathfrak{g}, \mathfrak{l})_q$ , the map

$$P(V) \mapsto V, \qquad \xi \otimes v \mapsto \xi v, \qquad \xi \in U_q \mathfrak{g}, \quad v \in V,$$

is a surjective morphism of  $U_q \mathfrak{g}$ -modules.

<sup>&</sup>lt;sup>4</sup>In what follows  $U_q\mathfrak{g}$  and  $F(\mathfrak{l})_q$  are identified to their images under the embedding into  $F(\mathfrak{g},\mathfrak{l})_q$ .

The projective objects P(V) of the category  $C(\mathfrak{g}, \mathfrak{l})_q$  are called standard projective objects.

Introduce the  $U_q \mathfrak{g}$ -module  $V^{\text{univ}} = P(L^{\text{univ}})$ .

**Corollary 2.2** In the category of  $U_q \mathfrak{l}$ -modules one has

$$V^{\mathrm{univ}} = \bigoplus_{\lambda \in P_+^{\mathbb{L}}} V_{\lambda}^{\mathrm{univ}},$$

where the  $U_q \mathfrak{l}$ -modules  $V_{\lambda}^{\text{univ}}$  are multiples of  $L(\mathfrak{l}, \lambda)$ .

**Corollary 2.3** For any module  $M \in C(\mathfrak{g}, \mathfrak{l})_q$  there exists a surjective morphism  $\widetilde{V}^{\text{univ}} \to M$  in the category  $C(\mathfrak{g}, \mathfrak{l})_q$ , with  $\widetilde{V}^{\text{univ}}$  being a multiple of the module  $V^{\text{univ}}$ .

Let us introduce the notation  $\mathcal{P}_{\lambda}$  for the projection in  $V^{\text{univ}}$  onto the  $U_q$ l-isotypic component  $V_{\lambda}^{\text{univ}}$  parallel to the sum of all other  $U_q$ l-isotypic components.

The category  $C(\mathfrak{g}, \mathfrak{l})_q$  and the category of approximately unital  $R(\mathfrak{g}, \mathfrak{l})_q$ -modules are closed under the operations of direct sums, passage to submodules and factor-modules. This allows one, using Lemma 2.1 and Corollary 2.3, to prove the following statement, which implies uniqueness of  $F(\mathfrak{g}, \mathfrak{l})_q$  and the embeddings  $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ ,  $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ .

**Proposition 2.4** Consider the algebra  $F(\mathfrak{g},\mathfrak{l})_q$  and the embeddings  $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g},\mathfrak{l})_q$ ,  $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g},\mathfrak{l})_q$  with the properties (F1) - (F4). Then the following statements hold.

1. The representation of  $U_q \mathfrak{g}$  in  $V^{\text{univ}}$  determines an approximately unital representation of  $R(\mathfrak{g}, \mathfrak{l})_q$  and a faithful representation of  $F(\mathfrak{g}, \mathfrak{l})_q$  in  $V^{\text{univ}}$ .

2. The image of  $F(\mathfrak{g},\mathfrak{l})_q$  under the embedding into  $\operatorname{End} V^{\operatorname{univ}}$  is the subalgebra of  $\operatorname{End} V^{\operatorname{univ}}$  generated by elements of  $U_q\mathfrak{g} \subset \operatorname{End} V^{\operatorname{univ}}$  and projections  $\mathfrak{P}_{\lambda}, \lambda \in P_+^{\mathbb{L}}$ .

Now we return to the construction of the algebra  $F(\mathfrak{g}, \mathfrak{l})_q$ . In what follows we will identify the algebras  $U_q\mathfrak{g}$ ,  $F(\mathfrak{l})_q$  with their images under the embeddings into End  $V^{\text{univ},5}$ . In particular,

 $P_{\lambda} \mapsto \mathcal{P}_{\lambda}, \quad \lambda \in P_{+}^{\mathbb{L}}.$ 

The following auxiliary statement is proved in the Appendix.

**Lemma 2.5** Given any  $\xi \in U_q \mathfrak{g}$ ,  $\lambda \in P_+^{\mathbb{L}}$ , there exists a finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  such that

$$\xi \mathfrak{P}_{\lambda} = \chi_{\Lambda} \xi \mathfrak{P}_{\lambda}, \qquad \mathfrak{P}_{\lambda} \xi = \mathfrak{P}_{\lambda} \xi \chi_{\Lambda}.$$

Consider the smallest subalgebra  $F(\mathfrak{g}, \mathfrak{l})_q \subset \operatorname{End} V^{\operatorname{univ}}$  that contains all the elements of  $U_q\mathfrak{g}$  and all the projections  $\mathcal{P}_{\lambda}, \lambda \in P_+^{\mathbb{L}}$ .

**Theorem 2.6** The algebra  $F(\mathfrak{g}, \mathfrak{l})_q$  along with the embeddings  $U_q\mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ ,  $F(\mathfrak{l})_q \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q$ , satisfy (F1) - (F4).

<sup>&</sup>lt;sup>5</sup>The representation of  $U_q\mathfrak{g}$  in  $V^{\text{univ}}$  is faithful since for any weight finite dimensional  $U_q\mathfrak{g}$ -module W there exists a surjective morphism of  $U_q\mathfrak{g}$ -modules  $V^{\text{univ}} \to W$  (Corollary 2.3). The representation of  $F(\mathfrak{l})_q$  in  $V^{\text{univ}}$  is faithful since  $L^{\text{univ}} \hookrightarrow V^{\text{univ}}$ .

**Proof.** (F1) is obviously satisfied.

Let us prove that  $\{\chi_{\Lambda}\}$  is a left approximate identity in  $R(\mathfrak{g},\mathfrak{l})_q$ . Every  $a \in R(\mathfrak{g},\mathfrak{l})_q$ has the form  $a = \sum_{i=1}^{N(a)} \xi_i \mathcal{P}_{\lambda_i} \eta_i$ , with  $\xi_i \in U_q \mathfrak{g}, \eta_i \in F(\mathfrak{g}, \mathfrak{l})_q, \lambda_i \in P_+^{\mathbb{L}}$ . It follows from Lemma 2.5 that  $\chi_{\Lambda}\xi_i\mathcal{P}_{\lambda_i} = \xi_i\mathcal{P}_{\lambda_i}$  for some finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  and all  $i = 1, 2, \ldots, N(a)$ . Hence  $\chi_{\Lambda}a = a$ . One can prove in a similar way that  $\{\chi_{\Lambda}\}$  is a right approximate identity in  $R(\mathfrak{g},\mathfrak{l})_q$ . Hence (F2) is satisfied.

Consider an approximately unital  $R(\mathfrak{g},\mathfrak{l})_q$ -module V. Given any vector  $v \in V$ , there exists such finite subset  $\Lambda \subset P^{\mathbb{L}}_+$  that  $\chi_{\Lambda} v = v$ . Thus for all  $\Lambda' \supset \Lambda$  we have

$$(\xi\chi_{\Lambda'})v = (\xi\chi_{\Lambda'})\chi_{\Lambda}v = (\xi\chi_{\Lambda'}\chi_{\Lambda})v = (\xi\chi_{\Lambda})v,$$

and hence the element

$$\xi v \stackrel{\text{def}}{=} \lim_{\Lambda \uparrow P_+^{\mathbb{L}}} (\xi \chi_\Lambda) v$$

is well defined. It is easy to prove that  $(\xi\eta)v = \xi(\eta v)$  for any  $\xi, \eta \in F(\mathfrak{g}, \mathfrak{l})_q, v \in V$ . In fact, there exist such finite subsets  $\Lambda', \Lambda'' \subset P_+^{\mathbb{L}}$  that

$$\chi_{\Lambda'}v = v, \qquad \chi_{\Lambda''}\eta\chi_{\Lambda'} = \eta\chi_{\Lambda'},$$

because  $\eta \chi_{\Lambda'} \in R(\mathfrak{g}, \mathfrak{l})_q$  and  $\{\chi_{\Lambda}\}$  is an approximate identity. Hence,

$$\begin{aligned} (\xi\eta)v &= \lim_{\Lambda\uparrow P^{\mathbb{L}}_{+}} (\xi\eta\chi_{\Lambda})v = \xi(\eta\chi_{\Lambda'})v = (\xi\chi_{\Lambda''})(\eta\chi_{\Lambda'})v = \lim_{\Lambda\uparrow P^{\mathbb{L}}_{+}} (\xi\chi_{\Lambda})(\eta\chi_{\Lambda'})v = \\ &= \xi(\eta\chi_{\Lambda'}v) = \xi(\eta v). \end{aligned}$$

It follows that V is an  $F(\mathfrak{g},\mathfrak{l})_{\mathfrak{g}}$ -module, and the first of the conditions (F3) is satisfied. The second condition, which requires the existence of a faithful representation as above, obviously holds.

The arguments below show that the  $F(\mathfrak{g},\mathfrak{l})_q$ -modules in question are locally  $U_q\mathfrak{l}$ -finite dimensional and weight:

$$\dim(U_q\mathfrak{l}\,v) = \dim U_q\mathfrak{l}(\chi_{\Lambda'}v) = \dim(U_q\mathfrak{l}\,\chi_{\Lambda'})v \le \dim R(\mathfrak{l})_q v < \infty.$$

Thus we have constructed a functor from the category of approximately unital  $R(\mathfrak{g}, \mathfrak{l})_{q}$ modules into  $C(\mathfrak{g},\mathfrak{l})_q$ . What remains is to construct an inverse functor.

Suppose  $V \in C(\mathfrak{g}, \mathfrak{l})_q$  and let  $V = \bigoplus V_{\mu}$  be a decomposition of V into a sum of  $\mu \in P_{\perp}^{\mathbb{L}}$ 

 $U_q$ l-isotypic components. Let  $\pi$  be the representation of  $U_q \mathfrak{g}$  in V, and denote by  $\Pi_{\lambda}$  the projection in V onto the isotypic component of  $V_{\lambda}$  parallel to  $\bigoplus V_{\mu}$ .

 $\substack{\mu \in P_+^{\mathbb{L}} \\ \mu \neq \lambda }$ It suffices to prove existence and uniqueness of an extension  $\tilde{\pi}$  of  $\pi$  to  $F(\mathfrak{g},\mathfrak{l})_q$  that possesses the following properties:

- i)  $\widetilde{\pi}(\mathcal{P}_{\lambda}) = \Pi_{\lambda}, \ \lambda \in P_{+}^{\mathbb{L}},$
- ii)  $\widetilde{\pi}|_{R(\mathfrak{g},\mathfrak{l})_q}$  is an approximately unital representation of  $R(\mathfrak{g},\mathfrak{l})_q$ , iii)  $\pi(\xi)v = \lim_{\Lambda\uparrow P_+^{\mathbb{L}}} \widetilde{\pi}(\xi\chi_{\Lambda})v, v \in V, \xi \in U_q\mathfrak{g}$ .

Uniqueness of such extension is obvious. To prove its existence, we consider subsequently the following cases:

1.  $V = V^{\text{univ}};$ 

2.  $V = \widetilde{V}^{\text{univ}}$ , with  $\widetilde{V}^{\text{univ}}$  being a multiple of the  $U_q \mathfrak{g}$ -module  $V^{\text{univ}}$ ;

3. V is a submodule of  $\widetilde{V}^{\text{univ}}$ ;

- 4. V is a standard projective object in  $C(\mathfrak{g}, \mathfrak{l})_q$ ;
- 5. (the general case)  $V \in C(\mathfrak{g}, \mathfrak{l})_q$ .

In the case 1) the desired statement follows from the definition of  $R(\mathfrak{g}, \mathfrak{l})_q$ . A passage from 1) to 2) and to 3) is obvious. To pass from 3) to 4), it is sufficient to use the fact that every  $U_q\mathfrak{g}$ -module  $P(V), V \in C(\mathfrak{l}, \mathfrak{l})_q$ , admits an embedding into a multiple  $\widetilde{V}^{\mathrm{univ}}$  of the  $U_q\mathfrak{g}$ -module  $V^{\mathrm{univ}}$ . This is because every locally finite dimensional  $U_q\mathfrak{l}$ -module admits an embedding into a  $U_q\mathfrak{l}$ -module  $\widetilde{L}^{\mathrm{univ}}$ , which is a multiple of  $L^{\mathrm{univ}}$  and has the property  $P(\widetilde{L}^{\mathrm{univ}}) = \widetilde{V}^{\mathrm{univ}}$ . Finally, to pass from 4) to 5), one can apply the existence of a surjective morphism  $P(V) \to V$  in the category  $C(\mathfrak{g}, \mathfrak{l})_q$ .  $\Box$ 

Recall that the Weyl group W is generated by simple reflections  $s_1, s_2, \ldots, s_l$  and acts on the weight lattice P.

**Proposition 2.7** 1. There exists a unique one-dimensional representation  $\tilde{\varepsilon}$  of  $F(\mathfrak{g}, \mathfrak{l})_q$  with

$$\widetilde{\varepsilon}(\xi) = \varepsilon(\xi), \quad \xi \in U_q \mathfrak{g}; \qquad \widetilde{\varepsilon}(P_\lambda) = \begin{cases} 1, & \lambda = 0, \\ 0, & \lambda \in P_+^{\mathbb{L}} \setminus \{0\} \end{cases}$$

2. There exists a unique anti-automorphism  $\widetilde{S}$  of the algebra  $F(\mathfrak{g},\mathfrak{l})_q$  such that

$$\widetilde{S}(\xi) = S(\xi), \quad \xi \in U_q \mathfrak{g}; \qquad \widetilde{S}(P_{\lambda}) = P_{-w_0^{\mathbb{L}}\lambda}, \quad \lambda \in P_+^{\mathbb{L}},$$

where  $w_0^{\mathbb{L}}$  is the longest element of the Weyl group  $W_{\mathbb{L}} \subset W$  generated by the simple reflections  $s_i, i \in \mathbb{L}$ .

**Proof.** Uniqueness of  $\tilde{\varepsilon}$  and  $\tilde{S}$  is obvious. Existence of  $\tilde{\varepsilon}$  has been demonstrated in the proof of Theorem 2.6. Turn to proving existence of  $\tilde{S}$ . In the special case  $\mathbb{G} = \mathbb{L}$  it follows from the definition of the algebra  $F(\mathfrak{l})_q \hookrightarrow \operatorname{End} L^{\operatorname{univ}}$  and the well-known isomorphism [2, p. 168]

$$L(\mathfrak{l},\lambda)^* \xrightarrow{\sim} L(\mathfrak{l},-w_0^{\mathbb{L}}\lambda), \qquad v(\mathfrak{l},\lambda) \mapsto v(\mathfrak{l},w_0^{\mathbb{L}}\lambda).$$

To pass from the special case  $\mathbb{G} = \mathbb{L}$  to the general case  $\mathbb{G} \supset \mathbb{L}$ , consider the algebra  $F(\mathfrak{g}, \mathfrak{l})_q^{\mathrm{op}}$  which is derived from  $F(\mathfrak{g}, \mathfrak{l})_q$  by replacement of the multiplication by the opposite one.  $U_q\mathfrak{g}$  can be embedded into  $F(\mathfrak{g}, \mathfrak{l})_q^{\mathrm{op}}$ :

$$U_q \mathfrak{g} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q^{\mathrm{op}}, \qquad \xi \mapsto S(\xi).$$

The algebra  $F(\mathfrak{l})_q$  also admits an embedding into  $F(\mathfrak{g},\mathfrak{l})_q^{\mathrm{op}}$ :

$$U_q \mathfrak{l} \hookrightarrow F(\mathfrak{g}, \mathfrak{l})_q^{\mathrm{op}}, \quad \xi \mapsto S(\xi); \qquad P_\lambda \mapsto P_{-w_0^{\mathbb{L}}\lambda}.$$

It is easy to verify (F1) - (F4) for this pair of embeddings. What remains is to use the uniqueness of such pair, Proposition 2.4.

The category  $C(\mathfrak{g}, \mathfrak{l})_q$  is canonically isomorphic to the category of approximately unital  $R(\mathfrak{g}, \mathfrak{l})_q$ -modules. In what follows we identify these categories.

Consider the Hopf subalgebra  $U_q\mathfrak{q}_{\mathbb{L}}^+ \subset U_q\mathfrak{g}$  generated by

 $E_i, \quad i \in \mathbb{G}; \qquad F_j, \quad j \in \mathbb{L}; \qquad K_m^{\pm 1}, \quad m = 1, 2, \dots, l.$ 

and the Hopf subalgebra  $U_q \mathfrak{q}_{\mathbb{L}}^- \subset U_q \mathfrak{g}$  generated by

$$E_i, \quad i \in \mathbb{L}; \qquad F_j, \quad j \in \mathbb{G}; \qquad K_m^{\pm 1}, \quad m = 1, 2, \dots, l.$$

The following statement is well known in the classical case q = 1 (see [7, p. 90]).

### **Proposition 2.8** The linear maps

$$U_{q}\mathfrak{g} \otimes_{U_{q}\mathfrak{l}} R(\mathfrak{l})_{q} \to R(\mathfrak{g},\mathfrak{l})_{q}, \qquad \xi \otimes r \mapsto \xi r,$$

$$R(\mathfrak{l})_{q} \otimes_{U_{q}\mathfrak{l}} U_{q}\mathfrak{g} \to R(\mathfrak{g},\mathfrak{l})_{q}, \qquad r \otimes \xi \mapsto r\xi,$$

$$(2.2)$$

are bijective.

**Proof.** By Proposition 2.7, it suffices to consider the linear map (2.2). Let us prove that it is onto. Note that for any  $\lambda \in P_+^{\mathbb{L}}$  and any finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  containing  $\lambda$  there exists an element z in the center  $Z(U_q\mathfrak{l})$  of the algebra  $U_q\mathfrak{l}$  such that  $P_{\lambda} = z\chi_{\Lambda}$ . On the other hand, for any  $\lambda \in P_+^{\mathbb{L}}$ ,  $\xi \in U_q\mathfrak{g}$  there exist finite subsets  $\Lambda', \Lambda'' \ni \lambda$  such that

$$P_{\lambda}\xi = P_{\lambda}\xi\chi_{\Lambda'}, \qquad \xi\chi_{\Lambda'} = \chi_{\Lambda''}\xi\chi_{\Lambda'}.$$

Hence  $P_{\lambda}\xi = (P_{\lambda}\chi_{\Lambda''})\xi\chi_{\Lambda'} = z\chi_{\Lambda''}\xi\chi_{\Lambda'} = z\xi\chi_{\Lambda'}$ . Thus for any  $\xi \in U_q\mathfrak{g}$  and any finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  there exist such  $\tilde{\xi} \in U_q\mathfrak{g}$  and a finite subset  $\tilde{\Lambda} \subset P_+^{\mathbb{L}}$  that  $\chi_{\Lambda}\xi = \tilde{\xi}\chi_{\tilde{\Lambda}}$ . This can be readily used to prove that the linear map (2.2) is onto.

Prove that it is injective. One can use well-known results on bases in quantum universal enveloping algebras [6, Chapter 8] to derive a decomposition  $U_q \mathfrak{g} = U_q \mathfrak{q}_{\mathbb{L}}^- \otimes_{U_q \mathfrak{l}} U_q \mathfrak{q}_{\mathbb{L}}^+$ . It follows from this decomposition and Lemma A.1 that  $U_q \mathfrak{g}$  is a free right  $U_q \mathfrak{l}$ -module. Choose a free basis  $\{e_i\}$ . Let *a* be a non-zero element of  $U_q \mathfrak{g} \otimes_{U_q \mathfrak{l}} R(\mathfrak{l})_q$ . It has the form

$$a = \sum_{i} e_i \otimes r_i, \qquad r_i \in R(\mathfrak{l})_q$$

Let  $\tilde{a}$  be the image of a under the map (2.2). We treat it as an element of End  $V^{\text{univ}}$ . It suffices to prove that the restriction of the linear map  $\tilde{a}$  to the subspace  $\{1 \otimes v | v \in L^{\text{univ}}\}$ is non-zero. It is easy to verify that  $\tilde{a}(1 \otimes v) = \sum_{i} e_i \otimes r_i v$  for any  $v \in L^{\text{univ}}$ . It remains to use our choice of  $\{e_i\}$  and the fact that the representation of  $U_q \mathfrak{l}$  in  $L^{\text{univ}}$  is faithful.  $\Box$ 

### 3 The functors ind and $\Pi$

Consider two pairs of subsets

$$\mathbb{L} \subset \mathbb{G} \subset \{1, 2, \dots, l\}, \qquad \mathbb{L}_1 \subset \mathbb{G}_1 \subset \{1, 2, \dots, l\},\$$

with  $\mathbb{L}_1 \subset \mathbb{L}$ ,  $\mathbb{G}_1 \subset \mathbb{G}$ . Obviously, one has embeddings of the associated Hopf subalgebras

$$U_q\mathfrak{l}_1 \subset U_q\mathfrak{l}, \qquad U_q\mathfrak{g}_1 \subset U_q\mathfrak{g}.$$

 $R(\mathfrak{g},\mathfrak{l})_q$  is a left ideal in  $F(\mathfrak{g},\mathfrak{l})_q$  and hence is a left  $U_q\mathfrak{g}_1$ -module. We claim that  $R(\mathfrak{g},\mathfrak{l})_q$ is an approximately unital left  $R(\mathfrak{g}_1,\mathfrak{l}_1)_q$ -module. In fact,  $R(\mathfrak{g},\mathfrak{l})_q$  is an approximately unital left  $R(\mathfrak{g},\mathfrak{l})_q$ -module, hence a  $U_q\mathfrak{g}$ -module of the category  $C(\mathfrak{g},\mathfrak{l})_q$  given by (1.1). Since  $U_q\mathfrak{l}_1 \subset U_q\mathfrak{l}$ , the module  $R(\mathfrak{g},\mathfrak{l})_q$  is also in the category  $C(\mathfrak{g},\mathfrak{l}_1)_q$ . Finally, in view of  $U_q\mathfrak{g}_1 \subset U_q\mathfrak{g}$ , we are inside the category  $C(\mathfrak{g}_1,\mathfrak{l}_1)_q$ , which is equivalent to our claim. In a similar way, one can prove that  $R(\mathfrak{g},\mathfrak{l})_q$  is a right  $R(\mathfrak{g}_1,\mathfrak{l}_1)_q$ -module.<sup>6</sup>

Throughout the rest of this section we follow [7] and replace the groups involved therein with the corresponding quantum universal enveloping algebras. Introduce the functor  $P_{\mathfrak{g}_1,\mathfrak{l}_1}^{\mathfrak{g},\mathfrak{l}}$  from the category  $C(\mathfrak{g}_1,\mathfrak{l}_1)_q$  to the category  $C(\mathfrak{g},\mathfrak{l})_q$  by defining it on objects from  $C(\mathfrak{g}_1,\mathfrak{l}_1)_q$  as follows:

$$P_{\mathfrak{g}_1,\mathfrak{l}_1}^{\mathfrak{g},\mathfrak{l}}(Z)=R(\mathfrak{g},\mathfrak{l})_q\otimes_{R(\mathfrak{g}_1,\mathfrak{l}_1)_q}Z.$$

The action on morphisms is defined in an obvious way.

 $C(\mathfrak{g}_1,\mathfrak{l}_1)_q$  has enough projectives and the functor  $P_{\mathfrak{g}_1,\mathfrak{l}_1}^{\mathfrak{g},\mathfrak{l}}$  is covariant and right exact.<sup>7</sup> Hence one has well defined derived functors:  $\left(P_{\mathfrak{g}_1,\mathfrak{l}_1}^{\mathfrak{g},\mathfrak{l}}\right)_j, j \in \mathbb{Z}_+$ , from  $C(\mathfrak{g}_1,\mathfrak{l}_1)_q$  to  $C(\mathfrak{g},\mathfrak{l})_q$ .

Consider the two special cases:  $\mathfrak{l}_1 = \mathfrak{l}$  and  $\mathfrak{g}_1 = \mathfrak{g}$ . Start from the first one. Let  $\operatorname{ind}_{\mathfrak{g}_1,\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}$  be the functor from the category  $C(\mathfrak{g}_1,\mathfrak{l})_q$  to the category  $C(\mathfrak{g},\mathfrak{l})_q$  defined on objects as follows:

$$\operatorname{ind}_{\mathfrak{g}_1,\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}(Z) = U_q \mathfrak{g} \otimes_{U_q \mathfrak{g}_1} Z.$$

The action on morphisms is defined in an obvious way. Just as in the classical case q = 1, one gets an isomorphism of functors  $P_{\mathfrak{g}_1,\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}$  and  $\operatorname{ind}_{\mathfrak{g}_1,\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}$ . Describe briefly this construction. The functor  $\operatorname{ind}_{\mathfrak{g}_1,\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}$  is left adjoint to the forgetful functor  $\mathcal{F}_{\mathfrak{g},\mathfrak{l}}^{\mathfrak{g}_1,\mathfrak{l}} : C(\mathfrak{g},\mathfrak{l})_q \to C(\mathfrak{g}_1,\mathfrak{l})_q$ determined by the embedding  $U_q\mathfrak{g}_1 \to U_q\mathfrak{g}$ . On the other hand,  $P_{\mathfrak{g}_1,\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}$  is left adjoint to the functor  $(\mathcal{F}^{\vee})_{\mathfrak{g},\mathfrak{l}}^{\mathfrak{g}_1,\mathfrak{l}}$ , to be defined below (cf. [7, Proposition 2.34]). The functor  $(\mathcal{F}^{\vee})_{\mathfrak{g},\mathfrak{l}}^{\mathfrak{g}_1,\mathfrak{l}} : C(\mathfrak{g},\mathfrak{l})_q \to C(\mathfrak{g}_1,\mathfrak{l})_q$  is defined on objects of the category  $C(\mathfrak{g},\mathfrak{l})_q$  as follows:

$$(\mathfrak{F}^{\vee})^{\mathfrak{g}_1,\mathfrak{l}}_{\mathfrak{g},\mathfrak{l}}(X) = \operatorname{Hom}_{R(\mathfrak{g},\mathfrak{l})_q}(R(\mathfrak{g},\mathfrak{l})_q,X)_{\mathfrak{l}}.$$

A structure of  $U_q \mathfrak{g}_1$ -module in  $\operatorname{Hom}_{R(\mathfrak{g},\mathfrak{l})_q}(R(\mathfrak{g},\mathfrak{l})_q,X)$  is imposed via the structure of right  $U_q \mathfrak{g}_1$ -module in  $R(\mathfrak{g},\mathfrak{l})_q$ , and the subscript  $\mathfrak{l}$  stands for distinguishing the maximal submodule in the category  $C(\mathfrak{g}_1,\mathfrak{l})_q$ . The action of  $(\mathcal{F}^{\vee})^{\mathfrak{g}_1,\mathfrak{l}}_{\mathfrak{g},\mathfrak{l}}$  on morphisms is defined in an obvious way. What remains is to construct an isomorphism of functors  $\mathcal{F}^{\mathfrak{g}_1,\mathfrak{l}}_{\mathfrak{g},\mathfrak{l}} \xrightarrow{\sim} (\mathcal{F}^{\vee})^{\mathfrak{g}_1,\mathfrak{l}}_{\mathfrak{g},\mathfrak{l}}$ (cf. [7, Proposition 2.33]). Let  $X \in C(\mathfrak{g},\mathfrak{l})_q$  and  $x \in X$ . Associate to every x a morphism of  $R(\mathfrak{g},\mathfrak{l})_q$ -modules given by

$$R(\mathfrak{g},\mathfrak{l})_q \to X, \qquad r \mapsto rx.$$

One can verify that the map  $X \to \operatorname{Hom}_{R(\mathfrak{g},\mathfrak{l})_q}(R(\mathfrak{g},\mathfrak{l})_q,X)$ , which arises this way, is a morphism of  $U_q\mathfrak{g}_1$ -modules and provides the desired isomorphism of functors.

An important consequence is the observation that for any  $V \in C(\mathfrak{l}, \mathfrak{l})_q$  the standard projective object P(V) in the category  $C(\mathfrak{g}, \mathfrak{l})_q$  is canonically isomorphic to  $P_{\mathfrak{l}\mathfrak{l}}^{\mathfrak{g},\mathfrak{l}}(V)$ .

In the second special case  $\mathfrak{g}_1 = \mathfrak{g}$  the functor in question is called the Bernstein functor and it is denoted by  $\Pi_{\mathfrak{g},\mathfrak{l}_1}^{\mathfrak{g},\mathfrak{l}}$ :

$$\Pi \equiv \Pi_{\mathfrak{g},\mathfrak{l}_1}^{\mathfrak{g},\mathfrak{l}}(Z) = R(\mathfrak{g},\mathfrak{l})_q \otimes_{R(\mathfrak{g},\mathfrak{l}_1)_q} Z, \qquad Z \in C(\mathfrak{g}_1,\mathfrak{l}_1)_q.$$

<sup>&</sup>lt;sup>6</sup>One can see from the proof that  $R(\mathfrak{g},\mathfrak{l})_q$  is a  $R(\mathfrak{g}_1,\mathfrak{l}_1)_q$ -bimodule. <sup>7</sup>See [7, p. 840]

In the classical case q = 1 the derived functors  $\Pi_j$  are crucial in constructing unitarizable Harish-Chandra modules via the Vogan-Zuckerman cohomological induction [7]. Turn to describing a quantum analogue for this method.

### 4 A quantum analogue for cohomological induction

We assume in the sequel that  $\mathbb{G} = \{1, 2, \dots, l\}$ , and hence  $U_q \mathfrak{g} = U_q$ .

Recall (see [4, 8]) a definition of the pairs  $(\mathfrak{g}, \mathfrak{k})$  used in a construction of bounded symmetric domains via the Harish-Chandra embedding. Here  $\mathfrak{g}$  is a simple complex Lie algebra of type  $A, B, C, D, E_6, E_7$ . A Hopf subalgebra  $U_q \mathfrak{k} \subset U_q \mathfrak{g}$  is generated by  $E_i, F_i$ ,  $i \in \mathbb{K}; K_j^{\pm 1}, j = 1, 2, \ldots, l$ , with  $\mathbb{K} = \{1, 2, \ldots, l\} \setminus \{l_0\}$ . We assume also that the simple root  $\alpha_{l_0}$  has coefficient 1 in the decomposition of the maximal root

$$\delta = \sum_{i=1}^{l} n_i \alpha_i, \qquad n_{l_0} = 1.$$
(4.1)

In what follows the notation  $(\mathfrak{g}, \mathfrak{k})$  will stand only for such pairs.

To clarify the precise nature of condition (4.1), equip the Lie algebra  $\mathfrak{g}$  with a grading as follows: deg  $E_{l_0} = 1$ , deg  $F_{l_0} = -1$ , deg  $H_{l_0} = 0$ , deg  $E_j = \deg F_j = \deg H_j = 0$  for  $j \neq l_0$ . The automorphism  $\theta$  of  $\mathfrak{g}$  given by the formula

$$\theta(\xi) = (-1)^{\deg \xi} \xi$$

is an involution. Now (4.1) implies that

$$\mathfrak{g} = \mathfrak{p}^- \oplus \mathfrak{k} \oplus \mathfrak{p}^+, \qquad \mathfrak{p}^\pm = \{\xi \in \mathfrak{g} | \deg \xi = \pm 1\}.$$

It is worthwhile to note that the pairs  $(\mathfrak{g}, \mathfrak{k})$  in question are complexifications of the pairs  $(\mathfrak{g}_0, \mathfrak{k}_0)$ , with  $\mathfrak{g}_0$  being the Lie subalgebra of the automorphism group of an irreducible bounded symmetric domain, and  $\mathfrak{k}_0$  being the Lie algebra of the stabilizer of a point in this domain [4].

Equip  $U_q \mathfrak{g}$  with a structure of Hopf \*-algebra via the involution \* defined as follows:

$$E_j^* = \begin{cases} -K_j F_j, & j = l_0 \\ K_j F_j, & j \neq l_0 \end{cases}; \qquad F_j^* = \begin{cases} -E_j K_j^{-1}, & j = l_0 \\ E_j K_j^{-1}, & j \neq l_0 \end{cases};$$
$$(K_j^{\pm 1})^* = K_j^{\pm 1}, \qquad j = 1, 2, \dots, l.$$

A module V over a Hopf \*-algebra A is said to be unitarizable if it admits a positive definite invariant form  $(\cdot, \cdot)$ :

$$(av_1, v_2) = (v_1, a^*v_2), \qquad a \in A, \ v_1, v_2 \in V.$$

The cohomological induction is among the tools for constructing unitarizable modules of the category  $C(\mathfrak{g}, \mathfrak{k})$  in the case q = 1. Describe a q-analogue for this method.

Let  $\mathbb{L} \subset \{1, 2, \ldots, l\}$ ,  $\mathbb{L} \not\subset \mathbb{K}$ , and  $U_q \mathfrak{l}$  is the Hopf subalgebra corresponding to the subset  $\mathbb{L}$ . Obviously,  $U_q \mathfrak{l}$  inherits the structure of Hopf \*-algebra. We intend to derive a unitarizable  $U_q \mathfrak{g}$ -module of the category  $C(\mathfrak{g}, \mathfrak{k})_q$  starting from a unitarizable  $U_q \mathfrak{l}$ -module of the category  $C(\mathfrak{g}, \mathfrak{k})_q$  starting from a unitarizable  $U_q \mathfrak{l}$ -module of the category  $C(\mathfrak{g}, \mathfrak{k})_q$  starting from a unitarizable  $U_q \mathfrak{l}$ -module of the category  $C(\mathfrak{g}, \mathfrak{k})_q$  starting from a unitarizable  $U_q \mathfrak{l}$ -module of the category  $C(\mathfrak{g}, \mathfrak{k})_q$ .

 $<sup>{}^{8}</sup>U_{q}(\mathfrak{l} \cap \mathfrak{k})$  is a Hopf subalgebra corresponding to the subset  $\mathbb{L} \cap \mathbb{K}$ .

If  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l) \in P$  and  $\lambda_j = 0$  for all  $j \in \mathbb{L}$ , one has the following well defined one-dimensional  $U_q$ I-module  $\mathbb{C}_{\lambda}$ :

$$E_j \mathbf{1} = F_j \mathbf{1} = 0, \quad j \in \mathbb{L}; \qquad K_j^{\pm 1} \mathbf{1} = q_j^{\pm \lambda_j} \mathbf{1}, \quad j = 1, 2, \dots, l.$$

As an important example of such linear functional one should mention the difference  $\rho_{\mathfrak{u}} = \rho - \rho_{\mathfrak{l}}$  between the half-sum  $\rho$  of positive roots of the Lie algebra  $\mathfrak{g}$  and the half-sum  $\rho_{\mathfrak{l}}$  of positive roots of the Lie algebra  $\mathfrak{l}$ :

$$\rho_{\mathfrak{u}}(H_j) = \begin{cases} 1, & j \notin \mathbb{L}, \\ 0, & j \in \mathbb{L}. \end{cases}$$

Every  $U_q\mathfrak{l}$ -module  $Z \in C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$  determines a  $U_q\mathfrak{l}$ -module  $Z^{\#} = Z \otimes \mathbb{C}_{2\rho_{\mathfrak{u}}} \in C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$ . Equip it with a structure of  $U_q\mathfrak{q}_{\mathbb{L}}^-$ -module via the surjective morphism of algebras  $U_q\mathfrak{q}_{\mathbb{L}}^- \to U_q\mathfrak{l}$ :

$$F_i \mapsto \begin{cases} F_i, & i \in \mathbb{L} \\ 0, & i \in \{1, 2, \dots, l\} \setminus \mathbb{L} \end{cases}, \quad E_j \mapsto E_j, \quad K_j^{\pm 1} \mapsto K_j^{\pm 1}, \ j \in \{1, 2, \dots, l\}.$$

The generalized Verma module  $\operatorname{ind}_{\mathfrak{q}_{\mathbb{L}}^{-}}^{\mathfrak{g}} Z^{\#} \stackrel{\text{def}}{=} U_q \mathfrak{g} \otimes_{U_q \mathfrak{q}_{\mathbb{L}}^{-}} Z^{\#}$  belongs to the category  $C(\mathfrak{g}, \mathfrak{l})_q$  (see Lemma A.1). Hence the equality

$$\mathcal{L}_{j}(Z) = \left(\Pi_{\mathfrak{g},\mathfrak{l}\cap\mathfrak{k}}^{\mathfrak{g},\mathfrak{k}}\right)_{j} \left( \operatorname{ind}_{\mathfrak{q}_{\mathbb{L}}^{-}}^{\mathfrak{g}}(Z^{\#}) \right)$$

determines functors  $\mathcal{L}_j, j \in \mathbb{Z}_+$ , from  $C(\mathfrak{l}, \mathfrak{l} \cap \mathfrak{k})_q$  to  $C(\mathfrak{g}, \mathfrak{k})_q$ .

In the classical case q = 1 under suitable dominance assumptions on Z the only nonzero modules  $\mathcal{L}_s(Z)$  are those with  $s = \frac{1}{2}(\dim \mathfrak{k} - \dim(\mathfrak{k} \cap \mathfrak{l}))$  [7, p. 369]. So, a particular interest is in considering the  $U_q\mathfrak{g}$ -modules

$$A_{\mathfrak{q}}(\lambda) \stackrel{\text{def}}{=} \mathcal{L}_{s}(\mathbb{C}_{\lambda}), \qquad s = \frac{1}{2} (\dim \mathfrak{k} - \dim(\mathfrak{k} \cap \mathfrak{l})),$$

with  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l) \in P$  and  $\lambda_j = 0$  for  $j \in \mathbb{L}$ .

### Appendix. Proofs of the Lemmas

Let

Let  $\mathbb{G} \supset \mathbb{L}$  be subsets of  $\{1, 2, \ldots, l\}$ . Equip  $U_q \mathfrak{g}$  with a grading by setting

$$\deg E_j = \begin{cases} 1, \quad j \in \mathbb{G} \setminus \mathbb{L} \\ 0, \quad j \in \mathbb{L} \end{cases}, \qquad \deg F_j = \begin{cases} -1, \quad j \in \mathbb{G} \setminus \mathbb{L} \\ 0, \quad j \in \mathbb{L} \end{cases}, \\ \deg K_j^{\pm 1} = 0, \qquad j = 1, 2, \dots, l. \end{cases}$$
$$(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j = \{\xi \in U_q \mathfrak{q}_{\mathbb{L}}^{\pm} | \deg \xi = j\}. \text{ Obviously, } (U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_0 = U_q \mathfrak{l}.\end{cases}$$

**Lemma A.1** The homogeneous components  $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j$  are free left and free right  $U_q \mathfrak{l}$ -modules of finite rank.

**Proof.** Recall that the subsets  $\mathbb{L} \subset \mathbb{G} \subset \{1, 2, ..., l\}$  determine the subgroups  $W_{\mathbb{L}} \subset W_{\mathbb{G}}$  of the Weyl group W. Those subgroups are exactly the Weyl groups of the Lie algebras  $\mathfrak{l}$  and  $\mathfrak{g}$ , respectively. Let  $w_{0,\mathbb{L}}$  (respectively,  $w_{0,\mathbb{G}}$ ) be the longest element of the Weyl group  $W_{\mathbb{L}}$  (respectively,  $W_{\mathbb{G}}$ ).

We start from the larger subset  $\mathbb{G}$ . Choose a reduced expression

$$w_{0,\mathbb{G}} = s_{i_1} s_{i_2} s_{i_3} \dots s_{i_M},$$
 (A.2)

with  $s_{i_k}$  being the reflection corresponding to the simple root  $\alpha_{i_k}$ . Consider the Lusztig automorphisms  $T_i, i \in \mathbb{G}$ , of the algebra  $U_q \mathfrak{g}'$  generated by  $E_i, F_i, K_i^{\pm 1}, i \in \mathbb{G}$ , [6, Chapter 8]. They have the form

$$T_i(K_j) = K_j K_i^{-a_{ij}},$$

$$T_i(E_i) = -F_i K_i, \quad T_i(F_i) = -K_i^{-1} E_i,$$

$$T_i(E_j) = \sum_{r+s=-a_{ij}} \operatorname{const} E_i^s E_j E_i^r, \quad i \neq j,$$

$$T_i(F_j) = \sum_{r+s=-a_{ij}} \operatorname{const} F_i^r F_j F_i^s, \quad i \neq j.$$
(A.3)

In a similar way, consider the algebra  $U_q l'$  associated to  $\mathbb{L}$ . Let  $W^{\mathbb{L}} = \{w \in W_{\mathbb{G}} | l(ws) > l(w)$  for all simple reflections  $s \in W_{\mathbb{L}}\}$ , where l(w) stands for the length of w. One has  $w_{0,\mathbb{G}} = w^{\mathbb{L}} w_{0,\mathbb{L}}$  with  $w^{\mathbb{L}} \in W^{\mathbb{L}}$  and  $l(w_{0,\mathbb{G}}) = l(w^{\mathbb{L}}) + l(w_{0,\mathbb{L}})$  [5, Proposition 1.10(c)]. This allows one to rewrite the reduced expression (A.2) in the form

$$w_{0,\mathbb{G}} = s_{i_1} s_{i_2} s_{i_3} \dots s_{i_{M'}} s_{i_{M'+1}} s_{i_{M'+2}} \dots s_{i_M}, \tag{A.4}$$

with  $w_{0,\mathbb{L}} = s_{i_{M'+1}}, s_{i_{M'+2}}, \dots, s_{i_M}$  and  $w^{\mathbb{L}} = s_{i_1} s_{i_2} s_{i_3} \dots s_{i_{M'}}$ .

We are about to apply the Lusztig theorem [6, Theorem 8.24]. Observe that the special reduced expression (A.4), in view of the explicit form of the Lusztig automorphisms (A.3) implies that all the monomials

$$T_{i_1}T_{i_2}\cdots T_{i_{M'-1}}(E_{i_{M'}}^{a_{M'}})\cdot T_{i_1}T_{i_2}\cdots T_{i_{M'-2}}(E_{i_{M'-1}}^{a_{M'-1}})\cdot \ldots \cdot T_{i_1}T_{i_2}(E_{i_3}^{a_3})\cdot T_{i_1}(E_{i_2}^{a_2})\cdot E_{i_1}^{a_1}$$

are in  $U_q l'$ . It follows from the Lusztig theorem [6, Theorem 8.24] that the monomials

$$T_{i_1}T_{i_2}\cdots T_{i_{M-1}}(E_{i_M}^{a_M})\cdot T_{i_1}T_{i_2}\cdots T_{i_{M-2}}(E_{i_{M-1}}^{a_{M-1}})\cdot \dots$$
$$\dots \cdot T_{i_1}T_{i_2}\cdots T_{i_{M'+1}}(E_{i_{M'+2}}^{a_{M'+2}})\cdot T_{i_1}T_{i_2}\cdots T_{i_{M'}}(E_{i_{M'+1}}^{a_{M'+1}}) \quad (A.5)$$

with all  $a_{i_k} \in \mathbb{Z}_+$ , form a free basis in the right  $U_q\mathfrak{l}$ -module  $U_q\mathfrak{q}_{\mathbb{L}}^+$ . As one can readily separate out for each  $j \in \mathbb{Z}_+$  finitely many such monomials that span the *j*-th homogeneous component, we thus get our claim for  $(U_q\mathfrak{q}_{\mathbb{L}}^+)_j$  as a right  $U_q\mathfrak{l}$ -module. All other claims can be proved in a similar way.

**Corollary A.2** Every  $U_q\mathfrak{g}$ -module V contains the largest submodule  $V_{\mathfrak{l}}$  of the category  $C(\mathfrak{g},\mathfrak{l})_q$ .

**Proof.** Obviously, V possesses the largest weight submodule  $V_{\mathfrak{h}}$ . What remains is to prove that the subspace  $V_{\mathfrak{l}} = \{v \in V_{\mathfrak{h}} | \dim(U_q \mathfrak{l} v) < \infty\}$  is a submodule of the  $U_q \mathfrak{g}$ -module V. Let  $\xi \in (U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_i, v \in V_{\mathfrak{l}}$ . It follows from Lemma A.1 that for some  $\{\eta_1, \eta_2, \ldots, \eta_{N(j)}\}$ 

$$\left(U_q \mathfrak{q}_{\mathbb{L}}^{\pm}\right)_j = \sum_{k=1}^{N(j)} \eta_k U_q \mathfrak{l}$$

Hence,

$$\dim(U_q\mathfrak{l}\,\xi v) \leq \dim\left(\left(U_q\mathfrak{q}_{\mathbb{L}}^{\pm}\right)_j v\right) = \dim\left(\sum_{k=1}^{N(j)} \eta_k U_q\mathfrak{l}\,v\right) < \infty$$

Thus  $\xi v \in V_{\mathfrak{l}}$ .

**Corollary A.3** For any  $V \in C(\mathfrak{l}, \mathfrak{l})_q$ , the module P(V) belongs to the category  $C(\mathfrak{g}, \mathfrak{l})_q$ .

**Proof.** Since  $U_q \mathfrak{g}$  is a free right  $U_q \mathfrak{l}$ -module, one has an embedding of  $U_q \mathfrak{l}$ -modules

$$i: V \hookrightarrow P(V), \qquad i: v \mapsto 1 \otimes v.$$

The relation  $P(V) = P(V)_{\mathfrak{l}}$  is due to

$$P(V) = U_q \mathfrak{g} \, i(V) \subset U_q \mathfrak{g} \, P(V)_{\mathfrak{l}} = P(V)_{\mathfrak{l}}. \qquad \Box$$

The rest of the statements of Lemma 2.1 can be proved in the same way as in the classical case q = 1.

To prove Lemma 2.5 we need the following auxiliary statement.

**Lemma A.4** Let  $\lambda \in P_+^{\mathbb{L}}$  and  $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j$  be the homogeneous components of the graded algebras  $U_q \mathfrak{q}_{\mathbb{L}}^{\pm}$ .

- 1. The vector spaces  $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j \cdot \mathfrak{P}_{\lambda} \subset \operatorname{End} V^{\operatorname{univ}}$  are finite dimensional.
- 2. The vector spaces  $\mathfrak{P}_{\lambda} \cdot (U_{q}\mathfrak{q}_{\mathbb{L}}^{\pm})_{i} \subset \operatorname{End} V^{\operatorname{univ}}$  are finite dimensional.<sup>9</sup>

**Proof.** Prove the first statement. We consider End  $V^{\text{univ}}$  as a  $U_q \mathfrak{l}$ -module with respect to the action as follows:

$$(\xi a): v \mapsto \xi(av), \qquad v \in V^{\text{univ}}, \quad a \in \text{End}\, V^{\text{univ}}, \quad \xi \in U_q\mathfrak{l}.$$

Obviously,  $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_i \cdot \mathfrak{P}_{\lambda}$  is a submodule of the  $U_q \mathfrak{l}$ -module End  $V^{\text{univ}}$ .

Let  $\pi_{\lambda} : U_q \mathfrak{l} \to \text{End } L(\mathfrak{l}, \lambda)$  be the representation of the Hopf subalgebra  $U_q \mathfrak{l}$  corresponding to the  $U_q \mathfrak{l}$ -module  $L(\mathfrak{l}, \lambda)$ . If  $\xi \in \text{Ker } \pi_{\lambda}$ , then  $\xi \cdot \mathfrak{P}_{\lambda} = 0$ . Hence the diagram



 $<sup>^{9}</sup>$ A dot is used here to denote the product of elements in End  $V^{\text{univ}}$ .

can be completed up to a commutative one, and

$$\dim(U_q\mathfrak{l}\cdot\mathfrak{P}_{\lambda}) \leq (\dim(L(\mathfrak{l},\lambda)))^2.$$

Thus the first statement of the Lemma is proved in the case j = 0. It remains to elaborate the fact that  $(U_q \mathbf{q}_{\mathbb{L}}^{\pm})_j$  is a right  $U_q \mathbf{l}$ -module of finite rank (see Lemma A.1).

Now the first statement is proved. The second one can be proved in a similar way using the commutative diagram



in the category of  $U_q l^{\text{op}}$ -modules.<sup>10</sup>

Turn to the proof of Lemma 2.5. Consider the subspace  $A \subset U_q \mathfrak{g}$  of such elements  $\xi \in U_q \mathfrak{g}$  that for any  $\lambda \in P_+^{\mathbb{L}}$  there exists a finite subset  $\Lambda \subset P_+^{\mathbb{L}}$  with

$$\xi \cdot \mathcal{P}_{\lambda} = \chi_{\Lambda} \cdot \xi \cdot \mathcal{P}_{\lambda}, \tag{A.6}$$

$$\mathcal{P}_{\lambda} \cdot \xi = \mathcal{P}_{\lambda} \cdot \xi \cdot \chi_{\Lambda}. \tag{A.7}$$

One has to prove that  $A = U_q \mathfrak{g}$ . For that, it suffices to demonstrate that A is a subalgebra of  $U_q \mathfrak{g}$  and  $A \supset (U_q \mathfrak{g}_{\mathbb{L}}^{\pm})_i$  for all j.

Prove the first statement. It is obvious that A is a vector subspace. Now let  $\xi, \eta \in A$ , and consider the product  $\xi\eta$ . A double application of (A.6) first w.r.t.  $\eta$  and then  $\xi$  allows one to deduce that the image of the linear operator  $\xi \cdot \eta \cdot \mathcal{P}_{\lambda} \in \text{End } V^{\text{univ}}$  is accommodated by the sum of finitely many isotypic components  $V_{\lambda}^{\text{univ}}$ . Let this sum be  $\bigoplus_{\lambda' \in \Lambda'} V_{\lambda'}^{\text{univ}}$ , then

 $\xi \cdot \eta \cdot \mathfrak{P}_{\lambda} = \chi_{\Lambda'} \cdot \xi \cdot \eta \cdot \mathfrak{P}_{\lambda}$ . Thus we get (A.6) for  $\xi \eta$ .

In a similar way, apply (A.7) twice to deduce that for some finite subset  $\Lambda'' \subset P_+^{\mathbb{L}}$  one has  $\operatorname{Ker}(\mathcal{P}_{\lambda} \cdot \xi \cdot \eta) \supset (\operatorname{id} - \chi_{\Lambda''}) V^{\operatorname{univ}}$ , hence  $\mathcal{P}_{\lambda} \cdot \xi \cdot \eta = \mathcal{P}_{\lambda} \cdot \xi \cdot \eta \cdot \chi_{\Lambda''}$ . Thus (A.7) holds for  $\xi\eta$ .

Turn to the second statement. We restrict ourselves to proving that (A.6) is valid for all  $\xi \in (U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j$ . It follows from Lemma A.4 that for every j,  $\lambda$ , the vector space  $(U_q \mathfrak{q}_{\mathbb{L}}^{\pm})_j \cdot \mathcal{P}_{\lambda}$  is finite dimensional. It is also a  $U_q\mathfrak{l}$ -module since  $U_q\mathfrak{l}(U_q\mathfrak{q}_{\mathbb{L}}^{\pm})_j \subset (U_q\mathfrak{q}_{\mathbb{L}}^{\pm})_j$ . Hence  $(U_q\mathfrak{q}_{\mathbb{L}}^{\pm})_j \cdot \mathcal{P}_{\lambda}$  is a sum of finitely many  $U_q\mathfrak{l}$ -isotypic components. On the other hand, if for some  $a \in \operatorname{End} V^{\operatorname{univ}}$ , the  $U_q\mathfrak{l}$ -module  $U_q\mathfrak{l} \cdot a \subset \operatorname{End} V^{\operatorname{univ}}$  is a multiple of  $L(\mathfrak{l}, \mu)$ ,  $\mu \in P_+^{\mathbb{L}}$ , one has  $aV^{\operatorname{univ}} \subset V_{\mu}^{\operatorname{univ}}$  and  $\mathcal{P}_{\mu} \cdot a = a$ . It follows that  $\mathcal{P}_{\lambda'} \cdot (U_q\mathfrak{q}_{\mathbb{L}}^{\pm})_j \cdot \mathcal{P}_{\lambda} = 0$  for all but finitely many  $\lambda' \in P_+^{\mathbb{L}}$ . Hence  $\xi \cdot \mathcal{P}_{\lambda} = \chi_{\Lambda} \cdot \xi \cdot \mathcal{P}_{\lambda}$  for some finite subset  $\Lambda \subset P_+^{\mathbb{L}}$ .

### References

- [1] R. J. Baston, M. G. Eastwood. The Penrose Transform. Its Interaction with Representation Theory. Clarendon Press, Oxford, 1989.
- [2] N. Bourbaki, Groupes et Algèbres de Lie, chapitres VII-VIII. Herman, 1975.

<sup>&</sup>lt;sup>10</sup>The algebra  $U_q l^{\text{op}}$  is derived from  $U_q l$  by replacing the multiplication with the opposite one.

- [3] C. De Concini, V. Kac. Representations of quantum groups at roots of 1. In: Operator Algebras, Unitary Representations, Enveloping Algebras and Invariant Theory, A. Connes, M. Duflo, R. Rentchler (eds.), Birkhäuser, Boston-Basel-Berlin, 1990, 472 – 506.
- [4] S. Helgason. Differential Geometry and Symmetric Spaces. Academic Press, New York, London, 1962.
- [5] J. E. Humphreys. Reflection Groups and Coxeter Groups. Cambridge Univ. Press, Cambridge, 1990.
- [6] J. C. Jantzen. Lectures on Quantum Groups. American Mathematical Society, Providence, R. I., 1996.
- [7] A. Knapp, D. A. Vogan, Jr. Cohomological Induction and Unitary Representations. Princeton Univ. Press, Princeton, 1995.
- [8] A. Korányi, Function spaces on bounded symmetric domains, In: Analysis and geometry on complex homogeneous domains. Birkhäuser Boston, Inc., Boston, MA, 2000, J. Faraut, S. Kaneyuki, A. Koranyi, Qi-keng Lu, Guy Roos, (eds), 183 – 281.
- [9] D. Shklyarov, S. Sinel'shchikov, A. Stolin, and L. Vaksman Non-compact quantum groups and quantum Harish-Chandra modules. In: Supersymmetry and Quantum Field Theory, North-Holland, D. Sorokin (ed.), the Netherlands, ser. Nucl. Phys. B (Proc. Suppl.), 102 & 103 (2001), 334 – 337.
- [10] D. Shklyarov, S. Sinel'shchikov, A. Stolin, and L. Vaksman, On a q-analogue of the Penrose transform, Ukrainian Phys. J. 47 (2002), No 3, 288 – 292.
- [11] S. Sinel'shchikov, L. Vaksman, On q-analogues of bounded symmetric domains and Dolbeault complexes, Math. Phys. Anal. Geom. 1 (1998), No 1, 75 – 100.