A NEW LIE BIALGEBRA STRUCTURE ON sl(2,1)

GIZEM KARAALI

ABSTRACT. We describe the Lie bialgebra structure on the Lie superalgebra sl(2,1) related to an r-matrix that cannot be obtained by a Belavin-Drinfeld type construction. This structure makes sl(2,1) into the Drinfeld double of a four-dimensional subalgebra.

It is well-known that non-degenerate r-matrices (describing quasitriangular Lie bialgebra structures) on simple Lie algebras are classified by Belavin-Drinfeld triples, (the original references are [BD1, BD2], more pedagogical presentations providing ample background can be found in [CP, ES]). A similar construction using Belavin-Drinfeld type triples is possible for simple Lie superalgebras with nondegenerate Killing form. Surprisingly, though, in the super setting, there are certain non-degenerate r-matrices that do not fit such a description, see [K].

The purpose of this note is to study the super Lie bialgebra structure associated to such an r-matrix on the simple Lie superalgebra sl(2, 1). We start in Section 1 with some background on super Lie bialgebra structures and some basic constructions related to them. In Section 2, we explicitly describe the r-matrix that we will be interested in. Section 3 describes in detail the associated super Lie bialgebra structure on sl(2, 1); this structure makes sl(2, 1) into the Drinfeld double of a four-dimensional subalgebra. A comparison with the standard super Lie bialgebra structure is also provided in this section. We end in Section 4 with a brief discussion of the results and further directions for investigation.

Acknowledgments. The author would like to thank N. Reshetikhin, V. Serganova and M. Yakimov for their comments and suggestions.

1. LIE BIALGEBRA STRUCTURES ON LIE SUPERALGEBRAS

1.1. Cohomology of Lie superalgebras. The cohomology theory of Lie superalgebras is more complicated than that of Lie algebras. Even for simple Lie superalgebras and for low dimensions, it is not yet completed. Here we summarize certain basic facts that we will use. For more on the cohomology theory of Lie superalgebras one can look at [F, SZ].

Recall that if \mathfrak{g} is a Lie algebra, then an *n*-cochain taking values in a \mathfrak{g} -module M is an alternating *n*-linear map $f(x_1, x_2, \dots, x_n)$ of *n* variables in \mathfrak{g} . We can view

²⁰⁰⁰ Mathematics Subject Classification. Primary 17B62, 17B20.

 $Key\ words\ and\ phrases.$ Classical r-matrices, Lie superalgebras, super Lie bialgebras, Drinfeld double.

each such n-cochain as a linear map $f : \bigwedge^n \mathfrak{g} \to M$. In this case, the coboundary df of an n-cochain f is the (n + 1)-cochain defined by:

$$df(x_1, \cdots, x_{n+1}) = \sum_{i=1}^{n+1} (-1)^{i+1} x_i f(x_1, \cdots, \hat{x}_i, \cdots, x_{n+1}) \\ + \sum_{1 \le i < j \le n+1} (-1)^{i+j} f([x_i, x_j], x_1, \cdots, \hat{x}_i, \cdots, \hat{x}_j, \cdots, x_{n+1})$$

If \mathfrak{g} is a Lie superalgebra, then the space of n-cochains with values in a \mathfrak{g} -module $M = M_{\overline{0}} \oplus M_{\overline{1}}$ is itself a graded space. Denoting this space by $C^n(\mathfrak{g}, M)$, we have:

$$C^{n}(\mathfrak{g}, M) = \bigoplus_{i+j=n} Hom(\wedge^{i}\mathfrak{g}_{\overline{0}} \otimes S^{j}\mathfrak{g}_{\overline{1}}, M).$$

The even part of $C^n(\mathfrak{g}, M)$ is:

$$C^{n}_{\overline{0}}(\mathfrak{g},M) = \bigoplus_{i+j=n} Hom(\wedge^{i}\mathfrak{g}_{\overline{0}} \otimes S^{j}\mathfrak{g}_{\overline{1}}, M_{\overline{j}}),$$

while the odd part is given by:

$$C^{n}_{\overline{1}}(\mathfrak{g}, M) = \bigoplus_{i+j=n} Hom(\wedge^{i}\mathfrak{g}_{\overline{0}} \otimes S^{j}\mathfrak{g}_{\overline{1}}, M_{\overline{j+1}}).$$

Equivalently we can view $C^n(\mathfrak{g}, M)$, for $n \ge 1$, as the $\mathbb{Z}/2\mathbb{Z}$ -graded vector space of all super alternating *n*-linear maps f of $\mathfrak{g}^n = \mathfrak{g} \times \mathfrak{g} \times \cdots \times \mathfrak{g}$ into M, i.e. maps f satisfying:

$$f(x_1, x_2, \cdots, x_i, x_{i+1}, \cdots, x_n) = -(-1)^{|x_i||x_{i+1}|} f(x_1, x_2, \cdots, x_{i+1}, x_i, \cdots, x_n).$$

We set $C^0(\mathfrak{g}, M) = M$. The differential *d* is defined as follows: For an *n*-cochain *f*, the coboundary *df* is an (n + 1)-cochain given by:

$$df(x_1, \cdots, x_{n+1}) = \sum_{i=1}^{n+1} \sigma_1(i, j) x_i f(x_1, \cdots, \hat{x_i}, \cdots, x_{n+1}) \\ + \sum_{1 \le i < j \le n+1} \sigma_2(i, j) f([x_i, x_j], x_1, \cdots, \hat{x_i}, \cdots, \hat{x_j}, \cdots, x_{n+1})$$

where the signs in the above sums are as follows:

$$\sigma_1(i,j) = (-1)^{i+1} (-1)^{|x_i|(|f|+|x_1|+|x_2|+\dots+|x_{i-1}|)}$$

$$\sigma_2(i,j) = (-1)^{i+j} (-1)^{|x_i||x_j|} (-1)^{|x_i|(|x_1|+\dots+|x_{i-1}|)} (-1)^{|x_j|(|x_1|+\dots+|x_{j-1}|)}$$

We note that this formula for d agrees with that of [SZ] when we use the super alternating property of f.

If $M = \mathfrak{g} \otimes \mathfrak{g}$, then \mathfrak{g} acts on M on the left by the following extension of the adjoint representation:

$$g \cdot (a \otimes b) = (g \cdot a) \otimes b + (-1)^{|g||a|} a \otimes (g \cdot b) = [g, a] \otimes b + (-1)^{|g||a|} a \otimes [g, b]$$

In this setup, a 0-cochain is a linear map $f_0 : \mathbb{C} \to \mathfrak{g} \otimes \mathfrak{g}$. Therefore it is determined uniquely by $f_0(1) \in \mathfrak{g} \otimes \mathfrak{g}$ and hence can be identified with an element r of $\mathfrak{g} \otimes \mathfrak{g}$. The coboundary dr of r is a 1-cochain defined by:

$$dr(a) = a \cdot r = [a \otimes 1 + 1 \otimes a, r].$$

A 1-cochain is a linear map $f : \mathfrak{g} \to \mathfrak{g} \otimes \mathfrak{g}$. It is a 1-cocycle if df = 0, or in other words:

$$\begin{array}{ll} 0 & = & df(a,b) = (-1)^{|a||f|} a \cdot f(b) - (-1)^{|b|(|f|+|a|)} b \cdot f(a) - f([a,b]) \\ & = & (-1)^{|a||f|} [a \otimes 1 + 1 \otimes a, f(b)] - (-1)^{|b|(|f|+|a|)} [b \otimes 1 + 1 \otimes b, f(a)] - f([a,b]) \end{array}$$

which we can rewrite as:

$$\begin{aligned} f([a,b]) &= (-1)^{|a||f|} [a \otimes 1 + 1 \otimes a, f(b)] - (-1)^{|b|(|f|+|a|)} [b \otimes 1 + 1 \otimes b, f(a)] \\ &= [f(a), b \otimes 1 + 1 \otimes b] + (-1)^{|a||f|} [a \otimes 1 + 1 \otimes a, f(b)] \\ &= (-1)^{|a||f|} a \cdot f(b) - (-1)^{|b||f(a)|} b \cdot f(a). \end{aligned}$$

We will call the resulting formula the *super cocycle condition*:

$$f([a,b]) = (-1)^{|a||f|} a \cdot f(b) - (-1)^{|b||f(a)|} b \cdot f(a).$$

1.2. Super Lie bialgebras. A super Lie bialgebra is a triple $(\mathfrak{g}, [\cdot, \cdot], \delta)$ such that:

- (1) \mathfrak{g} is a Lie superalgebra with the super bracket $[\cdot, \cdot]$;
- (2) $\delta: \mathfrak{g} \to \mathfrak{g} \otimes \mathfrak{g}$ is a skew-symmetric linear map whose dual $\delta^*: \mathfrak{g}^* \otimes \mathfrak{g}^* \to \mathfrak{g}^*$ defines a Lie superalgebra structure on \mathfrak{g}^* ;
- (3) δ and $[\cdot, \cdot]$ are compatible in the following sense:

$$\delta([a,b]) = [\delta(a), b \otimes 1 + 1 \otimes b] + [a \otimes 1 + 1 \otimes a, \delta(b)].$$

We will denote such a super Lie bialgebra by (\mathfrak{g}, δ) if the super bracket $[\cdot, \cdot]$ is unambiguous. Note that the last condition is equivalent to δ being a 1-cocycle on \mathfrak{g} with values in $\mathfrak{g} \otimes \mathfrak{g}$, for the cohomology theory of Lie superalgebras as summarized in 1.1. Since δ^* is a super bracket, δ is even, and the super cocycle condition above coincides with the non-graded version.

The Jacobi identity for δ^* is equivalent to the following *coJacobi identity* for δ which holds for any $x \in \mathfrak{g}$:

$$Alt_s(\delta \otimes Id) \cdot \delta(x) = 0.$$

Here $Alt_s : \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \to \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g}$ is defined on homogeneous basis vectors by:

$$Alt_s(a \otimes b \otimes c) = a \otimes b \otimes c + (-1)^{|a|(|b|+|c|)} b \otimes c \otimes a + (-1)^{|c|(|a|+|b|)} c \otimes a \otimes b.$$

A (finite dimensional) super Manin triple is a triple $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ of (finite dimensional) Lie superalgebras such that:

- (1) \mathfrak{g} is equipped with a non-degenerate super-symmetric invariant bilinear form (\cdot, \cdot) ;
- (2) \mathfrak{g}_+ and \mathfrak{g}_- are Lie subsuperalgebras of \mathfrak{g} and $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$ as vector spaces;
- (3) \mathfrak{g}_+ and \mathfrak{g}_- are isotropic with respect to (\cdot, \cdot) .

Since the bilinear form is non-degenerate, \mathfrak{g}_+ and \mathfrak{g}_- are in fact maximal isotropic or *Lagrangian* subsuperalgebras. We note here that in the infinite dimensional case,

GIZEM KARAALI

the definition of a super Manin triple will be the same as above. However, one needs to take into account the topology on vector spaces.

These two notions (i.e. super Lie bialgebras and super Manin triples) are related to one another in a way similar to the Lie algebra case:

Proposition 1. Let $(\mathfrak{p}, [,], \delta)$ be a super Lie bialgebra and set $\mathfrak{g}_+ = \mathfrak{p}$ and $\mathfrak{g}_- = \mathfrak{p}^*$ and define $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$. Then $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ is a super Manin triple. Conversely, any finite dimensional super Manin triple $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ gives rise to a super Lie bialgebra structure on \mathfrak{g}_+ .

Remark 1.1. This is Proposition 1 of [A] where it was proven modulo certain calculations left to the reader.

1.3. The Drinfeld double construction. Another related construction is that of the Drinfeld double. Here we will use a direct analogue of the non-graded version, in the spirit of [GZB]. Before explicitly presenting this approach, we should also mention that other superizations of the double construction exist. See for instance [V] for a more geometrically motivated development of the double.

Let $(\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{g}}, \delta_{\mathfrak{g}})$ be a finite dimensional super Lie bialgebra¹. Then clearly \mathfrak{g}^* is also a super Lie bialgebra with the associated structures defined by:

$$[\cdot,\cdot]_{\mathfrak{g}^*} = (\delta_{\mathfrak{g}})^* \qquad \delta_{\mathfrak{g}^*} = ([\cdot,\cdot]_{\mathfrak{g}})^*.$$

Let us fix a homogeneous basis $\{e_i\}$ for \mathfrak{g} and define the structure constants C_k^{ij} , D_k^{ij} of the relevant structures on \mathfrak{g} as follows:

$$[e_i, e_j]_{\mathfrak{g}} = \sum_k C_{ij}^k e_k \qquad \delta_{\mathfrak{g}}(e_k) = \sum_{i,j} D_k^{ij} e_i \wedge e_j$$

where we use the notation: $a \wedge b = a \otimes b - (-1)^{|a||b|} b \otimes a$ for any two homogeneous elements a, b. From these we can determine the structure constants of \mathfrak{g}^* ; if we let $\{e_i^*\}$ be the homogeneous basis for \mathfrak{g}^* dual to $\{e_i\}$, then we have:

$$[e_i^*, e_j^*]_{\mathfrak{g}^*} = \sum_k C_k^{ij} e_k^* \qquad \delta_{\mathfrak{g}^*}(e_k^*) = \sum_{i,j} D_{ij}^k e_i^* \wedge e_j^*$$

where:

$$C_k^{ij} = \begin{cases} (-1)^{|e_i||e_j|} D_k^{ij} & i \neq j \\ \\ -2D_k^{ii} & i = j \end{cases} \qquad D_{ij}^k = \begin{cases} (-1)^{|e_i||e_j|} C_{ij}^k & i \neq j \\ \\ -2C_{ii}^k & i = j \end{cases}$$

These will follow directly from the definitions of linear duality:

$$\langle [x^*, y^*]_{\mathfrak{g}^*}, z \rangle = \langle x^* \otimes y^*, \delta_{\mathfrak{g}}(z) \rangle \quad \langle \delta_{\mathfrak{g}^*}(z^*), x \otimes y \rangle = \langle z^*, [x, y]_{\mathfrak{g}} \rangle$$

where we assume $x, y, z \in \mathfrak{g}$ and $x^*, y^*, z^* \in \mathfrak{g}^*$ are homogeneous. Clearly $C_{ii}^k = D_k^{ii} = 0$ unless e_i (and hence e_i^*) is odd.

 $^{^{1}}$ We will be assuming finite dimensionality, as this will be sufficient for our purposes. Infinite dimensional analogues will be more technically involved, and since we do not need them here, will not be discussed any further.

In this setup, the opposite super Lie bialgebra structure on \mathfrak{g}^* can be defined as follows ²:

$$[e_i^*, e_j^*]_{(\mathfrak{g}^*)^{op}} = [e_j^*, e_i^*]_{\mathfrak{g}^*} \qquad \delta_{(\mathfrak{g}^*)^{op}}(e_k^*) = \delta_{\mathfrak{g}^*}(e_k^*)$$

or equivalently:

$$[\cdot,\cdot]_{(\mathfrak{g}^*)^{op}} = (-\delta_{\mathfrak{g}})^* \qquad \delta_{(\mathfrak{g}^*)^{op}} = \delta_{\mathfrak{g}^*}$$

Note that we are only taking the opposite in terms of the Lie superalgebra structure.

The Drinfeld double \mathfrak{d} of \mathfrak{g} will be defined as the super Lie bialgebra with the underlying graded vector space identified with $\mathfrak{g} \oplus \mathfrak{g}^* \cong \mathfrak{g} \oplus (\mathfrak{g}^*)^{op}$. In order to define a Lie superalgebra structure on \mathfrak{d} , we first define a non-degenerate inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{d} by asserting super-symmetry:

$$\langle x^*, y \rangle = (-1)^{|x^*||y|} \langle y, x^* \rangle$$

and the isotropy of the subspaces \mathfrak{g} and \mathfrak{g}^* :

$$\langle \mathfrak{g}, \mathfrak{g} \rangle = \langle \mathfrak{g}^*, \mathfrak{g}^* \rangle = 0$$

(This choice of notation is intentional, and is meant to agree with that for the duality). We will require invariance of this form, which in terms of the bracket on \mathfrak{d} translates to:

$$\langle [x^*, y]_{\mathfrak{d}}, z \rangle = \langle x^*, [y, z]_{\mathfrak{d}} \rangle \qquad \langle x^*, [y^*, z]_{\mathfrak{d}} \rangle = \langle [x^*, y^*]_{\mathfrak{d}}, z \rangle$$

Then the condition that $[\cdot, \cdot]_{\mathfrak{d}}$ restricts to $[\cdot, \cdot]_{\mathfrak{g}}$ and $[\cdot, \cdot]_{(\mathfrak{g}^*)^{op}}$, respectively, on \mathfrak{g} and \mathfrak{g}^* yields the following description of $[\cdot, \cdot]_{\mathfrak{d}}$ in terms of the structure constants of \mathfrak{g} and \mathfrak{g}^* :

$$\begin{split} [e_i, e_j]_{\mathfrak{d}} &= \sum_k C_{ij}^k e_k \\ [e_i^*, e_j^*]_{\mathfrak{d}} &= \sum_k C_k^{ji} e_k^* \\ [e_i^*, e_j]_{\mathfrak{d}} &= \sum_k C_j^{ik} e_k + \sum_k C_{jk}^i e_k^* \end{split}$$

The super Lie bialgebra structure on \mathfrak{d} is defined to make the natural injections $\mathfrak{g} \to \mathfrak{d}$ and $(\mathfrak{g}^*)^{op} \to \mathfrak{d}$ embeddings of super Lie bialgebras, and hence is given by:

$$\delta_{\mathfrak{d}} = \delta_{\mathfrak{g}} + \delta_{(\mathfrak{g}^*)^{op}} = \delta_{\mathfrak{g}} + \delta_{\mathfrak{g}^*}$$

With the given structures, it can be shown (see [GZB] for details) that \mathfrak{d} is a quasitriangular super Lie bialgebra, with the *r*-matrix:

$$\sum_{i} (-1)^{|e_i|} e_i^* \otimes e_i$$

Although the superization of the main concepts we are using may seem straightforward, it can be shown that several unexpected situations come up during the process. For instance, see [L] for some interesting examples of Manin triples and a few such unexpected phenomena.

²To compare with [GZB], note that $T_s(a \wedge b) = -(a \wedge b)$. Here T_s is defined on homogeneous elements by $T_s(a \otimes b) = (-1)^{|a||b|} b \otimes a$. In other words, it is the permutation map in the category of super vector spaces.

2. Defining the r-matrix r(f)

From now on, let $\mathfrak{g} = sl(2,1)$. Define:

$$f(E_{11} + E_{33}) = 0 \qquad f(E_{22} + E_{33}) = E_{22} + E_{33}$$

$$f(E_{21}) = 0 \qquad f(E_{12}) = E_{12}$$

$$f(E_{23}) = 0 \qquad f(E_{13}) = E_{13}$$

$$f(E_{31}) = -E_{13} \qquad f(E_{32}) = E_{23} + E_{32}$$

and extend f to a linear map on \mathfrak{g} . It is easy to check that for any $x, y \in \mathfrak{g}$, this function satisfies:

$$(f-1)[f(x), f(y)] = f([(f-1)(x), (f-1)(y)])$$

which is equivalent to the associated 2-tensor being an r-matrix, (see Lemma 1 of [K]).

We write the quadratic Casimir element of \mathfrak{g} or equivalently the invariant tensor in $\mathfrak{g} \otimes \mathfrak{g}$:

$$\Omega = (E_{11} + E_{33}) \otimes (-E_{22} - E_{33}) + (-E_{22} - E_{33}) \otimes (E_{11} + E_{33}) + (E_{12} \otimes E_{21} + E_{21} \otimes E_{12}) + (-E_{13} \otimes E_{31} + E_{31} \otimes E_{13}) + (-E_{23} \otimes E_{32} + E_{32} \otimes E_{23}).$$

Defining r(f) to be the 2-tensor $(f \otimes 1)\Omega$, we get:

$$r(f) = r_0 + E_{12} \otimes E_{21} - E_{13} \otimes E_{31} + E_{32} \otimes E_{23} - E_{13} \otimes E_{13} + E_{23} \otimes E_{23}$$

where $r_0 = (-E_{22} - E_{33}) \otimes (E_{11} + E_{33})$. It is easy to see that r(f) satisfies:

(2.1)
$$r + T_s(r) = \Omega$$

Recall here that T_s is the permutation map in the category of super vector spaces.

r(f) does not allow a straightforward Belavin-Drinfeld type description. In fact we can prove that the two subsuperalgebras Im(f) and Im(f-1) will never be simultaneously isomorphic to root subsuperalgebras. The corresponding subsuperalgebras for functions constructible by Belavin-Drinfeld type data are always root subsuperalgebras.

All r-matrices on a simple Lie algebra satisfying Eqn.(2.1) are constructible by Belavin-Drinfeld type data. Thus, the existence in the super case of an r-matrix satisfying this equation but not allowing a Belavin-Drinfeld type description provides us with yet another example when the graded case is more involved than the non-graded case.

3. The Super Lie Bialgebra Structure Associated to r(f)

In the rest of this note, we will concentrate on the Lie bialgebra structure on \mathfrak{g} associated to $r(f) = (f \otimes 1)\Omega$, where f is the linear map introduced in Section 2 above. After explicitly describing this super Lie bialgebra structure we will compare it with the standard structure.

3.1. The cocommutator δ_f . We first describe this structure in terms of a cocommutator δ_f , the coboundary d(r(f)) of the *r*-matrix r(f). We have:

$$r(f) = (-E_{22} - E_{33}) \otimes (E_{11} + E_{33}) + E_{12} \otimes E_{21} - E_{13} \otimes E_{31} + E_{32} \otimes E_{23} - E_{13} \otimes E_{13} + E_{23} \otimes E_{23}$$

To compute δ_f we use $\delta_f(g) = d(r(f))(g) = g \cdot r(f) = [g \otimes 1 + 1 \otimes g, r(f)]$, and this gives us:

$$\begin{split} \delta_f(E_{11} + E_{33}) &= -E_{23} \wedge E_{23} \\ \delta_f(E_{22} + E_{33}) &= E_{13} \wedge E_{13} \\ \delta_f(E_{21}) &= E_{21} \wedge (E_{11} + E_{33}) - E_{23} \wedge (E_{13} + E_{31}) \\ \delta_f(E_{12}) &= E_{12} \wedge (-E_{22} - E_{33}) - (-E_{13}) \wedge (E_{23} + E_{32}) \\ \delta_f(E_{23}) &= 0 \\ \delta_f(E_{13}) &= 0 \\ \delta_f(E_{31}) &= (E_{13} + E_{31}) \wedge (E_{11} + E_{33}) + E_{21} \wedge E_{23} \\ \delta_f(E_{32}) &= (E_{23} + E_{32}) \wedge (-E_{22} - E_{33}) + (-E_{12}) \wedge E_{13} \end{split}$$

where we use the notation: $a \wedge b = a \otimes b - (-1)^{|a||b|} b \otimes a$ for any two homogeneous elements a, b.

3.2. Two subalgebras of g. Consider the following subspaces defined by f:

$$S_1 = \operatorname{Im}(f - 1) = \langle E_{11} + E_{33}, E_{21}, E_{23}, E_{13} + E_{31} \rangle$$

$$S_2 = \operatorname{Im}(f) = \langle E_{22} + E_{33}, E_{12}, E_{13}, E_{23} + E_{32} \rangle$$

The fact that r(f) is an r-matrix satisfying Eqn.(2.1) implies that these image subspaces are indeed Lie subsuperalgebras of \mathfrak{g} , (see Lemma 4 of [K]). In fact it is not difficult to see that both S_i are isomorphic as Lie superalgebras to a fourdimensional Lie superalgebra:

$$\mathfrak{s} = \mathfrak{s}_{\overline{0}} \oplus \mathfrak{s}_{\overline{1}}; \qquad \mathfrak{s}_{\overline{0}} = \langle h, x \rangle \quad \mathfrak{s}_{\overline{1}} = \langle y_1, y_2 \rangle$$

with the following relations:

$$\begin{aligned} [h,x] &= -x, & [h,y_1] &= -y_1, & [x,y_2] &= y_1, \\ [y_1,y_2] &= x, & [y_2,y_2] &= 2h, \end{aligned}$$

(any other commutator will be equal to zero). Therefore, we can write:

$$\mathfrak{g}\cong\mathfrak{s}\oplus\mathfrak{s}$$

where the direct sum is the direct sum of graded vector spaces. Note that ${\mathfrak s}$ is solvable.

Next we compute the restriction of δ_f onto the S_i . This is straightforward; on S_1 we get:

$$\delta_f(E_{11} + E_{33}) = -E_{23} \wedge E_{23}$$

$$\delta_f(E_{21}) = E_{21} \wedge (E_{11} + E_{33}) - E_{23} \wedge (E_{13} + E_{31})$$

$$\delta_f(E_{23}) = 0$$

$$\delta_f(E_{13} + E_{31}) = (E_{13} + E_{31}) \wedge (E_{11} + E_{33}) + E_{21} \wedge E_{23}$$

and the restriction onto S_2 is given by:

$$\delta_f(E_{22} + E_{33}) = E_{13} \wedge E_{13}$$

$$\delta_f(E_{12}) = E_{12} \wedge (-E_{22} - E_{33}) - (-E_{13}) \wedge (E_{23} + E_{32})$$

$$\delta_f(E_{13}) = 0$$

$$\delta_f(E_{23} + E_{32}) = (E_{23} + E_{32}) \wedge (-E_{22} - E_{33}) + (-E_{12}) \wedge E_{13}$$

In fact we can see that this gives Lie bialgebra structures to the S_i . Hence $(S_i, \delta_f|_{S_i})$ are actually Lie subbialgebras of \mathfrak{g} .

Let us now compute the Lie brackets defined on S_i^* by $\delta_f|_{S_i}$. For simplicity, we will work with the isomorphic super Lie bialgebras on \mathfrak{s} . Denote the associated cocommutators on \mathfrak{s} by δ_i ; in other words, define δ_1 and δ_2 so that $(S_1, \delta_f|_{S_1})$ is isomorphic to the super Lie bialgebra (\mathfrak{s}, δ_1) and $(S_2, \delta_f|_{S_2})$ is isomorphic to the super Lie bialgebra (\mathfrak{s}, δ_2) . Clearly we will have:

$$\begin{split} \delta_1(h) &= -y_1 \wedge y_1 & \delta_2(h) = y_1 \wedge y_1 \\ \delta_1(x) &= x \wedge h - y_1 \wedge y_2 & \delta_2(x) = -(x \wedge h - y_1 \wedge y_2) \\ \delta_1(y_1) &= 0 & \delta_2(y_1) = 0 \\ \delta_1(y_2) &= y_2 \wedge h + x \wedge y_1 & \delta_2(y_2) = -(y_2 \wedge h + x \wedge y_1) \end{split}$$

and we have $\delta_1 = -\delta_2$. In particular, we see that $(S_2, \delta_f|_{S_2})$ is (isomorphic to) the opposite super Lie bialgebra of $(S_1, \delta_f|_{S_1})$.

Recall that the Lie bracket $[\cdot, \cdot]_1$ on the dual \mathfrak{s}^* associated to δ_1 can uniquely be determined by the following: For any two elements $\alpha, \beta \in \mathfrak{s}^*$ and any element $s \in \mathfrak{s}$, we have:

$$\langle [\alpha,\beta]_1,s\rangle = \langle \alpha \otimes \beta, \delta_1(s)\rangle$$

(where $\langle \alpha, s \rangle = \alpha(s)$ is the pairing of \mathfrak{s} with its dual \mathfrak{s}^*). For instance we have:

$$\langle [y_1^*, y_1^*]_1, h \rangle = \langle y_1^* \otimes y_1^*, \delta_1(h) \rangle = - \langle y_1^* \otimes y_1^*, y_1 \wedge y_1 \rangle = -2 \langle y_1^* \otimes y_1^*, y_1 \otimes y_1 \rangle$$

= $-2(-1)^{|y_1||y_1^*|} (y_1^*(y_1))^2 = 2$

and $\langle [y_1^*, y_1^*]_1, s \rangle = 0$ for any other basis vectors of \mathfrak{s} . Therefore we get:

$$[y_1^*, y_1^*]_1 = 2h^*$$

Similarly we have:

$$\langle [y_1^*, y_2^*]_1, x \rangle = \langle y_1^* \otimes y_2^*, \delta_1(x) \rangle = \langle y_1^* \otimes y_2^*, x \wedge h - y_1 \wedge y_2 \rangle = - \langle y_1^* \otimes y_2^*, y_1 \otimes y_2 \rangle$$

= $-(-1)^{|y_1||y_2^*|} y_1^*(y_1) y_2^*(y_2) = 1$

and $\langle [y_1^*, y_2^*]_1, s \rangle = 0$ for any other basis vectors of \mathfrak{s} . Therefore we get:

$$[y_1^*, y_2^*]_1 = x^*$$

Likewise, we compute the other brackets on \mathfrak{s}^* . The nonzero brackets are:

$$\begin{split} & [h^*,x^*]_1 = -x^*, \qquad \qquad [h^*,y_2^*]_1 = -y_2^*, \qquad \qquad [x^*,y_1^*]_1 = y_2^* \\ & [y_1^*,y_2^*]_1 = x^*, \qquad \qquad [y_1^*,y_1^*]_1 = 2h^*, \end{split}$$

At this point, it is easy to notice that this is actually isomorphic to the Lie super-algebra \mathfrak{s} itself (via the map:

$$h^* \mapsto h, \qquad x^* \mapsto x, \qquad y_1^* \mapsto y_2, \qquad y_2^* \mapsto y_1).$$

Similar computations on $(S_2, \delta_f|_{S_2})$ show that the super Lie bialgebra (\mathfrak{s}, δ_2) is also self-dual. In particular, the Lie bracket $[\cdot, \cdot]_2$ on the dual \mathfrak{s}^* associated to δ_2 is given by:

$$\begin{split} & [h^*,x^*]_2 = x^*, & [h^*,y_2^*]_2 = y_2^*, & [x^*,y_1^*]_2 = -y_2^*, \\ & [y_1^*,y_2^*]_2 = -x^*, & [y_1^*,y_1^*]_2 = -2h^*, \end{split}$$

and this is isomorphic to the Lie superalgebra \mathfrak{s} itself (via the map:

 $h^* \mapsto -h, \qquad x^* \mapsto x, \qquad y_1^* \mapsto y_2, \qquad y_2^* \mapsto -y_1).$

We have seen earlier that $(S_2, \delta_f|_{S_2})$ is (isomorphic to) the opposite super Lie bialgebra of $(S_1, \delta_f|_{S_1})$. Therefore:

$$\mathfrak{g} \cong \mathfrak{s} \oplus (\mathfrak{s}^*)^{op}$$

where the direct sum is that of graded vector spaces. Here we may assume \mathfrak{s} is equipped with the super Lie bialgebra structure given by δ_1 or δ_2 , as both are self-dual.

3.3. The Drinfeld double of \mathfrak{s} . Let \mathfrak{d} be the Drinfeld double of (\mathfrak{s}, δ_2) , (we can carry out the following using (\mathfrak{s}, δ_1) instead, our choice is in fact arbitrary). The computations of 3.2 can be used to conclude that $\mathfrak{d} \cong \mathfrak{g}$ as a Lie superalgebra. Explicitly the map $i_1 : \mathfrak{s} \to \mathfrak{g}$ given on the generators of \mathfrak{s} as:

$h \mapsto E_{22} + E_{33}$	$x \mapsto E_{12}$
$y_1 \mapsto E_{13}$	$y_2 \mapsto E_{23} + E_{32}$

and the map $i_2: (\mathfrak{s}^*)^{op} \to \mathfrak{g}$ given on the generators as:

$h^* \mapsto -(E_{11} + E_{33})$	$x^* \mapsto E_{21}$
$y_1^* \mapsto -E_{13} - E_{31}$	$y_2^* \mapsto E_{23}$

are both super Lie bialgebra homomorphisms, and $\text{Im}(i_1) = S_2$ and $\text{Im}(i_2) = S_1$.

The inner product defined on \mathfrak{d} is given by:

$$\langle s_1 + \alpha_1, s_2 + \alpha_2 \rangle =_{def} \alpha_1(s_2) + (-1)^{|\alpha_2||s_1|} \alpha_2(s_1)$$

Clearly S_i are both isotropic with respect to this form. We only need to consider $\langle \alpha_1, s_2 \rangle$ and $\langle s_1, \alpha_2 \rangle$ where $s_i \in \text{Im}(i_1)$ and $\alpha_i \in \text{Im}(i_2)$. Now,

$$\langle \alpha_1, s_2 \rangle = \alpha_1(s_2), \qquad \langle s_1, \alpha_2 \rangle = (-1)^{|\alpha_2||s_1|} \alpha_2(s_1)$$

and we see that this form actually coincides with the super trace form on \mathfrak{g} . For example we can compute:

$$\langle E_{23}, E_{32} \rangle = \langle E_{23}, (E_{23} + E_{32}) - E_{23} \rangle = \langle E_{23}, (E_{23} + E_{32}) \rangle$$

$$= \langle i_2(y_2^*), i_1(y_2) \rangle = y_2^*(y_2) = 1$$

$$\langle E_{13}, E_{31} \rangle = \langle E_{13}, -(-E_{13} - E_{31}) - E_{13} \rangle = \langle E_{13}, -(-E_{13} - E_{31}) \rangle$$

$$= \langle i_1(y_1), -i_2(y_1^*) \rangle = -(-1)^{|y_1||y_1^*|} y_1^*(y_1) = 1$$

We know the super Lie algebra structure when restricted to S_1 and S_2 . From the invariance of the form we can find the mixed brackets. In other words we use:

$$\langle [\alpha_1, s_1], s_2 \rangle = \langle \alpha_1, [s_1, s_2] \rangle \qquad \langle [s_1, \alpha_1], \alpha_2 \rangle = \langle s_1, [\alpha_1, \alpha_2] \rangle$$

GIZEM KARAALI

where $\alpha_i \in \text{Im}(i_2) = S_1$ and $s_i \in \text{Im}(i_1) = S_2$. Some more computation shows that indeed the Lie superalgebra structure on \mathfrak{d} is the usual one, in other words, we find that $\mathfrak{d} \cong \mathfrak{g}$ as a Lie superalgebra.

The super Lie bialgebra structure on \mathfrak{d} is given by $\delta|_{\mathrm{Im}(\mathfrak{s})} + \delta|_{\mathrm{Im}(\mathfrak{s}^*)^{op}}$ or equivalently by $\delta|_{\mathrm{Im}(\mathfrak{s})} + \delta|_{\mathrm{Im}(\mathfrak{s}^*)}$. But this is equal to $\delta_f|_{S_1} + \delta_f|_{S_2} = \delta_f$. Hence, $\mathfrak{g} = sl(2,1)$ with the super Lie bialgebra structure δ_f is the Drinfeld double of the four dimensional solvable Lie superalgebra \mathfrak{s} .

3.4. The standard structure. It may be interesting to compare the above with the standard Lie bialgebra structure on \mathfrak{g} . Note that

$$r_s = (-E_{22} - E_{33}) \otimes (E_{11} + E_{33}) + E_{12} \otimes E_{21} - E_{13} \otimes E_{31} + E_{32} \otimes E_{23}$$

gives us an r-matrix satisfying Eqn.(2.1) and hence is a standard r-matrix constructed by Belavin-Drinfeld type data. Using $\delta_s = d(r_s)$ to find the associated cocommutator δ_s we see that:

$$\begin{split} \delta_s(E_{11} + E_{33}) &= 0\\ \delta_s(E_{22} + E_{33}) &= 0\\ \delta_s(E_{21}) &= E_{21} \wedge (E_{11} + E_{33}) - E_{23} \wedge E_{31}\\ \delta_s(E_{12}) &= E_{12} \wedge (-E_{22} - E_{33}) - (-E_{13}) \wedge E_{32}\\ \delta_s(E_{23}) &= 0\\ \delta_s(E_{13}) &= 0\\ \delta_s(E_{13}) &= E_{31} \wedge (E_{11} + E_{33})\\ \delta_s(E_{32}) &= E_{32} \wedge (-E_{22} - E_{33}) \end{split}$$

Restriction of δ_s onto the S_i would not give us well-defined maps on the S_i . Instead, we may consider its restriction onto two other solvable subalgebras of \mathfrak{g} : Define

$$T_1 = \langle E_{11} + E_{33}, E_{21}, E_{23}, E_{31} \rangle,$$

$$T_2 = \langle E_{22} + E_{33}, E_{12}, E_{13}, E_{32} \rangle.$$

Then on T_1 we get:

$$\delta_s(E_{11} + E_{33}) = 0$$

$$\delta_s(E_{21}) = E_{21} \wedge (E_{11} + E_{33}) - E_{23} \wedge E_{31}$$

$$\delta_s(E_{23}) = 0$$

$$\delta_s(E_{31}) = E_{31} \wedge (E_{11} + E_{33})$$

and the restriction of δ_s onto T_2 is given by:

$$\delta_s(E_{22} + E_{33}) = 0$$

$$\delta_s(E_{12}) = E_{12} \wedge (-E_{22} - E_{33}) - (-E_{13}) \wedge E_{32}$$

$$\delta_s(E_{13}) = 0$$

$$\delta_s(E_{32}) = E_{32} \wedge (-E_{22} - E_{33})$$

It is easy to see that $(T_i, \delta_s|_{T_i})$ is a Lie subbialgebra of (\mathfrak{g}, δ_s) for each *i*.

It is clear that $T_i \cong \mathfrak{t}$ where \mathfrak{t} is the four dimensional solvable Lie superalgebra

$$\mathfrak{t} = \mathfrak{t}_{\overline{0}} \oplus \mathfrak{t}_{\overline{1}}; \qquad \qquad \mathfrak{t}_{\overline{0}} = \langle h, x \rangle \quad \mathfrak{t}_{\overline{1}} = \langle y_1, y_2 \rangle$$

with the following relations:

$$[h, x] = -x,$$
 $[h, y_1] = -y_1,$ $[y_1, y_2] = x_1,$

(any other commutator will be equal to zero). Therefore, we can write:

$$\mathfrak{g}\cong\mathfrak{t}\oplus\mathfrak{t}$$

where the direct sum is that of graded vector spaces.

Let us now compute the Lie brackets defined on T_i^* by $\delta_s|_{T_i}$. For simplicity, we will work with the isomorphic super Lie bialgebras on t. Denote the associated cocommutators on t by δ_{si} ; in other words, define δ_{s1} and δ_{s2} so that $(T_1, \delta_s|_{T_1})$ is isomorphic to the super Lie bialgebra (t, δ_{s1}) and $(T_2, \delta_s|_{T_2})$ is isomorphic to the super Lie bialgebra (t, δ_{s2}) . Clearly we will have:

$$\begin{split} \delta_{s1}(h) &= 0 & \delta_{s2}(h) = 0 \\ \delta_{s1}(x) &= x \wedge h - y_1 \wedge y_2 & \delta_{s2}(x) = -(x \wedge h - y_1 \wedge y_2) \\ \delta_{s1}(y_1) &= 0 & \delta_{s2}(y_1) = 0 \\ \delta_{s1}(y_2) &= y_2 \wedge h & \delta_{s2}(y_2) = -y_2 \wedge h \end{split}$$

and we have $\delta_{s1} = -\delta_{s2}$. In particular, we see that $(T_2, \delta_s|_{T_2})$ is (isomorphic to) the opposite super Lie bialgebra of $(T_1, \delta_s|_{T_1})$.

At this point, the Lie bracket $[\cdot, \cdot]_1$ on the dual \mathfrak{t}^* associated to δ_{s1} can be easily determined. The nonzero brackets are:

$$[h^*, x^*]_1 = -x^*,$$
 $[h^*, y_2^*]_1 = -y_2^*,$ $[y_1^*, y_2^*]_1 = x^*$

Notice that this is actually isomorphic to the Lie superalgebra t itself (via the map:

$$h^* \mapsto h, \qquad x^* \mapsto x, \qquad y_1^* \mapsto y_2, \qquad y_2^* \mapsto y_1).$$

As $(T_2, \delta_s|_{T_2})$ is (isomorphic to) the opposite super Lie bialgebra of $(T_1, \delta_s|_{T_1})$, we get:

$$\mathfrak{g}\cong\mathfrak{t}\oplus(\mathfrak{t}^*)^{op}$$

where the direct sum is that of graded vector spaces, and we are considering \mathfrak{t} with the super Lie bialgebra structure given by δ_{s1} . Of course, we can show that $(\mathfrak{t}, \delta_{s2})$ is self-dual as well, and hence the above identity would still hold if we assumed that the super Lie bialgebra structure on \mathfrak{t} is the one associated to δ_{s2} .

Now arguments similar to those in 3.2 and 3.3 can be used to conclude that \mathfrak{g} is isomorphic to the double of \mathfrak{t} as a Lie superalgebra. Explicitly the map $i_{s_1} : \mathfrak{t} \to \mathfrak{g}$ given on the generators of \mathfrak{t} as:

$$\begin{aligned} h \mapsto E_{22} + E_{33} & x \mapsto E_{12} \\ y_1 \mapsto E_{13} & y_2 \mapsto E_{32} \end{aligned}$$

and the map $i_{s2}: (\mathfrak{t}^*)^{op} \to \mathfrak{g}$ given on the generators as:

$$\begin{aligned} h^* &\mapsto -(E_{11} + E_{33}) & x^* &\mapsto E_{21} \\ y_1^* &\mapsto -E_{31} & y_2^* &\mapsto E_{23} \end{aligned}$$

are both super Lie bialgebra homomorphisms, and $\text{Im}(i_{s1}) = T_2$ and $\text{Im}(i_{s2}) = T_1$.

GIZEM KARAALI

The inner product defined on \mathfrak{g} by this double structure is given by:

$$\langle s_1 + \alpha_1, s_2 + \alpha_2 \rangle =_{def} \alpha_1(s_2) + (-1)^{|\alpha_2||s_1|} \alpha_2(s_1)$$

Clearly T_i are both isotropic with respect to this form. We only need to consider $\langle \alpha_1, t_2 \rangle$ and $\langle t_1, \alpha_2 \rangle$ where $t_i \in \text{Im}(i_{s_1})$ and $\alpha_i \in \text{Im}(i_{s_2})$, but

$$\langle \alpha_1, s_2 \rangle = \alpha_1(s_2), \qquad \langle s_1, \alpha_2 \rangle = (-1)^{|\alpha_2||s_1|} \alpha_2(s_1)$$

and we see that this form also coincides with the super trace form on \mathfrak{g} .

We know the super Lie algebra structure when restricted to T_1 and T_2 . From the invariance of the form we can find the mixed brackets. In other words we use:

$$\langle [\alpha_1, t_1], t_2 \rangle = \langle \alpha_1, [t_1, t_2] \rangle \qquad \langle [t_1, \alpha_1], \alpha_2 \rangle = \langle t_1, [\alpha_1, \alpha_2] \rangle$$

where $\alpha_i \in \text{Im}(i_{s2}) = T_1$ and $t_i \in \text{Im}(i_{s1}) = T_2$. Some more computation shows that indeed the Lie superalgebra structure on \mathfrak{g} is the usual one.

The super Lie bialgebra structure on \mathfrak{g} coming from this double construction is given by $\delta|_{\mathrm{Im}(\mathfrak{t})} + \delta|_{\mathrm{Im}(\mathfrak{t}^*)^{op}}$ or equivalently by $\delta|_{\mathrm{Im}(\mathfrak{t})} + \delta|_{\mathrm{Im}(\mathfrak{t}^*)}$. But this is equal to $\delta_s|_{T_1} + \delta_s|_{T_2} = \delta_s$. Therefore, $\mathfrak{g} = sl(2, 1)$ with the standard super Lie bialgebra structure is the Drinfeld double of the four dimensional solvable Lie superalgebra \mathfrak{t} (equipped with the super Lie bialgebra structure given either by δ_{s_1} or δ_{s_2}).

4. Conclusion

Unlike in the non-graded case, super r-matrices satisfying Eqn.(2.1) may not be obtained via a simple modification of the Belavin-Drinfeld construction. In this note, we have studied the Lie bialgebra structure associated to one such r-matrix on sl(2,1), and we have shown that it has a nice description as the double of a four dimensional subalgebra. In the non-graded case, such structures only arise from twists of the standard r-matrix. Our r-matrix is not of this form, but shows similarities to such. These similarities may lead to an understanding of these special types of r-matrices that do not fit a Belavin-Drinfeld type description.

References

- [A] Andruskiewitsch, N.; "Lie superbialgebras and Poisson-Lie supergroups", Abh. Math. Sem. Univ. Hamburg 63 (1993), pp.147–163.
- [BD1] Belavin, A. A., Drinfeld, V. G.; "Solutions of the classical Yang-Baxter equation and simple Lie algebras", Funct. Anal. Appl. 16 (1982), pp.159–180.
- [BD2] Belavin, A. A., Drinfeld, V. G.; "Triangle equation and simple Lie algebras", Soviet Scientific Reviews Sect. C 4 (1984), pp.93–165.
- [CP] Chari, V., Pressley, A.; A Guide to Quantum Groups, Cambridge University Press, 1995.
- [ES] Etingof, P., Schiffmann, O.; Lectures on Quantum Groups, International Press, 1998.
- [F] Fuks, D. B.; Cohomology of Infinite-Dimensional Lie Algebras, (Contemporary Soviet Mathematics), 1986.
- [GZB] Gould, M. D., Zhang, R. B., Bracken, A. J.; "Lie bi-superalgebras and the graded classical Yang-Baxter equation", Rev. Math. Phys. 3 (1991), no. 2, pp.223–240.
- [K] Karaali, G.; "Constructing r-matrices on simple Lie superalgebras", J. Algebra 282 (2004), no.1, pp.83–102.
- [L] Leites, D., Shapovalov, A.; "Manin-Olshansky triples for Lie superalgebras", J. Nonlinear Math. Phys. 7 (2000), no. 2, pp.120–125.

- [SZ] Scheunert, M., Zhang, R. B.; "Cohomology of Lie superalgebras and their generalizations", J. Math. Phys. 39 (1998), pp.5024–5061.
- [V] Voronov, T.; "Graded manifolds and Drinfeld doubles for Lie bialgebroids", In: Quantization, Poisson Brackets and Beyond, Contemp. Math. 315, Amer. Math. Soc., Providence, RI, 2002, pp.131–168, arXiv:math.DG/0105237.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106

 $E\text{-}mail\ address: \texttt{gizem@math.ucsb.edu}$