

Characterization of quasi-Banach spaces which coarsely embed into a Hilbert space

N. Lovasoa RANDRIANARIVONY ¹

A (not necessarily continuous) map f between two metric spaces (X, d) and (Y, δ) is called a *coarse embedding* (see [G, 7.G]) if there exist two non-decreasing functions $\varphi_1 : [0, \infty) \rightarrow [0, \infty)$ and $\varphi_2 : [0, \infty) \rightarrow [0, \infty)$ such that

1. $\varphi_1(d(x, y)) \leq \delta(f(x), f(y)) \leq \varphi_2(d(x, y))$
2. $\varphi_1(t) \rightarrow \infty$ as $t \rightarrow \infty$.

It was proved in [JR] that ℓ_p does not coarsely embed into a Hilbert space when $p > 2$. The present article is a strengthening of that result by giving a full characterization of quasi-Banach spaces that coarsely embed into a Hilbert space. This result, as well as its proof, mirrors the theorem in [AMM] which characterizes spaces that uniformly embed into a Hilbert space. The combination of Theorem 1 and the theorem in [AMM] yield that a quasi-Banach space uniformly embeds into a Hilbert space if and only if it coarsely embeds into a Hilbert space. This is counterintuitive in that a uniform embedding gives information only on small distances while a coarse embedding gives information only on large distances.

Theorem 1 *A quasi-Banach space X coarsely embeds into a Hilbert space if and only if there is a probability space $(\Omega, \mathcal{B}, \mu)$ such that X is linearly isomorphic to a subspace of $L_0(\mu)$.*

To prove Theorem 1, we use a result essentially contained in [JR] as is recalled in the following Proposition:

¹Supported in part by NSF 0200690 and Texas Advanced Research Program 010366-0033-20013.

This paper represents a portion of the author's dissertation being prepared at Texas A&M University under the direction of W. B. Johnson.

Proposition 1 *Let X be a quasi-Banach space which coarsely embeds into a Hilbert space. Then there exists on X a continuous negative definite function g which satisfies $g(0) = 0$ and $\phi_1(\|x\|) \leq g(x) \leq \|x\|^{2\alpha}$ where $\phi_1 : [0, \infty) \rightarrow [0, \infty)$ is a nondecreasing function satisfying $\phi_1(t) \rightarrow \infty$ as $t \rightarrow \infty$, and $\alpha > 0$.*

Proof: Steps 0, 1, 2 (and a piece of Step 3 on the continuity of g) in [JR] extend to the case of quasi-Banach spaces.

Proof of Theorem 1:

Let X be a quasi-Banach space. A theorem of Aoki-Rolewicz gives an equivalent quasi-norm $\|\cdot\|$ on X which is also p -additive for some $0 < p \leq 1$, i.e. $\|x+y\|^p \leq \|x\|^p + \|y\|^p$ for all x, y in X . In particular X under this norm has type p .

Say X is linearly isomorphic to a subspace of $L_0(\mu)$ for some probability space $(\Omega, \mathcal{B}, \mu)$. Then since X has type p , X is isomorphic to a subspace of $L_r(\mu)$ for every $r < p$. (See [BL], Theorem 8.15.)

Now since $r < 2$, Nowak's result [N] implies that X coarsely embeds into a Hilbert space.

Conversely, let X be a quasi-Banach space which coarsely embeds into a Hilbert space. Let g be the negative definite function on X given by Proposition 1, and let f be the continuous positive definite function given by $f = e^{-g}$. Use Proposition 8.7 in [BL] to get a probability space $(\Omega, \mathcal{B}, \mu)$ and a continuous linear operator $U : X \rightarrow L_0(\mu)$ such that the characteristic function $\mathbb{E} \exp(itUx)$ of Ux is equal to $f(tx)$ for every $x \in X$ and $t \in \mathbb{R}$. We show that U is an isomorphism into.

Let $(x_n)_n$ be a sequence in X such that $U(x_n) \rightarrow 0$ in $L_0(\mu)$, i.e. in measure. Then $f(tx_n) = \mathbb{E}(\exp(itUx_n)) \rightarrow 1$ for each fixed t in \mathbb{R} .

If $(x_n)_n$ does not converge to 0, then by passing to a subsequence we can assume without loss of generality that $\|x_n\| \geq \epsilon$ for all n and for some $\epsilon > 0$. But since ϕ_1 is nondecreasing, we get for every $t > 0$:

$$e^{-\phi_1(t\|x_n\|)} \leq e^{-\phi_1(t\epsilon)}.$$

Since $\phi_1(s) \rightarrow \infty$ as $s \rightarrow \infty$, we can pick $t_0 > 0$ so that $e^{-\phi_1(t_0\epsilon)} < \frac{1}{2}$. For that t_0 , we have for every n :

$$f(t_0x_n) \leq e^{-\phi_1(t_0\epsilon)} < \frac{1}{2}.$$

This contradicts the fact that $f(t_0x_n) \rightarrow 1$.

Thus $x_n \rightarrow 0$, and hence U is one-to-one and its inverse is continuous.

Acknowledgement: I would like to thank W. B. Johnson for the constructive conversations that led to the production of this article.

References

- [AMM] I. Aharoni, B. Maurey, and B. S. Mityagin, Uniform embeddings of metric spaces and of Banach spaces into Hilbert spaces, *Israel J. Math.* 52(3) (1985), 251–265.
- [BL] Y. Benyamini and J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, Coll. Pub. 48, Amer. Math. Soc., Providence, RI (2000).
- [G] M. Gromov, Asymptotic invariants for infinite groups, *Geometric Group Theory*, G. A. Nibto and M. A. Roller, ed., Cambridge University Press (1993), 1–295.
- [JR] W. B. Johnson, N. L. Randrianarivony, ℓ_p ($p > 2$) does not coarsely embed into a Hilbert space, (preprint)
- [N] P. Nowak, Coarse embeddings of metric spaces into Banach spaces, (preprint).

N. Lovasoa Randrianarivony
Department of Mathematics
Texas A&M University
College Station, TX, USA
E-mail: nirina@math.tamu.edu