CYCLE CLASSES OF THE E-O STRATIFICATION ON THE MODULI OF ABELIAN VARIETIES

TORSTEN EKEDAHL AND GERARD VAN DER GEER

ABSTRACT. We introduce a stratification on the space of symplectic flags on the de Rham bundle of the universal principally polarized abelian variety in positive characteristic and study its geometric properties like irreducibility of the strata and we calculate the cycle classes. When the characteristic p is treated as a formal variable these classes can be seen as a deformation of the classes of the Schubert varieties for the corresponding classical flag variety (the classical case is recovered by putting p equal to 0). We relate our stratification with the E-O stratification on the moduli space of principally polarized abelian varieties of a fixed dimension and derive properties of the latter. Our results are strongly linked with the combinatorics of the Weyl group of the symplectic group.

1. INTRODUCTION

The moduli space \mathcal{A}_g of principally polarized abelian varieties of dimension g is defined over the integers. For the characteristic zero fibre $\mathcal{A}_q \otimes \mathbb{C}$ we have an explicit description as an orbifold $\operatorname{Sp}_{2q}(\mathbb{Z})\setminus\mathcal{H}_g$ with \mathcal{H}_g the Siegel upper half space of degree g. It is a recent insight, though, that perhaps the positive characteristic fibres $\mathcal{A}_g \otimes \mathbb{F}_p$ are more accessible than the characteristic zero one. A good illustration of this is provided by the E-O stratification of $\mathcal{A}_g \otimes \mathbb{F}_p$, a stratification consisting of 2^{g} strata, each of which is quasi-affine. It was originally defined by Ekedahl and Oort (see [Oo01]) by analyzing the structure of the kernel of multiplication by p of an abelian variety. It turns out that this group scheme can assume 2^g forms only, and this led to the strata. For g = 1 the two strata are the locus of ordinary and that of supersingular elliptic curves. Some strata possess intriguing properties. For example, the stratum of abelian varieties of p-rank 0 is a complete subvariety of $\mathcal{A}_q \otimes \mathbb{F}_p$ of codimension g, the smallest codimension possible. No analogue in characteristic 0 of this stratum nor of the stratification is known, and in fact, Keel and Sadun ([KS03]) proved that complete subvarieties of $\mathcal{A}_q \otimes \mathbb{C}$ of codimension g do not exist for $q \geq 3$.

While trying to find cycle classes for the E-O strata we realised that the strata could be described as degeneration loci for maps between vector bundles and as such loci are indexed by Young diagrams our attention was turned towards the combinatorics of the Weyl group. When considered in this light it is clear that much of the combinatorics of [Oo01] is closely related to the Weyl group W_g of Sp_{2g}, which is the group relevant for the analytic description of $\mathcal{A}_g \otimes \mathbb{C}$. The main idea of this paper is to try to make this connection more explicit. More precisely, the combinatorics of the E-O strata is most closely related to the combinatorics

¹⁹⁹¹ Mathematics Subject Classification. 14K10.

CLASSES.TEXOctober 22, 2018

associated to W_g and the sub-Weyl group corresponding to the maximal parabolic subgroup P of elements of Sp_{2g} fixing a maximal isotropic subspace in the 2gdimensional symplectic vector space. Indeed, this sub-Weyl group is S_g , the group of permutations on g letters (embedded as a sub-Weyl group in W_g), and the E-O strata are in bijection with the cosets W_g/S_g ; we shall use the notation \mathcal{V}_{ν} for the (open) stratum of $\mathcal{A}_g \otimes \mathbb{F}_p$ corresponding to $\nu \in W_g/S_g$ (and $\overline{\mathcal{V}}_{\nu}$ for its closure). The coset space W_g/S_g is also in bijection with the set of Bruhat cells in the space of maximal totally isotropic flags Sp_{2g}/P and we believe this to be no accident. (The formal relation between \mathcal{A}_g and Sp_{2g}/P is that Sp_{2g}/P is the *compact dual* of \mathcal{H}_g .)

In order to push the analogy further we introduce a "flag space" $\mathcal{F}_q \to \mathcal{A}_q$ whose fibres are isomorphic to the fibres of the map $\operatorname{Sp}_{2q}/B \to \operatorname{Sp}_{2q}/P$, where B is a Borel subgroup of P. In positive characteristic we define (and this definition clearly makes sense only in positive characteristic) a stratification of \mathcal{F}_{g} whose open strata \mathcal{U}_w and corresponding closures $\overline{\mathcal{U}}_w$ are parametrised by the elements of W_q . This is very similar to the Bruhat cells of Sp_{2q}/B and their closures, the Schubert strata, which are also parametrised by the elements of W_q . Our first main result is that this is more than a similarity when one is working locally; we show that for each point of \mathcal{F}_g there is a stratum preserving local isomorphism (in the étale topology) taking the point to some point of Sp_{2q}/B . As a lot is known about the local structure of the Schubert varieties we immediately get a lot of information about the local structure of our strata. The first consequence is that the dimension of $\overline{\mathcal{U}}_w$ is everywhere equal to the length of w. A very important consequence is that the $\overline{\mathcal{U}}_w$ are all normal; this situation differs markedly from the case of the closed E-O strata which in general are not normal. Another consequence is that the inclusion relation between the strata is given exactly by the Bruhat-Chevalley order on W_q . (A much more sophisticated consequence is that the local structure of the ℓ -adic intersection complex for a closed stratum is the same as for the Schubert varieties and in particular that the dimensions of its fibres over the open strata of the closed stratum are given by the Kazhdan-Lusztig polynomials. We shall not however pursue that in this article.)

We make several applications of our results on the structure of the strata $\overline{\mathcal{U}}_w$. The first, and most important, is that by construction the strata \mathcal{U}_w are defined as the locus where two symplectic flags on the same vector bundle are in relative position given by w. As we have shown that they have the expected codimension and are reduced we can use formulas of Fulton, and those of Pragacz and Ratajski as crystallized in the formulas of Kresch and Tamvakis to get formulas for the cycle classes of the strata. A result of Fulton give such formulas for all strata but in terms of a recursion formula that we have not been able to turn into a closed formula; even though these formulas should have independent interest we can use them to get formulas for the E-O strata as follows. If $w \in W_g$ is minimal for the Bruhat-Chevalley order in its coset wS_q , then \mathcal{U}_w maps by a finite étale map to the open E-O stratum \mathcal{V}_{ν} corresponding to the coset $\nu := wS_q$. We can compute the degree of this map in terms of the combinatorics of the element w and we then can push down our formula for \mathcal{U}_w to obtain formulas for the cycle classes of the E-O strata. Also the formulas of Kresch and Tamvakis can be used to give the classes of E-O strata. One interesting general consequence is that each class is a polynomial in the Chern classes λ_i of the Hodge bundle whose coefficients are polynomials in p.

This is a phenomenon already visible in the special cases of our formula that were known previously; the oldest such example being Deuring's mass formula for the number of supersingular elliptic curves (weighted by one over the cardinalities of their automorphism groups) that says that this mass is (p-1)/12. This appears in our context as the combination of the formula $(p-1)\lambda_1$ for the class of the supersingular locus and the formula deg $\lambda_1 = 1/12$. We interpret these results as giving rise to elements in the *p*-tautological ring; this is the ring obtained from the usual *tautological ring*, the ring generated by the Chern classes of the Hodge bundle, by extending the scalars to $\mathbb{Z}\{p\}$, the localisation of the polynomial ring $\mathbb{Z}[p]$ at the polynomials with constant coefficient 1. Hence we get elements parametrised by W_q/S_q in the p-tautological ring and we show that they form a basis for the p-tautological ring. Putting p equal to 0 maps these elements to elements of the ordinary tautological ring which can identified with the Chow ring of Sp_{2q}/P and these elements are the usual classes of the Schubert varieties. It seems that these results call for a p-Schubert calculus in the sense of a better understanding of these elements of the *p*-tautological ring and for instance their behaviour under multiplication.

However, there seems be a more intriguing problem. We have for each $w \in W_g$ a stratum in our flag space and they push down to elements of the *p*-tautological ring under the projection map to $\tilde{\mathcal{A}}_g$ (a toroidal compactification of \mathcal{A}_g). When setting *p* to 0 these elements specialise to the classes of the images of the Schubert varieties of $\operatorname{Sp}_{2g}/B_g$ in Sp_{2g}/P and for them the situation is very simple, either *w* is minimal in its S_g coset and then the Schubert variety maps birationally to the corresponding Schubert variety of Sp_{2g}/P or it is not and then it maps to 0. However, when it comes to the elements of the *p*-tautological ring this only allows us to conclude – in the non-minimal case – that the coefficients are divisible by *p* and indeed in general they are not zero. We show that unless they map to 0 they will always map to a multiple of a class of an E-O stratum. When the element, but our considerations give an extension of this map from elements minimal in their cosets to a larger class of elements. We give some examples of this map but in general it seems a very mysterious construction.

Another application is to the irreducibility of our strata (and hence also to the strata of the E-O stratification as they are images of some of our strata). As the strata are normal this is equivalent to the connectedness of a stratum and this connectedness can sometimes be proved by an arithmetic argument. It is natural to ask if this method produces all the irreducible strata and for the characteristic large enough (the size depending on g) we can show that indeed it does. This is done using a Pieri type formula for our strata obtained by applying a result of Pittie and Ram. A Pieri type formula for multiplying the class of a connected cycle by an ample line bundle has as a consequence that a part of boundary is supported by an ample line bundle and hence that it is connected. Exploiting that together with the fact that λ_1 is an ample line bundle on \mathcal{A}_g allows to show the desired converse. We are forced to assume that the characteristic is large (and are unable to specify how large) as we need to know how high a power of λ_1 one needs to twist the exterior powers of the dual of the Hodge bundle to make it generated by global sections.

There is a particular element of $w_{\emptyset} \in W_g$ which is the largest of the elements that are minimal in their right S_g -cosets which has the property that $\overline{\mathcal{U}}_{w_{\emptyset}}$ maps birationally onto \mathcal{A}_q . It is really the strata that are contained in this stratum that seem geometrically related to \mathcal{A}_g and indeed the elements $w \in W_g$ lying below w_{\emptyset} are the ones of most interest to us. (The rest of \mathcal{F}_g appears mostly as a technical device for relating our strata to the Schubert varieties.) It should be of particular interest to understand the map $\overline{\mathcal{U}}_{w_{\emptyset}} \subset \mathcal{F}_g \to \mathcal{A}_g$. It follows from a result of Oort on Dieudonné modules that the inverse image of an open E-O stratum under this map is a locally constant fibration which focuses particular interest on the fibres of the map, this fibre depending only on the element of $\nu \in W_q/S_q$ that specifies the E-O stratum. We call these fibres punctual flag spaces, using the notation \mathcal{F}_{ν} for the one associated to ν , and establish some basic properties of them: We determine their connected components, showing in particular that two points in the same connected component can be connected by a sequence of quite simple rational curves. We also show that knowing which strata \mathcal{U}_w have non-empty intersections with a given punctual flag space would determine the inclusion relations between the E-O strata. We intend to make a more detailed study of the punctual flag spaces and show how that knowledge will give more information on these inclusion relations in a sequel to this paper.

The geometric points of the stratum $\overline{\mathcal{U}}_{w_{\emptyset}}$ correspond to symplectic flags of subgroup schemes of the kernel of multiplication by p on the principally polarized abelian variety that is given by the image of the point in \mathcal{A}_g . This is of course is reminiscent of de Jong's moduli stack $\mathcal{S}(g,p)$ of $\Gamma_0(p)$ -structures. The major difference (apart from the fact that $\overline{\mathcal{U}}_{w_{\emptyset}}$ only makes sense in positive characteristic) is that the g-dimensional element of the flag is determined by the abelian variety in our case. We shall indeed identify $\overline{\mathcal{U}}_{w_{\emptyset}}$ with the component of $\mathcal{S}(g,p)$ that is the closure of the ordinary abelian varieties provided with a flag on the local part of the kernel of multiplication by p. As a consequence we get that that component of $\mathcal{S}(g,p)$ is normal and Cohen-Macaulay.

This paper is clearly heavily inspired by [Oo01]. The attentive reader will notice that we reprove some of the results of that paper, sometimes with proofs that are very close to the proofs used by Oort. We justify such duplications by our desire to emphasize the relations with the combinatorics of W_g and the flag spaces. Hence, we start with (a rather long) combinatorial section where the combinatorial aspects have been separated from the geometric ones. We hope that this way of presenting the material will be as clarifying to the reader as it has been to us. We intend to continue to exploit the relations with the flag spaces in a future paper that will deal with K3-surfaces and one on Dieudonné modules. Since its announcement in [Ge99] our idea of connecting the E-O stratification on \mathcal{A}_g with the Weyl group and filtrations on the de Rham cohomology has been taken up in other work. In this connection we want to draw attention to papers by Moonen and Wedhorn, cf., [Mo01, MW04].

We would like to thank Piotr Pragacz for some useful comments.

Conventions. We shall exclusively work in positive characteristic p > 0. After having identified final types and final elements in Section 2 we shall often use the same notation for the final type (which is a function on $\{1, \ldots, 2g\}$) and the corresponding final element (which is an element of the Weyl group W_g). In Sections 10 and 11 our strata will be considered in flag spaces over not just \mathcal{A}_g and $\tilde{\mathcal{A}}_g$ but also over the corresponding moduli stacks with a level structure.

Contents

1. Introduction	1
2. Combinatorics	5
3. The Flag Space	16
4. Strata on the Flag Space	19
5. Extension to the boundary	27
6. Existence of boundary compo	nents 29
7. Superspecial fibres	30
8. Local structure of strata	33
9. Punctual flag spaces	35
10. Pieri formulas	39
11. Irreducibility properties	44
12. The Cycle Classes	47
13. Tautological rings	53
14. Comparison with $\mathcal{S}(g, p)$.	55
15. Appendix $g = 3$	57
References	58

2. Combinatorics

This section is of a preparatory nature and deals with the combinatorial aspects of the E-O stratification. The combinatorics is determined by the Weyl group of the symplectic group of degree g. A general reference for the combinatorics of Weyl groups is [BL00]. We start by recalling some general notations and facts about W_g and its Bruhat-Chevalley order. We then go on to give various descriptions of the minimal elements in the S_g cosets (which we presume are well-known). The short subsection on shuffles will be used to understand the rôle that the multiplicative and étale part of the Barsotti-Tate group play in our stratification in the case of positive p-rank.

2.1. Final Elements in The Weyl Group. The Weyl group W_g of type C_g in Cartan's terminology is isomorphic to the semi-direct product $S_g \ltimes (\mathbb{Z}/2\mathbb{Z})^g$, where the symmetric group S_g on g letters acts on $(\mathbb{Z}/2\mathbb{Z})^g$ by permuting the g factors. Another description of this group, and the one we shall use here, is as the subgroup of the symmetric group S_{2g} of elements which map any symmetric 2-element subset of $\{1, \ldots, 2g\}$ of the form $\{i, 2g + 1 - i\}$ to a subset of the same type:

$$W_g = \{ \sigma \in S_{2g} : \sigma(i) + \sigma(2g+1-i) = 2g+1 \text{ for } i = 1, \dots, g \}.$$

The function $i \mapsto 2g + 1 - i$ on the set $\{1, \ldots, 2g\}$ will occur frequently. We shall sometimes use the notation \overline{i} for 2g + 1 - i. Using it we can say that $\sigma \in S_{2g}$ is an element of W_g precisely when $\sigma(\overline{i}) = \overline{\sigma(i)}$ for all *i*. This makes the connection with another standard description of W_g , namely as a group of signed permutations. An element in this Weyl group has a length and a codimension defined by:

$$\ell(w) = \#\{i < j \le g : w(i) > w(j)\} + \#\{i \le j \le g : w(i) + w(j) > 2g + 1\}$$

and

$$\operatorname{codim}(w) = \#\{i < j \le g \colon w(i) < w(j)\} + \#\{i \le j \le g \colon w(i) + w(j) < 2g + 1\}$$

and these satisfy the equality

$$\ell(w) + \operatorname{codim}(w) = g^2.$$

We shall use the following notation for elements in W_g . By $[a_1, a_2, \ldots, a_{2g}]$ we mean the permutation of $\{1, 2, \ldots, 2g\}$ with $\sigma(i) = a_i$. Since $\sigma(i)$ determines $\sigma(2g+1-i)$ for $1 \leq i \leq g$ sometimes we use the notation $[a_1, \ldots, a_g]$ instead (when $g \leq 4$ we shall even dispence with the commas and write $[a_1 \ldots a_g]$ which should cause no confusion as all the a_i must be single digit numbers. We shall also use cycle notation for permutations. In particular, for $1 \leq i < g$ we let $s_i \in S_{2g}$ be the permutation (i, i+1)(2g-i, 2g+1-i) in W_g which interchanges i and i+1 and we let $s_g = (g, g+1) \in S_{2g}$. Then $(W = W_g, S = \{s_1, \ldots, s_g\})$ is a Coxeter system.

Let (W, S) be a Coxeter system and $a \in W$. If X is a subset of S we denote by W_X the subgroup of W generated by X. It is well-known that for any subset X of S there exists precisely one element w of minimal length in aW_X and it has the property that every element $w' \in aW_X$ can be written in the form w' = wx with $x \in W_X$ and $\ell(w') = \ell(w) + \ell(x)$. Such an element w is called a X-reduced element. (Cf. [GrLie4-6], Ch. IV, Excercises §1, No 3).)

Let $W = W_g$ be the Weyl group and S be the set of simple reflections. If we take $X = S \setminus \{s_g\}$ then we find

$$W_X = \{ \sigma \in W_g : \sigma\{1, 2, \dots, g\} = \{1, 2, \dots, g\} \} \cong S_g$$

There is a natural partial order on W_g with respect W_X , the Bruhat-Chevalley order. It is defined in terms of Schubert cells $X(w_i)$ by:

$$w_1 \ge w_2 \iff X(w_1) \supseteq X(w_2).$$

Equivalently, if we define

(2.1)
$$r_w(i,j) := \#\{a \le i : w(a) \le j\}$$

then we have the combinatorial characterization

$$w_1 \le w_2 \iff r_{w_1}(i,j) \ge r_{w_2}(i,j) \quad \text{for all } 1 \le i,j \le 2g.$$

(Indeed it is easy to see that it is enough to check this for all $1 \leq i \leq g$ and $1 \leq j \leq 2g$.) Chevalley has shown that $w_1 \geq w_2$ if and only if any (hence every) X-reduced expression for w_1 contains a subexpression (obtained by just deleting elements) that is a reduced expression for w_2 ; here reduced means that w_2 is written as a product of $\ell(w_2)$ elements from S. Again, a reference for these facts is [BL00].

We now restrict to the following case. Let V be a symplectic vector space over \mathbb{Q} and consider the associated algebraic group $G = \operatorname{Sp}(V)$. If $E \subset V$ is a maximal isotropic subspace then the stabilizer of the flag $(0) \subset E \subset V$ is a parabolic subgroup conjugate to the standard parabolic corresponding to $H := S \setminus \{s_g\} \subset S$. Hence W_H will denote the subgroup of W_g generated by the elements of H and we will also use the notation P_H for the parabolic subgroup corresponding to W_H , i.e., the subgroup of the symplectic group stabilizing a maximal totally isotropic subgroup. As W_H consists of the permutations of W_g that stabilise the subsets $\{1, \ldots, g\}$ and $\{g+1, \ldots, 2g\}$ and the restriction of its action to $\{1, \ldots, g\}$ determine the full permutation we may identify W_H with S_g , the group of permutations of $\{1, \ldots, g\}$ and we shall do so without further mention. (This is of course compatible with the fact that H spans an A_{g-1} -subdiagram of the Dynkin diagram of G.) There are $2^g = |W_g|/|W_H|$ elements in W_g which are H-reduced elements. These 2^g elements will be called final elements of W_q . The Bruhat-Chevalley order between elements in W_g as well as the condition for being *H*-reduced can be conveniently expressed in terms of the concrete representation of elements of W_g as permutations in the following way.

Let A, B two finite subsets of $\{1, 2, ..., g\}$ of the same cardinality. We shall write $A \prec B$ if the *i*'th largest element of A is \leq the *i*'th largest element of B for all $1 \leq i \leq |A|$.

Lemma 2.1. *i)* If $w = [a_1a_2...a_g]$ and $w' = [b_1b_2...b_g]$ are two elements of W_g , then $w \leq w'$ in the Bruhat-Chevalley order precisely when for all $1 \leq d \leq g$ we have $\{a_1, a_2, ..., a_d\} \prec \{b_1, b_2, ..., b_d\}$.

ii) Let $w = [a_1a_2...a_g]$ and w^f be the final element of wW_H and let $w' = [b_1b_2...b_g]$ then $w^f \leq w'$ in the Bruhat-Chevalley order precisely when $\{a_1, a_2, ..., a_g\} \prec \{b_1, b_2, ..., b_g\}$.

iii) An element $\sigma \in W_g$ is H-reduced (or final) if and only if $\sigma(i) < \sigma(j)$ for all $1 \leq i < j \leq g$. Also, σ is H-reduced if and only if σ sends the first g - 1 simple roots into positive roots.

Proof. See for instance [BL00, p. 30].

2.2. Final Types and Young Diagrams. There are other descriptions of final elements that are sometimes equally useful. These involve maps of $\{1, 2, ..., 2g\}$ to $\{1, 2, ..., g\}$ and certain Young diagrams. We begin with the maps.

Definition 2.2. A final type (of degree g) is an increasing surjective map

$$\nu \colon \{0, 1, 2, \dots, 2g\} \to \{0, 1, 2, \dots, g\}$$

satisfying

$$\nu(2g-i) = \nu(i) - i + g \quad \text{for} \quad 0 \le i \le g$$

Note that we either have $\nu(i+1) = \nu(i)$ and then $\nu(2g-i) = \nu(2g-i-1) + 1$ or $\nu(i+1) = \nu(i) + 1$ and then $\nu(2g-i) = \nu(2g-i-1)$. A final type is determined by its values on $\{0, 1, \ldots, g\}$. There are 2^g final types of degree g corresponding to the vectors $(\nu(i+1) - \nu(i))_{i=0}^{g-1} \in \{0,1\}^g$. The notion of a final type was introduced by Oort (cf. [Oo01]).

To an element $w \in W_g$ we can associate the final type ν_w defined by

$$\nu_w(i) = i - r_w(g, i).$$

This is a final type because of the rule $r_w(g, 2g - i) - r_w(g, i) = g - i$ which follows by induction on *i* from the fact that w(2g + 1 - a) = 2g + 1 - w(a). It depends only on the coset wW_H of *w* since a permutation of the $a \leq g$ does not change the definition of $r_w(g, i) = \#\{a \leq g : w(a) \leq i\}$.

Conversely, to a final type ν we now associate the following element w_{ν} of the Weyl group, a permutation of $\{1, 2, \ldots, 2g\}$ as follows. Let

$$\beta = \{i_1, i_2, \dots, i_k\} = \{1 \le i \le g \colon \nu(i) = \nu(i-1)\}$$

with $i_1 < i_2 < \dots$ given in *increasing* order and let

$$\beta^c = \{j_1, j_2, \dots, j_{g-k}\}$$

be the elements of $\{1, 2, \ldots, g\}$ not in ξ , in *decreasing* order. We then define a permutation w_{ν} by mapping $1 \leq s \leq k$ to i_s and $k+1 \leq s \leq g$ to $2g+1-j_{s-k}$. The requirement that w_{ν} belong to W_g now completely specifies w_{ν} and by construction $w_{\nu}(i) < w_{\nu}(j)$ if $1 \leq i < j \leq g$ so it is a final element of W_g . It is clear from 2.1

that we get in this way all final elements of W_g . The Bruhat-Chevalley order for final elements can also be read off from the final type ν . We have $w \ge w'$ if and only if $\nu_w \ge \nu_{w'}$. This follows from 2.1, ii).

We summarize:

Lemma 2.3. By associating to a final type ν the element w_{ν} and to a final element $w \in W_g$ the final type ν_w we get an order preserving bijection between the set of 2^g final types and the set of final elements of W_g .

The final types are in bijection with certain Young diagrams: Our Young diagrams will be put in a position that is opposite to the usual positioning, i.e., larger rows will be below smaller ones and the rows will be lined up to the right. Furthermore, we shall make Young diagrams correspond to partitions by associating to one the parts that are the lengths of the rows of the diagram. We shall say that a Young diagram is *final of degree g* if its parts are $\leq g$ and no two parts are equal. They therefore correspond to subsets $\xi = \{g \geq \xi_1 > \xi_2 > \ldots > \xi_r\}$ of $\{1, 2, \ldots, g\}$.

To a final type ν we now associate the Young diagram Y_{ν} whose associated subset ξ is defined by

$$\xi_i = \#\{i \colon 1 \le i \le g, \, \nu(i) \le i - j\}.$$

A pictorial way of describing the Young diagram is by putting a stack of $i - \nu(i)$ squares in vertical position i for $1 \le i \le g$.

Example 2.4. This example corresponds to

 $\{\nu(i): i = 1, ..., g\} = \{1, 2, \dots, g - 5, g - 5, g - 4, g - 4, g - 3, g - 3\}$

and hence $\xi = \{5, 3, 1\}.$



The final elements w in W_g are in 1 - 1-correspondence with the elements of W_g/W_H . The group W_g acts on W_g/W_H by multiplication on the left, i.e., by the permutation representation. Therefore W_g also acts on the set of final types and the set of final Young diagrams. To describe this action we need the notion of a break point.

By a break point of a final type ν we mean an integer i with $1 \leq i \leq g$ such that either

- (1) $\nu(i-1) = \nu(i) \neq \nu(i+1)$, or
- (2) $\nu(i-1) \neq \nu(i) = \nu(i+1).$

If *i* is not a break point of $\nu = \nu_w$ then $\nu(i+1) = \nu(i-1)$ or $\nu(i+1) = \nu(i-1)+2$ and then $\nu_{s_iw} = \nu_w$. In particular, *g* is always a break point. The set of break points of ν equals

$$1 \le i \le g \colon \nu_{s_i w} \ne \nu_w \}.$$

Since $\nu = \nu_w$ determines a coset wW_H we have that *i* is not a break point of ν if and only if $w^{-1}s_iw \in W_H$, i.e., if and only if wW_H is a fixed point of s_i acting on W_q/W_H . The action of s_i on a final type ν is as follows: if *i* is not a break point then ν is fixed, otherwise replace the value of ν at i by $\nu(i) + 1$ if $\nu(i - 1) = \nu(i)$ and $\nu(i) - 1$ otherwise.

If w is a final element given by the permutation $[a_1, a_2, \ldots, a_g]$ then it defines a second final element, called the *complementary permutation*, defined by the permutation $[b_1, b_2, \ldots, b_g]$, where $b_1 < b_2 < \ldots < b_g$ are the elements of the complement $\{1, 2, \ldots, 2g\} \setminus \{a_1, \ldots, a_g\}$. If ξ is the partition defining the Young diagram of w then ξ^c defines the Young diagram of the complementary permutation. The set of break points of w and its complementary element are the same.

Lemma 2.5. Let w be a final element with associated final type ν and complementary element v.

i) We have that $v = \sigma_1 w \sigma_0 = w \sigma_1 \sigma_0$, where σ_0 (resp. σ_1) is the element of S_g (resp. W_g) that maps $1 \le i \le g$ to g + 1 - i resp. to 2g + 1 - i.

ii) If $1 \le i \le g$ has $\nu(i-1) \ne \nu(i)$ then $v^{-1}(i) = \nu(i)$ and if it has $\nu(i-1) = \nu(i)$ then $v^{-1}(i) = 2g + 1 - \nu(2g + 1 - i)$.

Proof. As w maps i to a_i we have that $2g + 1 - a_i$ is not among the a_j and hence $b_i = 2g + 1 - a_{g+1-i}$ (using that both the a_i and b_i are increasing sequences). This gives $\sigma_1 w \sigma_0(i) = \sigma_1(a_{g+1-i}) = 2g + 1 - a_{g+1-i} = b_i$ but we note that as $w \in W_g$ it commutes with σ_1 .

If $\nu(i-1) \neq \nu(i)$ and, say, $\nu(i) = i - k$ then we have k natural numbers $1 \leq i_1 < i_2 < \cdots < i_k < i$ such that $\nu(i_j - 1) = \nu(i_j)$. By the definition of v we then have $\nu(i-k) = (i-k) + k$ since the k values i_j $(j = 1, \ldots, k)$ are values for v, hence not of w. The second part is similar.

Remark 2.6. The permutations σ_1 and σ_0 of course have clear root theoretic relevance, they are respectively the longest element of W_g and S_g . Multiplication by them reverses the Bruhat-Chevalley order. Similarly it is clear that going from a final element to its complementary element also reverses the Bruhat-Chevalley order among the final elements and the first part of our statement says that that operation is obtained by multiplying by σ_1 and σ_0 , a fact that we presume generalises. Somewhat curiously our use of the complementary permutation seems unrelated to these facts.

In terms of Young diagrams the description is analogous and gives us a way to write the element w_{ν} as a reduced product of simple reflections. To each s_i we can associate an operator on final Young diagrams. If Y is a final diagram, s_i is defined on Y by adding or deleting a box in the *i*'th column if this gives a final diagram (only one of the two can give a final diagram) and then $s_i Y$ will be that new diagram; if neither adding nor deleting such a box gives a final Youg diagram we do nothing. In terms of the description as subsets ξ adding a box corresponds to $g+1-i \in \xi$ and $g+2-i \notin \xi$ and then $s_i \xi = (\xi \setminus \{g+1-i\}) \cup \{g+2-i\}$. It is then clear that for any final Young diagram Y there is a word $s_{i_1}s_{i_2}s_{i_3}\cdots s_{i_k}$ such that $Y = s_{i_1} s_{i_2} s_{i_3} \cdots s_{i_k} \emptyset$, where \emptyset denotes the empty Young diagram. Comparison with the action of s_i on final types and the correspondence between final types and Young diagrams shows that the action of s_i on diagrams is indeed obtained from that on final types. If we now have a word $t = s_{i_1} \dots s_{i_k}$ in the s_i we can make it act on Young diagram by letting each individual s_i act as specified. Note that this action only depends on the image of t in W_g but for the moment we want to consider the action by words. We define the *area* of a Young diagram Y to be the number of boxes it contains. We shall say that the word t is *building* if the area of $t\emptyset$ is equal to k, the length of the *word* (not the resulting element). This is equivalent to the action of s_{i_r} adding a box to $s_{i_{r+1}} \dots s_{i_k} \emptyset$ for all r.

Lemma 2.7. i) If ν is a final type and t is a word in the s_i such that $Y_{\nu}^c = t\emptyset$ then $w_{\nu} = w$, where w is the image of t in W_g and $\ell(w_{\nu}) = g(g+1)/2 - area(Y_{\nu})$.

ii) t is H-reduced if and only if t is building.

Proof. To prove i) we begin by noting that $t\emptyset$ only depends on the image of t in W_q so that i) is independent of the choice of t. Hence we may prove it by choosing a particular t using induction on the area of Y_{ν}^{c} . Note that $g(g+1)/2 - \operatorname{area}(Y_{\nu}) =$ $\operatorname{area}(Y_{\nu}^{c})$ so that the last part of i) says that $\ell(w_{\nu}) = \operatorname{area}(Y_{\nu}^{c})$. The final type ν with $\nu(i) = 0$ for $i \leq g$ corresponds to final diagram Y_{ν} with empty complementary diagram. We have $w_{\nu} = 1 \in W_q$, the empty product and it has length 0. This proves the base case of the induction. Suppose we proved the statement for diagram Y_{ν} with area $(Y_{\nu}^{c}) \leq a$. Adding one block to Y_{ν}^{c} to obtain $Y_{\nu'}^{c}$ means that for some i we have $g + 1 - i \in \xi^c$ and $g + 2 - i \notin \xi^c$, where ξ^c is the subset corresponding to Y_{ν}^{c} , the new subset is $(\xi')^{c} = (\xi \setminus \{g-i\}) \cup \{g-i+1\}$. This means that if i < gthere are $b < a \leq g$ such that $w_{\nu}(b) = i$ and $w_{\nu}(a) = 2g - i$ and $w_{\nu'}(b) = i + 1$ and $w_{\nu'}(a) = 2q + 1 - i$ and the rest of the integers between 1 and q mapped to the same elements. (The case when i = g is similar and left to the reader.) This makes it clear that we have $w_{\nu'} = s_i w_{\nu}$ so by the induction t maps to w_{ν} . It remains to show the formula for $\ell(w_{\nu})$. In the definition of $\ell(w)$ only the second term contributes as $w_{\nu}(i) < w_{\nu}(j)$ if $i < j \leq g$. Now, the only difference in the collections of sums w(i) + w(j) for $i \leq j$ and w equal to w_{ν} and $w_{\nu'}$ appears for (i, j) = (b, a) and we have $w_{\nu}(b) + w_{\nu}(a) = 2q$ and $w_{\nu'}(b) + w_{\nu'}(a) = 2q + 2$ so that the length of $w_{\nu'}$ is indeed one larger than that of $w_{\nu'}$.

As for ii), we have that $t\emptyset = Y_{\nu}^{c}$, where ν is the final type of w and then ii) is equivalent to t being H-reduced if and only $\operatorname{area}(Y_{\nu}^{c})$ is equal to the length of t. However by i) we know that $\operatorname{area}(Y_{\nu}^{c})$ is equal to $\ell(w_{\nu})$ and t is indeed H-reduced precisely when its length is equal to $\ell(w_{\nu})$.

Example 2.8. Consider again the Young diagram of the previous example but now for g = 5: We have $\xi = \{5, 3, 1\}$ and thus $\xi^c = \{2, 4\}$, so $w_{\nu} = [13579]$ and



 w_{ν} can be written as $s_4s_5s_2s_3s_4s_5$ (notice that permutations act from the left on diagrams).

We now characterize final types. Besides the function ν_w defined by

$$\nu(i) = i - \#\{a \le g \colon w(a) \le i\} = i - r_w(g, i)$$

and extended by $\nu(2g - i) = \nu(i) - i + g$ for $i = 0, \ldots, g$, we define a function $\mu = \mu_w$ on the integers $1 \le i \le 2g$ by

$$\mu(i) := \left(\max\{ w^{-1}(a) \colon 1 \le a \le i\} - g \right)^+,$$

where $(x)^+ := \max(x, 0)$. Alternatively, we have

$$\mu(i) = \min \{ 0 \le j \le g : r_w(g+j,i) = i \}.$$

Note that both μ and ν are non-decreasing functions taking values between 0 and g. Also ν can increase by 1 only and $\nu(2g) = \mu(2g) = g$. If w is final then ν_w is the final type associated to w. For an arbitrary w the function ν is the final type of the final element in the coset wS_q .

Lemma 2.9. We have $\mu_w(i) \ge \nu_w(i)$ for $1 \le i \le 2g$ with equality precisely when w is a final element and then ν_w is the final type of w.

Proof. We first prove the inequality $\mu \geq \nu$. Let $1 \leq i \leq g$. Suppose that $\mu(i) = m$, i.e., the maximal j with w(j) in [1, i] is g + m. Then there are at most m elements from [g + 1, 2g] which map into [1, i] and so there are at least i - m elements from [1, g] with their image under w in [1, i], so $i - \nu(i) \geq i - m$, in other words $\nu(i) \leq \mu(i)$. For i in the interval [g + 1, 2g] we consider $\nu(2g - i) = \#\{a \leq g : w(a) > i\}$. If $\mu(2g - i) = m$ then there are at least g - m elements from [1, g] mapping into [1, 2g - i] so $\nu(2g - i)$ is at most equal to m.

If w is final then w respects the order on [1, g] and this implies that if $\#\{a \leq g: w(a) \leq t\} = n$ then t - n elements from [g+1, 2g] map to [1, t], so the maximum element from [g+1, 2g] mapping into [1, t] is g + t - n. Hence $\mu(t) = t - n = \nu(t)$.

Conversely, if $\mu(i) = \nu(i)$ then this guarantees that w(i) < w(j) for all pairs $1 \le i < j \le g$.

We get as an immediate corollary.

Corollary 2.10. Let $w \in W_g$. We have that $r_w(g + \nu_w(i), i) = i$ for all $1 \le i \le g$ precisely when w is a final element.

Proof. The lemma says that if w is final then we have

 $\nu(i) = \mu(i) = \min \{ 0 \le j \le g : r_w(g+j,i) = i \}$

and in particular that $\nu(i) \in \{0 \le j \le g : r_w(g+j,i) = i\}$ which gives one direction.

Conversely, if we have $r_w(g + \nu_w(i), i) = i$, then $\nu_w(i) \ge \mu_w(i)$ and then the lemma gives that w is final.

2.3. Canonical Types. We now deal with an iterative way of constructing the function ν starting from its values on the endpoints and applying it repeatedly.

A final type ν is given by specifying $\nu(j)$ for $j = 1, \ldots, 2g$. But it suffices to specify the values of ν for the break points of ν . Under ν an interval $[i_1, i_2]$ between two consecutive break points of ν is mapped to an interval of length $i_2 - i_1$ or is mapped to one point. However, the image points $\nu(i_1)$ and $\nu(i_2)$ need not be break points of ν . Therefore we enlarge the set of break points to a larger set C_{ν} , called the *canonical domain*. We define C_{ν} to be the smallest subset of $\{0, 1, \ldots, 2g\}$ containing 0 and 2g such that if $j \in C_{\nu}$ then also $2g - j \in C_{\nu}$ and if $j \in C_{\nu}$ then $\nu(j) \in C_{\nu}$. It is obtained by starting from $R = \{0, 2g\}$ and adding the values $\nu(k)$ and $\nu(2g - k)$ for $k \in R$ and continuing till this stabilizes. The restriction of ν to C_{ν} is called a *canonical type*. We wish to see that the canonical domain C_{ν} contains the break points of ν and hence that we can retrieve ν from the canonical type of ν . To see this we need a technical lemma (its formulation is somewhat obscured by the fact that we also want to use it in another slightly different context). **Definition-Lemma 2.11.** We shall say that a subset $S \subseteq \{0, 1, \ldots, 2g\}$ is stable if it has the property that it contains 0 and is stable under $i \mapsto i^{\perp} := 2g - i$. For a stable subset S a map $f: S \to S \cap \{0, 1, \ldots, g\}$ is adapted to S if f(0) = 0and f(2g) = g, if it is contracting, i.e., it is increasing and $f(j) - f(i) \leq j - i$ for i < j and if it fulfills the following complementarity condition: For any two consecutive $i, j \in S$ (i.e., i < j and there are no $k \in S$ with i < k < j) we have $f(j) - f(i) = j - i \implies f(j^{\perp}) = f(i^{\perp})$.

i) If S is stable and f is a non-surjective function adapted to S then there is a proper subset $T \subset S$ such that $f_{|T}$ is adapted to T.

ii) If S is stable and f is a surjective function adapted to S then for any two consecutive $i, j \in S$ we have either f(i) = f(j) or f(j) - f(i) = j - i.

iii) We say that (S, f) is minimally stable if S is stable and f is adapted to S and furthermore there is no proper stable subset $T \subset S$ for which $f_{|T}$ is adapted to it, then the function $\nu : \{1, 2, \ldots, 2g\} \rightarrow \{1, 2, \ldots, g\}$ obtained from f by extending it linearly between any two consecutive $i, j \in S$ is a final type, $S = C_{\nu}$ and ν is the unique final extension of f. Conversely, if f is the canonical type of a final type ν then (C_{ν}, f) is minimally stable and in particular ν is the linear extension of its canonical type.

Proof. For i) consider $T = f(S) \cup (f(S))^{\perp}$. It is clearly stable under f and \perp and contains 0. If f is not surjective T is a proper subset of S.

Assume now that we are in situation of ii). We show that if $i < j \in S$ are consecutive then either f(j) - f(i) = j - i or f(i) = f(j) by descending induction on j - i.

By induction we are going to construct a sequence $i_k < j_k \in S$ k = 1, 2, ...of consecutive elements such that either $(i_{k-1}, j_{k-1}) = (j_k^{\perp}, i_k^{\perp})$ or $(f(i_k), f(j_k)) =$ (i_{k-1}, j_{k-1}) but not both $(i_{k-1}, j_{k-1}) = (j_k^{\perp}, i_k^{\perp})$ and $(i_{k-2}, j_{k-2}) = (j_{k-1}^{\perp}, i_{k-1}^{\perp})$ and in any case $j_k - i_k = j - i$. We start by putting $i_1 := i, j_1 := j$. Assume now that $i_k < j_k$ have been constructed. If we do not have $i_k, j_k \leq g$, then as $g \in S$, we must have $j_k^{\perp}, i_k^{\perp} \leq g$ and then we put $(i_{k+1}, j_{k+1}) = (j_k^{\perp}, i_k^{\perp})$. If we do have $i_k, j_k \leq g$ then by the surjectivity of f there are $i_{k+1}, j_{k+1} \in S$ such that $f(i_{k+1}) = i_k$ and $f(j_{k+1}) = j_k$. As f is increasing $i_{k+1} < j_{k+1}$ and by choosing i_{k+1} to be maximal and j_{k+1} to be minimal we may assume that they are neighbours. We must have that $j_{k+1} - i_{k+1} = j_1 - i_1$. Indeed, we have $f(j_{k+1}) - f(i_{k+1}) \leq j_k - i_k$ as f is contracting. If we have strict inequality we have $j - i = j_k - i_k < j_{k+1} - i_{k+1}$ and hence by the induction assumption we have either that $j_k - i_k = f(j_{k+1}) - f(i_{k+1}) = j_{k+1} - i_{k+1}$ which is a contradiction or $j_k = f(j_{k+1}) = f(i_{k+1}) = i_k$ which is also a contradiction. Hence we have $j_{k+1} - i_{k+1} = j_k - i_k = j - i$ and we have verified the required properties of $(i_{k+1}, j_{k+1}).$

There must now exist $1 \leq k < \ell$ such that $(i_k, j_k) = (i_\ell, j_\ell)$ and we pick k minimal for this property. If k = 1 we have either $j - i = j_{\ell-1} - i_{\ell-1} = f(j_\ell) - f(i_\ell) =$ f(j) - f(i) or $j - i = j_{\ell-2} - i_{\ell-2} = f(j_{\ell-1}) - f(i_{\ell-1}) = f(i^{\perp}) - f(j^{\perp})$ which implies that f(i) = f(j) by assumptions on f. We may hence assume that k > 1. We can not have both $(i_{k-1}, j_{k-1}) = (j_k^{\perp}, i_k^{\perp})$ and $(i_{\ell-1}, j_{\ell-1}) = (j_\ell^{\perp}, i_\ell^{\perp})$ as that would contradict the minimality of k. If $(i_{k-1}, j_{k-1}) = (j_k^{\perp}, i_k^{\perp})$ and $i_{\ell-1}, j_{\ell-1}) =$ $(f(i_\ell), f(j_\ell))$ then we get $j_{k-1} - i_{k-1} = j - i = j_{\ell-1} - i_{\ell-1} = f(i_{k-1}^{\perp}) - f(j_{k-1}^{\perp})$ which implies $f(j_{k-1}) = f(i_{k-1})$ which is either what we want in case k = 2or a contradiction. Similarly the case $(i_{\ell-1}, j_{\ell-1}) = (j_\ell^{\perp}, i_\ell^{\perp})$ and $i_{k-1}, j_{k-1}) =$ $(f(i_k), f(j_k))$ leads to a contradiction as does the case $i_{\ell-1}, j_{\ell-1} = (f(i_\ell), f(j_\ell))$ and $i_{k-1}, j_{k-1} = (f(i_k), f(j_k))$.

Finally, to prove iii) we note that by ii) for $i < j \in S$ consecutive we either have f(i) = f(j) or f(j) - f(i) = j - i. This means that the linear extension ν has the property that for $1 \leq i \leq 2g$ we have either $\nu(i) = \nu(i-1)$ or $\nu(i) = \nu(i-1)+1$ and if $\nu(i) = \nu(i-1)+1$ we get by the conditions on f that $\nu(2g-i+1) = \nu(2g-i)$. If for some $i \nu(i) = \nu(i-1)$ and $\nu(2g-i+1) = \nu(2g-i)$ we get that $g = f(2g) = \nu(2g) < g$ which is impossible by assumption and hence ν is indeed a final type. It is clear that S fulfills the defining property of C_{ν} so that $S = C_{\nu}$. The conditions on a final element implies that $\nu(j) - \nu(i) \leq j - i$ for j < i which implies that f has a unique final extension.

Conversely, if ν is a final type then C_{ν} clearly fulfills the required conditions and we have just noted that $\nu(j) - \nu(i) \leq j - i$ for j < i. The complementarity condition follows from the condition $\nu(i) = \nu(i-1) \iff \nu(2g-i+1) = \nu(2g-i) + 1$. \Box

We now give an interpretation of the canonical domain in terms of the Weyl group. Let $v \in W_g$ be a final element. A canonical fragment of v is a maximal interval $\{i < a \leq j\} =:]i, j] \subseteq \{1, 2, \ldots, 2g\}$ such that $v^k([i, j])$ remains an interval for all k.

Proposition 2.12. Let $v \in W_g$ be a final element, w its the complementary element, and ν the final type of w.

i) $\{1, 2, ..., 2g\}$ is the disjoint union of the canonical fragments of v and they are permuted by v.

ii) If [i, j] is a canonical fragment for v, and if $\nu(j) \neq \nu(j-1)$, then ν maps [i, j] bijectively to $|\nu(i), \nu(j)|$.

iii) If [i, j] is a canonical fragment, then so is $[\overline{j}, \overline{i}]$.

iv) The upper endpoints of the canonical fragments together with 0 comprise exactly the canonical domain for w.

Proof. If two canonical fragments I and J meet, their union K will be an interval and as $v^k(K) = v^k(I) \cup v^k(J)$, we see that $v^k(K)$ will be an interval for all k. By the maximality we get that I = J. On the other hand]i - 1, i] fulfills the stability condition so that i lies in a fragment. Hence $\{1, 2, \ldots, 2g\}$ is the disjoint union of fragments.

Let now R be the set of upper endpoints of fragments together with 0. As $\{1, 2, \ldots, 2g\}$ is the disjoint union of the fragments of v it follows that if [i, j] is a fragment, then i is also the upper end point of a fragment. Thus it follows from iii) that R is stable under $i \mapsto \overline{i}$. Let now i be an upper endpoint of a fragment. We want to show that $\nu(i) \in R$ and we may certainly assume that $\nu(i) \neq 0$ and we may also, by way of contradiction assume that i is a minimal upper endpoint for which $\nu(i)$ is not an upper endpoint. If $\nu(i) \neq \nu(i-1)$, then $v^{-1}(i) = \nu(i)$ and hence $\nu(i)$ is an upper endpoint of a fragment. Hence we may pick j < i such that $\nu(i) = \nu(i-1) = \cdots = \nu(j) \neq \nu(j-1)$. Then j can not belong to the same fragment as i and thus there must be an upper endpoint $j \leq k < i$. Then $\nu(k) = \nu(i)$ and by minimality of i we see that $\nu(k)$ is an upper endpoint which is a contradiction.

We therefore have shown that R contains 0 and is stable under $i \mapsto \overline{i}$ and ν . Hence it contains the canonical domain. Let now $j \in C_{\nu} \setminus \{0\}$ and let i be the largest $j \in C_{\nu}$ such that i < j. We now want show by induction on k that $v^{-k}(I)$, I := [i, j], remains an interval for all k and that also $v^{-k}(j)$ is one of its endpoints. Now, it follows from Lemma 2.5 that $C_{\nu} \setminus \{0\}$ is stable under v and hence $v^{-k}(j)$ will be the only element of C_{ν} in $v^{-k}(I)$. Under the induction assumption $v^{-k}(I)$ is an interval with $v^{-k}(j)$ as one of its endpoints and hence ν is constant on $v^{-k}(I)$ by Lemma 2.11. By Lemma 2.5 v^{-1} maps $v^{-k}(I)$ to interval with $v^{-k-1}(j)$ as one of its endpoints. This means that I is contained in a fragment and of the elements of I it is only j that can be contained in R. This means that there are no elements of R between consecutive elements of C_{ν} and hence $R \subseteq C_{\nu}$.

Corollary 2.13. By associating to a final type ν its canonical type, its Young diagram and the element w_{ν} we obtain a 1-1 correspondence between the following sets of cardinality 2^{g} : the set of canonical types, the set of final types, the set of final Young diagrams and the set of final elements of W_{q} .

2.4. Admissible Elements. The longest final element of W_g is the element

$$w_{\emptyset} := s_g s_{g-1} s_g s_{g-2} s_{g-1} s_g \dots s_g s_1 s_2 s_3 \dots s_g,$$

which as a permutation equals [g + 1, g + 2, ..., 2g]. Elements of W_g that satisfy $w \leq w_{\emptyset}$ are called *admissible*. We now characterize these.

Lemma 2.14. *i*) An element $w \in W_g$ fulfills $w \leq w_{\emptyset}$ if and only if $w(i) \leq g + i$ for all $1 \leq i \leq g$.

ii) The condition that $w \leq w_{\emptyset}$ is equivalent to $r_w(i, g+i) = i$ for all $1 \leq i \leq g$.

Proof. The first part follows immediately from the description of the Bruhat-Chevalley order (2.1) and the presentation of w_{\emptyset} .

For the second part one easily shows that $w(i) \leq g + i$ for all $1 \leq i \leq g$ is equivalent to $r_w(i, g + i) = i$ for all $1 \leq i \leq g$ which gives the first equivalence. \Box

Remark 2.15. The number of elements $w \in W_g$ with $w \leq w_{\emptyset}$ seems to be given by

$$\left(x\frac{d}{dx}\right)^g \left(\frac{1}{1-x}\right)_{\mid x=1/2}$$

This is the number of necklaces of sets of g labeled beads, see [S, Seq. A000629]. We leave this as a problem for the combinatorially inclined reader.

We give an illustration of the various notions for the case g = 2.

Example 2.16. g = 2. The Weyl group W_2 consists of 8 elements. We list the element, a reduced expression as a word (i.e. a decomposition $w = s_{i_1} \cdots c_{i_k}$ with $k = \ell(w)$), its length, the functions ν and μ and for final elements we also give the partition defining the Young diagram.

w	s	ℓ	ν	μ	Y
			6 3	6 3	
[4, 3, 2, 1]	$s_1 s_2 s_1 s_2$	4	$\{1, 2\}$	$\{2, 2\}$	
[4, 2, 3, 1]	$s_1 s_2 s_1$	3	$\{1, 1\}$	$\{2, 2\}$	
[3, 4, 1, 2]	$s_2 s_1 s_2$	3	$\{1, 2\}$	$\{1, 2\}$	Ø
[2, 4, 1, 3]	$s_{1}s_{2}$	2	$\{1, 1\}$	$\{1, 1\}$	$\{1\}$
[3, 1, 4, 2]	$s_{2}s_{1}$	2	$\{0, 1\}$	$\{0, 2\}$	
[2, 1, 4, 3]	s_1	1	$\{0, 0\}$	$\{0, 0\}$	
[1, 3, 2, 4]	s_2	1	$\{0, 1\}$	$\{0, 1\}$	$\{2\}$
[1, 2, 3, 4]	1	0	$\{0,0\}$	$\{0,0\}$	$\{1, 2\}$

The orbits of the complementary element will be play an important rôle in our discussion of the canonical flag. Here we introduce some definitions related to them.

Definition 2.17. Let $w \in W_g$ be a final and let v be its complementary element. Assume that S is an orbit of the action of v on its fragments. As v commutes with $i \mapsto \overline{i}$ we have that S is either invariant under $i \mapsto \overline{i}$ in which case we say that it is an *odd orbit* or that \overline{S} is another orbit in which case we say $\{S, \overline{S}\}$ is an *even orbit pair*.

2.5. Shuffles. Recall that a (p,q)-shuffle is a permutation σ of $\{1,2,\ldots,p+q\}$ for which $\sigma(i) < \sigma(j)$ whenever $i < j \leq p$ or p < i < j. It is clear that for each subset I of $\{1,2,\ldots,g\}$ there is a unique (|I|,g-|I|)-shuffle σ^I such that $I = \{\sigma^I(1), \sigma^I(2), \ldots, \sigma^I(|I|)\}$ and we will call it the shuffle associated to I. We will use the same notation for the corresponding element in W_g (i.e., fulfilling $\sigma^I(2g+1-i) = 2g+1-\sigma^I(i)$ for $1 \leq i \leq g$). By doing the shuffling from above instead of from below we get another shuffle σ_I given by $\sigma_I(i) = g+1-\sigma^I(g+1-i)$. We will use the same notation for its extension to W_g . Note that σ_I will shuffle the elements $\{g+1,g+2,\ldots,2g\}$ in the same way that σ^I shuffles $\{1,2,\ldots,g\}$, i.e., $\sigma^I(g+i) = g + \sigma_I(i)$, which is the relation with σ^I that motivates the definition. Note that if $I = \{i_1 < \cdots < i_r\}$ and if we assume that $i_r > r$ (if not then σ^I and σ_I are the identity elements) and we let k be the smallest index such that $i_k > k$, then $\sigma^I = s_{i_k-1}\sigma^{I'}$ and $\sigma_I = s_{g+1-(i_k-1)}\sigma_{I'}$, where $I' = \{i_1, \ldots, i_k - 1, \ldots, i_r\}$. We call the $s_{i-1}ws_{g+1-(i-1)}$ for $w \in W_g$ the *i*'th elementary shuffle of w, say that I' is the elementary reduction of I whose reduction index is i_k .

We define the *height* of a shuffle associated to a subset $\{i_1, i_2, \ldots, i_k\} \subseteq \{1, 2, \ldots, g\}$ to be $\sum_s (i_s - s)$. Using $w' = s_i w s_{g+1-t} \iff s_i w' s_{g+1-t} = w$, we see that starting with a shuffle w of a final element we arrive at a final element after ht(w) elementary shuffles.

Definition 2.18. Let Y be a final Young diagram of degree g. The shuffles of Y are the elements of W_g of the form $\sigma^I w_Y \sigma_I^{-1}$ for $I \subseteq \{1, 2, \ldots, g\}$.

If $w \leq w_{\emptyset}$ we say that $1 \leq i \leq g$ is a *semi-simple index* for w if w(i) = g+i (note that as $w \leq w_{\emptyset}$ we always have $w(i) \leq g+i$). The set of semi-simple indices will be called the *semi-simple index set* and its cardinality the *semi-simple rank*. We say that w is *semi-simply final* if the semi-simple index set has the form [g - f + 1, g] (where then f is the semi-simple rank). This is equivalent to w having the form $[\ldots, 2g - f + 1, 2g - f + 2, \ldots, 2g]$. If $w = w_Y$, Y a final Young diagram, then w is semi-simply final and the semi-simple rank is equal to g minus the length of the largest row of Y (defined to be zero if Y is empty).

Proposition 2.19. Let $w \leq w_{\emptyset}$ be a semi-simply final element of semi-simple rank f and let $I \subseteq \{1, 2, \ldots, 2g\}$ be a subset with #I = f. Put $\tilde{I} := \{g + 1 - i : i \in I\}$. Then $w' := \sigma^{I} w \sigma_{I}^{-1}$ is an element with $w' \leq w_{\emptyset}$ of semi-simple rank f and semi-simple index set \tilde{I} . Conversely, all $w' \leq w_{\emptyset}$ whose semi-simple index set is equal to I are of this form.

Proof. Put $j := \sigma_I^{-1}(i)$. Note that $j > g - f \iff i \in \tilde{I}$. If j > g - f we have w(j) = g + j and hence $\sigma^I w_Y \sigma_I^{-1}(i) = \sigma^I(g + j) = g + \sigma_I(j) = g + i$. If on the other hand $j \leq g - f$, then if $w_Y(j) \leq g$ there is nothing to prove, if not we may write $w_Y(j) = g + k$ and as the semi-simple rank of Y is f we have k < j. Then

 $\sigma^{I} w_{Y} \sigma_{I}^{-1}(i) = \sigma^{I}(g+j) = g + \sigma_{I}(k)$ and as $k < j \le g - f$ we have $\sigma_{I}(k) < \sigma_{I}(j) = i$ which gives $\sigma^{I} w_{Y} \sigma_{I}^{-1}(i) < g + i$.

The converse is easy and left to the reader.

Finally, we define the *a*-number for any element $w \in W_q$ by

$$a(w) := r_w(g,g)$$

If w is final with associated Young diagram Y then its a-number, also denoted a_Y , is the largest integer a with $0 \le a \le g$ such that Y contains the diagram with $\xi = \{a, a - 1, a - 2, ..., 1\}$.

3. The Flag Space

3.1. The Flag space of the Hodge bundle. In this section we introduce the flag space of a principally polarized abelian scheme over a base scheme of characteristic p. We use the Frobenius morphism to produce from a chosen flag on the de Rham cohomology a second flag, the position of which with respect to the first flag will be the object of study.

We let S be a scheme (or Deligne-Mumford stack) in characteristic p and let $\mathcal{X} \to S$ be an abelian variety over S with principal polarization (everything would go through using a polarization of degree prime to p but we shall stick to the principally polarized case). We consider the de Rham cohomology sheaf $\mathcal{H}_{dR}^1(\mathcal{X}/S)$. It is defined as the hyper-direct image $\mathcal{R}^1\pi_*(O_{\mathcal{X}} \to \Omega^1_{\mathcal{X}/S})$. It is a locally free sheaf of rank 2g on S. The polarization (locally in the étale topology given by a relatively ample line bundle on \mathcal{X}/S) provides us with a symmetric homomorphism $\rho: \mathcal{X} \to \hat{\mathcal{X}}$ and the Poincaré bundle defines a perfect pairing between $\mathcal{H}_{dR}^1(\mathcal{X}/S)$ and $\mathcal{H}_{dR}^1(\hat{\mathcal{X}}/S)$ and thus $\mathcal{H}_{dR}(\mathcal{X}/S)$ comes equipped with a non-degenerate alternating form (cf. [Oo95])

$$\langle , \rangle \colon \mathcal{H}^1_{dR}(\mathcal{X}/S) \times \mathcal{H}^1_{dR}(\mathcal{X}/S) \to O_S.$$

Moreover, we have an exact sequence of locally free sheaves on S

$$0 \to \pi_*(\Omega^1_{\mathcal{X}/S}) \to \mathcal{H}^1_{dR}(\mathcal{X}/S) \to R^1\pi_*O_{\mathcal{X}} \to 0.$$

We shall write \mathbb{H} for the sheaf $\mathcal{H}^1_{dR}(\mathcal{X}/S)$ and \mathbb{E} for the Hodge bundle $\pi_*(\Omega^1_{\mathcal{X}/S})$. We thus have an exact sequence

$$0 \to \mathbb{E} \to \mathbb{H} \to \mathbb{E}^{\vee} \to 0$$

of locally free sheaves on S. The relative Frobenius $F: \mathcal{X} \to \mathcal{X}^{(p)}$ and the Verschiebung $V: \mathcal{X}^{(p)} \to \mathcal{X}$ satisfy $F \cdot V = p \cdot \mathrm{id}_{\mathcal{X}^{(p)}}$ and $V \cdot F = p \cdot \mathrm{id}_{\mathcal{X}}$ and they induce maps, also denoted F resp.V, in cohomology:

$$F: \mathbb{H}^{(p)} \to \mathbb{H}$$
 and $V: \mathbb{H} \to \mathbb{H}^{(p)}$.

Of course, we have FV = 0 and VF = 0 and F and V are adjoints (with respect to the alternating form). This implies that $\operatorname{Im}(F) = \ker(V)$ and $\operatorname{Im}(V) = \ker(F)$ are maximally isotropic subbundles of \mathbb{H} and $\mathbb{H}^{(p)}$. Moreover, since dF = 0 on $\operatorname{Lie}(\mathcal{X})$ it follows that F = 0 on \mathbb{E} and thus $\operatorname{Im}(V) = \ker(F) = \mathbb{E}^{(p)}$. Verschiebung thus provides us with a bundle map (again denoted by V): $V \colon \mathbb{H} \to \mathbb{E}^{(p)}$.

16

Consider the space $\mathcal{F} = \operatorname{Flag}(\mathbb{H})$ of symplectic flags on the bundle \mathbb{H} consisting of flags of subbundles $\{\mathbb{E}_i\}_{i=1}^{2g}$ satisfying $\operatorname{rk}(\mathbb{E}_i) = i$, $\mathbb{E}_{g+i} = E_{g-i}^{\perp}$, and $\mathbb{E}_g = \mathbb{E}$. This space is a scheme over S and it is fibred by the spaces $\mathcal{F}^{(i)}$ of partial flags

$$\mathbb{E}_i \subsetneq \mathbb{E}_{i+1} \subsetneq \ldots \subsetneq \mathbb{E}_g$$

So $\mathcal{F} = \mathcal{F}^{(1)} = \operatorname{Flag}(\mathbb{H})$ and $\mathcal{F}^{(g)} = S$ and there are natural maps

$$\pi_{i,i+1}\colon \mathcal{F}^{(i)} \to \mathcal{F}^{(i+1)}$$

the fibres of which are Grassmann varieties of dimension *i*. So the relative dimension of \mathcal{F} is g(g-1)/2. The space $\mathcal{F}^{(i)}$ is equipped with a universal partial flag. On \mathcal{F} the Chern classes of the bundle \mathbb{E} decompose into their roots:

$$\lambda_i = \sigma_i(\ell_1, \dots, \ell_g)$$
 with $\ell_i = c_1(\mathbb{E}_i/\mathbb{E}_{i-1}),$

where σ_i is the *i*-th elementary symmetric function.

On $\mathcal{F}^{(i)}$ we have the Chern classes $\ell_{i+1}, \ldots, \ell_q$ and

$$\lambda_j(i) := c_j(\mathbb{E}_i) \qquad j = 0, 1, \dots, i.$$

Its Chow ring is generated over that of \mathcal{A}_g by the monomials $\ell_1^{m_1} \cdots \ell_g^{m_g}$ with $0 \leq m_j \leq j-1$. For later use we record the following Gysin formula.

Formula 3.1. We have $(\pi_{i,i+1})_* \ell_{i+1}^k = s_{k-i}(i+1)$, where $s_j(i+1)$ denotes the *j*th Segre class of \mathbb{E}_{i+1} (*j*th complete symmetric function in the Chern roots $\ell_1, \ldots, \ell_{i+1}$).

Given an arbitrary flag of subbundles

$$=\mathbb{E}_0 \subsetneq \mathbb{E}_1 \subsetneq \ldots \subsetneq \mathbb{E}_q = \mathbb{E}_q$$

with rank $(\mathbb{E}_i) = i$ we can extend this uniquely to a symplectic filtration on \mathbb{H} by putting

$$\mathbb{E}_{g+i} = (\mathbb{E}_{g-i})^{\perp}.$$

By base change we can transport this filtration to $\mathbb{H}^{(p)}$.

0

We introduce a second filtration by starting with the isotropic subbundle

$$\mathbb{D}_g := \ker(V) = V^{-1}(0) \subset \mathbb{H}$$

and continuing with

$$\mathbb{D}_{g+i} = V^{-1}(\mathbb{E}_i^{(p)}).$$

We extend it to a symplectic filtration by setting $\mathbb{D}_{g-i} = (\mathbb{D}_{g+i})^{\perp}$. We thus have two filtrations \mathbb{E}_{\bullet} and \mathbb{D}_{\bullet} on the pullback of \mathbb{H} to \mathcal{F} .

We shall use the following notation

$$\mathcal{L}_i = \mathbb{E}_i / \mathbb{E}_{i-1}$$
 and $\mathcal{M}_i = \mathbb{D}_i / \mathbb{D}_{i-1}$ for $1 \le i \le 2g$.

For ease of reference we formulate a lemma which follows immediately from definitions.

Lemma 3.2. We have $\mathcal{M}_{g+i} \cong \mathcal{L}_i^{(p)}$, $\mathcal{L}_{2g+1-i} \cong \mathcal{L}_i^{\vee}$ and $\mathcal{M}_{2g+1-i} \cong \mathcal{M}_i^{\vee}$.

We shall more generally for a family $X \to S$ of principally polarized abelian varieties say that a *Hodge flag* for the family is a complete symplectic flag $\{\mathbb{E}_i\}$ of \mathbb{H} for which \mathbb{E}_g is equal to the Hodge bundle. By construction this is the same thing as a section of $\mathcal{F}_g \to S$. We shall also call the associated flag $\{\mathbb{D}_i\}$ the *conjugate flag* of the Hodge flag. 3.2. The canonical flag of an abelian variety. In this section we shall confirm that the canonical filtration of X[p] (kernel of multiplication by p) by subgroup schemes of a principally polarized abelian variety X as defined by Ekedahl and Oort [Oo01] has its analogue for de Rham cohomology. Just as in [Oo01] we do this in a family $\mathcal{X} \to S$. It is the coarsest flag which is isotropic (i.e., if \mathbb{D} is a member of the flag then so is \mathbb{D}^{\perp}) and stable under F (i.e., if \mathbb{D} is a member of the flag then so is $F(\mathbb{D}^{(p)})$). The existence of such a minimal flag is proven by adding elements \mathcal{F}^{\perp} and $F(\mathbb{D}^{(p)})$ for \mathbb{D} already in the flag in a controlled fashion. We start by adding 0 to the flag. We then insist on three rules:

- (1) If we added $\mathbb{D} \subseteq \mathbb{D}_g$, then we immediately add \mathbb{D}^{\perp} (unless it is already in the flag constructed so far).
- (2) If we added $\mathbb{D}_g \subseteq \mathbb{D}$, then we immediately add $F(\mathbb{D}^{(p)})$ (unless it is already in the flag constructed so far).
- (3) If neither rule 1) nor rule 2) applies, then we add $F(\mathbb{D}^{(p)})$ for the largest element \mathbb{D} of the flag for which $F(\mathbb{D}^{(p)})$ is not already in the flag.

We should not however do this construction on S; we want to insure that we get a filtration by vector bundles: At each stage when we want to add the image $F(\mathbb{D}^{(p)})$, we have maps $F: \mathbb{D}^{(p)} \to \mathbb{H}$ of vector bundles and we then have a unique minimal decomposition of the base as a disjoint union of subschemes such that on each subscheme this map has constant rank. At the same time as we add $F(\mathbb{D}^{(p)})$ to the flag we replace the base by this disjoint union. On this disjoint union $F(\mathbb{D}^{(p)})$ then becomes a subbundle of \mathbb{H} and whether or not it is equal to the one of the previously defined subbundles is a locally constant condition. A simple induction then shows that we get a flag, i.e., for any two elements constructed one is included in the other, on a disjoint union of subschemes of S. As each element added is either the image under V of an element previously constructed or the orthogonal of such an element it is clear that this flag is the coarsening of any isotropic flag stable under V and it is equally clear that the decomposition of S is the coarsest possible decomposition. We shall call the (partial) flag obtained in this way the canonical flag of \mathcal{X}/S and the decomposition of S the canonical decomposition of the base.

To each stratum S' of the canonical decomposition of S we associate a canonical type as follows: Let $T \subseteq \{1, 2, \dots, 2q\}$ be the set of ranks of the elements of the canonical flag and let $f: T \to T \cap \{1, \dots, g\}$ be the function which to t associates $\operatorname{rk}(F(\mathbb{D}^{(p)}))$, where \mathbb{D} is the element of the canonical flag of rank t. We now claim that T and f fulfills the conditions of Lemma 2.11. Clearly T contains 0 and by construction it is invariant under $i \mapsto 2g - i$. Again by construction f is increasing and has f(0) = 0 and f(2g) = g. Furthermore, if $i, j \in T$ with i < j then F induces a surjective map $(\mathbb{D}/\mathbb{D}')^{(p)} \to F(\mathbb{D})/F(\mathbb{D}')$, where \mathbb{D} resp. \mathbb{D}' are the elements of the canonical flag for which the rank is j resp. i and hence f(j) - f(i) = $\operatorname{rk}(F(\mathbb{D})/F(\mathbb{D}')) \leq \operatorname{rk}(\mathbb{D}/\mathbb{D}') = j - i$. Finally, assume that f(j) - f(i) = j - i and let \mathbb{D} and \mathbb{D}' be as before. Putting $\mathbb{D}_1 := F(\mathbb{D}^{(p)})$ and $\mathbb{D}'_1 := F(\mathbb{D}'^{(p)})$ these are also elements of the canonical filtration and we by assumption F induces an isomorphism $F: (\mathbb{D}/\mathbb{D}')^{(p)} \to \mathbb{D}_1/\mathbb{D}'_1$. The fact that it is injective means that $\mathbb{D} \cap \ker F = \mathbb{D}' \cap \ker F$ which by taking annihilators, and using that ker F is its own annihilator, gives $\mathbb{D}^{\perp} + \ker F = \mathbb{D}' \perp + \ker F$ which implies $F(\mathbb{D}^{\perp}) = (\mathbb{D}'^{\perp})$ and hence that f(2q-i) = f(2q-j). Now, if f is not surjective then by Lemma 2.11 there is a proper subset of T fulfilling the conditions of the lemma. This is not possible as T by construction is a minimal subset with these conditions. Hence T and f fulfills the conditions of Lemma 2.11 and hence by it we get that (f,T) is a canonical type. Let ν be its associated final type. If $0 = \mathbb{D}_0 \subset \mathbb{D}_{i_1} \subset \cdots \subset \mathbb{D}_{2g} = \mathbb{H}$ is the canonical flag with $\operatorname{rk} \mathbb{D}_i = i$ we have also proved that F induces an isomorphism $(\mathbb{D}_j/\mathbb{D}_i)^{(p)} \to \mathbb{D}_{\nu(j)}/\mathbb{D}_{\nu(i)}$ which can be rephrased as an isomorphism

$$F: \mathbb{D}_{v(I)}^{(p)} \longrightarrow \mathbb{D}_{I}$$

where we have used the notation $\mathbb{D}_J := \mathbb{D}_i/\mathbb{D}_j$ for an interval J =]j, i] and $v \in W_g$ is the complementary element of (the final element of) ν . We shall say that ν (or more properly f) is the *canonical type* of the principally polarised abelian variety $\mathcal{X}_{S'} \to S'$. (We could consider the canonical type as a locally constant function on the canonical decomposition to the set of canonical (final) types.)

Remark 3.3. Note that the canonical flag is a flag containing \mathbb{D}_g and not \mathbb{E}_g . That will later mean that the canonical flag will be a coarsening of a conjugate flag which is derived from a Hodge flag. On the one hand this is to be expected. As the canonical flag is just that; it will be constructed in a canonical fashion from the family of principally polarized abelian varieties. Hence it is to be expected, and it is clearly true, that the canonical flag is horizontal with respect to the Gauss-Manin connection. On the other hand we do not want to just consider conjugate flags (or make constructions starting only with conjugate flags). The reason is essentially the same; as \mathbb{D}_g (or more generally the elements of the canonical flag) is horizontal it will not reflect first order deformations whereas \mathbb{E}_g isn't and does. This will turn out to be of crucial importance to us and is the reason why the Hodge flags will be the primary objects and the conjugate flags derived. On the other hand, when working pointwise, over an algebraically closed field say, we may recover the Hodge flag from the conjugate flag and then it is usually most convenient to work with the conjugate flag.

Example 3.4. Let X be an abelian variety with p-rank f and a(X) = 1 (equivalently, on G_g the operator V has rank g - 1 and semi-simple rank g - f). Then the canonical type is given by the numbers $\{\operatorname{rk}(C_i)\}$, i.e.,

$$\{0, f, f+1, \dots, 2g - f - 1, 2g - f, 2g\}$$

and ν is given by $\nu(f) = f$, $\nu(f+1) = f$, $\nu(f+2) = f+1, \ldots, \nu(g) = g-1, \ldots, \nu(2g-f-1) = g-1, \nu(2g-f) = g$ and $\nu(2g) = g$. The corresponding element $w \in W_g$ is $[f+1, g+1, \ldots, 2g-f-1, 2g-f+1, \ldots, 2g]$.

4. Strata on the Flag Space

4.1. The Stratification. The respective positions of two symplectic flags are encoded by a combinatorial datum, an element of a Weyl group. We shall now define strata on the flag space \mathcal{F} over the base S of a principally polarized abelian scheme $X \to S$ that mark the respective position of the two filtrations \mathbb{E}_{\bullet} and \mathbb{D}_{\bullet} that we have on the de Rham bundle over \mathcal{F} .

Intuitively, the stratum $\overline{\mathcal{U}}_w$ is defined as the locus of points x such that at x we have

$$\dim(\mathbb{E}_i \cap \mathbb{D}_j) \ge r_w(i,j) = \#\{a \le i \colon w(a) \le j\} \quad \text{for all} \quad 1 \le i, j \le 2g.$$

A more precise definition would be as degeneracy loci for some appropriate bundle maps. While this definition would work fine in our situation where we are dealing with flag spaces for the symplectic group it would not quite work when the symplectic group is replaced by the orthogonal group on an even-dimensional space (cf., [FP98]). With a view on future extensions of the ideas of this paper to other situations we therefore adopt the definition that would work in general. Hence assume that we have a semi-simple group G, a Borel group B of it, a G/B-bundle $T \to Y$ (with G as structure group) over some scheme Y, and two sections $s, t: Y \to T$ of it. Then for any element w of the Weyl group of G we define a (locally) closed subscheme \mathcal{U}_w resp. $\overline{\mathcal{U}}_w$ of Y in the following way. We choose locally (possibly in the étale topology) a trivialization of T for which t is a constant section. Then s corresponds to a map $Y \to G/B$ and we let \mathcal{U}_w (resp. $\overline{\mathcal{U}}_w$) be the inverse image of the B-orbit BwB (resp. of its closure). Another trivialization will differ by a map $Y \to B$ and as BwB and its closure are B-invariant these definitions are consistent and hence give global subschemes on Y. If s and t have the property that $X = \mathcal{U}_w$, then we shall say that s and t are in *relative position* w and if $Y = \overline{\mathcal{U}}_w$ we shall say that s and t are in *relative position* w.

Remark 4.1. The notation is somewhat misleading as it suggests that $\overline{\mathcal{U}}_w$ is the closure of \mathcal{U}_w which may not be the case in general. In the situation that we shall meet it will however be the case.

The situation to which we will apply this construction is when the base scheme is the space \mathcal{F} of symplectic flags \mathbb{E}_{\bullet} as above, s is the tautological section of the flag space of \mathbb{H} over \mathcal{F} , and t is the section given by the conjugate flag \mathbb{D}_{\bullet} . From now on we shall, unless otherwise mentioned, let \mathcal{U}_w and $\overline{\mathcal{U}}_w$ denote the subschemes of \mathcal{F} coming from the given s and t and $w \in W_g$. In this case it is actually often more convenient to use the language of flags rather than sections of G/B-bundles and we shall do so without further mention. We shall also say that a Hodge flag \mathbb{E}_{\bullet} is of stamp w resp. stamp $\leq w$ if \mathbb{E}_{\bullet} and its conjugate flag \mathbb{D}_{\bullet} are in relative position w resp. $\leq w$.

Lemma 4.2. Over \mathcal{U}_w we have an isomorphism $\mathcal{L}_i \cong \mathcal{M}_{w(i)}$ for all $1 \leq i \leq 2g$.

Proof. By the definition of the strata we have that the image of $\mathbb{E}_i \cap \mathbb{D}_{w(i)}$ has rank one greater than the ranks of $\mathbb{E}_{i-1} \cap \mathbb{D}_{w(i)}$, $\mathbb{E}_i \cap \mathbb{D}_{w(i)-1}$, and $\mathbb{E}_{i-1} \cap \mathbb{D}_{w(i)-1}$. So the maps $\mathbb{E}_i/\mathbb{E}_{i-1} \leftarrow \mathbb{E}_i \cap \mathbb{D}_{w(i)}/\mathbb{E}_{i-1} \cap \mathbb{D}_{w(i)-1} \rightarrow \mathbb{D}_{w(i)}/\mathbb{D}_{w(i)-1}$ give the isomorphism.

When the base of the principally polarized abelian scheme is \mathcal{A}_g we shall use the notation \mathcal{F}_g for the space of Hodge flags. Note that a Hodge flag with respect to $X \to S$ is the same thing as a lifting over $\mathcal{F}_g \to \mathcal{A}_g$ of the classifying map $S \to \mathcal{A}_g$. The conjugate flag as well as the strata \mathcal{U}_g and $\overline{\mathcal{U}}_g$ on S are then the pullbacks of the conjugate flag resp. the strata on \mathcal{F}_g .

4.2. Some Important Strata. We now give an interpretation for some of the most important strata. To begin with, if one thinks in terms instead of filtrations of X[p] by subgroup schemes it becomes clear that the condition $F(\mathbb{D}_i) \subseteq \mathbb{D}_i$ should be of interest. It can almost be characterized in terms of the strata $\overline{\mathcal{U}}_w$.

Proposition 4.3. Let $X \to S$ be a family of principally polarized abelian varieties and \mathbb{E}_{\bullet} a Hodge flag such that the flag is of stamp $\leq w$ and w is the smallest element with that property.

i) For $j \leq g$ we have that $r_w(i, g+j) = i$ precisely when $V(\mathbb{E}_i) \subseteq \mathbb{E}_i^{(p)}$.

ii) For $j \leq g$ we have that $r_w(g+j,i) = i$ implies that $F(\mathbb{D}_i^{(p)}) \subseteq \mathbb{D}_j$ and the converse is true if S is reduced.

iii) We have that $V(\mathbb{E}_i) \subseteq \mathbb{E}_i^{(p)}$ for all *i* precisely when $w \leq w_{\emptyset}$. If *S* is reduced $F(\mathbb{D}_i^{(p)}) \subseteq \mathbb{D}_i$ for all *i* precisely when $w \leq w_{\emptyset}$.

Proof. We have that $V(\mathbb{E}_i) \subseteq \mathbb{E}_j^{(p)}$ if and only if $\mathbb{E}_i \subseteq V^{-1}(\mathbb{E}_j^{(p)}) = \mathbb{D}_{g+j}$. On the other hand, by definition $\operatorname{rk} \mathbb{E}_i \cap \mathbb{D}_{g+j} \leq r_w(i, g+j)$ with equality for at least one point of S. As $\operatorname{rk} \mathbb{E}_i \cap \mathbb{D}_{g+j} = i \iff \mathbb{E}_i \subseteq \mathbb{D}_{g+j}$ we get the first part.

For the second part we start by claiming that $\mathbb{E}_{i}^{(p)} \subseteq \mathbb{D}_{j}$ is implied by $F(\mathbb{D}_{i}^{(p)}) \subseteq \mathbb{D}_{j}$. \mathbb{D}_{j} . Indeed, $F(\mathbb{D}_{i}^{(p)}) \subseteq \mathbb{D}_{j}$ is equivalent to $F(\mathbb{D}_{i}^{(p)})$ being orthogonal to \mathbb{D}_{2g-j} , i.e., to the condition that for $u \in \mathbb{D}_{i}^{(p)}$ and $v \in \mathbb{D}_{2g-j}$ we have $\langle Fu, v \rangle = 0$. This implies that $0 = \langle Fu, v \rangle = \langle u, Vv \rangle^{p}$ and hence $\langle u, Vv \rangle = 0$ as S is reduced which means that $\mathbb{D}_{i}^{(p)} \subseteq (V(\mathbb{D}_{2g-j}))^{\perp} = (\mathbb{E}_{g-j}^{(p)})^{\perp} = \mathbb{E}_{g+j}^{(p)}$. As S is reduced this implies that $\mathbb{D}_{i} \subseteq \mathbb{E}_{g+j}$ and this in turn is equivalent to $r_{w}(g+j,i) = i$. The argument can be reversed and then does not require S to be reduced.

Finally, we have from the first part that $V(\mathbb{E}_i) \subseteq \mathbb{E}_i^{(p)}$ for all $i \leq g$ precisely when $r_w(i, g + i) = i$ for all $i \leq g$ but by induction on i that is easily seen to be equivalent to $w(i) \leq g + i$ for all $i \leq g$ which by definition means that $w \leq w_{\emptyset}$. As $\mathbb{E}_g^{(p)} = V(\mathbb{H})$ the condition for i > g is trivially fulfilled.

The proof of the second equivalence is analogous in that using ii) the condition that $F(\mathbb{D}_i^{(p)}) \subseteq \mathbb{D}_i$ is equivalent to $r_w(g+i,i) = i$. In general, $r_u(i,j) = r_{u^{-1}}(j,i)$ so that this condition is equivalent to $r_{w^{-1}}(i,g+i) = i$ and hence by the same argument as before this condition for all i is equivalent to $w^{-1} \leq w_{\emptyset}$. Now, Chevalley's characterisation of the Bruhat-Chevalley order makes it clear that $u \leq v \iff u^{-1} \leq v^{-1}$ and hence get $w^{-1} \leq w_{\emptyset} \iff w \leq w_{\emptyset}^{-1}$. However, w_{\emptyset} is an involution.

Remark 4.4. i) As we shall see the strata $\overline{\mathcal{U}}_w$ in the universal case of \mathcal{F}_g are reduced. ii) Flags of stamp $w \leq w_{\emptyset}$ are called admissible.

We can also show that the relations between final and canonical types are reflected for flags. We say that a Hodge flag is a *final flag* if it is of stamp w for a final element w. Also if $I =]i, j] \subseteq \{1, 2, ..., 2g\}$ is an interval and \mathbb{F}_{\bullet} is a complete flag of a vector bundle of rank 2g then we define \mathbb{F}_I to be $\mathbb{F}_j/\mathbb{F}_i$.

Proposition 4.5. Let $X \to S$ be a principally polarized abelian scheme over S and \mathbb{E}_{\bullet} is a final flag for it of stamp w.

i) The conjugate flag \mathbb{D}_{\bullet} is a refinement of the canonical flag. In particular w is determined by $X \to S$. More directly, we have that the final type ν associated to w is given by

$$\operatorname{rk}(\mathbb{E}_q \cap \mathbb{D}_i) = i - \nu(i)$$

for all i. In particular the canonical decomposition of S with respect to $X \to S$ consists of a single stratum and its canonical type is the canonical type associated to w.

ii) Conversely, assume that S is reduced and that canonical decomposition of S consists of a single stratum, and let ν be the final type associated to the canonical type of the canonical flag. Then any Hodge flag \mathbb{E}_{\bullet} whose conjugate flag \mathbb{D}_{\bullet} is a refinement of the canonical flag and for which we have $F(\mathbb{D}_{i}^{(p)}) \subseteq \mathbb{D}_{\nu(i)}$ for all i, is a final flag.

ii) If I is a canonical fragment for v, the complementary element to w, then F induces a bijection $(\mathbb{D}_{v(I)})^{(p)} \cong \mathbb{D}_I$.

Proof. We start by showing that $F(\mathbb{D}_i) = \mathbb{D}_{\nu(i)}$ for all *i*. Indeed, this is equivalent to $F(\mathbb{D}_i) \subseteq \mathbb{D}_{\nu(i)}$ and $\operatorname{rk}((\ker(F) = \mathbb{E}_g^{(p)}) \cap \mathbb{D}_i^{(p)}) = i - \nu(i)$ as the second condition says that $F(\mathbb{D}_i)$ has rank $\nu(i)$. Now, the condition $F(\mathbb{D}_i) \subseteq \mathbb{D}_{\nu(i)}$ is by Proposition 4.3 implied by $r_w(g + \nu(i), i) = \nu(i)$ which is true for a final element by Corollary 2.10. On the other hand, the condition $\operatorname{rk}(\mathbb{E}_g^{(p)} \cap \mathbb{D}_i^{(p)}) = i - \nu(i)$ which is implied by $r_w(g, i) = \operatorname{rk}(\mathbb{E}_g \cap \mathbb{D}_i) = i - \nu(i)$ which is true by definition of ν . Now, the fact that $F(\mathbb{D}_i) = \mathbb{D}_j$ for all *i* and some *j* (depending on *i*) implies by induction on the steps of the construction of the canonical flag that \mathbb{D} is a refinement of the canonical flag. The rest of the first part then follows from what we have proved.

As for ii), assume first that \mathbb{E}_{\bullet} has a fixed stamp w' and let ν' be its final type. As \mathbb{D}_{\bullet} is an extension of the Hodge flag we get that when i is in the canonical domain of ν that $i - \nu'(i) = \operatorname{rk}(\mathbb{E}_g \cap \mathbb{D}_i) = i - \nu(i)$ so that ν and ν' coincide on the canonical domain of ν and hence they coincide by Lemma 2.11. The assumption that $F(\mathbb{D}_i^{(p)}) \subseteq \mathbb{D}_{\nu(i)} = \mathbb{D}_{\nu'(i)}$ for all i is by Proposition 4.3 equivalent to $r_w(g + \nu'(i), i) = i$ for all i and hence by Corollary 2.10 gives that w is final of type $\nu' = \nu$.

Finally, assume that I = [i, j]. The induced map $\mathbb{D}_{j}^{(p)}/\mathbb{D}_{i}^{(p)} \to \mathbb{D}_{\nu(j)}/\mathbb{D}_{\nu(i)}$ is always surjective but it follows from Lemma 2.11 that either the right hand side has dimension 0 or it has the same dimension as the right hand side. If they have the same dimension it induces an isomorphism $\mathbb{D}_{I}^{(p)} \longrightarrow \mathbb{D}_{v^{-1}(I)}$. If the right hand side has dimension zero, then again from Lemma 2.11 the two sides of $\mathbb{D}_{\overline{J}}^{(p)}/\mathbb{D}_{\overline{\tau}}^{(p)} \to \mathbb{D}_{\nu(\overline{J})}/\mathbb{D}_{\nu(\overline{\tau})}$ have the same dimension and hence this map is an isomorphism and again is an isomorphism $\mathbb{D}_{I}^{(p)} \longrightarrow \mathbb{D}_{v^{-1}(I)}$. As I is an arbitrary fragment we conclude. \square

The number of final extensions of a canonical flag will now be expressed in the familiar terms of the number of flags resp. self-dual flags in a vector space over a finite field (resp. a vector space with a unitary form). Hence we let $\gamma_n^e(m)$ be the number of complete \mathbb{F}_{p^m} -flags in $\mathbb{F}_{p^m}^n$ and let $\gamma_n^o(m)$ be the number of complete $\mathbb{F}_{p^{2m}}$ -flags self-dual under the unitary form $\langle (u_1, \ldots, u_n), (v_1, \ldots, v_n) \rangle := u_1 v_1^{p^m} + \cdots + u_n v_n^{p^m}$.

Lemma 4.6. Let X be a principally polarized abelian variety over an algebraically closed field and $w \in W_g$ the element whose canonical type is the canonical type of X. Put

$$\gamma(w) = \gamma_g(w) := \prod_{S=\overline{S}} \gamma^o_{\#I}(\#S/2) \prod_{\{S,\overline{S}\}} \gamma^e_{\#I}(\#S),$$

where the first product runs over the odd orbits and the second over the even orbit pairs and in both cases I is a typical member of S.

The number of final flags for X is then equal to $\gamma(w)$.

Proof. As we are over a perfect field, any symplectic flag extending \mathbb{D}_g is the conjugate flag of a unique Hodge flag. Hence we get from Proposition 4.5 that a final flag is the same thing as a flag \mathbb{D}_{\bullet} extending the canonical flag and for which $F(\mathbb{D}_i^{(p)}) \subseteq \mathbb{D}_{\nu(i)}$. The condition that $\dim(\mathbb{E}_g \cap \mathbb{D}_i) = i - v(i)$ then gives that we

actually have $F(\mathbb{D}_i^{(p)}) = \mathbb{D}_{\nu(i)}$. However, as \mathbb{D}_{\bullet} refines the canonical flag it is determined by the induced flags of the \mathbb{D}_I for v-fragments I and the stability condition $F(\mathbb{D}_i^{(p)}) = \mathbb{D}_{\nu(i)}$ transfers into the stability under the isomorphisms $F : \mathbb{D}_{\nu(I)}^{(p)} \longrightarrow \mathbb{D}_I$ of Subsection 3.2.

Hence the problem splits up into separate problems for each orbit under v and $i \mapsto \overline{\imath}$ (the map $i \mapsto \overline{\imath}$ transfers into the isomorphism $\mathbb{D}_{\overline{I}} \cong \mathbb{D}_{I}^{\vee}$ induced by the symplectic form). Given any fragment $I \in S$, S a v-orbit of fragments under v, the flag of $\mathbb{D}_{v^{k}(I)}$ for any k is then determined by the flag corresponding to I by the condition of F^{k} -stability takes the k'th Frobenius pullback of the flag on $\mathbb{D}_{v^{k}(I)}$ to the one of \mathbb{D}_{I} . Furthermore, the flag on \mathbb{D}_{I} then has to satisfy the consistency condition of being stable under $F_{S} := F^{\#S}$.

If now $\{S, \overline{S}\}$ is an even orbit pair the self-duality requirement for the flag means that the flags for the elements of \overline{S} are determined by those for the elements of Sand given $I \in S$ there is no other constraint on the flag on \mathbb{D}_I than the condition of stability under F_S . Hence we have the situation of a vector bundle \mathcal{D} over our base field \mathbf{k} and an isomorphism $F_S \colon \mathcal{D}^{(p^m)} \longrightarrow \mathcal{D}$, where m = #S, and we want to count the number of flags stable under F_S . Now, as \mathbf{k} is algebraically closed $\mathcal{D}_p := \{v \in \mathcal{D} : F_S(v) = v\}$ is an \mathbb{F}_{p^m} -vector space for which the inclusion map induces an isomorphism $\mathbf{k} \otimes \mathcal{D}_p \longrightarrow \mathcal{D}$. It then follows that F_S -stable flags correspond to \mathbb{F}_{p^m} -flags of \mathcal{D}_p .

If instead S is an odd orbit we will together with I also have \overline{I} in S and then, if #S = 2m, we will have that the flag on \mathbb{D}_I must be mapped to the dual flag on $\mathbb{D}_{\overline{I}}$ by $F_S := F^m$. The situation will be similar to the even orbit pair situation but with a "unitary twist" as in Proposition 7.2 and we get instead a correspondence with self-dual flags.

Example 4.7. For the canonical type associated to the final type of example 3.4 we have

$$\gamma_q(C_{\bullet}) = (p+1)(p^2 + p + 1) \cdots (p^{f-1} + p^{f-2} + \ldots + 1).$$

Definition 4.8. We let $w = u_f$ be the final element

$$u_f = s_g s_{g-1} s_g \cdots s_{g-f+1} \cdots s_g s_{g-f-1} \cdots s_g \cdots s_1 \cdots s_g,$$

i.e., if we introduce $\tau_j = s_j s_{j+1} \cdots s_g$ we have $u_f = \tau_g \tau_{g-1} \cdots \hat{\tau}_f \cdots \tau_1$. It corresponds to the Young diagram consisting of one row with g - f blocks and equals the element w given in 3.4.

Recall the notion of *a*-number a(X) for an abelian variety X of an (algebraically closed) field k of characteristic p, cf. [Oo01]. It equals the dimension over k of the vector space $\operatorname{Hom}(\alpha_p, X)$ of maps of the group scheme α_p to X. Equivalently, a(X) equals the dimension of the kernel of V on $H^0(X, \Omega^1_X)$. In our terms $H^0(X, \Omega^1_X) = \mathbb{E}_g$ and ker $V = \mathbb{D}_g$ so that $a(X) = \dim \mathbb{E}_g \cap \mathbb{D}_g$. The *p*-rank or *semi-simple rank* f on the other hand can be characterised by the condition that the $\dim_{\mathbf{k}} \cap_i V^i \mathbb{H}$ of V is equal to f.

Lemma 4.9. *i)* Let Y be a final Young diagram and $w \in W_g$ its final element and assume that $x = (X, \mathbb{E}_{\bullet}, \mathbb{D}_{\bullet}) \in \mathcal{U}_w(\mathbf{k})$, **k** a field. Then the p-rank of X equals the p-rank of Y.

ii) Let $w = u_f$ and $x = (X, \mathbb{E}_{\bullet}, \mathbb{D}_{\bullet}) \in \mathcal{F}_g(\mathbf{k})$. Then we have $x \in \mathcal{U}_w$ (resp. $\overline{\mathcal{U}}_w$) if and only if the filtration is V-stable and the p-rank of X is f and the a-number

of X is 1 (resp. the p-rank of X is $\leq f$). The image of \mathcal{U}_w in \mathcal{A}_g is the locus of abelian varieties of p-rank f and a-number 1.

Proof. By definition we have $r_w(i, g+i) = i$ for all $1 \le i \le g$ and $r_w(i, g+i-1) = i$ precisely when $i \le g - f$. Hence by Proposition 4.3 $V(\mathbb{E}_i) \subseteq \mathbb{E}_i^{(p)}$ for all $1 \le i \le g$, $V(\mathbb{E}_i) \subseteq \mathbb{E}_{i-1}^{(p)}$ for $i \le g - f$ and $V(\mathbb{E}_i) \subsetneq \mathbb{E}_{i-1}^{(p)}$ for i > g - f. The first and last conditions means that V induces an isomorphism $V \colon \mathbb{E}_{\{i\}} \longrightarrow \mathbb{E}_{\{i\}}$. On the other hand, the second condition gives $V^{g-f}(\mathbb{E}_{g-f}) = 0$. Together this gives that the semi-simple rank of X is f.

For the second part, since we have $w \leq w_{\emptyset}$ we must check the condition on the *p*-rank and the *a*-number. By the definition on \mathcal{U}_w we have $x \in \mathcal{U}_w$ if and only if $\operatorname{rk}(\mathbb{E}_g \cap \mathbb{D}_j) = 1$ for $f + 1 \leq j \leq g$ and 0 for $1 \leq j \leq f$. This implies that the kernel of $V (= \mathbb{D}_g \cap \mathbb{E}_g)$ has rank 1 and the semi-simple rank of V on \mathbb{E}_g is f. For $x \in \overline{\mathcal{U}}_w$ we get instead $\operatorname{rk}(\mathbb{E}_g \cap \mathbb{D}_j) \leq 1$ for $f + 1 \leq j \leq g$ and 0 for $1 \leq j \leq f$. \Box

Also the strata \mathcal{U}_w with $w \in S_g$ admit a relatively simple interpretation. Recall that an abelian variety is called *superspecial* if its *a*-number is equal to its dimension. This happens if and only if the abelian variety (without polarization) is isomorphic to a product of supersingular elliptic curves.

Lemma 4.10. Let x be a point of \mathcal{F}_g lying over $[X] \in \mathcal{A}_g$. The following are equivalent:

- (1) $x \in \bigcup_{w \in S_q} \mathcal{U}_w$.
- (2) $\dim(\mathbb{E}_g \cap \mathbb{D}_g) \ge g.$
- (3) $\ker(V) = \mathbb{E}_g$.
- (4) The underlying abelian variety X is superspecial.

Proof. It is well-known that X is superspecial if and only if V vanishes on \mathbb{E}_g . But this is equivalent to dim $(\mathbb{E}_g \cap \mathbb{D}_g) \geq g$. This explains the equivalences of (2), (3) and (4). Clearly, if $x \in \mathcal{U}_w$ with $w \in S_g$ then $r_w(g,g) = g$, hence (2) holds. Conversely, if X is superspecial then any filtration \mathbb{E}_{\bullet} on \mathbb{E}_g is V-stable and can be extended to a symplectic filtration. Since the degeneracy strata for $w \in S_g$, the Weyl group of GL_g , cover the flag space of flags on \mathbb{E}_g the lemma follows.

Lemma 4.11. Let x be a point of \mathcal{U}_w with underlying abelian variety X. Then the a-number of X equals a(w). Moreover, if $Y = \{1, 2, ..., a\}$ with corresponding final element $w_Y \in W_g$ then the image of \mathcal{U}_{w_Y} in \mathcal{A}_g is the locus T_a of abelian varieties with a-number a.

Proof. The *a*-number of an abelian variety is by definition the dimension of the kernel of V on $H^0(X, \Omega^1_X)$. But this is equal to $r_w(g,g) = a(w)$. The condition that a(X) = a implies that $r_w(g,g) = a$, hence $\nu(g) = g - a$. This implies that $\nu(g-a+i) \ge i$ for $i = 1, \ldots, a$. Therefore the 'smallest' ν satisfying these conditions is ν_{w_Y} .

4.3. Shuffling flags. Our first result on the stratification will concern the case when the p-rank is positive. All in all the étale and multiplicative parts of the kernel of multiplication by p on the abelian variety have very little effect on the space of flags on its de Rham cohomology. There is however one exception to this. The most natural thing to do is to put the multiplicative part at the bottom (and thus, by self-duality, the étale part at the top), which is what automatically happens

for a final filtration (on the conjugate filtration, that is). We may however start with a final filtration and then "move" the μ_p -factors up-wards. Note, that over a perfect field the kernel of multiplication by p is the direct sum of its multiplicative, local-local, and étale parts so that this is always possible. In general however it is possible only after a purely inseparable extension. This means that we get an inseparable map from a stratum where not all the μ_p -factors are at the bottom to a stratum where they all are. We intend to first give a combinatorial description of the strata that can be obtained in this way from a final stratum and then to compute the degree of the inseparable maps involved. However, as we have to compute an inseparable degree we should work with Hodge filtrations instead of conjugate filtrations as conjugate filtrations kill some infinitesimal information. This causes a slight conceptual problem as the V-simple parts in a final filtration are to be found "in the middle" rather than at the top and bottom (recall that V maps the top part of the conjugate filtration to the bottom of the Hodge filtration). This will not be a technical problem but the reader will probably be helped by keeping it in mind.

It turns out that the arguments used do not change if instead of considering shuffles of final elements we consider shuffles of semi-simply final elements. We shall treat the more general case as we shall need it later.

Hence we pick a subset $\tilde{I} \subseteq \{1, 2, \ldots, g\}$ and let $\overline{\mathcal{U}}_{\tilde{I}}^{ss}$ be the closed subscheme of \mathcal{F}_g defined by the conditions that V maps $\mathbb{E}_i^{(p)}$ to $\mathbb{E}_i^{(p)}$ for all $1 \leq i \leq g$ and to $\mathbb{E}_{i-1}^{(p)}$ for $i \notin \tilde{I}$. Hence $\overline{\mathcal{U}}_w \subseteq \overline{\mathcal{U}}_{\tilde{I}}^{ss}$ precisely when $w \leq w_{\emptyset}$ and the semi-simple index set of w is a subset of \tilde{I} . We also put

$$\mathcal{U}^{ss}_{\tilde{I}} := \overline{\mathcal{U}}^{ss}_{\tilde{I}} \setminus \cup_{\tilde{I}' \subset \tilde{I}} \overline{\mathcal{U}}^{ss}_{\tilde{I}'}$$

so that $\mathcal{U}_w \subseteq \mathcal{U}_{\tilde{I}}^{ss}$ precisely when $w \leq w_{\emptyset}$ and its semi-simple index set is equal to \tilde{I} . If $I := \{g + 1 - i : i \in \tilde{I}\}$ we get from Proposition 2.19 that these w are precisely those of the form $\sigma^I w' \sigma_I^{-1}$ for the semi-simply final w'.

We are now going construct, for every $I \subseteq \{1, 2, ..., g\}$, a morphism $S_I : \mathcal{U}_{\tilde{I}}^{ss} \to \mathcal{U}_{\{g-f+1,...,g\}}^{ss}$, where $\tilde{I} := \{g+1-i : i \in \tilde{I}\}$ and $\#\tilde{I} = f$. Let \tilde{i} be the reduction index of the elementary reduction I' of I and put i :=

Let \tilde{i} be the reduction index of the elementary reduction I' of I and put $i := g+1-\tilde{i}$. By Proposition 2.19 we have that $r_w(i+1,g+i) = i+1$ and w(i) = g+i. This means that if \mathbb{E}_{\bullet} is the (tautological) Hodge flag on $\mathcal{U}_{\tilde{I}}^{ss}$, then $V(\mathbb{E}_{i+1}) \subseteq \mathbb{E}_{i}^{(p)}$ and $V(\mathbb{E}_{i}) \subsetneq \mathbb{E}_{i-1}^{(p)}$ everywhere on $\mathcal{U}_{\tilde{I}}^{ss}$. This means that V gives a bijection on $\mathbb{E}_{\{i\}}$ and is zero on $\mathbb{E}_{\{i+1\}}$. Hence if Id denotes the map $s \mapsto 1 \otimes s$ from \mathcal{T} to $\mathcal{T}^{(p)}$ for the sheaves involved, the map induced by the quotient map

$$\operatorname{Ker}(\operatorname{Id} - V)_{\mathbb{E}_{\{i+1,i\}}} \to \operatorname{Ker}(\operatorname{Id} - V)_{\mathbb{E}_{\{i\}}}$$

where the kernel is computed in the étale topology on \mathcal{U}_w , is an isomorphism. Also for a sheaf \mathcal{T} with a linear isomorphism $V: \mathcal{T} \to \mathcal{T}^{(p)}$ we have an isomorphism $\operatorname{Ker}(V-I) \bigotimes \mathcal{O} \to \mathcal{T}$. It follows that the short exact sequence

$$0 \to \mathbb{E}_{\{i\}} \longrightarrow \mathbb{E}_{\{i+1,i\}} \longrightarrow \mathbb{E}_{\{i+1\}} \to 0$$

splits uniquely in a way compatible with V. This means that we may define a new flag where $\mathbb{E}'_j = \mathbb{E}_j$ for $j \neq i$ and $\mathbb{E}'_i/\mathbb{E}'_{i-1} = \mathbb{E}_{i+1}/\mathbb{E}_i$. Then the classifying map $\mathcal{U}_{\tilde{I}}^{ss} \to \mathcal{F}_g$ for this new flag will have its image in $\mathcal{U}_{\tilde{I}'}^{ss}$. Repeating this process we

end up with a flag whose classifying map will have its image in $\mathcal{U}_{\{g-f+1,\ldots,g\}}$ which by definition is our map S_I

Proposition 4.12. If $I \subseteq \{1, \ldots, g\}$ then the map $S_I : \mathcal{U}_{\tilde{I}}^{ss} \to \mathcal{U}_{\{g-f+1,\ldots,g\}}^{ss}$ is finite, radicial, and surjective.

Proof. To get from a point of $\mathcal{U}_{\{g-f+1,\ldots,g\}}^{ss}$ to one of $\mathcal{U}_{\overline{I}}^{ss}$ one has to find a *V*-invariant complement to some $\mathbb{E}_i/\mathbb{E}_{i-1}$ in $\mathbb{E}_{i+1}/\mathbb{E}_{i-1}$. As *V* will be zero on $\mathbb{E}_{i+1}/\mathbb{E}_i$ and bijective on $\mathbb{E}_i/\mathbb{E}_{i-1}$, a complement over the fraction field of a discrete valuation ring will extend to a complement over the discrete valuation ring (as the complement can not meet $\mathbb{E}_i/\mathbb{E}_{i-1}$ over the special fibre) so that the map is proper. It then remains to show that the map is a bijection over an algebraically closed field. In that case \mathbb{E}_g splits canonically as a sum of a *V*-nilpotent part and a *V*-semisimple part and the bijectivity is clear.

In order to determine the degree (necessarily of inseparability) we shall do the same factorisation as in the definition of S_I so that we may consider the situation of \tilde{I} with \tilde{i} and I' being the reduction index resp. elementary reduction of I. For the tautological flag \mathbb{E}_{\bullet} on $\mathcal{U}_{\tilde{I}}^{ss}$ we have that V is an isomorphism on $\mathbb{E}_{\{i\}}$ and zero on $\mathbb{E}_{\{i+1\}}$ while the opposite is true on $\mathcal{U}_{\tilde{I}'}^{ss}$

Lemma 4.13. The map $\mathcal{U}_{\tilde{i}}^{ss} \to \mathcal{U}_{\tilde{i}'}^{ss}$ is flat of degree p.

Proof. We consider the partial symplectic flag space $\mathcal{F}_g(i)$ consisting of the flags of \mathcal{F}_g by removing the *i*'th member \mathbb{D}_i and its annihilator. This means that we have a \mathbb{P}^1 -bundle $\mathcal{F}_g \to \mathcal{F}_g(i)$. Now, under this map $\mathcal{U}_{\overline{I}^s}^{ss}$ and $\mathcal{U}_{\overline{I}'}^{ss}$ map to the same subscheme $\mathcal{U} \subseteq \mathcal{F}_g(i)$ and the map $\mathcal{U}_{\overline{I}}^{ss} \to \mathcal{U}_{\overline{I}'}^{ss}$ is compatible with these projections. Over \mathcal{U} put $\mathcal{E} := \mathbb{E}_{\{i,i+1\}}$, $\mathcal{M} := \ker(V : \mathcal{E} \to \mathcal{E}^{(p)})$, and $\mathcal{L} := \operatorname{Im}(V : \mathcal{E} \to \mathcal{E}^{(p)})$. Then on the \mathbb{P}^1 -bundle $\pi : \mathcal{F}_g \to \mathcal{F}_g(i)$, the subscheme $\mathcal{U}_{\overline{I}'}^{ss}$ is defined by the vanishing of the composite $\mathcal{O}(-1) \to \pi^* \mathcal{E} \to \pi^*(\mathcal{E}/\mathcal{M})$ and in fact gives a section of \mathcal{F}_g over \mathcal{U} given by the sub-line bundle $\mathcal{M} \subset \mathcal{E}$. Hence it is enough to show that the projection map $\mathcal{U}_{\overline{I}}^{ss} \to \mathcal{F}_g(i)$ is flat of degree p. We have that $\mathcal{U}_{\overline{I}'}^{ss} \subseteq \mathcal{F}_g$ is defined by the vanishing of the composite $\mathcal{O}(-1)^{(p)} \to \pi^* \mathcal{E}^{(p)} \to \pi^*(\mathcal{E}^{(p)}/\mathcal{M})$. It is then enough to show that $\mathcal{U}_{\overline{I}'}^{ss} \subseteq \mathcal{F}_g$ is a relative Cartier divisor and for that is enough to show that is a proper subset with each fibre of $\mathcal{F}_g \to \mathcal{F}_g(i)$. This however is clear as for a geometric point of $\mathcal{F}_g(i)$ there just two points that lie in $\overline{\mathcal{U}}_{\emptyset}$, given by \mathcal{M} and $\mathcal{L}^{p^{-1}}$.

Composing these maps we get

Proposition 4.14. Let $I \subseteq \{1, \ldots, g\}$ and $\tilde{I} := \{g + 1 - i : i \in I\}$. Then the map $S_I : \mathcal{U}_I^{ss} \to \mathcal{U}_{\{1,\ldots,g\}}^{ss}$ is a finite purely inseparable map of degree $p^{\operatorname{ht}(I)}$.

Proof. The flatness and the degree of S_I follows by factoring it by maps as in Lemma 4.13 and noting that the number of maps is ht(I). The rest then follows from Proposition 4.12.

Remark 4.15. The result implies in particular that if w' is a shuffle of w by I, then $S_I: \mathcal{U}_{w'} \to \mathcal{U}_w$ is flat and purely inseparable of degree $p^{\operatorname{ht}(I)}$. We shall later (see Corollary 8.4) show that $\mathcal{U}_{w'}$ and \mathcal{U}_w are reduced. This shows that over the generic point of \mathcal{U}_w each simple shuffle towards w' really requires a finite inseparable extension of degree p. This is a kind of non-degeneracy statement which is the inseparable analogue of maximal monodromy (of which we shall also some examples later on). It can also be seen as saying that a certain Kodaira-Spencer map is injective.

4.4. The E-O strata on $\mathcal{A}_q \otimes \mathbb{F}_p$.

Definition 4.16. Let $w \in W_g$ be a final type. Then the E-O stratum \mathcal{V}_w associated to w is the closed subset of \mathcal{A}_g of points x for which the canonical type of the underlying variety is equal to the canonical type of w. We let $\overline{\mathcal{V}}_w$ be the closure of \mathcal{V}_w .

It is known that the dimension of \mathcal{V}_w is equal to dim(w), [Oo01]. This and the fact that the E-O strata form a stratification will also follow from our results in 8 and 9.3.

5. EXTENSION TO THE BOUNDARY

The moduli space \mathcal{A}_g admits several compactifications. The Satake or Baily-Borel compactification \mathcal{A}_g^* is in some sense minimal, cf. [FC90]. It is a stratified space

$$\mathcal{A}_q^* = \cup_{i=0}^g \mathcal{A}_i.$$

Chai and Faltings define in [F-C] a class of smooth toroidal compactifications. If $\tilde{\mathcal{A}}_g$ is such a toroidal compactification then there is a natural map $q \colon \tilde{\mathcal{A}}_g \to \mathcal{A}_g^*$. This induces a stratification of $\tilde{\mathcal{A}}_q$:

$$\tilde{\mathcal{A}}_g = \bigcup_{i=0}^g q^{-1}(\mathcal{A}_{g-i}) = \bigcup_{i=0}^g \mathcal{A}_g^{\langle i \rangle}.$$

The stratum $\mathcal{A}_{g}^{\langle i \rangle}$ parametrizes the semi-abelian varieties of torus rank *i*.

The Hodge bundle \mathbb{E} on \mathcal{A}_g can be extended to a rank g vector bundle, again denoted by \mathbb{E} , on $\tilde{\mathcal{A}}_g$. On $\mathcal{A}_g^{\langle i \rangle}$ the Hodge bundle fits into an exact sequence

$$0 \to \mathbb{E}' \to \mathbb{E} \to \mathbb{E}'' \to 0,$$

where the \mathbb{E}' is a rank g - i bundle and \mathbb{E}'' can be identified with cotangent bundle along the identity section of the toric part of the semi-abelian variety over $\mathcal{A}_{g}^{\langle i \rangle}$.

The bundle \mathbb{E}' is the pull back under $q: \mathcal{A}_g^{\langle i \rangle} \to \mathcal{A}_g^*$ of the Hodge bundle on \mathcal{A}_{g-i} . The Verschiebung V acts in a natural way on the extension \mathbb{E} and it preserves \mathbb{E}' . It induces an action on \mathbb{E}'' with trivial kernel because \mathbb{E}'' comes from the toric part and is generated by logarithmic forms.

The de Rham bundle \mathbb{H} on \mathcal{A}_g also admits an extension. In characteristic 0 this is the logarithmic de Rham sheaf $R^1\pi_*(\Omega^{\bullet}_{\tilde{\mathcal{X}}_g/\tilde{\mathcal{A}}_g}(\log))$, where the log refers to the acceptance of logarithmic singularities along the divisor at infinity, cf. [FC90, Thm. VI:1.1]. We have a short exact sequence

$$0 \to \mathbb{E} \to \mathbb{H} \to \mathbb{E}^{\vee} \to 0$$

extending the earlier mentioned sequence on \mathcal{A}_g .

We now want to compare the E-O strata on \mathcal{A}_g and \mathcal{A}_g and for this we introduce some notation. For a given integer $1 \leq i \leq g$ we can consider the Weyl group W_{g-i} as a subgroup of W_g by letting it act on the set $\{i + 1, i + 2, \ldots, g, \ldots, 2g - i\}$ via the bijection $j \longleftrightarrow i + j$ for $1 \le j \le g - i$. More precisely, define $\rho_i \colon W_{g-i} \to W_g$ via

$$\rho_i(w)(l) = \begin{cases} i+w(l) & \text{for } 1 \le l \le g-i \\ g+l & \text{for } g-i+1 \le l \le g. \end{cases}$$

This map respects the Bruhat-Chevalley order and final elements are mapped to final elements.

As symplectic flags on \mathbb{H} are determined by their restriction to \mathbb{E} and as we can extend \mathbb{E} to $\tilde{\mathcal{A}}_g$, we can extend \mathcal{F}_g to a flag bundle $\tilde{\mathcal{F}}_g$ on $\tilde{\mathcal{A}}_g$. Then we can also extend the degeneracy loci \mathcal{U}_w and $\overline{\mathcal{U}}_w$ to $\tilde{\mathcal{F}}_g$ and we shall use the same notation for these extensions.

Similarly, we can define the notion of a canonical filtration for a semi-abelian variety. If $1 \to T \to A \to A' \to 0$ is a semi-abelian variety with abelian part A' and toric part T of rank t and if the function ν' on $\{0, c_1, \ldots, c_r, c_{r+1}, \ldots, c_{2r} = 2 \dim(A')\}$ is the canonical type of A' then we define the canonical type of A to be the function ν on

$$\{0, t, t + c_1, \dots, t + c_r, t + c_{r+1}, \dots, t + c_{2r}, 2g - t, 2g\}$$

defined by $\nu(t + c_i) = t + \nu'(c_i)$. Using this definition we can extend the E-O stratification to $\tilde{\mathcal{A}}_q$.

The stratification $\mathcal{A}^{\langle i \rangle}$ of $\tilde{\mathcal{A}}_g$ induces a stratification $\tilde{\mathcal{F}}_g^{\langle i \rangle}$ by flag spaces $\mathcal{F}_g^{\langle i \rangle}$ over $\mathcal{A}_g^{\langle i \rangle}$. Recall that $\mathcal{A}_g^{\langle i \rangle}$ admits a map $q: \mathcal{A}_g^{\langle i \rangle} \to \mathcal{A}_{g-i}$ induced by the natural map $\tilde{\mathcal{A}}_g \to \mathcal{A}_g^*$. Similarly, we have a natural map $\pi_i = \pi: \mathcal{F}_g^{\langle i \rangle} \to \mathcal{F}_{g-i}$ given by restricting the filtration on \mathbb{E} to \mathbb{E}' .

We now describe the interplay between the two stratifications $(\mathcal{F}_g^{\langle i \rangle})_{i=1}^g$ and $(\mathcal{U}_w)_{w \in W_g}$.

Lemma 5.1. Let $w \in W_q$ be an element with $w \leq w_{\emptyset}$.

i) We have: $\mathcal{U}_w \cap \mathcal{F}_g^{\langle i \rangle} \neq \emptyset$ if and only if w is a shuffle of an element in $\rho_i(W_{g-i})$. ii) If $w = \rho_i(w')$ with associated degeneracy loci $\mathcal{U}_w \subset \mathcal{F}_g$ and $\mathcal{U}_{w'} \subset \mathcal{F}_{g-i}$ then we have $\mathcal{U}_w \cap \mathcal{F}_g^{\langle i \rangle} = \pi_i^{-1}(\mathcal{U}_{w'})$.

iii) In particular \mathcal{U}_w is the closure of its intersection with \mathcal{A}_q .

iv) At a point of $\tilde{\mathcal{A}}_g$ for which the torus part of the "universal" semi-abelian variety has rank r there is a smooth map from the formal completion of $\tilde{\mathcal{A}}_g$ at the point to the formal multiplicative group $\hat{\mathbb{G}}_m^r$ such that the locus where the torus rank of the universal semi-abelian variety is $s \leq r$ the inverse image of the locus of points of $\hat{\mathbb{A}}_m^r$ where r-s coordinates are 1. The restriction of this map to any $\overline{\mathcal{U}}_w$ is smooth.

Proof. A V-stable filtration on \mathbb{E} restricts to a V-stable filtration on \mathbb{E}' . If $\mathcal{U}_w \cap \mathcal{F}_g^{\langle i \rangle}$ is not empty then it determines a $w' \in W_{g-i}$ such that $\mathcal{U}_w \cap \mathcal{F}_g^{\langle i \rangle} \subseteq \pi_i^{-1}(\mathcal{U}_{w'})$. Since V is invertible on \mathbb{E}'' one sees that w is a shuffle of $\rho_i(w')$ and that $\mathcal{U}_w \cap \mathcal{F}_g^{\langle i \rangle} = \pi_i^{-1}(\mathcal{U}_{w'})$.

For the third part we note that the dimension of the intersection of \mathcal{U}_w with the boundary has dimension one lower than that of \mathcal{U}_w itself.

The last part is a direct consequence of the local construction of $\hat{\mathcal{A}}_g$ using toroidal compactifications and of the universal semi-abelian variety using Mumford's construction where it is defined by taking the quotient of a semi-abelian variety by

a subgroup of the torus part, the subgroup being generated by the coordinate functions of $\hat{\mathbb{G}}_m^r$ (see [FC90] for details). As \mathbb{H} and \mathbb{E} only depend on that fixed semi-abelian variety it is clear that the restriction of the map to a $\overline{\mathcal{U}}_w$ is smooth. \Box

Note also that this lemma is compatible with shuffling. It also results from the lemma that we can define the E-O stratification on the Satake compactification by either considering the closure of the strata \mathcal{V}_w on \mathcal{A}_g or by considering the images of the final strata \mathcal{V}_w on $\tilde{\mathcal{A}}_g$.

6. EXISTENCE OF BOUNDARY COMPONENTS

Our intent in this section is to show the existence of points in the smallest possible stratum $\overline{\mathcal{U}}_1$, the stratum associated to the identity element of W_q .

Proposition 6.1. Let X be an irreducible component of any $\overline{\mathcal{U}}_w$ in \mathcal{F}_g . Then X contains a point of $\overline{\mathcal{U}}_1$.

Proof. We prove this by induction over g and over the Bruhat-Chevalley order of w. The statement is clear for g = 1. We start off by choosing a Chai-Faltings compactification $\tilde{\mathcal{A}}_g$ of \mathcal{A}_g with a semi-abelian family over it (and a "principal" cubical structure so that we get a principal polarization on the semi-abelian variety modulo its toroidal part).

What we now actually want to prove is the same statement as in the proposition but for $\tilde{\mathcal{F}}_g$ instead. Since $\overline{\mathcal{U}}_1$ is contained in \mathcal{F}_g the result will follow. We start off by considering the intersection of X with the boundary $\tilde{\mathcal{F}}_g - \mathcal{F}_g$ to get Y. Assume that Y is non-empty and irreducible by replacing it by an irreducible component of Y if necessary. Then Y is contained in $\pi_1^{-1}(\overline{\mathcal{U}}_{w'})$ with $\rho_1(w') = w$ for some $w' \in W_{g-1}$. We claim that Y now is an irreducible component of $\pi_1^{-1}(\overline{\mathcal{U}}_{w'})$. This follows from the fact that "we can freely move the toroidal part into an abelian variety" which is Lemma 5.1.

By induction on g we can assume that $\overline{\mathcal{U}}_{w'}$ in \mathcal{F}_{g-1} contains $\overline{\mathcal{U}}_{1'}$, where 1' is the identity element of W_{g-1} . Any component Z of $\overline{\mathcal{U}}_{\rho_1(1')}$ which lies in X and meets Y does not lie completely in the boundary $\tilde{\mathcal{F}}_g - \mathcal{F}_g$. By induction on the Bruhat-Chevalley order we can assume that $w = \rho_1(1')$ and X = Z. Note also that for any w'' < w we have that $\overline{\mathcal{U}}_{w''}$ does not meet the boundary and by induction on the Bruhat-Chevalley order we get that $X = \mathcal{U}_w \cap X$. On the other hand if Xdoes not meet the boundary we immediately get the same conclusion.

Hence we may and shall assume that Y has the property that it lies completely inside \mathcal{U}_w and that it is proper. Lemma 6.2 now shows that it has an ample line bundle of finite order which together with properness forces Y to zero-dimensional. Now we always have that dim $Y \ge \ell(w)$ (the proof is analogous to [Ful, Them 14.3]) which gives $\ell(w) = 0$ and so w = 1 which clearly is a trivial case.

Lemma 6.2. Suppose that X is a proper irreducible component of $\overline{\mathcal{U}}_w$ inside \mathcal{F}_g such that $X \cap \mathcal{U}_w = X$. Then X is 0-dimensional.

Proof. This is a version of the Raynaud trick. We have the variety X and two symplectic flags \mathbb{E}_{\bullet} and \mathbb{D}_{\bullet} which at all points of X are in the same relative position w. It follows from 4.2 that we have an isomorphism between $\mathcal{L}_i := \mathbb{E}_i / \mathbb{E}_{i-1}$ and $\mathcal{M}_{w(i)} := \mathbb{D}_{w(i)} / \mathbb{D}_{w(i)-1}$ over X and then as we also have isomorphisms between \mathcal{L}_i^p and \mathcal{M}_{g+i} and \mathcal{L}_i and $\mathcal{L}_{2g+1-i}^{-1}$ we conclude that all the \mathcal{L}_i have finite order. On

the other hand we know that $\mathcal{L}_{2g} \bigotimes \mathcal{L}_{2g-1} \bigotimes \cdots \bigotimes \mathcal{L}_{g+1}$ is relatively ample and $\mathcal{L}_g \bigotimes \mathcal{L}_{g-1} \bigotimes \cdots \bigotimes \mathcal{L}_1$ is ample on the base \mathcal{A}_g we conclude. \Box

7. Superspecial fibres

We shall now discuss the fibre of $\mathcal{F}_g \to \mathcal{A}_g$ over superspecial points. The superspecial abelian varieties are characterized by the condition that $\mathbb{E}_g = \mathbb{D}_g$, i.e., the strata \mathcal{U}_w for which $w \in S_g$. Furthermore, V induces an isomorphism $\mathbb{E}/\mathbb{E}_g \Longrightarrow$ $\mathbb{E}_g^{(p)}$. On the other hand the polarization gives an isomorphism $(\mathbb{E}_g)^* \longrightarrow \mathbb{E}/\mathbb{E}_g$. This leads to the following definition.

Definition 7.1. i) Let S be a scheme in characteristic p. A p-unitary vector bundle is a vector bundle \mathcal{E} over S together with an isomorphism $F^*\mathcal{E} \longrightarrow \mathcal{E}^*$, where $F: S \to S$ is the (absolute) Frobenius map.

ii) Let \mathcal{E} be a *p*-unitary vector bundle over S and let $P \to S$ be the bundle of complete flags on \mathcal{E} . The *p*-unitary Schubert strata of P is the stratification given by letting $\mathcal{U}_w, w \in S_g$, consist of the points for which the universal flag \mathcal{F} and the dual of the Frobenius pullback $(F^*\mathcal{F})^*$ are in position corresponding to w.

A map $F^*\mathcal{E} \to \mathcal{E}^*$ of vector bundles is the same thing as a map $F^*\mathcal{E} \bigotimes_{\mathcal{O}_S} \mathcal{E} \to \mathcal{O}_S$, which in turns corresponds to a bi-additive map $\langle -, - \rangle \colon \mathcal{E} \times \mathcal{E} \to \mathcal{O}_S$ fulfilling $\langle fa, b \rangle = f^p \langle a, b \rangle$ and $\langle a, fb \rangle = f \langle a, b \rangle$. We shall normally use this latter description.

All *p*-unitary vector bundles are trivial in the étale topology as the following proposition shows.

Proposition 7.2. If $\langle -, - \rangle$ is a p-unitary structure on the vector bundle \mathcal{E} then

 $E := \{ a \in \mathcal{E} : \forall b \in \mathcal{E} : \langle b, a \rangle = \langle a, b \rangle^p \}$

is a local (in the étale topology) system of \mathbb{F}_{p^2} -vector spaces. Furthermore, $\langle -, - \rangle$ induces a unitary (wrt the involution $(-)^p$ on \mathbb{F}_{p^2}) structure on E. Conversely, if Eis a local system of \mathbb{F}_{p^2} -vector spaces with a unitary structure, then $\mathcal{E} := E \bigotimes_{\mathbb{F}_{p^2}} \mathcal{O}_S$ is a p-unitary vector bundle.

These two constructions establish an equivalence between the categories of punitary vector bundles and that of local systems of unitary \mathbb{F}_{p^2} -vector spaces. In particular all p-unitary vector bundles (of the same rank) are locally isomorphic in the étale topology.

Proof. The pairing $\langle -, - \rangle$ gives rise to an isomorphism $\psi' \colon \mathcal{E} \longrightarrow (F^*\mathcal{E})^*$ by $a \mapsto b \mapsto \langle b, a \rangle$ and an isomorphism $\psi \colon (F^2)^* \colon \mathcal{E} \longrightarrow (F^*\mathcal{E})^*$ by $a \mapsto b \mapsto \langle a, b \rangle^p$. The composite $\rho := \psi^{-1} \circ \psi'$ thus gives an isomorphism $\mathcal{E} \longrightarrow (F^2)^*\mathcal{E}$. Then E is simply the kernel of $\rho - 1 \otimes id$ and the fact that $\mathcal{E} = E \bigotimes_{\mathbb{F}_{p^2}} \mathcal{O}_S$ follows from [HW61]. The pairing $\langle -, - \rangle$ then induces a unitary pairing on E which is perfect as $\langle -, - \rangle$ is. Conversely, it is clear that a unitary pairing on E translates to one on \mathcal{E} .

Finally, as all (perfect) unitary mappings on \mathbb{F}_{p^2} -vector spaces of fixed dimension are isomorphic we get the local isomorphism.

This proposition has the following immediate corollary.

Corollary 7.3. The flag variety fibrations of two p-unitary vector bundles of the same rank on the same base are locally isomorphic by an isomorphism preserving the p-unitary Schubert strata.

Let us now consider the situation where the base scheme is $\operatorname{Spec}(\mathbb{F}_{p^2})$ and \mathcal{E} is a \mathbb{F}_{p^2} -vector space given a unitary perfect pairing. If we consider the smallest unitary Schubert stratum then it corresponds to flags that coincide with their unitary dual. Taking duals once more we see that they are taken to themselves after pullback by the square of the Frobenius, hence are defined over \mathbb{F}_{p^2} . Furthermore, they are selfdual with respect to the unitary pairing. This should come as no surprise as that stratum corresponds to final filtrations on superspecial abelian varieties. The next to lowest strata are somewhat more interesting.

Lemma 7.4. Let V be a g-dimensional \mathbb{F}_{p^2} unitary vector space and let \mathbb{P} be projective space based on V. If $s = (i, i + 1) \in S_g$ for some $1 \leq i < g$ then the closed Schubert stratum $\mathcal{U}_s \subseteq \mathbb{P}$ consists of the flags $0 = \mathbb{E}_0 \subset \mathbb{E}_1 \subset \cdots \subset \mathbb{E}_g$ where the \mathbb{E}_j for $j \neq i, g - i$ are \mathbb{F}_{p^2} -rational with $\mathbb{E}_j^{\perp} = \mathbb{E}_{g-j}, \mathbb{E}_{g-i} = (\mathbb{E}_i^{(p)})^{\perp}$ unless i = 2g, and $\mathbb{E}_i \neq (\mathbb{E}_{g-i}^{(p)})^{\perp}$

Proof. If \mathbb{E}_{\bullet} and \mathbb{D}_{\bullet} are two flags in $V \bigotimes R$ (R some \mathbb{F}_{p^2} -algebra) and they are in position s, then $\dim(\mathbb{E}_j \cap \mathbb{D}_j) = r_s(j, j) = j$ for $i \neq j$, i.e., $\mathbb{E}_j = \mathbb{D}_j$ and for i the conditions give us $\mathbb{E}_i \cap \mathbb{D}_i = \mathbb{E}_{i-1}$. In our case, where $\mathbb{D}_j = F^* \mathbb{E}_{g-j}^{\perp}$, this means $\mathbb{E}_j = F^* \mathbb{E}_{g-j}^{\perp}$ for $j \neq i$ and $\mathbb{E}_i \neq (\mathbb{E}_{g-i}^{(p)})^{\perp}$. If also $j \neq g - i$ we can use this twice and get that $\mathbb{E}_j = \mathbb{E}_j^{(p^2)}$, i.e., \mathbb{E}_j is \mathbb{F}_{p^2} -rational. \Box

As a result we get the following connectedness result, analogous to [Oo01, Prop. 7.3].

Theorem 7.5. Let V be a g-dimensional \mathbb{F}_{p^2} unitary vector space and let \mathbb{P} be projective space based on V. Let $S \subseteq \{1, \ldots, g-1\}$. Let $\overline{\mathcal{U}}$ be the union of the $\overline{\mathcal{U}}_{s_i}$ for $i \in S$. Then two flags $0 = A_0 \subset A_1 \subset \cdots \subset A_{g-1} \subset A_g$ and $0 = B_0 \subset B_1 \subset$ $\cdots \subset B_{g-1} \subset B_g$ in $\overline{\mathcal{U}}_1$ lie in the same component of $\overline{\mathcal{U}}$ precisely when $B_i = A_i$ for all $i \notin S$. Furthermore, every connected component of $\overline{\mathcal{U}}$ contains an element of $\overline{\mathcal{U}}_1$.

Proof. The last statement is clear as every irreducible component of any $\overline{\mathcal{U}}_{s_i}$ contains a point of $\overline{\mathcal{U}}_1$. This follows from (6.1) but can also easily be seen directly.

We start by describing looking at the locus $\overline{\mathcal{U}}_{F}^{i}$ of a $\overline{\mathcal{U}}_{s_{i}}$ with $1 \leq i < g$ of flags for which all the components of the flag except the dimension i and dimension g-iparts are equal to a fixed (partial) $\mathbb{F}_{p^{2}}$ -rational self-dual flag F_{\bullet} , the claims are easily proved using Lemma 7.4.

- i) For any $1 \leq i \leq (g-2)/2$ or $(g+2)/2 \leq i < g$ we get an element in $\overline{\mathcal{U}}_F^i$ by picking any $\mathbb{E}_{i-1} \subset \mathbb{E}_i \subset \mathbb{E}_{i+1}$ and then letting \mathbb{E}_{g-i} be determined by Lemma 7.4. Hence the locus is isomorphic to \mathbb{P}^1 and the intersection with $\overline{\mathcal{U}}_1$ consists of the points for which \mathbb{E}_i and \mathbb{E}_{g-i} are \mathbb{F}_{p^2} -rational.
- ii) When g is even we get an element in $\overline{\mathcal{U}}_F^i$ by picking $\mathbb{E}_{g/2-1} \subset \mathbb{E}_{g/2} \subset \mathbb{E}_{g/2+1}$. Hence the locus is isomorphic to \mathbb{P}^1 and the intersection with $\overline{\mathcal{U}}_1$ consists of the points for which $\mathbb{E}_{g/2}$ is \mathbb{F}_{p^2} -rational.
- iii) When g is odd we get an element in $\overline{\mathcal{U}}_{F}^{i}$ by picking $\mathbb{E}_{(g-3)/2} \subset \mathbb{E}_{(g-1)/2} \subset \mathbb{E}_{(g-1)/2} \subset \mathbb{E}_{(g-1)/2}$ for which $\overline{\mathbb{E}}_{(g-1)/2} \subset F^* \overline{\mathbb{E}}_{(g-1)/2}^{\perp}$, where $\overline{\mathbb{E}}_{(g-1)/2} = \mathbb{E}_{(g-1)/2}/\mathbb{E}_{(g-3)/2}$ and F^* comes from the \mathbb{F}_{p^2} -rational structure on $\mathbb{E}_{(g+3)/2}/\mathbb{E}_{(g-3)/2}$ and the scalar product is inducted from that on \mathbb{E}_g , and then define $\mathbb{E}_{(g+1)/2}$ by the

condition that $\mathbb{E}_{(g+1)/2}/\mathbb{E}_{(g-3)/2} = F^*\overline{\mathbb{E}}_{(g-1)/2}^{\perp}$. As all non-degenerate unitary forms are equivalent, choosing a basis of $\mathbb{E}_{(g+3)/2}/\mathbb{E}_{(g-3)/2}$ for which the form has the standard form $\langle (x, y, z), (x, y, z) \rangle = x^{p+1} + y^{p+1} + z^{p+1}$ and hence $\overline{\mathcal{U}}_F^i$ is isomorphic to the Fermat curve of degree p+1 and is hence is irreducible. The intersection with $\overline{\mathcal{U}}_1$ consists of the points for which $\mathbb{E}_{(g-1)/2}$ is \mathbb{F}_{p^2} -rational and then $\mathbb{E}_{(g+1)/2} = \mathbb{E}_{(g-1)/2}^{\perp}$.

iv) When g is odd we get an element in $\overline{\mathcal{U}}_{F}^{i}$ by picking $\mathbb{E}_{(g+3)/2}$ fulfilling conditions dual to those of iii). Hence again $\overline{\mathcal{U}}_{F}^{i}$ is irreducible and the intersection with $\overline{\mathcal{U}}_{1}$ consists of the points for which $\mathbb{E}_{(g+1)/2}$ is $\mathbb{F}_{p^{2}}$ -rational and then $\mathbb{E}_{(g-1)/2} = \mathbb{E}_{(g+1)/2}^{\perp}$.

It follows from this description that two flags in $\overline{\mathcal{U}}_1$ lie in the same component of $\overline{\mathcal{U}}$ if they are equivalent under the equivalence relation generated by the relations that for any unitary \mathbb{F}_{p^2} -flag $0 = A_0 \subset A_1 \subset \cdots \subset A_{g-1} \subset A_g$ we may replace it by any flag that is the same except for A_i and A_{g-i} for $i \in S$. The theorem then follows from Lemma 7.6.

Lemma 7.6. *i)* Let **k** be a field and $\mathcal{F}\ell_n$ the set of complete flags of vector spaces in a finite dimensional vector space. The equivalence relation generated by the operations of modifying a flag E_{\bullet} by, for any *i*, replacing E_i by any *i*-dimensional subspace of E_{i+1} containing E_{i-1} contains just one equivalence class.

ii) Let $\mathcal{F}\ell_n$ the set of complete flags of vector spaces in an n-dimensional \mathbb{F}_{p^2} -vector space, selfdual with respect to a perfect unitary pairing. An elementary modification of such a flag E_{\bullet} is obtained by either, for any $1 \leq i \leq (n-1)/2$, replacing E_i by any isotropic i-dimensional subspace of E_{i+1} containing E_{i-1} and E_{n-i} by its annihilator or, when n is even, replacing $E_{n/2}$ by any maximal totally isotropic subspace contained in $E_{n/2+1}$ and contains $E_{n/2-1}$. Then the equivalence relation generated by all elementary operations contains just one equivalence class.

Proof. Starting with i), we prove it by induction on n, the dimension of the vector space V. Given two flags E_{\bullet} and F_{\bullet} , if E_1 and F_1 are equal we may use induction applied to E_{\bullet}/E_1 and F_{\bullet}/E_1 . We now use induction on the smallest j such that $E_1 \subseteq F_j$, j = 1 already being taken care of. We now get new flag F'_{\bullet} by replacing F_{j-1} by $F_{j-2} \bigoplus E_1$, which works as $E_1 \subsetneq F_{j-1}$ and we then have $E_1 \subseteq F'_{j-1}$.

Continuing with ii) we again use induction on n and start with two selfdual flags E_{\bullet} and F_{\bullet} . Let us first assume that n is even, n = 2k. Then E_k and F_k are isotropic subspaces. If they have non-trivial intersection then we may pick a 1-dimensional subspace contained in it and then use i) to replace E_{\bullet} and F_{\bullet} by flags for which E_k and F_k are the same and $E_1 = F_1$. This implies that also $E_{n-1} = F_{n-1}$ and we may consider E_{n-1}/E_1 with its two flags induced from E_{\bullet} and F_{\bullet} and use induction to conclude. Assuming $F_k \cap E_k = \{0\}$ we may again use i) to modify F_{\bullet} , keeping F_k fixed, so that $E_1 \subseteq F_{k+1}$. This means that $E_1 \bigoplus F_{k-1}$ is totally isotropic and we may replace $F_{n/2}$ by it to obtain a new flag F'_{\bullet} for which $F'_{n/2}$ and $E_{n/2}$ intersect non-trivially.

When n is odd, n = 2k + 1, we may again use induction on n to finish if E_k and F_k intersect non-trivially. If not we may again use i) to reduce to the case when $E_1 \subseteq F_{k+2}$ and then we may replace F_k by $E_1 \bigoplus F_{k-1}$ and F_{k+1} by its annihilator.

8. LOCAL STRUCTURE OF STRATA

8.1. Stratified Spaces. We now want to show that our moduli space locally looks like the space of complete symplectic flags (in 2g-dimensional space). This isomorphism will on the one hand preserve the degeneration strata and on the other hand will be compatible with all the forgetful maps to partial flag spaces (all the way down to that of totally isotropic g-dimensional spaces). In order to have a convenient way of formulating such a result we introduce the two following notions:

By a *stratified space* we will mean a scheme together with a collection of closed subschemes, called *strata*. A map between stratified spaces is said to be stratified if it maps strata into strata.

If P is a partially ordered set then a diagram of spaces over P associates to each element p of P a scheme X_p and to each relation p > p' a map $X_p \to X_{p'}$ fulfilling the condition that the composite $X_p \to X_{p'} \to X_{p''}$ equals the map $X_p \to X_{p''}$ for any p > p' > p''. We shall also similarly speak about a diagram of stratified spaces where both the schemes and the maps are assumed to be stratified. Given a field k and a k-point x of a diagram X. we may speak of its (strict) Henselization at x, which at each $p \in S$ is the Henselization at x of X_p .

For a positive integer g we now consider the partially ordered set P_g whose elements are the subsets of $\{1, 2, \ldots, g-1\}$ and with ordering that of inclusion. We have two diagrams of stratified spaces over this set: The first, $\mathcal{F}\ell_g$, associates to the subset S the flag space of a maximal totally isotropic subspace E of a symplectic 2g-dimensional vector space and partial flags of subspaces of E whose dimensions form the set S. The map associated to an inclusion $S \subset S'$ is simply the map forgetting some of the elements of the flag. Similarly, we let \mathcal{F}_g be the diagram that to a subset S associates the space of flags over the moduli space \mathcal{A}_g of principally polarized abelian varieties that associates to an abelian variety the space of flags on the Hodge bundle whose dimensions form the subset S.

The diagram $\mathcal{F}\ell_g$ becomes a stratified diagram by considering the stratifications given by the (closed) Schubert cells with respect to some fixed complete flag. In positive characteristic p the diagram \mathcal{F}_g becomes a stratified diagram by considering the degeneracy loci given by the relative positions of the Hodge flag \mathbb{E} . and the conjugate flag \mathbb{D} .

8.2. Height 1-Maps. For schemes in a fixed positive characteristic p we shall say that a closed immersion $S \hookrightarrow S'$ defined by the ideal sheaf \mathcal{I} is a *height* 1-map if $\mathcal{I}_{S}^{(p)} = 0$, where for an ideal I, we let $I^{(p)}$ be the ideal generated by the p'th powers of elements of I. If R is a local ring in characteristic p, the *height* 1-*hull* of R is the quotient ring $R/m_R^{(p)}$ (so that the spectrum of it is the largest closed subscheme of Spec R for which the map from Spec R/m_R to Spec $R/m_R^{(p)}$ is a height 1-map). If k is a field of characteristic p and x: Spec $k \to S$ a k-map to a k-scheme S of characteristic p, then by the *height* 1-*neighbourhood* of x we will mean the spectrum of the height 1-hull of the local ring of S at x. It is clear that taking height 1-neighbourhoods of k-points is functorial under maps between k-pointed kschemes. Finally, we shall say that two local rings are *height* 1-*isomorphic* if their respective height 1-hulls are isomorphic and that the height 1-hull of a k-point is *height* 1-*smooth* if its is isomorphic to the height 1-hull of a smooth k-point (i.e., is of the form $k[[t_1, \ldots, t_n]]/m^{(p)})$. **Theorem 8.1.** For each perfect field k of positive characteristic p and each kpoint x of \mathcal{F}_g there is a k-point y of $\mathcal{F}\ell_g$ such that the height 1-neighbourhood of x is isomorphic to the height 1-neighbourhood of y by a stratified isomorphism of diagrams.

Proof. Denote also by x the point of \mathcal{F}_{g} , the space of complete flags of the Hodge bundle, associated to x as a point of the diagram \mathcal{F}_q^{\cdot} . Let X be the height 1neighbourhood of x in \mathcal{F}_{g} and X the height 1-neighbourhood of x in \mathcal{F}_{g} . Now the ideal of the closed point of x in X has a divided power structure for which all the divided powers of order $\geq p$ are zero. This allows us get a trivialization of the restriction of the de Rham cohomology $\mathbb{H}_{|X} \longrightarrow X \times W$ which is horizontal (i.e., compatible with the Gauss-Manin connection on the left and the trivial connection on the right). Now, as the absolute Frobenius map on X factors through the closed point we get that the pullback $\mathbb{E}^{(p)}_{\bullet}$ is a horizontal flag and then so is \mathbb{D}_{\bullet} , its elements being either inverse images of horizontal subbundles by the horizontal map V or duals of horizontal subbundles. We now get a map from X to the space $\mathcal{F}\ell_q$ of complete symplectic flags on W such that the pullback of the universal flag equals \mathbb{E} .. We may furthermore, choose a symplectic isomorphism of W and the standard symplectic space such that \mathbb{D}_{\bullet} is taken to the fixed complete flag. We can extend this map in a compatible fashion for all partial flag spaces giving a map from the diagram X. to $\mathcal{F}\ell_q^i$ and we will denote by y the k-point which is the composite of x and this map. This map is clearly a stratified map and by the infinitesimal Torelli theorem it induces an isomorphism from X to Y, the first height 1-neighbourhood of y in $\mathcal{F}\ell_{q}^{\cdot}$.

Corollary 8.2. For each perfect field k of positive characteristic p and each k-point x of \mathcal{F}_g there is a k-point y of \mathcal{F}_g such that the Henselization of x is isomorphic to the Henselization of y by a stratified isomorphism.

Proof. The theorem provides such an isomorphism over the height 1-hull X of x. Now, over $\mathcal{O}_{\mathcal{F}_g,x}$ we may extend the trivialization of $\mathbb{H}_{|X}$ to a trivialization of $\mathbb{H}_{\mathcal{F}_g,x}$ which also extends the trivialization of \mathbb{D} (making, of course, no requirements of horizontality). This gives a map from the localization, \tilde{X} , of \mathcal{F}_g at x to $\mathcal{F}\ell_g$ that extends the map from X to $\mathcal{F}\ell_g$. It thus induces a map from \tilde{X} to \tilde{Y} , the localization $\mathcal{F}\ell_g$ at y. Now, this map induces an isomorphism on tangent spaces and \mathcal{F}_g is smooth. This implies that we get an induced isomorphism on Henselizations and proves the theorem.

Lemma 8.3. Let A be a principally polarised abelian variety over an algebraically closed field. If a flag \mathbb{D}_{\bullet} for it has type w' which \leq its canonical type then w' is the final element corresponding to the canonical type of A.

Proof. The flag \mathbb{D}_{\bullet} has the property, as it is of a type \leq to the canonical type, that F maps $\mathbb{D}_{i}^{(p)}$ into $\mathbb{D}_{\nu_{w}(i)}$. Consider now the set I of i's for which \mathbb{D}_{i} is a member of the canonical flag. It clearly contains 0 and is closed under $i \mapsto \overline{i}$. Furthermore, if $i \in I$, then $F(\mathbb{D}_{i}^{(p)})$ has dimension $\nu_{w}(i)$ but is then equal to $\mathbb{D}_{\nu_{w}(i)}$ as it is contained in it. Hence I fulfills the conditions of Corollary 2.10 and hence contains the canonical domain which means that \mathbb{D}_{\bullet} is a refinement of the canonical flag and hence ν , the final type of A, and $\nu_{w'}$ coincide on the canonical domain of ν and

hence are equal by Corollary 2.10 which means that w' is the final element of the canonical type.

Corollary 8.4. *i)* Each stratum \mathcal{U}_w is smooth of dimension $\ell(w)$.

ii) The closed stratum $\overline{\mathcal{U}}_w$ is Cohen-Macaulay, reduced, and normal of dimension $\ell(w)$ and $\overline{\mathcal{U}}_w$ is the closure of \mathcal{U}_w for all $w \in W_q$.

iii) If w is final then the restriction of the projection $\mathcal{F}_g \to \mathcal{A}_g$ to \mathcal{U}_w is a finite surjective étale covering from \mathcal{U}_w to \mathcal{V}_w of degree $\gamma(w)$.

Proof. We know that each open Schubert cell of $\mathcal{F}\ell_g$ is smooth and each closed is Cohen-Macaulay by a proof completely along the lines of [Ful, Thm 14.3] and by a theorem of Chevalley (cf., [Ch94, Cor. of Prop 3]) they are smooth in codimension 1 so by Serre's criterion they are normal and reduced. The same statement for the stratification of \mathcal{F}_g then follows form the theorem. Finishing ii), the fact that $\overline{\mathcal{U}}_w = \overline{\mathcal{U}}_w$ follows more or less formally from the rest: If $x \in \overline{\mathcal{U}}_w$ then we know that the dimensions of all $\mathcal{U}_{w'}$ with w' < w that pass through are $\ell(w') < \ell(w)$ but the dimension of $\overline{\mathcal{U}}_w$ at that point is $\ell(w)$ and hence x must lie in the closure of \mathcal{U}_w .

As for iii), that \mathcal{U}_w maps into \mathcal{V}_w follows from the fact that the restriction of a final filtration to its canonical domain is a canonical filtration (Proposition 4.5). That the map $\mathcal{U}_w \to \mathcal{A}_g$ is unramified follows from the same statement for Schubert cells which is [BGG73, Prop. 5.1]. We next prove that $\mathcal{U}_w \to \mathcal{V}_w$ is proper. Note that by Proposition 4.5 and by the fact that by definition \mathcal{V}_w is the image of \mathcal{U}_w we get that the geometric points of \mathcal{V}_w consists of the principally polarised abelian varieties with a canonical filtration whose canonical type corresponds to the final type of w. Hence for properness we may assume that we have a principally polarised abelian variety over a discrete valuation ring R such that both its generic and special point are of type w and we suppose given a final flag over the generic point. Hence the canonical decomposition of Spec R for the abelian variety is equal to Spec Rand we have a canonical flag over Spec R. As $\overline{\mathcal{U}}_w$ is proper the map to it from the generic point of Spec R extends to a map from Spec R to $\overline{\mathcal{U}}_w$ hence giving a flag over its special point. This flag is then of a type $\leq w$ and hence by Lemma 8.3 it is equal to w and the image of Spec R lies in \mathcal{U}_w which proves properness.

Now, \mathcal{V}_w being by definition the schematic image of \mathcal{U}_w is reduced as \mathcal{V}_w is. As $\mathcal{U}_w \to \mathcal{V}_w$ is unramified it has reduced geometric fibres and as it is finite and \mathcal{V}_w is reduced, to show that it is flat it is enough to show that the cardinalities of the geometric fibres are the same for all geometric points of \mathcal{V}_w . This however is Lemma 4.6. Being finite, flat, and unramified it is étale. That its degree is $\gamma(w)$ follows from Lemma 4.6.

Remark 8.5. Note that the degree of the map $\mathcal{U}_w \to \mathcal{V}_w$ is $\gamma_g(w)$. It is not difficult by looking at the proof of Lemma 4.6 to show that it is a covering with structure group a product of linear and unitary groups over finite fields of characteristic p.

9. PUNCTUAL FLAG SPACES

Let M be the Dieudonné module of a truncated Barsotti-Tate group of level 1 over an algebraically closed field of characteristic p provided with an alternating perfect pairing (of Dieudonné modules). We let \mathcal{F}_M , the *punctual flag space* for M, be the scheme of self-dual admissible complete flags in M for which the middle element equals $\operatorname{Im}(V)$. It is well-known that every such M occurs as the Dieudonné

module of the kernel of multiplication by p on a principally polarised abelian variety and then \mathcal{F}_M is the intersection of $\overline{\mathcal{U}}_{\emptyset}$ and the fibre over a point of \mathcal{A}_q giving rise to M. Also by a result of Oort (cf., [Oo01]) the canonical type of it determines it (over an algebraically closed field) and hence we shall also use the notation \mathcal{F}_{ν} where ν is a final type. For $\Gamma = (I, \mathcal{S})$ where $I \subseteq \{1, \ldots, g\}$ with #I equal to the semi-simple rank of M and S a complete V stable flag of the V-semi-simple part of M, we then let \mathcal{F}_M^{Γ} be the part of $\mathcal{F}_M \cap \mathcal{U}_I^{ss}$ (which clearly can be defined directly in terms of M) for which the flag induces S on the V-semi-simple part. As the *p*-rank of M is f we easily see that \mathcal{F}_M is the disjoint union of the \mathcal{F}_M^{Γ} and putting $\mathcal{F}_{M}^{\mathcal{S}} := \mathcal{F}_{M}^{\{\{1,\ldots,g-f+1\},\Gamma\}}$ we have maps $S_{I} : \mathcal{F}_{M}^{I} \to \mathcal{F}_{M}^{\mathcal{S}}$. These maps are homeomorphisms by Proposition 4.14. This can be seen directly by decomposing M as $M^{mul} \bigoplus M^{\ell\ell} \bigoplus M^{et}$, where V is bijective on M^{mul} , F on M^{et} and F and V nilpotent on $M^{\ell\ell}$. Any element of an admissible flag over a perfect field will decompose in the same way (as that element is stable under F by definition and V by duality) and is hence determined by its intersection by M^{mul} , $M^{\ell\ell}$, and M^{et} . By self-duality the intersection of all the elements of the flag with M^{et} is determined by that with M^{mul} and that part is given by an arbitrary full flag of sub-modules of M^{mul} which is our S. That means that we may indeed reconstitute the whole flag from Γ and the induced flag on $M^{\ell\ell}$ and that any choice of flag on $M^{\ell\ell}$ gives rise to a flag in \mathcal{F}_M^{Γ} . That means that the map $\mathcal{F}_M^{\Gamma} \to \mathcal{F}_{M^{\ell\ell}}$ is a homeomorphism and we may for all practical purposes focus our attention on the case when F and V are nilpotent on M (i.e., M is local-local). Hence in this section unless otherwise mentioned the Dieudonné modules considered will be local-local. Note that the principal interest in this section will be focused on the question of which \mathcal{U}_w have a non-empty intersection with \mathcal{F}_M and that that problem is indeed by the above considerations immediately reduced to the local-local case.

We shall make extensive use of one way to move in each \mathcal{F}_M :

Consider a $w_{\emptyset} \ge w \in W_g$. Assume that we have index $1 \le i \le g-1$ for which $r_w(g+i-1,i+1) \ge i+1$. This means that for a flag \mathbb{D}_{\bullet} in $\overline{\mathcal{U}}_w$ we have that $F(\mathbb{D}_{i+1}) \subseteq \mathbb{D}_{i-1}$ or equivalently that F is zero on $\mathbb{D}_{i+1}/\mathbb{D}_{i-1}$. Hence if we replace \mathbb{D}_i by any $\mathbb{D}_{i-1} \subset \mathbb{D} \subset \mathbb{D}_{i+1}$ (replacing also \mathbb{D}_{2g-i} to make the flag self-dual) we shall still have an admissible flag as $V(\mathbb{D}) \subseteq \mathbb{D}_{i-1}$. In order to construct the \mathbb{E} -flag we apply V to the \mathbb{D} -flag which gives us half of the \mathbb{E} -flag and we complement by taking orthogonal spaces. In the \mathbb{E} -flag now $\mathbb{E}_{g-i+1}/\mathbb{E}_{g-i-1}$) to \mathcal{F}_M and we shall therefore call this family the simple family of index i and we shall write $P_{w,i}$ for this simple family. The condition $r_w(g+i-1,i+1) \ge i+1$ is equivalent to $r_w(g-i+1,i-1) = g-i-1$ and when it is fulfilled we shall say that g-i is movable for w.

Proposition 9.1. Any two points of the local flag space \mathcal{F}_M^{Γ} can be connected by a sequence of simple families.

Proof. We immediately reduce to the case when M is local (in which case the statement is about \mathcal{F}_M). We are going to identify \mathcal{F}_M with the scheme of V-stable flags in $\mathrm{Im}(V)$ and we prove the statement for any Dieudonné module N with F = 0 and V nilpotent. Let E and F be two V-stable flags in N. If $E_1 = F_1$ then we may consider N/E_1 and use induction on the length of N to conclude. If not we use induction on the smallest i such that $F_1 \subseteq E_i$ which we thus may assume to
be > 1. We now have $F_1 \subsetneq E_{i-1}$ and hence that F_1 is a complement to E_{i-1} in E_i so that in particular $E_i/E_{i-2} = E_{i-1}/E_{i-2} \bigoplus (F_1 + E_{i-2})/E_{i-2}$ which has a consequence that V is zero on E_i/E_{i-2} which means that every subspace of it is stable under V so that we get a \mathbb{P}^1 -family of flags in E_i/E_{i-2} in which both E_{i-1}/E_{i-2} and $(F_1 + E_{i-2})/E_{i-2}$ are members so that we may move E_{i-1} so that it contains F_1 .

Recall (cf, [Oo01, 14.3]) that one defines the partial order relation on final types $\nu_1 \subseteq \nu_2$ (resp. $\nu_1 \subset \nu_2$) by the condition that $\mathcal{V}_{\nu_1} \subseteq \overline{\mathcal{V}}_{\nu_2}$ (resp. $\mathcal{V}_{\nu_1} \subsetneq \overline{\mathcal{V}}_{\nu_2}$). We shall now see that this relation can be expressed in terms of local flag spaces. For this we let M_{ν} be a Dieudonné module of a principally polarised Barsotti-Tate group of level 1 with final type ν (there is up to isomorphism only one such M_{ν} , [Oo01, Thm. 9.4]).

Theorem 9.2. i) We have that $\nu' \subset \nu$ precisely when there is a $w \in W_g$ such that $w \leq \nu$ and there is a flag of type w in $\mathcal{F}_{M,\prime}$.

ii) If there is a flag of type w in $\mathcal{F}_{M_{\nu'}}$ then there is a $w' \leq w$ such that the intersection $\mathcal{U}_{w'} \cap \mathcal{F}_{M_{\nu'}}$ is finite.

Proof. Consider the image in \mathcal{A}_g of $\overline{\mathcal{U}}_{\nu}$. It is a closed subset containing \mathcal{V}_{ν} and hence contains $\overline{\mathcal{V}}_{\nu}$ and in particular it meets each fibre over a point of $\overline{\mathcal{V}}_{\nu}$ and consequently there is a point *s* in the intersection of $\overline{\mathcal{U}}_{\nu}$ and the fibre over a point *t* of $\mathcal{V}_{\nu'}$. Now, *s* lies in some $\mathcal{U}_w \subseteq \overline{\mathcal{U}}_{\nu}$ and consequently fulfills $w \leq \nu$ and as $\nu \leq w_{\emptyset}$, *s* also lies in the local flag space of *t* and as has been noted this is the "same" as $\mathcal{F}_{M_{\nu'}}$. The converse is clear.

As for the second part, the proof of Lemma 6.2 shows that a $w' \leq w$ which is minimal for the condition that $\mathcal{U}_{w'} \cap \mathcal{F}_{M_{\nu'}}$ is non-empty has $\mathcal{U}_{w'} \cap \mathcal{F}_{M_{\nu'}}$ finite. \Box

The theorem allows us to reprove a result of Oort (cf., [Oo01]).

Corollary 9.3. The E-O stratification on \mathcal{A}_q is a stratification.

Proof. The condition in 9.2 says that $\nu' \subset \nu$ if and only if the closure $\overline{\mathcal{U}}_{\nu}$ of \mathcal{U}_{ν} has a non-empty intersection with the punctual flag space $\mathcal{F}_{\nu'}$. The proof gives more precisely that a given point s of $\mathcal{V}_{\nu'}$ lies in $\overline{\mathcal{V}}_{\nu}$ precisely when $\overline{\mathcal{U}}_{\nu}$ intersects the fibre over s of the map $\mathcal{F}_g \to \mathcal{A}_g$. This condition does not depend on point s by a result of Oort on Dieudonné modules (cf., [Oo01]).

From this theorem it is clear that the condition that $\mathcal{U}_w \cap \mathcal{F}_M \neq \emptyset$ is important. We shall say that an admissible $w \in W_g$ occurs in ν , with ν a final type, if $\mathcal{U}_w \cap \mathcal{F}_M \neq \emptyset$ and we shall write it symbolically as $w \to \nu$.

Remark 9.4. It is important to realise that a priori this relation depends on the characteristic that is implicit in all of this article. Hence the notation $w \xrightarrow{p} \nu$ would be more appropriate. It is our hope that the relation will a posteriori turn out to be independent of p. If not and if one is working with several p the more precise notation will have to be used.

Hence we can formulate the theorem as saying that $\nu' \subset \nu$ precisely when there exists an admissible w with $w \to \nu'$ and $w \leq \nu$. Suppose final types ν and ν' given. An element w of minimal length in the set of minimal elements of $\{w \in W_g : \nu > w, w \to \nu'\}$ in the Bruhat-Chevalley order has the property that $\mathcal{U}_w \cap \mathcal{F}_{M_{\nu'}}$ has

dimension 0 for the generic point of $\mathcal{V}_{\nu'}$. Clearly, then $\ell(w) \ge \ell(\nu')$ for every w as in 9.2.

Example 9.5. Since E-O strata on \mathcal{A}_g are defined using the projection from the flag space, the closure of an E-O stratum on \mathcal{A}_g need not be given by the Bruhat-Chevalley order on the set of final elements and indeed it isn't. Oort gave the first counterexample for g = 7 based on products of abelian varieties. We reproduce his example and give two other ones, one for g = 5 and one for g = 6 that do not come from products.

i) Let g = 7 and let $w_1 = [1, 2, 4, 6, 7, 10, 12]$ and $w_2 = [1, 2, 3, 7, 9, 10, 11]$. Then w_1 and w_2 are final elements of W_7 and have length $\ell(w_1) = 8$ and $\ell(w_2) = 9$. In the Bruhat-Chevalley order neither $w_1 \leq w_2$ nor $w_2 \leq w_1$ holds. Despite this we have $\overline{\mathcal{V}}_{w_1} \subset \overline{\mathcal{V}}_{w_2}$. The explanation for this lies in the fact that the simple family $P_{w_1,4}$ hits the stratum U_{w_3} with w_3 the element $[1, 2, 3, 7, 6, 10, 11] = s_3 w_1 s_4$, with $w_2 > w_3$ and $w_3 \to w_1$, so by 9.2 it follows that $\overline{\mathcal{V}}_{w_1} \subset \overline{\mathcal{V}}_{w_2}$. (That there is such a simple family can be proved directly but for now we leave it is an unsupported claim as a proof "by hand" would be somewhat messy. A more systematic study of these phenomena will appear in a subsequent paper.) This explains the phenomenon observed in [Oo01], p. 406 (but note the misprints there). Also the element $w_2 > 0$ $w_4 = [1, 2, 3, 7, 9, 5, 11] \rightarrow w_1$ will work for w_1 . The element w_1 is the final element corresponding to taking the product of a Dieudonné module with final element [135] and a Dieudonné module with final element [1246] whereas similarly w_2 appears as the "product" of the final elements [135] and [1256]. As [1246] < [1256] there is a degeneration of a Dieudonné module of type [1256] to one of type [1246]. This shows that this example simply expresses the fact that \subset must be stable under products whereas the Bruhat-Chevalley order isn't. (We'd like to thank Ben Moonen for pointing this out to us.)

ii) For g = 5 we consider the final elements $w_1 = [1,3,4,6,9]$ and $w_2 = [1,2,6,7,8]$ of length 5 and 6 and the non-final element $w_3 = [1,2,6,4,8]$ in W_5 . Then $w_3 < w_2$ and $w_3 \rightarrow w_1$, so that \mathcal{V}_{w_1} lies in the closure of \mathcal{V}_{w_2} . But in the Bruhat-Chevalley order neither $w_1 < w_2$ nor $w_2 < w_1$ holds.

iii) Let g = 6 and consider the final elements $w_1 = [1, 3, 5, 6, 9, 11]$ and $w_2 := [1, 2, 6, 8, 9, 10]$ of length $\ell(w_1) = 8$ and $\ell(w_2) = 9$. In the Bruhat-Chevalley order we do not have $w_1 \leq w_2$. Nevertheless, \mathcal{V}_{w_1} occurs in the closure of the E-O stratum \mathcal{V}_{w_2} . Indeed, the admissible element $w_3 = [1, 2, 6, 8, 4, 10]$ satisfies $w_2 \geq w_3 \rightarrow w_1$: \mathcal{U}_{w_3} has a non-empty intersection with the punctual flag space \mathcal{F}_{w_1} . This time neither of the elements w_1 and w_2 are products in the sense of i). Furthermore, as \mathcal{V}_{w_1} is of codimension 1 in \mathcal{V}_{w_2} this example can not derived by taking the transitive closure of the closure under products of the Bruhat-Chevalley relation. The claim that we have $w_3 \rightarrow w_1$ and the two preceding ones will be substantiated in a subsequent paper.

There is an approach to the study of the relation of the E-O strata and the strata on \mathcal{F}_g which is in some sense "dual" to the study of punctual flag spaces; that of considering the image in \mathcal{A}_g of the \mathcal{U}_w . The following result gives a compatibility result on these images and the E-O stratification.

Proposition 9.6. i) The image of any \mathcal{U}_w , $w \in W_g$, is a union of strata \mathcal{V}_{ν} . In particular the image of a $\overline{\mathcal{U}}_w$ is equal to some $\overline{\mathcal{V}}_{\nu}$.

ii) For any final ν and $w \in W_g$, the maps $\mathcal{U}_w \cap \pi^{-1} \mathcal{V}_\nu \to \mathcal{V}_\nu$ and $\overline{\mathcal{U}}_w \cap \pi^{-1} \mathcal{V}_\nu \to \mathcal{V}_\nu$, where π is the projection $\mathcal{F}_g \to \mathcal{A}_g$, have the property that there is a surjective flat map $X \to \mathcal{V}_\nu$ such that the pullback of them to X is isomorphic to the product $X \times (\mathcal{F}_\nu \cap \mathcal{U}_w)$ resp. $X \times (\mathcal{F}_\nu \cap \overline{\mathcal{U}}_w)$.

iii) A generic point of a component of \mathcal{U}_w maps to the generic point of some \mathcal{V}_{ν} and that ν is independent of the chosen component of \mathcal{U}_w .

Proof. The first part follows directly from Oort's result on the uniqueness for the Dieudonné module in a stratum \mathcal{V}_{ν} as it implies that if one fibre of $\pi^{-1}(\mathcal{V}_{\nu}) \to \mathcal{V}_{\nu}$ meets \mathcal{U}_w then they all do. As for the second part it would follow if we could prove that there is a surjective flat map $X \to \mathcal{V}_{\nu}$ such that the pullback of $(\mathbb{H}, \mathbb{E}, F, V, \langle -, -\rangle)$ is isomorphic to a the constant data (provided by the Dieudonné module of type ν). For this we first pass to the space X_{ν} of bases of \mathbb{H} for which the first g elements form a basis of \mathbb{E} which is flat surjective over \mathcal{V}_{ν} . Over X_{ν} the data is the pullback from a universal situation where F, V, and $\langle -, -\rangle$ are given by matrices. In this universal situation we have an action of the group G of base changes and two points over an algebraically closed field give rise to isomorphic ($\mathbb{H}, \mathbb{E}, F, V, \langle -, -\rangle$) precisely when they are in the same orbit. By assumption (and Oort's theorem) the image of X_{ν} lies in an orbit so it is enough to show that the data over an orbit can be made constant by a flat surjective map. However, the map from G to the orbit obtained by letting g act on a fixed point of the orbit has this property.

The third part follows directly from the second.

The proposition gives us a map $\tau_p \colon W_g \to W_g/S_g$ which to w associates the final type of the open stratum into which each generic point of \mathcal{U}_w maps. We shall return to this map in Section 13.

Example 9.7. Note that the punctual flag space is in general rather easy to understand as it only depends on the image of V and we are almost talking about the space of flags stable under a nilpotent endomorphism (remember that we have reduced to the local-local case). Almost but not quite as the endomorphism is semi-linear rather than linear. What is complicated is the induced stratification. Already the case of $\nu = s_3 \in W_3$ is an illustrative example. We have then that $\ker V \cap \operatorname{Im} V$ is of dimension 2 (we have on Jordan block for V on $\operatorname{Im} V$ is size 2 and one of size 1). The first element, \mathbb{E}_1 , of the flag must lie in ker $V \cap \operatorname{Im} V$ so we get a \mathbb{P}^1 of possibilities for it. If $\mathbb{E}_1 = \operatorname{Im} V^2$, then V is zero on $\mathbb{E}_3/\mathbb{E}_1$ and we can choose $\mathbb{E}_2/\mathbb{E}_1$ as an arbitrary subspace of \mathbb{E}_3 giving us a \mathbb{P}^1 of choices for \mathbb{E}_2 . On the other, if $\mathbb{E}_1 \neq \text{Im} V^2$, then $\mathbb{E}_3/\mathbb{E}_1$ has a Jordan block of size 2 and hence there is only one V-stable 1-dimensional subspace and thus the flag is determined by \mathbb{E}_1 . The conclusion is that the punctual flag space is the union of two \mathbb{P}^1 's meeting at a single point. The intersection point is the canonical filtration (which is a full flag) and one can show that the rest of the points on one component are flags of type [241] and the rest of the points on the other are flags of type [315].

10. Pieri formulas

In this section we are going to apply a theorem of Pittie and Ram ([PR99]) to obtain a Pieri type formula for our strata. (It seems to be historically more correct to speak of Pieri-Chevalley type formulas, cf,[Ch94].) The main application of it will not be to obtain cycle class formulas, as Pieri formulas usually do it will not

give formulas for individual strata but only for certain linear combinations. To us the principal use of these formulas will be that it shows that a certain strictly positive linear combination of the boundary components will be a section of an ample line bundle (or close to ample as one of the contributors to ampleness will be λ_1 which is ample only on the Satake compactification). This will have as consequence affineness for the open strata as well as a connectivity result for the boundary of the closed ones. We shall see in Section 13 that there is also a Pieri formula for the classes of the E-O strata though we know very little about it.

In this section we are going to work with level structures. There are two reasons for this. The first one is that we are going to exploit the ampleness λ_1 and even formulating the notion of ampleness for a Deligne-Mumford stack is somewhat awkward. The second is that one of the consequences of our considerations will be an irreducibility criterion for strata. Irreducibility for a stratum on \mathcal{A}_g does not imply irreducibility for the same stratum on the space $\mathcal{A}_{g,n}$ of principally polarised abelian varieties with level *n*-structure, where always $p \not| n$. In fact irreducibility for the level *n* case means irreducibility on \mathcal{A}_g together with the fact that the monodromy group of the level *n* cover is the maximum possible. Hence in this section we shall use $\mathcal{A}_{g,n}$ but also some toroidal compactification $\tilde{\mathcal{A}}_{g,n}$ (cf., [FC90]). Everything we have said so far applies to this situation giving us in particular $\mathcal{F}_{g,n}$ and $\tilde{\mathcal{F}}_{g,n}$ but we have the extra property that for $n \geq 3$ then $\tilde{\mathcal{A}}_{g,n}$ and hence $\tilde{\mathcal{F}}_{g,n}$ are smooth projective varieties.

We now introduce the classes $\ell_i := c_1(\mathbb{E}_{\{i\}})$ for $1 \leq i \leq 2g$ in the Chow ring $CH^*(\mathcal{F}_g)$. By self-duality of the flag \mathbb{E}_{\bullet} we have that $\ell_{2g+1-i} = -\ell_i$ and by construction $c_1(\mathbb{D}_{\{i\}}) = p\ell_{i-g} = -p\ell_{3g+1-i}$ for $g+1 \leq i \leq 2g$. Furthermore, $\ell_1 + \cdots + \ell_g$ is the pullback from $\tilde{\mathcal{A}}_q$ of λ_1 , the first Chern class of the Hodge bundle.

Now we let $M_i := c_1(\mathbb{D}_{[2g-i,2g]}), 1 \leq i \leq g$, and start by noting that if $n = (n_1, \ldots, n_g)$ then $n \cdot M := \sum_i n_i M_i$ is relatively ample for $\tilde{\mathcal{F}}_g \to \tilde{\mathcal{A}}_g$ if $n_i > 0$ for $1 \leq i < g$. Indeed, by construction $L_i := \ell_{2g} + \cdots + \ell_{2g-i+1}, 1 \leq i \leq g$ is the pullback from the partial flag space $\tilde{\mathcal{F}}_g[i]$ of flags with elements of rank i, 2g - i, and g (and with the rank g-component equal to \mathbb{E}_g) and on $\tilde{\mathcal{F}}_g[i]$ we have that $\ell_{2g} + \cdots + \ell_{2g-i+1}$ is ample. It is then well-known that any strictly positive linear combination of these elements is relatively ample. From the formulas above we get that $M_i = p(L_{g-i} + \lambda_1)$ (where we put $L_0 = 0$). On the other hand λ_1 is almost ample; it is the pullback from \mathcal{A}_g^* of an ample line bundle.

Now we identify the L_i with the fundamental weights of the root system of C_g . Note that W_g acts on the ℓ_i considered as parts of the weight lattice by $\sigma(\ell_i) = \ell_{\sigma(i)}$ (keeping in mind that $\ell_{2g+1-i} = -\ell_i$) and then acts accordingly on the L_i . Let us also note that (by Chevalley's characterisation of the Bruhat-Chevalley order) if w' < w with $\ell(w') = \ell(w) - 1$ and if $w = s_{i_1} \cdots s_{i_k}$, then w' is of the form $s_{i_1} \cdots \widehat{s_{i_r}} \cdots s_{i_k}$ which can be rewritten as ws_{α} , where $s_{\alpha} = (s_{i_{r+1}} \cdots s_{i_k})^{-1} s_{i_r} (s_{i_{r+1}} \cdots s_{i_k})$ which thus is the reflection with respect to a unique positive root.

Theorem 10.1. For each $1 \leq i \leq g$ and $w \in W_g$ we have that

$$(p\lambda_1 + pL_{g-i} - wL_i)[\overline{\mathcal{U}}_w] = \sum_{w' \prec w} c^i_{w,w'}[\overline{\mathcal{U}}_{w'}] \in \mathrm{CH}^1_{\mathbb{Q}}(\overline{\mathcal{U}}_w),$$

where $c_{w,w'}^i \ge 0$ and $w' \prec w$ means $w' \le w$ and $\ell(w) = \ell(w') + 1$. Furthermore, $c_{w,w'}^i > 0$ precisely when $w' = ws_{\alpha}$ for α a positive root for which the simple root α_i

appears with a strictly positive coefficient when α is written as a linear combination of the simple roots.

Proof. We shall use [PR99] which has the following setup: We fix a semi-simple algebraic group G (which in our case is the symplectic group Sp_{2g} but using this in the notation will only confuse) with Borel group B and fix a principal B-bundle $E \to X$ over an algebraic variety X. Letting $E(G/B) \to X$ be the associated G/B-bundle we have, because its structure group is B and not just G, Schubert varieties $\Omega_w \to X$ (which fibre by fibre are the usual Schubert varieties). For every weight $\lambda \in P$, P being the group of weights for G, we have two line bundles on E(G/B); on the one hand y^{λ} obtained by regarding λ as a character of B which gives a G-equivariant line bundle on G/B and hence a line bundle on E(G/B), on the other hand the character λ can also be used to construct, with the aid of the principal B-bundle E, a line bundle x^{λ} on X and then by pullback to E(G/B) a line bundle also denoted by x^{λ} . A result of [PR99, Corollary] then says that if λ is a dominant weight then

$$y^{\lambda}[\mathcal{O}_{\Omega_w}] = \sum_{\eta \in \mathcal{T}_w^{\lambda}} x^{\eta(1)}[\mathcal{O}_{\Omega_{v(\eta,w)}}] \in K_0(E(G/B)).$$

Here \mathcal{T}_w^{λ} is a certain set of piecewise linear paths $\eta \colon [0,1] \to P \bigotimes \mathbb{R}$ in the real vector space spanned by P and $v(\eta, w)$ is a certain element in the Weyl group of G which is always $\leq w$ and \mathcal{T}_w^{λ} has the property that $\eta(1) \in P$ for all its elements η . An important property of \mathcal{T}_w^{λ} is that it only depends on w and λ and not on E. It follows immediately from the description of [PR99], that $v(\eta, w) = w$ in only one case, namely when η is the straight line $\eta(t) = tw\lambda$. Hence we can rewrite the formula as

$$(y^{\lambda} - x^{w\lambda})[\mathcal{O}_{\Omega_w}] = \sum_{\eta \in \mathcal{T}_w^{\lambda}} x^{\eta(1)}[\mathcal{O}_{\Omega_{v(\eta,w)}}],$$

where the sum now runs over all elements of \mathcal{T}_w^{λ} for which $v(\eta, w) < w$. Taking Chern characters and looking at the top term which appears in codimension $\operatorname{codim}(w) + 1$ we get

$$(c_1(y^{\lambda}) - c_1(x^{w\lambda}))[\Omega_w] = \sum_{\eta \in \mathcal{T}_w^{\lambda}} [\Omega_{v(\eta,w)}],$$

where the sum is now over the elements of $\mathcal{T}_{w}^{\lambda}$ for which $\ell(v(\eta, w)) = \ell(w) - 1$. To determine the multiplicity with which a given $[\Omega_{w'}]$ appears in the right hand side we could no doubt use the definition of $\mathcal{T}_{w}^{\lambda}$. However, it seems easier to note that that multiplicity is independent of E and hence we may assume that X is a point and by additivity in λ that λ is a fundamental weight λ_i . In that case one can use a result of Chevalley ([Ch94, Prop. 10]) to get the description of the theorem. However, we want this formula to be true not in the Chow group of E(G/B) but instead in the Chow group of the relative Schubert subvariety of index w of E(G/B). This however is no problem as the (relative) cell decomposition shows that this Chow group injects into the Chow group of E(G/B).

In our case we now use $X = \mathcal{F}_g$ and \mathbb{E}_{\bullet} as *B*-bundle and then pull back this formula along the section provided by \mathbb{D}_{\bullet} (over $\overline{\mathcal{U}}_w$ so that the section takes values in the relative Schubert variety of index *w*). To make the pullback possible (note that the relative Schubert variety will in general not be smooth over the base) we remove the relative Schubert varieties of codimension 2 in the relative Schubert variety in question. This forces us to remove the part of $\overline{\mathcal{U}}_w$ where the section encounters the removed locus. This is however a codimension 2 subset by Corollary 8.4 so its removal will not affect $\operatorname{CH}^1_{\mathbb{O}}(\overline{\mathcal{U}}_w)$. There is a small problem however. The result of [PR99] is formulated under the assumption of a smooth variety X over the complex numbers. In our case the base is an algebraic stack in positive characteristic. We do not see any problems in extending their proof to our situation; the assumption of a spatial base seems irrelevant and characteristic zero is used for a vanishing theorem ([PR99, Proof of Proposition]) which seems to hold in all characteristics for the situation at hand (by for instance Frobenius splitting). However, we do not need to verify this. Instead we can use that we can first introduce a level structure to make the base spatial and then pull back the vector bundle from a Grassmannian (and hence the flag from a flag space over such a Grassmannian). This case can be lifted to characteristic zero and then the result is obtained from the characteristic zero result by using the specialisation map in K-theory. (A somewhat more sophisticated reduction would no doubt prove the result for any X, E, and G.) \square

To apply the theorem we start by some preliminary results which will be used to exploit the positivity of the involved line bundles.

Lemma 10.2. Let X be a proper (irreducible) variety of dimension > 1 and \mathcal{L} a line bundle on X which is ample on some open subset $U \subseteq X$. Let $D := X \setminus U$ and $H \subset U$ the zero set of a section of $\mathcal{L}_{|U}$. If D is connected then so is $D \cup H$.

Proof. By replacing the section by a power of it, we may assume that \mathcal{L} is very ample giving an embedding $U \hookrightarrow \mathbb{P}^n$. Let Z be the closure of the graph of this map in $X \times \mathbb{P}^n$ and let Y the image of Z under the projection on the second factor giving us two surjective maps $X \leftarrow Z \to Y$ and let D' be the inverse image in Z of D. Assume that $D \cup H$ is the disjoint union of the non-empty closed subsets A and B and let A' and B' be their inverse images in Z. Now, Y is irreducible of dimension > 1 and hence H'' is connected, where H'' is the hyperplane section of Y corresponding to H, so that the images of A' and B' in Y must meet. However, outside of D' the map $Z \to Y$ is a bijection and hence the meeting point must lie below a point of D' and hence A' and B' both meet D'. This implies that A and B both meet D which is a contradiction as D is assumed to be connected.

Proposition 10.3. Let \mathcal{L} be the determinant det \mathbb{E} of the Hodge bundle over $\mathcal{A}_{g,n}$, $n \geq 3$ (and prime to p).

i) There is, for each $1 \leq i < g$, an integer m_i such that the global sections of $\Lambda^{g-i}(\mathbb{H}/\mathbb{E}) \bigotimes \mathcal{L}^{\otimes m}$ generate this bundle over $\mathcal{A}_{g,n}$ whenever $m \geq m_i$. These m_i can be chosen independently of p (but depending on g and n).

ii) Putting $N_i := L_i + n_i \lambda_1$ for $1 \le i < g$ and $N_g := \lambda_1$, then $\sum_i m_i N_i$ is ample on \mathcal{F}_g if $m_i > 0$ for all $1 \le i \le g$.

iii) Fix $w \in W_g$ and put $L := \sum_{i < g} L_i$, $N := \sum_{i < g} N_i$, and $m = \sum_i m_i$. Choose r, s, t, and u such that $rN + t\lambda_1 - wL$ resp. $sN + u\lambda - wL_g$ can be written as a positive linear combination of the N_i (using that $L_g = -\lambda_1$). Then if p > r + sm and (g-1)p > t + um we have that $p(L + (g-1)\lambda_1) + pm\lambda_1 - wL - mwL_g$ is ample on $\mathcal{F}_{g,n}$. The constants r, s, t, and u can be chosen independently of p.

Proof. Starting with i) it follows directly from the fact that λ_1 is ample on $\mathcal{A}_{g,n}$. The independence on p follows from the existence of a model of $\mathcal{A}_{g,n}$ that exists over Spec $\mathbb{Z}[1/n, \zeta_n]$.

As for ii) we have that $\pi_* \mathcal{O}(L_i) = \Lambda^{g-i}(\mathbb{H}/\mathbb{E}), 1 \leq i < g$, as $\pi : \mathcal{F}_{g,n} \to \mathcal{A}_{g,n}$ can be identified with the space of flags on \mathbb{H}/\mathbb{E} and $\mathcal{O}(L_i)$ is $\det(\mathbb{H}/\mathbb{E}_{2g-i})$. By definition we then have that $\pi_* \mathcal{O}(N_i)$ is generated by global sections on $\mathcal{A}_{g,n}$. We know that on the flag space SL_g/B we have that the canonical ring $\bigoplus_{\lambda} H^0(\operatorname{SL}_g/B, \mathcal{L}_{\lambda})$, where λ runs over the dominant weights and \mathcal{L}_{λ} is the corresponding line bundle, is generated by the $H^0(\operatorname{SL}_g/B, \mathcal{L}_{\lambda_i}), 1 \leq i < g$, where λ_i is the *i*'th fundamental weight (see for instance [RR85]). Also $\mathcal{O}(\sum_{i < g} m_i N_i)$ is relatively very ample and we have just shown that $\pi_* \mathcal{O}(\sum_{i < g} m_i N_i)$ is generated by global sections. As λ_1 is ample on $\mathcal{A}_{g,n}$ we get that $\pi_* \mathcal{O}(\sum_{i \leq g} m_i N_i)$ is ample. Continuing with iii) We have that $N = L + n\lambda_1$ which gives $p(L + (g - 1)\lambda_1) +$

Continuing with iii) We have that $N = L + n\lambda_1$ which gives $p(L + (g - 1)\lambda_1) + pm\lambda_1 - wL - mwL_g = (p - (r + sm))N + (p(g - 1) - (t + um)) + (rN + t\lambda_1 - wL) + m(sN + u\lambda_1 - wL_g)$. We then conclude by the definitions of r, s, t, and u and ii).

Remark 10.4. i) The constants r, s, t, and u are quite small and easy to compute. We know nothing about the m_i but imagine that they would not be too large.

ii) It would seem that the last part would not be applicable for g = 1 but it can be easily modified to do so. On the other hand for g = 1 everything is trivial anyway.

We are now ready for the first application of the Pieri formula.

Proposition 10.5. *i)* There is a bound depending only on g and n such that if p is larger than that bound, then for an irreducible component Z of some $\mathcal{U}_w \subseteq \tilde{\mathcal{F}}_{g,n}$, $w \in W_g$, the union of the complement of Z in \overline{Z} and the intersection of \overline{Z} and $\tilde{\mathcal{F}}_{g,n} \setminus \mathcal{F}_{g,n}$ is connected if the intersection of \overline{Z} with the boundary $\tilde{\mathcal{F}}_{g,n} \setminus \mathcal{F}_{g,n}$ is connected or empty.

ii) There is a bound depending only on g and n such that if p is larger than that bound, then for $w \in W_q$ of semi-simple rank 0 we have that \mathcal{U}_w is affine.

Proof. By Proposition 10.3 there is a bound depending only on g and n such that if p is larger than it, then $M := p(L + (g - 1)\lambda_1) + pm\lambda_1 - wL - mwL_g$ is ample on $\mathcal{F}_{g,n}$. Summing up Pieri's formula (Theorem 10.1) for $1 \leq i < g$ and m times the formula for i = g we get that $M[\overline{Z}]$ is supported on \overline{Z} intersected with smaller strata. (Note that the Pieri formula is a priori – and quite likely in reality – only true modulo torsion. We may however simply multiply it with a highly divisible integer and that doesn't change the support.) We then conclude by Lemma 10.2 once we have .

As for the second part we argue as in the first part and conclude that $\overline{\mathcal{U}}_w \setminus \mathcal{U}_w$ is the support of an ample divisor (as in the theorem each component must appear as we are summing up for $1 \leq i \leq g$ and some α_i must appear in the expansion of α) and hence \mathcal{U}_w is affine.

Remark 10.6. i) The condition on the intersection with the boundary somewhat cuts down on possible applications of the first part. In the applications of the next section it turns out that we only need to apply it when the intersection is empty.

ii) For the second part we would like to say more generally that the image of \mathcal{U}_w is affine in $\mathcal{F}_{g,n}^*$ for some appropriate definition of $\mathcal{F}_{g,n}^*$ analogous to the Satake

compactification. The problem is that it doesn't seem as if some power of M would normally be generated by its global sections so that we can not define $\mathcal{F}_{g,n}^*$ as the image of $\tilde{\mathcal{F}}_{g,n}$.

11. IRREDUCIBILITY PROPERTIES

In this section we shall prove irreducibility of a large class of strata and also that if the characteristic is large enough and our irreducibility criterion is not fulfilled then (with some extra conditions on the stratum) the stratum is reducible. Our proofs show two advantages of working on the flag spaces. The major one is that our strata are normal so that irreducibility follows from connectedness. The connectedness of the closed E-O strata except $\overline{\mathcal{V}}_1$ is proven in [Oo01] but the $\overline{\mathcal{V}}_w$ are most definitely not locally connected and hence that does not say very much about the irreducibility. In the converse direction we also make use of the Pieri formula.

Definition-Lemma 11.1. Let $\{Z_{\alpha}\}$ be a stratification of a Deligne-Mumford stack X of finite type over a field by which we mean that the strata Z_{α} are locally closed reduced substacks of X such that the closure $\overline{Z_{\alpha}}$ of a stratum is the union of strata. By the k-skeleton of the stratification we mean the union of the strata of dimension $\leq k$ (which is a closed substack). The boundary of a stratum Z_{α} is the complement of Z_{α} in its closure. Assume furthermore that each Z_{α} is irreducible and that for Z_{α} of dimension strictly greater than some fixed N we have that its boundary is connected (and in particular non-empty). Then the intersection of a connected union Z of closed strata $\overline{Z_{\alpha}}$ with the N-skeleton is connected.

Proof. It is enough, by induction to prove that if we remove a stratum Z' of maximal dimension which is > N from Z, then the result remains connected. Assume that $Z \setminus Z'$ is the disjoint union of two closed subset Z_1 and Z_2 . By assumption the boundary of Z' is connected and hence lies in Z_1 say. That means that $Z' \cup Z_1$ is closed and disjoint from Z_2 which by the connectedness of Z implies that Z_2 is connected.

Proposition 11.2. There is a bound depending only on g and n such that the following is true if p is larger than that bound:

Let $X \subseteq \tilde{\mathcal{F}}_{g,n}$ be a connected union of irreducible components of closed strata $\overline{\mathcal{U}}_w$ (for possibly different w) which lie inside of $\mathcal{F}_{g,n}$. Then the intersection of X with the 1-skeleton is connected.

Proof. This follows directly from Proposition 10.5 and Lemma 11.1. (Note that for level 1 or 2 we may pass to a higher level in order to apply the proposition.) \Box

We now want to interpret this Proposition (and its converse which will be true for any p) in arithmetical terms. Hence we define the 1-skeleton graph of level n as the following edge-colored graph: Its vertices are the points of $\overline{\mathcal{U}}_1 \subset \mathcal{F}_{g,n}$, i.e., isomorphism classes of principally polarised superspecial g-dimensional abelian varieties A together with a level n structure and a complete flag $0 = \mathbb{D}_0 \subset \mathbb{D}_1 \subset$ $\cdots \subset \mathbb{D}_g = H^0(A, \Omega^1_A)$ on $H^0(A, \Omega^1_A)$ for which $\mathbb{D}_{g-i}^\perp = V^{-1}\mathbb{D}_i$. For each $1 \leq i \leq g$ we connect two vertices by an edge of color i if there is an irreducible component of $\overline{\mathcal{U}}_{s_i}$ which contains them. **Lemma 11.3.** If $S \subseteq \{1, \ldots, g\}$ has the property that it contains g and for every $1 \leq i < g$ we have that either i or g - i belongs to S, then the subgraph of the 1-skeleton graph consisting of all vertices and all edges of colours $i \in S$ is connected.

Proof. This follows from [Oo01, Prop. 7.3] and Theorem 7.5.

For a subset $S \subseteq \{1, \ldots, g\}$ the *S*-subgraph of the 1-skeleton graph is the subgraph with the same vertices and with only the edges whose colour is in *S*. This definition allows us to formulate our irreducibility conditions.

Theorem 11.4. i) Let $w \in W_g$ and let $S := \{1 \le i \le g : s_i \le w\}$. If the S-subgraph of the 1-skeleton graph is connected then $\overline{\mathcal{U}}_w \subseteq \mathcal{A}_g$ is irreducible.

ii) There is a bound depending only on g such that if p is larger than that bound the following is true: If $w \in W_g$ is admissible and either final or of semi-simple rank 0 and if $S := \{1 \le i \le g : s_i \le w\}$, then there is a bijection between the irreducible components of $\overline{\mathcal{U}}_w \subseteq \mathcal{A}_q$ and the connected components of the 1-skeleton graph.

Proof. The first part is clear as Proposition 6.1 says that each component meets $\overline{\mathcal{U}}_1$ and then by the assumption on connectedness of the S-subgraph $\overline{\mathcal{U}}_w$ is connected but by Corollary 8.4 it is normal and hence is irreducible.

As for the second part assume first that w is final but of positive semi-simple rank. That means that its Young diagram does not contain a row of length g and hence by Lemma 2.7 and the Chevalley characterisation of the Bruhat-Chevalley order we have that $s_i \leq w$ for all and hence the S-subgraph is connected by Lemma 11.3 which make the statement trivially true. We may therefore assume that the semi-simple rank is 0 and hence that $\overline{\mathcal{U}}_w$ lies entirely in $\mathcal{F}_{g,n}$. In that case the result follows from Proposition 11.2 and the fact that two irreducible components of two $\overline{\mathcal{U}}_{s_i}$ meet only at $\overline{\mathcal{U}}_1$.

Projecting down to \mathcal{A}_g we get the following corollary which shows irreducibility for many E-O strata.

Theorem 11.5. Let $w \in W_g$ be a final element whose Young diagram Y does not contain all rows of length i with $\lceil (g+1)/2 \rceil \leq i \leq g$. Then $\overline{\mathcal{V}}_w$ is irreducible and the étale cover $\mathcal{U}_w \to \mathcal{V}_w$ is connected.

Proof. This follow from Theorem 11.4 and Lemmas 2.7 and 11.3.

Example 11.6. For g = 2 the locus of abelian surfaces of p-rank ≤ 1 is irreducible. For g = 3 all E-O strata except the superspecial locus $(Y = \{1, 2, 3\})$ and the Moret-Bailly locus $(Y = \{2, 3\})$ are irreducible.

In [Ha04] S. Harashita has proved that the number of irreducible components of an E-O stratum that is contained in the supersingular locus is given as a class number and as a consequence that, except possibly for small p, these strata are reducible. As has been proved by Oort (cf., [Ha04, Prop. 5.2]) these strata are exactly the ones to which Theorem 11.5 does not apply.

We shall finish this section by showing that the 1-skeleton graph can be described in purely arithmetic terms very strongly reminiscent of the results Harashita. Note that even for final elements our results are not formally equivalent to Harashita's as we are dealing with the set of components of the final strata in \mathcal{F}_g whereas Harashita is dealing with their images in \mathcal{A}_q . In any case our counting of the

number of components uses 11.4 and hence is valid only for sufficiently large p whereas Harashita's are true unconditionally.

We start by giving a well-known description of the vertices of the 1-skeleton graph (see for instance [Ek87]) valid when q > 1. We fix a super-singular elliptic curve E and its endomorphism ring \mathbf{D} which is provided with the Rosati involution *. To simplify life we assume, as we may, that E is defined over \mathbb{F}_p and hence **D** contains the Frobenius map F. It has the property that $\mathbf{D}F = F\mathbf{D} = \mathbf{D}f\mathbf{D}$, the unique maximal ideal containing p. Furthermore we have that $\mathbf{D}/\mathbf{D}F \cong \mathbb{F}_{p^2}$. There is then a bijection between isomorphism classes of \mathbf{D} -lattices M (i.e., right modules torsion free and finitely generated as abelian groups) of rank g (i.e., of rank 4g as abelian groups) and q-dimensional abelian varieties A. The correspondence associates to the abelian variety A the **D**-module $\operatorname{Hom}(E, A)$. Polarisations on A then correspond to positive definite unitary forms, i.e., a bi-additive map $\langle -, - \rangle \colon M \times M \to \mathbf{D}$ such that $\langle md,n\rangle = \langle m,n\rangle dn, \ \langle n,m\rangle = (\langle m,n\rangle)^*, \ \text{and} \ m\neq 0 \implies \langle m,m\rangle > 0.$ The polarisation is principal precisely when the form is *perfect*, i.e., the induced map of right **D**-modules $M \to \operatorname{Hom}_{\mathbf{D}}(M, \mathbf{D})$ given $n \mapsto (m \mapsto \langle m, n \rangle)$ is an isomorphism. In general we put $M^* := \operatorname{Hom}_{\mathbf{D}}(M, \mathbf{D})$ and then the form induces an embedding $M \to M^*$ which makes the image of finite index. More precisely we on $M \bigotimes \mathbb{Q}$ we get a non-degenerate pairing with values in $\mathbf{D} \bigotimes \mathbb{Q}$ and then we may identify M^* with the set $\{n \in M \bigotimes \mathbb{Q} : \forall m \in M : \langle m, n \rangle \in \mathbf{D} \}$. Using this we get a $\mathbf{D} \bigotimes \mathbb{Q}/\mathbb{Z}$ valued unitary perfect form on M^*/M given by $\langle \overline{m}, \overline{n} \rangle := \langle m, n \rangle \mod \mathbf{D}$. As usual **D**-valued super-lattices $M \subseteq N$ corresponds to totally isotropic submodules of $M^*/M.$

If now $S \subseteq \{0, \ldots, g\}$ is stable under $i \mapsto g-i$ then an *arithmetic S-flag* consists of the choice of unitary forms on **D**-modules M_i of rank g for $i \in S$ and compatible isometric embeddings $M_i \hookrightarrow M_j$ whenever i < j fulfilling the following conditions:

- For all $i \in S$ with $i \geq g/2$ we have that M_i^*/M_i is killed by F of \mathbf{D} and can hence be considered as an $\mathbf{D}/m = \mathbb{F}_{p^2}$ -vector space with a perfect unitary form.
- We have that $FM_i^* = M_{g-i}$ for all $i \in S$.
- The length of M_j/M_i for i < j is equal to j i.

Remark 11.7. i) Note that we allow S to be empty in which case there is exactly one S-flag.

ii) As follows (implicitly) from the proof of next proposition, the isomorphism class of an element of an arithmetic S-flag tensored with \mathbb{Q} is independent of the flag. Hence we may consider only lattices in a fixed unitary form over $\mathbf{D} \otimes \mathbb{Q}$ and then think of M_i as a sublattice of M_j .

Proposition 11.8. Let $S \subseteq \{1, \ldots, g\}$ and let $\overline{S} \subseteq \{0, \ldots, g\}$ be the set of integers of the form i or g - i for $i \in S$. Then the set of isomorphism classes of \overline{S} -flags is in bijection with the set of the set of connected components of the S-subgraph of the 1-skeleton graph.

Proof. This follows from the discussion above and Theorem 7.5 once we have proven that an S-flag can be extended to an $\{0, \ldots, g\}$ -flag. Assume first that $g \notin S$ and let $i \in S$ be the largest element in S. By assumption we have $FM_i^* = M_{g-i} \subseteq M_i \subset M_i^*$ and we have that the length of M_i^*/FM_i^* is g whereas by assumption that of M_i/M_{g-i} is 2i - g. Together this gives that the length of M_i^*/M_i is g - (2i - g) = 2(g-i). As the form on M_i^*/M_i is a non-degenerate unitary \mathbb{F}_{p^2} -form and as all such

forms are equivalent we get that there is a g - i-dimensional totally isotropic (and hence its own orthogonal) subspace of M_i^*/M_i which then gives an $M_i \subset M_g \subset M_i^*$ and as M_g/M_i is its own orthogonal we get that the pairing on M_g is perfect. We then put $M_0 := FM_g$ and the rest of extension is immediate.

12. The Cycle Classes

If one wishes to exploit our stratification on \mathcal{F}_g and the E-O stratification on \mathcal{A}_g fully, one needs to know the cohomology classes (or Chow classes) of the (closed) strata. In this section we show how to calculate these classes. The original idea for the determination of the cycle classes can be illustrated well by the *p*-rank strata. If X is a principally polarized abelian variety of dimension g which is generic in the sense that its *p*-rank is g, then its kernel of multiplication by p contains a direct sum of g copies of μ_p , the multiplicative group scheme of order p. Viewing μ_p as a fat point at the origin we get a tangent vector to X at the origin, and after dualizing, a section of the Hodge bundle. If the abelian variety looses p-rank under specialization the g sections thus obtained become dependent and the loci where this happens have classes represented by a multiple of the Chern classes of the Hodge bundle.

To calculate the cycle classes of the E-O strata on $\mathcal{A}_g \otimes \mathbb{F}_p$ we shall use the theory of degeneration cycles of maps between vector bundles. To this end we shall apply formulas of Fulton for degeneracy loci of symplectic bundle maps to calculate the classes of the $\overline{\mathcal{U}}_w$ and formulas of Pragacz and Ratajski and of Kresch and Tamvakis for calculating those of $\overline{\mathcal{V}}_{\nu}$.

12.1. Fulton's formulas. Over the flag space \mathcal{F}_g we have the pull back of the de Rham bundle and the two flags \mathbb{E}_{\bullet} and \mathbb{D}_{\bullet} on it. We denote by ℓ_i the roots of the Chern classes of \mathbb{E} so that $c_1(\mathbb{E}_i) = \ell_1 + \ell_2 + \ldots + \ell_i$. We then have $c_1(\mathbb{D}_{g+i}) - c_1(\mathbb{D}_{g+i-1}) = p\ell_i$.

Recall that for each element $w \in W_g$ we have a degeneracy locus $\overline{\mathcal{U}}_w$ in \mathcal{F}_g and $\tilde{\mathcal{F}}_g$. Its codimension equals the length $\ell(w)$ and it thus makes sense to consider the cycle class $u_w = [\overline{\mathcal{U}}_w]$ in $\mathrm{CH}^{\mathrm{codim}(w)}(\tilde{\mathcal{F}}_g)$, where we write \mathcal{F}_g instead of $\mathcal{F}_g \otimes \mathbb{F}_p$.

Fulton's setup in [Fu96] is the following (or more precisely the part that interests us): We have a symplectic vector bundle H over some scheme X and two full symplectic flags $0 \subset \cdots \subset E_2 \subset E_1 = H$ and $0 \subset \cdots \subset D_2 \subset D_1 = H$. For each $w \in W_g$ on defines the degeneracy locus $\overline{\mathcal{U}}_w$ by $\{x \in X : \forall i, j: \dim \mathbb{E}_{i,x} \cap \mathbb{D}_{j,x} \leq r_w(i,j)\}$ (of course this closed subset is given a scheme structure by considering these conditions as rank conditions for maps of vector bundles). Fulton then defines a polynomial in two sets of variables x_i and y_j , $i, j = 1, \ldots, g$, such that if this polynomial is evaluated as $x_i = c_1(E_i/E_{i+1})$ and $y_j = c_1(D_j/D_{j+1})$ then it gives the class of $\overline{\mathcal{U}}_w$ provided that $\overline{\mathcal{U}}_w$ has the expected codimension $\operatorname{codim}(w)$ (and Xis Cohen-Macaulay). The precise definition of these polynomials are as follows: For a partition $\mu = \{\mu_1 > \mu_2 > \ldots > \mu_r > 0\}$ with $r \leq g$ and $\mu_1 \leq g$ one defines a Schur function

$$\Delta_{\mu}(x) := \det(x_{\mu_i+j-i})_{1 \le i,j \le r}$$

in the variables x_i and puts

$$\Delta(x,y) := \Delta_{(g,g-1,\ldots,1)}(\sigma_i(x_1,\ldots,x_g) + \sigma_i(y_1,\ldots,y_g)),$$

where the σ_i is the *i*-th elementary symmetric function. One then considers the "divided difference operators" ∂_i on the polynomial ring $\mathbb{Z}[x_1, \ldots, x_q]$ by

$$\partial_i(F(x)) = \begin{cases} \frac{F(x) - F(s_i x)}{x_i - x_{i+1}} & \text{if } i < g, \\ \frac{F(x) - F(s'_g x)}{2x_g} & \text{if } i = g, \end{cases}$$

where s_i interchanges x_i and x_{i+1} for $i = 1, \ldots, g-1$, but s'_g sends x_g to $-x_g$ and leaves the other x_i unchanged. We write an element $w \in W_g$ as a product $w = s_{i_\ell} s_{i_{\ell-1}} \ldots s_{i_1}$ with $\ell = \ell(w)$ and set

(12.1)
$$P_w := \partial_{i_\ell} \cdots \partial_{i_1} (\prod_{i+j \le g} (x_i - y_j) \cdot \Delta).$$

An application of Fulton's formulas gives the following.

Theorem 12.1. Let $w = s_{i_{\ell}} s_{i_{\ell-1}} \dots s_{i_1}$ with $\ell = \ell(w)$ be an element of the Weyl group W_g . Then the cycle class $u_w := [\overline{\mathcal{U}}_w]$ in $\operatorname{CH}_{\mathbb{Q}}^{\operatorname{codim}(w)}(\tilde{\mathcal{F}}_g)$ is given by

$$u_w = \partial_{i_1} \cdots \partial_{i_\ell} \left(\prod_{i+j \le g} (x_i - y_j) \cdot \Delta(x, y) \right)_{|x_i = -\ell_i, y_j = p\ell_j}$$

Proof. By construction $\overline{\mathcal{U}}_w$ is the degeneracy locus of the flags \mathbb{E}_{\bullet} and \mathbb{D}_{\bullet} . By Corollary 8.4 they have the expected dimension and hence the degeneracy cycle class is equal to the class of $\overline{\mathcal{U}}_w$.

For a final element $w \in W_g$ the map $\overline{\mathcal{U}}_w \to \overline{\mathcal{V}}_w$ is generically finite of degree $\gamma_g(w)$. By applying the Gysin map to the formula of the theorem using formula 3.1 we can in principle calculate the cohomology classes of all the push-downs of final strata, hence of the E-O strata.

Example 12.2. g = 2.

The Weyl group W_2 consists of 8 elements; we give the cycle classes in $\tilde{\mathcal{F}}_2$ and the push-downs on $\tilde{\mathcal{A}}_2$.

 w	s	ℓ	$[\overline{\mathcal{U}}_w]$	$\pi_*([\overline{\mathcal{U}}_w])$
f 1				
[4,3]	$s_1 s_2 s_1 s_2$	4	1	0
[4, 2]	$s_1 s_2 s_1$	3	$(p-1)\lambda_1$	0
[3, 4]	$s_2 s_1 s_2$	3	$-\ell_1 + p\ell_2$	1+p
[2, 4]	$s_{1}s_{2}$	2	$(1-p)\ell_1^2 + (p^2 - p)\lambda_2$	$(p-1)\lambda_1$
[3, 1]	$s_{2}s_{1}$	2	$(1-p^2)\ell_1^2 + (1-p)\ell_1\ell_2 + (1-p)\ell_2^2$	$p(p-1)\lambda_1$
[2, 1]	s_1	1	$(p-1)(p^2+1)\lambda_1\lambda_2$	0
[1, 3]	s_2	1	$(p^2 - 1)\ell_1^2(\ell_1 - p\ell_2)$	$(p-1)(p^2-1)\lambda_2$
[1, 2]	1	0	$-(p^4-1)\lambda_1\lambda_2\ell_1$	$(p^4-1)\lambda_1\lambda_2$

In the flag space \mathcal{F}_2 the stratum corresponding to the empty diagram is $U_{s_2s_1s_2}$ and the strata contained in its closure are the four final ones $U_{s_2s_1s_2}$, $U_{s_1s_2}$, U_{s_2} and U_1 and the two non-final ones U_{s_1} and $U_{s_2s_1}$. The Bruhat-Chevalley order on these is given by the diagram:



The four final strata $U_{s_2s_1s_2}$, $U_{s_1s_2}$, U_{s_2} and U_1 lie étale of degree 1 over the *p*-rank 2 locus, the *p*-rank 1 locus of abelian surfaces with *p*-rank 0 and *a*-number 1, and the locus of superspecial surfaces (a = 2). The locus U_{s_1} is an open part of the fibres over the superspecial points. The locus $U_{s_2s_1}$ is of dimension 2 and lies finite but inseparably of degree *p* over the *p*-rank 1 locus. Then E_1 corresponds to an α_p and E_2/E_1 to a μ_p . In the final type locus $U_{s_1s_2}$ the fibrration is $\mu_p \subset \mu_p \oplus \alpha_p$. Note that this description is compatible with the calculated classes of the loci.

We have implemented the calculation of the Gysin map in Macaulay2 (cf., [M2]) and calculated all cycle classes for $g \leq 5$. For g = 3, 4 the reader will find the classes in Appendix. (The Macaulay2 code for performing the calculations can be found at http://www.math.su.se/~teke/strata.m2.) We shall return to the qualitative consequences one can draw from Theorem 12.1 in the next section.

12.2. The *p*-rank strata. It is very useful to have closed formulas for the cycle classes of important strata. We give the formulas for the strata defined by the *p*-rank and by the *a*-number. The formulas for the *p*-rank strata can be derived immediately from the definition of the strata.

Let V_f be the closed E-O stratum of \mathcal{A}_g of semi-abelian varieties of p-rank $\leq f$. It has codimension g - f. To calculate its class we consider the element w_{\emptyset} , the longest final element. The corresponding locus $\overline{\mathcal{U}}_{\emptyset}$ is a generically finite cover of \mathcal{A}_g of degree $\gamma_g(w_{\emptyset}) = \prod_{i=1}^{g-1} (p^i + p^{i-1} + \ldots + 1)$. The map of \mathcal{U}_{\emptyset} to the p-rank g locus is finite. The space $\overline{\mathcal{U}}_{\emptyset}$ contains the degeneracy loci \mathcal{U}_w for all final elements $w \in W_g$. The condition that a point x of \mathcal{F}_g lies in $\overline{\mathcal{U}}_{\emptyset}$ is that the filtration \mathbb{E}_i for $i = 1, \ldots, g$ is stable under V. By forgetting part of the flag and considering flags \mathbb{E}_i with $j = i, \ldots, g$ we find that $\overline{\mathcal{U}}_{\emptyset} \to \mathcal{A}_g$ is fibered by generically finite morphisms

$$\overline{\mathcal{U}}_{\emptyset} = \overline{\mathcal{U}}^{(1)} \xrightarrow{\pi_1} \overline{\mathcal{U}}^{(2)} \xrightarrow{\pi_2} \dots \xrightarrow{\pi_{g-1}} \overline{\mathcal{U}}^{(g)} = \mathcal{A}_g.$$

We shall write $\pi_{i,j}$ for the composition $\pi_j \pi_{j-1} \cdots \pi_i : \overline{\mathcal{U}}^{(i)} \to \overline{\mathcal{U}}^{(j)}$ and $\pi_{\emptyset} = \pi_{1,g}$. Since V_{g-1} is given by the vanishing of the map $\det(V) : \det(\mathbb{E}_g) \to \det(\mathbb{E}^{(p)})$ the

Since V_{g-1} is given by the vanishing of the map $\det(V)$: $\det(\mathbb{E}_g) \to \det(\mathbb{E}^{(p)})$ the class of V_{g-1} is $(p-1)\lambda_1$. The pull-back of V_{g-1} to $\overline{\mathcal{U}}_{\emptyset}$ decomposes in g irreducible components

$$\pi_{\emptyset}^{-1}(V_{g-1}) = \bigcup_{i=1}^{g} Z_i,$$

where Z_i is the degeneracy locus of the induced map $\phi_i = V_{|\mathcal{L}_i} \colon \mathcal{L}_i \to \mathcal{L}_i^{(p)}$. Note that the Z_i are the $\overline{\mathcal{U}}_w$ for the w which are shuffles of the final element u_{g-1} (see subsection 4.2) defining the E-O stratum of p-rank f and Z_g is the stratum corresponding to the element $u_{g-1} \in W_g$. An abelian variety of p-rank g-1 and a-number 1 has a unique subgroup scheme α_p . The index i of Z_i indicates where this subgroup scheme can be found (i.e. its Dieudonné module lies in \mathbb{E}_i , but not in \mathbb{E}_{i-1}).

It follows from the definition of Z_i as degeneracy set that the class of Z_i on $\overline{\mathcal{U}}_{\emptyset}$ equals $(p-1)\ell_i$ as ϕ_i can be interpreted as a section of $\mathcal{L}_i^{(p)} \otimes \mathcal{L}_i^{-1}$. We also know by 4.3 that the map $Z_i \to Z_{i+1}$ is inseparable. Therefore $\pi_{\emptyset}([Z_i]) = p^{n(i)}\pi_{\emptyset}([Z_g])$ for some integer $n(i) \ge g - i$. Using the fact that $(\pi_{\emptyset})_*([Z_g]) = \gamma_g(u_{g-1})[V_{g-1}] =$ $\deg(\pi_{1,g-1})[V_{g-1}]$ we see that

$$(\pi_{\emptyset})_*(\pi_{\emptyset}^*([V_{g-1}])) = \sum_{i=1}^g (\pi_{\emptyset})_*(Z_i) = \sum_{i=1}^g p^{n(i)} \deg(\pi_{1,g-1})[V_{g-1}],$$

while on the other hand

$$(\pi_{\emptyset})_*(\pi_{\emptyset}^*([V_{g-1}])) = \deg(\pi_{\emptyset})[V_{g-1}] = (1+p+\ldots+p^{g-1})\deg(\pi_{1,g-1})[V_{g-1}].$$

Comparison yields that n(i) = g - i and so we find

$$(\pi_{\emptyset})_*(\ell_i) = p^{g-i} \operatorname{deg}(\pi_{1,g-1})\lambda_1$$

and

$$(\pi_{\emptyset})_*([Z_i]) = (p-1)p^{g-i} \deg(\pi_{1,g-1})\lambda_1$$

Lemma 12.3. In the Chow groups with rational coefficients of $\overline{\mathcal{U}}^{(i)}$ and $\overline{\mathcal{U}}^{(i+1)}$ we have for the push down of the j-th Chern class $\lambda_j(i)$ of \mathbb{E}_i the relations:

$$\pi_*^i \lambda_j(i) = p^j (p^{i-j} + p^{i-j-1} + \dots + p + 1) \lambda_j(i+1)$$

and

$$p^{f(g-f)}(\pi_{1,g})_*(\ell_g\ell_{g-1}\cdots\ell_{f+1}) = (\pi_{1,g})_*(\ell_1\ell_2\cdots\ell_{g-f}).$$

Proof. The relation $(\pi_1)_*([Z_1]) = p[Z_2]$ translates into the case j = 1 and i = 1. Using the push-pull formula and the relations $\pi_i^*(\lambda_j(i+1) = \ell_{i+1}\lambda_{j-1}(i) + \lambda_j(i))$ the formulas for the push downs of the $\lambda_j(i)$ follow by induction on j and i. \Box

We now calculate the class of all *p*-rank strata V_f .

Theorem 12.4. The class of the locus V_f of semi-abelian varieties of p-rank $\leq f$ in the Chow ring $CH_{\mathbb{Q}}(\tilde{\mathcal{A}}_q)$ equals

$$[V_f] = (p-1)(p^2 - 1) \cdots (p^{g-f} - 1) \lambda_{g-f}$$

Proof. The class of the final stratum $\overline{\mathcal{U}}_{u_f}$ on $\overline{\mathcal{U}}_{\emptyset}$ is given by the formula

$$(p-1)^{g-f}\ell_g\ell_{g-1}\cdots\ell_{f+1}$$

as it is the simultaneous degeneracy class of the maps ϕ_j for $j = f + 1, \ldots, g$. By pushing down this class under $\pi_{\emptyset} = \pi_{1,g}$ we find using 12.3 and the notation $\lambda_j(i) = c_j(\mathbb{E}_i)$ that

$$(\pi_{1,g})_*(\ell_g\ell_{g-1}\cdots\ell_{f+1}) = p^{-f(g-f)}(\pi_{1,g})_*(\ell_1\ell_2\cdots\ell_{g-f})$$

= $p^{-f(g-f)}(\pi_{1,g})_*(\pi_{1,g-f}^*(\lambda_{g-f}(g-f)))$
= $p^{-f(g-f)} \deg(\pi_{1,g-f})(\pi_{g-f,g})_*(\lambda_{g-f}(g-f)).$

Applying Lemma 12.3 repeatedly we find

$$(\pi_{g-f,g})_*(\lambda_{g-f}(g-f)) = p^{f(g-f)}(1+p)(1+p+p^2)\cdots(1+\ldots+p^{f-1})\lambda_g$$
$$= p^{f(g-f)}\gamma_g(u_f)\lambda_g,$$

with $\gamma_g(u_f)$ the number of final filtrations refining the canonical filtration of u_f . Hence we get $(\pi_{\emptyset})_*(\overline{\mathcal{U}}_{u_f}) = (p-1)^{g-f} \deg(\pi_{1,g-f})\gamma_g(u_f)\lambda_g$. On the other hand, we have that $(\pi_{\emptyset})_*(\overline{\mathcal{U}}_{u_f}) = \gamma_g(u_f)[V_f]$. Together this proves the result. \Box

12.3. The *a*-number Strata. Another case where we can find attractive explicit formulas is that of the E-O strata \mathcal{V}_w with w the element of W_g associated to $Y = \{1, 2, \ldots, a\}$. We denote these by T_a . Here we can work directly on \mathcal{A}_g . The locus T_a on \mathcal{A}_g may be defined as the locus $\{x \in \mathcal{A}_g : \operatorname{rank}(V)|_{\mathbb{E}_g} \leq g - a\}$. We have $T_{a+1} \subset T_a$ and $\dim(T_g) = 0$. We apply now formulas of Pragacz and Ratajski, cf. [PR97] for the degeneracy locus for the rank of a self-adjoint bundle map of symplectic bundles globalizing the results in isotropic Schubert calculus from [Pr91]. Before we apply their result to our case we have to introduce some notation.

Define for a vector bundle A with Chern classes a_i the expression

$$Q_{ij}(A) := a_i a_j + 2 \sum_{k=1}^{j} (-1)^k a_{i+k} a_{j-k}$$
 for $i > j$.

A subset $\beta = \{g \ge \beta_1 > \ldots > \beta_r \ge 0\}$ of $\{1, 2, \ldots, g\}$ (with r even, note that β_r may be zero) is called admissible and for such subsets we set

$$Q_{\beta} = \text{Pfaffian}(x_{ij}),$$

where the matrix (x_{ij}) is anti-symmetric with entries $x_{ij} = Q_{\beta_i,\beta_j}$. Applying the formula of Pragacz-Ratajski to our situation gives the following result:

Theorem 12.5. The cycle class $[T_a]$ of the reduced locus T_a of abelian varieties with a-number $\geq a$ is given by

$$[T_a] = \sum_{\beta} Q_{\beta}(\mathbb{E}^{(p)}) \cdot Q_{\rho(a)-\beta}(\mathbb{E}^*),$$

where the sum is over the admissible subsets β contained in the subset $\rho(a) = \{a, a - 1, a - 2, ..., 1\}.$

Example 12.6.

$$[T_1] = (p-1)\lambda_1$$

$$[T_2] = (p-1)(p^2+1)\lambda_1\lambda_2 - (p^3-1)2\lambda_3$$

$$[T_3] =$$

...

$$[T_g] = (p-1)(p^2+1)\dots(p^g+(-1)^g)\lambda_1\lambda_2\dots\lambda_g.$$

As a corollary we find a result of one of us (cf. [Ek87]) on the number of principally polarized abelian varieties with a = g.

Corollary 12.7. We have

$$\sum_{X} \frac{1}{\#\operatorname{Aut}(X)} = (-1)^{g(g+1)/2} 2^{-g} \left[\prod_{j=1}^{g} (p^j + (-1)^j) \right] \cdot \zeta(-1)\zeta(-3) \dots \zeta(1-2g),$$

where the sum is over the isomorphism classes (over \mathbb{F}_p) of principally polarized abelian varieties of dimension g with a = g, and $\zeta(s)$ is the Riemann zeta function.

Proof. Combine the formula for T_g with the Hirzebruch-Mumford Proportionality Theorem (see [Ge99]) which says that

$$\deg(\lambda_1\lambda_2\cdots\lambda_g)=(-1)^{\frac{g(g+1)}{2}}\prod_{j=1}^g\frac{\zeta(1-2j)}{2},$$

when interpreted for the stack \mathcal{A}_q .

The formulas for the cycles classes of the *p*-rank strata and the *a*-number strata can be seen as generalizations of the classical formula of Deuring (known as Deuring's Mass Formula) which states that

$$\sum_{E} \frac{1}{\#\operatorname{Aut}(E)} = \frac{p-1}{24},$$

where the sum is over the isomorphism classes over \mathbb{F}_p of supersingular elliptic curves. It is obtained from the formula for V_{g-1} or T_1 for g = 1, i.e., $[V_1] = (p-1)\lambda_1$, by remarking that the degree of λ_1 is 1/12 times the degree of a generic point of the stack $\tilde{\mathcal{A}}_1$.

One can obtain formulas for all the E-O strata by applying the formulas of Pragacz-Ratajski or those of Kresch-Tamvakis [KT02, Cor. 4]. If Y is a Young diagram given by a subset $\{\xi_1, \ldots, \xi_r\}$ we call $|\xi| = \sum_{i=1}^r \xi_i$ the weight and r the length of ξ . Moreover, we need the excess $e(\xi) = |\xi| - r(r+1)/2$ and the *intertwining number* $e(\xi, \eta)$ of two strict partitions with $\xi \cap \eta = \emptyset$ by

$$e(\xi,\eta) = \sum_{i\geq 1} i \,\#\{j:\xi_i > \eta_j > \xi_{i+1}\}$$

(where we use $\xi_k = 0$ if k > r). We put $\rho_g = \{g, g - 1, ..., 1\}$ and $\xi' = \rho_g - \xi$ and have then $e(\xi, \xi') = e(\xi)$. The formula obtained by applying the result of Kresch and Tamvakis interpolates between the formulas for the two special cases, the *p*-rank strata and *a*-number strata, as follows:

Theorem 12.8. For a Young diagram given by a partition ξ we have

$$[\overline{\mathcal{V}}_Y] = (-1)^{e(\xi) + |\xi'|} \sum_{\alpha} Q_{\alpha}(\mathbb{E}^{(p)}) \sum_{\beta} (-1)^{e(\alpha,\beta)} Q_{(\alpha \cup \beta)'}(\mathbb{E}^*) \det(c_{\beta_i - \xi'_j}(\mathbb{E}^*_{g - \xi'_j})),$$

where the sum is over all admissible α and all admissible β that contain ξ' with length $\ell(\beta) = \ell(\xi')$ and $\alpha \cap \beta = \emptyset$.

12.4. Positivity of tautological classes. The Hodge bundle possesses certain positivity properties. It is well-known that the determinant of the Hodge bundle (represented by the class λ_1) is ample on \mathcal{A}_g . Over \mathbb{C} this is a classical result, while in positive characteristic this was proven by Moret-Bailly [MB85]. On the other hand, the Hodge bundle itself is not positive in positive characteristic. For example, for g = 2 the restriction of \mathbb{E} to a line from the *p*-rank 0 locus is $O(-1) \oplus O(p)$, [MB81]. But our results 12.4 imply the following non-negativity result.

Theorem 12.9. The Chern classes $\lambda_i \in CH_{\mathbb{Q}}(\mathcal{A}_g \otimes \mathbb{F}_p)$ (i = 1, ..., g) of the Hodge bundle \mathbb{E} are represented by effective classes with \mathbb{Q} -coefficients.

13. TAUTOLOGICAL RINGS

We shall now interpret the results of previous sections in terms of tautological rings. Recall that the tautological ring of \mathcal{A}_g is the subring of $\mathrm{CH}_{\mathbb{Q}}(\mathcal{A}_g)$ generated by the Chern classes λ . To obtain maximal precision we shall use the subring and not the Q-subalgebra (but note that is still a subring of $CH_{\mathbb{Q}}(\bar{\mathcal{A}}_q)$ not of the integral Chow ring $CH^*(\mathcal{A}_q)$). As a graded ring it is isomorphic to the Chow ring $\operatorname{CH}^*(\operatorname{Sp}_{2q}/P_H)$ and as an abstract graded ring it is generated by the λ_i with relations coming from the identity $1 = (1 + \lambda_1 + \dots + \lambda_g)(1 - \lambda_1 + \dots + (-1)^g \lambda_g).$ This implies that it has a \mathbb{Z} -basis consisting of the square free monomials in the λ_i . (Note however, that the degree maps from the degree g(g+1)/2 part are not the same; on $\operatorname{Sp}_{2g}/P_H$ the degree of $\lambda_1 \dots \lambda_g$ is ± 1 , whereas for \mathcal{A}_g it is given by the Hirzebruch-Mumford proportionality theorem as in the previous section.) As $\mathcal{F}_q \to \mathcal{A}_q$ is an SL_q/B -bundle we can express $\mathrm{CH}_{\mathbb{Q}}(\mathcal{F}_q)$ as an algebra over $CH_{\mathbb{Q}}(\dot{\mathcal{A}}_{q})$; it is the algebra generated by the ℓ_{i} and the relations are that the elementary symmetric functions in them are equal to the λ_i . This makes it natural to define the *tautological ring* of \mathcal{F}_g to be the subring of $\mathrm{CH}_{\mathbb{Q}}(\mathcal{F}_g)$ generated by the ℓ_i . It will then be the algebra over tautological ring of \mathcal{A}_q generated by the ℓ_i and with the relations that say that the elementary symmetric functions in the ℓ_i are equal to the λ_i . Again this means that the tautological ring for \mathcal{F}_g is isomorphic to the integral Chow ring of $\operatorname{Sp}_{2g}/B_g$, the space of full symplectic flags in a 2gdimensional symplectic vector space. Note furthermore that the Gysin maps for $\operatorname{Sp}_{2g}/B_g \to \operatorname{Sp}_{2g}/P_H$ and $\tilde{\mathcal{F}}_g \to \tilde{\mathcal{A}}$ are both given by Formula 3.1.

Theorem 12.1 shows in particular that the classes of the $\overline{\mathcal{U}}_w$ and $\overline{\mathcal{V}}_{\nu}$ lie in the respective tautological rings. However, we want to both compare the formulas for these classes with the classical formulas for the Schubert varieties and take into account the variation of the coefficients of the classes when expressed in a fixed basis for the tautological ring. To do this we introduce the ring $\mathbb{Z}\{p\}$ which is the localisation of the polynomial ring $\mathbb{Z}[p]$ at the multiplicative subset of polynomials with constant coefficient equal to 1. Hence evaluation at 0 extends to a ring homomorphism $\mathbb{Z}\{p\} \to \mathbb{Z}$ which we shall call the *classical specialisation*. Hence, an element of $\mathbb{Z}\{p\}$ is invertible precisely when its classical specialisation is invertible. By a modulo n consideration we see that an integer polynomial with 1 as constant coefficient can have no integer zero $n \neq \pm 1$. That means in particular that evaluation at a prime p induces a ring homomorphism $\mathbb{Z}\{p\} \to \mathbb{Q}$ taking the variable p to the integer p (this dual use of p should hopefully not cause confusion) which we shall call the *characteristic* p specialisation. We now extend scalars of the two tautological rings from \mathbb{Z} to $\mathbb{Z}\{p\}$ and we shall call them the *p*-tautological rings. We shall also need to express the condition that an element is in the subring obtained by extension to a subring of $\mathbb{Z}\{p\}$ and we shall then say that the element has *coefficients* in the subring. We may consider the Fulton polynomial P_w of (12.1) as a polynomial with coefficients in $\mathbb{Z}\{p\}$ and when we evaluate them on elements of the tautological ring as in Theorem 12.1 we get elements $[\overline{\mathcal{U}}_w]$ of the *p*-tautological ring of \mathcal{F}_q . If ν is a final element we can push down the formula for $[\overline{\mathcal{U}}_{\nu}]$ using (3.1) and then we get an element in the *p*-tautological ring of \mathcal{A}_{g} . We then note that $\gamma(w)$ is a polynomial in p with constant coefficient equal to 1 and hence we can define $[\overline{\mathcal{V}}_{\nu}] := \gamma(w)^{-1} \pi_*[\overline{\mathcal{U}}_{\nu}]$, where $\pi \colon \tilde{\mathcal{F}}_g \to \tilde{\mathcal{A}}_g$ is the projection map. By construction these elements map to the classes of $\overline{\mathcal{U}}_w$ resp. $\overline{\mathcal{V}}_{\nu}$ under specialisation to characteristic p. We shall need to compare them with the classes of the Schubert varieties. To be specific we shall define the Schubert varieties of $\operatorname{Sp}_{2g}/B_g$ by the condition dim $E_i \cap D_j \geq r_w(i, j)$, where D_{\bullet} is a fixed reference flag (and then the Schubert varieties of $\operatorname{Sp}_{2g}/P_H$ are the images of the Schubert varieties of $\operatorname{Sp}_{2g}/B_g$ for final elements of W_g).

Theorem 13.1. i) The classes $[\overline{\mathcal{U}}_w]$ and $[\overline{\mathcal{V}}_\nu]$ map to the classes of the corresponding Schubert varieties under classical specialisation.

ii) The classes $[\overline{\mathcal{U}}_w]$ and $[\overline{\mathcal{V}}_\nu]$ form a $\mathbb{Z}\{p\}$ -basis for the respective p-tautological rings.

iii) The coefficients of $[\overline{\mathcal{U}}_w]$ and $[\overline{\mathcal{V}}_\nu]$ are in $\mathbb{Z}[p]$.

iv) For $w \in W_g$ we have that $\ell(w) = \ell(\tau_p(w))$ (see Section 9 for the definition of τ_p) precisely when the specialisation to characteristic p of $\pi_*[\overline{\mathcal{U}}_w]$ is non-zero. In particular there is a unique map $\tau \colon W_g \to W_g/S_g \bigcup \{0\}$ such that $\tau(w) = 0$ precisely when $\pi_*[\overline{\mathcal{U}}_w] = 0$ which implies that $\ell(w) \neq \ell(\tau_p(w))$ and is implied by $\ell(w) \neq \ell(\tau_p(w))$ for all sufficiently large p. Furthermore if $\tau(w) \neq 0$ then $\ell(w) =$ $\ell(\tau(w))$ and $\pi_*[\overline{\mathcal{U}}_w]$ is a non-zero multiple of $[\overline{\mathcal{V}}_{\tau(w)}]$.

Proof. The first part is clear as putting p = 0 gives the Fulton formulas for $x_i = -\ell_i$ and $y_i = 0$ which are the Fulton formulas for the Schubert varieties in $\operatorname{Sp}_{2g}/B_g$. One then obtains the formulas for the Schubert varieties of $\operatorname{Sp}_{2g}/P_H$ by pushing down by Gysin formulas. The remaining compatibility needed is that the classical specialisation of $\gamma(w)$ is the degree of the map from the Schubert variety of $\operatorname{Sp}_{2g}/B_g$ for a final element to the corresponding Schubert variety of $\operatorname{Sp}_{2g}/P_H$. However, the classical specialisation of $\gamma(w)$ is 1 and the map between Schubert varieties is an isomorphism between Bruhat cells.

As for the second part we need to prove that the determinant of the matrix expressing the classes of the strata in terms of a basis of the tautological ring (say given by monomials in the ℓ_i resp. the λ_i) is invertible. Given that an element of $\mathbb{Z}\{p\}$ is invertible precisely when its classical specialisation is, we are reduced to proving the corresponding statement in the classical case. However, there it follows from the cell decomposition given by the Bruhat cells which give that the classes of the Schubert cells form a basis for the integral Chow groups.

To prove iii) it is enough to verify the conditions of Proposition 13.2. Hence let $w \in W_g$ resp. a final element ν and consider a an element m in the tautological ring of degree complementary to that of $[\overline{\mathcal{U}}_w]$ resp. $[\overline{\mathcal{V}}_\nu]$. By the projection formula the degree of $m[\overline{\mathcal{U}}_w]$ resp. $m[\overline{\mathcal{V}}_\nu]$ is the degree of the restriction of m to $\overline{\mathcal{U}}_w$ resp. $\overline{\mathcal{V}}_\nu$ and it is enough to show that the denominators of these degrees are only divisible by a finite number of primes (independently of the characteristic p). However, if the characteristic is different from 3 we may pull back to the moduli space with a level 3 structure and there the degree is an integer as the corresponding strata are schemes. Hence the denominator divides the degree of the level 3 structure covering $\tilde{\mathcal{A}}_{g,3} \to \tilde{\mathcal{A}}_g$ which is independent of p.

Finally for iv), it is clear that in the Chow ring of $\overline{\mathcal{A}}_g$ we have that $\pi_*[\overline{\mathcal{U}}_w]$ is non-zero precisely when $\pi: [\overline{\mathcal{U}}_w] \to \overline{\mathcal{V}}_{\tau_p(w)}$ is generically finite as all fibres have the same dimension by Proposition 9.6. This latter fact also gives that it is generically finite precisely when $\overline{\mathcal{U}}_w$ and $\overline{\mathcal{V}}_{\tau_p(w)}$ have the same dimension which is equivalent to $\ell(w) = \ell(\tau_p(w))$. When this is the case we get that $\pi_*[\overline{\mathcal{U}}_w]$ is a non-zero multiple of $[\overline{\mathcal{V}}_{\tau_p(w)}]$ again as the degree over each component of $\overline{\mathcal{V}}_{\tau_p(w)}$ is the same by Proposition 9.6. Consider now instead $\pi_*[\overline{\mathcal{U}}_w]$ in the *p*-tautological ring and expand $\pi_*[\overline{\mathcal{U}}_w]$ as a linear combination of the $[\overline{\mathcal{V}}_\nu]$. Then what we have just shown is that for every specialisation to characteristic *p* at most one of the coefficients are non-zero. This implies that in the *p*-tautological ring at most one of the coefficients is non-zero. If it is zero then $\pi_*[\overline{\mathcal{U}}_w]$ is always zero in all characteristic *p* specialisations and we get $\ell(w) \neq \ell(\tau_p(w))$ for all *p*. If it is non-zero, then the coefficient is non-zero for all sufficiently large *p*. This proves iv).

To complete the proof of the theorem we need to prove the following proposition.

Proposition 13.2. Let a be an element of the p-tautological ring for $\tilde{\mathcal{F}}_g$ or $\tilde{\mathcal{A}}_g$. Assume that there exists an $n \neq 0$ such that for all elements b of the tautological ring of complementary degree and all sufficiently large primes p we have that $\deg(\overline{ab}) \in \mathbb{Z}[n^{-1}]$, where \overline{a} and \overline{b} are the specialisations to characteristic p of a resp. b. Then the coefficients of a are in $\mathbb{Z}[p]$.

Proof. If r(x) is one of the coefficients of a, then the assumptions say that $r(p) \in \mathbb{Z}[n^{-1}]$ for all sufficiently large primes p. Write r as g(x)/f(x) where f and g are integer polynomials with no common factor. Thus there are integer polynomials s(x) and t(x) such that s(x)f(x) + t(x)g(x) = m, where m is a non-zero integer. If g is non-constant there are arbitrarily large primes ℓ such that there is an integer k such that $\ell|f(k)$ (by for instance the fact that there is a prime which splits completely in a splitting field of f). By Dirichlet's theorem on primes in arithmetic progressions there are arbitrarily large primes p such that $f(p) \neq 0$ and $f(p) \equiv f(k) \equiv 0 \mod \ell$. By making ℓ so large so that $\ell \not\mid m$ we get that $\ell \not\mid g(q)$ (as s(q)f(q) + t(q)g(q) = m) and hence ℓ appears in the denominator of r(q). By making ℓ so large so that $\ell \not\mid n$ we conclude.

Example 13.3. If w is a shuffle of a final element ν we have $\tau(w) = \nu$.

We can combine this theorem with our results on the punctual flag spaces to give an algebraic criterion for inclusion between E-O strata.

Corollary 13.4. Let ν' and ν be final elements. Then for sufficiently large p we have that $\nu' \subseteq \nu$ if there are $w, w' \in W_g$ for which $w' \leq w$, $\tau(w) = \nu$, and $\tau(w') = \nu'$.

Proof. Assume there are $w, w' \in W_g$ for which $w' \leq w, \tau(w) = \nu$, and $\tau(w') = \nu'$. By Proposition 9.6 we have that for $\pi \colon \tilde{\mathcal{F}}_g \to \tilde{\mathcal{A}}_g$ the following image relations $\pi(\overline{\mathcal{U}}_w) = \overline{\mathcal{V}}_{\nu_1}$ and $\pi(\overline{\mathcal{U}}_{w'}) = \overline{\mathcal{V}}_{\nu'_1}$ for some ν_1 and ν'_1 and by the theorem $\nu_1 = \tau(w)$ and $\nu'_1 = \tau(w')$ for sufficiently large p. As $w' \leq w$ we have that $\overline{\mathcal{U}}_{w'} \subseteq \overline{\mathcal{U}}_w$ which implies that $\pi \overline{\mathcal{U}}_{w'} \subseteq \pi(\overline{\mathcal{U}}_w)$.

14. Comparison with $\mathcal{S}(g, p)$.

We shall now make a comparison with de Jong's moduli stack of $\Gamma_0(p)$ -structures (cf, [Jo93]), $\mathcal{S}(g, p)$. Recall that for a family $\mathcal{A} \to S$ of principally polarised gdimensional abelian varieties a $\Gamma_0(p)$ -structure consists of the choice of a flag $0 \subset$ $H_1 \subset \cdots \subset H_g \subset \mathcal{A}[p]$ of flat subgroup schemes with H_i of order p^i and H_g totally isotropic with respect to the Weil pairing. We shall work exclusively in characteristic p and denote by $\overline{\mathcal{S}(g,p)}$ the mod p fibre of $\mathcal{S}(g,p)$. We now let $\mathcal{S}(g,p)^0$ be the closed subscheme of $\mathcal{S}(g,p)$ defined by the condition that H_g is of height 1, i.e., that the (relative) Frobenius map, $F_{\mathcal{A}/\mathcal{S}(g,p)}$ where $\pi \colon \mathcal{A} \to \mathcal{S}(g,p)$ is the universal abelian variety, on it is zero. For degree reasons we then get that H_g equals the kernel of $F_{\mathcal{A}/\mathcal{S}(g,p)}$. Using the principal polarisation we may identify the Lie algebra of π with $R^1\pi_*\mathcal{O}_{\mathcal{A}}$ and hence we get a flag $0 \subset \text{Lie}(H_1) \subset$ $\text{Lie}(H_2) \subset \cdots \subset \text{Lie}(H_g) = R^1\pi_*\mathcal{O}_{\mathcal{A}}$. By functoriality this is stable under V. Completing this flag by taking its annihilator in \mathcal{E} gives a flag in $\overline{\mathcal{U}}_{w_{\emptyset}}$ thus giving a map $\mathcal{S}(g,p)^0 \to \overline{\mathcal{U}}_{w_{\emptyset}}$.

Theorem 14.1. The canonical map $S(g,p)^0 \to \overline{\mathcal{U}}_{w_{\emptyset}}$ is an isomorphism. In particular, $S(g,p)^0$ is the closure of its intersection with the locus of ordinary abelian varieties and is normal and Cohen-Macaulay.

Proof. Starting with the tautological flag $\{\mathbb{E}_i\}$ on $\overline{\mathcal{U}}_{w_{\emptyset}}$ we consider the induced flag $\{\mathbb{E}_{g+i}/\mathbb{E}_g\}$ in $R^1\pi_*\mathcal{O}_{\mathcal{A}}$. This is a V-stable flag of the Lie algebra of a height 1 group scheme so by, for instance [Mu70, Thm. §14], any V-stable subbundle comes from a subgroup scheme of the kernel of $F_{\mathcal{A}/\mathcal{S}(g,p)}$ and thus the flag $\{\mathbb{E}_{g+i}/\mathbb{E}_g\}$ gives rise to a complete flag of subgroup schemes with H_g equal to the kernel of the Frobenius map and hence a map from $\overline{\mathcal{U}}_{w_{\emptyset}}$ to $\mathcal{S}(g,p)^0$ which clearly is the inverse of the canonical map.

The rest of the theorem now follows from Corollary 8.4.

15. Appendix g = 3

15.1. Admissible Strata for g = 3. In the following matrix one finds the loci lying in $\overline{\mathcal{U}}_{w_{\emptyset}}$. In the sixth column we give for each w an example of a final ν such that $w \to \nu$.

ℓ	Y	w	u	word	$w \to \nu$	
0	$\{1, 2, 3\}$	[123]	$\{0, 0, 0\}$	Id	[123]	Superspecial
1		[132]	$\{0, 0, 0\}$	s_2	[123]	Fibre over s.s.
1		[213]	$\{0, 0, 0\}$	s_1	[123]	Fibre over s.s.
1	$\{2, 3\}$	[124]	$\{0, 0, 1\}$	s_3	[124]	Moret-Bailly
2		[142]	$\{0, 0, 1\}$	$s_{3}s_{2}$	[135]	
2		[214]	$\{0, 0, 1\}$	s_3s_1	[135]	
2		[231]	$\{0, 0, 0\}$	$s_{1}s_{2}$	[123]	Fibre over s.s.
2		[312]	$\{0, 0, 0\}$	$s_{2}s_{1}$	[123]	Fibre over s.s.
2	$\{1, 3\}$	[135]	$\{0, 1, 1\}$	$s_{2}s_{3}$	[135]	f = 0, a = 2
3		[153]	$\{0, 1, 1\}$	$s_2 s_3 s_2$	[236]	Shuffle of $\{1, 2\}$
3		[241]	$\{0, 0, 1\}$	$s_3 s_1 s_2$	[124]	
3		[315]	$\{0, 1, 1\}$	$s_2 s_3 s_1$	[124]	
3		[321]	$\{0, 0, 0\}$	$s_1 s_2 s_1$	[123]	Fibre over s.s.
3		[412]	$\{0, 0, 1\}$	$s_3 s_2 s_1$	[236]	Shuffle of $\{1, 2\}$
3	$\{3\}$	[145]	$\{0, 1, 2\}$	$s_3 s_2 s_3$	[145]	f = 0
3	$\{1, 2\}$	[236]	$\{1, 1, 1\}$	$s_1 s_2 s_3$	[236]	a=2
4		[154]	$\{0, 1, 2\}$	$s_3 s_2 s_3 s_2$	[246]	Shuffle of $\{2\}$
4		[326]	$\{1, 1, 1\}$	$s_1 s_2 s_3 s_1$	[236]	
4		[351]	$\{0, 1, 1\}$	$s_2 s_3 s_1 s_2$	[236]	
4		[415]	$\{0, 1, 2\}$	$s_3 s_2 s_3 s_1$	[246]	Shuffle of $\{2\}$
4		[421]	$\{0, 0, 1\}$	$s_3 s_1 s_2 s_1$	[236]	
4	$\{2\}$	[246]	$\{1, 1, 2\}$	$s_3 s_1 s_2 s_3$	[246]	f = 1
5		[426]	$\{1, 1, 2\}$	$s_3 s_1 s_2 s_3 s_1$	[356]	Shuffle of $\{1\}$
5		[451]	$\{0, 1, 2\}$	$s_3 s_2 s_3 s_1 s_2$	[356]	Shuffle of $\{1\}$
5	$\{1\}$	[356]	$\{1, 2, 2\}$	$s_2 s_3 s_1 s_2 s_3$	[356]	f = 2
6	{}	[456]	$\{1, 2, 3\}$	$s_3s_2s_3s_1s_2s_3$	[456]	f = 3

15.2. **E-O Cycle Classes for** g = 3. We give the cycle classes of the (reduced) E-O strata for g = 3.

Y	class
đ	1
\emptyset {1}	$(p-1)\lambda_1$
${2}{1}$	$(p-1)\lambda_1 (p-1)(p^2-1)\lambda_2$
$\{1,2\}$	$(p-1)(p^2+1)\lambda_1\lambda_2 - 2(p^3-1)\lambda_3$
{3}	$(p-1)(p^2-1)(p^3-1)\lambda_3$
$\{1, 3\}$	$(p-1)^2(p^3+1)\lambda_1\lambda_3$
$\{2,3\}$	$(p-1)^2(p^6-1)\lambda_2\lambda_3$
$\{1, 2, 3\}$	$(p-1)(p^2+1)(p^3-1)\lambda_1\lambda_2\lambda_3$

15.3. E-O Cycle Classes for $g = 4$.	We give the cycle classes of the (reduced)
E-O strata for $g = 4$.	

Y	class
Ø	1
{1}	$(p-1)\lambda_1$
$\{2\}$	$(p-1)(p^2-1)\lambda_2$
$\{1, 2\}$	$(p-1)(p^2+1)\lambda_1\lambda_2-2(p^3-1)\lambda_3$
{3}	$(p-1)(p^2-1)(p^3-1)\lambda_3$
$\{1,3\}$	$(p-1)^2(p+1)((p^2-p+1)\lambda_1\lambda_3-2(p^2+1)\lambda_4)$
$\{2,3\}$	$(p-1)^{2}((p^{6}-1)\lambda_{2}\lambda_{3}-(2p^{6}+p^{5}-p-2)\lambda_{1}\lambda_{4})$
$\{1, 2, 3\}\$ $\{4\}$	$ (p-1)(p^2+1)((p^3+1)((p^3-1)\lambda_1\lambda_2\lambda_3 - 2(3p^3+p^2-p+3)\lambda_2\lambda_4) (p-1)(p^2-1)(p^3-1)(p^4-1)\lambda_4 $
$\{1,4\}$	$(p-1)(p-1)(p-1)(p-1)\lambda_4$ $(p-1)^3(p+1)(p^4+1)\lambda_1\lambda_4$
$\{2,4\}$	$(p-1)^3(p^8-1)\lambda_2\lambda_4$
$\{1, 2, 4\}$	$(p-1)^2(p^4-1)((p^2+1)\lambda_1\lambda_2 - 2(p^2+p+1)\lambda_3)\lambda_4$
$\{3, 4\}$	$(p-1)^2(p^2+1)(p^3-1)(p^2-p+1)((p+1)^2\lambda_3-p\lambda_1\lambda_2)\lambda_4$
$\{1, 3, 4\}$	$(p-1)^2(p^4-1)(p^6-1)\lambda_1\lambda_3\lambda_4$
$\{2, 3, 4\}$	$(p-1)(p^6-1)(p^8-1)\lambda_2\lambda_3\lambda_4$
$\{1, 2, 3, 4\}$	$(p-1)(p^2+1)(p^3-1)(p^4+1)\lambda_1\lambda_2\lambda_3\lambda_4$

References

- [BGG73] I. N. Bernstein, I. M. Gelfand, and S. I. Gelfand, Schubert cells, and the cohomology of the spaces G/P, Uspehi Mat. Nauk 28 (1973), no. 3(171), 3–26.
- [BL00] S. Billey and V. Lakshmibai, Singular loci of Schubert varieties, Progress in Mathematics, vol. 182, Birkhäuser Boston Inc., Boston, MA, 2000.
- [Ch94] C. Chevalley, Sur les décompositions cellulaires des espaces G/B, Algebraic groups and their generalizations: classical methods (University Park, PA, 1991), Proc. Sympos. Pure Math., vol. 56, Amer. Math. Soc., Providence, RI, 1994, With a foreword by Armand Borel, pp. 1–23.
- [Ek87] T. Ekedahl, On supersingular curves and abelian varieties, Math. Scand. 60 (1987), 151–178.
- [FC90] G. Faltings and C.-L. Chai, Degeneration of abelian varieties., Ergebnisse der Math., no. 22, Springer Verlag, 1990.
- [FP98] W. Fulton and P. Pragacz, Schubert varieties and degeneracy loci, SLN, vol. 1689, Springer-Verlag, Berlin, 1998, Appendix J by the authors in collaboration with I. Ciocan-Fontanine.
- [Fu96] W. Fulton, Determinantal formulas for orthogonal and symplectic degeneracy loci, J. Differential Geom. 43 (1996), no. 2, 276–290.
- [Ful] _____, Intersection theory, vol. 3, Ergebnisse der Mathematik und ihrer Grenzgebiete, no. 2, Springer Verlag, 1984.
- [Ge99] G. van der Geer, Cycles on the moduli space of abelian varieties, Moduli of curves and abelian varieties (C. Faber and E. Looijenga, eds.), Aspects Math., E33, Vieweg, Braunschweig, 1999, pp. 65–89.
- [GrLie4-6] N. Bourbaki, Groupes et algèbres de Lie. IV-VI, Hermann, Paris, 1968.
- [HW61] H. Hasse and E. Witt, Zyklische unverzweigte Erweiterungskörper von primzahlgrade p über einem algebraischen Funktionenkörper der Charakteristik p, Monatsh. Math. Phys. 43 (1963), 477–492.
- [Ha04] $\mathbf{S}.$ Harashita, Ekedahl-Oortcontainedthestratainsupersingularlocus, Tech. Report 15,University \mathbf{of} Tokyo, 2004,http://faculty.ms.u-tokyo.ac.jp/users/preprint/preprint2004.html.

- [Jo93] A. J. de Jong, The moduli spaces of principally polarized abelian varieties with $\Gamma_0(p)$ level structure, J. of Algebraic Geom. **2** (1993), no. 4, 667–688.
- [KS03] S. Keel and L. Sadun, *Oort's conjecture for* $A_g \otimes \mathbb{C}$, J. Amer. Math. Soc. **16** (2003), no. 4, 887–900 (electronic).
- [KT02] A. Kresch and H. Tamvakis, Double Schubert polynomials and degeneracy loci for the classical groups, Ann. Inst. Fourier (Grenoble) 52 (2002), no. 6, 1681–1727.
- [M2] D. R. Grayson and M. E. Stillman, Macaulay 2, a software system for research in algebraic geometry, Available at http://www.math.uiuc.edu/Macaulay2.
- [MB81] L. Moret-Bailly, Familles de courbes et variétés abéliennes sur P¹., Séminaire sur les pinceaux de courbes de genre au moins deux. (L. Szpiro, ed.), Astérisque, vol. 86, Soc. math. de France, 1981, pp. 109−140.
- [MB85] _____, Pinceaux de variétés abéliennes, Astérisque (1985), no. 129, 266.
- [MW04] B. Moonen and T. Wedhorn, Discrete invariants of varieties in positive characteristic, IMRN (2004).
- [Mo01] B. Moonen, Group schemes with additional structures and Weyl group cosets, Moduli of abelian varieties (Texel Island, 1999), Progr. Math., vol. 195, Birkhäuser, Basel, 2001, pp. 255–298.
- [Mu70] D. Mumford, Abelian varieties, Tata Institute of Fundamental Research Studies in Mathematics, vol. 5, Oxford University Press., London, 1970.
- [Oo01] F. Oort, A stratification of a moduli space of abelian varieties, Moduli of abelian varieties (Texel Island, 1999), Progr. Math., vol. 195, Birkhäuser, Basel, 2001, pp. 345– 416.
- [Oo95] _____, Complete subvarieties of moduli spaces, Abelian varieties (Egloffstein, 1993), de Gruyter, Berlin, 1995, pp. 225–235.
- [PR97] P. Pragacz and J. Ratajski, Formulas for Lagrangian and orthogonal degeneracy loci; Q-polynomial approach, Compositio Math. 107 (1997), no. 1, 11–87.
- [PR99] H. Pittie and A. Ram, A Pieri-Chevalley formula in the K-theory of a G/B-bundle, Electron. Res. Announc. Amer. Math. Soc. 5 (1999), 102–107 (electronic).
- [Pr91] P. Pragacz, Algebro-geometric applications of Schur S- and Q-polynomials, Topics in invariant theory (Paris, 1989/1990), Lecture Notes in Math., vol. 1478, Springer, Berlin, 1991, pp. 130–191.
- [RR85] S. Ramanan and A. Ramanathan, Projective normality of flag varieties and Schubert varieties, Invent. Math. 79 (1985), no. 2, 217–224.
- [S] N. Sloane, The on-line encyclopedia of integer sequences, http://www.research.att.com/~njas/sequences.

Department of Mathematics, Stockholm University, SE-106 91 Stockholm, Sweden E-mail address: teke@math.su.se

FACULTEIT WISKUNDE EN INFORMATICA, UNIVERSITY OF AMSTERDAM, PLANTAGE MUIDER-GRACHT 24, 1018 TV AMSTERDAM, THE NETHERLANDS

E-mail address: geer@science.uva.nl