# ENDOMORPHISMS OF RELATIVELY HYPERBOLIC GROUPS

### IGOR BELEGRADEK AND ANDRZEJ SZCZEPAŃSKI

ABSTRACT. We generalize some results of Paulin and Rips-Sela on endomorphisms of hyperbolic groups to relatively hyperbolic groups, and in particular prove the following.

• If G is a finitely generated non-elementary relatively hyperbolic group with slender parabolic subgroups, and either G is not co-Hopfian or Out(G) is infinite, then G splits over a slender group.

• If a finitely generated non-parabolic subgroup H of a non-elementary relatively hyperbolic group is not Hopfian, then H acts non-trivially on an  $\mathbb{R}$ -tree.

• Any finitely presented group is isomorphic to a finite index subgroup of Out(H) for some Kazhdan group H. (This sharpens a result of Ollivier-Wise).

#### 1. INTRODUCTION

The Bestvina-Paulin method [Pau88, Pau91, Bes88] further developed by Sela has been a key ingredient in much of recent work on endomorphisms of hyperbolic groups [Pau91, RS94, Sel95, Sel97, Sel99], on endomorphisms of geometrically finite Kleinian groups [Sel97, OP98, DP03], and in Sela's work on Tarski problem [Sel02].

Very recently, Groves [Gro] generalized various results on endomorphisms of hyperbolic groups, notably the Sela's shortening argument, to relatively hyperbolic groups with finitely generated free abelian parabolic subgroups.

The point of this paper is to show that some of the easier applications of the Bestvina-Paulin method extend, with little effort, to relatively hyperbolic groups with fairly arbitrary parabolic subgroups. By "easier" we mean the results that do not require JSJ-decompositions or the shortening argument.

Relatively hyperbolic groups were introduced by Gromov [Gro87] and since then various characterizations of relatively hyperbolic groups have been obtained by

<sup>2000</sup> Mathematics Subject classification. Primary 20F65. Keywords: relatively hyperbolic, automorphism group, Hopfian, co-Hopfian, Kazhdan, splitting, actions on trees.

Farb [Far98], Bowditch [Bow], Yaman [Yam04], Osin [Osic], Drutu-Sapir [DS]. See also [Szc98, Dah03a, Bum].

In this paper we follow Gromov's definition of a relatively hyperbolic group, as developed by Bowditch. More precisely, a group G is hyperbolic relative to a family of subgroups  $\mathcal{G}$  if G acts properly discontinuously and isometrically on a proper, geodesic, hyperbolic metric space X so that G acts on the ideal boundary of X as a geometrically finite convergence group whose maximal parabolic subgroups are precisely the elements of  $\mathcal{G}$ . We do not assume that maximal parabolic subgroups are finitely generated. We refer to [Bow, Bow99, Yam04] for background.

A subgroup of a relatively hyperbolic group G is called *elementary* if the limit set of its action as a convergence group on the Bowditch's boundary of Gcontains at most two points, which happens exactly if the subgroup is finite, virtually- $\mathbb{Z}$ , or parabolic. Otherwise, the subgroup is called *non-elementary*. The Tits Alternative for convergence groups [Tuk94] implies that any small subgroup of a relatively hyperbolic group is elementary, where a subgroup is called *small* if it contains no non-cyclic free subgroup.

Examples of relatively hyperbolic groups are:

- the free products of finitely many (arbitrary) groups are hyperbolic relative to the factors (because the free product action on the corresponding Bass-Serre tree satisfies the Bowditch's definition [Bow] of a relatively hyperbolic group).
- hyperbolic groups are hyperbolic relative to any conjugacy invariant collection  $\mathcal{G}$  of quasi-convex subgroups such that any element of  $\mathcal{G}$  is equal to its normalizer, and any two distinct elements of  $\mathcal{G}$  have finite intersection [Bow, Theorem 7.11].
- geometrically finite isometry groups of Hadamard manifolds of negatively pinched sectional curvature as defined in [Bow95] are hyperbolic relative to maximal parabolic subgroups. This includes complete finite volume manifolds of negatively pinched sectional curvature.
- Osin [Osid] developed small cancellation theory for relatively hyperbolic groups, and proved that small cancellation quotients of relatively hyperbolic groups are hyperbolic relative to the images of maximal parabolic subgroups. Osin's methods imply [Osia] that any non-elementary relatively hyperbolic group has a Kazhdan quotient that is non-elementary relatively hyperbolic.
- combinations theorems for relatively hyperbolic groups were proved in [Dah03b, Osib], e.g. the amalgamation of relatively hyperbolic groups over parabolic subgroups is relatively hyperbolic, when the parabolic subgroup is maximal in at least one of the factors.

- Kleiner-Hruska [HK] proved that CAT(0)-groups with isolated flats are hyperbolic relative to the flat stabilizers. Examples of CAT(0)-groups with isolated flats are listed in [HK].
- Sela's limit group are hyperbolic relative to non-cyclic maximal abelian subgroups. This was first proved in [Dah03b] using a combination theorem, and in fact according to [AB] limit groups are CAT(0)-groups with isolated flats.

The main observation is as follows.

**Theorem 1.1.** Let H be a finitely generated group and G be a non-elementary relatively hyperbolic group.

(i) If  $\rho_k \colon H \to G$  is an arbitrary sequence of pairwise G-non-conjugate homomorphisms whose images are not parabolic, then H admits a nontrivial action on an  $\mathbb{R}$ -tree T.

(ii) Furthermore, if all  $\rho_k$  are injective, and  $H_a$  is an arbitrary arc stabilizer of the H-action on T, then each finitely generated subgroup of  $H_a$  is isomorphic to a elementary subgroup of G.

Proof of Theorem 1.1. Let (X, d) be a proper, geodesic,  $\delta$ -hyperbolic metric space from the definition of a relative hyperbolic group. Choose a fundamental domain F for the G-action on X. Fix a finite generating set S for H. For  $x \in X$  we let

$$\mu_k(x) = \max_{s \in S} d(x, \rho_k(s)(x)),$$

and let  $\mu_k := \inf_{x \in X} \mu_k(x)$ . Choose  $x_k \in X$  with  $\mu_k(x_k) \leq \mu_k + \frac{1}{k}$ . First we show that  $\mu_k$  has no bounded subsequence. Arguing by contradiction, consider a bounded subsequence, which we still denote  $\mu_k$ . Let  $R = 1 + \sup_k \mu_k$ . Choose a *G*-invariant system of horoballs in *X* such that any two horoballs are at least *R* apart. There exists a sequence  $g_k \in G$  such that  $g_k(x_k) \in F$ . If the sequence  $g_k(x_k)$  is precompact, then since *G* acts properly discontinuously,  $g_k \rho_k g_k^{-1}$  fall into finitely many *G*-conjugacy classes, which contradicts the assumption. If  $g_k(x_k)$  is not precompact, then after passing to a subsequence  $g_k(x_k)$  lie in a horoball *H*, and since horoballs are at least *R* apart,  $g_k \rho_k(s) g_k^{-1}(g_k(x_k)) \in H$ for each  $s \in S$ . Hence  $g_k \rho_k(H) g_k^{-1}$  lies in the parabolic subgroup that stabilizes *H*, which contradicts the assumption that  $\rho_k(H)$  is not parabolic.

Then by the work of Bestvina-Paulin (see e.g. Sections 3.1-3.5 of [Bes02]), the pointed spaces  $(\frac{1}{\mu_k}X, x_k)$  subconverges to an  $\mathbb{R}$ -tree T, and  $\rho_k$ 's converge to a non-trivial isometric H-action on T, which proves (i).

Assume  $\rho_k$  are all injective. Let *a* be an arc of *T* with the stabilizer  $H_a$  in *H*. Let  $\check{H}_a$  be the subgroup of  $H_a$  of index at most 2 that fixes *a* pointwise. Approximate a by geodesic segments  $a_k$  in  $\frac{1}{\mu_k}X$ , denote the midpoint of  $a_k$  by  $m_k$ . Changing  $\rho_k$  within its G-conjugacy class, we can assume  $m_k \in F$ .

By [Pau91, p. 341] (cf. [BS94, p. 284]), there exists a constant  $C(\delta)$ , such that for each  $h, f \in \check{H}_a$  and all large k, we have  $d(m_k, \rho_k([h, f])(m_k) < C(\delta)$ .

First, we prove that if  $m_k$  is precompact in F, then  $H_a$  is small, hence elementary. Indeed, since  $m_k$  is precompact, only finitely elements of G can satisfy  $d(m_k, g(m_k)) < C(\delta)$ , because G acts isometrically and properly discontinuously. We denote the number of such elements by M. If  $H_a$  is not small, then  $\check{H}_a$  contains a rank two free subgroup generated by  $h_1$ ,  $h_2$ . By the previous paragraph, for all large enough k,  $d(m_k, \rho_k([h_1, h_2^s])(m_k) < C(\delta)$  for  $s = 1, \ldots, M + 1$ . So  $\rho_k([h_1, h_2^{s_1}] = \rho_k([h_1, h_2^{s_2}])$  for some  $s_1 \neq s_2$ . Hence  $\rho_k([h_1, h_2^{s_1-s_2}]) = 1$ , which is impossible since  $\rho_k$  is injective and  $h_1, h_2$  generate a free subgroup.

Now assume that  $m_k$  is not precompact. Fix a *G*-invariant system of horoballs in *X* such that any two horoballs are at least  $C(\delta) + 1$  apart. Passing to a subsequence we can assume that  $m_k$  belong to one horoball. Denote by *P* the maximal parabolic subgroup of *G* stabilizing this horoball, and let  $p \in \partial X$  be the unique fixed point of *P*. By above, for each  $h, f \in \check{H}_a$  and all large *k*, we have  $\rho_k([h, f])$ ,  $\rho_k([f^{-1}, h])$ ,  $\rho_k([f, h^{-1}])$  belong to *P*. Since  $h^{-1}[h, f]h = [f, h^{-1}]$  and  $f^{-1}[h, f]f = [f^{-1}, h]$  and  $\rho_k$  is injective, we conclude that  $\rho_k(h)(p) = p = \rho_k(f)(p)$ , hence  $\rho_k(h), \rho_k(f) \in P$  for all large *k*. So any finitely generated subgroup of  $\check{H}_a$  is mapped into *P* by  $\rho_k$  for large *k* in the subsequence.

Finally, given an arbitrary finitely generated subgroup  $\Gamma$  of  $H_a$ , let  $\dot{\Gamma} := \Gamma \cap \dot{H}_a$ ; the index of  $\check{\Gamma}$  in  $\Gamma$  is  $\leq 2$  so  $\Gamma$  normalizes  $\check{\Gamma}$ . By the previous paragraph  $\rho_k(\check{\Gamma}) \subset P$  for some k. Since  $\rho_k(\Gamma)$  normalizes  $\rho_k(\check{\Gamma})$ , we get  $\rho_k(\Gamma) \subset P$ , as promised.

**Remark 1.2.** For example, if parabolic subgroups of G are small, then each  $H_a$  is small (and hence finite, virtually- $\mathbb{Z}$  or parabolic). The same is true if small is replaced by amenable, slender, finitely generated virtually nilpotent, finitely generated virtually abelian, etc. Also if parabolic subgroups of G are free, and G is torsion-free, then each  $H_a$  is locally free.

### 2. Endomorphisms and actions on $\mathbb{R}$ -trees

Theorem 1.1 immediately implies the following corollary, which for hyperbolic groups is due to Paulin [Pau91].

**Corollary 2.1.** If a finitely generated non-elementary relatively hyperbolic group G has infinite Out(G), then G acts non-trivially on an  $\mathbb{R}$ -tree such that any

finitely generated subgroup of each arc stabilizer is isomorphic to an elementary subgroup of G.

A group is called *co-Hopfian* if every injective endomorphism of the group is surjective.

**Corollary 2.2.** Suppose G is a finitely generated non-elementary relatively hyperbolic group that contains no infinite torsion group and is not isomorphic to a parabolic subgroup of G. If G is not co-Hopfian, then G acts non-trivially on an  $\mathbb{R}$ -tree such that any finitely generated subgroup of each arc stabilizer is isomorphic to an elementary subgroup of G.

Corollary 2.2 is deduced from Theorem 1.1, by repeating the proof of Rips-Sela [RS94, Theorems 3.1] where the same result was proved for hyperbolic groups (in which case all parabolic subgroups are trivial, and the assumption "finitely generated, contains no infinite torsion group, and is not isomorphic to a parabolic subgroup" holds automatically). Later Sela [Sel97] proved that a torsion-free hyperbolic group is co-Hopfian if and only if it is freely indecomposable. A similar characterization of co-Hopfian geometrically finite Kleinian groups was obtained in [DP03].

**Remark 2.3.** We do not know whether the assumption in Corollary 2.2 that "G contains no infinite torsion group" can be dropped. What is actually used in the proof is that if  $\phi$  is a non-surjective monomorphism of  $G \to G$  and  $A_k$  is the (necessarily finite) centralizer of the subgroup  $\phi^k(G)$ , then the torsion group  $A = \bigcup_k A_k$  is finite.

The assumption "G is not isomorphic to a parabolic subgroup of G" in Corollary 2.2 holds e.g. if parabolic subgroups are small and G is non-elementary, or if parabolic subgroups are co-Hopfian. This assumption cannot be dropped.

**Example 2.4.** There exists a non-elementary relatively hyperbolic group that is Kazhdan and not co-Hopfian.

Proof. Osin's small cancellation methods allow to construct relatively hyperbolic Kazhdan groups with prescribed parabolic groups. Specifically, if H be a finitely generated group and K be a torsion-free hyperbolic Kazhdan group, then  $G := H * \mathbb{Z} * K$  is hyperbolic relative to the factors H,  $\mathbb{Z}$ . If follows from properties of free products that K is a suitable subgroup of G (see [Osid] for a definition). By [Osid, Theorem 2.4] there is an epimorphism  $\eta \colon G \to \overline{G}$ such that  $\eta(K) = \overline{G}$ , the restriction of  $\eta$  to  $H \cup \mathbb{Z}$  is injective,  $\overline{G}$  is hyperbolic relative to  $\eta(H)$ ,  $\eta(\mathbb{Z})$ , and  $\overline{G}$  is obtained from G by adding finitely many relations. In particular,  $\overline{G}$  is Kazhdan and non-elementary, and furthermore if H is finitely presented, then so is  $\overline{G}$ . Now suppose H is a Higman's universal finitely presented group [Hig61], i.e. H is a finitely presented group such that any finitely presented group embeds into H. Then  $\overline{G}$  embeds into  $H \cong \eta(H) \subset \overline{G}$ , hence  $\overline{G}$  is not co-Hopfian.

**Remark 2.5.** Corollaries 2.1, 2.2 generally fail for finitely generated subgroups of hyperbolic groups: a non-co-Hopfian finitely generated subgroup of a hyperbolic group was constructed in [KW01], and by [OW] there exist Kazhdan groups that have infinite outer automorphism groups yet are embedded into hyperbolic group. However, the examples in [KW01, OW] are not finitely presented, and it seems no finitely presented examples are known.

## 3. Splitting over slender groups

A group is called *slender* if all its subgroups are finitely generated. The class of slender group is closed under extensions [DS99], in particular, virtually polycyclic groups are slender. Slender groups are small because non-abelian free groups contain infinitely generated subgroups.

If G is a relatively hyperbolic group with slender parabolic subgroups, then small subgroups of G are slender. Then since in any group an ascending chain of small subgroups is small, every ascending chain of slender subgroups of G stabilizes. Hence, any nontrivial minimal G-action on an  $\mathbb{R}$ -tree with slender arc stabilizers is stable [BF95], and therefore [BF95] if G is finitely presented, then there exists a nontrivial G-action on a simplicial tree slender edge stabilizers, i.e. a splitting of G over a slender subgroup. (Note that the splitting is nontrivial provided G is non-elementary). In fact, according to Dunwoody [Dun, Theorem 2], the assumption "G is finitely presented" can be replaced by "G is finitely generated". Thus combining [Dun, Theorem 2] with Corollaries 2.1, 2.2, we get the results stated in the abstract about co-Hopf property and infinite outer automorphism group, as well as the following.

**Corollary 3.1.** Let H be a finitely generated group that is not slender, and let G be a relatively hyperbolic group with slender parabolic subgroups. If  $\rho_k: H \to G$  is an arbitrary sequence of pairwise G-non-conjugate injective homomorphisms, then H admits a nontrivial splitting over a slender group.

**Remark 3.2.** It is interesting to compare Corollary 3.1 with a result of Dahmani [Dah] who showed that if H is a finitely presented group and G is a relatively hyperbolic group, then up to G-conjugacy there are only finitely many subgroups of G that are non-parabolic, do not split over parabolic subgroups, and are the images of homomorphisms  $H \to G$ .

A group H is said to have property  $F\mathbb{R}$  if any H-action on an  $\mathbb{R}$ -tree fixes a point, i.e. is not *non-trivial*. By Theorem 1.1 if H is finitely generated and has property  $F\mathbb{R}$ , then there are only finitely many G-conjugacy classes of homomorphisms  $H \to G$  with non-parabolic images.

The class of groups with property  $F\mathbb{R}$  is closed under finite extensions [Ser77] and quotients, and contains

- Kazhdan groups (this was proved in [Wat82] for simplicial trees, and extended to  $\mathbb{R}$ -trees in [Nos93]);
- Aut $(F_n)$  and Out $(F_n)$  for n > 2 [CV96], where  $F_n$  is a free group of rank n;
- mapping class groups  $M_{q,r}$  for  $g \ge 2$  [CV96];
- finitely generated Coxeter groups with no  $\infty$  labels in their Coxeter diagrams (this is an exercise in [Ser77, p. 93]),
- fundamental groups of closed orientable irreducible non-Haken 3-manifolds. See [MS88, Proposition 2.1], where it is also proved that if a finitely generated orientable 3-manifold group acts non-trivially on an ℝ-tree, it acts non-trivially on an simplicial tree,
- small groups that do not surject onto  $\mathbb{Z}$ ,  $\mathbb{Z}_2 * \mathbb{Z}_2$ . (This is well-known, but we could not find a reference so here is a proof. If a small group G acts nontrivially on an  $\mathbb{R}$ -tree, then G stabilizes a line, or fixes an end, as proved e.g. in [Nos93]. In the former case, G surject onto  $\mathbb{Z}$ , or  $\mathbb{Z}_2 * \mathbb{Z}_2$ . In the latter case, the minimal G-subtree has an abelian hyperbolic length function, and the function is the absolute value of a homomorphism  $G \to \mathbb{R}$ , which is nontrivial because the G-action is nontrivial. Thus in this case G surjects onto  $\mathbb{Z}$ .)

Recall that group is called *Hopfian* if every surjective endomorphism of the group is injective.

**Corollary 4.1.** If a finitely generated non-parabolic subgroup H of a relatively hyperbolic group has property  $F\mathbb{R}$ , then H is Hopfian.

Proof. If  $\rho$  is a non-bijective epimorphism of H, then precomposing the inclusion  $H \hookrightarrow G$  with the powers  $\rho^k$ , we get the homomorphisms  $H \to G$  with image H, which is non-parabolic by assumption. To show that  $\rho$  is injective, it is enough to prove that some power of  $\rho$  is injective. Look at the sequence  $\psi_k := \rho^{2^k}$ , so that for any two endomorphisms in the sequence  $\{\psi_k\}$  the one with the larger index is a power of the other. By Theorem 1.1, after passing to a subsequence, we can assume that  $\psi_k$ 's are conjugate in G. In particular, if  $\psi$  denotes the first endomorphism in the subsequence, we get  $\psi^{s_k} = i_{q_k} \circ \psi$ ,

where  $g_k \in G$  and  $i_{g_k}$  is the corresponding inner automorphism of G. Then  $\psi^{s_k}$  and  $\psi$  have equal kernels. Take  $\gamma \in \ker(\psi^{s_k-1})$ . Since  $\psi$  is onto, we can find  $\tilde{\gamma} \in H$  with  $\psi(\tilde{\gamma}) = \gamma$ . So  $\tilde{\gamma} \in \ker(\psi^{s_k}) = \ker(\psi)$ , which implies  $\gamma = 1$ , as wanted.

That hyperbolic groups with property  $F\mathbb{R}$  are Hopfian was noted to [RS94, Theorem 2.1], and Corollary 4.1 extends this result to relatively hyperbolic case. Later Sela [Sel99] showed that any torsion-free hyperbolic group is Hopfian, and more recently Bumagin proved [Bum04] that any finitely generated subgroup of a torsion-free hyperbolic group is Hopfian. Thus the assumption "H has property  $F\mathbb{R}$ " in Corollary 4.1 is probably far from optimal.

Ollivier-Wise [OW] constructed a non-Hopfian Kazhdan group. Since Kazhdan groups are finitely generated and have property  $F\mathbb{R}$ , we conclude by Corollary 4.1 that their group cannot be embedded in a relatively hyperbolic group as a non-parabolic subgroup.

## 5. Automorphism groups of Kazhdan groups

Ollivier-Wise [OW] showed that any finitely presented group is isomorphic to the quotient of a torsion-free hyperbolic group G by a normal subgroup H that is Kazhdan. In case G is non-elementary, the canonical homomorphism  $G/H \rightarrow \text{Out}(H)$  is injective, thus Ollivier-Wise concluded that any finitely presented group embeds into Out(H) for some Kazhdan group H. We improve this statement as follows.

**Corollary 5.1.** Any finitely presented group is isomorphic to a finite index normal subgroup of the group Out(H), where H is a Kazhdan group with the property that the group Aut(H) is hyperbolic.

By contrast, any hyperbolic Kazhdan group has finite outer automorphism group [Pau91]. Corollary 5.1 is a particular case of the following corollary, and the remark that  $\operatorname{Aut}(H)$  is hyperbolic because it contains a hyperbolic subgroup G of finite index.

**Corollary 5.2.** If H has property  $F\mathbb{R}$  and is a finitely generated normal subgroup of a non-elementary torsion-free relatively hyperbolic group G, then the canonical homomorphisms  $G/H \cong \text{Inn}(G)/\text{Inn}(H) \to \text{Out}(H)$  and  $G \cong \text{Inn}(G) \to \text{Aut}(H)$  are embeddings onto finite index subgroups.

*Proof.* We think of G as a discrete convergence group. Infinite elementary subgroups have elementary normalizers because the normalizer stabilizes the limit set of the subgroup. Since H is normal and G is non-elementary, we conclude that H is non-elementary.

Now we show that non-elementary subgroups have finite centralizers. Indeed, if the centralizer  $C_G(H)$  of H in G contains an infinite order element c, then H fixes the limit set of the cyclic subgroup generated by c, so H cannot be non-elementary. Thus  $C_G(H)$  must be a torsion group, in particular,  $C_G(H)$ is elementary, else  $C_G(H)$  would have to contain a non-abelian free subgroup. If  $C_G(H)$  is infinite, it has a nonempty limit set that must be fixed by H, contradicting the assumption that H is non-elementary.

Since G is torsion free,  $C_G(H)$  is trivial. Hence the canonical homomorphism  $\operatorname{Inn}(G) \to \operatorname{Aut}(H)$  in injective, and we identify its image with  $\operatorname{Inn}(G)$ . The kernel of the surjection of  $\operatorname{Out}(H) = \operatorname{Aut}(H)/\operatorname{Inn}(H)$  onto  $\operatorname{Aut}(H)/\operatorname{Inn}(G)$  equals to  $\operatorname{Inn}(G)/\operatorname{Inn}(H) \cong G/H$ . By Theorem 1.1 applied to the inclusion  $H \hookrightarrow G$  precomposed with automorphisms of H, the group  $\operatorname{Aut}(H)/\operatorname{Inn}(G)$  is finite.  $\Box$ 

#### 6. Acknowledgements

The first author was partially supported by the NSF grant # DMS-0352576. We are grateful to Oleg V. Belegradek, François Dahmani, Ilya Kapovich, Misha Kapovich, and Denis Osin for helpful communications.

#### References

- [AB] E. Alibegovic and M. Bestvina, *Limit groups are CAT(0)*, arXiv:math.GR/0410198.
- [Bes88] M. Bestvina, Degenerations of the hyperbolic space, Duke Math. J. 56 (1988), no. 1, 143–161.
- [Bes02] \_\_\_\_\_, ℝ-trees in topology, geometry, and group theory, Handbook of geometric topology, North-Holland, Amsterdam, 2002, pp. 55–91.
- [BF95] M. Bestvina and M. Feighn, Stable actions of groups on real trees, Invent. Math. 121 (1995), no. 2, 287–321.
- [Bow] B. H. Bowditch, *Relatively hyperbolic groups*, Southampton preprint, 1999, www.maths.soton.ac.uk/staff/Bowditch/preprints.html.
- [Bow95] B. H. Bowditch, Geometrical finiteness with variable negative curvature, Duke Math. J. 77 (1995), no. 1, 229–274.
- [Bow99] B. H. Bowditch, *Convergence groups and configuration spaces*, Geometric group theory down under (Canberra, 1996), de Gruyter, Berlin, 1999, pp. 23–54.
- [BS94] M. R. Bridson and G. A. Swarup, On Hausdorff-Gromov convergence and a theorem of Paulin, Enseign. Math. (2) 40 (1994), no. 3-4, 267–289.
- [Bum] I. Bumagin, On definitions of relatively hyperbolic groups, arXiv:math.GR/0402072.
- [Bum04] I. Bumagin, Subgroups of torsion-free hyperbolic groups have the Hopf property, Geom. Dedicata 106 (2004), no. 1, 211–230.
- [CV96] M. Culler and K. Vogtmann, A group-theoretic criterion for property FA, Proc. Amer. Math. Soc. 124 (1996), no. 3, 677–683.
- [Dah] F. Dahmani, Accidental parabolics and relatively hyperbolic groups, arXiv:math.GR/0210421.
- [Dah03a] F. Dahmani, Classifying spaces and boundaries for relatively hyperbolic groups, Proc. London Math. Soc. (3) 86 (2003), no. 3, 666–684.

- [Dah03b] \_\_\_\_\_, Combination of convergence groups, Geom. Topol. 7 (2003), 933–963 (electronic).
- [DP03] T. Delzant and L. Potyagailo, Endomorphisms of Kleinian groups, Geom. Funct. Anal. 13 (2003), no. 2, 396–436.
- [DS] C. Drutu and M. Sapir, *Tree-graded spaces and asymptotic cones of groups*, arXiv:math.GT/0405030.
- [DS99] M. J. Dunwoody and M. E. Sageev, JSJ-splittings for finitely presented groups over slender groups, Invent. Math. 135 (1999), no. 1, 25–44.
- [Dun] M. L. Dunwoody, *Groups acting on real trees*, Southampton preprint, 2004, www.maths.soton.ac.uk/staff/Dunwoody/Tess11.dvi.
- [Far98] B. Farb, *Relatively hyperbolic groups*, Geom. Funct. Anal. 8 (1998), no. 5, 810–840.
- [Gro] D. Groves, *Limit groups for relatively hyperbolic groups, I: The basic tools,* arXiv:math.GR/0412492.
- [Gro87] M. Gromov, Hyperbolic groups, Essays in group theory, Math. Sci. Res. Inst. Publ., vol. 8, Springer, 1987, pp. 75–263.
- [Hig61] G. Higman, Subgroups of finitely presented groups, Proc. Roy. Soc. Ser. A 262 (1961), 455–475.
- [HK] G. C. Hruska and B. Kleiner, *Hadamard spaces with isolated flats*, arXiv:math.GR/0411232.
- [KW01] I. Kapovich and D. T. Wise, On the failure of the co-Hopf property for subgroups of word-hyperbolic groups, Israel J. Math. 122 (2001), 125–147.
- [MS88] J. W. Morgan and P. B. Shalen, Degenerations of hyperbolic structures. III. Actions of 3-manifold groups on trees and Thurston's compactness theorem, Ann. of Math. (2) 127 (1988), no. 3, 457–519.
- [Nos93] G. A. Noskov, The T,FA, amenability and continuity properties for groups acting on R-trees, Algebra i Analiz 5 (1993), no. 3, 238–251.
- [OP98] K. Ohshika and L. Potyagailo, Self-embeddings of Kleinian groups, Ann. Sci. École Norm. Sup. (4) 31 (1998), no. 3, 329–343.
- [Osia] D.V. Osin, in preparation.
- [Osib] \_\_\_\_\_, Relative Dehn functions of amalgamated products and HNN-extensions, arXiv:math.GR/0411027.
- [Osic] \_\_\_\_\_, Relatively hyperbolic groups: Intrinsic geometry, algebraic properties, and algorithmic problems, arXiv:math.GR/0404040.
- [Osid] \_\_\_\_\_, Small cancellations over relatively hyperbolic groups and embedding theorems, arXiv:math.GR/0411039.
- [OW] Y. Ollivier and D. T. Wise, Kazhdan groups with infinite outer automorphism group, arXiv:math.GR/0409203.
- [Pau88] F. Paulin, Topologie de Gromov équivariante, structures hyperboliques et arbres réels, Invent. Math. 94 (1988), no. 1, 53–80.
- [Pau91] \_\_\_\_\_, Outer automorphisms of hyperbolic groups and small actions on R-trees, Arboreal group theory (Berkeley, CA, 1988), Math. Sci. Res. Inst. Publ., vol. 19, Springer, 1991, pp. 331–343.
- [RS94] E. Rips and Z. Sela, Structure and rigidity in hyperbolic groups. I, Geom. Funct. Anal. 4 (1994), no. 3, 337–371.
- [Sel95] Z. Sela, The isomorphism problem for hyperbolic groups. I, Ann. of Math. (2) 141 (1995), no. 2, 217–283.
- [Sel97] \_\_\_\_\_, Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups. II, Geom. Funct. Anal. 7 (1997), no. 3, 561–593.

# ENDOMORPHISMS OF RELATIVELY HYPERBOLIC GROUPS

- [Sel99] \_\_\_\_\_, Endomorphisms of hyperbolic groups. I. The Hopf property, Topology 38 (1999), no. 2, 301–321.
- [Sel02] \_\_\_\_\_, Diophantine geometry over groups and the elementary theory of free and hyperbolic groups, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 2002, pp. 87–92.
- [Ser77] J.-P. Serre, Arbres, amalgames, SL<sub>2</sub>, Société Mathématique de France, Paris, 1977, Avec un sommaire anglais, Rédigé avec la collaboration de Hyman Bass, Astérisque, No. 46.
- [Szc98] A. Szczepański, Relatively hyperbolic groups, Michigan Math. J. 45 (1998), no. 3, 611–618.
- [Tuk94] P. Tukia, Convergence groups and Gromov's metric hyperbolic spaces, New Zealand J. Math. 23 (1994), no. 2, 157–187.
- [Wat82] Y. Watatani, Property T of Kazhdan implies property FA of Serre, Math. Japon. 27 (1982), no. 1, 97–103.
- [Yam04] A. Yaman, A topological characterisation of relatively hyperbolic groups, J. Reine Angew. Math. 566 (2004), 41–89.

IGOR BELEGRADEK, SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA30332-0160

*E-mail address*: ib@math.gatech.edu

Andrzej Szczepański, Uniwersytet Gdanski, Instytut Matematyki, Wita Stwosza 57, 80-952 Gdansk, Poland

*E-mail address*: matas@paula.univ.gda.pl