A Sharp Inequality for Conditional Distribution of the First Exit Time of Brownian Motion^{*}

Majid Hosseini Department of Mathematics, State University of New York at New Paltz, 75 S. Manheim Blvd. Suite 9, New Paltz, NY 12561 hosseinm@newpaltz.edu

September 7, 2004

*AMS Subject Classification (2000): 60J65, 60K99

Abstract

Let U be a domain, convex in x and symmetric about the y-axis, which is contained in a centered and oriented rectangle R. If τ_A is the first exit time of Brownian motion from A and $A^+ = A \cap \{(x,y) : x > 0\}$, it is proved that $P^z(\tau_{U^+} > s \mid \tau_{R^+} > t) \leq P^z(\tau_U > s \mid \tau_R > t)$ for every s, t > 0 and every $z \in U^+$.

Keywords: Brownian Motion, First Exit Time, Conditional Distribution

1 Introduction

Let A be a subset of \mathbb{R}^2 . The set A is convex in x if its intersection with every line parallel to the x-axis is a single interval or empty. We put $A^+ = A \cap \{(x, y) \mid x > 0\}$. Also, let $B_t = (B_{1,t}, B_{2,t}), t \ge 0$ be standard two dimensional Brownian motion and $\tau_A = \inf\{t > 0 : B_t \notin A\}$. We will prove the following.

Theorem 1 Let U be an open, bounded, and connected set in \mathbb{R}^2 which is symmetric about the y-axis and convex in x. Also, let R be an open rectangle containing U, that is symmetric with respect to the y-axis, and has sides parallel to the axes. If z is a point in U^+ , then for every s, t > 0,

$$P^{z}(\tau_{U^{+}} > s \mid \tau_{R^{+}} > t) \le P^{z}(\tau_{U} > s \mid \tau_{R} > t).$$
(1)

Recently, inequalities of this type, wherein the values of various quantities related to U, U^+ , R, and R^+ are compared, have been studied extensively. Davis [2] proved the first inequality of this kind for the heat kernel of Laplacian. Bañuelos and Méndez-Hernández [1] extended Davis's result to the heat kernel of Schrödinger operators and integrals of these kernels. You [4] proved an inequality of this type for the trace of Schrödinger operators. Davis and Hosseini [3] proved the extension to the heat content. The inequalities studied in [1, 2, 3, 4] are "ratio inequalities", in the sense that the left side of the inequality is the ratio of some functional of U^+ and the same functional for R^+ , and the right side is the the corresponding ratio for U and R. Inequality (1) is not strictly a ratio inequality, but rather a "ratio-like" inequality.

The proof of Theorem 1 is based on the idea of conditioning on zeros. Since the zeros of Brownian motion are uncountable, it is not possible to use this approach for Brownian motion directly. Therefore, we first prove a discrete analog of (1), and then use scaling. The discrete analog is stated and proved in the following section. Its proof is an application and modification of the techniques introduced in [3]. We will repeat some of the material in [3] so that we can refer to them and modify them for our purpose. We will point them out as we go through the proof. The method of deriving (1) from its discrete counterpart is a standard application of the invariance principle. We will omit this derivation for the sake of brevity. See [3] for a detailed description of an almost identical derivation.

The following example shows that if the convexity condition in Theorem 1 is removed, we can find a domain U for which Theorem 1 fails. Let 0 < d < 1/2 and take $U = (-1,1) \times (-1,1) \setminus \{(0,y) : |y| \ge d/2\}$. Let $R = (-1,1) \times (-1,1)$ and z = (d,1/2). Also, put s = t = 1. We will show at the end of Section 2 that, for this example, the right side of (1) converges to zero as $d \to 0$ while the left side is equal to 1 for all values of d.

Note that since

$$E^{z}\left(\tau_{U} \mid \tau_{R} > t\right) = \int_{0}^{\infty} P^{z}\left(\tau_{U} > s \mid \tau_{R} > t\right) \, ds,$$

and a similar equation holds for $E^{z}(\tau_{U^{+}} | \tau_{R^{+}} > t)$, the following is an immediate consequence of Theorem 1.

Corollary 2 Let U, R, and z be as in Theorem 1. Then for every t > 0,

$$E^{z}(\tau_{U^{+}} \mid \tau_{R^{+}} > t) \le E^{z}(\tau_{U} \mid \tau_{R} > t).$$
 (2)

Corollary 2 and the ratio inequalities proved in [1, 2, 3, 4] lead to the following conjecture.

Conjecture 3 Let U, R, and z be as in Theorem 1. Then

$$\frac{E^{z}\left(\tau_{U^{+}}\right)}{E^{z}\left(\tau_{R^{+}}\right)} \leq \frac{E^{z}\left(\tau_{U}\right)}{E^{z}\left(\tau_{R}\right)}.$$
(3)

2 Discrete-Time Inequalities

Let $\{X_i\}_{i\geq 0}$ and $\{Y_i\}_{i\geq 0}$ be independent sequences of random variables such that both sequences $\{X_{i+1} - X_i\}$ and $\{Y_{i+1} - Y_i\}$ are i.i.d. sequences of random variables, each taking values 0, 1, and -1 with probability 1/3. Let

 $Z_i = (X_i, Y_i)$. Thus Z_i is a random walk on \mathbb{Z}^2 started at Z_0 . Consider $\Lambda \subset \mathbb{Z}^2$ and let z = (x, y) be a point in Λ . For any set A let τ_A be the first exit time of $Z_i, i \geq 0$ from A.

Proposition 4 Let Λ be a bounded and connected subset of \mathbb{Z}^2 which is symmetric about the y-axis and convex in x. Let T be a rectangle containing Λ with sides parallel to the axes. Then for all nonnegative integers m and n,

$$P^{z}\left(\tau_{\Lambda^{+}} > m \mid \tau_{T^{+}} > n\right) \leq P^{z}\left(\tau_{\Lambda} > m \mid \tau_{T} > n\right).$$

$$\tag{4}$$

We will prove the equivalent statement, that for all nonnegative integers m and n,

$$\frac{P^{z}\left(\tau_{\Lambda^{+}}>m,\tau_{T^{+}}>n\right)}{P^{z}\left(\tau_{\Lambda}>m,\tau_{T}>n\right)} \leq \frac{P^{z}\left(\tau_{T^{+}}>n\right)}{P^{z}\left(\tau_{T}>n\right)}.$$
(5)

Let $l = \max(m, n)$ and let $\boldsymbol{y} = (y_0, \dots, y_l)$ be a sequence such that $y_0 = y$, and $|y_i - y_{i-1}| \leq 1$ for $1 \leq i \leq l$. In addition, assume that

$$P^{z}\left(Y_{i}=y_{i}, Z_{i}\in\Lambda, 0\leq i\leq l\right)>0.$$

We will call such \boldsymbol{y} admissible. Let \boldsymbol{y} be an admissible sequence. For an event A, define

$$P^{y,z}(A) = P^{z}(A \mid Y_{i} = y_{i}, 0 \le i \le l).$$

We will prove the following.

Lemma 5 Let Λ , T, m, n and z be as in Proposition 4. Let \boldsymbol{y} be an admissible sequence. Then

$$\frac{P^{\boldsymbol{y},z}\left(\tau_{\Lambda^{+}} > m, \tau_{T^{+}} > n\right)}{P^{\boldsymbol{y},z}\left(\tau_{\Lambda} > m, \tau_{T} > n\right)} \le \frac{P^{\boldsymbol{y},z}\left(\tau_{T^{+}} > n\right)}{P^{\boldsymbol{y},z}\left(\tau_{T} > n\right)}.$$
(6)

Note that the right side of (6) is independent of \boldsymbol{y} and equals the right side of (5). Therefore Lemma 5 implies Proposition 4.

For an admissible \boldsymbol{y} , let $\gamma_{\boldsymbol{y}}^{\Lambda}(k) = \max\{i : (i, y_k) \in \Lambda\}$. Since $\{X_i\}_{i \geq 0}$ and $\{Y_i\}_{i \geq 0}$ are independent, if z = (x, y), then

$$P^{\boldsymbol{y},z}\left(\tau_{T^+} > n\right) = P^x\left(0 < X_i \le \gamma_{\boldsymbol{y}}^T(i), 0 \le i \le n\right),$$

and

$$P^{\boldsymbol{y},z}\left(\tau_T > n\right) = P^x\left(|X_i| \le \gamma_{\boldsymbol{y}}^T(i), 0 \le i \le n\right),$$

and similar equalities hold for the remaining quantities in (6).

Let $f: \{0, \ldots, n\} \to \mathbb{Z}^+$ and $h: \{0, \ldots, m\} \to \mathbb{Z}^+$ be such that $h(i) \leq f(i)$ for all $i \in \{0, \ldots, \min(m, n)\}$. Furthermore assume x is an integer such that $x \leq h(0)$. Also, put $F^+ = \{0 < X_i \leq f(i), 0 \leq i \leq n\}$ and $F = \{|X_i| \leq f(i), 0 \leq i \leq n\}$. Define H^+ and H similarly, by replacing f with h and nwith m. We will show that

$$P^{x}\left(H^{+} \mid F^{+}\right) \leq P^{x}\left(H \mid F\right),\tag{7}$$

which, by the above discussion, implies (6).

In order to prove (7), we need to investigate the properties of the joint distribution of $\{X_i\}_{i=0}^m$ and $\{|X_i|\}_{i=0}^m$, given F^+ and F respectively. For the case $m \leq n$, this was done in [3]. In particular, Lemmas 7 and 8 in [3] are the main tools for this task. These two Lemmas are restated as Lemmas 6 and 7 in this note. Their proofs and the discussion immediately preceding them are repeated from [3], both for completeness and for the modifications we will make in these proofs to prove the case m > n.

Let a, b, α , and β be integers such that $0 \le a < b \le n$ and that $0 < \alpha \le f(a)$, and $0 < \beta \le f(b)$. Define the probability measures $P_{a,b}^{\alpha}$ and $R_{a,b}^{\alpha,\beta}$ on the set of all finite sequences $(x_a, x_{a+1}, \ldots, x_b)$ of integers by

$$P_{a,b}^{\alpha}(x_a, x_{a+1}, \dots, x_b)$$

$$\stackrel{\text{def}}{=} P(X_k = x_k, a \le k \le b | X_a = \alpha, 0 < X_i \le f(i), a \le i \le b),$$

and

$$R_{a,b}^{\alpha,\beta}(x_a, x_{a+1}, \dots, x_b) \overset{\text{def}}{=} P(X_k = x_k, a \le k \le b \mid X_a = \alpha, X_b = \beta, 0 < X_i \le f(i), a \le i \le b).$$

Let π_j be the coordinate maps: $\pi_j(x_a, x_{a+1}, \ldots, x_b) = x_j$. Under $P_{a,b}^{\alpha}$ the finite sequence of random variables $\pi_a, \pi_{a+1}, \ldots, \pi_b$ is a Markov chain started at α with (non-stationary) transition probabilities, which do not depend on α , given by

$$P_{a,b}^{\alpha}(\pi_{k+1} = v | \pi_k = u) = \frac{p_v^{k+1}}{\sum_{j=u-1}^{u+1} p_j^{k+1}} \quad \text{for } v = u - 1, u, u + 1, \quad (8)$$

where

$$p_j^{k+1} = P(0 < X_i \le f(i), k+1 \le i \le b | X_{k+1} = j).$$

Also, under $R_{a,b}^{\alpha,\beta}$ the sequence $\pi_a, \pi_{a+1}, \ldots, \pi_b$ is a Markov chain started at α with transition probabilities given by

$$R_{a,b}^{\alpha,\beta}(\pi_{k+1} = v | \pi_k = u) = \frac{r_v^{k+1}}{\sum_{j=u-1}^{u+1} r_j^{k+1}} \quad \text{for } v = u - 1, u, u + 1, \quad (9)$$

where

$$r_j^{k+1} = P(X_b = \beta, 0 < X_i \le f(i), k+1 \le i \le b | X_{k+1} = j).$$

Lemma 6 Let $0 < \alpha_0 \leq \alpha_1$. Consider Markov chains $\psi_a(=\alpha_0), \psi_{a+1}, \ldots, \psi_b$ and $\zeta_a(=\alpha_1), \zeta_{a+1}, \ldots, \zeta_b$ that have transition probabilities given by (8) with α replaced by α_0 and α_1 respectively. Then there are Markov chains $\tilde{\psi}_a, \ldots, \tilde{\psi}_b$ and $\tilde{\zeta}_a, \ldots, \tilde{\zeta}_b$, defined on a common probability space Ω , and having the same transition probabilities and initial distributions as ψ_a, \ldots, ψ_b and ζ_a, \ldots, ζ_b respectively, such that

$$\tilde{\psi}_i(\omega) \leq \tilde{\zeta}_i(\omega) \quad \text{for every } \omega \in \Omega \text{ and } a \leq i \leq b.$$
(10)

Proof We will use induction. For each k satisfying $a \leq k < b$, all $r \in \mathbb{R}$, and each u such that $1 \leq u \leq f(k)$, let $F_{k+1}(r, u) = P_{a,b}^{\alpha_0}(\pi_{k+1} \leq r | \pi_k = u)$ and $G_{k+1}(r, u) = P_{a,b}^{\alpha_1}(\pi_{k+1} \leq r | \pi_k = u)$. Let $\tilde{\psi}_a = \alpha_0$ and $\tilde{\zeta}_a = \alpha_1$ on Ω . Assume that $\tilde{\psi}_a, \ldots, \tilde{\psi}_k$ and $\tilde{\zeta}_a, \ldots, \tilde{\zeta}_k$ have been defined on Ω . Let T_{k+1} be a random variable uniformly distributed over [0, 1], defined on Ω , and independent of $\tilde{\psi}_a, \ldots, \tilde{\psi}_k$ and $\tilde{\zeta}_a, \ldots, \tilde{\zeta}_k$. For each u such that $1 \leq u \leq f(k)$ and all $t \in [0, 1]$ define $F_{k+1}^{-1}(t, u) = \inf\{r : F_{k+1}(r, u) \geq t\}$ and define $G_{k+1}^{-1}(t, u)$ similarly. On the event $\{\tilde{\psi}_k = u\}$ let $\tilde{\psi}_{k+1} = F_{k+1}^{-1}(T_{k+1}, u)$. Define $\tilde{\zeta}_{k+1}$ in a similar manner. It is routine to check that these are indeed Markov chains with the desired transition probabilities and initial distributions. We use induction to prove $\tilde{\psi}_k(\omega) \leq \tilde{\zeta}_k(\omega)$ for all $\omega \in \Omega$. We have $\tilde{\psi}_a = \alpha_0 \leq \alpha_1 = \tilde{\zeta}_a$. Now assume that $\tilde{\psi}_k(\omega) \leq \tilde{\zeta}_k(\omega)$ for all $\omega \in \Omega$. Consider $1 \leq u \leq u'$ and $\omega \in \{\tilde{\psi}_k = u, \tilde{\zeta}_k = u'\}$. For all $r \in \mathbb{R}$ and each $1 \leq u \leq u'$ we have

$$F_{k+1}(r,u) \ge G_{k+1}(r,u'). \tag{11}$$

For if $u' \geq u + 1$, (11) follows directly from the form of the transition probabilities for ζ and ψ . When u = u' we have $F_{k+1}(r, u) = G_{k+1}(r, u')$. Hence by (11), we have $F_{k+1}^{-1}(t, u) \leq G_{k+1}^{-1}(t, u')$ for each $t \in [0, 1]$, and so $\tilde{\psi}_{k+1}(\omega) \leq \tilde{\zeta}_{k+1}(\omega)$. This completes the proof of Lemma 6. **Lemma 7** Let $0 < \alpha_0 \leq \alpha_1$ and $0 < \beta_0 \leq \beta_1$. Consider Markov chains $\psi_a(=\alpha_0), \ldots, \psi_b$ and $\zeta_a(=\alpha_1), \ldots, \zeta_b$ that have transition probabilities given by (9) with (α, β) replaced by (α_0, β_0) and (α_1, β_1) respectively. Then there are Markov chains $\tilde{\psi}_a, \ldots, \tilde{\psi}_b$ and $\tilde{\zeta}_a, \ldots, \tilde{\zeta}_b$, defined on a common probability space Ω , and having the same transition probabilities and initial distributions as ψ_a, \ldots, ψ_b and ζ_a, \ldots, ζ_b respectively, such that

$$\psi_i(\omega) \le \zeta_i(\omega)$$
 for every $\omega \in \Omega$ and $a \le i \le b$. (12)

Proof For each k satisfying $a \leq k < b$, all $r \in \mathbb{R}$, and each u such that $1 \leq u \leq f(k)$, let $F_{k+1}(r, u) = R_{a,b}^{\alpha_0,\beta_0}(\pi_{k+1} \leq r | \pi_k = u)$ and $G_{k+1}(r, u) = R_{a,b}^{\alpha_1,\beta_1}(\pi_{k+1} \leq r | \pi_k = u)$. The rest of the proof follows the proof of Lemma 6 closely. The only difference is in the proof of the statement that for all $r \in \mathbb{R}$ and each $1 \leq u \leq u'$ we have

$$F_{k+1}(r,u) \ge G_{k+1}(r,u'). \tag{13}$$

To show this first assume that $\beta_0 = \beta_1$ and u' = u + 1. Then (13) follows directly from the form of the transition probabilities for ζ and ψ . When u = u' and $\beta_0 = \beta_1$, we have $F_{k+1}(r, u) = G_{k+1}(r, u')$. These facts also imply that for fixed r, the function $G_{k+1}(r, u)$ is decreasing in u. Put together, the special case $\beta_0 = \beta_1$ follows. Therefore, Lemma 7 holds in this case. The special case $\alpha_0 = \alpha_1$ and u' = u follows from the fact that running our conditioned walks backwards in time still gives a conditioned walk (we are just counting paths) and the special case of Lemma 7 for $\beta_0 = \beta_1$. The general case follows from these two special cases by first considering the pairs (α_0, β_0) and (α_1, β_0) followed by the pairs (α_1, β_0) and (α_1, β_1). This completes the proof of (13), and therefore, of Lemma 7.

We can now prove (7). We consider two cases: (1) $m \leq n$ and (2) m > n. Let e_0 be the left side of (7). In both cases, we will find a partition Ψ of F such that

$$e_0 \le P^x \left(H \mid A \right), \quad A \in \Psi. \tag{14}$$

For the case $m \leq n$ the proof is essentially the same as the proof of the case m = n which was done in [3]. Since we will use the ideas in this proof for proving the case m > n, we bring this proof here with the minor adjustments that are necessary.

We start with the case $m \leq n$. Let $N = \operatorname{card}\{i : 0 < i \leq n, X_i = 0\}$ and and let M_1, M_2, \ldots, M_N be the indices $i \leq n$ such that $X_i = 0$. Define

$$Q_{2,c,d} = \{ N = 2, M_1 = c, M_2 = d \} \cap F.$$

Let k be a positive integer. For a sequence i_1, \ldots, i_k of positive integers, define Q_{k,i_1,\ldots,i_k} in the same manner as $Q_{2,c,d}$. Also, put $Q = \{N = 0\} \cap F$.

Clearly $e_0 = P^x(H|Q)$. To make the argument easier to follow, assume that k = 2, $M_1 = c$, and $M_2 = d$ and consider $Q_{2,c,d}$. Given $Q_{2,c,d}$, the random variables $|X_0|, \ldots, |X_n|$ form a nonhomogeneous Markov chain ψ_0, \ldots, ψ_n . Note that the Markov chain $\psi_0, \ldots, \psi_{c-1}$ has the same transition probabilities as the sequence π_0, \ldots, π_{c-1} under $R_{0,c-1}^{x,1}$. Also the Markov chain $\psi_{c+1}, \ldots, \psi_{d-1}$ has the same transition probabilities as the sequence $\pi_{c+1}, \ldots, \pi_{d-1}$ under $R_{c+1,d-1}^{1,1}$ and the Markov chain $\psi_{d+1}, \ldots, \psi_n$ has the same transition probabilities as the sequence π_{d+1}, \ldots, π_n under $P_{d+1,n}^1$.

Now consider $s_1, s_2, s_3 \geq 1$ and $t_1, t_2, t_3 \geq 1$ for which it is possible to condition on F^+ , $X_{c-1} = s_1$, $X_c = s_2$, $X_{c+1} = s_3$, $X_{d-1} = t_1$, $X_d = t_2$, and $X_{d+1} = t_3$. Fix the s_i and the t_i $(1 \leq i \leq 3$.) Then given the above condition the sequence X_0, \ldots, X_n is a non-homogeneous Markov chain ζ_0, \ldots, ζ_n such that $\zeta_0, \ldots, \zeta_{c-1}$ will have the same transition probabilities as π_0, \ldots, π_{c-1} under $R_{0,c-1}^{s,s_1}$. Also $\zeta_{c+1}, \ldots, \zeta_{d-1}$ has the same transition probabilities as $\pi_{c+1}, \ldots, \pi_{d-1}$ under $R_{c+1,d-1}^{s,s_1}$ and $\zeta_{d+1}, \ldots, \zeta_n$ has the same transition probabilities as π_{d+1}, \ldots, π_n under $P_{d+1,n}^{t_3}$. For the Markov chains $\{\psi_i\}_{i=0}^{c-1}$ and $\{\zeta_i\}_{i=0}^{c-1}$, consider $\tilde{\psi}_0, \ldots, \tilde{\psi}_{c-1}$ and $\tilde{\zeta}_0, \ldots, \tilde{\zeta}_{c-1}$ as constructed in Lemma 7. Do the same for the time frame [c + 1, d - 1], making sure that $\{T_i\}_{i=1}^{d-1}$ considered in Lemma 7 are independent of $\{T_i\}_{i=d+1}^c, \{\tilde{\psi}_i\}_{i=1}^c, \ldots, \tilde{\psi}_n$ and $\tilde{\zeta}_{d+1}, \ldots, \tilde{\zeta}_n$ as constructed in Lemma 6, again, making sure that T_{d+1}, \ldots, π_n are independent of all previous $T_i, \tilde{\psi}_i$, and $\tilde{\zeta}_i$. We also have $s_2 = \zeta_c > \psi_c = 0$ and $t_2 = \zeta_d > \psi_d = 0$. Define $\tilde{\zeta}_c = s_2, \tilde{\psi}_c = 0, \tilde{\zeta}_d = t_2$, and $\tilde{\psi}_d = 0$ on Ω .

Put $\hat{H}^+ = \{\zeta_i \leq h(i), 0 \leq i \leq m\}$ and $\hat{H} = \{\psi_i \leq h(i), 0 \leq i \leq m\}$. Hence for every $\omega \in \Omega$ we have $I_{\tilde{H}^+}(\omega) \leq I_{\tilde{H}}(\omega)$.

The Markov chain ψ has the same distribution as the Markov chain $|X_0|, \ldots, |X_n|$ given $Q_{2,c,d}$. The Markov chain $\tilde{\zeta}$ has the same distribution as X_0, \ldots, X_n given F^+ and $X_{c-1} = s_1, X_c = s_2, X_{c+1} = s_3, X_{d-1} = t_1, X_d = t_2$, and $X_{d+1} = t_3$.

Therefore for all possible values of s_1 , s_2 , s_3 , t_1 , t_2 , and $t_3 \ge 1$,

$$P^{x} \Big(H^{+} \mid F^{+} \text{ and } X_{c-1} = s_{1}, X_{c} = s_{2}, X_{c+1} = s_{3}, X_{d-1} = t_{1}, X_{d} = t_{2}, X_{d+1} = t_{3} \Big)$$
$$\leq P^{x} \Big(H \mid Q_{2,c,d} \Big).$$

Since this is true for all possible values of $s_1, s_2, s_3, t_1, t_2, t_3 \ge 1$, we have that

$$P^{x}\left(H^{+} \mid F^{+}\right) \leq P^{x}\left(H \mid Q_{2,c,d}\right).$$

This proves (14) for $Q_{2,c,d}$. The argument for all other Q_{k,i_1,\ldots,i_k} is the same as above. Therefore the proof of (14) for the case $m \leq n$ is complete.

To prove (14) for the case m > n, let N, M_1, \ldots, M_N, Q and Q_{k,i_1,\ldots,i_k} be as in the proof of the case $m \le n$. Again, to make the proof easier to follow, we focus on $Q_{2,c,d}$. Given $Q_{2,c,d}$, the random variables $|X_0|, \ldots, |X_m|$ form a nonhomogeneous Markov chain ψ_0, \ldots, ψ_m . The part ψ_0, \ldots, ψ_n has the same transition probabilities as discussed in the proof of the case $m \le n$. The part ψ_n, \ldots, ψ_m has transition probabilities

$$P(\psi_{k+1} = v | \psi_k = u) = \frac{1}{3} \quad \text{for } u > 0 \text{ and } v = u - 1, u, u + 1, \quad (15)$$

and

$$P(\psi_{k+1} = 1 | \psi_k = 0) = \frac{2}{3} = 1 - P(\psi_{k+1} = 0 | \psi_k = 0).$$
(16)

Now consider $s_1, s_2, s_3 \ge 1$ and $t_1, t_2, t_3 \ge 1$ for which it is possible to condition on F^+ , $X_{c-1} = s_1$, $X_c = s_2$, $X_{c+1} = s_3$, $X_{d-1} = t_1$, $X_d = t_2$, and $X_{d+1} = t_3$. Fix the s_i and the t_i . Then given the above condition the sequence X_0, \ldots, X_m is a non-homogeneous Markov chain ζ_0, \ldots, ζ_m . The part ζ_0, \ldots, ζ_n has the same transition probabilities as discussed in the proof of the case $m \le n$. The part ζ_n, \ldots, ζ_m has transition probabilities

$$P(\zeta_{k+1} = v \mid \zeta_k = u) = \frac{1}{3} \quad \text{for } v = u - 1, u, u + 1.$$
 (17)

Construct $\tilde{\psi}_0, \ldots, \tilde{\psi}_n$ and $\tilde{\zeta}_0, \ldots, \tilde{\zeta}_n$ as in the proof of the case $m \leq n$. For $n \leq k \leq m-1$, put $F_{k+1}(r, u) = P(\psi_{k+1} \leq r \mid \psi_k = u)$ and $G_{k+1}(r, u) = P(\zeta_{k+1} \leq r \mid \zeta_k = u)$. It is easy to see that for $0 < u \leq v$ with v > 0,

$$F_{k+1}(r,u) \ge G_{k+1}(r,v)$$
 for $n \le k \le m-1$. (18)

Next, we will construct $\tilde{\psi}_{n+1}, \ldots, \tilde{\psi}_m$ and $\tilde{\zeta}_{n+1}, \ldots, \tilde{\zeta}_m$ on Ω by the method used in Lemma 6. Assume that $\tilde{\psi}_n, \ldots, \tilde{\psi}_k$ and $\tilde{\zeta}_n, \ldots, \tilde{\zeta}_k$ have been defined on Ω . Let T_{k+1} be a random variable uniformly distributed over [0, 1], defined on Ω , independent of T_0, \ldots, T_k , as well as independent of $\tilde{\psi}_0, \ldots, \tilde{\psi}_k$ and $\tilde{\zeta}_0, \ldots, \tilde{\zeta}_k$. For each u such that $0 \leq u$ and all $t \in [0, 1]$ define $F_{k+1}^{-1}(t, u) = \inf\{r : F_{k+1}(r, u) \geq t\}$ and for $u \in \mathbb{Z}$ define $G_{k+1}^{-1}(t, u)$ similarly. On the event $\{\tilde{\psi}_k = u\}$ let $\tilde{\psi}_{k+1} = F_{k+1}^{-1}(T_{k+1}, u)$. Define $\tilde{\zeta}_{k+1}$ in a similar manner. Again, it is easy to check that $\tilde{\psi}_n, \ldots, \tilde{\psi}_m$ and $\tilde{\zeta}_n, \ldots, \tilde{\zeta}_m$ are Markov chains with the same transition probabilities as ψ_n, \ldots, ψ_m and ζ_n, \ldots, ζ_m respectively.

We will show that

$$\{0 < \tilde{\zeta}_k \le h(k), 0 \le k \le m\} \subseteq \{|\tilde{\psi}_k| \le h(k), 0 \le k \le m\}$$

$$= \{\tilde{\psi}_k \le h(k), 0 \le k \le m\}.$$
(19)

Note that, by construction, for $0 \leq k \leq n$ and for all $\omega \in \Omega$, we have $\tilde{\psi}_k(\omega) \leq \tilde{\zeta}_k(\omega)$ and $0 < \tilde{\zeta}_k(\omega)$. We will show, by induction, that for $n+1 \leq k \leq m$, if $0 < \tilde{\zeta}_k(\omega)$, then $\tilde{\psi}_k(\omega) \leq \tilde{\zeta}_k(\omega)$. First note that $\tilde{\psi}_n(\omega) \leq \tilde{\zeta}_n(\omega)$ and $0 < \tilde{\zeta}_n(\omega)$ for all $\omega \in \Omega$.

Now assume that $\omega \in \{\tilde{\psi}_k \leq \tilde{\zeta}_k, \tilde{\zeta}_k > 0\}$. We will show that $\tilde{\psi}_{k+1}(\omega) \leq \tilde{\zeta}_{k+1}(\omega)$. Consider integers u and v with $0 \leq u \leq v$ and v > 0 and assume that $\omega \in \{\tilde{\psi}_k = u, \tilde{\zeta}_k = v\}$. By (18), we have $F_{k+1}^{-1}(t, u) \leq G_{k+1}^{-1}(t, v)$ for each $t \in [0, 1]$. Therefore, if $\omega \in \{0 < \tilde{\zeta}_k \leq h(k), 0 \leq k \leq m\}$, then $\omega \in \{\tilde{\psi}_k \leq h(k), 0 \leq k \leq m\}$. Hence for all $\omega \in \Omega$,

$$I_{\{0<\tilde{\zeta}_k\leq h(k),0\leq k\leq m\}}(\omega)\leq I_{\{\tilde{\psi}_k\leq h(k),0\leq k\leq m\}}(\omega).$$

The Markov chain $\tilde{\psi}_0, \ldots, \tilde{\psi}_m$ has the same distribution as the Markov chain $|X_0|, \ldots, |X_m|$ given $Q_{2,c,d}$. The Markov chain $\tilde{\zeta}_0, \ldots, \tilde{\zeta}_m$ has the same distribution as X_0, \ldots, X_m given F^+ and $X_{c-1} = s_1, X_c = s_2, X_{c+1} = s_3, X_{d-1} = t_1, X_d = t_2$, and $X_{d+1} = t_3$.

Therefore for all possible values of $s_1, s_2, s_3, t_1, t_2, t_3 \ge 1$,

$$P^{x}\Big(H^{+} \mid F^{+} \text{ and } X_{c-1} = s_{1}, X_{c} = s_{2}, \quad X_{c+1} = s_{3}, X_{d-1} = t_{1}, X_{d} = t_{2}, X_{d+1} = t_{3}\Big)$$
$$\leq P^{x}\Big(H \mid Q_{2,c,d}\Big).$$

Since this is true for all possible values of $s_1, s_2, s_3, t_1, t_2, t_3 \ge 1$, we have that

$$P^{x}(H^{+} | F^{+}) \leq P^{x}(H | Q_{2,c,d}).$$

This proves for $Q_{2,c,d}$. The argument for Q and all other Q_{k,i_1,\ldots,i_k} is the same as above. Therefore the proof of (14) and of Lemma 5 is complete. As we

pointed out earlier, Lemma 5 implies Proposition 4. Finally, the functional central limit theorem implies that Theorem 1 follows from Proposition 4. This completes the proof of Theorem 1.

Now we will show that for the example in the introduction inequality (1) fails. Recall that $B_t = (B_{1,t}, B_{2,t}), t \ge 0$, is the standard two dimensional Brownian motion. Note that $R^+ = U^+$ and therefore the left side of (1) equals 1. On the other hand for any $\theta < 1$ we have,

$$P^{z}(\tau_{U} > 1, \tau_{R} > 1) \leq P^{z}(\tau_{U} > 1) \leq P^{z}(B_{1,t} > 0 \text{ for all } t \in (0,\theta))$$
(20)
+ $P^{z}(B_{2,t} < d/2 \text{ for some } t \in (0,\theta)).$

Given $\epsilon > 0$, by the continuity of Brownian motion paths, there exists $\theta_0 \in (0, 1)$ such that

$$P^{z} (B_{2,t} < d/2 \text{ for some } t \in (0, \theta_{0})) \leq P^{z} (B_{2,t} < 1/4 \text{ for some } t \in (0, \theta_{0})) < \frac{\epsilon}{2}.$$
(21)

Also, by the Law of iterated logarithm, for d small enough, we have

$$P^{z}\left(B_{1,t} > 0 \text{ for all } t \in (0,\theta_{0})\right) < \frac{\epsilon}{2}.$$
(22)

Put together, (20), (21), and (22) imply that

$$\lim_{d \to 0} P^{z}(\tau_{U} > 1, \tau_{R} > 1) = 0.$$
(23)

On the other hand,

$$\lim_{d \to 0} P^{z}(\tau_{R} > 1) = P^{(0,1/2)}(\tau_{R} > 1) > 0.$$
(24)

It follows immediately from (23) and (24) that

$$\lim_{d \to 0} P^{z}(\tau_{U} > 1 \mid \tau_{R} > 1) = 0.$$

3 Higher Dimensions

The analog of Theorem 1 holds for an arbitrary dimension k. We state it as Theorem 8 and we will show how the proof of Theorem 1 can be modified to prove it.

Represent a point in \mathbb{R}^k by $z = (z_1, \dots, z_k)$. Also for $A \subseteq \mathbb{R}^k$ put $A^+ = A \cap \{z \in \mathbb{R}^k \mid z_1 > 0\}$. Call A convex in z_1 if the intersection of A with every line parallel to z_1 -axis is a connected interval or empty.

Theorem 8 Let k be a positive integer and let U be a bounded, connected, and open subset of \mathbb{R}^k which is symmetric about $\{z_1 = 0\}$ and convex in z_1 . Also, let $R = (-L_1, L_1) \times \ldots \times (-L_k, L_k)$ be a k-dimensional rectangle, containing U. Then for every $z \in U^+$ and every s, t > 0,

$$P^{z}(\tau_{U^{+}} > s \mid \tau_{R^{+}} > t) \le P^{z}(\tau_{U} > s \mid \tau_{R} > t).$$
(25)

The proof of Theorem 8 is very similar to the proof of Theorem 1. Let $\{X_i^1\}_{i\geq 0}, \ldots, \{X_i^k\}_{i\geq 0}$ be k independent one-dimensional random walks, each constructed as $\{X_i\}_{i\geq 0}$ in Section 2. Then the analog of Proposition 4 holds by conditioning on $\{(X_i^2, \cdots, X_i^k)\}_{i=0}^l$.

Acknowledgement I am most grateful to Professor Burgess Davis for his valuable comments on the earlier drafts of this paper, and in particular for pointing out the example in the introduction.

References

- Bañuelos, R. and Méndez-Hernández, P.J.: 'Sharp inequalities for heat kernels of Schrödinger operators and applications to spectral gaps', J. Funct. Anal. 176 (2000) 368–399.
- [2] Davis, B.: 'On the spectral gap for fixed membranes', Ark. Mat.
 (1) 39 (2001) 65–74.
- [3] Davis, B. and Hosseini, M.: 'On Ratio Inequalities for Heat Content', J. London Math. Soc. (2) 69 (2004) 97–106.
- [4] You, D.: 'Sharp inequalities for ratios of partition functions of Schrödinger operators', *Potential Anal.* 18 (2003) 219–250.