

q -ANALOGUES OF THE SUMS OF POWERS OF CONSECUTIVE INTEGERS

TAEKYUN KIM

Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim@kongju.ac.kr (or tkim64@hanmail.net)

ABSTRACT. Let n, k be the positive integers ($k > 1$), and let $S_{n,q}(k)$ be the sums of the n -th powers of positive q -integers up to $k - 1$: $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l l^n$. Following an idea due to J. Bernoulli, we explore a formula for $S_{n,q}(k)$.

1. INTRODUCTION

In the early of 17th century Faulhaber [3, 6, 11] computed the sums of powers $1^m + 2^m + \cdots + n^m$ up to $m = 17$ and realized that for odd m , it is not just a polynomial in n but a polynomial in the triangular number $N = n(n+1)/2$. A good account of Faulhaber's work was given by Knuth [3, 6, 11]. In 1713, J. Bernoulli first discovered the method which one can produce those formulae for the sum $\sum_{l=1}^n l^k$ for any natural numbers k , cf. [1, 2, 4, 5, 13]. Let q be an indeterminate which can be considered in complex number field, and for any integer k define the q -integer as $[k]_q = \frac{q^k - 1}{q - 1}$, cf. [2, 7, 8, 9, 10]. Note that $\lim_{q \rightarrow 1} [k]_q = k$. Recently, many authors are studying the problems of q -analogues of sums of powers of consecutive integers [3, 4, 5, 6, 7, 8, 11, 12, 13, 14]. In this paper we consider the q -analogues of the sums of powers of consecutive integers due to J. Bernoulli. For any positive integers $n, k (> 1)$,

2000 Mathematics Subject Classification 11S80, 11B68

Key words and phrases: Sums of powers, Bernoulli number, Bernoulli polynomials

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

let $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l l^n$. Following an idea due to J. Bernoulli, we explore a formula for $S_{n,q}(k)$ as follows:

$$q^{-k} S_{l-1,q}(k) + (q^{-k} \log q) S_{l,q}(k) = \frac{1}{l} \sum_{i=0}^{l-1} \binom{l}{i} B_{i,q} k^{l-i} + \frac{(1 - q^{-k}) B_{l,q}(0)}{l},$$

where $B_{i,q}$ are the q -analogue of Bernoulli numbers.

2. q -ANALOGUE OF THE SUMS OF THE n -TH POWERS OF POSITIVE INTEGERS UP TO $k - 1$

Let j be the positive integers. Then we easily see that

$$(1) \quad q^{j+1}(j+1) - q^j j = (q-1)q^j j + q^{j+1}.$$

From Eq.(1), we note that

$$(2) \quad (q-1) \sum_{j=0}^{k-1} q^j j + q[k]_q = q^k k.$$

Thus, we have the following:

$$(3) \quad S_{1,q}(k) = \sum_{j=0}^{k-1} q^j j = \frac{q^k k - q[k]_q}{q-1}.$$

Note that $\frac{k(k-1)}{2} = \lim_{q \rightarrow 1} \frac{q^k k - q[k]_q}{q-1} = S_1(k)$. By the same method of Eq.(1), we easily see that

$$(4) \quad q^{j+1}(j+1)^2 - q^j j^2 = (q-1)q^j j^2 + 2q^{j+1}j + q^{j+1}.$$

Thus, we obtain

$$(5) \quad S_{2,q}(k) = \sum_{j=0}^{k-1} q^j j^2 = \frac{q^k k^2}{q-1} - 2q \frac{q^k k - q[k]_q}{(q-1)^2} - \frac{q[k]_q}{q-1}.$$

From the simple calculation, we note that

$$(6) \quad \begin{aligned} q^{j+1}(j+1)^3 - q^j j^3 &= (q-1)q^j j^3 + 3q q^j j^2 + 3q q^j j + q^{j+1}, \\ q^k k^3 &= (q-1)S_{3,q}(k) + 3q S_{2,q}(k) + 3q S_{1,q}(k) + q[k]_q. \end{aligned}$$

Thus, we see that

$$(7) \quad S_{3,q}(k) = \frac{q^k k^3}{q-1} - \frac{3q}{q-1} S_{2,q}(k) - \frac{3q}{q-1} S_{1,q}(k) - \frac{q[k]_q}{q-1}.$$

Let n, k be the positive integers ($k > 1$) and let $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l l^n$. Then we have

$$q^{l+1}(l+1)^n - l^{n+1}q^l = q^{l+1} \sum_{i=0}^n \binom{n+1}{i} l^i + (q-1)l^{n+1}q^l.$$

Summing over l from 0 to $k-1$, the left-hand side becomes $q^k k^{n+1}$, but the right-hand side is a linear combination of $S_{i,q}(k)$ as follows:

$$(8) \quad q^k k^{n+1} = q(n+1)S_{n,q}(k) + q \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q}(k) + (q-1)S_{n+1,q}(k).$$

Thus, we obtain the following :

Theorem A. *Let $n, k(> 1)$ be the positive integers. Then we have*

$$(9) \quad S_{n,q}(k) = \frac{k^{n+1}}{n+1} q^{k-1} - \frac{1}{n+1} \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q}(k) + \frac{q-1}{q(n+1)} S_{n+1,q}(k).$$

Remark. Note that

$$\begin{aligned} \lim_{q \rightarrow 1} S_{n,q}(k) &= \lim_{q \rightarrow 1} \left(\frac{k^{n+1}}{n+1} q^{k-1} - \frac{1}{n+1} \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q}(k) + \frac{q-1}{q(n+1)} S_{n+1,q}(k) \right) \\ &= \frac{k^{n+1}}{n+1} - \frac{1}{n+1} \sum_{i=0}^{n-1} \binom{n+1}{i} S_i(k) = S_n(k), \text{ cf. [13].} \end{aligned}$$

3. A FORMULA FOR $S_{n,q}(k)$

In this section, we assume $q \in \mathbb{C}$ with $|q| < 1$. Now, we consider the q -Bernoulli polynomials as follows:

$$(10) \quad \frac{\log q + t}{qe^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q}(x) \frac{t^n}{n!}, \text{ see [7, 8].}$$

In the case $x = 0$, $B_{n,q}(= B_{n,q}(0))$ will be called the q -Bernoulli numbers, see [8, 9]. Let $F_q(t, x) = \frac{\log q + t}{qe^t - 1} e^{xt}$. Then we see that

$$(11) \quad F_q(t, x) = -(t + \log q) \sum_{n=0}^{\infty} e^{(n+x)t} q^n, \quad t \in \mathbb{C} \text{ with } |t| < 2\pi, \text{ cf. [8, 9, 10].}$$

In [8], it was known that

$$(12) \quad B_{n,q}(x) = \sum_{j=0}^n \binom{n}{j} B_{j,q} x^{n-j} = m^{n-1} \sum_{i=0}^{m-1} q^i B_{n,q^m} \left(\frac{x+i}{m} \right), \text{ for } n \geq 0.$$

By (10) and (11), we easily see that the q -Bernoulli numbers can be rewritten as

$$B_{0,q} = \frac{q-1}{\log q}, \quad q(B_q + 1)^k - B_{k,q} = \delta_{k,1}, \text{ for } k \geq 1,$$

where $\delta_{k,1}$ is Kronecker symbol and we use the usual convention about replacing B_q^i by $B_{i,q}$, cf. [8, 9, 10]. From the simple calculation, we note that

$$(13) \quad -\sum_{n=0}^{\infty} e^{(n+k)t} q^n + \sum_{n=0}^{\infty} e^{nt} q^{n-k} = \sum_{l=0}^{\infty} (q^{-k} \sum_{n=0}^{k-1} n^l q^n) \frac{t^l}{l!} = \sum_{l=1}^{\infty} (q^{-k} l \sum_{n=0}^{k-1} n^{l-1} q^n) \frac{t^{l-1}}{l!}.$$

Thus, we have

$$(14) \quad \begin{aligned} \sum_{l=0}^{\infty} (B_{l,q}(k) - q^{-k} B_{k,q}(0)) \frac{t^l}{l!} &= -(t + \log q) \sum_{n=0}^{\infty} e^{(n+k)t} q^n + (t + \log q) \sum_{n=0}^{\infty} e^{nt} q^{n-k} \\ &= \sum_{l=0}^{\infty} (q^{-k} l S_{l-1,q}(k) + (q^{-k} \log q) S_{l,q}(k)) \frac{t^l}{l!}. \end{aligned}$$

By comparing the coefficient on both side in Eq.(14), we obtain the following:

Theorem B. *Let l, k be positive integers ($k > 1$). Then we see that*

$$q^{-k} S_{l-1,q}(k) + (q^{-k} \log q) S_{l,q}(k) = \frac{B_{l,q}(k) - q^{-k} B_{l,q}(0)}{l}.$$

Remark. Note that

$$S_{l-1}(k) = \lim_{q \rightarrow 1} (q^{-k} S_{l-1,q}(k) + (q^{-k} \log q) S_{l,q}(k)) = \frac{B_l(k) - B_l(0)}{l}, \text{ cf. [13],}$$

where $B_l(k)$ are called ordinary Bernoulli polynomials.

REFERENCES

1. T. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Math., Springer-Verlag, New York, 1986.
2. L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
3. J. Faulhaber, *Academia Algebrae, Darinnen die miraculosische inventiones zu den höchsten Cossen weiters continuirt und profitiert werden*, Augspurg, bey Johann Ulrich Schönigs, 1631.
4. K. C. Garrett and K. Hummel, *A combinatorial proof of the sum of q-cubes*, Electronic J. Combinatorics **11** (2004), R 9.
5. G. Gasper, M. Rahman, *Basic hypergeometric series*, Encyclopedia of Mathematics and Its Applications, Vol 96, Secod Edition, Cambridge Univ. Press, 2004.
6. V. J. W. Guo, J. Zeng, *A q-analogue of Faulhaber's formula for sums of powers*, arXiv:math.CO/0501441 (2005).
7. T. Kim, *Sums of powers of consecutive q-integers*, Advan. Stud. Contemp. Math. **9** (2004), 15-18.
8. T. Kim, *A new approach to q-zeta function*, arXiv:math.NT/0502005 (2005).
9. T. Kim, *q-Volkenborn Integration*, Russian J. Math. Phys. **9** (2002), 288-299.
10. T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.
11. D. E. Knuth, *Johann Faulhaber and sums of powers*, Math. Comput. **61** (1993), 277-294.
12. M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts and quints*, Electronic J. Combinatorics **11** (2004), R 71.
13. Y.-Y. Shen, *A note on the sums of powers of consecutive integers*, Tunghai Science **5** (2003), 101-106.
14. S. O. Warnaar, *On the q-analogues of the sums of cubes*, Electronic J. Combinatorics **11** (2004), N 13.