## Noncommutative Geometry and Quiver algebras

WILLIAM CRAWLEY-BOEVEY, PAVEL ETINGOF, AND VICTOR GINZBURG

#### Abstract

We develop a new framework for noncommutative differential geometry based on double derivations. This leads to the notion of moment map and of Hamiltonian reduction in noncommutative symplectic geometry. For any smooth associative algebra B, we define its noncommutative cotangent bundle  $T^*B$ , which is a basic example of noncommutative symplectic manifold. Applying Hamiltonian reduction to noncommutative cotangent bundles gives an interesting class of associative algebras,  $\Pi = \Pi(B)$ , that includes preprojective algebras associated with quivers. Our formalism of noncommutative Hamiltonian reduction provides the space  $\Pi/[\Pi,\Pi]$  with a Lie algebra structure, analogous to the Poisson bracket on the zero fiber of the moment map. In the special case where  $\Pi$  is the preprojective algebra associated with a quiver of non-Dynkin type, we give a complete description of the Gerstenhaber algebra structure on the Hochschild cohomology of  $\Pi$  in terms of the Lie algebra  $\Pi/[\Pi,\Pi]$ .

Table of Contents

- 1. Introduction
- 2. Calculus of double derivations
- 3. The derivation  $\Delta$
- 4. Hamilton operators
- 5. Noncommutative cotangent bundle
- 6. The representation functor
- 7. Hamiltonian reduction in Noncommutative Geometry
- 8. The necklace Lie algebra
- 9. Hochschild cohomology of preprojective algebras
- $10. \hspace{0.5cm} {\tt Deformations} \hspace{0.1cm} {\tt of} \hspace{0.1cm} {\tt preprojective} \hspace{0.1cm} {\tt algebras} \\$
- 11. Representation schemes

## 1 Introduction

1.1 This paper is devoted to the general formalism of noncommutative differential and symplectic geometry. Noncommutative symplectic geometry was introduced by Kontsevich [Ko] and further studied in [BLB], [Gi], where connections with quivers were discovered.

In the present paper, we develop a different 'double version' of noncommutative geometry. This version has much richer structure; it allows, in particular, to deal with noncommutative analogues of the Gerstenhaber algebra structure on polyvector fields. Roughly speaking, Kontsevich's version is obtained from the double version by applying the commutator quotient construction. This procedure is somewhat analogous to replacing a chain complex by its homology, so a lot of information is lost in this process.

It may also be mentioned that the theory developed below plays an important role in the mathematical formalism of 'open string' theory, cf. [La].

1.2 Let A be an arbitrary associative, not necessarily commutative, algebra. We extend the standard constructions of differential geometry like Lie derivative or contraction of a differential form with respect to a vector field, to noncommutative geometry. The role of differential forms is played by the so-called Karoubi-de Rham complex  $DR^{\bullet}A$ . Derivations  $A \to A$  act naturally on  $DR^{\bullet}A$  via Lie derivative and contraction operators, cf. [Lo]. In our 'double version' approach, the role of vector fields is played by double derivations  $A \to A \otimes A$  rather than ordinary derivations  $A \to A$ . One of the key technical points of the present paper is a construction of Lie derivative and contraction operators for double derivations.

For any integer  $n \geq 1$ , one has a representation functor

 $\operatorname{Rep}_n: \operatorname{Associative algebras} \longrightarrow \operatorname{Affine schemes},$ 

that assigns to an associative algebra A the scheme  $\operatorname{Rep}_n A$  of all n-dimensional A-modules, that is, of all homomorphisms of A into the associative algebra of  $n \times n$ -matrices. A general philosophy, due in particular to Kontsevich, says that meaningful concepts of noncommutative geometry should go, under the representation functor, to their commutative counterparts. Thus, elements of the Karoubi-de Rham complex  $\operatorname{DR}^{\bullet} A$  go to differential forms on the scheme  $\operatorname{Rep}_n A$ , and derivations  $A \to A$  go to vector fields on  $\operatorname{Rep}_n A$ . The meaning of double derivations is more subtle, it is discussed in Sect. 6.3.

In Section 3 we develop a 'double' version of noncommutative symplectic geometry. Thus, our notion of symplectic 2-form involves double derivations and is different from the one introduced by Kontsevich [Ko] and used in [Gi], [BLB]. Given an algebra A equipped with a symplectic 2-form, one has a natural Lie algebra structure on the vector space A/[A,A], analogous to the Poisson bracket on the space of functions on a symplectic manifold. If A is smooth, then  $\operatorname{Rep}_n A$  turns out to be a symplectic manifold. Elements of A/[A,A] go, under the representation functor, to regular functions on  $\operatorname{Rep}_n A$ , and this map is compatible with the Lie brackets.

For any algebra A, there is a distinguished double derivation  $\Delta: a \longmapsto a \otimes 1 - 1 \otimes a$ . Contraction with  $\Delta$  gives rise to a very interesting new operation on the Karoubi-de Rham complex  $\mathrm{DR}^{\bullet}A$ . This operation is closely related to the cyclic homology of A, see [Gi2]. Using contraction with  $\Delta$ , we also define a map sending closed 2-forms in  $\mathrm{DR}^2A$  to elements of A (up to a constant summand from the ground field). This map is referred to as noncommutative moment map.

Given a symplectic 2-form  $\omega \in \mathrm{DR}^2 A$ , we let  $\mathbf{w}$  be the image of  $\omega$  under the noncommutative moment map. We show that the function on  $\mathrm{Rep}_n A$  corresponding to the element  $\mathbf{w}$  gives rise to a moment map  $\mathrm{Rep}_n A \to (\mathfrak{gl}_n)^*$ , see Theorem 6.4.3. The algebra  $A_{\mathbf{w}} = A/A\mathbf{w}A$ , the quotient of A by the two-sided ideal generated by  $\mathbf{w}$ , is called a Hamiltonian reduction of A at  $\mathbf{w}$  since the scheme  $\mathrm{Rep}_n(A_{\mathbf{w}})$  may be identified with the zero fiber of the moment map.

In general, the symplectic 2-form on A does not give a symplectic form on the Hamiltonian reduction. The resulting structure on  $A_{\mathbf{w}}$  is weaker; it is

conveniently expressed in terms of the notion of  $Hamilton\ operator$ , see Sect. 4.3, borrowed from the works of Gelfand and Dorfman [GD]. One of the main results of our paper says that the Lie bracket on A/[A,A] descends to  $A_{\mathbf{w}}/[A_{\mathbf{w}},A_{\mathbf{w}}]$ , see Proposition 4.4.3 and Theorem 7.2.3. It should be emphasized that the 'double version' approach seems to be absolutely indispensable for the above constructions and results.

For any smooth associative algebra B, we define its noncommutative cotangent bundle. This is an associative algebra  $T^*B$  that comes equipped with a canonical symplectic 2-form  $\omega \in \mathrm{DR}^2(T^*B)$ . Each element of  $T^*B$  gives rise to a regular function on  $T^*(\mathrm{Rep}_n B)$ , the cotangent bundle on the scheme  $\mathrm{Rep}_n B$ . Furthermore, the 2-form  $\omega$  goes, under the representation functor, to the standard symplectic 2-form on  $T^*(\mathrm{Rep}_n B)$ .

Applying Hamiltonian reduction to noncommutative cotangent bundles gives an interesting class of associative algebras,  $\Pi = \Pi(B)$ , that includes preprojective algebras associated with quivers. This special case is considered in more detail in section 8. By our general result, the space  $\Pi/[\Pi,\Pi]$  acquires a canonical Lie algebra structure. Theorem 8.4.1, which is the second main result of the paper, gives, for quivers of non-Dynkin and not extended Dynkin type, a complete description of the Gerstenhaber algebra structure on the Hochschild cohomology of  $\Pi$  in terms of the Lie algebra  $\Pi/[\Pi,\Pi]$ .

Our last important result is a computation of the center of the corresponding Poisson algebra  $\operatorname{Sym}(\Pi/[\Pi,\Pi])$ , see Theorem 8.6.1. This should be viewed as an 'additive analogue' of a similar result for Goldman's Lie algebra associated to a Riemann surface, see [E]. The proof depends heavily on the results from [CB1] about *quiver varieties*.

Our approach to noncommutative geometry was partly motivated by [CB2], and is quite close to Van den Bergh's work [VB2]. Specifically, Van den Bergh considered a noncommutative analogue of Poisson geometry while in the present paper we mostly deal with a noncommutative analogue of symplectic geometry. Although symplectic geometry is a special case of Poisson geometry, the two papers have almost no overlap and (apart from ideological motivation originating from quiver theory) are quite independent.

For more details about the connection between Van den Bergh's work and the present paper the reader is referred to the Appendix to [VB2].

1.3 Acknowledgements. We are very grateful to Michel Van den Bergh for useful discussions and for a careful reading of the manuscript. The second and third author are partially supported by the NSF grants DMS-9988796 and DMS-0303465, respectively, and also by the CRDF grant RM1-2545-MO-03.

## 2 Calculus of double derivations

**2.1 Notation.** Throughout, we fix a field k of characteristic 0 and write 0 = 0, Hom = Hom<sub>k</sub>, etc.

Given an associative k-algebra A, let  $A^{\mathrm{op}}$  denote the opposite algebra, and write  $A^e = A \otimes A^{\mathrm{op}}$ . There is a canonical isomorphism  $(A^e)^{\mathrm{op}} \cong A^e$ . Thus, an A-bimodule is the same thing as a left  $A^e$ -module, and also the same thing as a right  $A^e$ -module.

The space  $A \otimes A$  has two *commuting* A-bimodule structures, called the *outer*, resp., *inner*, bimodule structure. These two bimodule structures are given by

$$b(a'\otimes a'')c:=(ba')\otimes (a''c), \quad \text{resp.}, \quad b(a'\otimes a'')c:=(a'c)\otimes (ba''), \quad a',a'',b,c\in A.$$

Fix a unital associative k-algebra R. Throughout, by an R-algebra we mean an associative unital k-algebra equipped with a unit preserving k-algebra imbedding  $R \to A$ . A morphism of R-algebra is meant to be an algebra homomorphism compatible with the identity map  $R \to R$ ; in particular, any R-algebra morphism is unit preserving.

Let A be an R-algebra and  $A \otimes_R A \xrightarrow{\mathbf{m}} A$ ,  $a' \otimes a'' \mapsto a'a''$ , the multiplication map. Let  $\Omega_R^1 A := \mathrm{Ker}(\mathbf{m})$  be the A-bimodule of noncommutative relative 1-forms on A (with respect to the subalgebra R), see [CQ, Sect. 2]. If A is finitely generated as an R-algebra then  $\Omega_R^1 A$  is finitely generated as a left  $A^e$ -module.

**Definition 2.1.1.** An R-algebra A is called *smooth over* R if it is finitely generated as an R-algebra and  $\Omega_R^1 A$  is projective as a left  $A^e$ -module.

Path algebras associated to quivers, to be studied later in this paper, are typical examples of smooth algebras.

Given an A-bimodule M, write  $\operatorname{Der}_R(A,M)$  for the space of R-linear derivations, that is, derivations  $\theta:A\to M$  such that  $\theta(R)=0$ . There is a canonical 'universal' derivation  $d:A\to \Omega^1_RA$ ,  $a\mapsto da:=a\otimes 1-1\otimes a$ , such that, for any A-bimodule M, we have a bijection

$$\operatorname{Der}_R(A, M) \xrightarrow{\sim} \operatorname{Hom}_{A^e}(\Omega^1_R A, M), \ \theta \mapsto i_{\theta} \ \text{where} \ i_{\theta}(u \, dv) := u \cdot \theta(v). \ (2.1.2)$$

In the 'absolute' case  $R = \mathbb{k}$ , we will use unadorned notation  $\operatorname{Der} A = \operatorname{Der}_{\mathbb{k}} A$ ,  $\Omega^1 A = \Omega^1_B A$ , etc. We say that A is *smooth* if A is smooth over  $\mathbb{k}$ .

**2.2 Separability elements.** In this subsection, as well as in most sections of the paper, we assume R to be a finite dimensional semisimple k-algebra, e.g., a direct sum of matrix algebras.

For such an algebra, there exists a canonical symmetric separability element  $\mathbf{e} = \sum_i e_i \otimes e^i \in R \otimes R$  such that the following holds, cf. [CQ, Prop. 4.2]:

$$(\mathrm{i}) \ r \cdot \mathbf{e} = \mathbf{e} \cdot r, \ \forall r \in R; \qquad (\mathrm{ii}) \ \sum\nolimits_i e_i \cdot e^i = 1; \qquad (\mathrm{iii}) \ \mathbf{e} = \mathbf{e}^\mathrm{op}, \qquad (2.2.1)$$

where in equation (i) we write  $r \cdot \mathbf{e} := \sum_i (r \cdot e_i) \otimes e^i$ , resp.,  $\mathbf{e} \cdot r := \sum_i e_i \otimes (e^i \cdot r)$ , and in equation (iii) we let  $(x \otimes y)^{\text{op}} := y \otimes x$  denote the flip involution on  $R \otimes R$ . We will often use Sweedler's notation and write  $\mathbf{e} = \mathbf{e}' \otimes \mathbf{e}''$  for the sum  $\sum_i e_i \otimes e^i$ .

Given an R-bimodule M and a (not necessarily R-stable)  $\Bbbk$ -vector subspace  $V \subset M$ , we write  $V^R = \{v \in V \mid rv = vr, \forall r \in R\}$  for the centralizer of R in V,

and put  $\mathbf{e} \cdot V := \{\mathbf{e} \cdot v \mid v \in V\}$ , where  $\mathbf{e} \cdot v$  stands for the left  $R^e$ -action. Also, let  $[R, M] \subset M$  be the  $\mathbb{k}$ -linear span of the set  $\{rm - mr \mid r \in R, m \in M\}$ . Thus, [R, M] is a vector subspace in M.

Equation (2.2.1)(i) means that  $\mathbf{e} \in (R \otimes R)^R$ , where  $R \otimes R$  is viewed as an R-bimodule with respect to the outer bimodule structure. The symmetry property (2.2.1)(iii) insures that the same holds for the inner bimodule structure as well.

Properties (2.2.1) yield the following result, cf. [CQ, (43)].

**Proposition 2.2.2.** (i) For any R-bimodule M we have

$$M = M^R \oplus [R, M]$$
, and  $\mathbf{e} \cdot M = M^R \xrightarrow{\sim} M/[R, M]$ ,

where the isomorphism on the right is the composite  $M^R \hookrightarrow M \twoheadrightarrow M/[R, M]$ . (ii) The projection  $M \to M^R$  along [R, M] is given by the formula:

$$M \to \mathbf{e} \cdot M, \quad m \longmapsto \sum_{i} e_i \cdot m \cdot e^i = \mathbf{e}' \cdot m \cdot \mathbf{e}'' = \mathbf{e}'' \cdot m \cdot \mathbf{e}'.$$

Let A be an R-algebra and M an A-bimodule. Clearly,  $[R, M] \subset [A, M]$ . Applying Proposition 2.2.2 we deduce

$$M = M^R + [A, M], \text{ and } \mathbf{e} \cdot [A, M] = [A, M]^R.$$
 (2.2.3)

Observe further that, for R as above, any R-algebra A that is smooth over  $\mathbbm{k}$  is also smooth over R. To see this, note that the map  $r \mapsto r \cdot \mathbf{e} = \mathbf{e} \cdot r$  provides a section of the R-bimodule map  $R \otimes R \to R$ , of multiplication. Tensoring over R on each side with A, we deduce that the A-bimodule map  $A \otimes A \to A \otimes_R A$  has a section. Thus,  $\Omega^1_R A$  is a direct summand of  $\Omega^1 A$ , and our claim follows.

**2.3 Double derivations.** Fix an R-algebra A. We consider R-linear derivations  $\Theta: A \to A \otimes A$ , where we view  $A \otimes A$  as an A-bimodule with respect to the outer bimodule structure. Put  $\mathbb{D}\mathbf{er}_R A := \mathrm{Der}_R(A, A \otimes A)$ . The inner bimodule structure on  $A \otimes A$  gives  $\mathbb{D}\mathbf{er}_R A$  a natural left  $A^e$ -module structure. Equivalently, we may view a double derivation  $\Theta \in \mathbb{D}\mathbf{er}_R A$  as a derivation  $A \to A^e$ , where  $A^e$  is viewed as an A-bimodule corresponding to left multiplication of  $A^e$  on itself. From this point of view, the  $A^e$ -module structure on  $\mathbb{D}\mathbf{er}_R A$  described above comes from right multiplication of  $A^e$  on itself.

Let M be an A-bimodule and  $u \in M$ . The map  $u_*: A \otimes A \to M, a' \otimes a'' \to a'ua''$  is a morphism of A-bimodules. Applying the functor  $\operatorname{Der}_R(A, -)$  to this morphism yields a map  $\mathbb{D}\mathbf{er}_R A = \operatorname{Der}_R(A, A \otimes A) \to \operatorname{Der}_R(A, M)$ . It is straightforward to verify that this way one obtains a well-defined map

$$(\mathbb{D}\mathbf{er}_R A) \otimes_{A^e} M \longrightarrow \mathrm{Der}_R(A, M), \quad \Theta \otimes u \longmapsto u_* \circ \Theta, \tag{2.3.1}$$

where the tensor product on the left is taken with respect to the A-bimodule structure on  $\mathbb{D}\mathbf{er}_R A$ .

In the special case M=A, we put  $\operatorname{Der}_R A:=\operatorname{Der}_R(A,A)$ . The multiplication map  $\mathbf{m}:A\otimes A\to A, a'\otimes a''\mapsto a'a''$ , is a morphism of A-bimodules

with respect to the outer bimodule structure on  $A \otimes A$  (but *not* with respect to the inner structure), hence, induces a map  $\mathbf{m}_* : \mathbb{D}\mathbf{er}_R A \to \mathrm{Der}_R A$ . Using the notation of (2.3.1), one can write  $\mathbf{m}_* : \Theta \mapsto 1_* \circ \Theta$ .

**Proposition 2.3.2.** Let R be a finite dimensional semisimple algebra. For any smooth R-algebra A the map (2.3.1) is a bijection,  $\mathbb{D}\mathbf{er}_R A$  is a finitely generated projective  $A^e$ -module, and there is a canonical short exact sequence

$$0 \longrightarrow \operatorname{Hom}_{A^e}(\Omega^1_R A, \Omega^1 A) \longrightarrow \mathbb{D}\mathbf{er}_R A \xrightarrow{\mathbf{m}_*} \operatorname{Der}_R A \longrightarrow 0.$$

*Proof.* For any R-algebra A and A-bimodule M, we have a canonical map

$$\operatorname{Hom}_{A^e}(\Omega^1_R A, A \otimes A) \otimes_{A^e} M \to \operatorname{Hom}_{A^e}(\Omega^1_R A, M).$$
 (2.3.3)

Using the universal property of  $\Omega_R^1 A$  we rewrite this map as  $(\mathbb{D}\mathbf{er}_R A) \otimes_{A^e} M \to \mathrm{Der}_R(A, M)$ . It is easy to check that the latter map is nothing but (2.3.1).

Now, if A is smooth over R, then  $\Omega_R^1 A$  is a finitely generated projective left  $A^e$ -module. Thus,  $\Omega_R^1 A$  is a direct summand of a finite rank free  $A^e$ -module. It follows that  $\operatorname{Hom}_{A^e}(\Omega_R^1 A, A \otimes A) = \mathbb{D}\mathbf{er}_R A$  is also a direct summand of a finite rank free  $A^e$ -module, hence, projective. Furthermore, we conclude that the map in (2.3.3) is a bijection because a similar map for  $\Omega_R^1 A$  being replaced by a finite rank free  $A^e$ -module is clearly bijective.

Finally, if A is smooth then, applying exact functor  $\operatorname{Hom}_{A^e}(\Omega^1_R A, -)$  to the short exact sequence of A-bimodules  $\Omega^1 A \hookrightarrow A \otimes A \stackrel{\mathbf{m}}{\twoheadrightarrow} A$  yields the short exact sequence of the Proposition.

**2.4** Double-derivations for a free algebra. Let  $R = \mathbb{k}$ , fix an integer  $n \geq 1$  and set  $A = \mathbb{k}\langle x_1, \ldots, x_n \rangle$ , a free associative  $\mathbb{k}$ -algebra on n generators. An element  $f \in \mathbb{k}\langle x_1, \ldots, x_n \rangle$  may be thought of as a polynomial function in n non-commuting variables  $x_1, \ldots, x_n$ .

For each i = 1, ..., n, we introduce a double derivation  $\partial_i \in \mathbb{D}\mathbf{er} A$  defined on generators by the formula:

$$\partial_i(x_j) = \begin{cases} 1 \otimes 1 & \text{if } i = j \\ 0 & \text{if } i \neq j. \end{cases}$$

It is easy to see that  $\mathbb{D}$ er A is a free left  $A^e$ -module with basis  $\{\partial_i, i = 1, \ldots, n\}$ . Next, let  $f_1, \ldots, f_n \in \mathbb{k}\langle x_1, \ldots, x_n \rangle$  be an n-tuple of elements of A. We write  $F = (f_1, \ldots, f_n)$  and think of F as a self-map of an n-dimensional 'noncommutative affine space'.

We may identify the space  $A \otimes A$  with  $A^e$ , and define the *Jacobi matrix* for the map F to be the following  $A^e$ -valued  $n \times n$ -matrix

$$DF = \|\partial_i(f_i)\|_{i,i=1,...,n} \in \text{Mat}_n(A^e).$$
 (2.4.1)

Now, let  $F=(f_1,\ldots,f_n)$  and  $G=(g_1,\ldots,g_n)$  be two *n*-tuples of elements of A. Let  $G\circ F$  be the 'composite' *n*-tuple obtained by substituting the elements  $f_1,\ldots,f_n\in A$  into the arguments of the noncommutative polynomials

 $g_1, \ldots, g_n$ . Equivalently put, giving an n-tuple  $F = (f_1, \ldots, f_n)$  is the same thing as giving an algebra homomorphism  $F: A \to A$  such that  $x_i \mapsto f_i, i = 1, \ldots, n$ . In this interpretation, the composite  $G \circ F$  corresponds to composing algebra homomorphisms.

Let  $P, Q \mapsto P \star Q$  denote multiplication in the algebra  $\operatorname{Mat}_n(A^e)$ ; note that it differs from multiplication in  $\operatorname{Mat}_n(A \otimes A)$ .

The following result is proved by a straightforward computation.

**Proposition 2.4.2 (Chain rule).** For any two algebra homomorphisms 
$$F, G : A \to A$$
, in  $\operatorname{Mat}_n(A^e)$  one has  $D(G \circ F) = (DG)(F) \star DF$ .

In the right-hand side of the equation above, the matrix (DG)(F) is obtained by applying the homomorphism  $F \otimes F : A \otimes A \to A \otimes A$  to each entry of the matrix  $DG \in \operatorname{Mat}_{n \times n}(A^e)$ .

We remark that the above Proposition has no analogue for ordinary derivations  $A \to A$  instead of double derivations; cf. also [Vo] for closely related constructions.

**2.5** Karoubi-de Rham complex. Let A be an R-algebra. The tensor algebra,  $T_A^{\bullet}(\Omega_R^1 A) = \bigoplus_{n \geq 0} T_A^n(\Omega_R^1 A)$  (tensor product over A), of the A-bimodule  $\Omega_R^1 A$  is a DG algebra  $(\Omega_R^{\bullet} A, d)$ , called the algebra of noncommutative relative differential forms on A. We have an isomorphism of left A-modules  $\Omega_R^n A = A \otimes_R T_R^n(A/R)$ , see [CQ]; usually, one writes  $a_0 da_1 da_2 \dots da_n \in \Omega_R^n A$  for the n-form corresponding to an element  $a_0 \otimes (a_1 \otimes \ldots \otimes a_n) \in A \otimes_R T_R^n(A/R)$  under this isomorphism.

If R is a finite dimesional semisimple algebra, then the DG algebra  $(\Omega_R^{\bullet}A, d)$  is known to be acyclic in positive degrees. Indeed, let  $\pi: A \to A/R$  denote the projection. The differential  $d_n: \Omega_R^n A \to \Omega_R^{n+1} A$  corresponds, under the isomorphism  $\Omega_R^{\bullet}A = A \otimes_R T_R^{\bullet}(A/R)$ , to the composite

$$A \otimes_R T_R^n(A/R) \xrightarrow{\pi \otimes \operatorname{Id}^{\otimes n}} T_R^{n+1}(A/R) \xrightarrow{\sim} R \otimes_R T_R^{n+1}(A/R) \hookrightarrow A \otimes_R T_R^{n+1}(A/R),$$

(we have used here that tensoring over R is an exact functor since R is a finite dimensional semisimple algebra).

From this definition of the differential, it is clear that  $Im(d_n) = Ker(d_{n+1})$ ; thus, we have

$$(\Omega_R^k A)_{\text{closed}} = (\Omega_R^k A)_{\text{exact}} := d(\Omega_R^{k-1} A), \quad \forall k \ge 1.$$
 (2.5.1)

Next, following Karoubi, we define

$$\mathrm{DR}_R^\bullet A := \Omega_R^\bullet A / [\Omega_R^\bullet A, \Omega_R^\bullet A]_{\mathrm{super}},$$

where  $[-,-]_{\text{super}}$  denotes the  $\mathbb{k}$ -linear span of all supercommutators. This is the noncommutative de Rham complex of A with de Rham differential  $d: DR_R^j A \to DR_R^{j+1} A$ , cf. also [Lo] for more details (in the case  $R = \mathbb{k}$ ).

For any A-bimodule M, we have the commutator space [A, M], and we let  $M_{\natural} := M/[A, M] = A \otimes_{A^e} M$  denote the corresponding commutator quotient.

We have a natural projection  $(\Omega_R^{\bullet}A)_{\natural} \to DR_R^{\bullet}A$  that becomes an isomorphism:

 $DR_R^j A = (\Omega_R^j A)_{\natural}, \text{ for } j = 0, 1.$ 

For  $j \geq 2$ , the projection  $(\Omega_R^j A)_{\natural} \to \mathrm{DR}_R^j A$  is not an isomorphism, in general. Sometimes, to distinguish between  $\Omega_R^{\bullet} A$  and  $\mathrm{DR}_R^{\bullet} A$  we will write  $\llbracket \omega \rrbracket$  for the image of an element  $\omega \in \Omega_R^{\bullet} A$  under the projection  $\Omega_R^{\bullet} A \to \mathrm{DR}_R^{\bullet} A$ .

The following standard result will be proved at the end of Sect. 2.7.

**Lemma 2.5.2 (Homotopy invariance).** Let  $A = \bigoplus_{k \geq 0} A_k$  be a graded R-algebra such that  $R \subset A_0$ . Then, the algebra imbedding  $A_0 \hookrightarrow A$  induces an isomorphism of de R-ham cohomology  $H^{\bullet}(\mathrm{DR}_R^{\bullet}A_0, d) \stackrel{\sim}{\to} H^{\bullet}(\mathrm{DR}_R^{\bullet}A, d)$ .

For any vector space  $V \subset A$ , we write  $dV = \{dv, v \in V\}$ , a subspace in  $\Omega^1_R A$ . Given an R-subalgebra  $B \subset A$ , there is a canonical DG algebra isomorphism  $\Omega^{\bullet}_B A \cong \Omega^{\bullet}_R A / \Omega^{\bullet}_R A \cdot dB \cdot \Omega^{\bullet}_R A$ . Similarly, given a two-sided ideal  $I \subset A$ , one has a DG algebra isomorphism, see [CQ]:

$$\Omega_R^{\bullet}(A/I) \cong \Omega_R^{\bullet} A / (\Omega_R^{\bullet} A \cdot I \cdot \Omega_R^{\bullet} A + \Omega_R^{\bullet} A \cdot dI \cdot \Omega_R^{\bullet} A). \tag{2.5.3}$$

Remark 2.5.4. In general, the induced map  $R \to A/I$  is not necessarily injective. So, according to our definition, A/I may not be an R-algebra. Such a situation may arise when we consider Hamiltonian reduction in noncommutative geometry (section 6.4 below). However, the only reason for insisting that R be injectively mapped into any R-algebra A is a frequent use of the notation A/R. So, in those rare occasions where the map  $R \to A$  fails to be injective, the symbol A/R should be understood as the quotient of A by the image of R in A.  $\Diamond$ 

**2.6 Contraction with a double derivation.** We will use lower case Greek letters  $\theta, \xi, \ldots$  to denote derivations  $A \to A$ , and upper case Greek letters,  $\Theta, \Xi, \ldots$  to denote double derivations  $A \to A \otimes A$ .

Any derivation  $\theta \in \operatorname{Der}_R A$  gives rise to contraction (with  $\theta$ ) maps  $i_{\theta}: \Omega_R^k A \to \Omega_R^{k-1} A$ , resp.,  $\operatorname{DR}_R^k A \to \operatorname{DR}_R^{k-1} A$ . The map  $i_{\theta}$  is defined on 1-forms by formula (2.1.2) and is extended to a map  $\Omega_R^{\bullet} A \to \Omega_R^{\bullet -1} A$  as a *super*-derivation.

Now let  $\Theta \in \mathbb{D}\mathbf{er}_R A$ . For any 1-form  $\alpha \in \Omega^1_R A$ , contraction with  $\Theta$  gives an A-bimodule map, see (2.1.2):

$$i_{\Theta}: \ \Omega^1_R A \longrightarrow A \otimes A, \quad \alpha \longmapsto i_{\Theta} \alpha = i'_{\Theta} \alpha \otimes i''_{\Theta} \alpha.$$
 (2.6.1)

Here and below, we will systematically use symbolic Sweedler's notation to write  $i_{\Theta}\alpha = i'_{\Theta}\alpha \otimes i''_{\Theta}\alpha$  (omitting the summation symbol) for an element in the tensor product. Similarly, we write the map  $\Theta: A \to A \otimes A$  as  $a \mapsto \Theta'(a) \otimes \Theta''(a)$ .

As usual, one may uniquely extend the map  $i_{\Theta}$  in (2.6.1) to higher degree differential forms by requiring that  $i_{\Theta}$  be a *super-derivation* of degree (-1). This way, we obtain a map  $i_{\Theta}: \Omega_R^{\bullet}A \to \Omega_R^{\bullet}A \otimes \Omega_R^{\bullet}A$ , which is a super-derivation of the graded algebra  $\Omega_R^{\bullet}A$  with coefficients in  $\Omega_R^{\bullet}A \otimes \Omega_R^{\bullet}A$ , viewed as an  $\Omega_R^{\bullet}A$ -bimodule with respect to the outer bimodule structure.

Explicitly, for any n = 1, 2, ..., and  $\alpha_1, ..., \alpha_n \in \Omega^1_R A$ , we get

$$i_{\Theta}(\alpha_1 \alpha_2 \dots \alpha_n) = \sum_{1 \le k \le n} (-1)^{k-1} \cdot (\alpha_1 \dots \alpha_{k-1} (i'_{\Theta} \alpha_k)) \otimes ((i''_{\Theta} \alpha_k) \alpha_{k+1} \dots \alpha_n).$$

$$(2.6.2)$$

It is often convenient to view the contraction map  $i_{\Theta}$  as a map  $\Omega_R^{\bullet}A \to T_{\Bbbk}^{\bullet}(\Omega_R^{\bullet}A)$ , and to extend the latter map further to get a super-derivation of the tensor algebra  $T_{\Bbbk}^{\bullet}(\Omega_R^{\bullet}A)$ .

With this understood, one has the following standard identities.

### Lemma 2.6.3. We have

$$i_{\Theta}(\alpha\beta) = (i_{\Theta}\alpha)\beta + (-1)^{\deg \alpha}\alpha (i_{\Theta}\beta), \quad \forall \alpha, \beta \in \Omega_R^{\bullet}A;$$
  
$$i_{\Phi} \circ i_{\Theta} + i_{\Theta} \circ i_{\Phi} = 0, \quad \forall \Phi, \Theta \in \mathbb{D}\mathbf{er}_R A.$$

The meaning as well as the proof of the second identity of the Lemma is illustrated by the following computation,

$$i_{\Phi} \circ i_{\Theta}(\alpha_{1}\alpha_{2}) = i_{\Phi} \left( i'_{\Theta}\alpha_{1} \otimes (i''_{\Theta}\alpha_{1}) \alpha_{2} - \alpha_{1} (i'_{\Theta}\alpha_{2}) \otimes i''_{\Theta}\alpha_{2} \right)$$
$$= i'_{\Theta}\alpha_{1} \otimes (i''_{\Theta}\alpha_{1}) (i'_{\Phi}\alpha_{2}) \otimes i''_{\Phi}\alpha_{2} - i'_{\Phi}\alpha_{1} \otimes (i''_{\Phi}\alpha_{1}) (i'_{\Theta}\alpha_{2}) \otimes i''_{\Theta}\alpha_{2}.$$

The expression for  $i_{\Theta} \circ i_{\Phi}(\alpha_1 \alpha_2)$  is obtained from this by switching the roles of  $\Theta$  and  $\Phi$ . But the flip  $\Theta \leftrightarrow \Phi$  takes the last line in the displayed formula above to its negative, and (2.6.3) follows.

**2.7** Lie derivative. For any derivation  $\theta \in \operatorname{Der}_R A$ , there is a standard Lie derivative operator  $L_{\theta}$  acting on various objects associated naturally with the algebra A, e.g., an operator  $L_{\theta} : \mathbb{D}\mathbf{er}_R A \to \mathbb{D}\mathbf{er}_R A$ . There is also a derivation  $L_{\theta} : \Omega_R^{\bullet} A \to \Omega_R^{\bullet} A$ , and the corresponding induced map  $L_{\theta} : \operatorname{DR}_R^{\bullet} A \to \operatorname{DR}_R^{\bullet} A$ . All standard formulas involving de Rham differential, contraction and Lie derivative operators hold in the Karoubi-de Rham setting. In particular, one has the Cartan identity  $L_{\theta} = d \circ i_{\theta} + i_{\theta} \circ d$ , for any  $\theta \in \operatorname{Der}_R A$ .

Now, given a double derivation  $\Theta \in \mathbb{D}\mathbf{er}_R A$ , we define the corresponding Lie derivative map as follows:

$$L_{\Theta}: \Omega_{R}^{1}A \longrightarrow (A \otimes \Omega_{R}^{1}A) \bigoplus (\Omega_{R}^{1}A \otimes A), \quad \alpha \longmapsto L_{\Theta}\alpha, \text{ where}$$
  
$$L_{\Theta}(x dy) := \Theta'(x) \otimes \Theta''(x) dy + (x d\Theta'(y)) \otimes \Theta''(y) + x \Theta'(y) \otimes d\Theta''(y).$$

We may naturally extend the map  $L_{\Theta}$  to a degree preserving derivation

$$L_{\Theta}: \ \Omega_R^n A \to \bigoplus_{0 \le k \le n} \Omega_R^k A \otimes \Omega_R^{n-k} A,$$

of the graded algebra  $\Omega_R^{\bullet}A$  with coefficients in  $\Omega_R^{\bullet}A\otimes\Omega_R^{\bullet}A$ , viewed as an  $\Omega_R^{\bullet}A$ -

bimodule with respect to the outer bimodule structure. Explicitly, we have

$$L_{\Theta}(a_0 da_1 \dots da_n) := \Theta'(a_0) \otimes \Theta''(a_0) da_1 \dots da_n +$$

$$+ \sum_{1 \le k \le n} \left( a_0 da_1 \dots da_{k-1} d\Theta'(a_k) \otimes \Theta''(a_k) da_{k+1} \dots da_n \right)$$

$$+ a_0 da_1 \dots da_{k-1} \Theta'(a_k) \otimes d\Theta''(a_k) da_{k+1} \dots da_n$$

As in the case of contractions, it is often convenient to view the Lie derivative as a map  $L_{\Theta}: \Omega_R^{\bullet}A \to T_{\Bbbk}^{\bullet}(\Omega_R^{\bullet}A)$ , and to extend the latter map further as a derivation of the tensor algebra  $T_{\Bbbk}^{\bullet}(\Omega_R^{\bullet}A)$ . Similarly, given  $\theta \in \operatorname{Der}_R A$ , we extend the Lie derivative  $L_{\theta}$  as a derivation of the tensor algebra  $T_{\Bbbk}^{\bullet}(\Omega_R^{\bullet}A)$ ; we may also extend the differential  $d: \Omega_R^{\bullet}A \to \Omega_R^{\bullet+1}A$  as a super-derivation of the tensor algebra  $T_{\Bbbk}^{\bullet}(\Omega_R^{\bullet}A)$ .

With these definitions, it is straightforward to verify the identities

$$[L_{\xi}, i_{\Theta}] = i_{L_{\xi}(\Theta)}$$
 and  $[L_{\xi}, L_{\Theta}] = L_{L_{\xi}(\Theta)}$ ,  $\forall \Theta \in \mathbb{D}\mathbf{er}_{R} A, \xi \in \mathrm{Der}_{R} A$ .

One also has the following Cartan formula for double derivations:

$$d \circ i_{\Theta} + i_{\Theta} \circ d = L_{\Theta}, \quad \forall \Theta \in \mathbb{D}\mathbf{er}_{R} A.$$
 (2.7.2)

It follows in particular that the Lie derivative commutes with the de Rham differential:

$$d \circ L_{\Theta} = d \circ d \circ i_{\Theta} + d \circ i_{\Theta} \circ d = d \circ i_{\Theta} \circ d = d \circ i_{\Theta} \circ d + i_{\Theta} \circ d \circ d = L_{\Theta} \circ d.$$

Remark 2.7.3. It seems very likely that, given  $\Theta, \Phi \in \mathbb{D}\mathbf{er}_R A$ , the map

$$L_{\Theta} \circ L_{\Phi} - L_{\Phi} \circ L_{\Theta} : \Omega_{R}^{\bullet} A \longrightarrow \Omega_{R}^{\bullet} A \otimes \Omega_{R}^{\bullet} A \otimes \Omega_{R}^{\bullet} A$$

is equal to the (appropriately defined) Lie derivative with respect to  $\{\Theta, \Phi\}$ , the Schouten double bracket of  $\Theta$  and  $\Phi$ , introduced by Van den Bergh [VB2].  $\Diamond$ 

Proof of Lemma 2.5.2. The grading on A gives rise to the Euler derivation  $\mathsf{E}:A\to A$ , defined by  $\mathsf{E}|_{A_k}=k\cdot \mathrm{Id},\,k=0,1,\ldots$  The action of the corresponding Lie derivative operator  $L_\mathsf{E}:\mathrm{DR}_R^\bullet A\to \mathrm{DR}_R^\bullet A$  has nonnegative integral eigenvalues. It is clear that the zero weight subspace is equal to  $\mathrm{DR}_R^\bullet A_0$ , so we have a direct sum decomposition  $\mathrm{DR}_R^\bullet A=(\mathrm{DR}_R^\bullet A_0)\bigoplus (\mathrm{DR}_R^\bullet A)_{>0}$ , where the second summand is spanned by the eigenspaces corresponding to strictly positive eigenvalues.

The direct sum decomposition above is stable under the maps d and  $i_{\mathsf{E}}$ . Furthermore, the Cartan identity shows that an appropriate rescaling of contraction map  $i_{\mathsf{E}}$  provides a map  $h: (\mathrm{DR}_R^{\bullet}A)_{>0} \to (\mathrm{DR}_R^{\bullet-1}A)_{>0}$  such that we have  $\mathrm{Id} = d \circ h + h \circ d$ . Thus, the differential  $d: (\mathrm{DR}_R^{\bullet}A)_{>0} \to (\mathrm{DR}_R^{\bullet}A)_{>0}$  is homotopic to zero. It follows that the direct summand  $(\mathrm{DR}_R^{\bullet}A)_{>0}$  has trivial de Rham cohomology.

**2.8 Reduced contraction and Lie derivative.** Observe that the sign of the permutation  $(1, \ldots, k, k+1, \ldots, k+l) \mapsto (k+1, \ldots, k+l, 1, \ldots, k)$  equals  $(-1)^{kl}$ . For any  $\alpha \otimes \beta \in \Omega_R^k A \otimes \Omega_R^l A$ , we put

$$(\alpha \otimes \beta)^{\diamond} := (-1)^{kl} \beta \, \alpha \in \Omega_R^{k+l} A, \tag{2.8.1}$$

and extend the assignment  $\alpha \otimes \beta \longmapsto (\alpha \otimes \beta)^{\diamond}$  to a linear map  $\Omega_R^{\bullet} A \otimes \Omega_R^{\bullet} A \to \Omega_R^{\bullet} A$ . It is clear that, in  $\mathrm{DR}_R^{k+l} A$ , one has

$$\llbracket (\alpha \otimes \beta)^{\diamond} \rrbracket = \llbracket \alpha \beta \rrbracket, \quad \forall \alpha \in \Omega_R^k A, \ \beta \in \Omega_R^l A. \tag{2.8.2}$$

Let  $\Theta \in \mathbb{D}\mathbf{er}_R A$ . For any n = 1, 2, ..., we define reduced contraction  $\iota_{\Theta}$ , resp., reduced Lie derivative  $\mathcal{L}_{\Theta}$ , as the following maps, cf. (2.6.2), resp., (2.7.1):

$$i_{\Theta}: \Omega_{R}^{n}A \to \Omega_{R}^{n-1}A, \quad \alpha \longmapsto i_{\Theta}\alpha = (i_{\Theta}\alpha)^{\diamond}, \quad \text{resp.},$$

$$\mathcal{L}_{\Theta}: \Omega_{R}^{n}A \to \Omega_{R}^{n}A, \quad \alpha \longmapsto \mathcal{L}_{\Theta}\alpha = (L_{\Theta}\alpha)^{\diamond}. \tag{2.8.3}$$

Explicitly, for any  $\alpha_1, \alpha_2, \dots, \alpha_n \in \Omega^1_R A$ , using the definition of  $i_{\Theta}$ , we find

$$i_{\Theta}(\alpha_1 \alpha_2 \dots \alpha_n) = \sum_{k=1}^n (-1)^{(k-1)(n-k+1)} \cdot (i_{\Theta}'' \alpha_k) \cdot \alpha_{k+1} \dots \alpha_n \alpha_1 \dots \alpha_{k-1} \cdot (i_{\Theta}' \alpha_k).$$

$$(2.8.4)$$

Similarly, using the definition of  $L_{\Theta}$ , we find

$$\mathcal{L}_{\Theta}(a_0 \, da_1 \, da_2 \dots \, da_n) = \Theta''(a_0) \cdot da_1 \, da_2 \dots \, da_n \cdot \Theta'(a_0)$$

$$+ \sum_{k=1}^{n} (-1)^{k(n-k)} \cdot \Theta''(a_k) \cdot da_{k+1} \dots \, da_n \, a_0 \, da_1 \, da_2 \dots da_{k-1} \cdot d\Theta'(a_k) \quad (2.8.5)$$

$$+ \sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)} \cdot d\Theta''(a_k) \cdot da_{k+1} \dots \, da_n \, a_0 \, da_1 \, da_2 \dots da_{k-1} \cdot \Theta'(a_k).$$

The main properties of reduced contraction i may be summarized as follows.

- **Lemma 2.8.6.** (i) Fix  $\Theta \in \mathbb{D}\mathbf{er}_R A$ . Then, for any  $\omega \in \Omega_R^n A$ , the element  $\iota_{\Theta}\omega \in \Omega_R^{n-1} A$  depends only on the image of  $\omega$  in  $\mathrm{DR}_R^n A$ ; in other words, the assignment  $\omega \mapsto \iota_{\Theta}\omega$  descends to a well-defined map  $\iota_{\Theta} : \mathrm{DR}_R^n A \to \Omega_R^{n-1} A$ .
- (ii) For fixed  $\omega \in DR_R^n A$ , the assignment  $\Theta \mapsto \iota_\Theta \omega$  gives an A-bimodule morphism  $\iota(\omega) : \mathbb{D}\mathbf{er}_R A \to \Omega_R^{n-1} A$ , where  $\mathbb{D}\mathbf{er}_R A$  is equipped with the A-bimodule structure induced from the inner bimodule structure on  $A \otimes A$ .
  - (iii) For any  $\omega \in \Omega_R^n A$ , the following diagram commutes:

$$(\mathbb{D}\mathbf{er}_{R} A)_{\natural} \xrightarrow{\mathbf{m}_{*}} \operatorname{Der}_{R} A$$

$$\downarrow i: \theta \mapsto i_{\theta} \llbracket \omega \rrbracket$$

$$(\Omega_{R}^{n-1} A)_{\natural} \xrightarrow{\operatorname{proj}} \operatorname{DR}_{R}^{n-1} A.$$

*Proof.* Given a 1-form  $\beta \in \Omega^1_R A$  and  $a \in A$ , using definitions, we compute

$$i_{\Theta}(a\beta) = i''_{\Theta}(a\beta) \cdot i'_{\Theta}(a\beta) = (i''_{\Theta}\beta) \cdot a \cdot (i'_{\Theta}\beta) = i''_{\Theta}(\beta a) \cdot i'_{\Theta}(\beta a) = i_{\Theta}(\beta a).$$

It is now immediate that, for any  $\omega \in \Omega_R^n A$ , we have:

$$i_{\Theta}(a\,\omega) = i_{\Theta}(\omega\,a), \quad \text{and} \quad i_{a'\cdot\Theta\cdot a''}\omega = a'\cdot(i_{\Theta}\omega)\cdot a'', \quad \forall\, a, a', a'' \in A.$$

Further, it is clear from (2.8.4) that  $\iota_{\Theta}(\alpha_1 \dots \alpha_n)$  and  $(-1)^{n-1}\iota_{\Theta}(\alpha_n \alpha_1 \dots \alpha_{n-1})$  are given by the same formula. This yields parts (i) and (ii). Part (iii) follows from (2.8.2).

We have  $(\Omega_R^1 A)_{\natural} = DR_R^1 A$ . Hence, for  $\omega \in DR_R^2 A$ , the diagram of Lemma 2.8.6(iii), in the special case n = 2, reads

$$(\mathbb{D}\mathbf{er}_{R} A)_{\natural} \xrightarrow{\mathbf{m}_{\natural}} \mathbb{D}\mathbf{er}_{R} A \qquad (2.8.7)$$

$$DR_{R}^{1} A$$

As a general rule, any formula involving contraction  $i_{\Theta}$ , resp., Lie derivative  $L_{\Theta}$ , and some other natural map  $\Omega_R^{\bullet}A \to \Omega_R^{\bullet}A$  (as opposed to a map  $\Omega_R^{\bullet}A \to \Omega_R^{\bullet}A \otimes \Omega_R^{\bullet}A$ ) gives rise to an analogous formula involving reduced contraction  $i_{\Theta}$ , resp., reduced Lie derivative  $\mathcal{L}_{\Theta}$ . In particular, one has

**Lemma 2.8.8.** (i) For any  $\Theta \in \mathbb{D}\mathbf{er}_R A$ , we have

$$d \circ i_{\Theta} + i_{\Theta} \circ d = \mathcal{L}_{\Theta}, \quad d \circ \mathcal{L}_{\Theta} = \mathcal{L}_{\Theta} \circ d.$$

(ii) For any  $\xi \in \operatorname{Der}_R A$ , the maps  $i_{\Theta}$  and  $i_{\xi}$  anti-commute, i.e. for any  $\omega \in \operatorname{DR}_R^{\bullet} A$ , in  $\Omega_R^{\bullet - 2} A$ , we have  $(i_{\xi} \circ i_{\Theta} + i_{\Theta} \circ i_{\xi})(\omega) = 0$ .

Remark 2.8.9. Given  $\Theta, \Phi \in \mathbb{D}\mathbf{er}_R A$ , it is not true that  $\iota_{\Phi} \circ \iota_{\Theta} + \iota_{\Theta} \circ \iota_{\Phi} = 0$ , in general.  $\Diamond$ 

*Proof of Lemma*. Both statements will be verified by direct, somewhat tedious, computations. The reader is referred to [Gi2] for a more conceptual argument.

To prove (i), fix  $\Theta \in \mathbb{D}\mathbf{er}_R A$  and  $a_0, a_1, \dots, a_n \in A$ . Using formula (2.8.4), for  $d \circ \iota_{\Theta}(a_0 da_1 \dots da_n)$  we get the following expression:

$$d\left(\sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)} \cdot (i_{\Theta}^{"}\alpha_{k}) \cdot \alpha_{k+1} \dots \alpha_{n} \alpha_{0} \dots \alpha_{k-1} \cdot (i_{\Theta}^{'}\alpha_{k})\right)$$

$$= \sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)+n-k} \Theta^{"}(a_{k}) \cdot da_{k+1} \dots da_{n} da_{0} da_{1} \dots da_{k-1} \cdot \Theta^{'}(a_{k})$$

$$+ \sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)} d\Theta^{"}(a_{k}) \cdot da_{k+1} \dots da_{n} a_{0} da_{1} \dots da_{k-1} \cdot \Theta^{'}(a_{k})$$

$$+ \sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)+n-1} \cdot \Theta^{"}(a_{k}) \cdot da_{k+1} \dots da_{n} a_{0} da_{1} \dots da_{k-1} \cdot d\Theta^{'}(a_{k}).$$

After simplification of powers of (-1), we find that  $d \circ \iota_{\Theta}(a_0 da_1 \dots da_n)$  is a sum of three terms,  $S_0 + S_+ + S_-$ , where

$$S_{0} = \sum_{k=1}^{n} (-1)^{nk-1} \Theta''(a_{k}) \cdot da_{k+1} \dots da_{n} da_{0} da_{1} \dots da_{k-1} \cdot \Theta'(a_{k});$$

$$S_{+} = \sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)} d\Theta''(a_{k}) \cdot da_{k+1} \dots da_{n} a_{0} da_{1} \dots da_{k-1} \cdot \Theta'(a_{k});$$

$$S_{-} = \sum_{k=1}^{n} (-1)^{k(n-k)} \cdot \Theta''(a_{k}) \cdot da_{k+1} \dots da_{n} a_{0} da_{1} \dots da_{k-1} \cdot d\Theta'(a_{k}).$$

On the other hand, for  $\iota_{\Theta} \circ d(a_0 da_1 \dots da_n) = \iota_{\Theta}(da_0 da_1 \dots da_n)$ , we find

$$i_{\Theta} \circ d(a_0 \, da_1 \dots da_n) = \Theta''(a_0) \cdot da_1 \dots da_n \cdot \Theta'(a_0)$$

$$+ \sum_{k=1}^n (-1)^{k(n-k+1)} \cdot \Theta''(a_k) \cdot da_{k+1} \dots da_n \, da_0 \, da_1 \dots da_{k-1} \cdot \Theta'(a_k)$$

$$= \Theta''(a_0) \cdot da_1 \dots da_n \cdot \Theta'(a_0)$$

$$+ \sum_{k=1}^n (-1)^{nk} \cdot \Theta''(a_k) \cdot da_{k+1} \dots da_n \, da_0 \, da_1 \dots da_{k-1} \cdot \Theta'(a_k)$$

$$= \Theta''(a_0) \cdot da_1 \dots da_n \cdot \Theta'(a_0) - S_0.$$

Thus, we see that

$$(d \circ i_{\Theta} + i_{\Theta} \circ d)(a_0 da_1 \dots da_n) = \Theta''(a_0) \cdot da_1 \dots da_n \cdot \Theta'(a_0) + S_+ + S_-.$$

The expression on the right hand side is exactly the one given by formula (2.8.5), and part (i) is proved.

To prove (ii), fix  $\Theta \in \mathbb{D}\mathbf{er}_R A$  and  $\xi \in \mathrm{Der}_R A$ , and also 1-forms  $\alpha_1, \ldots, \alpha_n \in \Omega^1_R A$ . For any pair of integers  $1 \leq j, k \leq n$  such that  $j \neq k$ , let  $S_{j,k}$  be the (n-2)-form obtained from the form

$$(i''_{\Theta}\alpha_k)\cdot\alpha_{k+1}\ldots\alpha_n\alpha_1\ldots\alpha_{k-1}\cdot(i'_{\Theta}\alpha_k)\in\Omega_R^{n-1}A$$

by replacing the factor  $\alpha_j \in \Omega^1_R A$  by the element  $i_{\xi} \alpha_j \in A$ .

With this notation, we express  $i_{\xi} \circ i_{\Theta}(\alpha_1 \alpha_2 \dots \alpha_n)$ , using formula (2.8.4) as follows:

$$i_{\xi} \left( \sum_{1 \leq k \leq n} (-1)^{(k-1)(n-k+1)} (i_{\Theta}'' \alpha_k) \cdot \alpha_{k+1} \dots \alpha_n \alpha_1 \dots \alpha_{k-1} \cdot (i_{\Theta}' \alpha_k) \right)$$

$$= \sum_{k=1}^{n} (-1)^{(k-1)(n-k+1)} \left( \sum_{j=k+1}^{n} (-1)^{j-k+1} S_{j,k} + \sum_{j=1}^{k-1} (-1)^{n-k+j-1} S_{j,k} \right).$$

Simplifying, we obtain

$$i_{\xi} \circ i_{\Theta}(\alpha_1 \alpha_2 \dots \alpha_n) = \sum_{1 \le k < j \le n} (-1)^{nk-n+j} S_{j,k} + \sum_{1 \le j < k \le n} (-1)^{nk+j} S_{j,k}. \quad (2.8.10)$$

On the other hand, we have

$$i_{\Theta} \circ i_{\xi}(\alpha_{1}\alpha_{2} \dots \alpha_{n}) = i_{\Theta} \left( \sum_{j=1}^{n} (-1)^{j-1} \alpha_{1} \dots \alpha_{j-1} \cdot (i_{\xi}\alpha_{j}) \cdot \alpha_{j+1} \dots \alpha_{n} \right)$$

$$= \sum_{1 \leq k < j \leq n} (-1)^{j-1+(k-1)(n-k)} S_{j,k} + \sum_{1 \leq j < k \leq n} (-1)^{j-1+(k-2)((n-1)-(k-1)+1)} S_{j,k}$$

$$= \sum_{1 \leq k < j \leq n} (-1)^{nk-n+j-1} S_{j,k} + \sum_{1 \leq j < k \leq n} (-1)^{nk+j-1} S_{j,k}.$$
(2.8.11)

Comparison of formulas (2.8.10) and (2.8.11) completes the proof of part (ii).  $\square$ 

Observe further that, by Lemma 2.8.6, the maps  $i_{\Theta}$  and  $\mathcal{L}_{\Theta}$  descend to the following well-defined maps (the diagram below does *not* commute):

$$DR_{R}^{\bullet}A \xrightarrow{d} DR_{R}^{\bullet+1}A \qquad (2.8.12)$$

$$\downarrow \iota_{\Theta} \qquad \qquad \downarrow \iota_{\Theta}$$

Next, fix  $\Theta \in \mathbb{D}\mathbf{er}_R A$ , let  $\theta = \mathbf{m}_*(\Theta) \in \mathrm{Der}_R A$  be the corresponding derivation  $A \to A$ , and  $L_{\theta} : \mathrm{DR}_R^{\bullet} A \to \mathrm{DR}_R^{\bullet} A$  the Lie derivative operator induced by  $\theta$ . It is immediate from part (iii) of Lemma 2.8.6 that one has a commutative triangle

$$DR_{R}^{\bullet}A \xrightarrow{\mathscr{L}_{\Theta}} \Omega_{R}^{\bullet}A \qquad (2.8.13)$$

$$DR_{R}^{\bullet}A$$

In other words, the diagram says that, if  $\Theta \in \mathbb{D}\mathbf{er}_R A$  and  $\theta = \mathbf{m}_*(\Theta) \in \mathrm{Der}_R A$ , then  $\mathscr{L}_{\Theta} = L_{\theta}$  as maps  $\mathrm{DR}_R^{\bullet} A \to \mathrm{DR}_R^{\bullet} A$ .

Remark 2.8.14. The reader should be warned that, in the above setting, it is not true in general that  $\mathcal{L}_{\Theta} = L_{\theta}$  as maps  $\Omega_R^n A \to \Omega_R^n A$ . In particular, for n = 0 we have two different maps  $\mathcal{L}_{\Theta}$ ,  $L_{\theta} : A \to A$ , where  $L_{\theta}(a) = \Theta'(a)\Theta''(a)$  and  $\mathcal{L}_{\Theta}(a) = \Theta''(a)\Theta'(a)$ .

## 3 The derivation $\Delta$

**3.1** From now on, we assume that R is a finite dimensional semisimple algebra, and  $\mathbf{e} = \mathbf{e}' \otimes \mathbf{e}'' \in R \otimes R$  is a fixed separability element, see Definition (2.2.1).

Refining slightly an idea of Van den Bergh, see [VB2, §3.3], we introduce the following distinguished derivation

$$\Delta: A \to A \otimes A, \ a \mapsto \Delta(a) = \sum_{i} (ae_i \otimes e^i - e_i \otimes e^i a) = a\mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}'' a.$$

We observe that  $\Delta(R) = 0$  since  $\mathbf{e} \in (R \otimes R)^R$ . Thus,  $\Delta \in \mathbb{D}\mathbf{er}_R A$ .

**Lemma 3.1.1.** (i) For any  $\omega \in \Omega_R^{\bullet} A$ , we have

$$L_{\Lambda}\omega = \omega \mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}''\omega$$
, and  $\mathscr{L}_{\Lambda}\omega = 0$ .

(ii) Writing  $[a, \omega] := a \omega - \omega a$ , for any  $a_0, a_1, \ldots, a_n \in A$ , we have

$$i_{\Delta}(a_0 da_1 \dots da_n) = \sum_{k=1}^{n} (-1)^{(k-1)(n+1)} \mathbf{e}'[a_k, da_{k+1} \dots da_n a_0 da_1 \dots da_{k-1}] \mathbf{e}''.$$

In particular, we have  $\operatorname{Im}(i_{\Delta}) \subset [A, \Omega_{R}^{\bullet}A]^{R}$ .

*Proof.* We prove (i) in the special case  $\omega = x \, dy \, dz \in \Omega^2_R A$ . We compute

$$L_{\Delta}\omega = L_{\Delta}(x \, dy \, dz) = (x\mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}''x) \, dy \, dz$$

$$+ x \, d(y\mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}''y) \, dz + x \, dy \, d(z\mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}''z)$$

$$= x\mathbf{e}' \otimes \mathbf{e}'' \, dy \, dz - \mathbf{e}' \otimes \mathbf{e}''x \, dy \, dz$$

$$+ x \, dy \, \mathbf{e}' \otimes \mathbf{e}'' \, dz - x\mathbf{e}' \otimes \mathbf{e}'' \, dy \, dz + x \, dy \, dz \, \mathbf{e}' \otimes \mathbf{e}'' - x \, dy \, \mathbf{e}' \otimes \mathbf{e}'' \, dz$$

$$= x \, dy \, dz \, \mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}'' \, x \, dy \, dz = \omega \, \mathbf{e}' \otimes \mathbf{e}'' - \mathbf{e}' \otimes \mathbf{e}'' \, \omega.$$

The computation of  $L_{\Delta}\omega$  in the general case is very similar.

Further, it is clear from formula (2.8.1) that for any  $\omega \in \Omega_R^n A$ , we have  $(\sum_i \omega e_i \otimes e^i)^{\diamond} = \sum_i e^i \omega e_i = (e_i \otimes e^i \omega)^{\diamond}$ . We deduce that  $\mathcal{L}_{\Delta}\omega = 0$ . This proves part (i).

A similar direct computation based on (2.8.4) yields an analogue of the formula of part (ii), with the roles of  $\mathbf{e}'$  and  $\mathbf{e}''$  flipped. By the symmetry of the separability element, we have  $\mathbf{e}'' \otimes \mathbf{e}' = \mathbf{e}' \otimes \mathbf{e}''$ , and the formula of part (ii) follows. The last claim of the Lemma follows from that formula and (2.2.3).  $\square$ 

Observe that the projection  $\mathbf{m}_*: \mathbb{D}\mathbf{er}_R A \to \operatorname{Der}_R A$  clearly maps the derivation  $\Delta$  to zero. It follows that the composite  $\operatorname{DR}_R^{\bullet} A \xrightarrow{\imath_{\Delta}} \Omega_R^{\bullet-1} A \xrightarrow{\operatorname{proj}} \operatorname{DR}_R^{\bullet-1} A$ ,  $\alpha \mapsto \llbracket \imath_{\Delta} \alpha \rrbracket$ , is the zero map.

Corollary 3.1.2. (i) For any  $\omega \in \Omega_R^{\bullet} A$ , in  $\mathrm{DR}_R^{\bullet-1} A$ , we have  $\llbracket \imath_{\Delta} \omega \rrbracket = 0$ . Also, in  $\Omega_R^{\bullet-2} A$ , resp. in  $\mathrm{DR}_R^{\bullet-1} A$ , we have

$$i_{\Theta}(i_{\Delta}\omega) = 0$$
, resp.,  $\mathscr{L}_{\Theta}(i_{\Delta}\omega) = 0$ ,  $\forall \Theta \in \mathbb{D}\mathbf{er}_{R} A$ .

(ii) Furthermore, we have:

$$i_{\Delta} \circ d + d \circ i_{\Delta} = 0$$
, and  $L_{\theta} \circ i_{\Delta} = i_{\Delta} \circ L_{\theta}$ ,  $\forall \theta \in \operatorname{Der}_{R} A$ .

*Proof.* By Lemma 2.8.6(i), the element  $\iota_{\Theta}(\iota_{\Delta}\omega) \in \Omega_R^{\bullet-2}A$  depends only on  $\llbracket \iota_{\Delta}\omega \rrbracket$ , the image of  $\iota_{\Delta}\omega$  in  $\mathrm{DR}_R^{\bullet-1}A$ . Therefore, since  $\llbracket \iota_{\Delta}\omega \rrbracket = 0$ , we conclude that  $\iota_{\Theta}(\iota_{\Delta}\omega) = 0$ . The proof for  $\mathscr{L}_{\Theta}(\iota_{\Delta}\omega) = 0$  is entirely similar.

The first formula of part (ii) follows from Lemma 3.1.1(i) combined with the Cartan formula of Lemma 2.8.8(i). The second formula follows from the first one, using the standard Cartan formula  $L_{\theta} = d \circ i_{\theta} + i_{\theta} \circ d$  and Lemma 2.8.8.  $\square$ 

**3.2** Inner derivations. Given an A-bimodule M and  $m \in M$ , write ad m:  $A \to M$  for the *inner derivation* ad m(a) := am - ma. If  $m \in M^R$  then ad m(R) = 0, hence ad  $m \in \text{Der}_R(A, M)$ .

In the special case  $M = A \otimes A$ , the map  $M \to \operatorname{Der}(A, M)$ ,  $m \longmapsto \operatorname{ad} m$  is, in effect, a morphism of A-bimodules, provided  $\operatorname{Der}(A, M) = \operatorname{Der}(A, A \otimes A)$  is equipped with the bimodule structure induced by the inner bimodule structure on  $A \otimes A$ .

Let  $\mathbf{e} \in R \otimes R \subset A \otimes A$  be the separability element, see §2.2. By definition, we have  $\Delta = \operatorname{ad} \mathbf{e}$ .

Further, using Proposition 2.2.2 we see that, for any  $p = p' \otimes p'' \in (A \otimes A)^R$ , one has that  $p = p' \otimes p'' = \sum_{i \in I} e_i p' \otimes p'' e^i = \mathbf{e}' p' \otimes p'' \mathbf{e}''$ . The last expression is nothing but the result of the *inner* action of the element  $p^{\text{op}} = p'' \otimes p'$  on  $\mathbf{e} = \mathbf{e}' \otimes \mathbf{e}''$ . It follows that the inner derivation ad p corresponding to  $p = p' \otimes p''$  can be written, using the A-bimodule structure on  $\mathbb{D}\mathbf{er}_R A$ , as  $p'' \cdot \Delta \cdot p'$ .

This way, one obtains the following result.

**Lemma 3.2.1.** (i) For any  $p = p' \otimes p'' \in (A \otimes A)^R$ , we have  $\operatorname{ad} p = p'' \cdot \Delta \cdot p'$ ; hence,  $\mathbf{m}_*(p'' \cdot \Delta \cdot p') = \operatorname{ad} a$ , where  $a = p'p'' \in A^R$ .

(ii) Inner derivations form an  $A^e$ -submodule in  $\mathbb{D}\mathbf{er}_R A$ , and this  $A^e$ -submodule is generated by the derivation  $\Delta$ .

Corollary 3.2.2. For any  $\omega \in DR_R^n A$  and  $p = p' \otimes p'' \in (A \otimes A)^R$ , in  $\Omega_R^{\bullet} A$  we have

$$i_{\operatorname{ad} p}\omega = p''\left(i_{\Delta}\omega\right)p', \quad and \quad \mathscr{L}_{\operatorname{ad} p}\omega = dp''\left(i_{\Delta}\omega\right)p' - (-1)^{\operatorname{deg}\omega}p''\left(i_{\Delta}\omega\right)dp'.$$

In particular,  $\mathcal{L}_{\operatorname{ad} p}(a) = 0$ , for any  $a \in A = \Omega^0_R A$ .

*Proof.* The map  $\iota(\omega): \mathbb{D}\mathbf{er}_R A \longrightarrow \Omega_R^{n-1} A$  being a map of A-bimodules, for any  $p = p' \otimes p'' \in (A \otimes A)^R$ , we deduce

$$i_{\text{ad }p}\omega = i_{p''\cdot\Delta\cdot p'}\omega = p''(i_{\Delta}\omega)p'.$$

This proves the first formula.

Next, using the Cartan formula, we compute

$$\mathcal{L}_{\operatorname{ad} p}\omega = \iota_{\operatorname{ad} p} d\omega + d \iota_{\operatorname{ad} p}\omega = p'' (\iota_{\Delta} d\omega) p' + d(p'' (\iota_{\Delta} \omega) p')$$

$$= p'' (\iota_{\Delta} d\omega) p' + dp'' (\iota_{\Delta} \omega) p' + p'' (d\iota_{\Delta} \omega) p' - (-1)^{\operatorname{deg} \omega} p'' (\iota_{\Delta} \omega) dp'$$

$$= p'' (\mathcal{L}_{\Delta} \omega) p' + dp'' (\iota_{\Delta} \omega) p' - (-1)^{\operatorname{deg} \omega} p'' (\iota_{\Delta} \omega) dp'.$$

But  $\mathcal{L}_{\Delta}\omega = 0$  by Lemma 3.1.1(i), and the result follows.

П

Let  $\operatorname{Inn}_R A \subset \operatorname{Der}_R A$  be the subspace of inner derivations. This is a *Lie ideal* with respect to the commutator of derivations. From Corollary 3.2.2, using the commutative diagram of Lemma 2.8.6(iii) we deduce

Corollary 3.2.3. In  $DR_R^{\bullet}A$ , we have

$$i_{\operatorname{ad} a}\omega = a \cdot \imath_{\Delta}\omega$$
, and  $L_{\operatorname{ad} a}\omega = da \cdot \imath_{\Delta}\omega$ ,  $\forall \omega \in \operatorname{DR}^n_R A, \ a \in A^R$ .

**3.3** Geometric example. Let  $R = \mathbb{k}$  be an algebraically closed field (of characteristic zero as usual), and let  $A = \mathbb{k}[X]$  be the coordinate ring of a smooth affine algebraic variety. Then,  $A \otimes A = \mathbb{k}[X \times X]$ . In this case, it is well-known that the Ext-group  $\operatorname{Ext}_{A^e}^k(A, A \otimes A)$  vanishes for all  $k \neq \dim X$ . Thus, if  $\dim X > 1$  then, we have  $\operatorname{Der}(A, A \otimes A) / \operatorname{Inn}(A, A \otimes A) = \operatorname{Ext}_{A^e}^1(A, A \otimes A) = 0$ . We deduce that any derivation  $\Theta \in \mathbb{D}$ er A is inner.

Assume now that X is a smooth curve (dim X=1). Then, the bimodule  $\Omega^1 A \subset A \otimes A$  is the ideal of the diagonal divisor  $D \subset X \times X$ . Hence we have

$$\mathbb{D}\mathbf{er}\,A = \mathrm{Hom}_{A^e}(\Omega^1 A, A \otimes A) = \Gamma(X \times X, \, \mathcal{O}_{X \times X}(D)) \tag{3.3.1}$$

is the space of regular functions on  $(X \times X) \setminus D$  with at most simple poles along D. We also have algebra isomorphisms  $\operatorname{Hom}_{A^e}(\Omega^1 A, \Omega^1 A) = \Gamma(X \times X, \mathcal{O}_{X \times X}) = A \otimes A$ .

It is known that, for a smooth curve X, the algebra  $A = \mathbb{k}[X]$  is smooth in the sense of Definition 2.1.1, cf. [CQ]. This way, the short exact sequence of Proposition 2.3.2 becomes the top row of the following diagram

$$0 \longrightarrow A \otimes A \xrightarrow{j} \mathbb{D}\mathbf{er} A \longrightarrow \mathbb{D}\mathbf{er} A \longrightarrow 0$$

$$\parallel \qquad \qquad \parallel_{\Psi} \qquad \qquad \parallel_{\Phi}$$

$$0 \longrightarrow \Gamma(\mathcal{O}_{X \times X}) \longrightarrow \Gamma(\mathcal{O}_{X \times X}(D)) \longrightarrow \Gamma(\mathcal{O}_{X \times X}(D)/\mathcal{O}_{X \times X}) \longrightarrow 0$$

In the bottom row of the diagram above we have used shorthand notation  $\Gamma(-)$  for  $\Gamma(X \times X, -)$ ; this row is obtained by applying the global sections functor to the natural extension of sheaves on  $X \times X$ . The vertical isomorphism  $\Phi$ , in the diagram, follows from the identification  $\operatorname{Der} A = \mathcal{F}(X)$ , with the space of regular vector fields on X. The vertical isomorphism  $\Psi$  comes from (3.3.1).

Observe that the function  $1 \in A \otimes A$  corresponds under the above identifications to the element  $\mathrm{Id}_{\Omega} \in \mathrm{Hom}_{A^c}(\Omega^1 A, \Omega^1 A)$ . Therefore, in the diagram we have  $j(1) = \Delta$ , and the map j is nothing but the imbedding ad  $: A \otimes A \hookrightarrow \mathrm{Der}(A, A \otimes A)$ , of inner derivations.

# 4 Hamilton operators.

**4.1** Noncommutative moment map. We fix a semisimple finite-dimensional  $\mathbb{R}$ -algebra R and an R-algebra A. Let  $(A/R)^R$  be the centralizer of R in the R-bimodule A/R.

We are going to define a canonical linear map

$$\mu_{\rm nc}: ({\rm DR}_R^2 A)_{\rm closed} \longrightarrow (A/R)^R$$
, such that  $d\mu_{\rm nc}(\omega) = i\Delta\omega$  (4.1.1)

holds in  $\Omega_R^1 A$ , for any  $\omega \in (DR_R^2 A)_{closed}$ .

To this end, we observe that the de Rham differential d anti-commutes with  $i_{\Delta}$ , by Corollary 3.1.2(ii). Therefore, for any  $\omega \in \mathrm{DR}^2_R A$  such that  $d\omega = 0$ , we have  $d(i_{\Delta}\omega) = -i_{\Delta}(d\omega) = 0$ . Moreover, by the last statement of Lemma 3.1.1, we know that  $i_{\Delta}\omega \in (\Omega^1_R A)^R_{\mathrm{closed}}$ .

Recall further that the complex  $(\Omega_R^{\bullet}A, d)$  is acyclic in positive degrees, see (2.5.1). It follows that the de Rham differential yields an isomorphism  $d: A/R \xrightarrow{\sim} (\Omega_R^1 A)_{\text{closed}}$ . This isomorphism clearly commutes with the adjoint action of R, hence, induces an isomorphism  $(A/R)^R \xrightarrow{\sim} (\Omega_R^1 A)_{\text{closed}}^R$ . Thus, we may (and will) define the map  $\mu_{\text{nc}}$  as a composite:

$$\mu_{\rm nc}: (\mathrm{DR}_R^2 A)_{\rm closed} \xrightarrow{\imath_\Delta} (\Omega_R^1 A)_{\rm closed}^R = (\Omega_R^1 A)_{\rm exact}^R \xrightarrow{d^{-1}} (A/R)^R.$$
 (4.1.2)

The map  $\mu_{\rm nc}$  will play a crucial role in our approach to noncommutative Hamiltonian reduction. We will see in §6 that the object corresponding to  $\mu_{\rm nc}$  under the representation functor is closely related to the ordinary moment map used in (commutative) symplectic geometry.

Here are a few basic formulas involving the moment map  $\mu_{\rm nc}$ .

**Proposition 4.1.3.** Let  $\omega$  be a closed 2-form and  $\mathbf{w} \in A$  a representative of the class  $\mu_{nc}(\omega) \in A/R$ . Then,

- (i) We have  $\mathbf{w} \in R + A^R$ , and also  $d\mathbf{w} = 0$  in  $DR_R^1 A$ .
- (ii) For any  $\theta \in \operatorname{Der}_R A$  and  $\Theta \in \operatorname{\mathbb{D}er}_R A$ , in A/R, one has

$$\mu_{\rm nc}(L_{\theta}\omega) = L_{\theta}(\mu_{\rm nc}(\omega)), \text{ and } \mathscr{L}_{\Theta}(\mu_{\rm nc}(\omega)) = 0.$$

(iii) For any 
$$p = p' \otimes p'' \in (A \otimes A)^R$$
,  $a \in A^R$ , and  $u, v \in A$ , one has 
$$\begin{aligned} i_{\operatorname{ad} p} \omega &= p'' \cdot d\mathbf{w} \cdot p' & \operatorname{holds in } \Omega^1_R A, \\ i_{\operatorname{ad} a} \omega &= a \cdot d\mathbf{w} & \operatorname{holds in } \operatorname{DR}^1_R A; \\ \mu_{\operatorname{nc}}(du \, dv) &= \mathbf{e}' \cdot [u, v] \cdot \mathbf{e}'' \, \operatorname{mod} R. \end{aligned}$$

*Proof.* By Corollary 3.2.3, we know that  $i_{\Delta}\omega = 0$  in  $\mathrm{DR}^1_R A$ , hence  $d\mathbf{w} = i_{\Delta}\omega = 0$ . Observe next that, since  $\mu_{\mathrm{nc}}(\omega) \in (A/R)^R$ , we have  $[\mathbf{w}, R] \subset R$ . Hence the derivation ad  $\mathbf{w}$  preserves R. But any derivation of the finite dimensional semisimple algebra R is inner. Hence, there exists  $r \in R$  such that  $\mathrm{ad} \mathbf{w}|_R = \mathrm{ad} r|_R$ . Thus,  $\mathbf{w} - r$  commutes with R, and (i) follows.

Part (ii) and the first two equations of part (iii) are immediate from the properties of  $i_{\Delta}$  established earlier.

To prove the last formula, using Lemma 3.1.1(ii) we compute

$$i_{\Delta}(du\,dv) = \mathbf{e}'(-[dv,u] + [du,v])\mathbf{e}'' = \mathbf{e}'([du,v] + [u,dv])\mathbf{e}'' = \mathbf{e}'(d[u,v])\mathbf{e}''.$$

The last expression equals  $d(\mathbf{e}' \cdot [u, v] \cdot \mathbf{e}'')$ . Thus, in  $\Omega_R^1 A$ , we obtain  $d\mu_{\rm nc}(du \, dv) = i_{\Delta}(du \, dv) = d(\mathbf{e}' \cdot [u, v] \cdot \mathbf{e}'')$ , hence  $\mu_{\rm nc}(du \, dv) - \mathbf{e}' \cdot [u, v] \cdot \mathbf{e}'' \in R$ , and we are done.

Recall the notation  $V^R:=V\cap M^R$  for any vector subspace V in an R-bimodule M.

**Proposition 4.1.4.** Let A be an R-algebra such that the following sequence is exact

$$0 \longrightarrow R \longrightarrow DR_R^0 A \stackrel{d}{\longrightarrow} DR_R^1 A. \tag{4.1.5}$$

Then, the map  $\mu_{\rm nc}: ({\rm DR}^2_R A)_{\rm closed} \to (A/R)^R$  can be lifted canonically to a map  $\widetilde{\mu}_{\rm nc}$  in the diagram below

$$DR_{R}^{1}A \xrightarrow{d} (DR_{R}^{2}A)_{\text{closed}}$$

$$\downarrow^{i_{\Delta}} \qquad \downarrow^{i_{\Lambda}} \qquad \downarrow^{i_{\Lambda}} \qquad \downarrow^{i_{\Lambda}} \qquad (4.1.6)$$

$$[A, A]^{R} \xrightarrow{d} [A, \Omega_{R}^{1}A]^{R}.$$

The map  $\widetilde{\mu_{nc}}$  has the following properties:

- The above diagram commutes and we have  $\mu_{nc}(\omega) = \widetilde{\mu_{nc}}(\omega) \mod R$ ;
- The map  $\widetilde{\mu_{nc}}$  commutes with the Lie derivative  $L_{\theta}$ , for any  $\theta \in \operatorname{Der}_R A$ .

The exactness of (4.1.5) is equivalent to the following two equalities:

$$\operatorname{Ker}[d: \operatorname{DR}^0_R A \to \operatorname{DR}^1_R A] = R \quad \text{and} \quad [A, A] \cap R = 0.$$
 (4.1.7)

Proof of Proposition. Let  $\omega \in DR_R^2 A$  be a closed 2-form and  $\mathbf{w} \in A$  a representative of the class  $\mu_{\rm nc}(\omega) \in A/R$ . We know that  $d\mathbf{w} = 0$  in  $DR_R^1 A$ , by Proposition 4.1.3(i). Hence, the first equality in (4.1.7) yields  $\mathbf{w} \in R + [A, A]$ . Now, the second equality in (4.1.7) implies that there is a unique representative of the class  $\mu_{\rm nc}(\omega)$  that belongs to [A, A]. This provides a lift  $\widehat{\mu_{\rm nc}} : (DR_R^2 A)_{\rm closed} \to [A, A]$ .

Further, for any derivation  $\theta \in \operatorname{Der}_R A$ , we have  $\theta([A,A]) \subset [A,A]$  and  $\theta(R) = 0$ . Thus, for any choice of  $\mathbf{w} \in R + [A,A]$  we get  $\theta(\mathbf{w}) \in \theta(R + [A,A]) = \theta([A,A]) \subset [A,A]$ . Therefore, we deduce  $L_{\theta}\widetilde{\mu}_{\operatorname{nc}}(\omega) - \widetilde{\mu}_{\operatorname{nc}}(L_{\theta}\omega) \in [A,A]$ . On the other hand, since  $\mu_{\operatorname{nc}}$  commutes with  $L_{\theta}$ , we have  $L_{\theta}\widetilde{\mu}_{\operatorname{nc}}(\omega) - \widetilde{\mu}_{\operatorname{nc}}(L_{\theta}\omega) \in R$ . It follows that  $L_{\theta}\widetilde{\mu}_{\operatorname{nc}}(\omega) - \widetilde{\mu}_{\operatorname{nc}}(L_{\theta}\omega) \in R \cap [A,A] = 0$ , by the second equality in (4.1.7). We conclude that the map  $\widetilde{\mu}_{\operatorname{nc}}$  commutes with  $L_{\theta}$ .


Observe finally that the second equality in (4.1.7) implies, in particular, that R is commutative. Hence, from Proposition 4.1.3(i) we deduce  $[\mathbf{w}, R] \subset [R, R] = 0$ . Thus,  $\mathbf{w} \in A^R$ , and we get  $\widehat{\mu}_{\mathrm{nc}}(\omega) \in [A, A] \cap A^R = [A, A]^R$ . This completes the proof.

**4.2** Symplectic 2-forms. We say that a 2-form  $\omega \in DR_R^2 A$  is nondegenerate if the map  $i(\omega): Der_R A \to DR_R^1 A$ ,  $\theta \mapsto i_\theta \omega$ , is a bijection. Given a nondegenerate form  $\omega \in DR_R^2 A$ , we invert the bijection  $i(\omega)$  to obtain a bijection  $H_\omega = i(\omega)^{-1}: DR_R^1 A \xrightarrow{\sim} Der_R A$ .

We say that a 2-form  $\omega \in \mathrm{DR}^2_R A$  is bi-nondegenerate if the map  $\iota(\omega)$ :  $\mathbb{D}\mathbf{er}_R A \to \Omega^1_R A$ ,  $\Theta \mapsto \iota_\Theta \omega$  is a bijection, hence, an isomorphism of A-bimodules.

Let  $\omega \in \mathrm{DR}^2_R A$  be a bi-nondegenerate 2-form. Applying the functor  $(-)_{\natural}$  to the map  $\imath(\omega)$  yields the bijection  $\imath(\omega)_{\natural}: (\mathbb{D}\mathbf{er}_R A)_{\natural} \overset{\sim}{\to} (\Omega^1_R A)_{\natural} = \mathrm{DR}^1_R A$ .

Further, using the commutativity of diagram (2.8.7), we deduce that the map  $\mathbf{m}_{\natural} : (\mathbb{D}\mathbf{er}_R A)_{\natural} \to \mathrm{Der}_R A$  is *injective*; moreover, one has the following diagram



In this diagram, the vertical maps are the natural imbedding and projection (along  $\text{Ker}(i(\omega))$ ), respectively, and the horizontal maps  $\mathbf{m}_{\natural}$  and  $i(\omega)$  are both bijections whose composite equals  $i(\omega)_{\natural}$ . It follows in particular that, for a smooth algebra A, any bi-nondegenerate form is automatically nondegenerate.

Next, we invert the bijection  $i(\omega)_{\natural}$  and introduce the following composite

$$\mathsf{H}_{\omega}: \ \mathrm{DR}_{R}^{1}A \xrightarrow{\imath(\omega)_{\natural}^{-1}} (\mathbb{D}\mathbf{er}_{R}A)_{\natural} \xrightarrow{\mathbf{m}_{\natural}} \mathrm{Der}_{R}A.$$
 (4.2.1)

We see from the diagram above that the map  $H_{\omega}$  is injective, furthermore, it provides a section of the projection  $i(\omega)$ :  $\operatorname{Der}_R A \twoheadrightarrow \operatorname{DR}_R^1 A$ . It follows in particular that we have

$$i_{\mathsf{H}_{\omega}(\alpha)}\omega = \alpha, \quad \forall \alpha \in \mathrm{DR}^1_R A.$$
 (4.2.2)

Both in the nondegenerate and bi-nondegenerate cases, we also consider the composite

$$A_{\natural} \xrightarrow{d} \mathrm{DR}_{R}^{1} A \xrightarrow{\mathsf{H}_{\omega}} \mathrm{Der}_{R} A, \quad p \longmapsto \theta_{p} := \mathsf{H}_{\omega}(dp).$$
 (4.2.3)

**Proposition 4.2.4.** Let  $\omega$  be a closed 2-form and  $\mathbf{w} \in A$  a representative of the class  $\mu_{\mathrm{nc}}(\omega) \in A/R$ . Let  $\theta \in \mathrm{Der}_R A$  and put  $\alpha = i_{\theta}\omega$ . Then

- (i) We have  $\theta(\mathbf{w}) = -i_{\Delta}\alpha$ .
- (ii) The 1-form  $\alpha \in DR^1_RA$  is closed if and only if  $L_{\theta}\omega = 0$ .

Proof. Let  $\theta \in \operatorname{Der}_R A$ . The maps  $i_{\theta}$  and  $i_{\Delta}$  anti-commute, by Lemma 2.8.8(ii). Hence, using (4.2.2) we find  $\theta(\mathbf{w}) = i_{\theta}(i_{\Delta}\omega) = -i_{\Delta}(i_{\theta}\omega) = -i_{\Delta}\alpha$ , and (i) follows. To prove (ii), we write  $L_{\theta}\omega = di_{\theta}\omega + i_{\theta}d\omega = di_{\theta}\omega$ , since  $d\omega = 0$ . We see that  $L_{\theta}\omega = 0$  if and only if  $di_{\theta}\omega = 0$ .

Fix a closed 2-form  $\omega \in \mathrm{DR}^2_R A$ . A derivation  $\theta \in \mathrm{Der}_R A$  is said to be  $\omega$ -symplectic if in  $\mathrm{DR}^2_R A$  one has  $L_\theta \omega = 0$ . Clearly,  $\omega$ -symplectic derivations form a Lie subalgebra  $\mathrm{Der}^\omega_R A \subset \mathrm{Der}_R A$ .

**Definition 4.2.5.** A closed 2-form  $\omega \in DR_R^2 A$  is called *symplectic*, resp., *bi-symplectic*, if it is nondegenerate, resp., bi-nondegenerate.

**Lemma 4.2.6.** Let  $\omega$  be a closed 2-form and  $\mathbf{w} \in A$  a representative of the class  $\mu_{\rm nc}(\omega) \in A/R$ . Then

(i) For any  $\omega$ -symplectic derivation  $\theta \in \operatorname{Der}_R^{\omega} A$  we have  $\theta(\mathbf{w}) \in R$ .

- (ii) If  $\omega$  is either symplectic or bi-symplectic then, for any  $a \in A_{\natural}$ , we have  $\theta_a \in \operatorname{Der}_B^{\omega} A$ ; furthermore,  $\theta_a(\mathbf{w}) = 0$ .
  - (iii) If the sequence (4.1.5) is exact then  $\theta \in \operatorname{Der}_R^{\omega} A$  implies  $\theta(\mathbf{w}) = 0$ .

Proof. Let  $\theta \in \operatorname{Der}_R^{\omega} A$ . Since  $L_{\theta}$  commutes with  $\iota_{\Delta}$ , from Proposition 4.2.4(ii) we deduce  $L_{\theta}\iota_{\Delta}\omega = \iota_{\Delta}L_{\theta}\omega = 0$ . The map d, hence  $d^{-1}$ , also commutes with  $L_{\theta}$ . Thus, in A/R, we get  $L_{\theta}\mu_{\rm nc}(\omega) = L_{\theta}d^{-1}(\iota_{\Delta}\omega) = d^{-1}(L_{\theta}\iota_{\Delta}\omega) = 0$ . This means that  $L_{\theta}\mathbf{w} = 0 \mod R$ , hence  $\theta(\mathbf{w}) \in R$ , and (i) follows.

Since da is a closed 1-form for any  $a \in A$ , we deduce from Proposition 4.2.4(ii) that  $\theta_a \in \operatorname{Der}_R^{\omega} A$ . Further, in the special case n=1, the formula of Lemma 3.1.1(ii) says that the map  $\iota_{\Delta}: \Omega_R^1 A \to [A, \Omega_R^0 A]^R = [A, A]^R$  is given by  $\iota_{\Delta}(a_0 da_1) = \mathbf{e}'[a_1, a_0]\mathbf{e}''$ ; in particular, we have  $\iota_{\Delta}(da) = 0$ , for any  $a \in A$ . Therefore, since  $i_{\theta_a} \omega = da$ , we conclude that  $\theta_a(\mathbf{w}) = -\iota_{\Delta}(da) = 0$ , by Proposition 4.2.4(i). Part (ii) follows.

Finally, let  $\theta \in \operatorname{Der}_R^{\omega} A$ . Then  $\theta(\mathbf{w}) \in R$  by (i). Thus, in the setting of Proposition 4.1.4, for  $\mathbf{w} = \widetilde{\mu_{\rm nc}}(\omega)$ , we obtain  $\theta(\mathbf{w}) \in R \cap \theta([A,A]) \subset R \cap [A,A] = 0$ . This proves part (iii).

Remark 4.2.7. Let  $\mathbf{w} \in A$  be a representative of the class  $\mu_{\rm nc}(\omega) \in A/R$ . Then, for any derivation  $\theta \in \operatorname{Der}_R A$ , the element  $\theta(\mathbf{w}) = i_{\theta}(\imath_{\Delta}\omega)$  is independent of the choice of  $\mathbf{w}$ .

**4.3** Hamilton operators. We are going to introduce the notion of *Hamilton operator*, motivated by a similar construction used by Gelfand and Dorfman in their work on integrable systems, cf. [GD] and [Do].

Let  $\mathfrak{G}$  be a (possibly infinite dimensional) Lie algebra and  $(C^{\bullet}, d)$  a complex with differential  $d: C^{\bullet} \to C^{\bullet+1}$ . By a  $\mathfrak{G}$ -equivariant structure on  $C^{\bullet}$  we mean a pair of linear maps  $\mathfrak{G} \to \operatorname{Hom}_{\mathbb{k}}(C^{\bullet}, C^{\bullet}), x \mapsto L_x$ , and  $\mathfrak{G} \to \operatorname{Hom}_{\mathbb{k}}(C^{\bullet}, C^{\bullet-1}), x \mapsto i_x$ , which satisfy the standard commutation relations:

$$[L_x, L_y] = L_{[x,y]}, \quad L_x = d \circ i_x + i_x \circ d, \quad i_x \circ i_y + i_y \circ i_x = 0, \quad [L_x, i_y] = i_{[x,y]},$$

for any  $x, y \in \mathfrak{G}$ . In this case, we call  $(C^{\bullet}, d)$  a  $\mathfrak{G}$ -equivariant complex.

**Definition 4.3.1.** Let  $(C^{\bullet}, d)$  be a  $\mathfrak{G}$ -equivariant complex. A linear map  $H: C^1 \longrightarrow \mathfrak{G}$  is said to be skew-symmetric if one has

$$i_{\mathsf{H}(\alpha_1)}\alpha_2 + i_{\mathsf{H}(\alpha_2)}\alpha_1 = 0, \quad \forall \alpha_1, \alpha_2 \in C^1.$$
 (4.3.2)

A skew-symmetric map H is called a *Hamilton operator* if one has

$$i_{\mathsf{H}(\alpha_3)}L_{\mathsf{H}(\alpha_1)}\alpha_2 + \text{cyclic permutations of } \{1,2,3\} = 0, \quad \forall \alpha_1, \alpha_2, \alpha_3 \in C^1.$$

$$(4.3.3)$$

Given a skew-symmetric map H, we introduce the following bilinear pairings

$$(0) \quad \{-,-\}^0_{\mathsf{H}}: \ C^0 \times C^0 \, \longrightarrow \, C^0, \quad (p,q) \longmapsto \{p,q\}^0_{\mathsf{H}}:=i_{\mathsf{H}(dp)}dq;$$

(1) 
$$\{-,-\}_{\mathsf{H}}^1: C^1 \times C^1 \longrightarrow C^1,$$
 
$$(\alpha,\beta) \longmapsto \{\alpha,\beta\}_{\mathsf{H}}^1:=i_{\mathsf{H}(\alpha)} \circ d\beta - i_{\mathsf{H}(\beta)} \circ d\alpha + d \circ i_{\mathsf{H}(\alpha)}\beta$$
 (4.3.4)

The main results about Hamilton operators, due to Gelfand and Dorfman, may be summarized as follows.

**Proposition 4.3.5.** (i) A skew-symmetric map H is a Hamilton operator if and only if one has

$$[\mathsf{H}(\alpha),\mathsf{H}(\beta)]=\mathsf{H}\big(\{\alpha,\beta\}^1_\mathsf{H}\big),\quad\forall\alpha,\beta\in C^1.$$

- (ii) For any Hamilton operator H, the bilinear pairing given by the first, resp. second, formula in (4.3.4) makes  $C^0$ , resp.,  $C^1$ , a Lie algebra.
- (iii) With these Lie algebra structures, the maps  $C^0 \xrightarrow{d} C^1 \xrightarrow{\mathsf{H}} \mathfrak{G}$  are both Lie algebra homomorphisms.

The Lie bracket on  $C^0$  given by the first formula in (4.3.4) is usually referred to as the *Poisson bracket* on  $C^0$  induced by H.

A basic example of Hamilton operators is provided by Poisson bivectors. Specifically, let X be a smooth manifold, and write  $\mathcal{T}(X)$  for the Lie algebra of vector fields on X and  $\Omega^{\bullet}(X)$  for the graded space of differential forms on X. The de Rham complex,  $(\Omega^{\bullet}(X), d)$ , has an obvious structure of  $\mathcal{T}(X)$ -equivariant complex. Now, to each bivector  $\pi \in \Gamma(X, \wedge^2 \mathcal{T}_X)$  one associates a skew-symmetric map  $H_{\pi}: \Omega^1(X) \to \mathcal{T}(X)$ ,  $\alpha \mapsto i_{\pi}\alpha$ . This map is a Hamilton operator if and only if the bivector  $\pi$  defines a Poisson structure on X, which holds if and only if  $\pi$  has vanishing Schouten bracket:  $[\pi, \pi] = 0$ .

**4.4** Hamilton operator arising from a bi-symplectic form. The  $\mathcal{T}(X)$ -equivariant complex of differential forms on a manifold X, considered above, may be generalized to the noncommutative setup. Specifically, for any R-algebra A, the Karoubi-de Rham complex  $\mathrm{DR}_R^{\bullet}A$  has an obvious structure of  $\mathrm{Der}_R A$ -equivariant complex.

Any symplectic or bi-symplectic 2-form  $\omega \in DR_R^2 A$  gives rise to a Hamilton operator  $H_{\omega}: DR_R^1 A \to Der_R A$ . This is the map  $H_{\omega}$  introduced in Sect. 4.2, see (4.2.1). One can verify that the equation  $d\omega = 0$  implies condition (4.3.3) in the definition of Hamilton operator.

According to Proposition 4.3.5, the Hamilton operator  $H_{\omega}: DR_R^1 A \to Der_R A$  gives the space  $A_{\natural} = DR_R^0 A$  a Lie algebra structure. Thus, we obtain the following result, first discovered by Kontsevich, cf. [Ko],[BLB],[Gi]:

**Proposition 4.4.1.** For any symplectic or bi-symplectic 2-form  $\omega \in DR_R^2A$ , the pairing  $(p,q) \longmapsto \{p,q\} := \theta_p(q) \mod [A,A]$  gives  $A_{\natural}$  a Lie algebra structure.

With this structure, the map  $A_{\natural} \to \operatorname{Der}_R^{\omega} A$ ,  $p \longmapsto \theta_p$  becomes a Lie algebra homomorphism.

Recall that we have a moment map  $\mu_{\rm nc}:({\rm DR}_R^2A)_{\rm closed}\to A/R$ . Let  ${\bf w}\in A$  be a representative of the class  $\mu_{\rm nc}(\omega)\in A/R$ , and write  $R[{\bf w}]$  for the subalgebra of A generated by R and  ${\bf w}$ . Notice that the subalgebra  $R[{\bf w}]$  is independent of the choice of  ${\bf w}$ . Notice further that, by Proposition 4.1.3(i), we may always find a representative  ${\bf w}$  that commutes with R. This justifies the notation  $R[{\bf w}]$ .

**Lemma 4.4.2.** The image of the composite  $R[\mathbf{w}] \hookrightarrow A \twoheadrightarrow A_{\natural}$  is contained in the center of the Lie algebra  $A_{\natural}$ .

*Proof.* For any  $p \in A_{\natural}$ ,  $r', r'' \in R$ , and m = 1, 2, ..., we compute

$$\{p, r' \cdot \mathbf{w}^m \cdot r''\} = \theta_p(r' \cdot \mathbf{w}^m \cdot r'') = r' \cdot \theta_p(\mathbf{w}^m) \cdot r'' = r' \cdot \left(\sum_{k=0}^m \mathbf{w}^k \cdot \theta_p(\mathbf{w}) \cdot \mathbf{w}^{m-k}\right) \cdot r''.$$

Each term in the last sum vanishes since  $\theta_p(\mathbf{w}) = 0$ , by Lemma 4.2.6(ii).

Next, write  $A\mathbf{w}A\subset A$  for the two-sided ideal generated by  $\mathbf{w}$ . The algebra  $A_{\mathbf{w}}:=A/A\mathbf{w}A$  may be thought of as a Hamiltonian reduction of A.

The following important result will be proved in Sect. 7.1.

**Proposition 4.4.3.** (i) For any  $p \in A_{\natural}$  the derivation  $\theta_p$  preserves the ideal A**w**A, hence induces a well-defined derivation  $\bar{\theta}_p : A_{\mathbf{w}} \to A_{\mathbf{w}}$ .

- (ii) The Lie bracket on  $A_{\natural}$  descends to a well-defined Lie bracket on  $(A_{\mathbf{w}})_{\natural}$ .
- (iii) There is a Lie algebra map  $\bar{\varphi}$  making the following diagram commute

$$\begin{array}{ccc} A_{\natural} & \xrightarrow{p \mapsto \bar{\theta}_p} & \operatorname{Der}_R(A_{\mathbf{w}}) \\ & & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & \\ & & & \\ & & & \\ & & \\ & & & \\ & & \\ & & & \\ & \\ & & \\ & \\ & & \\ & & \\ & & \\ & \\ & & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ &$$

**4.5** Hamilton operators via Hochshild homology. Another class of examples of equivariant complexes arises from Hochschild homology.

In more detail, given an R-algebra A and an A-bimodule M, let  $H_R^k(A,M)$ , resp.  $H_k^R(A,M)$ , denote the k-th Hochschild cohomology, resp. Hochschild homology, group (with coefficients in M) of the algebra A relative to the subalgebra R. In particular, we have  $H_R^0(A,M)=M^A$ , the centralizer of A in M, and  $H_0^R(A,M)=M/[A,M]$ , the commutator quotient of M. Also, we have  $H_R^1(A,M)=\operatorname{Der}_R(A,M)/\operatorname{Inn}_R(A,M)$ .

Remark 4.5.1. If the algebra R is semisimple, then one has a canonical isomorphism  $H_R^{\bullet}(A,M) \cong H^{\bullet}(A,M)$ , between relative and absolute Hochschild cohomology of A, cf. [Lo, §1.2.13]. However, this isomorphism will play no role below.

In the special case M=A, we shall use simplified notation  $HH^k_R(A):=H^k_R(A,A)$ , resp.  $HH^R_k(A)=H^R_k(A,A)$ .

We recall that Hochschild cohomology of an associative algebra has a natural Gerstenhaber bracket  $HH_R^{k-1}(A) \times HH_R^{l-1}(A) \longrightarrow HH_R^{k+l-1}(A)$ . This bracket, together with cup-product, give  $HH_R^{\bullet}(A)$  the structure of a Gerstenhaber algebra. The Gerstenhaber bracket on  $HH_R^{l}(A)$  is the one induced by the commutator bracket on derivations.

Further, there is a natural contraction-pairing  $i: HH^k_R(A) \times HH^R_l(A) \to HH^R_{l-k}(A)$  and Lie derivative pairing  $L: HH^k_R(A) \times HH^R_l(A) \to HH^R_{l-k+1}(A)$ 

which make Hochschild homology a module over the Gerstenhaber algebra  $HH_R^{\bullet}(A)$ . Furthermore, the resulting Lie derivative action of the Lie algebra  $HH_R^1(A)$  on Hochschild homology may be promoted to a structure of  $HH_R^1(A)$ -equivariant complex on  $HH_{\bullet}^R(A)$ , with the Connes differential  $B: HH_{\bullet}^R(A) \to HH_{\bullet+1}^R(A)$  playing the role of differential, see [GDT], [Lo, Ch. 3].

In particular, in the special case l=0 we get an action of the Lie algebra  $HH_R^1(A)$  on  $A_{\natural}$ . It is easy to verify that this action is induced by the tautological  $\operatorname{Der}_R A$ -action on A; the induced  $\operatorname{Der}_R A$ -action on  $A_{\natural}$  clearly descends to  $\operatorname{Der}_R A / \operatorname{Inn}_R A$  since any inner derivation maps A to [A, A].

To construct Hamilton operators for this equivariant complex, we exploit the main idea due to Van den Bergh [VB1].

To this end, observe that the inner bimodule structure on  $A\otimes A$  survives in  $H_R^{\bullet}(A,A\otimes A)$ , hence, makes each cohomology group  $HH_R^k(A,A\otimes A)$  an A-bimodule. Now, fix an integer d>0 and assume that  $HH_R^k(A,A\otimes A)=0$  unless k=d. Assume, in addition, that A has finite Hochschild dimension, cf. Correction to [VB1]. Then, according to (a relative version of) [VB1, Thm. 1], one has canonical duality isomorphisms

$$H_k^R(A, H_R^d(A, A \otimes A)) \cong HH_R^{d-k}(A), \quad \forall k = 0, 1, \dots$$
 (4.5.2)

For any *central* element  $\pi \in H_R^d(A, A \otimes A)^A$ , we introduce the following composition of A-bimodule maps

$$\mathsf{H}_{\pi}: HH_{k}^{R}(A) \xrightarrow{a \mapsto a \cdot \pi} H_{k}^{R}(A, H_{R}^{d}(A, A \otimes A)) \xrightarrow{(4.5.2)} HH_{R}^{d-k}(A). \tag{4.5.3}$$

A central element  $m \in M^A$  of an A-bimodule M is said to be a unit if the map  $a \mapsto a \cdot m = m \cdot a$  yields an A-bimodule isomorphism  $A \stackrel{\sim}{\to} M$ . Thus, any bimodule that has a unit is non-canonically isomorphic to A as an A-bimodule.

Now, assume that one has a unit  $\pi \in H^d_R(A, A \otimes A)^A$ . Then, the composite map in (4.5.3) provides an A-bimodule isomorphism  $H_{\pi} : HH^R_{\bullet}(A) \xrightarrow{\sim} HH^{d-\bullet}_R(A)$ . We may use this isomorphism to transport the Connes differential  $B : HH^R_{\bullet}(A) \to HH^R_{\bullet+1}(A)$  to a differential  $\partial_{\pi} : HH^R_{\bullet}(A) \to HH^{\bullet-1}_R(A)$ .

Claim 4.5.4. (i) For any  $\pi \in H^d_R(A, A \otimes A)^A$ , the morphism in (4.5.3) intertwines contraction and cup-product maps, i.e., we have

$$\mathsf{H}_{\pi}(i_{n}\alpha) = \eta \cup \mathsf{H}_{\pi}(\alpha), \quad \forall \alpha \in HH_{\bullet}(A), \ \eta \in HH^{\bullet}(A).$$

(ii) Assume in addition that the algebra A has finite Hochschild dimension and there is an integer d > 0 such that one has A-bimodule isomorphisms (Gorenstein type property):

$$H_R^k(A, A \otimes A) \cong \begin{cases} A & \text{if } k = d \\ 0 & else. \end{cases}$$
 (4.5.5)

Then, for any choice of unit  $\pi \in H^d_R(A, A \otimes A)^A$ , the differential  $\partial_{\pi}$  makes Hochschild cohomology, equipped with the standard Gerstenhaber algebra structure, into a Batalin-Vilkovisky algebra, in other words, for the Gerstenhaber

bracket on Hochschild cohomology one has

$$\{u,v\} = \partial_{\pi}(u \cdot v) - \partial_{\pi}(u) \cdot v - (-1)^{k} u \cdot \partial_{\pi}(v), \quad \forall u \in HH_{R}^{k}(A), v \in HH_{R}^{l}(A).$$

We defer the proof of this Claim to a separate publication.

Example 4.5.6. Let  $A = \mathbb{k}[X]$  be the coordinate ring of a smooth affine algebraic variety X of dimension  $d := \dim X$ . In this case we have  $HH^k_R(A, A \otimes A) = 0$  unless k = d. A choice of unit  $\pi \in H^d_R(A, A \otimes A)^A = \Gamma(X, \wedge^d \mathscr{T}_X)$  corresponds to a choice of a nowhere vanishing section of the canonical sheaf  $K_X = \Omega^d_X$ . This section gives a trivialization of  $K_X$ . Thus, X is a Calabi-Yau manifold, and the Batalin-Vilkovisky structure of Claim 4.5.4(ii) is nothing but the standard Batalin-Vilkovisky structure on the space  $\bigoplus_{k\geq 0} \Gamma(X, \wedge^k \mathscr{T}_X)$  associated with the chosen volume form  $\pi^{-1} \in \Omega^d_X$ , cf. e.g. [Sch].

The special case where property (4.5.5) holds for d=2 leads to a construction of Hamilton operators. Specifically, one has the following result that may be thought of as a homological analogue of the construction of Poisson brackets used in [VB2].

**Proposition 4.5.7.** Let A be an R-algebra of finite Hochschild dimension such that condition (4.5.5) holds for d = 2.

Then, for any choice of unit  $\pi \in H^2_R(A, A \otimes A)^A$ , the corresponding map  $H_{\pi}: HH_1^R(A) \xrightarrow{\sim} HH_R^1(A)$  in (4.5.3) is a Hamilton operator for  $HH_{\bullet}^R(A)$ , viewed as an  $HH_R^1(A)$ -equivariant complex.

In particular, the space A/[A, A] acquires a natural Lie bracket  $\{-, -\}_{\pi}$ , cf. Proposition 4.3.5(ii).

The proposition above easily follows from Claim 4.5.4. To see this, let  $\mathsf{H} := \mathsf{H}_{\pi}$ , pick  $\alpha, \beta \in HH_1(A)$ , and put  $u := \mathsf{H}(\alpha)$  and  $v = \mathsf{H}(\beta)$ . We apply the isomorphism  $\mathsf{H}^{-1} : HH^1(A) \xrightarrow{\sim} HH_1(A)$  to the displayed formula in Claim 4.5.4(ii). Using part (i) of Claim 4.5.4 and writing  $[-,-]_{\mathrm{Gerst}}$  for the commutator in  $HH^1(A)$ , we obtain

$$\mathsf{H}^{-1}\big([\mathsf{H}(\alpha),\mathsf{H}(\beta)]_{\mathrm{Gerst}}\big) = \mathsf{B}(i_{\mathsf{H}(\alpha)}\beta) + i_{\mathsf{H}(\alpha)}\mathsf{B}(\beta) - i_{\mathsf{H}(\beta)}\mathsf{B}(\alpha).$$

The right hand side here is nothing but the bracket  $\{\alpha, \beta\}_{\mathsf{H}}^1$  introduced in (4.3.4). We see that the above equation is exactly the one from Proposition 4.3.5(i).  $\square$ 

# 5 Noncommutative cotangent bundle.

**5.1** Following an idea of [CB2], for any R-algebra B, we define  $T^*B := T_B(\mathbb{D}\mathbf{er}_R B)$ , the tensor algebra of the B-bimodule  $\mathbb{D}\mathbf{er}_R B$ . Thus,  $T^*B = \bigoplus_{k\geq 0} T_k^* B$ , is a graded B-algebra such that  $T_0^*B = B$ . The algebra  $T^*B$  may be thought of as the coordinate ring of the 'noncommutative cotangent bundle' on Spec B, a 'noncommutative space'. Indeed, we will see later that the representation functor takes the algebra  $T^*B$  to the cotangent bundle on the representation scheme for the algebra B.

Write  $\Delta_B$  for the  $\Delta$ -derivation  $\Delta_B: B \to B \otimes B$ ; this derivation will be frequently viewed as an element of  $T_1^*B = \mathbb{D}\mathbf{er}_R B$ . We also have the  $\Delta$ -derivation  $\Delta_{T^*B}: T^*B \to T^*B \otimes T^*B$ , and an Euler derivation  $\mathsf{E}: T^*B \to T^*B$  arising from the grading on  $T^*B$ , thus,  $\mathsf{E}(a) = k \cdot a$ , for any  $a \in T_k^*B$ .

**Theorem 5.1.1.** Assume that B is smooth. Then, the algebra  $T^*B$  is also smooth and it has a canonical bi-symplectic 2-form  $\omega \in DR_R^2(T^*B)$ , such that  $L_{\mathsf{E}}\omega = \omega$ , and such that the derivation  $\Delta_B \in T_1^*B = \mathbb{D}\mathbf{er}_R B$  is a representative of the class  $\mu_{\mathrm{nc}}(\omega)$ .

Remark 5.1.2. The bi-symplectic structure on  $T^*B$  is closely related to the 'double bracket' on  $T^*B$  introduced by Van den Bergh [VB2].

In more detail, for any  $p \in T^*B$ , let  $\Theta_p : T^*B \to T^*B \otimes T^*B$  denote the double derivation corresponding to the 1-form dp under the isomorphism  $\mathbb{D}\mathbf{er}_R(T^*B) \cong \Omega^1_R(T^*B)$  provided by the bi-symplectic form of Theorem 5.1.1. Given  $\Phi, \Psi \in \mathbb{D}\mathbf{er}_R B$ , view these double derivations of B as two degree 1 elements of  $T^*B$ , and let  $p := \Phi$ . Then, we have a well-defined element  $\Theta_{\Phi}(\Psi) \in T^*B \otimes T^*B$ . This is again an element of degree 1 with respect to the induced grading on the tensor product, i.e., an element of  $(B \otimes \mathbb{D}\mathbf{er}_R B) \oplus (\mathbb{D}\mathbf{er}_R B \otimes B)$ . The map  $T^*B \otimes T^*B \to T^*B \otimes T^*B$ ,  $\Phi \otimes \Psi \longmapsto \Theta_{\Phi}(\Psi)$  can be shown, see Appendix to [VB2], to be equal to Van den Bergh's double bracket.

Theorem 5.1.1 combined with Proposition 4.4.1 provides the graded space  $(T^*B)_{\natural}$  with a natural Lie algebra structure such that the Lie bracket has degree (-1). Using the explicit construction of the bi-symplectic 2-form of the Theorem given in Sect. 5.4 below, one proves

**Corollary 5.1.3.** (i) For any R-algebra A, the space  $(\mathbb{D}\mathbf{er}_R A)_{\natural}$  has a natural Lie algebra structure.

(ii) With this Lie algebra structure on  $(\mathbb{D}\mathbf{er}_R A)_{\natural}$  and the standard Lie algebra structure on  $\mathrm{Der}_R A$  given by the commutator of derivations, the canonical map  $\mathbf{m}_* : (\mathbb{D}\mathbf{er}_R A)_{\natural} \to \mathrm{Der}_R A$ , cf. Proposition 2.3.2, becomes a Lie algebra morphism.

Next, following [CB2], for any  $r \in R$  we define an associative R-algebra  $\Pi^r(B)$  to be the quotient of  $T^*B$  by the two-sided ideal generated by the element  $\Delta_B - r \in T_1^*B + T_0^*B$ .

For r = 0, the grading on  $T^*B$  induces a grading  $\Pi^0(B) = \bigoplus_{k \geq 0} \Pi_k^0(B)$  on the algebra  $\Pi^0(B)$ . Therefore, the Euler derivation E descends to a derivation E:  $\Pi^0(B) \to \Pi^0(B)$ .

For a general  $r \in R$ , the grading on  $T^*B$  gives rise to an increasing filtration  $F_0\Pi^r(B) \subset F_1\Pi^r(B) \subset \ldots$  Writing  $\operatorname{gr} \Pi^r(B)$  for the corresponding associated graded algebra, one has a natural surjective graded algebra map  $\Pi^0(B) \twoheadrightarrow \operatorname{gr} \Pi^r(B)$ .

Remark 5.1.4. It has been shown in [CB2], that

- There is a canonical isomorphism  $\Pi^0(B) \cong T_B^{\bullet}(H_R^1(B, B \otimes B)).$
- If  $B = \mathbb{k}[X]$  is the coordinate ring of a smooth affine curve X, then we have  $\Pi^0(\mathbb{k}[X]) \cong \mathbb{k}[T^*X]$ , the coordinate ring of the total space of the cotangent bundle on X.

• If  $B = \mathbb{k}Q$  is the path algebra of a quiver Q, then  $\Pi^r(\mathbb{k}Q) \cong \Pi^r(Q)$  is the corresponding deformed preprojective algebra, as defined in [CBH].

For r=0, the grading on the algebra  $\Pi^0(B)$  induces a grading  $\Pi^0(B)_{\natural} = \bigoplus_{k\geq 0} \Pi^0_k(B)_{\natural}$ . With respect to this grading, the Lie bracket on  $\Pi^0(B)_{\natural}$  has degree (-1), that is, we have  $\{-,-\}:\Pi^0_k(B)_{\natural}\times\Pi^0_l(B)_{\natural}\to\Pi^0_{k+l-1}(B)_{\natural}$ , for any  $k,l\geq 0$ . Therefore, the map  $\mathsf{E}_{\mathrm{Lie}}:\Pi^0(B)_{\natural}\to\Pi^0(B)_{\natural}$ , such that  $\mathsf{E}_{\mathrm{Lie}}(a):=(k-1)\cdot a, \, \forall a\in\Pi^0_k(B)_{\natural}, \, k=0,1,\ldots$ , is a derivation of the Lie algebra  $\Pi^0(B)_{\natural}$ . We let  $\Bbbk\cdot\mathsf{E}_{\mathrm{Lie}}\ltimes\Pi^0(B)_{\natural}$  denote the corresponding semi-direct product Lie algebra.

The following result will play an important role in applications to preprojective algebras associated with quivers.

**Proposition 5.1.5.** (i) The Lie bracket on  $(T^*B)_{\natural}$  descends to a Lie algebra structure on  $\Pi^r(B)_{\natural}$ , and there is a natural Lie algebra morphism  $\Pi^r(B)_{\natural} \to HH^1_R(\Pi^r(B)), \ p \mapsto \overline{\theta}_p$ .

(ii) For r=0, the assignment  $p\mapsto \bar{\theta}_p$ ,  $\mathsf{E}_{\mathrm{Lie}}\mapsto \mathsf{E}$ , gives rise to a Lie algebra morphism  $\Bbbk\cdot\mathsf{E}_{\mathrm{Lie}}\ltimes(\Pi^0(B)_{\natural}/R)\to HH^1_R(\Pi^0(B))$ . This morphism is injective provided one has  $(\mathrm{DR}^0_RB)_{\mathrm{closed}}=R$ .

Part (i) of the Proposition is immediate from Proposition 4.4.3 applied to the algebra  $A = T^*B$ . Further, it is straightforward to see that the map of part (ii) is a Lie algebra morphism. The injectivity statement in part (ii) is more complicated; it will be proved later, in Sect. 7.4.

Note that condition  $(DR_R^0 B)_{closed} = R$  in part (ii) may be interpreted as saying that the algebra B is *connected*.

**5.2** The Liouville 1-form. We recall the structure of  $\Omega_R^1(T_BM)$ , the bimodule of noncommutative 1-forms on the tensor algebra  $A := T_BM$  of an arbitrary B-bimodule M. To this end, introduce the following A-bimodule

$$\widetilde{\Omega} := (A \otimes_B \Omega_R^1 B \otimes_B A) \bigoplus (A \otimes_R M \otimes_R A). \tag{5.2.1}$$

Abusing the notation slightly, for any  $a', a'' \in A$ ,  $m \in M, \beta \in \Omega^1_R B$ , we write  $a' \cdot \widetilde{m} \cdot a'' := 0 \oplus (a' \otimes m \otimes a'') \in \widetilde{\Omega}$  and  $a' \cdot \widetilde{\beta} \cdot a'' := (a' \otimes \beta \otimes a'') \oplus 0 \in \widetilde{\Omega}$ .

Let Q denote the A-subbimodule in  $\Omega$  generated by the following set

$$\{(\widetilde{b'mb''}) - \widetilde{db'} \cdot (mb'') - b' \cdot \widetilde{m} \cdot b'' - (b'm) \cdot \widetilde{db''}\}_{\{b',b'' \in B, m \in M\}}$$
 (5.2.2)

**Lemma 5.2.3.** Let M be a projective  $B^e$ -module, and  $A := T_BM$ . Then

- (i) There is an A-bimodule isomorphism  $\Omega^1_R A \cong \Omega/Q$ .
- (ii) The imbedding of the first direct summand in  $\widetilde{\Omega}$ , resp. the projection to the second direct summand in  $\widetilde{\Omega}$ , induces, via the isomorphism in (i), a canonical extension of A-bimodules

$$0 \longrightarrow A \otimes_B \Omega^1_R B \otimes_B A \stackrel{\varepsilon}{\longrightarrow} \Omega^1_R A \stackrel{\nu}{\longrightarrow} A \otimes_B M \otimes_B A \longrightarrow 0.$$

(iii) The assignment  $B \oplus M = T_B^0 M \oplus T_B^1 M \to \widetilde{\Omega}$ ,  $b \oplus m \mapsto d\widetilde{b} + \widetilde{m}$  extends uniquely to a derivation  $\widetilde{d}: A = T_B^0 M \to \widetilde{\Omega}/Q$ ; this derivation corresponds, via the isomorphism in (i), to the canonical universal derivation  $d: A \to \Omega_B^1 A$ .

(iv) If B is smooth and M is finitely generated (as a  $B^e$ -module), then the algebra  $A = T_BM$  is also smooth.

*Proof.* First of all, using the relations (5.2.2) it is straightforward to verify that the assignment  $b \oplus m \longmapsto \widetilde{db} + \widetilde{m}$  gives rise to a derivation  $\widetilde{d}: A \to \widetilde{\Omega}/Q$ .

Now, let E be an A-bimodule and let  $\delta: A \to E$  be a derivation. Again, the relations (5.2.2) insure that the assignment  $\widetilde{db} + \widetilde{m} \longmapsto \delta(b) + \delta(m)$  extends to a well-defined A-bimodule map  $\widetilde{\Omega}/Q \to E$ . It follows that the derivation  $\widetilde{d}: A \to \widetilde{\Omega}/Q$  enjoys the universal property for the bimodule of 1-forms. Thus, we conclude that  $\Omega^1_R A \cong \widetilde{\Omega}/Q$ . This proves (i) and (iii).

Part (ii) is clear from (5.2.2), see also [CQ, Corollary 2.10]; part (iv) is [CQ, Proposition 5.3(3)].

Recall that, associated with any 1-form  $\alpha \in DR_R^1B$ , there is a B-bimodule map  $\iota(\alpha): \mathbb{D}\mathbf{er}_R B \to B$ ,  $\Theta \mapsto \iota_{\Theta}\alpha$ . This map extends uniquely to an algebra homomorphism  $\iota(\alpha): T^*B = T_B(\mathbb{D}\mathbf{er}_R B) \longrightarrow B$ , such that  $\iota(\alpha)|_{T_0^*B} = \mathrm{Id}_B$  and  $\iota(\alpha)|_{T_1^*B} = \iota(\alpha)$ . Finally, the algebra homomorphism  $\iota(\alpha)$  induces, by functoriality, a DG algebra homomorphism  $\iota(\alpha)_*: \Omega_R^{\bullet}(T^*B) \to \Omega_R^{\bullet}B$ .

**Proposition 5.2.4.** Assume that B is smooth. Then, there is a canonical Liouville 1-form  $\lambda \in \mathrm{DR}^1_R(T^*B)$  such that the following holds:

- (i) One has  $i_{\mathsf{E}}\lambda = 0$ , and  $L_{\mathsf{E}}\lambda = \lambda$ .
- (ii) In  $T_1^*B = \mathbb{D}\mathbf{er}_R B$ , we have  $i_{\Delta_{T^*B}}\lambda = \Delta_B$ .
- (iii) For any 1-form  $\alpha \in DR_B^1 B$ , in  $DR_B^1 B$ , we have  $\iota(\alpha)_*(\lambda) = \alpha$ .

Remark 5.2.5. The Proposition above is a noncommutative analogue of a similar result for the standard Liouville 1-form on  $T^*X$ , the cotangent bundle on a manifold X. To explain this, observe that any 1-form  $\alpha$  on X is, by definition, a section  $\alpha: X \to T^*X$  of the cotangent bundle. Now, equation  $\iota(\alpha)_*(\lambda) = \alpha$  of the Proposition is an analogue of the equation  $\alpha^*(\lambda) = \alpha$ , for the pull-back of the Liouville 1-form on  $T^*X$  via the section  $\alpha: X \to T^*X$ .

To construct the Liouville 1-form, we consider the following natural maps

$$(\mathbb{D}\mathbf{er}_R B) \otimes_{B^e} \Omega_R^1 B = \mathrm{Hom}_{B^e}(\Omega_R^1 B, B \otimes B) \otimes_{B^e} \Omega_R^1 B \to \mathrm{Hom}_{B^e}(\Omega_R^1 B, \Omega_R^1 B).$$

Since B is smooth, the last map is a bijection, cf. Proposition 2.3.2. We invert this bijection and let  $\Lambda \in (\mathbb{D}\mathbf{er}_R B) \otimes_{B^e} \Omega^1_R B$  denote the canonical element corresponding to the identity  $\mathrm{Id} \in \mathrm{Hom}_{B^e}(\Omega^1_R B, \Omega^1_R B)$  under the composite (from right to left) of the inverse maps in the displayed formula above.

We can write  $\Lambda = \sum_s \wp_s \otimes \gamma_s$ , for some  $\wp_s \in \mathbb{D}\mathbf{er}_R B$  and  $\gamma_s = db_s$ , where  $b_s \in B$  and  $s = 1, \ldots, l$ . The map  $\Omega_R^1 B \to \Omega_R^1 B$  that corresponds to the element  $\sum_s \wp_s \otimes \gamma_s$  is given by the formula  $\alpha \longmapsto \sum_s (i'_{\wp_s} \alpha) \cdot \gamma_s \cdot (i''_{\wp_s} \alpha)$ . By construction, this should be the identity map; thus we must have

$$\sum_{s} (i'_{\wp_s} \alpha) \cdot \gamma_s \cdot (i''_{\wp_s} \alpha) = \alpha, \quad \forall \alpha \in \Omega^1_R B.$$
 (5.2.6)

Now, by Lemma 5.2.3(ii) applied to the *B*-bimodule  $M = \mathbb{D}\mathbf{er}_R B$ , there is a canonical  $T^*B$ -bimodule imbedding  $\varepsilon : T^*B \otimes_B \Omega^1_R B \otimes_B T^*B \hookrightarrow \Omega^1_R (T^*B)$ .

Applying the commutator quotient functor  $E \longmapsto E/[T^*B, E]$  to the  $T^*B$ -bimodule map  $\varepsilon$  we get the following chain of natural maps

$$\frac{T^*B\otimes_B\Omega^1_RB}{[B,\,T^*B\otimes_B\Omega^1_RB]} = \frac{T^*B\otimes_B\Omega^1_RB\otimes_BT^*B}{[T^*B,\,T^*B\otimes_B\Omega^1_RB\otimes_BT^*B]} \stackrel{\varepsilon}{\longrightarrow} \frac{\Omega^1_R(T^*B)}{[T^*B,\,\Omega^1_R(T^*B)]}.$$

The leftmost term in this formula is nothing but  $T^*B \otimes_{B^e} \Omega^1_R B$  and the rightmost term equals  $DR^1_R(T^*B)$ . Thus, we have canonical maps

$$(\mathbb{D}\mathbf{er}_R\,B)\otimes_{B^e}\Omega^1_RB=T_1^*B\otimes_{B^e}\Omega^1_RB\,\hookrightarrow\,T^*B\otimes_{B^e}\Omega^1_RB\to\mathrm{DR}^1_R(T^*B).$$

We define  $\lambda \in \mathrm{DR}^1_R(T^*B)$  to be the image of the canonical element  $\Lambda \in (\mathbb{D}\mathbf{er}_R B) \otimes_{B^e} \Omega^1_R B$  under the composite map above. Writing  $\Lambda = \sum_s \wp_s \otimes \gamma_s$  and using the same notation as in formulas (5.2.1)-(5.2.2), we get the following analogue of the classical expression ' $\lambda = p \, dq$ ' for the Liouville 1-form:

$$\lambda = \sum_{s=1}^{l} \wp_s \cdot \widetilde{\gamma}_s = \sum_{s=1}^{l} \wp_s \cdot \widetilde{db}_s, \quad \wp_s \in \mathbb{D}\mathbf{er}_R B = T_1^* B, \ \gamma_s = db_s, \ b_s \in B. \ (5.2.7)$$

**5.3** Given a B-bimodule M, let  $M^{\vee} := \operatorname{Hom}_{B^e}(M, B \otimes B)$  denote the dual B-bimodule equipped with the B-bimodule structure induced by the inner B-bimodule structure on  $B \otimes B$ . We have a canonical B-bimodule map **bidual**:  $M \to (M^{\vee})^{\vee}$ . This map is an isomorphism for any finitely generated projective  $B^e$ -module M.

In the special case  $M = \Omega_R^1 B$ , we have  $(\Omega_R^1 B)^{\vee} = \operatorname{Hom}_{B^e}(\Omega_R^1 B, B \otimes B) = \mathbb{D}\mathbf{er}_R B$ . Therefore, if  $\Omega_R^1 B$  is a finitely generated projective bimodule, one also has a canonical B-bimodule isomorphism bidual :  $\Omega_R^1 B \stackrel{\sim}{\to} (\mathbb{D}\mathbf{er}_R B)^{\vee}$ ,  $\alpha \mapsto \alpha^{\vee} = \operatorname{bidual}(\alpha)$ . Explicitly, for  $\alpha \in \Omega_R^1 B$ , the element  $\alpha^{\vee}$  is a map given by

$$\alpha^{\vee}: \mathbb{D}\mathbf{er}_{B} B \to B \otimes B, \quad \Theta \longmapsto -i_{\Theta}^{"}\alpha \otimes i_{\Theta}^{\prime}\alpha,$$
 (5.3.1)

where the flip of the order of tensor factors is due to the fact that the B-bimodule structure on  $\operatorname{Hom}_{B^e}(\mathbb{D}\mathbf{er}_R B, B \otimes B)$  comes from the *inner* bimodule structure on  $B \otimes B$  and the negative sign comes from the sign of the transposition permutation.

Further, we have canonical bijections

$$\mathbb{D}\mathbf{er}_R\, B \underset{\mathbb{R}^e}{\otimes} \, \Omega^1_R B \xrightarrow[\sim]{\mathrm{Id} \otimes \mathtt{bidual}} \, \mathbb{D}\mathbf{er}_R\, B \underset{\mathbb{R}^e}{\otimes} \, (\mathbb{D}\mathbf{er}_R\, B)^\vee \xrightarrow[\sim]{} \mathrm{Hom}_{B^e}(\mathbb{D}\mathbf{er}_R\, B, \mathbb{D}\mathbf{er}_R\, B).$$

The composite of these bijections sends the canonical element  $\Lambda = \sum_s \wp_s \otimes \gamma_s \in (\mathbb{D}\mathbf{er}_R B) \otimes_{B^e} \Omega_R^1 B$  to the identity map  $\mathrm{Id} : \mathbb{D}\mathbf{er}_R B \to \mathbb{D}\mathbf{er}_R B$ . This yields the following identity, dual in some sense to the identity (5.2.6):

$$\sum_{s} \Theta''(b_s) \cdot \wp_s \cdot \Theta'(b_s) = \sum_{s} i_{\Theta}''(\gamma_s) \cdot \wp_s \cdot i_{\Theta}'(\gamma_s) = \Theta, \quad \forall \Theta \in \mathbb{D}er_R B. \quad (5.3.2)$$

Proof of Proposition 5.2.4. To prove (i), we compute  $i_{\mathsf{E}}\lambda = \sum_s i_{\mathsf{E}}(\wp_s \cdot \widetilde{\gamma}_s) = \sum_s \wp_s \cdot (i_{\mathsf{E}}\widetilde{\gamma}_s) = 0$ , since  $i_{\mathsf{E}}\widetilde{\beta} = 0$  for any  $\beta \in \Omega^1_R B$ . Observe further that  $\lambda$  belongs to the image of  $T_1^*B \otimes_B \Omega^1_R B \otimes_B T_0^*B$  in  $\mathrm{DR}^1_R(T^*B)$ , cf. (5.2.7). We deduce that  $L_{\mathsf{E}}\lambda = \lambda$ .

To prove property (ii) in the Proposition, we apply formula (5.3.2) to  $\Theta = \Delta_B$  and observe that, for any  $b \in B$ , one has  $i_{\Delta_{T^*B}}\widetilde{db} = \Delta_{T^*B}(\widetilde{b}) = \widetilde{\Delta_B}b$ . Thus, in  $T^*B$ , we obtain a chain of equalities

$$i_{\Delta_{T^*B}}\lambda = \sum\nolimits_s (i_{\Delta_{T^*B}}''\widetilde{\gamma}_s) \cdot \wp_s \cdot (i_{\Delta_{T^*B}}'\widetilde{\gamma}_s) = \sum\nolimits_s \widetilde{\Delta_B''(b_s)} \cdot \wp_s \cdot \widetilde{\Delta_B'(b_s)} = \widetilde{\Delta_B}.$$

To prove (iii), fix  $\alpha \in DR_R^1 B$ . Then, we find

$$i(\alpha)_*(\lambda) = \sum_s i(\alpha)_* (\wp_s \cdot \widetilde{\gamma}_s) = \sum_s (i_{\wp_s} \alpha) \cdot \gamma_s.$$

But  $(i_{\wp_s}\alpha)\cdot\gamma_s=(i''_{\wp_s}\alpha)\cdot(i'_{\wp_s}\alpha)\cdot\gamma_s=(i'_{\wp_s}\alpha)\cdot\gamma_s\cdot(i''_{\wp_s}\alpha) \bmod [\Omega^1_RB,B]$ . Hence, using the identity in (5.2.6) we get  $\imath(\alpha)_*(\lambda)=\alpha$ , as required.

**5.4** We put  $\omega := d\lambda \in DR_R^2(T^*B)$ . This is a closed 2-form. To avoid confusion, we write  $\mathbf{w} := \Delta_B$ , viewed as an element of  $T_1^*B$ .

**Proposition 5.4.1.** (i) The 2-form  $\omega$  is bi-symplectic.

- (ii) We have  $i_{\Delta_{T^*B}}\omega = d\mathbf{w}$  and  $\lambda = i_{\mathsf{E}}\omega$ .
- (iii) For any  $p \in T_k^*B$ ,  $k = 0, 1, ..., in (T^*B)_{\natural}$ , we have:

$$i_{\text{ad }p}\lambda = p \cdot \mathbf{w}, \quad and \quad i_{\theta_n}\lambda = k \cdot p.$$

It is clear that the Proposition above implies Theorem 5.1.1.

In preparation for the proof of Proposition 5.4.1 we put  $A := T^*B$  and  $M := \mathbb{D}\mathbf{er}_R B$ . We apply the functor  $\operatorname{Hom}_{A^c}(-, A \otimes A)$  to the short exact sequence in Lemma 5.2.3(ii). For a smooth algebra B, one obtains this way the following short exact sequence

$$0 \to A \otimes_B (\mathbb{D}\mathbf{er}_R B)^{\vee} \otimes_B A \xrightarrow{\nu^{\vee}} (\Omega_R^1 A)^{\vee} \xrightarrow{\varepsilon^{\vee}} A \otimes_B (\Omega_R^1 B)^{\vee} \otimes_B A \to 0.$$

Note that the middle term in this short exact sequence may be identified with  $(\Omega_R^1(T^*B))^{\vee} \cong \mathbb{D}\mathbf{er}_R(T^*B)$ .

Now, contraction with  $\omega$  gives a  $T^*B$ -bimodule map  $\iota(\omega): \mathbb{D}\mathbf{er}_R(T^*B) \to \Omega^1_R(T^*B)$ , and we have

**Lemma 5.4.2.** Let B be a smooth algebra and  $A := T^*B$ . Then, the following diagram commutes

$$0 \to A \otimes_B (\mathbb{D}\mathbf{er}_R B)^{\vee} \otimes_B A \xrightarrow{\nu^{\vee}} \mathbb{D}\mathbf{er}_R A \xrightarrow{\varepsilon^{\vee}} A \otimes_B (\Omega_R^1 B)^{\vee} \otimes_B A \to 0$$

$$-\mathrm{Id} \otimes \mathrm{bidual} \otimes \mathrm{Id} \bigg\| \qquad \qquad \iota(\omega) \bigg| \qquad \qquad \mathrm{Id} \bigg\|$$

$$0 \to A \otimes_B (\Omega_R^1 B) \otimes_B A \xrightarrow{\varepsilon} \Omega_R^1 A \xrightarrow{\nu} A \otimes_B (\mathbb{D}\mathbf{er}_R B) \otimes_B A \to 0$$

*Proof.* We use the notation of Sect. 5.2. Thus, we have the canonical element  $\Lambda = \sum_s \wp_s \otimes \gamma_s$ , the Liouville 1-form  $\lambda = \sum_s \wp_s \cdot \widetilde{\gamma}_s$ , and the 2-form  $\omega = d\lambda = \sum_s d\wp_s \cdot \widetilde{\gamma}_s$ , where  $\gamma_s = db_s$ , see (5.2.7). Given  $\Theta \in \mathbb{D}\mathbf{er}_R(T^*B)$ , we find

$$i_{\Theta}\omega = \sum_{s} i_{\Theta}(d\wp_{s} \cdot \widetilde{\gamma}_{s})$$

$$= -\sum_{s} i_{\Theta}''(d\wp_{s}) \cdot \widetilde{\gamma}_{s} \cdot i_{\Theta}'(d\wp_{s}) + \sum_{s} i_{\Theta}''(\widetilde{\gamma}_{s}) \cdot d\wp_{s} \cdot i_{\Theta}'(\widetilde{\gamma}_{s})$$

$$= -\sum_{s} \Theta''(\wp_{s}) \cdot \widetilde{\gamma}_{s} \cdot \Theta'(\wp_{s}) + \sum_{s} \Theta''(\widetilde{b}_{s}) \cdot d\wp_{s} \cdot \Theta'(\widetilde{b}_{s}).$$
(5.4.3)

Now, let  $\alpha \in \Omega^1_R B$  and put

$$\vec{\alpha} := 1 \otimes \mathtt{bidual}(\alpha) \otimes 1 \in T^*B \otimes_B (\mathbb{D}\mathbf{er}_R B)^{\vee} \otimes_B T^*B \subset \mathbb{D}\mathbf{er}_R (T^*B).$$

Thus,  $\vec{\alpha}: T^*B \to T^*B \otimes T^*B$  is a derivation of degree (-1) that annihilates the space  $T_0^*B = B$  and acts on  $T_1^*B = \mathbb{D}\mathbf{er}_R B$  as the map  $\alpha^{\vee}: \mathbb{D}\mathbf{er}_R B \to B \otimes B = T_0^*B \otimes T_0^*B \subset T^*B \otimes T^*B$  given by formula (5.3.1). Therefore, for  $\Theta = \vec{\alpha}$ , the second sum in the last line of formula (5.4.3) vanishes, and we obtain

$$i_{\overrightarrow{\alpha}}\omega = -\sum_{s} (\overrightarrow{\alpha})''(\wp_{s}) \cdot \widetilde{\gamma}_{s} \cdot (\overrightarrow{\alpha})'(\wp_{s})$$

$$= -\sum_{s} (\alpha^{\vee})''(\wp_{s}) \cdot \widetilde{\gamma}_{s} \cdot (\alpha^{\vee})'(\wp_{s}) = \sum_{s} i'_{\wp_{s}}(\alpha) \cdot \widetilde{\gamma}_{s} \cdot i''_{\wp_{s}}(\alpha) = \widetilde{\alpha},$$

where in the third equality we used formula (5.3.1), and the last equality is due to identity (5.2.6). This proves that the left square in the diagram of the Lemma commutes.

Next, we prove the commutativity of the right square in the diagram of the Lemma. Since all maps in the diagram are  $T^*B$ -bimodule maps, it suffices to verify commutativity on any derivation  $\Theta \in \mathbb{D}\mathbf{er}_R(T^*B)$  such that  $\varepsilon^{\vee}(\Theta) \in 1 \otimes (\Omega_R^1 B) \otimes 1$ . This condition means that we have  $\Theta(T_0^* B) \subset T_0^* B \otimes T_0^* B$ . Thus,  $\Theta$  gives a derivation  $\Theta_B : B \to B \otimes B$ , and we have  $\varepsilon^{\vee}(\Theta) = 1 \otimes \Theta_B \otimes 1 \in T^*B \otimes_B (\Omega_B^1 B) \otimes_B T^*B$ . We deduce that one has  $\Theta(\widetilde{b}) = \Theta_B(b)$ , for any  $b \in B$ .

We now take  $\Theta$  as above and apply the map  $\nu$  (in the bottom row of the diagram of the Lemma) to the expression in the last line of (5.4.3). The first sum in this expression is annihilated by the map  $\nu$ . Also, we have  $\nu(d\wp_s) = \wp_s$ . Therefore, from (5.4.3) we get

$$\nu(\imath_\Theta\omega) = \sum\nolimits_s \nu\left(\Theta''(\widetilde{b}_s) \cdot d\wp_s \cdot \Theta'(\widetilde{b}_s)\right) = \sum\nolimits_s \widetilde{\Theta''_B(b_s)} \cdot \wp_s \cdot \widetilde{\Theta'_B(b_s)} = \widetilde{\Theta_B},$$

where the last equality is due to identity (5.3.2). This proves that the right square in the diagram of the Lemma commutes.  $\hfill\Box$ 

Proof of Proposition 5.4.1. To prove that  $\omega = d\lambda \in DR^2(T^*B)$  is a bi-symplectic form, we must show that the vertical map  $\iota(\omega)$  in the middle of the diagram of Lemma 5.4.2 is a bijection. But the vertical maps on the left and on the right of the diagram of Lemma 5.4.2 are both bijections. Hence, the statement follows from the commutativity of the diagram by diagram chase.

Next, by Proposition 5.2.4 we get  $d(\Delta_B) = d(i_{\Delta_{T^*B}}\lambda) = -i_{\Delta_{T^*B}}(d\lambda) = -i_{\Delta_{T^*B}}\omega$ . Also, we compute  $i_{\mathsf{E}}\omega = i_{\mathsf{E}}(d\lambda) = L_{\mathsf{E}}\lambda - d(i_{\mathsf{E}}\lambda) = \lambda$ , since  $i_{\mathsf{E}}\lambda = 0$ . Further, by Corollary 3.2.2, we know that  $i_{\mathsf{ad}\,p}\omega = p\,d\mathbf{w}$ . Thus we find

$$i_{\operatorname{ad} p}\lambda = -i_{\operatorname{ad} p}i_{\operatorname{E}}\omega = i_{\operatorname{E}}i_{\operatorname{ad} p}\omega = i_{\operatorname{E}}(p\,d\mathbf{w}) = p\cdot\operatorname{E}(\mathbf{w}) = p\cdot\mathbf{w},$$

where in the last equality we have used that the element **w** has degree 1 with respect to the grading on  $T^*B$ . Similarly, for any  $p \in T_k^*B$ , we have  $\mathsf{E}(p) = k \cdot p$ . Hence, we obtain

$$i_{\theta_n}\lambda = -i_{\theta_n}i_{\mathsf{E}}\omega = i_{\mathsf{E}}i_{\theta_n}\omega = i_{\mathsf{E}}dp = \mathsf{E}(p) = k \cdot p.$$

## 6 The representation functor

Throughout this section, k is an algebraically closed field of characteristic zero.

**6.1** Representation schemes. Let I be a finite set and  $R = \mathbb{k}I$ , the algebra of functions  $I \to \mathbb{k}$  with pointwise multiplication. For each  $i \in I$ , let  $e_i \in R$  be the characteristic function of the one point set  $\{i\} \subset I$ . We have  $1 = \sum_{i \in I} e_i$ , furthermore,  $\mathbf{e} := \sum_{i \in I} e_i \otimes e_i$  is the symmetric separability element for R.

Giving a left R-module V is the same thing as giving an I-graded vector space  $V = \bigoplus_{i \in I} V_i$ , where  $V_i = e_i V$ . For any left R-module V, the action of R gives an algebra map  $R \to \operatorname{Hom}_{\Bbbk}(V,V)$ . This makes  $\operatorname{Hom}_{\Bbbk}(V,V)$  an R-algebra, to be denoted End :=  $\operatorname{Hom}_{\Bbbk}(V,V)$  below.

From now on, let A be a *finitely presented* associative R-algebra. Given a finite dimensional left R-module  $V=\bigoplus_{i\in I}V_i$ , we may consider the set  $\operatorname{Hom}_{R\text{-alg}}(A,\operatorname{End})$  of all R-algebra maps  $\rho:A\to\operatorname{End}$ . More precisely, there is an affine seme of finite type over  $\Bbbk$ , to be denoted  $\operatorname{Rep}(A,V)$ , such that the set  $\operatorname{Hom}_{R\text{-alg}}(A,\operatorname{End})$  is the set of  $\Bbbk$ -points of  $\operatorname{Rep}(A,V)$ . The latter scheme, by definition, represents the functor  $B\longmapsto \operatorname{Hom}_{R\text{-alg}}(A,\operatorname{End}_B(B\otimes V))$ , from the category of finitely generated commutative  $\Bbbk$ -algebras to the category Sets.

Let  $\mathrm{GL}(V)^R = \prod_i \mathrm{GL}(V_i)$  be the group of R-module automorphisms of V, and let  $\mathbb{G}_m \subset \mathrm{GL}(V)^R$  be the 1-dimensional torus of scalar automorphisms of V. We put  $G := \mathrm{GL}(V)^R/\mathbb{G}_m$ , and let  $\mathfrak{g} = \mathrm{Lie}\,G$  be the Lie algebra of G. We may (and will) identify both  $\mathfrak{g}$  and its dual,  $\mathfrak{g}^*$ , with the codimension 1 subspace in  $\oplus_i \mathrm{End}_{\Bbbk} V_i$  formed by linear maps with the vanishing total trace.

From now on, we will assume that the scheme  $\operatorname{Rep}(A,V)$  is  $\operatorname{non-empty}$ , and write  $\mathbbm{k}[\operatorname{Rep}(A,V)]$  for its coordinate ring. We consider  $\mathbbm{k}[\operatorname{Rep}(A,V)] \otimes \operatorname{End}$ , a tensor product of associative algebras. The map  $R \to \operatorname{End}$  makes  $\mathbbm{k}[\operatorname{Rep}(A,V)] \otimes \operatorname{End}$  an R-algebra that may also be identified with the algebra of regular maps  $\operatorname{Rep}(A,V) \to \operatorname{End}$ , equipped with pointwise multiplication. The action of G on V makes  $\operatorname{Rep}(A,V)$  a G-scheme. This gives a G-action on  $\mathbbm{k}[\operatorname{Rep}(A,V)]$  by algebra automorphisms. We also have a G-action on End, by conjugation, and we let G act diagonally on  $\mathbbm{k}[\operatorname{Rep}(A,V)] \otimes \operatorname{End}$ . We write  $\mathbbm{k}[\operatorname{Rep}(A,V)]^G$ , resp.,  $(\mathbbm{k}[\operatorname{Rep}(A,V)] \otimes \operatorname{End})^G$ , for the corresponding subalgebra of G-invariants.

**6.2 Evaluation homomorphisms.** Write  $\operatorname{Sym} E = \bigoplus_{j\geq 0} \operatorname{Sym}_{\mathbb{k}}^{j} E$  for the symmetric algebra of a  $\mathbb{k}$ -vector space E.

Given an *I*-tuple  $\{d_i \in \mathbb{k}\}_{i \in I}$ , we define a linear function  $\mathbf{d} : R \to \mathbb{k}$  by the assignment  $e_i \longmapsto d_i$ . This function extends by multiplicativity to an algebra map  $\operatorname{Sym} R \to \mathbb{k}$ , and thus makes  $\mathbb{k}$  a 1-dimensional  $\operatorname{Sym} R$ -module to be denoted  $\mathbb{k}_{\mathbf{d}}$ .

Now let  $V = \bigoplus_{i \in I} V_i$  be a finite dimensional left R-module. We put  $d_i := \dim V_i$ , and write  $\mathbf{d} = \mathbf{d}(V) : e_i \mapsto d_i$  for the corresponding linear function  $\mathbf{d} : R \to \mathbb{k}$ 

Let A be an R-algebra. The composite map  $R \hookrightarrow A \twoheadrightarrow A_{\natural}$  induces a graded algebra homomorphism  $\operatorname{Sym} R \to \operatorname{Sym}(A_{\natural})$ . We introduce a commutative  $\Bbbk$ -algebra  $\mathcal{O}_{\mathbf{d}}(A) := \Bbbk_{\mathbf{d}} \bigotimes_{\operatorname{Sym} R} \operatorname{Sym}(A_{\natural})$ .

To each element  $a \in A$ , one associates the following evaluation function  $\widehat{a} : \operatorname{Rep}(A, V) \to \operatorname{End}, \ \rho \mapsto \widehat{a}(\rho) := \rho(a)$ . The assignment  $a \longmapsto \widehat{a}$  clearly gives an associative R-algebra homomorphism

ev: 
$$A \longrightarrow (\mathbb{k}[\operatorname{Rep}(A, V)] \otimes \operatorname{End})^G$$
,  $a \longmapsto \widehat{a}$ .

For any  $a \in A$ , composing the function  $\widehat{a}$  with the trace map  $\operatorname{Tr}: \operatorname{End} \to \Bbbk$ , applied to the second tensor factor above, we obtain a G-invariant element  $\operatorname{Tr} \widehat{a} \in \Bbbk[\operatorname{Rep}(A,V)]^G$ . If  $a \in [A,A]$ , then  $\operatorname{Tr} \widehat{a} = 0$ , due to symmetry of the trace. Thus, the assignment  $a \longmapsto \operatorname{Tr} \widehat{a}$  gives a well-defined  $\Bbbk$ -linear map  $\operatorname{Tr} \circ \operatorname{ev}: A_{\natural} \to \Bbbk[\operatorname{Rep}(A,V)]^G$ .

It is clear that the element  $\widehat{e}_i \in \mathbb{k}[\operatorname{Rep}(A, V)] \otimes \operatorname{End}$  corresponding to an idempotent  $e_i \in R = \mathbb{k}I$  is a constant function on  $\operatorname{Rep}(A, V)$  whose value equals the projector on the direct summand  $V_i$ . Therefore, we have  $\operatorname{Tr} \circ \operatorname{ev}(e_i) = \dim V_i = \operatorname{\mathbf{d}}(e_i)$ . Thus, we see that the linear map  $\operatorname{Tr} \circ \operatorname{ev} : A_{\natural} \to \mathbb{k}[\operatorname{Rep}(A, V)]^G$  may be uniquely extended, by multiplicativity, to a commutative algebra morphism

$$\psi_{\mathbf{d}}: \mathcal{O}_{\mathbf{d}}(A) \longrightarrow \mathbb{k}[\operatorname{Rep}(A, V)]^{G}, \quad \mathbf{d} = \mathbf{d}(V), 
a_{1} \& \dots \& a_{n} \longmapsto (\operatorname{Tr} \widehat{a}_{1}) \cdot \dots \cdot (\operatorname{Tr} \widehat{a}_{n}), \quad \forall a_{1}, \dots, a_{n} \in A_{\natural}.$$
(6.2.1)

There is also a natural evaluation map on differential forms that sends the Karoubi-de Rham complex of A to the ordinary de Rham complex of the representation scheme. In more detail, write  $\Omega^{\bullet}(\text{Rep}) := \Omega^{\bullet}(\text{Rep}(A, V)) = \bigwedge_{\mathbb{k}[\text{Rep}(A, V)]}^{\bullet} \Omega^{1}(\text{Rep}(A, V))$  for the DG algebra of algebraic differential forms on the scheme Rep(A, V) (in the ordinary sense of commutative algebra).

We have an algebra homomorphism

ev : 
$$\Omega^n A \longrightarrow (\Omega^n(\text{Rep}) \otimes \text{End})^G$$
,  $\alpha = a_0 da_1 \dots da_n \mapsto \widehat{\alpha} = \widehat{a}_0 d\widehat{a}_1 \dots d\widehat{a}_n$ ,

defined as a composite

$$\begin{split} \Omega^n A &= A \otimes (A/\Bbbk)^{\otimes n} \stackrel{\operatorname{ev}}{\longrightarrow} \left( \Bbbk[\operatorname{Rep}] \otimes \operatorname{End} \right) \otimes \left( \Omega^1(\operatorname{Rep}) \otimes \operatorname{End} \right)^{\otimes n} \\ &\to \left( \bigwedge\nolimits_{\Bbbk[\operatorname{Rep}]}^n \Omega^1(\operatorname{Rep}) \right) \otimes \operatorname{End}^{\otimes n+1} \stackrel{\operatorname{Id} \otimes m}{\longrightarrow} \Omega^n(\operatorname{Rep}) \otimes \operatorname{End}. \end{split}$$

Observe that since  $\widehat{r}$  is a constant function, for any  $r \in R$ , we have  $d\widehat{r} = 0$ . It follows easily that the map above induces a well-defined DG algebra morphism  $\Omega_R^{\bullet}A \to \left(\Omega^{\bullet}(\operatorname{Rep}(A,V)) \otimes \operatorname{End}\right)^G$ . Further, composing the latter morphism with the trace map  $\operatorname{Id} \otimes \operatorname{Tr} : \Omega^{\bullet}(\operatorname{Rep}(A,V)) \otimes \operatorname{End} \longrightarrow \Omega^{\bullet}(\operatorname{Rep}(A,V))$ , we thus obtain a linear map

$$\operatorname{Tr} \circ \operatorname{ev} : \operatorname{DR}_R^{\bullet} A \longrightarrow \Omega^{\bullet}(\operatorname{Rep}(A, V))^G, \quad \alpha \longmapsto \operatorname{Tr} \widehat{\alpha}.$$
 (6.2.2)

The map (6.2.2) clearly commutes with the de Rham differentials.

**6.3 Double tangent bundle.** Inspired by an idea of Kontsevich and Rosenberg [KR], we are going to discuss the effect of the action of the representation functor on double derivations.

Fix a finite dimensional left R-module V. To simplify notation, write  $\mathscr{R} := \mathbb{k}[\operatorname{Rep}(A,V)]$ . We identify the algebra  $\mathscr{R} \otimes \mathscr{R}$  with  $\mathbb{k}[\operatorname{Rep}(A,V) \times \operatorname{Rep}(A,V)]$ . We will also consider the associative R-algebra  $\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}$ . There are two algebra maps  $\operatorname{ev}_l, \operatorname{ev}_r : A \to \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}$ , given by the formulas  $\operatorname{ev}_l(a) := \widehat{a} \otimes 1$ , and  $\operatorname{ev}_r(a) := 1 \otimes \widehat{a}$ , respectively. Here,  $\widehat{a}$  is viewed either as an element of  $\mathscr{R} \otimes \operatorname{End}$  or as an element of  $\operatorname{End} \otimes \mathscr{R}$ . The two maps  $\operatorname{ev}_l, \operatorname{ev}_r$ , make  $\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}$  an A-bimodule, via  $a'(u \otimes F \otimes v)a'' := \operatorname{ev}_l(a') \cdot (u \otimes F \otimes v) \cdot \operatorname{ev}_r(a'')$ .

We define the double tangent  $\mathcal{R}$ -bimodule by

$$\mathscr{T}^e(A, V) := \operatorname{Der}_R(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}).$$
 (6.3.1)

We observe that the A-bimodule structure on  $\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}$  commutes with the obvious  $\mathscr{R}$ -bimodule structure, hence, the latter structure indeed makes  $\mathscr{T}^e(A,V)$  an  $\mathscr{R}$ -bimodule, that is, a  $\mathbb{K}[\operatorname{Rep}(A,V) \times \operatorname{Rep}(A,V)]$ -module (not necessarily free, in general).

Given  $\rho \in \operatorname{Rep}(A, V)$ , we let  $\mathbb{k}_{\rho}$  denote the 1-dimensional  $\mathscr{R}$ -module such that  $f \in \mathbb{k}[\operatorname{Rep}(A, V)]$  acts in  $\mathbb{k}_{\rho}$  via multiplication by the scalar  $f(\rho) \in \mathbb{k}$ . It is clear that the geometric fiber of the double tangent  $\mathscr{R}$ -bimodule  $\mathscr{T}^e(A, V)$  at a point  $(\rho, \varphi) \in \operatorname{Rep}(A, V) \times \operatorname{Rep}(A, V)$  is given by

$$\mathscr{T}^e(A,V)|_{(\rho,\varphi)}:=(\Bbbk_\rho\otimes \Bbbk_\varphi)\bigotimes_{\mathscr{R}\otimes\mathscr{R}}\mathscr{T}^e(A,V).$$

To get a better understanding of the vector space on the right of this formula, observe that any R-algebra homomorphism  $\rho:A\to \operatorname{End} V$  makes V a left A-module, to be denoted  $V_\rho$ . The action map  $A\otimes_R V_\rho\to V_\rho$  is surjective since A has a unit. This gives a surjective map of left A-modules  $A\otimes_R V \twoheadrightarrow V_\rho$ , where  $A\otimes_R V$  is regarded as a projective left A-module generated by the vector space V. We set  $K_\rho:=\operatorname{Ker}(A\otimes_R V\twoheadrightarrow V_\rho)$ , a left A-module.

**Lemma 6.3.2.** For any  $(\rho, \varphi) \in \text{Rep}(A, V) \times \text{Rep}(A, V)$ , one has a canonical isomorphism

$$\mathscr{T}^e(A,V)|_{(\rho,\varphi)} \simeq \operatorname{Hom}_A(K_\rho,V_\varphi).$$

Furthermore, this space is finite dimensional if A is finitely generated.

*Proof.* For any left A-module M, we have  $\operatorname{Hom}_A(A \otimes_R V, M) = \operatorname{Hom}_R(V_\rho, M)$  and  $\operatorname{Ext}_A^1(A \otimes_R V, M) = 0$ . Hence, the long exact sequence of Ext-groups arising from the short exact sequence  $K_\rho \hookrightarrow A \otimes_R V \twoheadrightarrow V_\rho$  reads:

$$\operatorname{Hom}_A(V_{\rho}, M) \hookrightarrow \operatorname{Hom}_R(V_{\rho}, M) \to \operatorname{Hom}_A(K_{\rho}, M) \to \operatorname{Ext}_A^1(V_{\rho}, M) \to 0.$$

Observe next that the  $\mathscr{R}$ -bimodule structure on  $\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}$  induces an  $\mathscr{R}$ -bimodule structure on each Hochschild cohomology group  $H_R^p(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R})$ ,  $p \geq 0$ . Note that  $(\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R})^R = \mathscr{R} \otimes \operatorname{End}^R \otimes \mathscr{R}$ , and  $H^0(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}) = (\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R})^A$ . In particular, we have the following exact sequence of  $\mathscr{R}$ -bimodules (to be compared with the displayed exact sequence above):

$$(\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R})^{A} \hookrightarrow (\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R})^{R} \xrightarrow{\operatorname{ad}} \operatorname{Der}_{R}(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}) \quad (6.3.3)$$
$$\longrightarrow H^{1}_{R}(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}) \to 0.$$

We leave to the reader to verify that, for any two representations  $\rho, \varphi \in \text{Rep}(A, V)$ , the geometric fibers at  $(\rho, \varphi)$  of the  $\mathscr{R}$ -bimodules occurring in (6.3.3) are given by

$$\begin{split} & \left(\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}\right)^{A} \big|_{(\rho,\varphi)} = \operatorname{Hom}_{A}(V_{\rho}, V_{\varphi}), \\ & \left(\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}\right)^{R} \big|_{(\rho,\varphi)} = \operatorname{Hom}_{R}(V_{\rho}, V_{\varphi}), \\ & \operatorname{Der}_{R}(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}) \big|_{(\rho,\varphi)} = \operatorname{Hom}_{A}(K_{\rho}, V_{\varphi}), \\ & \left. H^{1}_{R}(A, \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}) \right|_{(\rho,\varphi)} = \operatorname{Ext}^{1}_{A}(V_{\rho}, V_{\varphi}). \end{split}$$

The statement of the Lemma is now clear.

Next, we consider the following maps

$$A\otimes A \xrightarrow{\quad \text{ev}\otimes \text{ev} \quad} (\mathscr{R}\otimes \text{End})\otimes (\text{End}\otimes \mathscr{R}) \xrightarrow{\quad \text{id}_{\mathscr{R}}\otimes \mathbf{m}\otimes \text{id}_{\mathscr{R}}} \mathscr{R}\otimes \text{End}\otimes \mathscr{R},$$

where  $\mathbf{m} : \operatorname{End} \otimes \operatorname{End} \to \operatorname{End}$  denotes the multiplication in the R-algebra End. We observe that the two maps above are morphisms of A-bimodules, with respect to the outer bimodule structure on  $A \otimes A$ . Let  $\operatorname{ev}_{l,r} : A \otimes A \longrightarrow \mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}$  denote the composite map, which is an A-bimodule map again.

The "meaning" of the double derivation bimodule  $\mathbb{D}\mathbf{er}_R A$  is somewhat clarified by the following observation: The A-bimodule morphism  $\mathrm{ev}_{l,r}$  gives rise to a canonical map

$$\operatorname{ev}_{l,r}: \mathbb{D}\operatorname{er}_R A \longrightarrow \mathscr{T}^e(A,V) = \operatorname{Der}_R(A,\mathscr{R} \otimes \operatorname{End} \otimes \mathscr{R}).$$

**6.4** Moment maps. Let G be any algebraic group with Lie algebra  $\mathfrak{g}$ . Write  $\mathfrak{g}^*$  for the linear dual of  $\mathfrak{g}$ ; we often view  $x \in \mathfrak{g} = (\mathfrak{g}^*)^*$  as a linear function on  $\mathfrak{g}^*$ . The group G acts on  $\mathfrak{g}^*$  via the coadjoint action.

Let X be a G-scheme and  $\varpi \in \Omega^2(X)^G$  be a closed (not necessarily non-degenerate) G-invariant 2-form on X. Each element  $x \in \mathfrak{g}$  gives rise to a vector field,  $\xi_x$ , on X.

**Definition 6.4.1.** A G-equivariant map  $\mu: X \to \mathfrak{g}^*$  is said to be a moment map for  $(X, \varpi, G)$  if the pull-back morphism  $\mu^*: \mathbb{k}[\mathfrak{g}^*] \longrightarrow \mathbb{k}[X]$  has the following property:

$$d(\mu^*(x)) = i_{\xi_x} \varpi$$
, for any  $x \in \mathfrak{g}$ .

Remark 6.4.2. Moment map is not uniquely determined by the triple  $(X, \varpi, G)$ , in general.  $\Diamond$ 

Given a moment map  $\mu: X \to \mathfrak{g}^*$ , we write  $\mu^{-1}(0) \subset X$  for its scheme-theoretic zero fiber, a G-stable subscheme in X. If X is smooth and the 2-form  $\omega$  is nondegenerate, then  $\mathbb{k}[X]$  acquires a natural Poisson algebra structure. The latter is known to descend to a well-defined Poisson algebra structure on  $\mathbb{k}[\mu^{-1}(0)]^G$ .

Now, let A be a finitely generated associative R-algebra equipped with a closed 2-form  $\omega \in \mathrm{DR}^2_R A$ . Fix a finite dimensional R-module V. On  $\mathrm{Rep}(A,V)$ , we get a closed G-invariant 2-form  $\mathrm{Tr}\,\widehat{\omega} \in (\Omega^2\mathrm{Rep}(A,V))^G$ , cf. (6.2.2). We are interested in moment maps for the triple  $(\mathrm{Rep}(A,V),\mathrm{Tr}\,\widehat{\omega},G)$ .

Observe that the trace pairing  $(x,y) \mapsto \operatorname{Tr}(x \cdot y)$  on  $\mathfrak g$  provides a G-invariant isomorphism  $\mathfrak g^* \cong \mathfrak g$  and that the Lie algebra  $\mathfrak g = \operatorname{Hom}_R(V,V)$  may be viewed as a subspace in the associative algebra  $\operatorname{End} = \operatorname{Hom}_{\mathbb K}(V,V)$ . Thus, we have an imbedding  $\mathfrak g^* \hookrightarrow \operatorname{End}$  and any moment map  $\mu$  for  $(\operatorname{Rep}(A,V),\operatorname{Tr}\widehat{\omega},G)$  may be identified with a map  $\operatorname{Rep}(A,V) \longrightarrow \operatorname{End}$ , that is, with an element  $\mu \in (\mathbb k[\operatorname{Rep}(A,V)] \otimes \operatorname{End})^G$ .

Next, recall that we have defined a noncommutative moment map  $\mu_{\rm nc}$ :  $({\rm DR}_R^2A)_{\rm closed} \to (A/R)^R$ . Note that since  $R=\Bbbk I$  is commutative and semisimple, we have  $R\subset A^R$  and, moreover,  $(A/R)^R=A^R/R$ . Given a closed 2-form  $\omega\in{\rm DR}_R^2A$  and a representative  ${\bf w}\in A^R$  of the class  $\mu_{\rm nc}(\omega)\in A/R$ , we form the algebra  $A_{\bf w}=A/A{\bf w}A$ . The projection  $A\to A_{\bf w}$  gives a closed imbedding of schemes  ${\rm Rep}(A_{\bf w},V)\hookrightarrow {\rm Rep}(A,V)$ , hence, a natural restriction map  ${\mathbb k}[{\rm Rep}(A,V)]\to {\mathbb k}[{\rm Rep}(A_{\bf w},V)]$ .

Recall that if the form  $\omega$  is bi-symplectic then Propositions 4.4.1 and 4.4.3 provide the spaces  $A_{\natural}$  and  $(A_{\mathbf{w}})_{\natural}$  with compatible Lie brackets. These Lie brackets make the algebras  $\operatorname{Sym}(A_{\natural}), \operatorname{Sym}((A_{\mathbf{w}})_{\natural})$  into Poisson algebras, with  $\operatorname{Sym} R$  being a central subalgebra in each of them. Therefore, for any linear map  $\mathbf{d}: R \to \mathbb{k}$ , the corresponding Poisson algebra structures descend to the quotients  $\mathcal{O}_{\mathbf{d}}(A)$  and  $\mathcal{O}_{\mathbf{d}}(A_{\mathbf{w}})$ , respectively.

Our main result about representation functors is the following


**Theorem 6.4.3.** Fix a closed 2-form  $\omega \in DR_R^2 A$ .

(i) For any  $\mathbf{w} \in A^R$  in the preimage of the element  $\mu_{nc}(\omega) \in A^R/R$ , the element

$$\boldsymbol{\mu} := \widehat{\mathbf{w}} \in (\Bbbk[\operatorname{Rep}(A, V)] \otimes \operatorname{End})^G$$

gives a moment map for the triple  $(\operatorname{Rep}(A, V), \operatorname{Tr} \widehat{\omega}, G)$ .

Write  $\mu : \operatorname{Rep}(A, V) \to \mathfrak{g}^*$  for the corresponding map, and put  $\mathbf{d} = \mathbf{d}(V)$ . Then, we have  $\operatorname{Rep}(A_{\mathbf{w}}, V) = \mu^{-1}(0)$ , as subschemes in  $\operatorname{Rep}(A, V)$ ; thus, there is a commutative diagram



(ii) Assume in addition that A is smooth and  $\omega$  is a bi-symplectic form.

Then, the scheme  $\operatorname{Rep}(A,V)$  is smooth; furthermore, the 2-form  $\operatorname{Tr} \widehat{\omega}$  makes  $\operatorname{Rep}(A,V)$  a symplectic manifold, and all maps in the diagram above are Poisson algebra morphisms.

Remark 6.4.4. Different choices of elements **w** corresponding to the same class  $\mu_{\rm nc}(\omega) \in A^R/R$  give rise to different moment maps, in general.

Proof of Theorem 6.4.3. We set  $\widehat{A} := \mathbb{k}[\operatorname{Rep}(A, V)] \otimes \operatorname{End}$ , and view it as an R-algebra. The map  $a \mapsto \widehat{a}$  is an R-algebra morphism  $A \to \widehat{A}$ .

Given any  $x \in \mathfrak{g}$  we may (and will) view it as the element  $1 \otimes x \in \widehat{A}$ . In particular, for  $u \in \widehat{A}$ , we write  $x \cdot u := (1 \otimes x) \cdot u$ , resp.,  $u \cdot x := u \cdot (1 \otimes x)$ . Thus, associated to  $x \in \mathfrak{g}$ , we have an inner derivation ad  $x : \widehat{A} \to \widehat{A}$ ,  $u \mapsto \operatorname{ad} x(u) := (1 \otimes x) \cdot u - u \cdot (1 \otimes x)$ .

Further, differentiating the G-action on  $\operatorname{Rep}(A,V)$ , one associates to any element  $x \in \mathfrak{g}$  a vector field  $\xi_x$  on  $\operatorname{Rep}(A,V)$ , and we let  $\xi_x(f)$  denote the result of the  $\xi_x$ -action on a function  $f \in \mathbb{k}[\operatorname{Rep}(A,V)]$ . We will also consider  $\xi_x$ -action on elements of  $\widehat{A}$ ; for an End-valued function  $f \otimes y \in \mathbb{k}[\operatorname{Rep}(A,V)] \otimes \operatorname{End} = \widehat{A}$ , the action is defined by the formula  $\xi_x(f \otimes y) = \xi_x(f) \otimes y$ . This way, one gets a derivation  $\xi_x : \widehat{A} \to \widehat{A}$ .

Observe next that, for any  $a \in A$ , the function  $\widehat{a} : \operatorname{Rep}(A, V) \to \operatorname{End}$  is G-equivariant. Hence, in  $\widehat{A}$ , one has an equality

$$\xi_x \widehat{a} = \operatorname{ad} x(\widehat{a}), \quad \forall x \in \mathfrak{g}, \ a \in A.$$
 (6.4.5)

This implies a similar equation for differential forms. For instance, given a 2-form  $\omega = a_0 da_1 da_2 \in DR_R^2 A$ , we get

$$i_{\xi_x}\widehat{\omega} = i_{\xi_x}(\widehat{a}_0 \, d\widehat{a}_1 \, d\widehat{a}_2) = \widehat{a}_0 \xi_x(\widehat{a}_1) \, d\widehat{a}_2 - \widehat{a}_0 \, d\widehat{a}_1 \, \xi_x(\widehat{a}_2)$$

$$\stackrel{(6.4.5)}{=} \widehat{a}_0 \operatorname{ad} x(\widehat{a}_1) \, d\widehat{a}_2 - \widehat{a}_0 \, d\widehat{a}_1 \, \operatorname{ad} x(\widehat{a}_2) = i_{\operatorname{ad} x}(\widehat{a}_0 \, d\widehat{a}_1 \, d\widehat{a}_2) = i_{\operatorname{ad} x}\widehat{\omega}.$$

$$(6.4.6)$$

Now, with the notation introduced above, proving the theorem amounts to showing the identity:

$$i_{\mathcal{E}_x}(\operatorname{Tr}\widehat{\omega}) = d\operatorname{Tr}(x \cdot \widehat{\mathbf{w}}), \quad \forall x \in \mathfrak{g}.$$
 (6.4.7)

We have

$$i_{\xi_x}(\operatorname{Tr}\widehat{\omega}) = \operatorname{Tr}(i_{\xi_x}\widehat{\omega}) = \operatorname{Tr}(i_{\operatorname{ad} x}\widehat{\omega}),$$

by (6.4.6). Further, from Proposition 4.1.3 applied to the algebra  $\widehat{A}$  and to the 2-form  $\widehat{\omega}$  we deduce that, in  $\mathrm{DR}^1(\widehat{A})$ , one has

$$\operatorname{Tr}(i_{\operatorname{ad} x}\widehat{\omega}) = \operatorname{Tr}(x \cdot d\mu_{\operatorname{nc}}(\widehat{\omega})) = \operatorname{Tr}(x \cdot \widehat{d\mathbf{w}}) = \operatorname{Tr}(d(x \cdot \widehat{\mathbf{w}})) = d\operatorname{Tr}(x \cdot \widehat{\mathbf{w}}).$$

Hence, we obtain  $i_{\xi_x} \operatorname{Tr} \widehat{\omega} = \operatorname{Tr}(i_{\operatorname{ad} x} \widehat{\omega}) = d \operatorname{Tr}(x \cdot \widehat{\mathbf{w}})$ . This proves formula (6.4.7). We conclude that  $\widehat{\mathbf{w}}$  is indeed a moment map for the triple (Rep(A, V), Tr  $\widehat{\omega}$ , G).

Finally, surjectivity of the horizontal maps in the diagram of the Theorem follows from the well-known result due to Le Bruyn-Procesi [LBP]. Commutativity of the diagram is clear.

We now turn to part (ii) of the Theorem. First, recall that if B is any R-algebra then, for any left B-module Q, the space  $Q^* = \operatorname{Hom}_{\mathbb{K}}(Q, \mathbb{K})$  acquires a natural structure of right B-module, and vice versa. Further, given a right B-module P and a left B-module Q, one has a canonical linear map

$$\Psi: P \otimes_B Q \longrightarrow \operatorname{Hom}_B(P, Q^*)^*, \quad p \otimes q \longmapsto \Psi(p \otimes q),$$
where  $\Psi(p \otimes q): \operatorname{Hom}_B(P, Q^*) \longrightarrow \mathbb{k}, \quad f \mapsto \langle f(p), q \rangle.$  (6.4.8)

It is immediate to see that the map  $\Psi$  is a bijection if P is a finite rank free B-module and  $\dim_{\mathbb{R}} Q < \infty$ . We deduce that: For any finitely generated projective B-module P and finite dimensional module Q, the map  $\Psi$  becomes an isomorphism of (finite dimensional) vector spaces.

We now resume the proof of part (ii) of the Theorem. The implication A is  $smooth \Longrightarrow \operatorname{Rep}(A, V)$  is smooth is standard, cf. e.g. [Gi3].

Next, fix  $\rho \in \operatorname{Rep}(A, V)$ , view  $V = V_{\rho}$  as a left A-module and view  $\operatorname{End}_{\Bbbk} V_{\rho}$  as an A-bimodule. We put  $\operatorname{End}_{\rho} := \operatorname{End}_{\Bbbk} V_{\rho}$  and  $\operatorname{End}_{\rho}^* := (\operatorname{End}_{\Bbbk} V_{\rho})^*$ . The trace paring  $(u, v) \longmapsto \operatorname{Tr}(u \cdot v)$  gives an A-bimodule isomorphism  $tr : \operatorname{End}_{\rho}^* \stackrel{\sim}{\to} \operatorname{End}_{\rho}$ .

By a standard infinitesimal computation, the Zariski tangent space to the scheme  $\operatorname{Rep}(A,V)$  at the point  $\rho \in \operatorname{Rep}(A,V)$  equals

$$T_{\rho}\operatorname{Rep}(A, V) = \operatorname{Der}_{R}(A, \operatorname{End}_{\rho}) = \operatorname{Hom}_{A^{e}}(\Omega_{R}^{1}A, \operatorname{End}_{\rho}).$$

Therefore, for the cotangent space at  $\rho \in \text{Rep}(A, V)$ , we get  $T_{\rho}^* \text{Rep}(A, V) = \text{Der}_R(A, \text{End}_{\rho})^* = \text{Hom}_{A^c}(\Omega_R^1 A, \text{End}_{\rho})^*$ .

Further, let  $\omega \in DR_R^2 A$ . The 2-form  $\omega$  gives rise to the 'contraction map'  $i(\omega): \mathbb{D}\mathbf{er}_R A \longrightarrow \Omega_R^1 A$ , and the corresponding 2-form  $\operatorname{Tr}\widehat{\omega} \in \Omega^2(\operatorname{Rep}(A,V))$ , induces a similar contraction map  $\widehat{i} = i(\operatorname{Tr}\widehat{\omega}): T_{\rho}\operatorname{Rep}(A,V) \to T_{\rho}^*\operatorname{Rep}(A,V)$ .

It is straightforward to verify that the following diagram commutes:

where we have used simplified notation  $T_{\rho} = T_{\rho} \text{Rep}(A, V)$ ,  $\Omega^{1} = \Omega_{R}^{1} A$ , and  $\text{Der} = \text{Der}_{R}$ .

Assume now that A is smooth, so  $P := \Omega_R^1 A$  is a finitely generated projective  $A^e$ -module. Then, the map  $\Psi$  in (6.4.8) becomes an isomorphism, and the map (2.3.1) in the top row of the commutative diagram above also becomes an isomorphism, by Proposition 2.3.2. Further, if  $\omega$  is bi-nondegenerate, then the vertical map on the left of the diagram is a bijection. It follows that the vertical map on the right is a bijection as well. Thus, we have proved that the 2-form  $\operatorname{Tr}\widehat{\omega}$  is nondegenerate.

The last statement of the Theorem, compatibility with Poisson brackets, is the main result of [Gi].

# 7 Hamiltonian reduction in noncommutative geometry

Throughout this section, R stands for a finite dimensional semisimple k-algebra.

**7.1** Proof of Proposition 4.4.3. Let A be an R-algebra with a bi-symplectic form  $\omega \in DR_R^2 A$ , and  $\iota(\omega) : \mathbb{D}\mathbf{er}_R A \xrightarrow{\sim} \Omega_R^1 A$  the bijection induced by  $\omega$ . For any  $v \in A$ , let  $\Theta_v := \iota(\omega)^{-1}(dv)$  denote the corresponding double derivation and, using Sweedler's notation, write  $\Theta_v = \Theta_v' \otimes \Theta_v''$ .

With this understood, we have the following explicit formula for the map  $H_{\omega}: DR_R^1 A \to Der_R A$ , introduced in (4.2.1).

**Lemma 7.1.1.** Let  $u', u'', v \in A$ , and put  $\theta := \mathsf{H}_{\omega}(u' \cdot dv \cdot u'') \in \mathsf{Der}_R A$ . Then,

(i) We have  $\theta(q) = \Theta'_v(q) \cdot u'' \cdot u' \cdot \Theta''_v(q)$ ,  $\forall q \in A$ .

(ii) Let  $\mathbf{w} \in A$  be a representative of the class  $\mu_{nc}(\omega) \in (A/R)^R$ , let  $u = u'' \otimes u' \in (A \otimes A)^R$ , and put  $\theta := \mathsf{H}_{\omega}(u' \cdot d\mathbf{w} \cdot u'')$ . Then,

$$i(\omega)^{-1}(u' \cdot d\mathbf{w} \cdot u'') = u' \cdot \Delta \cdot u'' = \operatorname{ad} u, \quad and \quad \theta = \operatorname{ad}(u''u').$$

*Proof.* The map  $i := i(\omega)$  is an A-bimodule isomorphism, hence, so is  $i^{-1}$ . We deduce

$$\imath^{-1}(u'\cdot dv\cdot u'')=u'\cdot \imath^{-1}(dv)\cdot u''=u'\cdot \Theta_v\cdot u''=(\Theta_v'\cdot u'')\otimes (u'\cdot \Theta_v'').$$

Further, by definition one has  $\theta = \mathbf{m}_{\natural} \circ \imath^{-1}(u' \cdot dv \cdot u'')$ . Hence, we get  $\theta(q) = (\Theta'_v(q) \cdot u'') \cdot (u' \cdot \Theta''_v(q))$ , and (i) follows. Part (ii) is a reformulation of the first equation in Proposition 4.1.3(iii).

Next, we set  $I := A\mathbf{w}A \subset A$ , so  $A_{\mathbf{w}} = A/I$ .

**Lemma 7.1.2.** (i) For any  $p \in A_{\natural}$ , one has  $\theta_p(I) \subset I$ . Hence,  $\theta_p$  descends to a well-defined derivation  $\bar{\theta}_p : A/I \to A/I$ .

(ii) If  $p \in I + [A, A]$ , then  $\bar{\theta}_p : A/I \to A/I$  is an inner derivation.

*Proof.* To prove (i), we compute  $\theta_p(A\mathbf{w}A) \subset \theta_p(A)\mathbf{w} A + A \theta_p(\mathbf{w}) A + A \mathbf{w} \theta_p(A)$ . The summand in the middle vanishes since  $\theta_p(\mathbf{w}) = 0$ , by Lemma 4.2.6(ii). The other two summands clearly belong to  $I = A\mathbf{w}A$ , and (i) follows.

We turn to part (ii). It suffices to prove the result for elements of the form  $p = a\mathbf{w}b \in A\mathbf{w}A = I$ . For such an element, in  $\Omega_R^1 A$ , we have  $dp = d(a\mathbf{w}b) = (da)\mathbf{w}b + a(d\mathbf{w})b + a\mathbf{w}(db)$ . Hence, using the formulas and notation of Lemma 7.1.1, for  $\theta_p = \theta$  and any  $q \in A$ , we find

$$\theta_p(q) = \Theta'_a(q) \cdot \mathbf{w} \cdot b \cdot \Theta''_a(q) + \mathrm{ad}(ba)(q) + \Theta'_b \cdot a \cdot \mathbf{w} \cdot \Theta''_b(q).$$

In this sum, the first and third summands clearly belong to the ideal  $A\mathbf{w}A = I$ . Hence, these terms give no contribution to the induced map  $A/I \to A/I$ ,  $q \mapsto \bar{\theta}_p(q)$ . The summand in the middle gives an inner derivation  $\mathrm{ad}(ba)$ . We conclude that the induced derivation  $\bar{\theta}_p$  is an inner derivation  $A/I \to A/I$  corresponding to the image of ba in A/I.

Proof of Proposition 4.4.3. By part (i) of Lemma 7.1.2 we get a map  $\varphi: A \to \operatorname{Der}_R A_{\mathbf{w}}, a \longmapsto \bar{\theta}_a$ . This map vanishes on [A,A]. Furthermore, part (ii) of Lemma 7.1.2 implies that the composite map  $A \stackrel{\varphi}{\longrightarrow} \operatorname{Der}_R A_{\mathbf{w}} \to HH^1_R(A_{\mathbf{w}})$  vanishes on  $I = A\mathbf{w}A$ . Thus, we conclude that this composite vanishes on I + [A,A], hence, descends to a well-defined map

$$\bar{\varphi}: A_{\mathbf{w}}/[A_{\mathbf{w}}, A_{\mathbf{w}}] = A/(I + [A, A]) \longrightarrow HH^1_R(A_{\mathbf{w}}).$$
 (7.1.3)

We may now combine the map  $\bar{\varphi}$  with the natural action of  $HH_R^1(A_{\mathbf{w}})$  on  $(A_{\mathbf{w}})_{\natural}$ , see Sect. 4.5, to obtain a map

$$(A_{\mathbf{w}})_{\natural} \longrightarrow \operatorname{Hom}_{\Bbbk}((A_{\mathbf{w}})_{\natural}, (A_{\mathbf{w}})_{\natural}), \quad a \longmapsto \bar{\varphi}_{a}.$$

It is immediate from the construction that, writing  $\pi:A_{\natural} \to (A_{\mathbf{w}})_{\natural}$  for the natural projection, one has

$$\bar{\varphi}_{\pi(a)}(\pi(a')) = \pi(\theta_a(a')) = \pi(\{a, a'\}), \quad \forall a, a' \in A_{\natural}.$$

Therefore, skew-symmetry and Jacobi identity for the Poisson bracket on  $A_{\natural}$  imply similar properties for the bracket  $(A_{\mathbf{w}})_{\natural} \times (A_{\mathbf{w}})_{\natural} \to (A_{\mathbf{w}})_{\natural}, a \times a' \mapsto \bar{\varphi}_a(a')$ .

Finally, it is clear that the map  $\varphi: A_{\natural} \to \operatorname{Der}_R A_{\mathbf{w}}$ , considered above is a Lie algebra homomorphism. Therefore, we deduce that the map  $\bar{\varphi}: (A_{\mathbf{w}})_{\natural} \longrightarrow HH^1_R(A_{\mathbf{w}})$  in (7.1.3) is a Lie algebra homomorphism, cf. also Theorem 7.2.3(ii).

**7.2** Main result. Let  $DR_R^{\bullet}B$  be the Karoubi-de Rham complex of an R-algebra B. We define

$$K_R^{\bullet}B:=\mathrm{Ker}\left[\imath_{\Delta}:\mathrm{DR}_R^{\bullet}B\longrightarrow\Omega_R^{\bullet-1}B\right].$$

The graded subspace  $K_R^{\bullet}B \subset \operatorname{DR}_R^{\bullet}B$  is stable under d, as well as under the maps  $L_{\theta}, i_{\theta}$ , for any  $\theta \in \operatorname{Der}_R B$ , since  $i_{\Delta}$  either commutes or anti-commutes with these maps.

For any inner derivation ad  $a, a \in B^R$ , by Corollary 3.2.3 we have  $i_{\operatorname{ad} a}\omega = a \cdot i_{\Delta}\omega$ . Hence, we deduce  $i_{\operatorname{ad} a}\omega = 0$ , for any  $\omega \in K_{\mathbb{R}}^{\bullet}B$ . In this case, we

also have  $L_{\operatorname{ad} a}\omega = d \circ i_{\operatorname{ad} a}\omega + i_{\operatorname{ad} a} \circ d\omega = 0$ . We see that, when restricted to  $K_R^{\bullet}B$ , the operations  $i_{\theta}, L_{\theta}$  depend only on the cohomology class of  $\theta$  in  $HH_R^1(B) = \operatorname{Der}_R B / \operatorname{Inn}_R B$ . Thus, the complex  $K_R^{\bullet}B$  acquires a natural structure of  $HH_R^1(B)$ -equivariant complex in the sense of Sect. 4.3.

Remark 7.2.1. According to [Gi2], there is a natural graded space isomorphism  $K_R^{\bullet}B \cong HH_{\bullet}^RB$ . The above defined action of  $HH_R^1(B)$  on  $K_R^{\bullet}B$  then becomes part of the standard action of Hochschild cohomology on Hochschild homology mentioned in Sect. 4.5. This homological interpretation is not quite convenient, however, for performing Hamiltonian reduction. Thus, in the present paper, we will neither use nor prove the isomorphism  $K_R^{\bullet}B \cong HH_{\bullet}^RB$ .

Note that since  $DR_R^{-1}B = 0$ , for any  $\omega \in DR_R^0B = B_{\natural}$ , we have  $i_{\Delta}\omega = 0$ . Thus,  $K_R^0B = B_{\natural}$ , and the  $HH_R^1(B)$ -action on  $B_{\natural}$  is the action mentioned in Sect. 4.5, that has been already used in the proof of Proposition 4.4.3.

Now, let A be an R-algebra,  $\omega \in \mathrm{DR}^2_R A$  a closed 2-form,  $\mathbf{w} \in A$  a representative of the class  $\mu_{\mathrm{nc}}(\omega) \in A/R$ . Below, we will consider the complex  $K^{\bullet}_R B$  for the algebra  $B = A_{\mathbf{w}} := A/A\mathbf{w}A$ , the Hamiltonian reduction of A at  $\mathbf{w}$ . Let  $\omega_{\mathbf{w}} \in \mathrm{DR}^2_R A_{\mathbf{w}}$  denote the image of  $\omega$  under the projection  $\mathrm{DR}^2_R A \twoheadrightarrow \mathrm{DR}^2_R A_{\mathbf{w}}$ .

**Lemma 7.2.2.** For any  $a \in A_{\mathbf{w}}$  and  $\theta \in \operatorname{Der}_R A_{\mathbf{w}}$ , we have da,  $i_{\theta}\omega_{\mathbf{w}} \in K_R^1 A_{\mathbf{w}}$ . Furthermore, the map  $\theta \mapsto i_{\theta}\omega_{\mathbf{w}}$  kills inner derivations, hence, descends to a well-defined map  $\mathbf{i} : HH_R^1(A_{\mathbf{w}}) \to K_R^1(A_{\mathbf{w}})$ .

Proof. It is immediate that  $i_{\Delta}(da) = 0$ , hence  $da \in K_R^1 A_{\mathbf{w}}$ . It is also clear that we have  $i_{\Delta}\omega_{\mathbf{w}} = 0$  since  $\mathbf{w} = 0$  in  $A_{\mathbf{w}}$ . Therefore, for any derivation  $\theta \in \operatorname{Der}_R A_{\mathbf{w}}$ , we find  $i_{\Delta}(i_{\theta}\omega_{\mathbf{w}}) = -i_{\theta}(i_{\Delta}\omega_{\mathbf{w}}) = 0$ . Thus, we have proved that  $i_{\theta}\omega_{\mathbf{w}} \in K_R^1 A_{\mathbf{w}}$ . Further, for  $a \in A_{\mathbf{w}}$ , by Proposition 4.1.3(iii) we get  $i_{\operatorname{ad}} a\omega_{\mathbf{w}} = a \cdot d\mathbf{w}$ , which is equal to zero in  $\operatorname{DR}_R^1 A_{\mathbf{w}}$ .

Assume now that  $\omega$  is a bi-symplectic 2-form. Then we have the Lie bracket on  $A_{\natural}$  provided by Proposition 4.4.1 and we consider the projection  $A_{\natural} \twoheadrightarrow (A_{\mathbf{w}})_{\natural}$ . The following Theorem, which is a strengthening of Proposition 4.4.3, is our

main result about Hamiltonian reduction in noncommutative geometry.

**Theorem 7.2.3.** (i) The bi-symplectic form  $\omega$  gives rise, canonically, to a Hamilton operator  $\mathsf{H}_{\mathbf{w}}: K^1_R(A_{\mathbf{w}}) \to HH^1_R(A_{\mathbf{w}})$  such that  $i_{\mathsf{H}_{\mathbf{w}}(\alpha)}\omega = \alpha$ , for any  $\alpha \in K^1_R(A_{\mathbf{w}})$ , cf. (4.2.2).

- (ii) With the Lie bracket on  $(A_{\mathbf{w}})_{\natural}$  induced by the Hamilton operator  $\mathsf{H}_{\mathbf{w}}$  from (i), the natural projection  $A_{\natural} \twoheadrightarrow (A_{\mathbf{w}})_{\natural}$  becomes a Lie algebra morphism.
- (iii) If the algebra A is smooth then the maps  $\mathbf i$  and  $H_{\mathbf w}$  are mutually inverse bijections.

From part (i) we deduce

Corollary 7.2.4. The map  $H_{\mathbf{w}}$  is injective and the map  $\mathbf{i}$  is surjective.

**7.3 Proof of Theorem 7.2.3.** First of all, Lemma 7.1.1(ii) yields the following result.

**Lemma 7.3.1.** Let  $\omega \in DR_R^2 A$  be a bi-symplectic form on A. Then, the bijection  $i: \mathbb{D}\mathbf{er}_R A \longrightarrow \Omega_R^1 A$  induced by  $\omega$  restricts to an A-bimodule isomorphism  $i: \operatorname{Inn}_R(A, A \otimes A) \xrightarrow{\sim} A(d\mathbf{w}) A$ .

Next, in the setup of Theorem 7.2.3, we introduce the two-sided ideal  $I := A\mathbf{w}A \subset A$ , and put

$$\operatorname{Der}_{R}(A, I) := \{ \theta \in \operatorname{Der}_{R} A \mid \theta(I) \subset I \} = \{ \theta \in \operatorname{Der}_{R} A \mid \theta(\mathbf{w}) \in I \}. \tag{7.3.2}$$

Further, consider the canonical surjection  $\varrho: \operatorname{DR}_R^{\bullet}A \twoheadrightarrow \operatorname{DR}_R^{\bullet}(A_{\mathbf{w}})$ , cf. (2.5.3), and put  $\operatorname{DR}_R^{\bullet}(A,I) := \varrho^{-1}(K_R^{\bullet}(A_{\mathbf{w}}))$ . Clearly,  $\operatorname{DR}_R^{\bullet}(A,I)$  is a subcomplex of the de Rham complex  $\operatorname{DR}_R^{\bullet}A$ .

**Lemma 7.3.3.** For the map  $H_{\omega}: DR_R^1A \to Der_RA$  in (4.2.1) and any  $\alpha \in DR_R^1A$ , we have  $\alpha \in DR_R^1(A, I)$  if and only if  $H_{\omega}(\alpha) \in Der_R(A, I)$ .

In particular, the map  $\mathsf{H}_{\omega}$  restricts to an imbedding  $\mathrm{DR}^1_R(A,I) \hookrightarrow \mathrm{Der}_R(A,I)$ .

*Proof.* The map  $\varrho$  used in the definition of  $\mathrm{DR}^1_R(A,I)$  is induced by the natural DG algebra projection  $\varrho_{\Omega}: \Omega_R^{\bullet}A \twoheadrightarrow \Omega_R^{\bullet}(A_{\mathbf{w}}) = \Omega_R^{\bullet}(A/I)$ . Furthermore, for any  $k=1,2,\ldots$ , we have a commutative diagram

$$\begin{array}{ccc}
\operatorname{DR}_{R}^{k} A & \xrightarrow{\imath_{\Delta}} & \Omega_{R}^{k-1} A \\
\downarrow^{\varrho} & & \downarrow^{\varrho_{\Omega}} \\
\operatorname{DR}_{R}^{k} (A/I) & \xrightarrow{\imath_{\Delta}} & \Omega_{R}^{k-1} (A/I)
\end{array} (7.3.4)$$

For k = 1, from the commutativity of the diagram, we deduce

$$\mathrm{DR}_{R}^{1}(A,I) = \varrho^{-1}\left(K_{R}^{1}(A_{\mathbf{w}})\right) = \{\alpha \in \mathrm{DR}_{R}^{1}A \mid \imath_{\Delta}\alpha \in I\}. \tag{7.3.5}$$

Now, let  $\alpha \in DR_R^1 A$  and let  $\theta = \mathsf{H}_{\omega}(\alpha) \in Der_R A$  be the derivation corresponding to  $\alpha$  via the bi-symplectic form. Thus,  $\alpha = i_{\theta}\omega$ , by (4.2.2). Further, by Proposition 4.2.4(i) we have  $\theta(\mathbf{w}) = -i_{\Delta}\alpha$ . The proof of the Lemma is now completed by the following equivalences

$$\alpha \in \mathrm{DR}^1_R(A, I) \iff \imath_\Delta \alpha \in I \iff \theta(\mathbf{w}) \in I \iff \theta \in \mathrm{Der}_R(A, I).$$

Any derivation  $\theta \in \operatorname{Der}_R(A, I)$  induces a well-defined derivation  $\bar{\theta} : A_{\mathbf{w}} \to A_{\mathbf{w}}$ . It is clear that  $\operatorname{Der}_R(A, I)$  is a Lie subalgebra in  $\operatorname{Der}_R A$  and the assignment  $\theta \mapsto \bar{\theta}$  gives a well-defined Lie algebra morphism  $\operatorname{Der}_R(A, I) \to \operatorname{Der}_R A_{\mathbf{w}}$ . We form the following composite

$$\Phi: \operatorname{Der}_R(A,I) \longrightarrow \operatorname{Der}_R A_{\mathbf{w}} \twoheadrightarrow \operatorname{Der}_R A_{\mathbf{w}} / \operatorname{Inn}_R A_{\mathbf{w}} = HH^1_R(A_{\mathbf{w}}),$$

where the second map is the natural projection. Also, let  $\Psi$  be the restriction of the map  $\varrho$ , cf. (7.3.4), to the subspace  $DR_R^1(A, I) \subset DR_R^1A$ .

According to Lemma 7.3.3 we obtain a diagram

$$K_R^1(A_{\mathbf{w}}) \xrightarrow{\Psi} \mathrm{DR}_R^1(A,I) \xrightarrow{\mathsf{H}_{\omega}} \mathrm{Der}_R(A,I) \xrightarrow{\Phi} HH_R^1(A_{\mathbf{w}}).$$
 (7.3.6)

**Lemma 7.3.7.** For  $\alpha \in DR_R^1(A, I)$ , we have  $\Psi(\alpha) = 0 \iff \Phi \circ H_\omega(\alpha) = 0$ .

*Proof.* We need to introduce some auxiliary objects. Let

$$\mathbf{I} := \Omega_R^{\bullet} A \cdot (I + dI) \cdot \Omega_R^{\bullet} A \subset \Omega_R^{\bullet} A$$

be the two-sided graded ideal in the algebra  $\Omega_R^{\bullet}A$  generated by the spaces  $I\subset\Omega_R^0A$  and  $dI\subset\Omega_R^1A$ . Note that

$$\mathbf{I} \cap \Omega_R^0 A = I$$
, and  $\mathbf{I} \cap \Omega_R^1 A = A(d\mathbf{w})A + I \cdot dA \cdot A + A \cdot dA \cdot I$ . (7.3.8)

Clearly, **I** is the smallest *d*-stable two-sided ideal in  $\Omega_R^{\bullet}A$  that contains *I*. Further, formula (2.5.3) shows that, in the notation of diagram (7.3.4), we have  $\mathbf{I} = \operatorname{Ker} \varrho_{\Omega}$ . Thus, from the commutativity of the diagram, as in the proof of the previous lemma, we deduce

$$DR_R^{\bullet}(A, I) = \varrho^{-1}(K_R^{\bullet}(A_{\mathbf{w}})) = \{\alpha \in DR_R^{\bullet}A \mid \iota_{\Delta}\alpha \in \mathbf{I}\}.$$
 (7.3.9)

Given a vector subspace  $E \subset \Omega_R^{\bullet} A$ , let  $\llbracket E \rrbracket$  denote the image of E under the projection  $\Omega_R^{\bullet} A \twoheadrightarrow \mathrm{DR}_R^{\bullet} A$ . With this notation, we obtain

$$K_R^{\bullet}(A_{\mathbf{w}}) \cong \mathrm{DR}_R^{\bullet}(A, I) / \llbracket \mathbf{I} \rrbracket = \{ \alpha \in \mathrm{DR}_R^{\bullet} A \mid \imath_{\Delta} \alpha \in \mathbf{I} \} / \llbracket \mathbf{I} \rrbracket.$$
 (7.3.10)

In particular, we see that  $\llbracket \mathbf{I} \rrbracket \subset \mathrm{DR}_R^{\bullet}(A,I)$ , in other words, one has  $i_{\Delta}(\llbracket \mathbf{I} \rrbracket) \subset \mathbf{I}$ . We can now turn to the proof of the implication ' $\Rightarrow$ ' of the Lemma. The isomorphism in (7.3.10) and formula (7.3.8) show that

$$\operatorname{Ker} \Psi = \llbracket \mathbf{I} \cap \Omega_R^1 A \rrbracket = \llbracket A(d\mathbf{w})A + I \cdot dA \cdot A + A \cdot dA \cdot I \rrbracket = \llbracket I \cdot dA + A \cdot d\mathbf{w} \rrbracket. \tag{7.3.11}$$

Thus, it suffices to show that if  $\alpha \in [I \cdot dA + A \cdot d\mathbf{w}]$ , then  $\Phi \circ \mathsf{H}_{\omega}(\alpha) = 0$ .

Assume first that  $\alpha \in A \cdot d\mathbf{w}$ . Then, Lemma 7.3.1 implies that  $\mathsf{H}_{\omega}(\alpha)$  is an inner derivation. It follows that the induced map  $A/I \to A/I$  is also an inner derivation, hence  $\Phi \circ \mathsf{H}_{\omega}(\alpha) = 0$ .

Next let  $\alpha = u \, dv \in I \cdot dA$  and put  $\theta := \mathsf{H}_{\omega}(\alpha)$ . The explicit formula for  $\theta = \mathsf{H}_{\omega}(\alpha)$  provided by Lemma 7.1.1(i) says that, for any  $a \in A$ , we have  $\theta(a) = \Theta'_v(a) \cdot u \cdot \Theta''_v(a)$ . We see that  $u \in I$  implies  $\theta(a) \in I$ , for any  $a \in A$ . Hence, for  $\alpha \in I \cdot dA$ , the corresponding derivation  $\theta$  induces the zero map  $\bar{\theta} : A/I \to A/I$ , and we are done.

We prove the implication ' $\Leftarrow$ '. Let  $\theta \in \operatorname{Der}_R(A, I)$  be such that the induced derivation  $\bar{\theta}: A_{\mathbf{w}} \to A_{\mathbf{w}}$  gives the zero class in  $HH^1_R(A_{\mathbf{w}})$ . This means that  $\theta = \xi + \operatorname{ad} a$ , where  $a \in A^R$  and  $\xi \in \operatorname{Der}_R(A, I)$  is such that  $\xi(A) \subset I$ . (Note that for any  $a \in A$ , we have  $\operatorname{ad} a(I) = [a, I] \subset I$ , hence  $\operatorname{Inn}_R A \subset \operatorname{Der}_R(A, I)$ .)

Let  $\alpha = i_{\theta}\omega \in \mathrm{DR}^1_R A$  be the 1-form that corresponds to  $\theta$ . We can write  $\alpha = i_{\xi}\omega + i_{\mathrm{ad}\,a}\omega$ . By Lemma 7.3.1, we get  $i_{\mathrm{ad}\,a}\omega \in [\![A(d\mathbf{w})A]\!]$ . Since  $[\![A(d\mathbf{w})A]\!] \subset \mathrm{Ker}\,\Psi$ , by (7.3.11), we conclude that  $i_{\mathrm{ad}\,a}\omega \in \mathrm{Ker}\,\Psi$ .

Next, given any 2-form  $\beta = x\,dy\,dz$ , we deduce from the inclusion  $\xi(A) \subset I$  that in  $\mathrm{DR}^1_RA$  one has  $i_\xi(x\,dy\,dz) = x\,\xi(y)\,dz - x\,dy\,\xi(z) \in I\cdot dA$ . Therefore, we obtain  $i_\xi\omega \in \llbracket I\cdot dA \rrbracket \subset \mathrm{Ker}\,\Psi$ . Thus, we have  $\alpha = i_\xi\omega + i_{\mathrm{ad}\,a}\omega \in \mathrm{Ker}\,\Psi$ , and the Lemma is proved.

Proof of Theorem 7.2.3. Lemma 7.3.7 implies that the map  $\mathsf{H}_{\omega}: \mathrm{DR}^1_R(A,I) \to \mathrm{Der}_R(A,I)$  sends the subspace  $\mathrm{Ker}\,\Psi \subset \mathrm{DR}^1_R(A,I)$  into the subspace  $\mathrm{Ker}\,\Phi \subset \mathrm{Der}_R(A,I)$ . We see that this map descends to a well-defined and injective map, see diagram (7.3.6):

$$K_R^1(A_{\mathbf{w}}) = \operatorname{DR}_R^1(A, I) / \operatorname{Ker} \Psi \xrightarrow{\mathsf{H}_{\omega}} \operatorname{Der}_R(A, I) / \operatorname{Ker} \Phi \cong \operatorname{Im} \Phi.$$
 (7.3.12)

Composing the resulting map with the imbedding Im  $\Phi \hookrightarrow HH^1_R(A_{\mathbf{w}})$ , we thus obtain an injective map  $\mathsf{H}_{\mathbf{w}}: K^1_R(A_{\mathbf{w}}) \hookrightarrow HH^1_R(A_{\mathbf{w}})$ .

The map  $H_{\mathbf{w}}$  satisfies all the identities required for a Hamilton operator. To check these identities, one chooses representatives in  $\mathrm{DR}^1_R(A,I)$  for classes in  $K^1_R(A_{\mathbf{w}})$  and verifies the corresponding identities for those representatives. The latter satisfy the required identities due to the fact that the map  $H_{\omega}:\mathrm{DR}^1_RA\to\mathrm{Der}_RA$  in (4.2.1) is known to be a Hamilton operator. This completes the proof of parts (i)-(ii) of the Theorem.

Assume now that the algebra A is smooth. Then, the map  $\mathbf{m}_{\natural} : (\mathbb{D}\mathbf{er}_R A)_{\natural} \to \mathrm{Der}_R A$  is a bijection, by Proposition 2.3.2. It follows that the map  $\mathsf{H}_{\omega}$  in (4.2.1) is also a bijection. Now, Lemma 7.3.7 shows that the latter bijection induces a bijection  $\mathsf{H}_{\omega} : \mathrm{DR}^1_R(A,I) \xrightarrow{\sim} \mathrm{Der}_R(A,I)$ . We see that, for the map  $\mathsf{H}_{\mathbf{w}} : K^1_R(A_{\mathbf{w}}) \hookrightarrow HH^1_R(A_{\mathbf{w}})$  defined above, we have  $\mathrm{Im}(\mathsf{H}_{\mathbf{w}}) = \mathrm{Im}(\Phi)$ . Thus, we are reduced to proving surjectivity of the map  $\Phi$ , which is equivalent to surjectivity of the map  $\mathrm{Der}_R(A,I) \to \mathrm{Der}_R A_{\mathbf{w}}, \theta \longmapsto \bar{\theta}$ .

To prove this last statement, we let  $f: A \twoheadrightarrow A/I = A_{\mathbf{w}}$  denote the projection and consider the following diagram

$$\operatorname{Der}_{R}(A/I, A/I) \xrightarrow{f^{*}} \operatorname{Der}_{R}(A, A/I) \xleftarrow{f_{*}} \operatorname{Der}_{R}(A, A)$$

For a smooth algebra A the bimodule  $\Omega_R^1A$  is projective. Hence, the functor  $\operatorname{Der}_R(A,-)=\operatorname{Hom}_{A^e}(\Omega_R^1A,-)$  is exact. We deduce that the map  $f_*$  in the diagram above is surjective. Hence, for any  $\bar{\theta}\in\operatorname{Der}_R(A/I,A/I)$  there exists an R-linear derivation  $\theta:A\to A$  such that  $f^*\bar{\theta}=f_*\theta$ . From this equation we get  $f(\theta(I))=(f^*\bar{\theta})(I)=0$ . Therefore,  $\theta\in\operatorname{Der}_R(A,I)$  and, furthermore, the map  $\operatorname{Der}_R(A,I)\to\operatorname{Der}_RA_{\mathbf{w}}$  sends  $\theta\longmapsto\bar{\theta}$ . This completes the proof of surjectivity, hence, the proof of the Theorem.

**7.4** Proof of Proposition 5.1.5. Set  $\Pi := \Pi^0(B)$ . This is a graded algebra with degree zero component being equal to B.

To prove the injectivity statement of part (ii) of the Proposition, we use the homotopy invariance of de Rham cohomology, see Lemma 2.5.2. We deduce that  $(DR_R^0\Pi)_{closed} = (DR_R^0B)_{closed} = R$ , where the last equality holds by assumption.

Next, write  $\omega_{\mathbf{w}}$  for the image of  $\omega$  in  $\mathrm{DR}^2_R\Pi$ . Since  $T^*B$  is a smooth algebra, Theorem 7.2.3 implies that the map  $\mathbf{i}: HH^1_R(\Pi) \to K^1_R(\Pi), \bar{\theta} \mapsto i_{\bar{\theta}}\omega_{\mathbf{w}}$  is a bijection. The bijection  $\mathbf{i}$  clearly sends the derivation  $\bar{\theta}_p$  to the 1-form  $dp \in \mathrm{DR}^1(\Pi)$ . The latter is zero if and only if  $p \in R$ , since  $(\mathrm{DR}^0_R\Pi)_{\mathrm{closed}} = R$ . Thus, we have proved that  $\bar{\theta}_p$  is a nonzero element in  $HH^1_R(\Pi)$ , for any nonzero element  $p \in \Pi_{\mathbb{b}}/R$ .

Further, by Proposition 5.4.1(ii), we have  $\mathbf{i}(\mathsf{E}) = i_{\mathsf{E}}\omega_{\mathbf{w}} = \lambda_{\mathbf{w}}$ , the image in  $\mathrm{DR}^1_R A_{\mathbf{w}}$  of the Liouville 1-form  $\lambda$ . The form  $\lambda_{\mathbf{w}}$  is not exact. Indeed, we have  $d\lambda_{\mathbf{w}} = \omega_{\mathbf{w}}$ , and the latter form is nonzero since the map  $\bar{\theta} \mapsto i_{\bar{\theta}}\omega_{\mathbf{w}}$  is a bijection.

Thus, any nonzero element of  $\mathbb{k} \cdot \mathsf{E}_{\mathrm{Lie}} \ltimes (\Pi_{\natural}/R)$  goes to a class  $u \in HH^1_R(\Pi)$  such that  $\mathbf{i}(u) \neq 0$ . Hence,  $u \neq 0$ , and injectivity of the map in part (ii) of Proposition 5.1.5 follows.

# 8 The necklace Lie algebra

**8.1** Let Q be a quiver with vertex set I, and write  $\overline{Q}$  for the double of Q. Let  $\mathbf{P} := \mathbb{k} \overline{Q}$  be the path algebra of  $\overline{Q}$ . The subalgebra  $R \subset \mathbf{P}$  formed by trivial paths may be (and will be) identified with  $\mathbb{k} I$ .

Given an edge  $a \in \overline{Q}$  we write  $a^*$  for the corresponding reverse edge. We set  $\omega := \sum_{a \in Q} da \, da^* \in \mathrm{DR}^2_R \mathbf{P}$ . It is clear that  $d\omega = 0$ .

The following result allows to apply the general machinery developed in the previous sections to algebras associated with quivers.

**Proposition 8.1.1.** The path algebra **P** is smooth. Furthermore, we have

(i) The assumptions of Proposition 4.1.4 hold for  $A={\bf P},$  more precisely, one has

$$H^{k}(\mathrm{DR}_{R}^{\bullet}\mathbf{P}) = \begin{cases} R & \text{if } k = 0\\ 0 & \text{if } k > 0. \end{cases}$$

$$(8.1.2)$$

- (ii) The 2-form  $\omega$  is bi-symplectic.
- (iii) The following element gives a representative of the class  $\mu_{\rm nc}(\omega) \in \mathbf{P}/R$ :

$$\mathbf{w} := \sum_{a \in Q} (a \cdot a^* - a^* \cdot a) = \sum_{a \in Q} [a, a^*] \in [\mathbf{P}, \mathbf{P}]^R,$$

more precisely, we have  $\mathbf{w} = \widetilde{\mu_{\rm nc}}(\omega)$ .

It is convenient to introduce a function  $\epsilon : \overline{Q} \to \{\pm 1\}$  by setting  $\epsilon(a)$  to be 1 if  $a \in Q$  and -1 if  $a \in \overline{Q} \setminus Q$ . Then, we may write  $\mathbf{w} = \sum_{a \in \overline{Q}} \epsilon(a) \cdot aa^*$ .

Proof of Proposition 8.1.1. For any two vertices  $i, j \in I$ , let  $E_{ij}$  be a vector space with basis  $\{a\}_{a:i\to j}$ , formed by the set of edges (from i to j) of the quiver  $\overline{Q}$ . The assignment sending a pair of edges  $a, b \in \overline{Q}$  to  $\epsilon(a)$  if  $b = a^*$  and to zero

otherwise extends to a non-degenerate pairing  $E_{ij} \times E_{ji} \to \mathbb{k}$ . In particular, the pairing provides, for any  $i \in I$ , the space  $E_{ii}$  with a non-degenerate skew-symmetric bilinear form, and also gives rise to vector space isomorphisms  $E_{ji} \cong E_{ij}^*$ , for any pair  $i, j \in I$ ,  $i \neq j$ . We set  $E := \bigoplus_{i,j \in I} E_{ij}$  and extend the pairings above to a symplectic form  $\langle -, - \rangle : E \times E \to \mathbb{k}$ .

The space E has a natural R-bimodule structure, and the symplectic form  $\langle -, - \rangle$  yields an R-bimodule isomorphism  $S: E^* \xrightarrow{\sim} E, \langle -, a \rangle \longmapsto a^*$ , where  $E^* = \operatorname{Hom}_{\mathbb{K}}(E, \mathbb{K})$ , the  $\mathbb{K}$ -linear dual of E, is equipped with a natural R-bimodule structure.

We have  $\mathbf{P} = T_R E$ , the tensor algebra of an R-bimodule. This implies, in particular, that the algebra  $\mathbf{P}$  is smooth, as claimed in Proposition 8.1.1.

To prove Proposition 8.1.1(ii), we consider the following isomorphisms of  $\mathbf{P}$ -bimodules

$$\operatorname{Der}_{R}(\mathbf{P}, \mathbf{P} \otimes \mathbf{P}) \xrightarrow{\sim}_{F} \mathbf{P} \otimes_{R} E^{*} \otimes_{R} \mathbf{P} \xrightarrow{\operatorname{id} \otimes_{S} \otimes_{\operatorname{Id}}} \mathbf{P} \otimes_{R} E \otimes_{R} \mathbf{P} \xrightarrow{\sim}_{G} \Omega_{R}^{1}(\mathbf{P}), \quad (8.1.3)$$

where the maps F, G are given, respectively, by

$$\Theta \overset{F}{\longmapsto} \sum_{a \in \overline{Q}} (\Theta'' a \otimes \langle a, - \rangle \otimes \Theta' a), \quad \text{resp.}, \quad \sum_{a \in \overline{Q}} f_a \otimes a \otimes g_a \overset{G}{\longmapsto} \sum_{a \in \overline{Q}} f_a \, da \, g_a.$$

It is straightforward to verify that the composite bijection in (8.1.3) is nothing but the map  $\Theta \mapsto \iota_{\Theta} \omega$ . We conclude that  $\omega$  is a bi-symplectic 2-form on **P**.

Since **P** is smooth, we deduce that the map  $i : \operatorname{Der}_R \mathbf{P} \to \operatorname{DR}^1_R \mathbf{P}$ ,  $\theta \mapsto i_\theta \omega$ , is a bijection as well. This can also be seen directly from the following diagram

$$\operatorname{Der}_{R} \mathbf{P} \xrightarrow{\stackrel{\sim}{\longrightarrow}} \mathbf{P} \otimes_{R^{e}} E^{*} \xrightarrow{\stackrel{\sim}{\longrightarrow}} \mathbf{P} \otimes_{R^{e}} E \xrightarrow{\stackrel{\sim}{\longrightarrow}} \operatorname{DR}_{R}^{1} \mathbf{P}, \quad (8.1.4)$$

where the maps  $F_{\natural}$  and  $G_{\natural}$  are given by

$$F_{\natural}: \ \theta \longmapsto \sum_{a \in \overline{Q}} \theta(a) \otimes \langle a, - \rangle, \quad G_{\natural}: \ \sum_{a \in \overline{Q}} f_a \otimes a \longmapsto \sum_{a \in \overline{Q}} f_a \, da$$

In the formulas above, we view  $\mathbf{P}$  as a right  $R^e$ -module, hence we have  $\mathbf{P} \otimes_{R^e} E = (\mathbf{P} \otimes_R E)/[R, \mathbf{P} \otimes_R E]$ . The inverse isomorphism in (8.1.4) sends a 1-form  $\sum f_a da \in \mathrm{DR}^1_R \mathbf{P}$  to the derivation  $\theta$  such that  $\theta(a) = f_{a^*}$ .

The path algebra comes equipped with a natural grading  $\mathbf{P} = \bigoplus_{k \geq 0} \mathbf{P}_k$ , by length of the path. Using Lemma 2.5.2 on homotopy invariance of de Rham cohomology, we obtain formula (8.1.2). Observe further that  $\mathbf{P}_0 = R$  and  $[\mathbf{P}, \mathbf{P}] \subset \bigoplus_{k \geq 1} \mathbf{P}_k$ . Therefore, we have  $[\mathbf{P}, \mathbf{P}] \cap R = 0$ , hence the sequence (4.1.5) in Proposition 4.1.4 is exact for  $A = \mathbf{P}$ . Thus, Proposition 4.1.4 holds for  $A = \mathbf{P}$ . This completes the proof of part (i) of Proposition 8.1.1.

Part (iii) is verified by a straightforward computation based on the last formula in Proposition 4.1.3(iii).

**8.2** Preprojective algebras. For any  $c \in \mathbb{k}I$ , one clearly has  $(\mathbf{w}-c) \mod R = \mu_{\rm nc}(\omega)$ . Write  $\mathbf{P}(\mathbf{w}-c)\mathbf{P}$  for the two-sided ideal in  $\mathbf{P}$  generated by  $\mathbf{w}-c$ . The corresponding Hamiltonian reduction  $\mathbf{P}/\mathbf{P}(\mathbf{w}-c)\mathbf{P}$  is known as deformed preprojective algebra,  $\Pi^c(Q) := \mathbf{P}/\mathbf{P}(\mathbf{w}-c)\mathbf{P}$ .

Remark 8.2.1. Let  $B = \mathbb{k}Q$  be the path algebra of the quiver Q (not of  $\overline{Q}$ ). It is easy to see that there is a natural isomorphism  $\mathbf{P} \cong T^*B$ , where  $T^*B = T^*(\mathbb{k}Q)$  is the algebra considered in Sect. 5. Furthermore, the bi-symplectic form  $\sum_{a \in Q} da \, da^*$ , on  $\mathbf{P}$ , agrees with the bi-symplectic form on  $T^*(\mathbb{k}Q)$  constructed in Sect. 5. Note, however, that the grading on  $\mathbf{P}$  by length of the path differs from the grading on the algebra  $T^*B$  used in Sect. 5 (this latter grading assigns grade degree zero to any edge  $a \in Q$ ).

Further, it has been shown in [CB2] that, in the notation of Sect. 5 one has an algebra isomorphism  $\Pi^c(Q) \cong \Pi^c(\Bbbk Q)$ .

For any dimension vector  $\mathbf{d} = (d_i \in \mathbb{Z}_{\geq 0}, i \in I)$ , one associates with  $\mathbf{P}$ , resp., with  $\Pi^c(Q)$ , the corresponding commutative algebra  $\mathcal{O}_{\mathbf{d}}(\mathbf{P})$ , resp.,  $\mathcal{O}_{\mathbf{d}}(\Pi^c(Q))$ , see Sect. 6. The necklace Lie bracket induces Poison algebra structures on both  $\mathcal{O}_{\mathbf{d}}(\mathbf{P})$  and  $\mathcal{O}_{\mathbf{d}}(\Pi^c(Q))$ , see Sect. 6.4. Furthermore, according to Theorem 6.4, the scheme  $\operatorname{Rep}(\mathbf{P}, V)$ , dim  $V = \mathbf{d}$ , acquires a symplectic structure. This makes  $\mathbb{k}[\operatorname{Rep}(\mathbf{P}, V)]$  a Poisson algebra, and each of the following subalgebras is stable under the Poisson bracket

$$\mathbb{k}[\operatorname{Rep}(\mathbf{P}, V)] \supset \mathbb{k}[\operatorname{Rep}(\mathbf{P}, V)]^G \twoheadrightarrow \mathbb{k}[\operatorname{Rep}(\Pi^c(Q), V)]^G = \mathbb{k}[\mu^{-1}(c)]^G.$$

In the special case c = 0, we put  $\Pi := \mathbf{P}/\mathbf{PwP}$ , the ordinary preprojective algebra. In Section 11.5 we will prove

**Proposition 8.2.2.** If Q is connected and not Dynkin or extended Dynkin, then its preprojective algebra  $\Pi$  is prime, and has center  $Z(\Pi) = \mathbb{k}1$ .

Next, recall the R-bimodule E spanned by the edges of  $\overline{Q}$ . The degree 2 component of the algebra  $\mathbf{P}$  is equal to  $E \otimes_R E$ . Thus, using Sweedler's notation we may write  $\mathbf{w} = \mathbf{w}' \otimes \mathbf{w}'' \in E \otimes_R E$ . With this notation, one has the following explicit description of the space of 1-forms, resp., of derivations, for the algebra  $\Pi$ .

**Lemma 8.2.3.** (i) The assignment  $p'(da) p'' \mapsto p' \otimes a \otimes p''$  gives a  $\Pi$ -bimodule isomorphism

$$\Omega_R^1 \Pi \cong \frac{\Pi \otimes_R E \otimes_R \Pi}{\Pi \cdot (\mathbf{w}' \otimes \mathbf{w}'' \otimes 1 - 1 \otimes \mathbf{w}' \otimes \mathbf{w}'') \cdot \Pi}$$

(ii) The assignment  $\theta \longmapsto \sum_a \theta(a^*) \otimes a$  induces a vector space isomorphism

$$\operatorname{Der}_R\Pi\overset{\sim}{\to}\big\{\sum\nolimits_{a\in\overline{Q}}p_a\otimes a\in(\Pi\otimes_RE)^R\ \big|\ \sum\nolimits_{a\in\overline{Q}}[p_a,a]=0\big\}.$$

Proof of the Lemma is straightforward and is left to the reader.

**8.3** Necklace bracket. The bi-symplectic form  $\omega$  gives a Lie bracket on  $\mathbf{P}_{\natural}$ , known as the *necklace Lie bracket*, see [BLB],[Gi].

To write explicit formulas for this bracket it is convenient to introduce the following notation. Given a path p in  $\overline{Q}$  of length  $\ell(p)$ , for each integer  $1 \leq i \leq \ell(p)$ , let  $p_{< i}$ ,  $a_i$  and  $p_{> i}$  be the paths of lengths i-1, 1 and  $\ell(p)-i$  respectively, with  $p=(p_{< i})a_i(p_{> i})$ . For each edge  $a\in \overline{Q}$  we introduce a map

$$\partial_a : \mathbf{P}_{\natural} \longrightarrow \mathbf{P}, \ p \mapsto \partial_a p := \sum_{i=1}^{\ell(p)} \delta_{a,a_i} \cdot (p_{>i})(p_{< i})$$
 (8.3.1)

(where  $\delta$  is the Kronecker delta function).

The proof of the next lemma will be given at the end of this section.

**Lemma 8.3.2.** We have the following formulas:

$$dp = \sum_{a \in \overline{Q}} (\partial_a p) da \text{ holds in } DR_R^1 \mathbf{P}, \quad \forall p \in \mathbf{P}_{\natural}.$$

$$\theta_p(a) = \epsilon(a) \cdot \partial_{a^*} p \text{ holds in } \mathbf{P}, \quad \forall p \in \mathbf{P}_{\natural};$$

$$\{p, q\} = \sum_{a \in \overline{Q}} \epsilon(a^*) \cdot (\partial_a p) (\partial_{a^*} q), \quad \forall p, q \in \mathbf{P}_{\natural}.$$

By Proposition 4.4.3(ii), we know that, for any  $c \in R$ , the necklace Lie bracket on  $\mathbf{P}_{\natural}$  descends to a well-defined Lie bracket on  $\Pi^{c}(Q)_{\natural}$ , cf. also Proposition 5.1.5.

From now on, we will only consider the case c=0, so  $\Pi=\Pi^0(Q)$ . We set  $L:=\Pi_{\natural}=\Pi/[\Pi,\Pi]$ .

The grading on the path algebra  $\mathbf{P} = \bigoplus_{k \geq 0} \mathbf{P}_k$ , by length of the path, descends to natural gradings  $\Pi = \bigoplus_{k \geq 0} \Pi_k$  and  $L = \bigoplus_{k \geq 0} L_k$ . These gradings make  $\mathbf{P}_{\natural}$  and L graded Lie algebras, with necklace Lie bracket of degree (-2), e.g., for any  $k, m \geq 0$ , we have  $\{L_k, L_m\} \subset L_{k+m-2}$  (the bracket vanishes whenever k+l < 0).

Associated with the grading on  $\Pi$ , we have the corresponding *Euler derivation*  $\operatorname{eu}:\Pi\to\Pi$  and a Lie algebra derivation  $\operatorname{eu}_{\operatorname{Lie}}:L\to L$  defined by  $\operatorname{eu}|_{L_k}=(k-2)\cdot\operatorname{Id}_{L_k}$ , for any  $k=0,1,\ldots$  Our present notation for Euler derivations is different from the one used in Sect. 5, since the grading on  $\Pi$  that we are using now is *not* the one used in Sect. 5.

**8.4** Hochschild cohomology. We use the notation  $HH^{\bullet}(\Pi) := H_{R}^{\bullet}(\Pi, \Pi)$  for relative Hochschild cohomology of the preprojective algebra  $\Pi$ , cf. Remark 4.5.1. According to Proposition 4.4.3, we have a Lie algebra map  $L = \Pi_{\natural} \to HH^{1}(\Pi)$ . As in Proposition 5.1.5, this map gives rise to a Lie algebra map  $\mathbb{k} \cdot \operatorname{eu}_{\operatorname{Lie}} \ltimes (L/R) \to HH^{1}(\Pi)$ , where the semidirect product  $\mathbb{k} \cdot \operatorname{eu}_{\operatorname{Lie}} \ltimes (L/R)$  is viewed as a graded Lie algebra with  $\operatorname{eu}_{\operatorname{Lie}}$  assigned grade degree zero.

One of the main results of this paper, to be proved in §9, is the following

**Theorem 8.4.1.** Assume that Q is neither Dynkin nor extended Dynkin. Then

(i) The map  $p\mapsto \bar{\theta}_p,\ \mathrm{eu}_{\scriptscriptstyle\mathrm{Lie}}\mapsto \mathrm{eu},\ induces\ a\ graded\ Lie\ algebra\ isomorphism$ 

$$\mathbb{k} \cdot \mathsf{eu}_{\text{\tiny Lie}} \ltimes (L/R) \stackrel{\sim}{\to} \operatorname{Der}_R \Pi / \operatorname{Inn}_R \Pi = HH^1(\Pi).$$

(ii) The Hochschild cohomology of the preprojective algebra is given by

$$HH^{i}(\Pi) \cong \begin{cases} \mathbb{k} & \text{if } i = 0\\ \mathbb{k} \cdot \text{eu}_{\text{Lie}} \ltimes (L/R) & \text{if } i = 1\\ L & \text{if } i = 2\\ 0 & \text{if } i > 2. \end{cases}$$

$$(8.4.2)$$

The isomorphism  $HH^1(\Pi) \cong \mathbb{k} \cdot \operatorname{eu}_{\operatorname{Lie}} \ltimes (L/R)$  in (8.4.2) can be shown to map the (Gerstenhaber) Lie bracket on  $HH^1(\Pi)$  to the bracket in the Lie algebra  $\mathbb{k} \cdot \operatorname{eu}_{\operatorname{Lie}} \ltimes (L/R)$ . Furthermore, one verifies that the Gerstenhaber bracket  $HH^1(\Pi) \times HH^2(\Pi) \longrightarrow HH^2(\Pi)$  corresponds, via formula (8.4.2) for i = 1, 2, to the adjoint action of the Lie algebra L/R on L. Thus, Theorem 8.4.1 completely describes the Gerstenhaber bracket on the Hochschild cohomology of  $\Pi$ .

There is also an associative and graded-commutative cup-product  $HH^i \times HH^j \to HH^{i+j}$  on Hochschild cohomology. In the case of the preprojective algebra, the only non-trivial cup-product is the skew-symmetric pairing  $\cup : HH^1(\Pi) \times HH^1(\Pi) \to HH^2(\Pi)$ . It is easy to see that, under the isomorphism in (8.4.2), this pairing is given by the formulas

$$\operatorname{eu} \cup \operatorname{eu} = 0, \quad \operatorname{eu} \cup p = (\deg p) \cdot p, \quad p \cup q = \{p, q\},$$

for any homogeneous elements  $p, q \in L/R$ , (since R is central in L, the necklace bracket in the last formula is viewed as a map  $L/R \times L/R \to L$ ).

**8.5 Dynkin case.** In the case of (finite) Dynkin quiver Q, the corresponding preprojective algebra  $\Pi$  is *not* Koszul, in general, and there are nonzero cohomology groups  $HH^i(\Pi)$  for i > 2, see [ES], [BBK].

Now let Q be an extended Dynkin quiver and let  $\Gamma \subset SL_2(\mathbb{k})$  be the finite group associated to Q via the McKay correspondence. It is convenient to introduce a 2-dimensional vector space V and write  $\Gamma \subset SL(V) = SL_2(\mathbb{k})$ . Also, put  $\mathscr{S} := \Gamma \setminus \{1\}$  and let  $\mathbb{k}[\mathscr{S}]^{\Gamma}$  denote the space of class-functions on  $\mathscr{S}$ .

According to [CBH], the preprojective algebra  $\Pi = \Pi^0(Q)$  is Morita equivalent to  $\mathbb{k}[V]\sharp\Gamma$ , the cross product of  $\Gamma$  with the polynomial algebra on V. Thus, by Morita invariance of Hochschild cohomology, we have  $HH^{\bullet}(\Pi) \cong HH^{\bullet}(\mathbb{k}[V]\sharp\Gamma)$ . Further, the algebra  $\mathbb{k}[V]$ , hence the cross product algebra  $\mathbb{k}[V]\sharp\Gamma$ , is Koszul. Therefore, for Hochschild cohomology one easily finds

$$HH^{i}(\Pi) \cong HH^{i}(\mathbb{k}[V]\sharp\Gamma) = \begin{cases} \mathbb{k}[V]^{\Gamma} & \text{if } i = 0\\ (\mathbb{k}[V] \otimes V)^{\Gamma} & \text{if } i = 1\\ (\mathbb{k}[\mathscr{S}] \oplus \mathbb{k}[V])^{\Gamma} \otimes \wedge^{2}V & \text{if } i = 2\\ 0 & \text{if } i > 2. \end{cases}$$
(8.5.1)

In formula (8.5.1), the space  $(\Bbbk[V] \otimes V)^{\Gamma}$  stands for the Lie algebra of  $\Gamma$ -invariant polynomial vector fields on V. Note that we may interpret  $L/R = \Pi_{\natural}/R = (\Bbbk[V]\sharp\Gamma)_{\natural}/R$  as the Lie algebra of  $\Gamma$ -invariant Hamiltonian vector fields on V. Thus, we see that the semidirect product  $\Bbbk \cdot \operatorname{eu} \ltimes (L/R)$ , occurring in formula (8.4.2), is a proper Lie subalgebra in  $(\Bbbk[V] \otimes V)^{\Gamma}$ . Observe also that the group  $\Gamma$  acts trivially on  $\wedge^2 V$ . Thus,  $\Bbbk[V]^{\Gamma} \otimes \wedge^2 V = (\Bbbk[V] \otimes \wedge^2 V)^{\Gamma}$ , is the space of  $\Gamma$ -invariant polynomial bivector fields on V. This is clearly a rank 1 free  $\Bbbk[V]^{\Gamma}$ -module with generator  $\pi \in \wedge^2 V$ , the constant volume element.

Further, by the general duality between Hochschild homology and cohomology due to Van den Bergh [VB1], see (4.5.2)-(4.5.3), one has a canonical isomorphism  $\wedge^2 V \otimes HH_{\bullet}(\Bbbk[V]\sharp\Gamma) \cong HH^{2-\bullet}(\Bbbk[V]\sharp\Gamma)$ . Therefore, the isomorphisms in (8.5.1) yield formulas for Hochschild homology of the algebra  $\Pi$  as well. In particular, we get a chain of isomorphisms

$$(\mathbb{k}[\mathscr{S}] \oplus \mathbb{k}[V])^{\Gamma} \xrightarrow{\sim} \wedge^{2} V^{*} \otimes HH^{2}(\Pi) \xrightarrow{\sim} HH_{0}(\Pi) = \Pi_{h}. \tag{8.5.2}$$

The composite isomorphism in (8.5.2) can be described explicitly as follows. The restriction of this map to the direct summand  $\mathbb{k}[\mathscr{S}]^{\Gamma}$  is obtained as a composition  $\mathbb{k}[\mathscr{S}]^{\Gamma} \xrightarrow{\sim} R_0 \hookrightarrow R \hookrightarrow \Pi_{\natural}$ , where we have used the notation  $R_0 := \{r = \sum r_i e_i \in R \mid \sum r_i = 0\}$  and the first of the above maps is explained e.g. in [CBH]. The restriction of the composite in (8.5.2) to the direct summand  $\mathbb{k}[V]^{\Gamma}$  is equal to the composition

$$\Bbbk[V]^{\Gamma} \hookrightarrow \Bbbk[V] \sharp \Gamma \twoheadrightarrow (\Bbbk[V] \sharp \Gamma)_{\natural} \xrightarrow{\sim} \Pi_{\natural},$$

where the first two maps are the natural maps and the rightmost isomorphism is provided by Morita equivalence.

**8.6 Poisson algebras.** Let  $R[\mathbf{w}] \subset \mathbf{P}$  be the subalgebra generated by R and  $\mathbf{w}$ , and let  $\overline{R[\mathbf{w}]} \subset \mathbf{P}_{\natural}$  denote the image of  $R[\mathbf{w}]$  under the projection  $\eta: R[\mathbf{w}] \hookrightarrow \mathbf{P} \twoheadrightarrow \mathbf{P}/[\mathbf{P}, \mathbf{P}] = \mathbf{P}_{\natural}$ . The imbedding of vector spaces  $\overline{R[\mathbf{w}]} \hookrightarrow \mathbf{P}_{\natural}$  extends, by multiplicativity to an injective graded algebra homomorphism  $\operatorname{Sym}(\overline{R[\mathbf{w}]}) \hookrightarrow \operatorname{Sym}(\mathbf{P}_{\natural})$ , of the corresponding symmetric algebras (over  $\Bbbk$ ).

The Lie algebra structure on  $\mathbf{P}_{\natural}$ , resp., on  $\Pi_{\natural}$ , gives the symmetric algebra  $\mathrm{Sym}(\mathbf{P}_{\natural})$ , resp.,  $\mathrm{Sym}(\Pi_{\natural})$ , the structure of a Poisson algebra. By Lemma 4.4.2, the space  $\overline{R[\mathbf{w}]}$  is contained in the center of the Lie algebra  $\mathbf{P}_{\natural}$ , hence,  $\mathrm{Sym}(\overline{R[\mathbf{w}]})$  is a central subalgebra of the Poisson algebra  $\mathrm{Sym}(\mathbf{P}_{\natural})$ .

Similarly, the imbedding  $R \hookrightarrow \Pi_{\natural}$  makes  $\operatorname{Sym} R$  a central Poisson subalgebra in  $\operatorname{Sym}(\Pi_{\natural})$ .

The third important result of this paper is the following theorem, to be proved in Section 11 below.

**Theorem 8.6.1.** Assume that Q is neither Dynkin nor extended Dynkin. Then (i) The Poisson center of the algebra  $Sym(\mathbf{P}_{\natural})$  is equal to  $Sym(\overline{R[\mathbf{w}]})$ .

Furthermore, the kernel of the natural projection  $\operatorname{Sym}(R[\mathbf{w}]) \twoheadrightarrow \operatorname{Sym}(\overline{R[\mathbf{w}]})$  is a principal ideal in the algebra  $\operatorname{Sym}(R[\mathbf{w}])$  generated by the element  $\mathbf{w} \in \operatorname{Sym}^1(R[\mathbf{w}]) = R[\mathbf{w}]$ .

(ii) The Poisson center of the algebra  $\operatorname{Sym}(\Pi_h)$  is equal to  $\operatorname{Sym} R$ .

Corollary 8.6.2. If Q is neither Dynkin nor extended Dynkin, then we have

- (i) The center of the Lie algebra  $\mathbf{P}_{\natural}$  equals  $\overline{R[\mathbf{w}]}$ ; moreover, the kernel of the projection  $\eta: R[\mathbf{w}] \to \overline{R[\mathbf{w}]}$  is the line spanned by  $\mathbf{w}$ .
  - (ii) The center of the Lie algebra  $\Pi_{\natural}$  equals R.
- **8.7** Proof of Lemma 8.3.2. First, fix an arbitrary derivation  $\theta$ . We compute

$$i_{\theta}\omega = i_{\theta}\left(\sum_{a \in Q} da \, da^{*}\right) = \sum_{a \in Q} \left(i_{\theta} da\right) da^{*} - \sum_{a \in Q} da \left(i_{\theta} da^{*}\right)$$

$$= \sum_{a \in Q} \left(\theta(a) \, da^{*} - da \, \theta(a^{*})\right)$$

$$= \sum_{a \in \overline{Q}} \epsilon(a) \cdot \theta(a) \, da^{*} \, \operatorname{mod}\left[\mathbf{P}, \Omega_{R}^{1}\mathbf{P}\right].$$
(8.7.1)

Next, let  $p = (p_{< i})a_i(p_{> i})$  be a path in  $\overline{Q}$ . For the 1-form  $dp \in DR_R^1 \mathbf{P} = \Omega_R^1 \mathbf{P}/[\mathbf{P}, \Omega_R^1 \mathbf{P}]$ , we find

$$dp = \sum_{i=1}^{\ell(p)} (p_{< i}) \, da_i \, (p_{> i}) = \sum_{i=1}^{\ell(p)} (p_{> i}) (p_{< i}) \, da_i \, \operatorname{mod} \left[ \mathbf{P}, \Omega_R^1 \mathbf{P} \right]. \tag{8.7.2}$$

Now, let  $\theta : \mathbf{P} \to \mathbf{P}$  be the derivation that annihilates  $\mathbb{k}I \subset \mathbf{P}$  and acts on generators of  $\mathbf{P}$  by the formula  $a \longmapsto \epsilon(a) \cdot \partial_{a^*}p$ . For this derivation, we obtain

$$i_{\theta}\omega \stackrel{(8.7.1)}{=} \sum_{a \in \overline{Q}} \sum_{i=1}^{\ell(p)} \epsilon(a^*) \cdot \delta_{a,a_i} \cdot (p_{>i})(p_{< i}) da^* = \sum_{i=1}^{\ell(p)} (p_{>i})(p_{< i}) da_i \stackrel{(8.7.2)}{=} dp.$$

We see that the derivation  $\theta$  satisfies the same equation  $i_{\theta}\omega = dp$  as the derivation  $\theta_p$  attached to the image of p in  $\mathbf{P}_{\natural}$  via the map (4.2.3). Hence,  $\theta = \theta_p$  and, therefore,  $\theta_p(a) = \epsilon(a) \cdot \partial_{a^*} p$ .

The rest of the proof is straightforward and is left to the reader.  $\Box$ 

### 9 Proof of Theorem 8.4.1

**9.1** Given a  $\mathbb{Z}$ -graded vector space  $M = \bigoplus_{k \in \mathbb{Z}} M_k$ , let  $P(M,t) = \sum_{k \in \mathbb{Z}} \dim M_k \cdot t^k \in \mathbb{k}[[t,t^{-1}]]$  denote its Poincaré series. If each graded piece  $M_k$  is an R-bimodule, then one can define a refined matrix-valued Poincaré series  $P_{\text{matrix}}(M,t)$ , that is an  $I \times I$ -matrix whose entries are elements of  $\mathbb{k}[[t,t^{-1}]]$  given by  $P_{\text{matrix}}(M,t)_{ij} := \sum_{k \in \mathbb{Z}} \dim(e_i M e_j) \cdot t^k$ ,  $i,j \in I$ . With this notation, we have

$$P(M^R, t) = \operatorname{Tr} P_{\text{matrix}}(M, t). \tag{9.1.1}$$

**9.2** From now on, we assume that Q is neither Dynkin nor extended Dynkin. We are interested in the Hochschild cohomology  $HH^{\bullet}(\Pi)$  of the algebra  $\Pi$ . Observe that the natural grading on  $\Pi$  induces a grading on each Hochschild cohomology group.

**Lemma 9.2.1.** The groups  $HH^{i}(\Pi)$  vanish for all i > 2, and we have

$$P(HH^{0}(\Pi), t) - P(HH^{1}(\Pi), t) + P(HH^{2}(\Pi), t) = |I|.$$

*Proof.* Since Q is neither Dynkin nor extended Dynkin, the algebra  $\Pi$  is known to be a Koszul algebra, see [MV], [MOV]. Specifically, a  $\Pi$ -bimodule resolution of  $\Pi$  is provided by the following Koszul complex

$$\left(0 \to \Pi \otimes_R \Pi \longrightarrow \Pi \otimes_R E \otimes_R \Pi \longrightarrow \Pi \otimes_R \Pi\right) \stackrel{m}{\twoheadrightarrow} \Pi, \tag{9.2.2}$$

where E denotes the R-bimodule generated by the edges of  $\overline{Q}$ . Therefore, the Hochschild cohomology of  $\Pi$  may be computed by applying  $\operatorname{Hom}_{\Pi\text{-bimod}}(-,\Pi)$  to the complex above. This way, we get the following complex

$$\Pi^R \longrightarrow (\Pi \otimes E)^R \longrightarrow \Pi^R.$$
(9.2.3)

The vanishing part of the Lemma is now clear.

Let  $\chi_{\mathrm{matrix}}$  denote the matrix-valued graded Euler characteristic, that is, the alternating sum of the matrix-valued Poincaré series of the complex  $\Pi \to \Pi \otimes E \to \Pi$ . The matrix  $\chi_{\mathrm{matrix}}$  is easily expressed in terms of the matrix  $P := P_{\mathrm{matrix}}(\Pi, t)$  and the adjacency matrix C of the graph  $\overline{Q}$  by the formula  $\chi_{\mathrm{matrix}} = t^2 \cdot P - t \cdot P \cdot C + P$ .

Further, by formula (9.1.1), the Euler characteristic of the complex (9.2.3) is an element of  $\mathbb{k}[[t]]$  which is equal to  $\text{Tr}(\chi_{\text{matrix}})$ . Now, according to [MOV], one has a matrix identity  $P = 1/(1 - C \cdot t + t^2)$ . Thus, we compute

$$\operatorname{Tr}(\chi_{\operatorname{matrix}}) = \operatorname{Tr}(t^2 \cdot P - t \cdot P \cdot C + P) = \operatorname{Tr}(P \cdot (1 - C \cdot t + t^2)) = \operatorname{Tr}(\operatorname{Id}) = |I|,$$
 and we are done.

**9.3** Proof of formula (8.4.2). Case i = 0 of the formula follows from Proposition 8.2.2 (to be proved later, in Sect. 11.5).

Next, one observes that the complex inside the parenthesis in (9.2.2) is self-dual. In other words, applying the functor  $\operatorname{Hom}_{\Pi\text{-bimod}}(-,\Pi\otimes\Pi)$  to this complex one obtains the same complex again. Therefore, using (9.2.2) as a resolution for the computation of  $\operatorname{Ext}_{\Pi\text{-bimod}}^{\bullet}(\Pi,\Pi\otimes\Pi)$ , we see that the only nonzero cohomology group sits in degree two and we have  $\operatorname{Ext}_{\Pi\text{-bimod}}^{2}(\Pi,\Pi\otimes\Pi)\cong\Pi$ . We conclude that formula (4.5.5) holds for the algebra  $A=\Pi$  and d=2. Thus, from Van den Bergh's isomorphism (4.5.2) we deduce  $HH^{i}(\Pi)=HH_{2-i}(\Pi)$ , for any i. In particular, we get  $HH^{2}(\Pi)=HH_{0}(\Pi)=\Pi/[\Pi,\Pi]=L$ .

We have already computed  $HH^i(\Pi)$  for i = 0, 2. Furthermore, we know that  $HH^i(\Pi) = 0$  for all i > 2, by the first claim of Lemma 9.2.1. Therefore, from the formula for the Poincaré series given in Lemma 9.2.1 we deduce

$$\begin{split} P(HH^{1}(\Pi),t) &= -|I| + P(HH^{0}(\Pi),t) + P(HH^{2}(\Pi),t) \\ &= -|I| + P(Z(\Pi),t) + P(L,t) = P(L,t) - |I| + 1 \\ &= P(L/R,t) + 1 = P(\mathbb{k} \cdot \operatorname{eu}_{\operatorname{Lie}} \ltimes (L/R),t). \end{split}$$

By comparing the two sides, we see that the case i = 1 of (8.4.2), as well as part (i) of the Theorem, follows by dimension argument from the injectivity statement in Proposition 5.1.5. This completes the proof of formula (8.4.2).  $\square$ 

# 10 Deformations of the preprojective algebra

10.1 It is well-known that first order infinitesimal deformations of an associative algebra are controlled by the second Hochschild cohomology group, and obstructions to deformations are controlled by the third Hochschild cohomology group. Thus, according to formula (8.4.2), first order infinitesimal deformations of the preprojective algebra  $\Pi$  are unobstructed, and its versal deformation is parametrized by the vector space L.

We will show that, in the non-Dynkin case, any formal (infinite order) one-parameter deformation of the algebra  $\Pi$  is equivalent to one obtained by deforming the defining relation in the preprojective algebra as follows:

$$\sum_{a \in Q} \left[ a, a^* \right] = t \cdot f,$$

where t is the deformation parameter and f is an element of  $\mathbf{P}[[t]]$  that commutes with R.

In more detail, from the general formula (2.2.3) applied to the algebra  $A = \Pi$  we deduce that one can find a graded subspace  $\mathbf{P}_L \subset \mathbf{P}^R$  which is complementary to the graded space  $\mathrm{Ker}[\mathbf{P} \twoheadrightarrow L] = [\mathbf{P}, \mathbf{P}] + \mathbf{PwP} \subset \mathbf{P}$ , i.e., such that we have

$$\mathbf{P} = \mathbf{P}_L \bigoplus ([\mathbf{P}, \mathbf{P}] + \mathbf{P}\mathbf{w}\mathbf{P}), \text{ and } \mathbf{P}_L \subset \mathbf{P}^R.$$
 (10.1.1)

We fix such a subspace  $\mathbf{P}_L$  once and for all, and also choose and fix a graded  $\mathbb{k}$ -basis  $\{f_j\}_{j\in\mathbb{N}}$  of  $\mathbf{P}_L$ . For each  $j\in\mathbb{N}$ , introduce a formal variable  $t_j$  that may be thought of as the j-th coordinate function on  $\mathbf{P}_L$  with respect to the basis  $\{f_j\}_{j\in\mathbb{N}}$ .

Given a k-vector space V, write  $V[[\mathbf{t}]] := \lim_{\substack{n \to \infty \\ n \to \infty}} V[[t_1, ..., t_n]]$  for the vector space of V-valued formal power series in the (infinitely many) formal variables  $t_j$ . Thus,  $k[[\mathbf{t}]]$  is a complete topological local k-algebra with maximal ideal  $(\mathbf{t}) \subset k[[\mathbf{t}]]$  formed by the series without constant term. For any V, the space  $V[[\mathbf{t}]]$  acquires a natural structure of complete topological  $k[[\mathbf{t}]]$ -module. In particular, we have a  $k[[\mathbf{t}]]$ -module  $\mathbf{P}_L[[\mathbf{t}]]$ . Observe that the identity map  $\mathbf{I} : \mathbf{P}_L \to \mathbf{P}_L$  may be viewed as an element of  $\mathbf{P}_L[[\mathbf{t}]]$  that has the form  $\sum_{j \in \mathbb{N}} t_j \cdot f_j$ .

We define a formal deformation,  $\widehat{\Pi}$ , of the preprojective algebra  $\Pi$  as the quotient of the algebra  $\mathbf{P}[[\mathbf{t}]]$  modulo the *closed* two-sided ideal (topologically) generated by the element

$$\sum_{a \in Q} [a, a^*] - \sum_{j \in \mathbb{N}} t_j \cdot f_j. \tag{10.1.2}$$

Thus,  $\widehat{\mathbf{\Pi}}$  is a complete topological  $\mathbb{k}[[\mathbf{t}]]$ -algebra that may be thought of as a deformation of  $\Pi$  with formal parameters  $\{t_j\}$ , i.e., the base of the deformation is  $\operatorname{Spec}(\mathbb{k}[[\mathbf{t}]])$ , a formal scheme.

**Theorem 10.1.3.** If the quiver Q is not Dynkin (but possibly extended Dynkin) then the deformation  $\widehat{\Pi}$  is flat over  $\mathbb{k}[[\mathbf{t}]]$ , and is a versal deformation of  $\Pi$ .

#### 10.2 We begin the proof of Theorem 10.1.3 with an easy lemma.

Let the vector space  $\widetilde{\Pi} = \Pi \otimes \mathbb{k}[t]/(t^2)$  be equipped with a  $\mathbb{k}[t]/(t^2)$ -algebra structure given by a star-product of the form  $p, q \longmapsto p \star q = pq + t \cdot \beta(p, q)$ . Thus,  $\widetilde{\Pi}$  is a first order deformation of the algebra  $\Pi$ . We write  $[\beta] \in HH^2(\Pi)$  for the Hochschild cohomology class corresponding to this deformation.

Recall the notation  $\Pi^R$  for the centralizer of R in  $\Pi$ .

**Lemma 10.2.1.** Assume that in  $\Pi$  we have

$$\sum_{a \in Q} (a \star a^* - a^* \star a) = t \cdot c \mod(t^2), \quad \textit{for some} \quad c \in \Pi^R.$$

Then, the class  $[\beta] \in HH^2(\Pi)$  goes, via the isomorphism  $HH^2(\Pi) \xrightarrow{\sim} L$  of Theorem 8.4.1, cf. also Sect. 8.5, to the image of  $c \in \Pi$  under the projection  $\Pi \to \Pi/[\Pi,\Pi] = L$ .

*Proof.* Recall that the Koszul resolution (9.2.2) of the algebra  $\Pi$  admits a standard imbedding, as a subcomplex, into the Bar-resolution of  $\Pi$ :

Koszul: 
$$\Pi \otimes_{R} \Pi \longrightarrow \Pi \otimes_{R} E \otimes_{R} \Pi \longrightarrow \Pi \otimes_{R} \Pi \quad (10.2.2)$$

$$\downarrow^{i_{2}} \qquad \qquad \downarrow^{i_{1}} \qquad \qquad \parallel_{\mathrm{Id}}$$
Bar: 
$$\Pi \otimes_{R} T_{R}^{2} \Pi \otimes_{R} \Pi \longrightarrow \Pi \otimes_{R} \Pi \otimes_{R} \Pi \longrightarrow \Pi \otimes_{R} \Pi,$$

In this diagram,  $i_1$  is a  $\Pi$ -bimodule map induced by the natural imbedding  $E \hookrightarrow \Pi$  and  $i_2$  is a  $\Pi$ -bimodule map defined by the formula

$$i_2: p \otimes q \longmapsto \sum_{a \in Q} (p \otimes a \otimes a^* \otimes q - p \otimes a^* \otimes a \otimes q).$$

Now, given a deformation of  $\Pi$  as in the lemma, the class  $[\beta] \in HH^2(\Pi)$  corresponds to a 2-cocycle in  $\operatorname{Hom}_{\Pi\text{-bimod}}(\Pi \otimes_R T_R^2\Pi \otimes_R \Pi, \Pi)$  given by

$$\widetilde{\beta}: \Pi \otimes_R T_R^2 \Pi \otimes_R \Pi \longrightarrow \Pi, \quad p \otimes u \otimes v \otimes q \longmapsto p \cdot \beta(u, v) \cdot q.$$

It is clear that, under the imbedding of complexes (10.2.2), the cocycle  $\widetilde{\beta}$  restricts to a map  $i_2^*(\widetilde{\beta}) \in \operatorname{Hom}_{\Pi\text{-bimod}}(\Pi \otimes_R \Pi, \Pi)$  such that  $i_2^*(\widetilde{\beta}) : p \otimes q \longmapsto p \cdot c \cdot q$ . Hence, the cohomology class of  $i_2^*(\widetilde{\beta})$  in the complex (9.2.3) is represented by the element  $c \in \Pi^R$ . The lemma follows.

10.3 By Theorem 8.4.1, we know that deformations of  $\Pi$  are unobstructed. Thus, general deformation theory implies that there exists a formal flat multiparameter deformation  $\widetilde{\Pi}$  of the algebra  $\Pi$ , parametrized by the formal neighborhood of zero in  $HH^2(\Pi) = L$ .

Recall that we have chosen a vector space  $\mathbf{P}_L \subset \mathbf{P}$ . It is clear from (10.1.1) that the projection  $\mathbf{P} \twoheadrightarrow L$  induces an isomorphism  $\mathbf{P}_L \stackrel{\sim}{\to} L$ .

We use the last isomorphism to identify  $HH^2(\Pi)$  with  $\mathbf{P}_L$ . Thus, the deformation  $\widetilde{\Pi}$  becomes parametrized by the formal neighborhood of zero in  $\mathbf{P}_L$ . Therefore, there exists an  $R[[\mathbf{t}]]$ -bimodule isomorphism  $\widetilde{\Pi} \cong \Pi[[\mathbf{t}]]$ , equal to the identity modulo the ideal  $(\mathbf{t})$ , in the notation of §10.1. We transport the algebra structure on  $\widetilde{\Pi}$  to  $\Pi[[\mathbf{t}]]$  via that isomorphism. This way, we make  $\Pi[[\mathbf{t}]]$  a complete topological  $\mathbb{k}[[\mathbf{t}]]$ -algebra, to be denoted  ${}^*\Pi[[\mathbf{t}]]$ .

Further, for each  $k \geq 0$ , let  $\Pi[[\mathbf{t}]]_k$  denote the space of  $\Pi$ -valued series homogeneous in  $t_j$ 's of degree k. Writing a power series in the variables  $t_j$  as  $p = \sum_{k \geq 0} p_k$ ,  $p_k \in \Pi[[\mathbf{t}]]_k$ , yields a direct product decomposition  $\Pi[[\mathbf{t}]] = \prod_{k \geq 0} \Pi[[\mathbf{t}]]_k$ . Thus, multiplication in the algebra  $\Pi[[\mathbf{t}]]$  takes the form of star-product:

$$(p,q) \longmapsto p \star q = pq + \beta_1(p,q) + \beta_2(p,q) + \dots, \quad \beta_k : \Pi \times \Pi \longrightarrow \Pi[[\mathbf{t}]]_k.$$

10.4 We return now to the setup of Theorem 10.1.3 and consider the  $\mathbb{k}[[t]]$ -algebra  $\widehat{\Pi}$ . In Sect. 10.5 we will prove the following

**Lemma 10.4.1.** There exists a continuous algebra automorphism  $\varphi : \mathbb{k}[[\mathbf{t}]] \xrightarrow{\sim} \mathbb{k}[[\mathbf{t}]]$  and a continuous  $\mathbb{k}$ -algebra isomorphism  $\Phi : \widehat{\mathbf{\Pi}} \xrightarrow{\sim} {}^{\star}\Pi[[\mathbf{t}]]$  such that

- The map  $\Phi$  is  $\varphi$ -semilinear, i.e., for any  $f \in \mathbb{k}[[\mathbf{t}]]$  and  $p \in \widehat{\mathbf{\Pi}}$ , one has  $\Phi(f \cdot p) = \varphi(f) \cdot \Phi(p)$ ;
- The differential of  $\varphi$  at the origin  $0 \in \mathbf{P}_L$  equals  $d\varphi = \mathrm{Id} : \mathbf{P}_L \to \mathbf{P}_L$ , the identity map.

This Lemma easily implies Theorem 10.1.3. Indeed, the deformation  ${}^{\star}\Pi[[t]]$  being flat by construction, it follows from Lemma 10.4.1 that the deformation  $\widehat{\mathbf{\Pi}}$  is also flat and provides a versal deformation. The theorem is proved.

10.5 Proof of Lemma 10.4.1. Recall the vector space  $\mathbf{P}_L \subset \mathbf{P}$  and let  $\Pi_L$  denote its image under the projection  $\mathbf{P} \twoheadrightarrow \Pi$ . By (10.1.1), we have

$$\Pi = \Pi_L \bigoplus [\Pi, \Pi], \text{ and } \Pi_L \subset \Pi^R.$$
 (10.5.1)

Furthermore we have natural isomorphisms  $\mathbf{P}_L \xrightarrow{\sim} \Pi_L \xrightarrow{\sim} L$ , where the first map is induced by the projection  $\mathbf{P} \twoheadrightarrow \Pi$  and the second map by the projection  $\Pi \twoheadrightarrow L$ .

Next, we introduce the notation  $\mathbf{m}$  for a multi-index, that is for a function  $\mathbf{m}: \mathbb{N} \to \{0, 1, \ldots\}$  such that  $\mathbf{m}(j) = 0$  for all but finitely many j's. Given such a multi-index  $\mathbf{m}$ , write  $t^{\mathbf{m}} := t_1^{\mathbf{m}(1)} t_2^{\mathbf{m}(2)} \ldots$ , for the corresponding monomial in the variables  $t_j$ . Put  $|\mathbf{m}| = \sum_{j \in \mathbb{N}} \mathbf{m}(j)$ .

To prove the lemma, we must show that there exist algebra homomorphisms

$$\Phi: \widehat{\mathbf{\Pi}} \to {}^{\star}\Pi[[\mathbf{t}]], \quad \text{and} \quad \varphi: \mathbb{k}[[\mathbf{t}]] \to \mathbb{k}[[\mathbf{t}]], \ \mathbf{t} \longmapsto \varphi(\mathbf{t}) =: \mathbf{t}' = \{t'_i\}_{i \in \mathbb{N}},$$

where  $t'_j = t_j + \sum_{|\mathbf{m}|>1} \lambda_{i\mathbf{m}} t^{\mathbf{m}}, \lambda_{i\mathbf{m}} \in \mathbb{k}$  (formal change of variables), such that  $\Phi$  is the identity in degree 0 and  $\Phi(a) = a + \sum_{|\mathbf{m}|>0} t^{\mathbf{m}} z_{\mathbf{m}}(a)$  for each edge  $a \in \overline{Q}$ . (here if  $a \in e_i \Pi e_j$  then  $z_{\mathbf{m}}(a)$  also must belong to  $e_i \Pi e_j$  to ensure the relations between projectors  $e_i$  and a are respected by  $\Phi$ ).

We construct  $\Phi$  by induction in  $|\mathbf{m}|$ . First we consider the case  $|\mathbf{m}| = 1$ , i.e.,  $t^{\mathbf{m}} = t_j$  for some j.

In the algebra  $\Pi[[t]]$ , we have an expansion

$$\sum_{a \in Q} \left( a \star a^* - a^* \star a \right) = \sum_{|\mathbf{m}| > 0} t^{\mathbf{m}} \cdot y_{\mathbf{m}}, \quad y_{\mathbf{m}} \in \Pi^R.$$

Hence, Lemma 10.2.1 implies that the image of  $y_{\mathbf{m}} \in \Pi^R$  under the projection to L is equal to  $f_j$ . Therefore, the element  $z_j(a)$  in the expansion  $\Phi(a) = a + \sum_{|\mathbf{m}|>0} t^{\mathbf{m}} z_{\mathbf{m}}(a)$  must satisfy the equation

$$f_j - y_j = \sum_{a \in O} ([z_j(a), a^*] + [a, z_j(a^*)]).$$

Now, since  $y_j$  projects to  $f_j \in L$ , we have  $f_j - y_j \in [\Pi, \Pi]$ . Since  $[\Pi, \Pi]$  is spanned by elements of the form  $[a, z], [a^*, z]$  we see that the above equation does have some solution  $z_j(a)$ .

Next, let  $|\mathbf{m}| \geq 2$ . Then, the elements  $z_{\mathbf{m}}(a)$  are determined by the equation

$$y_{\mathbf{m}} + \sum_{a} ([a, z_{\mathbf{m}}(a^*)] + [z_{\mathbf{m}}(a), a^*]) + \sum_{\mathbf{k} + \mathbf{l} = \mathbf{m}, |\mathbf{k}| > 0, |\mathbf{l}| > 0} \sum_{a} [z_{\mathbf{k}}(a), z_{\mathbf{l}}(a^*)] \in \Pi_L.$$

This equation again has solutions, since  $\Pi = [\Pi, \Pi] \oplus \Pi_L$  and  $\Pi_L \subset \Pi^R$ , by (10.5.1). For such a solution, we have

$$y_{\mathbf{m}} + \sum_{a} ([a, z_{\mathbf{m}}(a^*)] + [z_{\mathbf{m}}(a), a^*]) + \sum_{\mathbf{k} + \mathbf{l} = \mathbf{m}, |\mathbf{k}| > 0, |\mathbf{l}| > 0} \sum_{a} [z_{\mathbf{k}}(a), z_{\mathbf{l}}(a^*)] = \sum_{a} \lambda_{j\mathbf{m}} f_j.$$

Continuing so, we obtain  $z_{\mathbf{m}}(a)$  for all  $\mathbf{m}$ , and we see that because of the construction of  $z_{\mathbf{m}}(a)$ ,  $\Phi$  indeed defines a homomorphism as required, with  $t'_j = t_j + \sum_{|\mathbf{m}|>1} \lambda_{i\mathbf{m}} t^{\mathbf{m}}$ .

## 11 Representation schemes

11.1 Stabilization. Fix a quiver Q with vertex set I and let  $R = \mathbb{k}I$ .

Let  $a_1, \ldots, a_p \in \mathbb{k}Q$  be a finite (possibly empty) collection of homogeneous elements in the path algebra of Q, of some degrees  $\deg a_j \geq 2$ . Let  $\mathfrak{A} \subset \mathbb{k}Q$  denote the two-sided ideal generated by the elements  $a_1, \ldots, a_p$ , and set  $A = \mathbb{k}Q/\mathfrak{A}$ . Thus,  $\mathfrak{A}$  is a graded ideal and  $A = \bigoplus_{r \geq 0} A(r)$  is a graded algebra such that A(0) = R.

Let  $\mathbf{d} = (d_i \in \mathbb{Z}_{\geq 0}, i \in I)$  be an I-tuple, to be referred to as dimension vector. Let  $V = \bigoplus_{i \in I} V_i$  be an R-module such that  $\dim V_i = d_i, \forall i \in I$ . We will write  $\operatorname{Rep}(A, \mathbf{d}) := \operatorname{Rep}(A, V)$  for the corresponding representation scheme and, following Sect. 6.1, put  $G_{\mathbf{d}} := (\prod_i \operatorname{GL}(V_i)) / \mathbb{G}_m$ .

We consider the algebras  $\mathcal{O}_{\mathbf{d}}(A)$  and  $\mathbb{k}[\operatorname{Rep}(A,\mathbf{d})]^{G_{\mathbf{d}}}$ , cf. Sect. 6.2. The grading on A makes  $\mathcal{O}_{\mathbf{d}}(A)$ , resp.,  $\mathbb{k}[\operatorname{Rep}(A,\mathbf{d})]^{G_{\mathbf{d}}}$ , a graded algebra with respect to the *total* grading  $\mathcal{O}_{\mathbf{d}}(A) = \bigoplus_{r \geq 0} \mathcal{O}_{\mathbf{d}}(A)(r)$ , resp.,  $\mathbb{k}[\operatorname{Rep}(A,\mathbf{d})]^{G_{\mathbf{d}}} = \bigoplus_{r \geq 0} \mathbb{k}[\operatorname{Rep}(A,\mathbf{d})]^{G_{\mathbf{d}}}(r)$ .

 $\bar{\text{In}}$  (6.2.1), we have defined an algebra map  $\psi_{\mathbf{d}} = \text{Tr} \circ \text{ev}_{\mathbf{d}} : \mathcal{O}_{\mathbf{d}}(A) \longrightarrow \mathbb{k}[\text{Rep}(A,\mathbf{d})]^{G_{\mathbf{d}}}$ . This map is clearly a graded algebra homomorphism.

The following result is a refined version of the 'stabilization phenomenon' observed in [Gi, §4].

**Proposition 11.1.1.** For any positive integer r, there exists  $N(r) \gg 0$  such that the following holds:

If  $\mathbf{d} = \{d_i\}_{i \in I}$  is such that  $d_i \geq N(r)$  for all  $i \in I$ , then the map  $\psi_{\mathbf{d}} : \mathcal{O}(A)(s) \to \mathbb{k}[\operatorname{Rep}_{\mathbf{d}} A]^{G_{\mathbf{d}}}(s)$ , the restriction of  $\psi_{\mathbf{d}}$  to the homogeneous component of degree s, is a bijection for all  $0 \leq s \leq r$ .

First, we are going to prove the proposition in the special case where  $\mathfrak{A} = 0$ , that is, for  $A = \mathbb{k}Q$ . In this case, we have  $\operatorname{Rep}(A, \mathbf{d}) = \operatorname{Rep}(Q, \mathbf{d})$  is the variety of **d**-dimensional representations of the quiver Q. This variety is a vector space.

#### **Lemma 11.1.2.** Proposition 11.1.1 holds for $A = \mathbb{k}Q$ .

Let **L** denote the direct sum of homogeneous components of  $(\Bbbk Q)_{\natural}$  of strictly positive degrees. Thus, we have a vector space direct sum decomposition  $(\Bbbk Q)_{\natural} = R \bigoplus \mathbf{L}$ . The composite  $\operatorname{Sym}(\mathbf{L}) \hookrightarrow \operatorname{Sym}((\Bbbk Q)_{\natural}) \twoheadrightarrow \mathcal{O}_{\mathbf{d}}(\Bbbk Q)$  is clearly a graded algebra isomorphism.

The proof of Lemma 11.1.2 given below copies the argument in the proof of [Gi, Proposition 4.2]; it is based on the fact that there are no polynomial identities which are satisfied in matrix algebras of all sizes.

Proof of Lemma 11.1.2. To simplify the exposition, we will present the proof in the case where Q has one vertex and two edge-loops at that vertex; the general case is entirely similar. Thus,  $\mathbf{d} = d$  is a single integer, so  $\operatorname{Rep}(Q, \mathbf{d})$  is the space of pairs of  $d \times d$ -matrices X, Y, and the vector space  $\mathbf{L}$  is spanned by nonempty cyclic words w in the alphabet with two letters x, y.

Assume the statement of the lemma is not true. Thus, there exists an element  $g \in \operatorname{Sym} \mathbf{L}$  such that  $\psi_{\mathbf{d}}(g) = 0$  for all d. In other words, we have a relation of degree r of the form

$$\sum_{J} c_{J} \cdot \prod_{w} \operatorname{Tr}(w(X, Y))^{J(w)} = 0, \ (c_{J} \in \mathbb{k})$$

which is satisfied for any matrices X and Y. Here J is a function on the set of cyclic words with nonnegative integer values and finite support.

We may assume without loss of generality that at least one cyclic word present in this relation involves x. Let us differentiate the relation with respect to X. We get

$$\sum_{J} \sum_{u} J(u) \cdot c_{J} \cdot \prod_{w} \text{Tr}(w(X,Y))^{J(w) - \delta_{wu}} \cdot \sum_{k: u_{k} = x} (u_{k+1} \dots u_{k-1})(X,Y) = 0,$$

where  $u_k$  is the k-th letter of the word u (we choose a representation of every cyclic word by a usual word, and agree that  $m \pm 1 = \pm 1$  if u has length m). But for large enough d, all words of length  $\leq r$  of matrices X, Y are linearly independent for generic X, Y (for example, take X, Y to be the operators by which x, y act in the quotient of the free algebra in x, y by the r-th power of the augmentation ideal). Thus, we have

$$\sum_{J} J(u) \cdot c_{J} \cdot \prod_{w} \operatorname{Tr}(w(X, Y))^{J(w) - \delta_{wu}} = 0$$

for each u which contains x. This gives a relation of degree smaller than r, so we are done by using induction.

Proof of Proposition 11.1.1. The algebra projection  $\mathbb{k}Q \to A$  makes  $\operatorname{Rep}(A, \mathbf{d})$  a closed subscheme in  $\operatorname{Rep}(\mathbb{k}Q, \mathbf{d})$ , not necessarily reduced in general. The defining ideal of this subscheme is generated by the matrix elements of the matrix valued functions  $\widehat{a}_1, \ldots, \widehat{a}_p$ .

To avoid confusion, we will use boldface notation for the map  $\psi_{\mathbf{d}} : \mathcal{O}_{\mathbf{d}}(\Bbbk Q) \to \mathbb{k}[\operatorname{Rep}(Q,\mathbf{d})]^{G_{\mathbf{d}}}$  and reserve the notation  $\psi_{\mathbf{d}}$  for the map  $\mathcal{O}_{\mathbf{d}}(A) \to \mathbb{k}[\operatorname{Rep}(A,\mathbf{d})]^{G_{\mathbf{d}}}$ .

Assume the statement of the proposition does not hold. Then there exists a homogeneous element  $g \in \mathcal{O}_{\mathbf{d}}(A)$ , of degree  $0 < s \leq r$ , and a dimension vector  $\mathbf{d}$  such that  $d_i \geq N(r)$ ,  $\forall i$ , and such that one has  $\psi_{\mathbf{d}}(g) = 0$ . Let  $\widetilde{g}$  be a representative of g in Sym(L). Thus,  $\psi_{\mathbf{d}}(\widetilde{g})$  is a degree r homogeneous polynomial on the vector space  $\text{Rep}(\mathbb{k}Q, \mathbf{d})$ .

By construction, the polynomial  $\psi_{\mathbf{d}}(\widetilde{g})$  vanishes on the subscheme  $\operatorname{Rep}(A, \mathbf{d})$ . Hence,  $\psi_{\mathbf{d}}(\widetilde{g})$  is a  $G_{\mathbf{d}}$ -invariant homogeneous polynomial that belongs to the ideal generated by the matrix elements of the functions  $\widehat{a}_1, \ldots, \widehat{a}_p$ . Therefore, using Weyl's first fundamental theorem of invariant theory we deduce that  $\psi_{\mathbf{d}}(\widetilde{g})$  is a linear combination of products of the form  $\operatorname{Tr}(\widehat{w}_1) \cdot \ldots \cdot \operatorname{Tr}(\widehat{w}_m) \cdot \operatorname{Tr}(\widehat{w}_{m+1}\widehat{a}_j)$ , where  $j \in [1, p]$  and  $w_1, \ldots, w_{m+1}$  are certain paths in Q of total length  $\ell(w_1) + \ldots + \ell(w_{m+1}) + \ell(a_j) = r$ .

Now, Lemma 11.1.2 says that the element  $\widetilde{g}$  must be equal to the corresponding linear combination of the elements  $w_1 \& \dots \& w_m \& (w_{m+1}a_j) \in \operatorname{Sym}^{m+1}(\mathbf{L})$ . It is clear that all elements of this form generate the kernel of the algebra projection  $\operatorname{Sym}(\mathbf{L}) \twoheadrightarrow \mathcal{O}_{\mathbf{d}}(A)$ . It follows that g = 0, as desired.

11.2 Quiver varieties. Let  $\overline{Q}$  be the double of a quiver Q, and write  $\mathbf{P} := \mathbb{k}\overline{Q}$  for the corresponding path algebra.

For any dimension vector  $\mathbf{d} \geq 0$ , the vector space  $\operatorname{Rep}(\overline{Q}, \mathbf{d}) = T^* \operatorname{Rep}(Q, \mathbf{d})$  is equipped with a natural symplectic structure. It is easy to see that the corresponding symplectic 2-form on  $\operatorname{Rep}(\overline{Q}, \mathbf{d})$  equals  $\operatorname{Tr} \widehat{\omega}$ , the image of the 2-form  $\omega \in \operatorname{DR}^2_R \mathbf{P}$  under the evaluation map, see Sect. 6.2.

Fix a dimension vector **d**. Recall that the Lie algebra  $\mathfrak{g}_{\mathbf{d}} = \operatorname{Lie} G_{\mathbf{d}}$ , as well as its dual  $\mathfrak{g}_{\mathbf{d}}^*$ , are both identified with the a codimension 1 hyperplane formed by the *I*-tuples  $(x_i \in \operatorname{Lie} \operatorname{GL}(d_i))_{i \in I}$  such that  $\sum_i \operatorname{Tr} x_i = 0$ .

The group  $G_{\mathbf{d}}$  acts linearly on  $\operatorname{Rep}(\overline{Q}, \mathbf{d})$ . This action is Hamiltonian, see Sect. 6.2, and the corresponding moment map reads, cf. [CB1]:

$$\mu : \operatorname{Rep}(\overline{Q}, \mathbf{d}) \longrightarrow \mathfrak{g}_{\mathbf{d}}, \quad \rho \longmapsto \mu(\rho) = \sum_{a \in Q} [\rho(a), \rho(a^*)].$$
 (11.2.1)

In this section, we are interested in the algebra  $\Pi$ , the preprojective algebra of Q. We have  $\Pi = \mathbf{P}/\mathfrak{A}$ , where  $\mathfrak{A}$  is the two-sided ideal generated by the single element  $\mathbf{w} = \sum_{a \in Q} [a, a^*]$ . The algebra  $\Pi$  is the Hamiltonian reduction of  $\mathbf{P}$  with respect to this element. The projection  $\mathbf{P} \to \Pi$  makes  $\operatorname{Rep}(\Pi, \mathbf{d})$ , the scheme of all  $\mathbf{d}$ -dimensional representations of the algebra  $\Pi$ , a closed subscheme in  $\operatorname{Rep}(\overline{Q}, \mathbf{d})$ . We have  $\operatorname{Rep}(\Pi, \mathbf{d}) = \mu^{-1}(0)$ , the scheme theoretic zero fiber of the moment map.

We let  $M(Q, \mathbf{d}) := \operatorname{Spec} \mathbb{k}[\operatorname{Rep}(\Pi, \mathbf{d})]^{G_{\mathbf{d}}}$  be the corresponding categorical quotient, an affine subscheme of  $\operatorname{Rep}(\overline{Q}, \mathbf{d})/\!/G_{\mathbf{d}} := \operatorname{Spec} \mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]^{G_{\mathbf{d}}}$  of finite type. The scheme  $M(Q, \mathbf{d})$  is called *quiver variety*, it is a Poisson scheme that may be thought of as a hamiltonian reduction of  $\operatorname{Rep}(\overline{Q}, \mathbf{d})$ . By Theorem 6.4.3, the map  $\psi_{\mathbf{d}} : \mathcal{O}_{\mathbf{d}}(\mathbf{P}) \longrightarrow \mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]^{G_{\mathbf{d}}}$  is a surjective morphism of graded Poisson algebras that descends to a surjective graded Poisson algebra homomorphism  $\psi_{\mathbf{d}} : \mathcal{O}_{\mathbf{d}}(\Pi) \twoheadrightarrow \mathbb{k}[M(Q, \mathbf{d})]^{G_{\mathbf{d}}}$ .

11.3 We need to recall some results from [CB1] about quiver varieties. Let C denote the adjacency matrix of the quiver Q and write  $A = 2\mathrm{Id} - C$  for the corresponding Cartan matrix. Define the Tits quadratic form on the vector space  $\mathbb{k}^{|I|}$  by the formula

$$q(\mathbf{d}) = \frac{1}{2}(\mathbf{d}, A\mathbf{d}).$$

Let  $p(\mathbf{d}) = 1 - q(\mathbf{d})$ .

Let  $R_+$  be the set of positive roots of Q. Let  $\Sigma_0$  denote the set of positive dimension vectors  $\beta$  such that whenever  $\beta$  is written as a sum of two or more positive roots,  $\beta = \beta_1 + ... + \beta_m$ , one has  $p(\beta) > p(\beta_1) + ... + p(\beta_m)$ .

**Theorem 11.3.1.** ([CB1]). (i)  $\mathbf{d} \in \Sigma_0$  if and only if there is a simple representation of  $\Pi$  with dimension vector  $\mathbf{d}$ .

If  $\mathbf{d} \in \Sigma_0$  then:

- (ii) the schemes  $Rep(\Pi, \mathbf{d})$  and  $M(Q, \mathbf{d})$  are reduced, i.e., they are affine algebraic varieties. Moreover, these varieties are irreducible, and their generic points correspond to simple representations;
  - (iii) Rep( $\Pi$ , **d**) is a complete intersection in Rep( $\overline{Q}$ , **d**).
  - (iv) The Poisson structure on  $M(Q, \mathbf{d})$  is generically symplectic.

One can pull-back polynomial functions via the moment map (11.2.1). This yields an algebra homomorphism  $\mu^* : \mathbb{k}[\mathfrak{g}_{\mathbf{d}}] \to \mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]$ , and we have

Corollary 11.3.2. Assume that  $\mathbf{d} \in \Sigma_0$ . Then the Poisson center of the algebra  $\mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]^{G_{\mathbf{d}}}$  equals  $\mu^* \left( \mathbb{k}[\mathfrak{g}_{\mathbf{d}}]^{G_{\mathbf{d}}} \right)$ , the pull-back of the algebra of  $\operatorname{Ad} G_{\mathbf{d}}$ -invariant polynomials on the Lie algebra  $\mathfrak{g}_{\mathbf{d}}$ .

Proof. The assumption on **d** insures that the moment map  $\mu : \operatorname{Rep}(\overline{Q}, \mathbf{d}) \to \mathfrak{g}_{\mathbf{d}}$  is flat, in particular, it is surjective. This follows from Theorem 11.3.1(iii), since the dimension of any fiber of  $\mu$  is not less the dimension of the zero fiber, and the latter is a complete intersection. Furthermore, all fibers of  $\mu$  are irreducible. To see this, let  $x \in \mathfrak{g}_{\mathbf{d}}$ . The standard increasing filtration on the algebra  $\mathbb{k}[\operatorname{Rep}(\overline{Q},\mathbf{d})]$ , by degree of the polynomial, induces an increasing filtration on the quotient algebra  $\mathbb{k}[\mu^{-1}(x)]$ . It is well known that, since  $\mu$  is flat and the zero fiber of  $\mu$  is reduced, for the corresponding associated graded algebra, one has  $\operatorname{gr} \mathbb{k}[\mu^{-1}(x)] \cong \mathbb{k}[\mu^{-1}(0)]$ . But, the scheme  $\mu^{-1}(0)$  is irreducible by Theorem 11.3.1(ii). It follows that the algebra  $\mathbb{k}[\mu^{-1}(0)]$ , hence also  $\mathbb{k}[\mu^{-1}(x)]$ , has no zero-divisors. Thus, the scheme  $\mu^{-1}(x)$  is irreducible as well.

Next, we claim that for any sufficiently general  $\operatorname{Ad} G_{\operatorname{\mathbf{d}}}$ -conjugacy class  $\operatorname{\mathsf{O}} \subset \mathfrak{g}_{\operatorname{\mathbf{d}}}$ , the preimage  $\mu^{-1}(\operatorname{\mathsf{O}})$  is a smooth submanifold in  $\operatorname{Rep}(\overline{Q},\operatorname{\mathbf{d}})$  and, moreover, the action of  $G_{\operatorname{\mathbf{d}}}$  on  $\mu^{-1}(\operatorname{\mathsf{O}})$  is free. Indeed, the former statement follows from the latter by general properties of moment maps. To prove the freeness, we let the conjugacy class  $\operatorname{\mathsf{O}}$  be such that any  $x \in \operatorname{\mathsf{O}}$  is a diagonalizable endomorphism, say  $x = \operatorname{diag}(s_1,\ldots,s_d)$ , and, in addition, such that for any proper subcollection  $S \subset \{s_1,\ldots,s_d\}$  one has  $0 \neq \sum_{s \in S} s$ . Let  $\rho \in \mu^{-1}(\operatorname{\mathsf{O}})$  be a representation of  $\overline{Q}$  and assume that  $E \subset \mathbb{k}^d$  is a nontrivial subrepresentation. Therefore, the vector space E is  $\rho(a)$ -stable for any endomorphism  $\rho(a), a \in \overline{Q}$ . Hence, we must have  $\operatorname{Tr} x|_E = \sum_{a \in Q} \operatorname{Tr}([\rho(a)|_E, \rho(a^*)|_E]) = 0$ , cf. (11.2.1). On the other hand, it is clear that E is the span of certain eigen-spaces of x and thus  $\operatorname{Tr} x|_E = \sum_{s \in S} s$ , for some subcollection  $S \subset \{s_1,\ldots,s_d\}$ , where  $|S| = \dim E$ . This contradicts our condition on the eigenvalues of x. We conclude that any point  $\rho \in \mu^{-1}(\operatorname{\mathsf{O}})$  is an irreducible representation of  $\overline{Q}$ , and our claim follows by the Schur lemma.

Thus, we have proved that for any sufficiently general Ad  $G_{\mathbf{d}}$ -conjugacy class  $O \subset \mathfrak{g}_{\mathbf{d}}$ , the preimage  $\mu^{-1}(O)$  is a smooth submanifold with a free  $G_{\mathbf{d}}$ -action. This implies that all  $G_{\mathbf{d}}$ -orbits in  $\mu^{-1}(O)$  are closed and correspond bijectively to points of the categorical quotient  $\mu^{-1}(O)/\!/G_{\mathbf{d}} = \operatorname{Spec} \mathbb{k}[\mu^{-1}(O)]^{G_{\mathbf{d}}}$ . Furthermore, the categorical quotient is a smooth affine symplectic algebraic variety.

Now, let F be a central element of the Poisson algebra  $\mathbb{k}[\operatorname{Rep}(\overline{Q},\mathbf{d})]^{G_{\mathbf{d}}}$ . Then, for any general conjugacy classes O, the restriction of F to  $\mu^{-1}(O)$  is a central element of the Poisson algebra  $\mathbb{k}[\mu^{-1}(O)]^{G_{\mathbf{d}}} = \mathbb{k}[\mu^{-1}(O)]/G_{\mathbf{d}}$ . The variety  $\mu^{-1}(O)//G_{\mathbf{d}}$  being symplectic, we deduce that the polynomial F must be constant on connected components of any general fiber of the moment map. Further, we know that any such fiber is irreducible, hence connected. Thus, F is constant on general fibers. We deduce, since  $\mathfrak{g}_{\mathbf{d}}$  is smooth, that  $F = \mu^*(\bar{F})$  for some  $\bar{F} \in \mathbb{k}[\mathfrak{g}_{\mathbf{d}}]$ . Finally, since F is  $G_{\mathbf{d}}$ -invariant, one may arrange that  $F \in \mu^*(\mathbb{k}[\mathfrak{g}_{\mathbf{d}}]^{G_{\mathbf{d}}})$ .

We will also use the following Lemma, which follows from [CB1]:

**Lemma 11.3.3.** Assume that Q is neither Dynkin nor extended Dynkin. Then for any integer  $N \geq 1$ , the set of vectors  $\mathbf{d} \in \Sigma_0$  such that  $d_i \geq N$  for all i is Zariski dense in kI.

*Proof.* According to Corollary 5.7 of [CB1],  $\Sigma_0$  is the set of all nonzero dimension vectors  $\mathbf{d}$  such that for any nonzero dimension vectors  $\alpha$ ,  $\beta$  such that  $\alpha + \beta = \mathbf{d}$ , one has  $(\alpha, A\beta) \leq -2$ .

Recall that the fundamental region is the set of nonzero dimension vectors **d** with connected support such that  $\sum_{i} a_{ij} d_{ij} \leq 0$  for all i.

Assume that **d** is in the fundamental region and  $(\mathbf{d}, A\mathbf{d}) < 0$ . Then for any nonzero dimension vectors  $\alpha, \beta$  such that  $\alpha + \beta = \mathbf{d}$ , by Lemma 8.2 of [CB1] one has  $(\alpha, A\beta) = (\mathbf{d} - \beta, A\beta) < 0$ . But it is clear from the definitions that  $(\beta, A\beta)$  is an *even* integer for any integral vector  $\beta$ . This implies that if in addition all  $d_i$  are even then  $(\alpha, A\beta) = (\mathbf{d} - \beta, A\beta) \le -2$ . Thus, if **d** is in the fundamental region,  $(\mathbf{d}, A\mathbf{d}) < 0$ , and  $d_i$  are even, then  $\mathbf{d} \in \Sigma_0$ .

Now consider the Perron-Frobenius eigenvector  $\mathbf{v} = (v_i)$  of the matrix C, with  $\sum v_i = 1$ . Since Q is not Dynkin and not affine, one has  $C\mathbf{v} = \lambda \mathbf{v}$ , where  $\lambda > 2$ . Thus  $(\mathbf{v}, A\mathbf{v}) = (2 - \lambda)|\mathbf{v}|^2 < 0$ , and  $\sum_j a_{ij}v_j = (2 - \lambda)v_i < 0$ . This means that there exists an open cone K in  $\mathbb{R}^I$  with axis of symmetry going through  $\mathbf{v}$  and such that any vector  $\mathbf{d} \in K$  with even integer coordinates  $\mathbf{d}_i$  is contained in  $\Sigma_0$ . This implies the lemma.

11.4 **Proof of Theorem 8.6.1.** Given an element  $x \in \operatorname{Sym} \mathbf{P}$ , we will denote by  $\bar{x}$  the image of that element under the projection  $\operatorname{Sym} \mathbf{P} \twoheadrightarrow \operatorname{Sym}(\mathbf{P}_{\flat})$ .

Let  $\mathbb{k}[w_{im}]$  be the polynomial algebra in an infinite set of variables  $w_{im}$ ,  $i \in I$ ,  $m = 1, 2, \ldots$  The assignment  $w_{im} \longmapsto e_i \mathbf{w}^m \in \mathbf{P}$  clearly induces a graded algebra isomorphism  $\operatorname{Sym} R \otimes \mathbb{k}[w_{im}] \stackrel{\sim}{\to} \operatorname{Sym}(R[\mathbf{w}])$ . Abusing the notation, we will often identify  $w_{im}$  with the corresponding element of  $\operatorname{Sym}(R[\mathbf{w}])$ .

Let **d** be a dimension vector and  $\psi_{\mathbf{d}} : \mathcal{O}(\mathbf{P}) \to \mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]^{G_{\mathbf{d}}}$  the corresponding algebra homomorphism. We define the following functions

$$\rho_{im} := \psi_{\mathbf{d}}(\bar{w}_{im}) = \widehat{\operatorname{Tr} e_i \mathbf{w}^m} \in \mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]^{G_{\mathbf{d}}}, \quad \forall i \in I, \ m = 1, 2, \dots$$

The relation between  $\mu$  and  $\mathbf{w}$  provided by Theorem 6.4.3 shows that the subalgebra  $\mu^*(\mathbb{k}[\mathfrak{g}_{\mathbf{d}}]) \subset \mathbb{k}[\operatorname{Rep}(\overline{Q}, \mathbf{d})]$  is generated by the matrix elements of

the matrix valued function  $\hat{\mathbf{w}}$ . Restricting attention to  $G_{\mathbf{d}}$ -invariants and using surjectivity of the homomorphism  $\psi_{\mathbf{d}}$ , we obtain

$$\mu^*(\mathbb{k}[\mathfrak{g}_{\mathbf{d}}]^{G_{\mathbf{d}}}) = \psi_{\mathbf{d}}(\operatorname{Sym}(R[\mathbf{w}])) = \mathbb{k}[\rho_{im}]. \tag{11.4.1}$$

We now prove part (i) of the Theorem. Let  $\bar{c} \in \operatorname{Sym}(\mathbf{P}_{\natural})$  be a homogeneous element of degree r > 0 which is a central element with respect to the Poisson bracket. Then, the element  $\psi_{\mathbf{d}}(\bar{c})$  has to be central in the Poisson algebra  $\mathbb{k}[\operatorname{Rep}(\overline{Q},\mathbf{d})]^{G_{\mathbf{d}}}$ , since the map  $\psi_{\mathbf{d}}$  is surjective. It follows from Corollary 11.3.2 that, for any  $\mathbf{d} \in \Sigma_0$ , we have  $\psi_{\mathbf{d}}(\bar{c}) \in \mu^*\left(\mathbb{k}[\mathfrak{g}_{\mathbf{d}}]^{G_{\mathbf{d}}}\right)$ . Hence, for such a  $\mathbf{d}$ , by (11.4.1), there exists a degree r homogeneous element  $z \in \mathbb{k}[\bar{w}_{im}]$  such that  $\psi_{\mathbf{d}}(\bar{c}) = \psi_{\mathbf{d}}(z)$ . Furthermore, if the dimension vector  $\mathbf{d} \in \Sigma_0$  is such that, in addition, one has  $d_i \geq N(r)$ , then, by Proposition 11.1.1, in  $\mathcal{O}_{\mathbf{d}}(\mathbf{P})$ , we get  $\bar{c} = z$ . Thus, we have shown that

$$\bar{c} \in \mathbb{k}[\bar{w}_{im}](r) \subset \mathcal{O}_{\mathbf{d}}(\mathbf{P})(r), \quad \forall \mathbf{d} \in \Sigma_0, \, d_i \ge N(r), i \in I.$$
 (11.4.2)

Next, identify Sym R with  $\mathbb{k}[R^*]$ , the polynomial algebra on the dual vector space  $R^* = \operatorname{Hom}_{\mathbb{k}}(R,\mathbb{k})$ . Any dimension vector may be viewed as a point  $\mathbf{d} \in R^*$ . Thus, for any  $f \in \operatorname{Sym} R = \mathbb{k}[R^*]$  there is a well-defined scalar  $f(\mathbf{d}) \in \mathbb{k}$ .

Now, write the direct sum decomposition  $\mathbf{P}_{\natural} = R \oplus \mathbf{L}$ , where  $\mathbf{L}$  is the sum of all homogeneous components of positive degrees. Thus, we have an algebra isomorphism  $\operatorname{Sym} R \otimes \operatorname{Sym} \mathbf{L} \xrightarrow{\sim} \operatorname{Sym} \mathbf{P}_{\natural}$ , and we can write our element  $\bar{c} \in \operatorname{Sym} \mathbf{P}_{\natural}$  in the form  $\bar{c} = \sum_{j=1}^{l} f_j \cdot \bar{c}_j$ , where  $f_j$  are elements of  $\operatorname{Sym} R$  and  $\bar{c}_j$  are linearly independent elements of  $\operatorname{Sym} \mathbf{L}$ . With this notation, the image of  $\bar{c}$  in  $\mathcal{O}_{\mathbf{d}}(\mathbf{P})$  equals  $\sum_{j=1}^{l} f_j(\mathbf{d}) \cdot \bar{c}_j$ .

Recall next that the imbedding  $\operatorname{Sym} \mathbf{L} \hookrightarrow \operatorname{Sym} \mathbf{P}_{\natural}$  induces an isomorphism  $\operatorname{Sym} \mathbf{L} \overset{\sim}{\to} \mathcal{O}_{\mathbf{d}}(\mathbf{P})$ . We see that formula (11.4.2) may be intrpreted as saying that the polynomial map  $R^* \to \operatorname{Sym} \mathbf{L}$ ,  $\mathbf{d} \longmapsto \sum_{j=1}^l f_j(\mathbf{d}) \cdot \bar{c}_j$ , takes values in the subspace  $\mathbb{k}[\bar{w}_{im}](r)$ , for every  $\mathbf{d} \in \Sigma_0$  such that  $d_i \geq N(r), \forall i \in I$ . By Lemma 11.3.3, this set of dimension vectors is Zariski dense. Hence, we deduce that  $\sum_{j=1}^l f_j(\mathbf{d}) \cdot \bar{c}_j \in \mathbb{k}[\bar{w}_{im}](r)$  for all  $\mathbf{d} \in R^*$ , which means that  $\sum_{j=1}^l f_j \cdot \bar{c}_j \in \operatorname{Sym} R \otimes \mathbb{k}[\bar{w}_{im}] = \operatorname{Sym}(\overline{R[\mathbf{w}]})$ . This completes the proof of the first statement of part (i) of the theorem.

To prove the second statement of part (i), let  $c \in \text{Sym}(R[\mathbf{w}])$  be a nonzero homogeneous element of degree r > 0 such that its image,  $\bar{c} \in \text{Sym}(\overline{R[\mathbf{w}]})$ , vanishes. We can write  $c = \sum_j f_j \cdot p_j(w_{im})$  where  $f_j \in \text{Sym } R$  and where  $p_j = p_j(w_{im}) \in \mathbb{k}[w_{im}]$  are some homogeneous linearly independent elements of positive degrees. Then, since  $\bar{c} = 0$ , applying the homomorphism  $\psi_{\mathbf{d}}$ , in  $\mathbb{k}[\text{Rep}(\overline{Q}, \mathbf{d})]$ , we find

$$0 = \psi_{\mathbf{d}}(\bar{c}) = \sum_{j} f_{j}(\mathbf{d}) \cdot p(\rho_{im}).$$

Now, let  $\mathbf{d} \in \Sigma_0$  be such that  $d_i \geq N(r), \forall i \in I$ . Then the only independent relation between the functions  $\rho_{im} \in \text{Rep}(\overline{Q}, \mathbf{d})$  with m < N is the relation  $\sum_i \rho_{i1} = \psi_{\mathbf{d}}(\mathbf{w}) = 0$ . Thus,  $\sum_j f_j(\mathbf{d}) \cdot p_j$ , thought of as an element of an abstract

polynomial algebra  $\mathbb{k}[w_{im}]$ , must belong to the principal ideal generated by the element  $\sum_i w_{i1}$ . Now, Lemma 11.3.3 and the linear independence of the polynomials  $p_j = p_j(w_{im})$  allows us to conclude that the element  $\sum_j f_j \otimes p_j \in \operatorname{Sym} R \otimes \mathbb{k}[w_{im}]$  is divisible by  $\sum_i w_{i1}$ . Part (i) of the theorem follows.

Now we prove part (ii) of the theorem by a very similar argument.

Let  $\bar{c}$  be a homogeneous central element of Sym  $\Pi_{\natural}$  of degree r > 0. The map  $\psi_{\mathbf{d}} : \mathcal{O}_{\mathbf{d}}(\Pi) \to \mathbb{k}[\operatorname{Rep}(\Pi, \mathbf{d})]^{G_{\mathbf{d}}}$  being surjective, we conclude that the element  $\bar{c}$  has to be a central element in the Poisson algebra  $\mathbb{k}[M(Q, \mathbf{d})]$ . By Theorem 11.3.1, if  $\mathbf{d} \in \Sigma_0$ , this scheme is in fact an irreducible algebraic variety which is, moreover, generically symplectic. Therefore, any Casimir on  $M(Q, \mathbf{d})$  has to be a constant function, and hence  $\psi_{\mathbf{d}}(\bar{c}) = 0$ .

Write  $\Pi_{\natural} = R \oplus L$ , where L is the positive degree part of  $\Pi_{\natural}$ . We can write  $\bar{c}$  in the form  $\bar{c} = \sum_{j=1}^{l} f_j \cdot \bar{c}_j$ , where  $f_j$  are elements of  $\operatorname{Sym} R$  and  $\bar{c}_j$  are linearly independent elements of  $\operatorname{Sym} L$ . Again, applying the homomorphism  $\psi_{\mathbf{d}}$  in our  $\Pi$ -setting, we get  $0 = \psi_{\mathbf{d}}(\bar{c}) = \sum_{j=1}^{l} f_j(\mathbf{d}) \cdot \psi_{\mathbf{d}}(\bar{c}_j)$ .

Let  $d_i \geq N(r)$  for all i. Then, by Proposition 11.1.1, we deduce that

Let  $d_i \geq N(r)$  for all i. Then, by Proposition 11.1.1, we deduce that  $\sum_j f_j(\mathbf{d}) \cdot \bar{c}_j = 0$ . This implies that  $f_j(\mathbf{d}) = 0$  for any j and any  $\mathbf{d} \in \Sigma_0$  such that  $d_i \geq N(r)$ . We know from Lemma 11.3.3 that the set of such  $\mathbf{d}$  is Zariski dense. Thus the polynomials  $f_j$  are all identically zero, i.e.  $\bar{c} = 0$ , which proves (ii).

# 11.5 The center of $\Pi$ . In this section we will prove Proposition 8.2.2. The argument is based on Lemma 11.3.3.

Let z be a central element of  $\Pi$ . To show that z is a constant it suffices to show that this is so in every finite dimensional representation of  $\Pi$ . Indeed, let  $\mathscr{J}_N$  be the ideal of elements of degree  $\geq N$  in  $\Pi$ ; then any element of  $\Pi$  acting by a scalar in  $\Pi/\mathscr{J}^N$  for all N is necessarily a scalar.

Let Y be a finite dimensional representation of  $\Pi$  with dimension vector d. We want to show that z acts by a scalar in Y. By Lemma 11.3.3, there exists a nonnegative dimension vector d' such that  $d+d' \in \Sigma_0$  (stable region). Let Y' be the direct sum of Y with the augmentation representation of dimension d'. Then  $Y' \in \text{Rep}(\bar{Q}, d+d')$ , which is an irreducible variety by [CB2]. So it suffices to show that z acts by a scalar on the generic representation of dimension d+d'. But the generic representation is irreducible, again by [CB2]. So we are done by Schur's lemma.

We now prove similarly that the algebra  $\Pi$  is prime. Suppose  $a, b \in \Pi$  are nonzero, but  $a\Pi b = 0$ . Choose a dimension vector  $\mathbf{d}$  such that there are representations of dimension  $\mathbf{d}$  which are not annihilated by a and representations which are not annihilated by b. By enlarging d if necessary we may assume that  $\mathbf{d} \in \Sigma_0$ . Let  $R_a$  and  $R_b$  be the closed subsets consisting of the representations in  $\operatorname{Rep}(\Pi, \mathbf{d})$  annihilated by a and b respectively. By assumption they are proper subsets. If X is a simple representation of  $\Pi$ , then  $a\Pi bX = 0$ , but  $\Pi bX = 0$  or X, so aX = 0 or bX = 0. Thus  $R_a \cup R_b$  contains all simple representations, and since they are dense in  $\operatorname{Rep}(\Pi, \mathbf{d})$  we have  $R_a \cup R_b = \operatorname{Rep}(\Pi, \mathbf{d})$ . But this contradicts the fact that  $\operatorname{Rep}(\Pi, \mathbf{d})$  is irreducible.

# References

- [BG] A. Braverman, and D. Gaitsgory, Poincaré-Birkhoff-Witt theorem for quadratic algebras of Koszul type. J. Algebra 181 (1996), 315–328.
- [BLB] R. Bocklandt, L. Le Bruyn, Necklace Lie algebras and noncommutative symplectic geometry. Math. Z. 240 (2002), 141–167.
- [BBK] S. Brenner, M. Butler, and A. King, Periodic algebras which are almost Koszul. Algebr. Represent. Theory 5 (2002), 331–367.
- [CB1] W. Crawley-Boevey, Geometry of the moment map for representations of quivers. Compositio Math. 126 (2001), 257–293.
- [CB2] W. Crawley-Boevey, Preprojective algebras, differential operators and a Conze embedding for deformations of Kleinian singularities. Comment. Math. Helv. 74 (1999), 548–574.
- [CBH] W. Crawley-Boevey, M. Holland: *Noncommutative deformations of Kleinian singularities*. Duke Math. J. **92** (1998), 605–635.
- [CQ] J. Cuntz, D. Quillen, Algebra extensions and nonsingularity. J. Amer. Math. Soc. 8 (1995), 251–289.
- [Do] I. Dorfman, Dirac structures and integrability of nonlinear evolution equations. Nonlinear Science: Theory and Applications. John Wiley & Sons, Ltd., Chichester, 1993.
- [E] P. Etingof, Casimirs of the Goldman Lie algebra of a closed surface. Preprint arXiv:math.QA/0407505
- [EM] P. Etingof, S. Montarani, Finite dimensional representations of symplectic reflection algebras associated to wreath products. Represent. Theory 9 (2005), 457–467. arXiv:math.RT/0403250.
- [EO] P. Etingof, V. Ostrik, Module categories over representations of  $SL_q(2)$  and graphs. Math. Res. Lett. 11 (2004), 103–114. arXiv:math.QA/0302130.
- [ES] K. Erdmann, N. Snashall, On Hochschild cohomology of preprojective algebras. I, II. J. Algebra 205 (1998), 391–412, 413–434.
- [GD] I. Gelfand, I. Dorfman, Hamiltonian operators and algebraic structures associated with them. (Russian) Funktsional. Anal. i Prilozhen. 13: 4 (1979), 13–30. English transl.: Funct. Anal. Appl. 13, 13–30 (1979).
- [GDT] I. Gelfand, Yu. Daletskiy, B. Tsygan, On a variant of noncommutative differential geometry. Soviet Math. Dokl. 40 (1990), 422–426.
- [Gi] V. Ginzburg, Non-commutative symplectic geometry, quiver varieties, and operads. Math. Res. Lett. 8 (2001), 377–400.
- [Gi2] V. Ginzburg, *Double derivations and cyclic homology*. arXiv:math.KT/0505236.
- [Gi3] V. Ginzburg, Lectures on Noncommutative geometry. arXiv:math.AG/0506603.
- [Ko] M. Kontsevich, Formal (non)commutative symplectic geometry. The Gelfand Mathematical Seminars, 1990–1992, 173–187, Birkhäuser Boston, 1993.
- [KR] M. Kontsevich, A. Rosenberg, Noncommutative smooth spaces. The Gelfand Mathematical Seminars, 1996–1999, 85–108, Birkhäuser Boston, 2000.

- [LBP] L. Le Bruyn, C. Procesi, Semisimple representations of quivers. Trans. Amer. Math. Soc. 317 (1990), 585–598.
- [La] C. I. Lazaroiu, On the non-commutative geometry of topological D-branes. arXiv:hep-th/0507222.
- [Lo] J.-L. Loday, Cyclic homology. Grundlehren der Mathematischen Wissenschaften, 301. Springer-Verlag, Berlin, 1998.
- [MV] R. Martínez-Villa, Applications of Koszul algebras: the preprojective algebra, in: Representation theory of algebras (Cocoyoc, 1994), 487–504, CMS Conf. Proc., 18, Amer. Math. Soc., Providence, RI, 1996.
- [MOV] A. Malkin, V. Ostrik, M. Vybornov, Quiver varieties and Lusztig's algebra. (to appear in Adv. Math.) arXiv:math.RT/0403222.
- [Sch] V. Schechtman, Remarks on formal deformations and Batalin-Vilkovisky algebras. arXiv:math.AG/9802006.
- [VB1] M. Van den Bergh, A relation between Hochschild homology and cohomology for Gorenstein rings. Proc. Amer. Math. Soc. 126 (1998), 1345–1348; Correction: Proc. Amer. Math. Soc. 130 (2002), 2809-2810.
- [VB2] M. Van den Bergh, Double Poisson algebras. arXiv:math.QA/0410528.
- [Vo] D. Voiculescu, A note on cyclic gradients. Indiana Univ. Math. J. 49 (2000), 837-841. arXiv:math.RA/0005045.
  - W. C-B.: Department of Pure Mathematics, University of Leeds, Leeds LS2 9JT, UK; W.Crawley-Boevey@leeds.ac.uk
  - P.E.: Department of Mathematics, Rm 2-165, MIT, 77 Mass. Ave, Cambridge, MA 02139, USA; etingof@math.mit.edu
  - V.G.: Department of Mathematics, University of Chicago, Chicago, IL 60637, USA; ginzburg@math.uchicago.edu