UNIFORM LINEAR BOUND IN CHEVALLEY'S LEMMA

J. ADAMUS, E. BIERSTONE AND P.D. MILMAN

ABSTRACT. We obtain a uniform linear bound for the Chevalley function at a point in the source of an analytic mapping that is regular in the sense of Gabrielov. There is a version of Chevalley's lemma also along a fibre, or at a point of the image of a proper analytic mapping. We get a uniform linear bound for the Chevalley function for a closed Nash (or formally Nash) subanalytic set.

Contents

1.	Introduction	1
2.	Techniques	3
3.	Ideals of relations and Chevalley functions	6
4.	Proofs of the main theorems	8
References		11

1. Introduction

Let $\varphi: M \to N$ denote an analytic mapping of analytic manifolds (over $\mathbb{K} = \mathbb{R}$ or \mathbb{C}). Let $a \in M$. Let $\varphi_a^*: \mathcal{O}_{\varphi(a)} \to \mathcal{O}_a$ or $\hat{\varphi}_a^*: \widehat{\mathcal{O}}_{\varphi(a)} \to \widehat{\mathcal{O}}_a$ denote the induced homorphisms of analytic local rings or their completions, respectively. (We write $\mathcal{O}_a = \mathcal{O}_{M,a}$, and \mathfrak{m}_a (or $\widehat{\mathfrak{m}}_a$) = maximal ideal of \mathcal{O}_a (or $\widehat{\mathcal{O}}_a$).) According to Chevalley's lemma (1943), there is an increasing function $l: \mathbb{N} \to \mathbb{N}$ (where \mathbb{N} denotes the nonnegative integers) such that

$$\hat{\varphi}_a^*(\widehat{\mathcal{O}}_{\varphi(a)}) \cap \widehat{\mathfrak{m}}_a^{l(k)+1} \ \subset \ \hat{\varphi}_a^*(\widehat{\mathfrak{m}}_{\varphi(a)}^{k+1}) \ ;$$

i.e., if $F \in \widehat{\mathcal{O}}_{\varphi(a)}$ and $\widehat{\varphi}_a^*(F)$ vanishes to order l(k), then F vanishes to order k, modulo an element of Ker $\widehat{\varphi}_a^*$ ([4]; cf. Lemma 3.2 below). Let $l_{\varphi^*}(a,k)$ denote the least l(k) satisfying Chevalley's lemma. We call $l_{\varphi^*}(a,k)$ the Chevalley function of $\widehat{\varphi}_a^*$.

Let $x=(x_1,\ldots,x_m)$ and $y=(y_1,\ldots,y_n)$ denote local coordinate systems for M and N at a and $\varphi(a)$, respectively. The local rings \mathcal{O}_a or $\widehat{\mathcal{O}}_a$ can be identified with the rings of convergent or formal power series $\mathbb{K}\{x\}=\mathbb{K}\{x_1,\ldots,x_m\}$ or $\mathbb{K}[\![x]\!]=\mathbb{K}[\![x_1,\ldots,x_m]\!]$, respectively. In the local coordinates, write $\varphi(x)=(\varphi_1(x),\ldots,\varphi_n(x))$. Then $\ker \widehat{\varphi}_a^*$ is the ideal of formal relations $\{F(y)\in\mathbb{K}[\![y]\!]:F(\varphi_1(x),\ldots,\varphi_n(x))=0\}$ (and $\ker \varphi_a^*$ is the analogous ideal of analytic relations).

Key words and phrases. Chevalley function, regular mapping, Nash subanalytic set. Research partially supported by NSERC Postdoctoral Fellowship PDF 267954-2003 (Adamus), and NSERC Discovery Grants OGP 0009070 (Bierstone), OGP 0008949 (Milman).

Chevalley's lemma is an analogue for such nonlinear relations of the Artin-Rees lemma. (See Remark 1.4.)

Let $r_a^1(\varphi)$ denote the generic rank of φ near a, and set

$$r_a^2(\varphi) := \dim \frac{\widehat{\mathcal{O}}_{\varphi(a)}}{\operatorname{Ker} \hat{\varphi}_a^*} , \qquad r_a^3(\varphi) := \dim \frac{\mathcal{O}_{\varphi(a)}}{\operatorname{Ker} \varphi_a^*}$$

(where dim denotes the Krull dimension). Then $r_a^1(\varphi) \leq r_a^2(\varphi) \leq r_a^3(\varphi)$. Gabrielov proved that if $r_a^1(\varphi) = r_a^2(\varphi)$, then $r_a^2(\varphi) = r_a^3(\varphi)$ [6]; i.e., if there are enough formal relations, then the ideal of formal relations is generated by convergent relations. The mapping φ is called regular at a if $r_a^1(\varphi) = r_a^3(\varphi)$. We say that φ is regular if it is regular at every point of M. Izumi [10] proved that φ is regular at a if and only if the Chevalley function of $\hat{\varphi}_a^*$ has a linear (upper) bound; i.e., there exist $\alpha, \beta \in \mathbb{N}$ such that

$$l_{\omega^*}(a,k) < \alpha k + \beta$$
,

for all $k \in \mathbb{N}$. On the other hand, Bierstone and Milman [2] proved that, if φ is regular, then $l_{\varphi^*}(a,k)$ has a *uniform bound*; i.e., for every compact $L \subset M$, there exists $l_L : \mathbb{N} \to \mathbb{N}$ such that

$$l_{\varphi^*}(a,k) \leq l_L(k)$$
,

for all $a \in L$ and $k \in \mathbb{N}$. In this article, we prove that the Chevalley function associated to a regular mapping has a *uniform linear bound*:

Theorem 1.1. Suppose that φ is regular. Then, for every compact $L \subset M$, there exist $\alpha_L, \beta_L \in \mathbb{N}$ such that

$$l_{\omega^*}(a,k) < \alpha_L k + \beta_L$$

for all $a \in L$ and $k \in \mathbb{N}$.

Chevalley's lemma can be used also to compare two notions of order of vanishing of a real-analytic function at a point of a subanalytic set. Let X denote a closed subanalytic subset of \mathbb{R}^n . Let $b \in X$ and let $\mathcal{F}_b(X) \subset \mathbb{R}[y-b]$ denote the formal local ideal of X at b. (See Lemma 3.6.) For all $F \in \widehat{\mathcal{O}}_b = \mathbb{R}[y-b]$, we define

(1.1)
$$\mu_{X,b}(F) := \max\{l \in \mathbb{N} : |T_b^l F(y)| \le \operatorname{const} |y - b|^l, \ y \in X\}, \\ \nu_{X,b}(F) := \max\{l \in \mathbb{N} : F \in \widehat{\mathfrak{m}}_b^l + \mathcal{F}_b(X)\},$$

where $T_b^l F(y)$ denotes the Taylor polynomial of order l of F at b. Then there exists $l: \mathbb{N} \to \mathbb{N}$ such that, for all $k \in \mathbb{N}$, if $F \in \widehat{\mathcal{O}}_b$ and $\mu_{X,b}(F) > l(k)$, then $\nu_{X,b}(F) > k$. (See Section 3.) For each k, let $l_X(b,k)$ denote the least such l(k). We call $l_X(b,k)$ the Chevalley function of X at b.

Theorem 1.2. Suppose that X is a Nash (or formally Nash) subanalytic subset of \mathbb{R}^n . Then the Chevalley function of X has a uniform linear bound; i.e., for every compact $K \subset X$, there exists $\alpha_K, \beta_K \in \mathbb{N}$ such that

$$l_X(b,k) \leq \alpha_K k + \beta_K$$

for all $b \in K$ and $k \in \mathbb{N}$.

Theorems 1.1 and 1.2 are the main new results in this article. They answer questions raised in [3, 1.28].

The closed Nash subanalytic subsets X of \mathbb{R}^n are the images of regular proper real-analytic mappings $\varphi: M \to \mathbb{R}^n$. In particular, a closed semianalytic set is

Nash. A closed subanalytic subset X of \mathbb{R}^n is formally Nash if, for every $b \in X$, there is a closed Nash subanalytic subset Y of X such that $\mathcal{F}_b(X) = \mathcal{F}_b(Y)$ [3]. Unlike the situation of Theorem 1.1, the converse of Theorem 1.2 is false [3, Example 12.8].

The main theorem of [3] (Theorem 1.13) asserts that, if X is a closed subanalytic subset of \mathbb{R}^n , then the existence of a uniform bound for $l_X(b,k)$ is equivalent to several other natural analytic and algebro-geometric conditions; for example, semi-coherence [3, Definition 1.2], stratification by the diagram of initial exponents of the ideal $\mathcal{F}_b(X)$, $b \in X$ [3, Theorem 8.1], and a \mathcal{C}^{∞} composite function property [3, §1.5]. A uniform bound for the Chevalley function measures loss of differentiability in a \mathcal{C}^r version of the composite function theorem. We use the techniques of [3] to prove Theorems 1.1 and 1.2 here.

Wang [12, Theorem 1.1] used [9, Theorem 1.2] to prove that the Chevalley function associated to a regular proper real-analytic mapping $\varphi \colon M \to \mathbb{R}^n$ has a uniform linear bound if and only if $X = \varphi(M)$ has a uniform linear product estimate; i.e., for every compact $K \subset X$, there exist $\alpha_K, \beta_K \in \mathbb{N}$ such that, for all $b \in K$ and $F, G \in \widehat{\mathcal{O}}_b$,

$$\nu_{X_i,b}(F \cdot G) \leq \alpha_K(\nu_{X_i,b}(F) + \nu_{X_i,b}(G)) + \beta_K ,$$

where $X_b = \bigcup_i X_i$ is a decomposition of the germ X_b into finitely many irreducible subanalytic components. We therefore obtain the following from Theorem 1.1:

Theorem 1.3. A closed Nash subanalytic subset of \mathbb{R}^n admits a uniform linear product estimate.

Remark 1.4. The Artin-Rees lemma can be viewed as a version of Chevalley's lemma for linear relations over a Noetherian ring R: Suppose that $\Psi: E \to G$ is a homomorphism of finitely-generated modules over R, and let $F \subset G$ denote the image of Ψ . Let \mathfrak{m} be a maximal ideal of R. Then $F \cap \mathfrak{m}^l G \subset \mathfrak{m}^k F$ if and only if $\Psi^{-1}(\mathfrak{m}^l G) \subset \operatorname{Ker} \Psi + \mathfrak{m}^k E$. The Artin-Rees lemma says that there exists $\beta \in \mathbb{N}$ such that $F \cap \mathfrak{m}^{k+\beta}G = \mathfrak{m}^k(F \cap \mathfrak{m}^\beta G)$, for all k. In particular, there is always a linear Artin-Rees exponent $l(k) = k + \beta$. Uniform versions of the Artin-Rees lemma were proved in [2, Theorem 7.4], [5], [8]. A uniform Artin-Rees exponent for a homomorphism of \mathcal{O}_M -modules, where M is a real-analytic manifold, measures loss of differentiability in Malgrange division, in the same way that a uniform bound for the Chevalley function relates to composite differentiable functions. (See [2].)

2. Techniques

2.1. **Linear algebra lemma.** Let R denote a commutative ring with identity, and let E and F be R-modules. If $B \in \operatorname{Hom}_R(E, F)$ and $r \in \mathbb{N}$, $r \geq 1$, we define

$$\operatorname{ad}{}^r B \ \in \ \operatorname{Hom}_R \left(F, \operatorname{Hom}_R \left(\bigwedge^r E, \bigwedge^{r+1} F \right) \right)$$

by the formula

$$(\operatorname{ad}^{r}B)(\omega)(\eta_{1}\wedge\cdots\wedge\eta_{r}) = \omega\wedge B\eta_{1}\wedge\cdots\wedge B\eta_{r} ,$$

where $\omega \in F$ and $\eta_1, \ldots, \eta_r \in E$. (ad ${}^0B := \operatorname{id}_F$, the identity mapping of F.) Clearly, if $r > \operatorname{rk} B$ then ad ${}^rB = 0$, and if $r = \operatorname{rk} B$ then ad ${}^rB \cdot B = 0$. (rk B means the smallest r such that $\bigwedge^s B = 0$ for all s > r.) If R is a field, then rk $B = \dim \operatorname{Im} B$, so we get:

Lemma 2.1 ([1, §6]). Let E and F be finite-dimensional vector spaces over a field \mathbb{K} . If $B: E \to F$ is a linear transformation and $r = \operatorname{rk} B$, then

$$\operatorname{Im} B = \operatorname{Ker} \operatorname{ad}^r B.$$

In particular, if A is another linear transformation with target F, then $A\xi + B\eta = 0$ (for some η) if and only if $\xi \in \text{Ker ad }^r B \cdot A$.

2.2. The diagram of initial exponents. Let A be a commutative ring with identity. Consider the total ordering of \mathbb{N}^n given by the lexicographic ordering of (n+1)-tuples $(|\beta|, \beta_1, \ldots, \beta_n)$, where $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{N}^n$ and $|\beta| = \beta_1 + \cdots + \beta_n$. For any formal power series $F(Y) = \sum_{\beta \in \mathbb{N}^n} F_{\beta} Y^{\beta} \in A[\![Y]\!] = A[\![Y_1, \ldots, Y_n]\!]$, we define the support supp $F := \{\beta \in \mathbb{N}^n : F_{\beta} \neq 0\}$ and the initial exponent $\exp F := \min \sup F$. ($\exp F := \infty \text{ if } F = 0$.)

Let I be an ideal in A[Y]. The diagram of initial exponents of I is defined as

$$\mathfrak{N}(I) := \{ \exp F \colon F \in I \setminus \{0\} \} \ .$$

Clearly, $\mathfrak{N}(I) + \mathbb{N}^n = \mathfrak{N}(I)$.

Suppose that A is a field \mathbb{K} . Then, by the formal division theorem of Hironaka [7] (see [2, Theorem 6.2]),

$$(2.1) \mathbb{K}\llbracket Y \rrbracket = I \oplus \mathbb{K}\llbracket Y \rrbracket^{\mathfrak{N}(I)},$$

where $\mathbb{K}[\![Y]\!]^{\mathfrak{N}}$ is defined as $\{F \in \mathbb{K}[\![Y]\!] : \operatorname{supp} F \subset \mathbb{N}^n \setminus \mathfrak{N}\}$, for any $\mathfrak{N} \in \mathbb{N}^n$ such that $\mathfrak{N} + \mathbb{N}^n = \mathfrak{N}$.

2.3. **Fibred product.** Let M denote an analytic manifold over \mathbb{K} , and let $s \in \mathbb{N}$, $s \geq 1$. Let $\varphi \colon M \to N$ be an analytic mapping. We denote by M_{φ}^s the s-fold fibred product of M with itself over N; i.e.,

$$M_{\varphi}^s := \{\underline{a} = (a^1, \dots, a^s) \in M^s \colon \varphi(a^1) = \dots = \varphi(a^s)\};$$

 M_{φ}^{s} is a closed analytic subset of M^{s} . There is a natural mapping $\underline{\varphi} = \underline{\varphi}^{s}$: $M_{\varphi}^{s} \to N$ given by $\underline{\varphi}(\underline{a}) = \varphi(a^{1})$; i.e., for each $i = 1, \ldots, s, \ \underline{\varphi} = \varphi \circ \rho^{i}$, where $\rho^{i} \colon M_{\varphi}^{s} \ni (x^{1}, \ldots, x^{s}) \mapsto x^{i} \in M$.

Suppose that $\mathbb{K}=\mathbb{R}$. Let E be a closed subanalytic subset of M, and let $\varphi\colon E\to\mathbb{R}^n$ be a continuous subanalytic mapping. Then the fibred product E^s_φ is a closed subanalytic subset of M^s , and the canonical mapping $\underline{\varphi}=\underline{\varphi}^s\colon E^s_\varphi\to\mathbb{R}^n$ is subanalytic.

Let \mathring{E}^s_{φ} denote the subset of E^s_{φ} consisting of points $\underline{x} = (x^1, \dots, x^s) \in E^s_{\varphi}$ such that each x^i lies in a distinct connected component of the fibre $\varphi^{-1}(\underline{\varphi}(\underline{x}))$. If φ is proper, then \mathring{E}^s_{φ} is a subanalytic subset of M^s [3, §7].

2.4. **Jets.** Let N denote an analytic manifold (over $\mathbb{K} = \mathbb{R}$ or \mathbb{C}), and let $b \in N$. Let $l \in \mathbb{N}$ and let $J^l(b)$ denote $\widehat{\mathcal{O}}_b/\widehat{\mathfrak{m}}_b^{l+1}$. If $F \in \widehat{\mathcal{O}}_b$, then $J^lF(b)$ denotes the image of F in $J^l(b)$. Let M be an analytic manifold, and let $\varphi \colon M \to N$ be an analytic mapping. If $a \in \varphi^{-1}(b)$, then the homomorphism $\widehat{\varphi}_a^* \colon \widehat{\mathcal{O}}_b \to \widehat{\mathcal{O}}_a$ induces a linear transformation $J^l\varphi(a) \colon J^l(b) \to J^l(a)$.

Suppose that $N = \mathbb{K}^n$. Let $y = (y_1, \dots, y_n)$ denote the affine coordinates of \mathbb{K}^n . Taylor series expansion induces an identification of $\widehat{\mathcal{O}}_b$ with the ring of formal power series $\mathbb{K}[y-b] = \mathbb{K}[y_1-b_1, \dots, y_n-b_n]$ (we write $F(y) = \sum_{\beta \in \mathbb{N}^n} F_{\beta}(y-b)^{\beta}$), and hence an identification of $J^l(b)$ with \mathbb{K}^q , $q = \binom{n+l}{l}$, with respect to which

 $J^l F(b) = (D^{\beta} F(b))_{|\beta| \leq l}$, where D^{β} denotes $1/\beta!$ times the formal derivative of order $\beta \in \mathbb{N}$.

Using a system of coordinates $x = (x_1, \ldots, x_m)$ for M in a neighbourhood of a, we can identify $J^l(a)$ with \mathbb{K}^p , $p = \binom{m+l}{l}$. Then

$$J^{l}\varphi(a) \colon (F_{\beta})_{|\beta| \leq l} \mapsto ((\hat{\varphi}_{a}^{*}(F))_{\alpha})_{|\alpha| \leq l} = \left(\sum_{|\beta| \leq l} F_{\beta} L_{\alpha}^{\beta}(a)\right)_{|\alpha| \leq l},$$

where $L_{\alpha}^{\beta}(a) = (\partial^{|\alpha|}\varphi^{\beta}/\partial x^{\alpha})(a)/\alpha!$ and $\varphi^{\beta} = \varphi_{1}^{\beta_{1}} \dots \varphi_{n}^{\beta_{n}} \ (\varphi = (\varphi_{1}, \dots, \varphi_{n})).$ Set $J_{b}^{l} := J^{l}(b) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{b} = \bigoplus_{|\beta| \leq l} \mathbb{K}[y - b].$ We put $J_{b}^{l}F(y) := (D^{\beta}F(y))_{|\beta| \leq l} \in J_{b}^{l}.$ (Evaluating at b transforms $J_{b}^{l}F$ to $J^{l}F(b)$.) The ring homomorphism $\hat{\varphi}_{a}^{*} : \widehat{\mathcal{O}}_{b} \to \widehat{\mathcal{O}}_{a}$ induces a homomorphism of $\mathbb{K}[x - a]$ -modules,

$$J_a^l \varphi \colon \quad J^l(b) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_a \qquad \longrightarrow \qquad J^l(a) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_a$$

$$\parallel \qquad \qquad \parallel$$

$$\bigoplus_{|\beta| \leq l} \mathbb{K} \llbracket x - a \rrbracket \qquad \qquad \bigoplus_{|\alpha| \leq l} \mathbb{K} \llbracket x - a \rrbracket$$

such that, if $F \in \widehat{\mathcal{O}}_b$, then

$$J_a^l \varphi \left((\hat{\varphi}_a^*(D^{\beta}F))_{|\beta| < l} \right) = (D^{\alpha}(\hat{\varphi}_a^*(F)))_{|\alpha| < l}.$$

By evaluation at a, $J_a^l \varphi$ induces $J^l \varphi(a) \colon J^l(b) \to J^l(a)$. $J_a^l \varphi$ identifies with the matrix (with rows indexed by $\alpha \in \mathbb{N}^m$, $|\alpha| \le l$, and columns indexed by $\beta \in \mathbb{N}^n$, $|\beta| \le l$) whose entries are the Taylor expansions at a of the $D^\alpha \varphi^\beta = (\partial^{|\alpha|} \varphi^\beta / \partial x^\alpha) / \alpha!$, $|\alpha| \le l$, $|\beta| \le l$.

Let $\underline{a}=(a^1,\ldots,a^s)\in M_\varphi^s$ and let $b=\underline{\varphi}(\underline{a})$. For each $i=1,\ldots,s$, the homomorphism $J_b^l=J^l(b)\otimes_{\mathbb K}\widehat{\mathcal O}_b\to J^l(a^i)\otimes_{\mathbb K}\widehat{\mathcal O}_{a^i}=J_{a^i}^l$ over $\widehat{\varphi}_{a^i}^*$, as defined above (using a coordinate system $x^i=(x_1^i,\ldots,x_m^i)$ for M in a neighbourhood of a^i), followed by the canonical homomorphism $J^l(a^i)\otimes_{\mathbb K}\widehat{\mathcal O}_{a^i}\to J^l(a^i)\otimes_{\mathbb K}\widehat{\mathcal O}_{M_\varphi^s,\underline{a}}$ over $(\widehat{\rho^i})_{\underline{a}}^*\colon\widehat{\mathcal O}_{a^i}\to\widehat{\mathcal O}_{M_\varphi^s,\underline{a}}$, induces an $\widehat{\mathcal O}_{M_\varphi^s,\underline{a}}$ -homomorphism $J^l(b)\otimes_{\mathbb K}\widehat{\mathcal O}_{M_\varphi^s,\underline{a}}\to J^l(a^i)\otimes_{\mathbb K}\widehat{\mathcal O}_{M_\varphi^s,\underline{a}}$. We thus obtain an $\widehat{\mathcal O}_{M_\varphi^s,\underline{a}}$ -homomorphism

$$J^{l}_{\underline{a}}\varphi \colon \quad J^{l}(b) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{M_{\varphi}^{s},\underline{a}} \quad \longrightarrow \quad \bigoplus_{i=1}^{s} J^{l}(a^{i}) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{M_{\varphi}^{s},\underline{a}}$$

$$\parallel \qquad \qquad \parallel$$

$$\bigoplus_{|\beta| \leq l} \widehat{\mathcal{O}}_{M_{\varphi}^{s},\underline{a}} \qquad \bigoplus_{i=1}^{s} \bigoplus_{|\alpha| \leq l} \widehat{\mathcal{O}}_{M_{\varphi}^{s},\underline{a}} \quad .$$

For any (germ at \underline{a} of an) analytic subspace L of M_{ω}^{s} , we also write

$$(2.2) J_{\underline{a}}^{l}\varphi \colon J^{l}(b) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{L,\underline{a}} \to \bigoplus_{i=1}^{s} J^{l}(a^{i}) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{L,\underline{a}}$$

for the induced $\widehat{\mathcal{O}}_{L,\underline{a}}$ -homomorphism. Evaluation at \underline{a} transforms $J_{\underline{a}}^{l}\varphi$ to

(2.3)
$$J^l \varphi(\underline{a}) = (J^l \varphi(a^1), \dots, J^l \varphi(a^s)) \colon J^l(b) \to \bigoplus_{i=1}^s J^l(a^i).$$

3. Ideals of relations and Chevalley functions

Let M denote an analytic manifold (over $\mathbb{K} = \mathbb{R}$ or \mathbb{C}), and let $\varphi = (\varphi_1, \dots, \varphi_n) \colon M \to \mathbb{K}^n$ be an analytic mapping. If $a \in M$, let \mathcal{R}_a denote the ideal of formal relations $\operatorname{Ker} \hat{\varphi}_a^*$.

Remark 3.1. \mathcal{R}_a is constant on connected components of the fibres of φ [3, Lemma 5.1].

Let s be a positive integer, and let $\underline{a} = (a^1, \dots, a^s) \in M^s_{\varphi}$. Put

(3.1)
$$\mathcal{R}_{\underline{a}} := \bigcap_{i=1}^{s} \mathcal{R}_{a^{i}} = \bigcap_{i=1}^{s} \operatorname{Ker} \hat{\varphi}_{a^{i}}^{*} \subset \widehat{\mathcal{O}}_{\underline{\varphi}(\underline{a})}.$$

If $k \in \mathbb{N}$, we also write

$$\mathcal{R}^k(\underline{a}) \ := \ \frac{\mathcal{R}_{\underline{a}} + \widehat{\mathfrak{m}}_{\underline{\varphi}(\underline{a})}^{k+1}}{\widehat{\mathfrak{m}}_{\underline{\varphi}(\underline{a})}^{k+1}} \ \subset \ J^k(\underline{\varphi}(\underline{a})) \ .$$

If $b \in \mathbb{K}^n$, let $\pi^k(b) \colon \widehat{\mathcal{O}}_b \to J^k(b)$ denote the canonical projection. For $l \geq k$, let $\pi^{lk}(b) \colon J^l(b) \to J^k(b)$ be the projection. Set

$$E^l(\underline{a}) := \operatorname{Ker} J^l \varphi(\underline{a}), \text{ and } E^{lk}(\underline{a}) := \pi^{lk}(\varphi(\underline{a})).E^l(\underline{a}).$$

3.1. Chevalley's lemma.

Lemma 3.2 ([2, Lemma 8.2.2]; cf. [4, § II, Lemma 7]). Let $\underline{a} \in M_{\varphi}^{s}$, $\underline{a} = (a^{1}, \ldots, a^{s})$. For all $k \in \mathbb{N}$, there exists $l \in \mathbb{N}$ such that $\mathcal{R}^{k}(\underline{a}) = E^{lk}(\underline{a})$; i.e., such that if $F \in \widehat{\mathcal{O}}_{\underline{\varphi}(\underline{a})}$ and $\widehat{\varphi}_{a^{i}}^{*}(F) \in \widehat{\mathfrak{m}}_{a^{i}}^{l+1}$, $i = 1, \ldots, s$, then $F \in \mathcal{R}_{\underline{a}} + \widehat{\mathfrak{m}}_{\underline{\varphi}(\underline{a})}^{k+1}$.

We write $l(\underline{a}, k) = l_{\varphi^*}(\underline{a}, k)$ for the least l satisfying the conclusion of the lemma.

Proof of Lemma 3.2. If $k \leq l_1 \leq l_2$, then

$$\mathcal{R}^k(\underline{a}) \subset E^{l_2,k}(\underline{a}) \subset E^{l_1,k}(\underline{a})$$
,

and the projection $\pi^{l_2,l_1}(\underline{\varphi}(\underline{a}))$ maps $\bigcap_{l\geq l_2} E^{ll_2}(\underline{a})$ onto $\bigcap_{l\geq l_1} E^{ll_1}(\underline{a})$. It follows that $\mathcal{R}^k(\underline{a}) = \bigcap_{l\geq k} E^{lk}(\underline{a})$. Since dim $J^k(\underline{\varphi}(\underline{a})) < \infty$, there exists $l \in \mathbb{N}$ such that $\mathcal{R}^k(\underline{a}) = E^{lk}(\underline{a})$.

3.2. Generic Chevalley function. Let $\underline{a} \in M^s_{\varphi}$ and $k \in \mathbb{N}$. Set

$$H_{\underline{a}}(k) := \dim_{\mathbb{K}} \frac{J^{k}(\underline{\varphi}(\underline{a}))}{\mathcal{R}^{k}(\underline{a})} , \qquad d^{lk}(\underline{a}) := \dim_{\mathbb{K}} \frac{J^{k}(\underline{\varphi}(\underline{a}))}{E^{lk}(\underline{a})} , \text{ if } l \geq k$$

 $(H_{\underline{a}} \text{ is the } \textit{Hilbert-Samuel function of } \widehat{\mathcal{O}}_{\varphi(\underline{a})}/\mathcal{R}_{\underline{a}}).$

Remark 3.3. $d^{lk}(\underline{a}) \leq H_{\underline{a}}(k)$ since $\mathcal{R}^k(\underline{a}) \subset E^{lk}(\underline{a})$. $\mathcal{R}^k(\underline{a}) = E^{lk}(\underline{a})$ (and $d^{lk}(\underline{a}) = H_{\underline{a}}(k)$) if and only if $l \geq l(\underline{a}, k)$.

Lemma 3.4 ([2, Lemma 8.3.3]). Let L be a subanalytic leaf in M_{φ}^s (i.e., a connected subanalytic subset of M_{φ}^s which is an analytic submanifold of M^s ; see Remark 4.4). Then there is a residual subset D of L such that, if $\underline{a},\underline{a}' \in D$, then $H_{\underline{a}}(k) = H_{\underline{a}'}(k)$ and $l(\underline{a},k) = l(\underline{a}',k)$, for all $k \in \mathbb{N}$.

Definition 3.5. We define the *generic Chevalley function* of L as $l(L, k) := l(\underline{a}, k)$ $(k \in \mathbb{N})$, where $\underline{a} \in D$.

Proof of Lemma 3.4. For $\underline{a} \in M_{\varphi}^s$ and $l \geq k$, write $J^l \varphi(\underline{a})$ (2.3) (using local coordinates for M^s as in §2.4, in a neighbourhood of a point of \overline{L}) as a block matrix

$$J^{l}\varphi(\underline{a}) = (S^{lk}(\underline{a}), T^{lk}(\underline{a}))$$
$$= \begin{pmatrix} J^{k}\varphi(\underline{a}) & 0 \\ * & * \end{pmatrix}$$

corresponding to the decomposition of vectors $\xi = (\xi_{\beta})_{\beta \in \mathbb{N}^{n}, |\beta| \leq l}$ in the source as $\xi = (\xi^{k}, \zeta^{lk})$, where $\xi^{k} = (\xi_{\beta})_{|\beta| < k}$ and $\zeta^{lk} = (\xi_{\beta})_{k < |\beta| < l}$. Then

$$E^{lk}(\underline{a}) = \{ \eta = (\eta_{\beta})_{|\beta| < k} \colon S^{lk}(\underline{a}) \cdot \eta \in \operatorname{Im} T^{lk}(\underline{a}) \} .$$

Thus, by Lemma 2.1

$$E^{lk}(\underline{a}) = \operatorname{Ker} \Theta^{lk}(\underline{a}), \text{ and } d^{lk}(\underline{a}) = \operatorname{rk} \Theta^{lk}(\underline{a}),$$

where

$$\Theta^{lk}(\underline{a}) \; := \; \operatorname{ad}^{\,r^{lk}(\underline{a})} T^{lk}(\underline{a}) \cdot S^{lk}(\underline{a}) \; , \quad r^{lk}(\underline{a}) \; := \; \operatorname{rk} T^{lk}(\underline{a}) \; .$$

Set

$$r^{lk}(L) \; := \; \max_{a \in L} r^{lk}(\underline{a}), \; \text{ and } \; d_L^{lk}(\underline{a}) \; := \; \operatorname{rk} \Theta_L^{lk}(\underline{a}), \; \; \underline{a} \in L \; ,$$

where

$$\Theta_L^{lk}(\underline{a}) := \operatorname{ad}^{r^{lk}(L)} T^{lk}(\underline{a}) \cdot S^{lk}(\underline{a})$$

(so that $\Theta^{lk}_L(\underline{a}) = 0$ if $r^{lk}(\underline{a}) < r^{lk}(L)$). Let $Y^{lk} := \{\underline{a} \in L \colon \ r^{lk}(\underline{a}) < r^{lk}(L)\}$. Set

$$d^{lk}(L) := \max_{\underline{a} \in L} d_L^{lk}(\underline{a}) .$$

Clearly, $d_L^{lk}(\underline{a}) = 0$ if $\underline{a} \in Y^{lk}$, and $d_L^{lk}(\underline{a}) = d^{lk}(\underline{a})$ if $\underline{a} \in L \setminus Y^{lk}$. Also set

$$Z^{lk} \ := \ Y^{lk} \cup \left\{\underline{a} \in L \colon d_L^{lk}(\underline{a}) < d^{lk}(L)\right\} \ .$$

Then Y^{lk} and Z^{lk} are proper closed analytic subsets of L. For all $\underline{a} \in L \setminus Z^{lk}$, $r^{lk}(\underline{a}) = r^{lk}(L)$ and $d^{lk}(\underline{a}) = d^{lk}_L(\underline{a}) = d^{lk}(L)$. Put

$$(3.2) D^k := L \setminus \bigcup_{l>k} Z^{lk} , \quad D := \bigcap_{k\geq 1} D^k .$$

By the Baire Category Theorem, the D^k (and hence also D) are residual subsets of L.

Fix $k \in \mathbb{N}$. If $\underline{a} \in D^k$, then $d^{lk}(\underline{a}) = d^{lk}(L)$, for all l > k. If, in addition, $l \geq l(\underline{a}, k)$, then $H_{\underline{a}}(k) = d^{lk}(L)$, by Remark 3.3. If $\underline{a}, \underline{a}' \in D^k$, then, choosing $l \geq l(\underline{a}, k)$ and $\geq l(\underline{a}', k)$, we get $H_{\underline{a}}(k) = H_{\underline{a}'}(k)$. For the second assertion of the lemma, suppose that $l \geq l(\underline{a}, k)$. Then $H_{\underline{a}'}(k) = H_{\underline{a}}(k) = d^{lk}(\underline{a}) = d^{lk}(L) = d^{lk}(\underline{a}')$, so that $l \geq l(\underline{a}', k)$, by Remark 3.3. In the same way, $l \geq l(\underline{a}', k)$ implies that $l \geq l(\underline{a}, k)$.

3.3. Chevalley function of a subanalytic set. Let N denote a real-analytic manifold, and let X be a closed subanalytic subset of N. If $b \in X$, then $\mathcal{F}_b(X)$ or $\mathcal{R}_b \subset \widehat{\mathcal{O}}_b$ denotes the *formal local ideal* of X at b, in the sense of the following simple lemma:

Lemma 3.6. Let $b \in X$. The following three definitions of $\mathcal{F}_b(X)$ are equivalent:

- (1) Let M be a real-analytic manifold and let $\varphi \colon M \to N$ be a proper real-analytic mapping such that $X = \varphi(M)$. Then $\mathcal{F}_b(X) = \bigcap_{a \in \varphi^{-1}(b)} \ker \hat{\varphi}_a^*$.
- (2) $\mathcal{F}_b(X) = \{ F \in \widehat{\mathcal{O}}_b : (F \circ \gamma)(t) \equiv 0 \text{ for every real-analytic arc } \gamma(t) \text{ in } X \text{ such that } \gamma(0) = b \}.$
- (3) $\mathcal{F}_b(X) = \{ F \in \widehat{\mathcal{O}}_b \colon T_b^k F(y) = o(|y-b|^k), \text{ where } y \in X, \text{ for all } k \in \mathbb{N} \}.$ Here $T_b^k F(y)$ denotes the Taylor polynomial of order k of F at b, in any local coordinate system.

Assume that $N = \mathbb{R}^n$, with coordinates $y = (y_1, \dots, y_n)$. Let $b \in X$. Recall (1.1).

Remark 3.7. $\nu_{X,b}(F) \leq \mu_{X,b}(F)$: Suppose that $F \in \widehat{\mathfrak{m}}_b^l + \mathcal{F}_b(X)$; say F = G + H, where $G \in \widehat{\mathfrak{m}}_b^l$ and $H \in \mathcal{F}_b(X)$. Then $|T_b^l G(y)| \leq c|y-b|^l$ and $T_b^l H(y) = o(|y-b|^l)$, $y \in X$, by Lemma 3.6. Hence $|T_b^l F(y)| \leq \operatorname{const}|y-b|^l$ on X.

Definition 3.8 (*Chevalley functions*). Let $b \in X$ and let $k \in \mathbb{N}$. Set

$$l_X(b,k) := \min\{l \in \mathbb{N} : \text{ If } F \in \widehat{\mathcal{O}}_b \text{ and } \mu_{X,b}(F) > l, \text{ then } \nu_{X,b}(F) > k\}$$
.

Let $\varphi \colon M \to N$ be a proper real-analytic mapping such that $X = \varphi(M)$. Set

$$l_{\varphi^*}(b,k) := \min\{l \in \mathbb{N} \colon \text{If } F \in \widehat{\mathcal{O}}_b \text{ and } \nu_{M,a}(\widehat{\varphi}_a^*(F)) > l$$
 for all $a \in \varphi^{-1}(b)$, then $\nu_{X,b}(F) > k\}$.

Remark 3.9. Suppose that $b = \underline{\varphi}(\underline{a})$, where $\underline{a} = (a^1, \dots, a^s) \in M_{\varphi}^s$, $s \geq 1$. By Lemma 3.2, $l_{\varphi^*}(\underline{a}, k) < \infty$. If \underline{a} includes a point a^i in every connected component of $\varphi^{-1}(b)$, then $\bigcap_{i=1}^s \operatorname{Ker} \hat{\varphi}_{a^i}^* = \mathcal{F}_b(X)$ (by Remark 3.1 and Lemma 3.6), so that $l_{\varphi^*}(b, k) \leq l_{\varphi^*}(\underline{a}, k)$.

Lemma 3.10 (see [3, Lemma 6.5]). Let $\varphi \colon M \to N$ be a proper real-analytic mapping such that $X = \varphi(M)$. Then $l_X(b,\cdot) \leq l_{\varphi^*}(b,\cdot)$ for all $b \in X$.

4. Proofs of the main theorems

Let $\varphi \colon M \to \mathbb{K}^n$ be an analytic mapping from a manifold M (over $\mathbb{K} = \mathbb{R}$ or \mathbb{C}). Let s be a positive integer. Let $\underline{a} = (a^1, \dots, a^s) \in M^s_{\varphi}$, and let $b = \varphi(\underline{a})$.

Remark 4.1. By (2.1), the Chevalley functions $l_{\varphi^*}(\underline{a},k)$ and $l_{\varphi^*}(b,k)$ (Definitions 3.8) can be defined using power series that are supported outside the diagram of initial exponents: Set $\mathfrak{N}_{\underline{a}} := \mathfrak{N}(\mathcal{R}_{\underline{a}})$ and $\mathfrak{N}_b := \mathfrak{N}(\mathcal{R}_b)$ (cf. 3.1 and Lemma 3.6). Then

$$\begin{split} l_{\varphi^*}(\underline{a},k) &= & \min\{l \in \mathbb{N} \colon \text{ If } F \in \widehat{\mathcal{O}}_b^{\mathfrak{N}_{\underline{a}}} \text{ and } \hat{\varphi}_{a^i}^*(F) \in \widehat{\mathfrak{m}}_{a^i}^{l+1}, \ i=1,\ldots,s, \\ & \text{ then } F \in \mathcal{R}_{\underline{a}} + \widehat{\mathfrak{m}}_b^{k+1}\} \ , \\ l_{\varphi^*}(b,k) &= & \min\{l \in \mathbb{N} \colon \text{ If } F \in \widehat{\mathcal{O}}_b^{\mathfrak{N}_b} \text{ and } \hat{\varphi}_a^*(F) \in \widehat{\mathfrak{m}}_a^{l+1}, \text{ for all } a \in \varphi^{-1}(b), \\ & \text{ then } F \in \mathcal{R}_b + \widehat{\mathfrak{m}}_b^{k+1}\} \ . \end{split}$$

(In the latter, we assume that φ is a proper real-analytic mapping.)

If $l \in \mathbb{N}$, set $J^l(b)^{\mathfrak{N}_{\underline{a}}} := \{ \xi = (\xi_\beta)_{|\beta| < l} \in J^l(b) \colon \xi_\beta = 0 \text{ if } \beta \in \mathfrak{N}_a \}$. Consider the linear mapping

$$\Phi^l(\underline{a}) \colon J^l(b)^{\mathfrak{N}_{\underline{a}}} \to \bigoplus_{i=1}^s J^l(a^i)$$

obtained by restriction of $J^l\varphi(\underline{a}): J^l(b) \to \bigoplus J^l(a^i)$ (2.3). Given $k \leq l$, write $\Phi^l(\underline{a})$ as a block matrix

$$\Phi^{l}(\underline{a}) = (A^{lk}(\underline{a}), B^{lk}(\underline{a})) ,$$

where $A^{lk}(\underline{a})$ is given by the restriction of $\Phi^l(a)$ to $J^k(b)^{\mathfrak{N}_{\underline{a}}}$.

Remark 4.2. If $\xi \in J^l(b)^{\mathfrak{N}_{\underline{a}}}$, write $\xi = (\eta, \zeta)$ corresponding to this block decomposition. Then $l \geq l_{\varphi^*}(\underline{a}, k)$ if and only if $A^{lk}(\underline{a})\eta + B^{lk}(\underline{a})\zeta = 0$ implies $\eta = 0$ [3, Lemma 8.13].

Lemma 4.3 ((cf. [3, Prop. 8.15]). Let $s \geq 1$ and consider $\varphi = \varphi^s \colon M_{\varphi}^s \to \mathbb{R}^n$. Let L be a relatively compact subanalytic leaf in M_{φ}^{s} (cf. Lemma 3.4) such that $\mathfrak{N}_{\underline{a}} = \mathfrak{N}(\mathcal{R}_{\underline{a}})$ is constant on L. Let l(k) = l(L,k) denote the generic Chevalley function of L. Then there exists $p \in \mathbb{N}$ such that $l_{\varphi^*}(\underline{a}, k) \leq l(k) + p$, for all $\underline{a} \in L$ and $k \in \mathbb{N}$.

Proof. Set $\mathfrak{N} = \mathfrak{N}_{\underline{a}}, \underline{a} \in L$. We can assume that \overline{L} lies in a coordinate chart for M^s as in §2.4. Let $k \in \mathbb{N}$ and let l = l(k). Let $\underline{a} = (a^1, \dots, a^s) \in L$, and set $b = \underline{\varphi}(\underline{a})$. Consider the linear mapping $\Phi^l(\underline{a}) = (A^{lk}(\underline{a}), B^{lk}(\underline{a})) : J^l(b)^{\mathfrak{N}} \to \bigoplus_{i=1}^s J^l(\overline{a^i})$ as above. The $\widehat{\mathcal{O}}_{L,\underline{a}}$ -homomorphism $J_{\underline{a}}^{l}\varphi\colon J^{l}(b)\otimes_{\mathbb{K}}\widehat{\mathcal{O}}_{L,\underline{a}}\to \bigoplus_{i=1}^{s}J^{l}(a^{i})\otimes_{\mathbb{K}}\widehat{\mathcal{O}}_{L,\underline{a}}$ (2.2) induces an $\mathcal{O}_{L,a}$ -homomorphism

$$\Phi_{\underline{a}}^{l} = (A_{\underline{a}}^{lk}, B_{\underline{a}}^{lk}) \colon J^{l}(b)^{\mathfrak{N}} \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{L,\underline{a}} \to \bigoplus_{i=1}^{s} J^{l}(a^{i}) \otimes_{\mathbb{K}} \widehat{\mathcal{O}}_{L,\underline{a}} ;$$

evaluating at \underline{a} transforms $\Phi^l_{\underline{a}}$ to $\Phi^l(\underline{a}) = (A^{lk}(\underline{a}), B^{lk}(\underline{a}))$. Let $r = \operatorname{rk} B^{lk}_{\underline{a}} = \text{ generic rank of } B^{lk}(\underline{x}), \ \underline{x} \in L$. Let $\Theta_{\underline{a}} = \operatorname{ad}^r B^{lk}_{\underline{a}} \cdot A^{lk}_{\underline{a}}$. Then $\operatorname{Ker} \Theta_{\underline{a}} = 0$ (i.e., $\operatorname{Ker} \Theta(\underline{x}) = 0$ generically on L, where $\Theta(\underline{x}) = \operatorname{ad}^r B^{lk}(\underline{x}) \cdot A^{lk}(\underline{x})$, by Remark 4.2). Let $d = \operatorname{rk} \Theta_{\underline{a}}$. Then there is a nonzero minor $\delta_{\underline{a}} \in \mathcal{O}_{L,\underline{a}}$ of $\Theta_{\underline{a}}$ of order d; $\delta_{\underline{a}}$ is induced by a minor $\delta(\underline{x})$ of order d of $\Theta(\underline{x})$, $\underline{x} \in L$, such that $\delta(\underline{x}) \neq 0$ on a residual subset of L. Since δ is a restriction to L of an analytic function defined in a neighbourhood of \overline{L} , the order of $\delta_{\underline{x}}$, $\underline{x} \in L$, is bounded on L; say, $\delta_{\underline{x}} \leq p$.

We claim that $l_{\varphi^*}(\underline{a}, k) \leq l(k) + p$ for all $\underline{a} \in L$: Let $\underline{a} = (a^1, \dots, a^s) \in L$, and let $b = \underline{\varphi}(\underline{a})$. Let l = l(k) and l' = l + p. Suppose that $F \in \widehat{\mathcal{O}}_b^{\mathfrak{N}}$ and $\widehat{\varphi}_{a^i}^*(F) \in \widehat{\mathfrak{m}}_{a^i}^{l'+1}$, $i=1,\ldots,s$. Let $\hat{\xi}_{\underline{a}}=(\hat{\eta}_{\underline{a}},\hat{\zeta}_{\underline{a}})$ denote the element of $J^l(b)^{\mathfrak{N}}\otimes_{\mathbb{K}}\widehat{\mathcal{O}}_{L,\underline{a}}$ induced by $J_b^l F \in J^l(b) \otimes_{\mathbb{K}} \widehat{\mathcal{O}_b}$ via the pull-back. Then each component of $A_{\underline{a}}^{lk} \hat{\eta}_{\underline{a}} + B_{\underline{a}}^{lk} \hat{\zeta}_{\underline{a}}$ belongs to $\widehat{\mathfrak{m}}_{L,\underline{a}}^{l'+1-l}$ (as we see by taking formal derivatives of order $\leq l$ of the $\widehat{\varphi}_{a^i}^*(F)$). It follows that each component of $\Theta_{\underline{a}}\widehat{\eta}_{\underline{a}}$ and therefore (by Cramer's rule) each component of $\delta_{\underline{a}} \cdot \hat{\eta}_{\underline{a}}$ belongs to $\widehat{\mathfrak{m}}_{L,\underline{a}}^{l'+1-l}$. Thus, each component of $\hat{\eta}_{\underline{a}}$ lies in $\widehat{\mathfrak{m}}_{L,a}^{l'+1-l-p} = \widehat{\mathfrak{m}}_{L,\underline{a}}$; i.e., $\widehat{\eta}_{\underline{a}}(\underline{a}) = 0$, so that F vanishes to order k at $b = \varphi(\underline{a})$.

Proof of Theorem 1.1. By [2, Theorems A,C], there is a locally finite partition of M into relatively compact subanalytic leaves L such that the diagram of initial exponents $\mathfrak{N}_a = \mathfrak{N}(\mathcal{R}_a)$ is constant on each L. Given L, let l(L,k) denote the generic Chevalley function. (In particular, $l(L,k) = l_{\varphi^*}(a,k)$, for all a in a residual subset of L.) Since φ is regular, there exist α_L, γ_L such that $l(L, k) \leq \alpha_L k + \gamma_L$, for all $k \in \mathbb{N}$ (by [10]). By Lemma 4.3 (in the case s = 1), there exists $p_L \in \mathbb{N}$ such that $l_{\varphi^*}(a, k) \leq \alpha_L k + \gamma_L + p_L$, for all $a \in L$ and all k. The result follows. \square

Remark 4.4. In the case $\mathbb{K}=\mathbb{C}$, we define "subanalytic leaf" using the underlying real structure. If φ is regular, then the diagram \mathfrak{N}_a is, in fact, an uppersemicontinuous function of a, with respect to the \mathbb{K} -analytic Zariski topology of M (and a natural total ordering of $\{\mathfrak{N}\in\mathbb{N}^n\colon\mathfrak{N}+\mathbb{N}^n=\mathfrak{N}\}$) [2, Theorem C], but we do not need the more precise result here.

Lemma 4.5. Let $s \ge 1$ and let $\underline{a} = (a^1, \dots, a^n) \in M_{\varphi}^s$. Suppose that φ is regular at a^1, \dots, a^n . Then there exist $\alpha, \beta \in \mathbb{R}$ such that $l_{\varphi^*}(\underline{a}, k) \le \alpha k + \beta$, for all $k \in \mathbb{N}$.

Proof. Let $b = \underline{\varphi}(\underline{a})$. For each $i = 1, \dots, s$, since φ is regular at a^i , there exist α^i, β^i such that

$$(4.1) l_{\varphi^*}(a^i, k) \leq \alpha^i k + \beta^i, for all k.$$

Of course, $\bigcap_{i=1}^s \operatorname{Ker} \hat{\varphi}_{a^i}^*$ is the kernel of the homomorphism $\widehat{\mathcal{O}}_b \to \bigoplus_{i=1}^s \widehat{\mathcal{O}}_b / \ker \hat{\varphi}_{a^i}^*$. By the Artin-Rees lemma (cf. Remark 1.4), there exists $\lambda \in \mathbb{N}$ such that, if $F \in \widehat{\mathfrak{m}}_b^{k+\lambda} + \ker \hat{\varphi}_{a^i}^*$, $i = 1, \ldots, s$, then

(4.2)
$$F \in \widehat{\mathfrak{m}}_b^k + \bigcap_{i=1}^s \operatorname{Ker} \widehat{\varphi}_{a^i}^*.$$

Now let $F \in \widehat{\mathcal{O}}_b$ and suppose that $\widehat{\varphi}_{a^i}^*(F) \in \widehat{\mathfrak{m}}_{a^i}^{\alpha^i(\lambda+k)+\beta^i+1}$, $i=1,\ldots,s$. Then $F \in \widehat{\mathfrak{m}}_b^{\lambda+k+1} + \operatorname{Ker} \widehat{\varphi}_{a^i}^*$, $i=1,\ldots,s$, by (4.1), so that $F \in \widehat{\mathfrak{m}}_b^{k+1} + \bigcap_{i=1}^s \operatorname{Ker} \widehat{\varphi}_{a^i}^*$, by (4.2). In other words, $l_{\varphi^*}(\underline{a},k) \leq \alpha k + \beta$, where $\alpha = \max \alpha^i$ and $\beta = \lambda \max \alpha^i + \max \beta^i$.

Proof of Theorem 1.2. Suppose that $\varphi: M \to \mathbb{R}^n$ is a real-analytic mapping, where M is compact. Let $X = \varphi(M)$. Let $s \geq 1$, $\underline{a} \in M_{\varphi}^s$, $b = \underline{\varphi}(\underline{a})$. If $\underline{a} = (a^1, \ldots, a^s)$ includes a point a^i in every connected component of $\varphi^{-1}(b)$, then

$$(4.3) l_X(b,k) \leq l_{\varphi^*}(\underline{a},k) ,$$

by Remark 3.9 and Lemma 3.10.

Let L be a relatively compact subanalytic leaf in M_{φ}^s , such that $\mathfrak{N}_{\underline{a}}=\mathfrak{N}(\mathcal{R}_{\underline{a}})$ is constant on L. Suppose that φ is regular at a^i , for all $\underline{a}=(a^1,\ldots,a^s)\in L$ and $i=1,\ldots,s$. Let l(L,k) denote the generic Chevalley function of L. By Lemma 4.5, there exist α,β such that $l(L,k)\leq \alpha k+\beta$. Therefore, by Lemma 4.3, there exist α_L,β_L such that

$$(4.4) l_{\varphi^*}(\underline{a}, k) \leq \alpha_L k + \beta_L, for all \underline{a} \in L.$$

To prove the theorem, we can assume that X is compact. Let φ be a mapping as above, such that $X = \varphi(M)$. We consider first the case that X is Nash. Then we can assume that φ is regular. Let s denote a bound on the number of connected components of a fibre $\varphi^{-1}(b)$, for all $b \in X$. Then there is a finite partition of M_{φ}^{s} into relatively compact subanalytic leaves L, such that $\mathfrak{N}_{\underline{a}} = \mathfrak{N}(\mathcal{R}_{\underline{a}})$ is constant on every L. By (4.3) and (4.4), for each L, there exist α_{L}, β_{L} such that $l_{X}(b,k) \leq \alpha_{L}k + \beta_{L}$, for all $b \in \underline{\varphi}(L)$ and all k. Therefore, $l_{X}(b,k)$ has a uniform linear bound.

Finally, we consider X formally Nash. Let $NR(\varphi) \subset M$ denote the set of points at which φ is not regular. Then $NR(\varphi)$ is a nowhere-dense closed analytic subset of M ([11, Theorem 1]). For each positive integer s, set

$$NR(\underline{\varphi}^s) := M_{\varphi}^s \cap \bigcup_{i=1}^s \{\underline{a} = (a^i, \dots, a^s) \in M^s : a^i \in NR(\varphi)\};$$

then $NR(\varphi^s)$ is a closed analytic subset of M^s_{φ} .

If $b \in X$ and a, a' belong to the same connected component of $\varphi^{-1}(b)$, then φ is regular at a if and only if φ is regular at a' (cf. Remark 3.1). Let t be a bound on the number of connected components of a fibre $\varphi^{-1}(b)$, for all $b \in X$. For each $s \leq t$, define $X_s := \{b \in X : \varphi^{-1}(b) \text{ has precisely } s \text{ regular components}\}$ and $Y_s := \{b \in X : \varphi^{-1}(b) \text{ has at least } s \text{ regular components}\}$. Then $X_s = Y_s \setminus Y_{s+1}$, and

$$Y_s = \underline{\varphi}^s(\mathring{M}_{\varphi}^s \setminus NR(\underline{\varphi}^s)) ;$$

in particular, all the X_s and Y_s are subanalytic (cf. §3.2).

The hypothesis of the theorem implies:

- (1) $X = \bigcup_{s=1}^{t} X_s;$
- (2) If $b \in X_s$ and $\underline{a} \in (\underline{\varphi}^s)^{-1}(b) \cap (\mathring{M}_{\varphi}^s \setminus NR(\underline{\varphi}^s))$, then $\mathcal{R}_{\underline{a}} = \mathcal{R}_b$.
- ((2) follows from the fact that $\mathcal{F}_b(X) = \mathcal{F}_b(Y_b)$, where Y_b is some closed Nash subanalytic subset of X, and (1) from the fact that the latter condition holds for all $b \in X$.)

By [11, Theorem 2], for each s, there is a finite stratification \mathcal{L}_s of M_{φ}^s compatible with $\mathrm{NR}(\underline{\varphi}^s)$ such that $\mathfrak{N}_{\underline{a}} = \mathfrak{N}(\mathcal{R}_{\underline{a}})$ is constant on every stratum $L \subset M_{\varphi}^s \backslash \mathrm{NR}(\underline{\varphi}^s)$, $L \in \mathcal{L}_s$. Clearly,

$$X_s = \bigcup_{\substack{L \in \mathcal{L}_s \\ L \subset M_{\varphi}^s \backslash \mathrm{NR}(\underline{\varphi}^s)}} \underline{\varphi}^s \left(L \cap \mathring{M}_{\varphi}^s \right) \cap X_s ;$$

hence

$$X \ = \ \bigcup_{s=1}^t \bigcup_{\substack{L \in \mathcal{L}_s \\ L \subset M_{\varphi}^s \backslash \mathrm{NR}(\underline{\varphi}^s)}} \underline{\varphi}^s \left(L \cap \mathring{M}_{\varphi}^s \right) \ .$$

Again by (4.3) and (4.4), for each L, there exist α_L, β_L such that $l_X(b, k) \leq \alpha_L k + \beta_L$, for all $b \in \underline{\varphi}(L)$ and all k. The result follows.

References

- E. Bierstone and P.D. Milman, Composite differentiable functions, Ann. of Math. (2) 116 (1982), 541–558.
- E. Bierstone and P.D. Milman, Relations among analytic functions I, Ann. Inst. Fourier (Grenoble) 37:1 (1987), 187–239; II, 37:2 (1987), 49–77.
- E. Bierstone and P.D. Milman, Geometric and differential properties of subanalytic sets, Ann. of Math. (2) 147 (1998), 731–785.
- 4. C. Chevalley, On the theory of local rings, Ann. of Math. (2) 44 (1943), 690-708.
- A.J. Duncan and L. O'Carroll, A full uniform Artin-Rees theorem, J. Reine Agnew. Math. 394 (1989), 203–207.
- A.M. Gabrielov, Formal relations between analytic functions, Math. USSR Izv. 7 (1973), 1056–1088.
- H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero: I, II, Ann. of Math. (2) 79 (1964), 109–326.

- 8. C. Huneke, Uniform bounds in Noetherian rings, Invent. Math. 107 (1992), 203-223.
- 9. S. Izumi, Linear complementary inequalities for orders of germs of analytic functions, Invent. Math. 65 (1982), 459–471.
- 10. S. Izumi, Gabrielov's rank condition is equivalent to an inequality of reduced orders, Math. Ann. 276 (1986), 81–89.
- W. Pawłucki, On Gabrielov's regularity condition for analytic mappings, Duke Math. J. 65 (1992), 299–311.
- 12. T. Wang, Linear Chevalley estimates, Trans. Amer. Math. Soc. 347 (1995), 4877–4898.
- J. Adamus, Institute of Mathematics of the Polish Academy of Sciences, 00-956 Warszawa 10, Sniadeckich 8, P.O. Box 21, Poland

 $E ext{-}mail\ address: adamus@impan.gov.pl}$

E. Bierstone, Department of Mathematics, University of Toronto, Toronto, Ontario, Canada M5S 3G3

 $E\text{-}mail\ address: \ \mathtt{bierston@math.toronto.edu}$

P.D. Milman, Department of Mathematics, University of Toronto, Toronto, Ontario, Canada M5S 3G3

 $E\text{-}mail\ address{:}\ \mathtt{milman@math.toronto.edu}$