

SPIN REPRESENTATIONS AND CENTRALIZER ALGEBRAS FOR THE SPINOR GROUPS

KAZUHIKO KOIKE

ABSTRACT. We pursue an analogy of the Schur-Weyl reciprocity for the spinor groups and pick up the irreducible spin representations in the tensor space $\Delta \otimes \otimes^k V$. Here Δ is the fundamental representation of $Pin(N)$ and V is the natural (vector) representation of the orthogonal group $O(N)$. We consider the centralizer algebra $\mathbf{CP}_k = \text{End}_{Pin(N)}(\Delta \otimes \otimes^k V)$ for $Pin(N)$, the double covering group of $O(N)$ and define two kinds of linear basis in \mathbf{CP}_k (one comes from invariant theory and the other from representation theory), both of which are parameterized by the 'generalized Brauer diagrams'. We develop analogous argument to the original Brauer centralizer algebra for $O(N)$ and determine the transformation matrices between the above two basis and give the multiplication rules of those basis. Finally we define the subspaces in $\Delta \otimes \otimes^k V$, on which the symmetric group \mathfrak{S}_k and $Pin(N)$ or $Spin(N)$ act as a dual pair.

1. INTRODUCTION

In this paper we consider analogs of the Brauer centralizer algebras for the spinor groups and define the subspace $T_{k,s}^0$ of the tensor space $\Delta \otimes \otimes^k V$, on which the symmetric group \mathfrak{S}_k and $Pin(N)$ act as the dual pair. (See Definition 10.1 and Theorem 10.3.) Here $V \cong \mathbb{R}^N$ is the vector (natural) representation of $O(N)$ and Δ is the fundamental representation of $Pin(N)$.

For the irreducible representations of the orthogonal groups, together with the other classical groups, this procedure is carried out by H. Weyl in his book *The Classical Groups*. Namely the centralizer algebra of the orthogonal group in $\otimes^k V$ is generated by the symmetric group \mathfrak{S}_k and the contractions and the immersions of the invariant form and is called the Brauer centralizer algebras. The basis of this algebra is parameterized by the diagrams, which is called the Brauer diagrams and the multiplication rules of the basis elements are given in [3]. H. Weyl defined the subspace consisting of the traceless tensors in $\otimes^k V$ and he showed that the symmetric group \mathfrak{S}_k and the orthogonal group act on

Date: March 26, 2019.

1991 *Mathematics Subject Classification.* 05E15, 17B10, 20G05, 22E46, 22E47.
Supported by the Grant-in-Aid for Scientific Research.

this space as the dual pair and he picked up the irreducible representations of $O(N)$ as the images of the Young symmetrizers corresponding to the highest weights of the irreducible representations.

So we follow this line and first determine the basis of the centralizer algebra

$$\mathbf{CP}_k = \text{Hom}_{Pin(N)}(\Delta \otimes^k V, \Delta \otimes^k V).$$

In this case, we can define two kinds of basis of the centralizer algebra \mathbf{CP}_k and the basis elements of both bases are parameterized by the same diagrams, which we call ‘the generalized Brauer diagrams’. (See figure 1.)

The generalized Brauer diagrams are, by definition, the graphs of two lines of vertices with k vertices in the upper row and l vertices in the lower row, in which vertices are connected with each other as in the usual Brauer diagrams except for admitting isolated vertices. Namely they are graphs with no loops in which the number of edges connected to each vertex is either 0 or 1.




FIGURE 1. The generalized Brauer diagrams of $k = 2$ and $l = 2$

Let us denote the set of the generalized Brauer diagrams with k vertices in the upper row and l vertices in the lower row by \mathbf{GB}_l^k .

Precisely speaking, for each element of \mathbf{GB}_l^k , we define two elements in $\mathbf{CP}_l^k = \text{Hom}_{Pin(N)}(\Delta \otimes^k V, \Delta \otimes^l V)$, one of which comes from the invariant theory (we call this element an invariant theoretic element.) and the other of which comes from the representation theory (we call this element a representation theoretic element.). The edges in the generalized Brauer diagram convey the same meaning as in the usual Brauer diagrams. As for the exact definitions of the above elements, see the part after Lemma 5.6 in the section 5 for the invariant theoretic elements and the part after Definition 6.5 in the section 6 for the representation theoretic elements.

For any element $d \in \mathbf{GB}_l^k$, let us denote the corresponding invariant theoretic element by d_{inv} and the corresponding representation theoretic element by d_{rt} respectively. Then we can show that both of the sets $\{d_{inv} | d \in \mathbf{GB}_l^k\}$ and $\{d_{rt} | d \in \mathbf{GB}_l^k\}$ are bases of the centralizer

space $\mathbf{CP}_1^k = \text{Hom}_{\text{Pin}(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V)$ if $k \leq n$ and $l \leq n$, here $N = 2n$ or $N = 2n + 1$. If $k = l$ and $k \leq n$, we obtain two kinds of basis of the centralizer algebra \mathbf{CP}_k .

In Theorem 7.2 and Theorem 7.6, we give the transformation matrices between the above two bases explicitly.

If $k > n$ or $l > n$, the sets $\{d_{inv}|d \in \mathbf{GB}_1^k\}$ and $\{d_{rt}|d \in \mathbf{GB}_1^k\}$ are not basis of \mathbf{CP}_1^k at all, but the invariant theory tells us that the elements of the set $\{d_{inv}|d \in \mathbf{GB}_1^k\}$ still span the whole space \mathbf{CP}_1^k . (Lemma 5.6.)

We also give a series of the formulas to calculate the products of the basis elements in Theorem 8.1, from which all the products of the basis elements can be calculated.

Those formulas tell us that we can define the ‘generic’ centralizer algebra of \mathbf{CP}_k if we put $N = X$ (X is an indeterminate.) as in the usual Brauer centralizer algebras.

From the definition, \mathbf{CP}_k naturally contains the ordinary Brauer centralizer algebra and the symmetric group of degree k .

Since we already know the decomposition rules (see the formula (2.2.3), (2.2.4), (2.4.2).) of the tensor products, all the irreducible spin representations (that is, the representations not coming from $O(N)$ or $SO(N)$) of $\text{Pin}(N)$ or $\text{Spin}(N)$ occur in this space $\Delta \otimes \otimes^k V$ for some k . (See Section 2.)

So we define the subspace $T_{k,s}^0$ ($s = 1, 2, \dots, \min(k, n)$) of $\Delta \otimes \otimes^k V$ to be the intersection of the kernels of all the projections from $\Delta \otimes \otimes^k V$ to Δ (See Definition 6.4.) and of the contractions and of the alternating operators of degrees greater than s . This space is an analog of the traceless tensors in the classical arguments ([21]).

Then on this space $T_{k,s}^0$, the symmetric group \mathfrak{S}_k of degree k and $\text{Pin}(N)$ act as a dual pair and we obtain an analogy of Schur-Weyl reciprocity for $\text{Pin}(N)$. (See Theorem 10.3.)

For odd spinor groups, the irreducible $\text{Pin}(2n + 1)$ modules are still irreducible as $\text{Spin}(2n + 1)$ -modules, hence $\text{Spin}(2n + 1)$ and \mathfrak{S}_k act on this space $T_{k,s}^0$ as a dual pair.

For $\text{Spin}(2n)$, if we define the ‘associator’ A of $\text{End}(\Delta)$ by $A = \text{id} \oplus -\text{id}$ on the direct sum $\Delta = \Delta^+ \oplus \Delta^-$ and consider the endomorphism $(A \otimes \text{id}) \in \text{End}(\Delta \otimes \otimes^k V)$, then $(A \otimes \text{id})$ is an involution and the action of $(A \otimes \text{id})$ commutes with those of \mathbf{CP}_k and $\text{Spin}(2n)$. (See Lemma 10.6.)

The centralizer algebra $\mathbf{CS}_k = \text{End}_{\text{Spin}(2n)}(\Delta \otimes \otimes^k V)$ for $\text{Spin}(2n)$ is generated by \mathbf{CP}_k and $(A \otimes \text{id})$.

We denote the ± 1 eigenspaces of $(A \otimes \text{id})$ in the spaces $T_{k,s}^0$ by $T_{k,s}^{0,\pm}$. On each of the spaces $T_{k,s}^{0,\pm}$, $\text{Spin}(2n)$ and \mathfrak{S}_k act as the dual pair. (See Theorem 10.8.)

The key theorems, which connect the fundamental pin or spin representations to the tensor calculus, are Theorem 4.1, 4.4 and Theorem 4.3, 4.6, which give the explicit $Pin(N)$ -equivariant (or $Spin(N)$ -equivariant) embeddings from $\bigwedge^\ell V$ to $\Delta \otimes \Delta^*$ under the specified basis of both spaces. The transformation matrices of the same weight spaces in the direct sum of the exterior products and $\Delta \otimes \Delta^*$ are given by the Hadamard matrices.

We note that the construction here goes over a field $\mathbb{Q}(\sqrt{2})$.

2. PARAMETRIZATION OF THE IRREDUCIBLE REPRESENTATIONS

We take a symmetric anti-diagonal matrix $S = (\delta_{i, N+1-i})$ of size N , where $\delta_{i,j}$ is Kronecker's delta and define the orthogonal group as

$$O(N, \mathbb{R}) = \{g \in GL(N); gS^t g = S\}.$$

Then its Lie algebra is given by

$$\mathfrak{so}(N, \mathbb{R}) = \{X \in M(N, \mathbb{R}); XS + S^t X = 0\}.$$

As a Cartan subalgebra \mathfrak{H} , we can take the intersection of the above Lie algebra and the diagonal matrices and let \mathbf{e}_i be a homomorphism from Cartan subalgebra to \mathbb{R} defined by taking the (i, i) -th component of the diagonal matrices.

Let P^+ be the set of the dominant integral weights for $\mathfrak{so}(N, \mathbb{R})$. Namely for $N = 2n + 1$, P^+ is given by

$$P^+ = \{\lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2 + \dots + \lambda_n \mathbf{e}_n; \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0\}$$

and for $N = 2n$, P^+ is given by

$$P^+ = \{\lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2 + \dots + \lambda_n \mathbf{e}_n; \lambda_1 \geq \lambda_2 \geq \dots \geq |\lambda_n| \geq 0\}.$$

Here all the λ_i 's are integers or half-integers (namely $1/2 + \mathbb{Z}$) simultaneously.

We put $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ and denote the corresponding irreducible character by $\lambda_{Spin(N)}$.

The set of the dominant integral weights of the group $SO(N)$ is the subset of the above, defined by the condition that all the λ_i 's are integers. So in that case we also write $\lambda_{SO(N)}$ for $\lambda_{Spin(N)}$. Then the vector representation (the natural representation) is denoted by $(1)_{SO(N)}$ and the representation $\bigwedge^i V$ of $\mathfrak{so}(N)$ ($i = 1, 2, \dots, n$) is denoted by e_i as usual. e_i is the irreducible representation $(1^i)_{SO(N)}$ if $i < n$ or if $i = n$ and $N = 2n + 1$.

If $i = n$ and $N = 2n$, e_n is not irreducible and is decomposed into the sum of two irreducible representations $e_n^+ = (1^n)_{SO(2n)}$ and $e_n^- = (1^{n-1}, -1)_{SO(2n)}$. Later we give an explicit description of e_n^+ and e_n^- in $\bigwedge^n V$.

We note that $\bigwedge^i V \cong \bigwedge^{N-i} V$ as $SO(N)$ module.

For a partition $\delta = (\delta_1, \delta_2, \dots, \delta_n)$, we denote the irreducible representations of $Spin(N)$ as follows.

For $Spin(2n+1)$, we put $\lambda = (1/2 + \delta_1, 1/2 + \delta_2, \dots, 1/2 + \delta_n)$ and denote this irreducible representation by $\lambda_{Spin(2n+1)} = [\Delta, \delta]_{Spin(2n+1)}$. Then $[\Delta, \delta]_{Spin(2n+1)}$ can be considered as an irreducible character of $Pin(2n+1)$. When we consider this as the representation of $Pin(2n+1)$, we write $[\Delta, \delta]_{Pin(2n+1)}$. (See also Remark 3.9.)

For $Spin(2n)$, we put

$$(1/2 + \delta)_{Spin(2n)}^+ = (1/2 + \delta_1, 1/2 + \delta_2, \dots, 1/2 + \delta_n)_{Spin(2n)}$$

and

$$(1/2 + \delta)_{Spin(2n)}^- = (1/2 + \delta_1, 1/2 + \delta_2, \dots, 1/2 + \delta_{n-1}, -1/2 - \delta_n)_{Spin(2n)}.$$

We denote simply by Δ^+ the irreducible representation $(1/2 + \emptyset)_{Spin(2n)}^+$ and by Δ^- the irreducible representation $(1/2 + \emptyset)_{Spin(2n)}^-$.

Then we have

$$\Delta = \Delta^+ + \Delta^- = \prod_{i=1}^n (x_i^{1/2} + x_i^{-1/2})$$

and

$$\Delta^+ - \Delta^- = \prod_{i=1}^n (x_i^{1/2} - x_i^{-1/2}).$$

Here x_i denotes the i -th diagonal component of the maximal torus (the diagonal matrices $= \{\text{diag}(x_1, x_2, \dots, x_n, x_n^{-1}, \dots, x_1^{-1})\}$) of the group $SO(2n)$. We put $\Delta' = \Delta^+ - \Delta^-$. Moreover we introduce the sum character and the difference character for $Spin(2n)$ and denote them by

$$[\Delta, \delta]_{Spin(2n)} = \lambda_{Spin(2n)}^{(+)} = (1/2 + \delta)_{Spin(2n)}^+ + (1/2 + \delta)_{Spin(2n)}^-$$

and

$$[\Delta', \delta]_{Spin(2n)} = \lambda_{Spin(2n)}^{(-)} = (1/2 + \delta)_{Spin(2n)}^+ - (1/2 + \delta)_{Spin(2n)}^-.$$

Then $[\Delta, \delta]_{Spin(2n)}$ becomes the irreducible character of $Pin(2n)$ and when we consider this as the representation of $Pin(2n)$, we write $[\Delta, \delta]_{Pin(2n)}$. (See the section 3.)

For a partition $\mu = (\mu_1, \mu_2, \dots, \mu_n)$, we also denote the sum character and the difference character for $SO(2n)$ by

$$\mu_{SO(2n)}^{(+)} = (\mu_1, \mu_2, \dots, \mu_n)_{SO(2n)} + (\mu_1, \dots, \mu_{n-1}, -\mu_n)_{SO(2n)}$$

and

$$\mu_{SO(2n)}^{(-)} = (\mu_1, \mu_2, \dots, \mu_n)_{SO(2n)} - (\mu_1, \dots, \mu_{n-1}, -\mu_n)_{SO(2n)}.$$

Then if $\ell(\mu) < n$, $\mu_{SO(2n)}^{(+)} = 2\mu_{SO(2n)}$ and $\mu_{SO(2n)}^{(-)} = 0$. For a partition μ with $\ell(\mu) < n$, $\mu_{SO(2n)}$ becomes an irreducible character of $O(2n)$, so we sometimes write $\mu_{O(2n)}$ for $\mu_{SO(2n)}$ in this case.

If $\ell(\mu) = n$, $\mu_{SO(2n)}^{(+)}$ is the irreducible character of $O(2n)$ and we sometimes write $\mu_{O(2n)}$ for $\mu_{SO(2n)}^{(+)}$ in this case.

First we give complementary formulas which are not dealt with in the previous paper [14]. We fix some notations. For a partition λ with its length at most n , we denote the irreducible character of the Symplectic group $Sp(2n)$ by $\lambda_{Sp(2n)}$. Then ${}^nC_{\mu,\nu}^\lambda$ denotes the multiplicity of $\lambda_{Sp(2n)}$ in the irreducible decomposition of the tensor product $\mu_{Sp(2n)}\nu_{Sp(2n)}$. Namely

$$\mu_{Sp(2n)}\nu_{Sp(2n)} = \sum_{\lambda} {}^nC_{\mu,\nu}^\lambda \lambda_{Sp(2n)}.$$

We note that we can calculate these structure constants ${}^nC_{\mu,\nu}^\lambda$ explicitly by using the Littlewood-Richardson rules. (See Theorem 3.1 in [12] and also [14], [15], [16].)

Theorem 2.1 (Decomposition of Tensor Products for $Spin(2n)$). (i)

If $\ell(\mu) = n$, we have

$$(2.1.1) \quad [\Delta, \delta]_{Spin(2n)} \mu_{SO(2n)}^{(-)} = \sum_{\substack{\nu, \lambda \\ \delta/\nu: \text{vertical strip}}} (-1)^{|\delta/\nu|} {}^nC_{\nu, (\mu - (1^n))}^\lambda \sum_{\substack{\kappa \supseteq \lambda, \ell(\kappa) \leq n \\ \kappa/\lambda: \text{vertical strip}}} [\Delta', \kappa]_{Spin(2n)}.$$

Here $(\mu - (1^n))$ denotes the partition $(\mu_1 - 1, \mu_2 - 1, \dots, \mu_n - 1)$ and δ/ν denotes the skew Young diagram and $|\delta/\nu|$ denotes the size of the skew Young diagram.

(ii) If $\ell(\mu) = n$, we have

$$(2.1.2) \quad [\Delta', \delta]_{Spin(2n)} \mu_{SO(2n)}^{(-)} = \sum_{\substack{\nu, \lambda \\ \delta/\nu: \text{vertical strip}}} (-1)^{|\mu|+|\nu|} {}^nC_{\nu, (\mu - (1^n))}^\lambda \sum_{\substack{\kappa \supseteq \lambda, \ell(\kappa) \leq n \\ \kappa/\lambda: \text{vertical strip}}} (-1)^{|\kappa|} [\Delta, \kappa]_{Spin(2n)}.$$

Proof. The proof is almost same as in the proof of Theorem 1.2 of [14].

From Theorem 10.1 in [14], we have

$$\begin{aligned} [\Delta, \delta]_{Spin(2n)}(x_1, x_2, \dots, x_n) &= \prod_{i=1}^n (x_i^{1/2} + x_i^{-1/2}) \sum_{\delta/\mu: \text{vertical strip}} (-1)^{|\delta/\mu|} \mu_{Sp(2n)}(x_1, x_2, \dots, x_n) \\ &= \prod_{i=1}^n (x_i^{1/2} + x_i^{-1/2}) (-1)^{|\delta|} \delta_{SO(2n+1)}(-x_1, -x_2, \dots, -x_n) \end{aligned}$$

and

$$\begin{aligned} [\Delta', \delta]_{Spin(2n)}(x_1, x_2, \dots, x_n) &= \prod_{i=1}^n (x_i^{1/2} - x_i^{-1/2}) \sum_{\delta/\mu: \text{vertical strip}} \mu_{Sp(2n)}(x_1, x_2, \dots, x_n) \\ &= \prod_{i=1}^n (x_i^{1/2} - x_i^{-1/2}) \delta_{SO(2n+1)}(x_1, x_2, \dots, x_n) \end{aligned}$$

and for a partition μ with $\ell(\mu) = n$, we have the following.

$$\mu_{SO(2n)}^{(-)}(x_1, x_2, \dots, x_n) = \prod_{i=1}^n (x_i - x_i^{-1}) (\mu - (1^n))_{Sp(2n)}(x_1, x_2, \dots, x_n).$$

Here x_i denotes the (i, i) -th component of the diagonal matrices which we take as a maximal torus for each of the classical groups.

Moreover from Lemma 10.2 in [14], we have

$$\prod_{i=1}^n (x_i^{1/2} + x_i^{-1/2})^2 \delta_{Sp(2n)}(x_1, x_2, \dots, x_n) = \sum_{\substack{\kappa/\delta: \text{vertical strip} \\ \ell(\kappa) \leq n}} \kappa_{SO(2n+1)}(x_1, x_2, \dots, x_n).$$

If we write the above in the form

$$\frac{\prod_{i=1}^n (1 + x_i)^2}{x_1 x_2 \cdots x_n} \delta_{Sp(2n)}(x_1, x_2, \dots, x_n) = \sum_{\substack{\kappa/\delta: \text{vertical strip} \\ \ell(\kappa) \leq n}} \kappa_{SO(2n+1)}(x_1, x_2, \dots, x_n).$$

and substitute $-x_i$ for x_i , we have

$$(-1)^{n+|\delta|} \prod_{i=1}^n (x_i^{1/2} - x_i^{-1/2})^2 \delta_{Sp(2n)}(x_1, x_2, \dots, x_n) = \sum_{\substack{\kappa/\delta: \text{vertical strip} \\ \ell(\kappa) \leq n}} \kappa_{SO(2n+1)}(-x_1, -x_2, \dots, -x_n).$$

We note that $\delta_{Sp(2n)}(-x_1, -x_2, \dots, -x_n) = (-1)^{|\delta|} \delta_{Sp(2n)}(x_1, x_2, \dots, x_n)$.

So we have (for abbreviation, we write \mathbf{x} for x_1, x_2, \dots, x_n .)

$$\begin{aligned} & [\Delta, \delta]_{Spin(2n)} \mu_{SO(2n)}^{(-)} \\ &= \prod_{i=1}^n \{(x_i^{1/2} + x_i^{-1/2})(x_i - x_i^{-1})\} (\mu - (1^n))_{Sp(2n)}(\mathbf{x}) \sum_{\delta/\nu: \text{vertical strip}} (-1)^{|\delta/\nu|} \nu_{Sp(2n)}(\mathbf{x}) \\ &= \prod_{i=1}^n \{(x_i^{1/2} + x_i^{-1/2})(x_i - x_i^{-1})\} \sum_{\delta/\nu: \text{vertical strip}} (-1)^{|\delta/\nu|} {}^n C_{\nu, (\mu - (1^n))}^\lambda \lambda_{Sp(2n)}(\mathbf{x}) \end{aligned}$$

$$\text{Since } \prod_{i=1}^n (x_i^{1/2} + x_i^{-1/2})^2 \lambda_{Sp(2n)}(\mathbf{x}) = \sum_{\substack{\kappa/\delta: \text{vertical strip} \\ \ell(\kappa) \leq n}} \kappa_{SO(2n+1)}(\mathbf{x}), \text{ we}$$

have the formula (2.1.1).

Also we have

$$\begin{aligned}
& [\Delta', \delta]_{Spin(2n)} \mu_{SO(2n)}^{(-)} \\
&= \prod_{i=1}^n \{(x_i^{1/2} - x_i^{-1/2})(x_i - x_i^{-1})\} (\mu - (1^n))_{Sp(2n)}(\mathbf{x}) \sum_{\delta/\nu: \text{vertical strip}} \nu_{Sp(2n)}(\mathbf{x}) \\
&= \prod_{i=1}^n \{(x_i^{1/2} + x_i^{-1/2})(x_i^{1/2} - x_i^{-1/2})^2\} \sum_{\delta/\nu: \text{vertical strip}} {}^nC_{\nu, (\mu - (1^n))}^\lambda \lambda_{Sp(2n)}(\mathbf{x}) \\
&\text{Since } (-1)^{n+|\lambda|} \prod_{i=1}^n (x_i^{1/2} - x_i^{-1/2})^2 \lambda_{Sp(2n)}(\mathbf{x}) = \sum_{\substack{\kappa/\delta: \text{vertical strip} \\ \ell(\kappa) \leq n}} \kappa_{SO(2n+1)}(-x_1, -x_2, \dots, -x_n),
\end{aligned}$$

and $|\lambda| \equiv |\nu| + |(\mu - (1^n))| \pmod{2}$, we have the formula (2.1.2). \square

From the above and Theorem 1.2 of [14], we can deduce the followings. For abbreviation, we introduce the following convention. For $\epsilon \in \{\pm 1\}$, if $\epsilon = 1$, we consider $\Delta^\epsilon = \Delta^+$, $e_n^\epsilon = e_n^+$, $(1/2 + \delta)_{Spin(2n)}^\epsilon = (1/2 + \delta)_{Spin(2n)}^+$, etc. and if $\epsilon = -1$, $\Delta^\epsilon = \Delta^-$, $e_n^\epsilon = e_n^-$, $(1/2 + \delta)_{Spin(2n)}^\epsilon = (1/2 + \delta)_{Spin(2n)}^-$, etc.

Theorem 2.2. *For $\epsilon, \epsilon_1, \epsilon_2 \in \{\pm 1\}$, we have the following formulas.*

(i)

$$(2.2.1) \quad (\Delta^\epsilon)^2 = e_n^\epsilon + \sum_{0 \leq 2i \leq n} e_{n-2i}.$$

$$(2.2.2) \quad \Delta^+ \Delta^- = \sum_{0 \leq 2i \leq n-1} e_{n-1-2i}.$$

(ii) *For a partition δ with its length $\ell(\delta) < n$, we have*

$$\begin{aligned}
(2.2.3) \quad & (1/2 + \delta)_{Spin(2n)}^\epsilon (1)_{SO(2n)} \\
&= (1/2 + \delta)_{Spin(2n)}^{-\epsilon} + \sum_{\substack{\mu \supset \delta \\ |\mu/\delta|=1}} (1/2 + \mu)_{Spin(2n)}^\epsilon + \sum_{\substack{\delta \supset \mu \\ |\delta/\mu|=1}} (1/2 + \mu)_{Spin(2n)}^\epsilon.
\end{aligned}$$

Here μ/δ denotes the skew diagram.

If $\ell(\delta) = n$, we have

$$\begin{aligned}
(2.2.4) \quad & (1/2 + \delta)_{Spin(2n)}^\epsilon (1)_{SO(2n)} = \sum_{\substack{\mu \supset \delta \\ |\mu/\delta|=1, \\ \ell(\mu) \leq n}} (1/2 + \mu)_{Spin(2n)}^\epsilon + \sum_{\substack{\delta \supset \mu \\ |\delta/\mu|=1}} (1/2 + \mu)_{Spin(2n)}^\epsilon.
\end{aligned}$$

(iii) For a partition μ with $\ell(\mu) < n$, we have

(2.2.5)

$$\Delta^\epsilon \mu_{SO(2n)} = \sum_{\substack{\mu \supseteq \nu \\ \mu/\nu \text{ vertical strip} \\ |\mu/\nu| \equiv 0 \pmod 2}} (1/2 + \nu)_{Spin(2n)}^\epsilon + \sum_{\substack{\mu \supseteq \nu \\ \mu/\nu \text{ vertical strip} \\ |\mu/\nu| \equiv 1 \pmod 2}} (1/2 + \nu)_{Spin(2n)}^{-\epsilon}.$$

(iv) For a partition $\mu = (\mu_1, \mu_2, \dots, \mu_n)$ with $\ell(\mu) = n$, we write $\mu_{SO(2n)}^+ = \mu_{SO(2n)}$ and $\mu_{SO(2n)}^- = (\mu_1, \dots, \mu_{n-1}, -\mu_n)_{SO(2n)}$. Then we have

$$(2.2.6) \quad \Delta^{\epsilon_1} \mu_{SO(2n)}^{\epsilon_2} = \sum_{\substack{\mu \supseteq \nu \\ \mu/\nu \text{ vertical strip} \\ (-1)^{|\mu/\nu|} = \epsilon_1 \epsilon_2}} (1/2 + \nu)_{Spin(2n)}^{\epsilon_2}.$$

Particularly, for the exterior products e_i 's if $i < n$, we have

$$\Delta^\epsilon (1^i)_{SO(2n)} = \sum_{0 \leq 2s \leq i} (1/2 + (1^{i-2s}))_{Spin(2n)}^\epsilon + \sum_{0 \leq 2s \leq i-1} (1/2 + (1^{i-1-2s}))_{Spin(2n)}^{-\epsilon}$$

and for e_n^ϵ , we have

$$\Delta^\epsilon e_n^\epsilon = \sum_{0 \leq 2s \leq n} (1/2 + (1^{n-2s}))_{Spin(2n)}^\epsilon$$

and

$$\Delta^{-\epsilon} e_n^\epsilon = \sum_{0 \leq 2s \leq n-1} (1/2 + (1^{n-1-2s}))_{Spin(2n)}^\epsilon.$$

(v)

$$(e_n^+)^2 = \sum_{\substack{s \equiv n \pmod 2 \\ t \equiv 0 \pmod 2 \\ 0 \leq s+t \leq n}} (2^s, 1^t)_{SO(2n)}, \quad (e_n^+)(e_n^-) = \sum_{\substack{s \equiv n-1 \pmod 2 \\ t \equiv 0 \pmod 2 \\ s+t \leq n-1}} (2^s, 1^t)_{SO(2n)}.$$

$$(e_n^-)^2 = (2^{n-1}, -2)_{SO(2n)} + \sum_{\substack{s \equiv n \pmod 2 \\ s \leq n-2}} (2^s, 1^{n-s-1}, -1)_{SO(2n)} + \sum_{\substack{s \equiv n \pmod 2 \\ t \equiv 0 \pmod 2 \\ s+t \leq n-2}} (2^s, 1^t)_{SO(2n)}.$$

Proof. All these formulas can be deduced easily from Theorem 2.1 and Theorem 1.2 of [14]. We note that the diagram obtained by attaching a vertical strip to $(\mu - (1^n))$ is equal to the diagram obtained by removing the vertical strip from μ . \square

Remark 2.3. In Example 10.3 of [14], the formulas (2.2.3), (2.2.4) are stated in different forms, since the author considered there that if we allow μ_n to take the value -1 , the formulas become slightly of unified form. But this notation is not compatible with the other formulas in [14]. So it might cause some misunderstandings. Also the author must

apologize for writing down the incorrect formula in Example 10.3 of [14],

$$\Delta' \Delta' = e_n - e_{n-1} + e_{n-2} - \cdots + (-1)^{n-1} e_1 + (-1)^n.$$

The correct formula deduced from the formula (2.2.1), (2.2.2) in the Theorem thm repeven is

$$\Delta' \Delta' = e_n - 2e_{n-1} + 2e_{n-2} - \cdots + (-1)^{n-1} 2e_1 + (-1)^n 2.$$

The rest of the formulas are correct in Example 10.3 of [14].

For $Spin(2n+1)$, we quote here several formulas, which we can deduce easily from the formulas in the Theorem 1.2 in [14] and Theorem 3.1 in [12].

Theorem 2.4. *We have the following formulas.*

(i)

$$(2.4.1) \quad \Delta^2 = e_0 + e_1 + e_2 + \cdots + e_n.$$

(ii)

$$(2.4.2) \quad [\Delta, \delta]_{Spin(2n+1)}(1)_{SO(2n+1)} \\ = [\Delta, \delta]_{Spin(2n+1)} + \sum_{\substack{\mu \supset \delta \\ |\mu/\delta|=1, \ell(\mu) \leq n}} [\Delta, \mu]_{Spin(2n+1)} + \sum_{\substack{\delta \supset \mu \\ |\delta/\mu|=1}} [\Delta, \mu]_{Spin(2n+1)}.$$

Here $\ell(\mu)$ denotes the length of μ and μ/δ denotes the skew diagram.

(iii) For any partition λ with $\ell(\lambda) \leq n$, we have

$$(2.4.3) \quad \Delta \otimes \lambda_{SO(2n+1)} = \sum_{\substack{\lambda \supseteq \mu \\ \lambda/\mu: \text{vertical strip}}} [\Delta, \mu]_{Spin(2n+1)}.$$

Hence $\Delta \otimes \lambda_{SO(2n+1)}$ contains Δ if and only if $\lambda = (1^k)$ for some k ($1 \leq k \leq n$). In this case, we have

$$\Delta \otimes (1^k)_{SO(2n+1)} = \sum_{i=0}^k [\Delta, (1^i)]_{Spin(2n+1)}.$$

3. CLIFFORD ALGEBRAS AND SPIN GROUPS

In this section we give explicit actions of the Lie group $Pin(N)$ and its Lie algebra $\mathfrak{so}(N\mathbb{R})$ on the spaces Δ and its dual Δ^* and $\bigwedge^i \mathbb{R}^N$ with respect to their specified bases, since later in the section 4, we need to show that the linear embedding from $\bigwedge^i \mathbb{R}^N$ to $\Delta \otimes \Delta^*$ given explicitly there is equivariant homomorphism by verifying the actions of generators of $Pin(N)$ or $Pin(N)$ on both sides. As for the reference, see [5].

We consider the spin groups and their Lie algebra mainly over \mathbb{R} .

Let V be a vector space of dimension N ($N = 2n+1$ or $N = 2n$) over the real number field \mathbb{R} and we take a basis $< u_1, u_2, \dots, u_n, (u_0), u_{\bar{n}}, \dots, u_{\bar{1}} >$

of V . Here $\{1, 2, \dots, n, (0), \bar{n}, \dots, \bar{1}\}$ is an index set and we consider that for the case $N = 2n$, we omit the index 0 and that for the case $N = 2n + 1$, we add the index 0. We introduce an order in the index set such that $1 < 2 < \dots < n (< 0) < \bar{n} < \bar{n} - 1 < \dots < \bar{1}$.

Let $B(\ , \)$ be a non-degenerate symmetric bilinear form on V defined by

$$B(u_k, u_\ell) = 0, \quad B(u_{\bar{k}}, u_{\bar{\ell}}) = 0, \quad B(u_k, u_{\bar{\ell}}) = \frac{1}{2} \delta_{k, \ell} (k, \ell > 0). \\ (B(u_0, u_k) = 0, \quad B(u_0, u_{\bar{k}}) = 0 \quad (k > 0), \quad B(u_0, u_0) = 1.)$$

In the tensor algebra $\mathfrak{T}(V) = \sum_{i=0}^{\infty} \bigotimes^i V$, let \mathfrak{I}_B be an ideal generated by $\langle v \otimes v - B(v, v) \rangle_{v \in V}$ (here v runs over all the elements in V .) and we define the Clifford algebra $Cl(V)$ by the quotient $Cl(V) = \mathfrak{T}(V)/\mathfrak{I}_B$.

We note that $\langle u_k + u_{\bar{k}}, \quad u_k - u_{\bar{k}} \quad (k > 0), \quad (u_0) \rangle$ is an orthogonal basis of V and $B(u_k + u_{\bar{k}}, u_k + u_{\bar{k}}) = 1, \quad B(u_k - u_{\bar{k}}, u_k - u_{\bar{k}}) = -1, \quad (B(u_0, u_0) = 1)$.

Then $Pin(N)$ and $Spin(N)$ are defined by

Definition 3.1.

$$Pin(N, B) := \{\pm v_1 v_2 \dots v_r; v_i \in V, B(v_i, v_i) = \pm 1\}$$

and

$$Spin(N, B) := \{\pm v_1 v_2 \dots v_{2r}; v_i \in V, B(v_i, v_i) = \pm 1\}.$$

Remark 3.2. Since the quadratic form B is the form of maximal index ([5]), that is, it has the signature (n, n) for $N = 2n$ and $(n + 1, n)$ for $N = 2n + 1$, $Pin(N, B)$ and $Spin(N, B)$ are not compact groups.

The Lie algebra $L(Pin(N)) = L(Spin(N)) (\cong \mathfrak{so}(N, B))$ in $Cl(V)$ is generated by the degree 2 elements $\langle u_k u_\ell \rangle, (k \neq \ell)$.

From now on, we simply write $Pin(N)$ and $Spin(N)$ for $Pin(N, B)$ and $Spin(N, B)$ because we fix the defining quadratic form B for the Clifford algebras.

We define the subspace $V_k = \mathbb{R}u_k + \mathbb{R}u_{\bar{k}} \quad (k = 1, 2, \dots, n)$ (and $V_0 = \mathbb{R}u_0$) of V . Then $B|_{V_k}$ is non-degenerate and $Cl(V_k)$ is isomorphic to all the 2×2 matrices $M(2, \mathbb{R})$. This isomorphism ϕ_2 is given by $\phi_2(u_k) = X$ and $\phi_2(u_{\bar{k}}) = Y$ and $\phi(u_k u_{\bar{k}} - u_{\bar{k}} u_k) = H$, here $\{X, Y, H\}$ is the standard triple in $\mathfrak{sl}(2)$ and is given by

$$X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Since $V = V_1 \oplus V_2 \oplus \dots \oplus V_n (\oplus V_0)$ is the orthogonal direct sum, we have the following isomorphism in this case.

$$Cl(V) \cong Cl(V_1) \otimes Cl(V_2) \otimes \dots \otimes Cl(V_n) (\otimes Cl(V_0)).$$

Here $Cl(V_k)$ is isomorphic to $M(2, \mathbb{R})$ for $k > 0$ and $Cl(V_0)$ is a two dimensional semisimple commutative algebra.

Remark 3.3. In [5] (see p38 Theorem 3.3.), the tensor products of the clifford algebras are considered in the category of graded algebras and there appear the signatures in the products. But here in the above we consider the tensor products in the usual (not graded) ways. So we need to say about the above isomorphism.

Lemma 3.4 ([20]). *Let (V, B) be a pair of vector space V over \mathbb{R} and a non-degenerate symmetric bilinear form B on V . Let V_1, V_2 be a vector subspace of V such that (V, B) is the orthogonal direct sum of $(V_1, B|_{V_1})$ and $(V_2, B|_{V_2})$. Let α be the involutive automorphism of $Cl(V_1)$ i.e., $\alpha(v_1) = -v_1$ for any elements $v_1 \in V_1$.*

If there exists an involutive element $\theta \in Cl(V_1)$ such that $Ad(\theta) = \alpha$, (i.e. $\theta^2 = 1$ and $\alpha(x) = \theta x \theta^{-1}$ for any $x \in Cl(V_1)$.) then we have

$$Cl(V) \cong Cl(V_1) \otimes Cl(V_2).$$

Here the tensor product is an ordinary tensor product (i.e., not graded one) and the isomorphism Φ from $Cl(V)$ to $Cl(V_1) \otimes Cl(V_2)$ is given by

$$\Phi(u) = \begin{cases} u \otimes 1 & \text{if } u \in V_1 \\ \theta \otimes u & \text{if } u \in V_2. \end{cases}$$

Proof. Since both the algebras have the same dimension, it is enough to show that Φ is surjective homomorphism.

For any $v = (v_1, v_2) \in V_1 \oplus V_2$ ($v_i \in V_i$), $\Phi(v)\Phi(v) = (v_1 \otimes 1 + \theta \otimes v_2)^2 = v_1^2 \otimes 1 + \theta^2 \otimes v_2^2 + v_1 \theta \otimes v_2 + \theta v_1 \otimes v_2 = (B(v_1, v_1) + B(v_2, v_2))1 \otimes 1$, since $\alpha(v_1) = -v_1 = \theta v_1 \theta^{-1}$. From the universality, Φ becomes a homomorphism and $Cl(V_1)$ is included in the image of Φ , so $\theta \otimes 1$ is in the image. Hence Φ is surjective. \square

For the orthogonal direct sum $V = V_1 \oplus V_2 \oplus \cdots \oplus V_n (\oplus V_0)$, the above θ in the lemma for each direct summand is given by $\theta = u_k u_{\bar{k}} - u_{\bar{k}} u_k$, $\phi(\theta) = H$.

From the above, $V_1 \otimes V_2 \otimes \cdots \otimes V_n (\otimes V_0)$ becomes a representation space of $Cl(V)$.

The central primitive idempotents in $Cl(V_0)$ are given by $\frac{1 \pm u_0}{2}$.

Then u_0 acts on $V_0^\pm = \mathbb{R} \frac{1 \pm u_0}{2}$ by ± 1 .

And so if $\dim V = 2n + 1$, for the Clifford algebra $Cl(V)$, we have the two different irreducible representations ϕ_\pm according to V_0^\pm .

Since the dimensions of V_0^\pm are one, so we omit the last tensor and define the representation space of ϕ_\pm by the same $\Delta = V_1 \otimes V_2 \otimes \cdots \otimes V_n$.

If $\dim V = 2n$, $Cl(V)$ is a central simple algebra over \mathbb{R} and the above Δ is the unique irreducible representation of $Cl(V)$ and we denote it by ϕ .

Explicit representation matrices of ϕ, ϕ_\pm are given in the following.

Lemma 3.5. *The representation matrices of ϕ, ϕ_{\pm} on $\Delta = V_1 \otimes V_2 \otimes \cdots \otimes V_n$ are given as follows.*

$$\begin{aligned} u_k &\longmapsto H \otimes \cdots \otimes H \otimes \overset{k}{X} \otimes 1_2 \cdots \otimes 1_2, \\ u_{\bar{k}} &\longmapsto H \otimes \cdots \otimes H \otimes \overset{k}{Y} \otimes 1_2 \cdots \otimes 1_2, \\ (u_0 &\longmapsto \pm H \otimes H \otimes H \otimes \cdots \otimes H,) \end{aligned}$$

, where $k \in \{1, 2, \dots, n\}$ and 1_2 denotes the identity matrix of size 2 and the sign of the image of u_0 is +1 for ϕ_+ and -1 for ϕ_- .

If $N = 2n$, Δ is not irreducible as $Spin(2n)$ module and is decomposed into two irreducible representations Δ^+ and Δ^- . We note that Δ is irreducible as a $Pin(2n)$ module.

If $N = 2n + 1$ and ϕ_{\pm} are restricted to $Spin(2n + 1)$, it is known that they give equivalent irreducible representations of $Spin(2n + 1)$. As $Pin(2n + 1)$ modules, they are different in the tensor product with the linear character \det . By ‘ \det ’, we denote the linear representation of $Pin(N)$ induced from the natural homomorphism $Pin(N) \rightarrow O(N) \xrightarrow{\det} \{\pm 1\}$.

We fix a base of the representation space $\Delta = V_1 \otimes V_2 \otimes \cdots \otimes V_n$. From the definition, the set of vectors $u_{p_1} \otimes u_{p_2} \otimes \cdots \otimes u_{p_n}$ ’s (here $p_1 \in \{1, \bar{1}\}, p_2 \in \{2, \bar{2}\}, \dots, p_n \in \{n, \bar{n}\}$) becomes a basis. We pick up the indices $\mathbf{I} = \{i_j | p_{i_j} = \bar{i}_j\}$ and denote the above vector $u_{p_1} \otimes \cdots \otimes u_{p_n}$ by $[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r] = [\mathbf{I}]$, where $\mathbf{I} = \{i_1, i_2, \dots, i_r\}$ and $i_1 < i_2 < \dots < i_r$.

From Lemma 3.5, we can write down the action u_k and $u_{\bar{k}}$ on this base. Since $Hu_k = u_k$, $Hu_{\bar{k}} = -u_{\bar{k}}$, $Xu_k = 0$, $Xu_{\bar{k}} = u_k$, and $Yu_k = u_{\bar{k}}$, $Yu_{\bar{k}} = 0$ (H, X, Y is the standard triple in $\mathfrak{sl}(2, \mathbb{R})$), we have the following lemma.

Lemma 3.6.

$$u_k[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r] = \begin{cases} (-1)^{s-1}[\mathbf{i}_1, \dots, \widehat{\mathbf{i}}_s, \dots, \mathbf{i}_r] & \text{if } k = i_s \text{ for some } \mathbf{i}_s \\ 0 & \text{otherwise,} \end{cases}$$

where $\widehat{\mathbf{i}}_s$ denotes the removal of the index i_s from the index set and

$$u_{\bar{k}}[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r] = \begin{cases} (-1)^{s-1}[\mathbf{i}_1, \dots, \mathbf{i}_{s-1}, \mathbf{k}, \mathbf{i}_s, \dots, \mathbf{i}_r] & \text{if } k \notin \mathbf{I} \text{ and } i_{s-1} < k < i_s \\ 0 & \text{otherwise.} \end{cases}$$

$$(u_0[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r] = \pm(-1)^r[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r] \quad \text{according to } \phi_{\pm}.)$$

This representation is sometimes called the Fock space representation since it is realized as the action on the exterior algebra of the isotropic subspace of V generated by $u_{\bar{1}}, u_{\bar{2}}, \dots, u_{\bar{n}}$. That is, under the correspondence of $[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r] = u_{\bar{i}_1} \wedge u_{\bar{i}_2} \wedge \dots \wedge u_{\bar{i}_r}$, u_k acts on

this space by the interior product (here we must adjust the inner product such as $(u_k, u_{\bar{\ell}}) = \delta_{k,\ell}$.) and $u_{\bar{k}}$ acts on this space by the exterior product $u_{\bar{k}} \wedge$ and u_0 acts by the degree operator $(-1)^{\deg}$.

Therefore for later convenience, we introduce the following convention. For the indices in the bracket $[i_1, i_2, i_3, \dots, i_r]$ which are not necessarily in increasing order, we correspond this element to 0 if the indices are not distinct, otherwise we correspond this to the signed base element $\epsilon(\sigma)[i_{\sigma(1)}, i_{\sigma(2)}, i_{\sigma(3)}, \dots, i_{\sigma(r)}]$, where σ is the permutation of $\{1, 2, \dots, r\}$ such that $i_{\sigma(1)} < i_{\sigma(2)} < i_{\sigma(3)} < \dots < i_{\sigma(r)}$ and $\epsilon(\sigma)$ denotes the signature of σ . This convention is compatible with the base of the Fock space representation.

The explicit action of the Lie algebra $L(\text{Spin}(N, B))$ on V is given by

$$\text{ad}(uv)w = [uv, w] = uvw - wuv = 2B(w, v)u - 2B(w, u)v,$$

where $u, v, w \in V$ and the product here is that of the Clifford algebra, therefore we can calculate the explicit actions of the standard root vectors of $\mathfrak{so}(N)$ with respect to the base $\langle u_1, \dots, u_n, (u_0), u_{\bar{n}}, \dots, u_{\bar{1}} \rangle$ of V . The standard root vectors of $L(\text{Spin}(N))$ for the simple roots are given by $X_k = u_k u_{\overline{k+1}}$, $Y_k = u_{k+1} u_{\bar{k}}$ ($1 \leq k \leq n-1$) and $h_i = \frac{u_i u_{\bar{i}} - u_{\bar{i}} u_i}{2}$ ($1 \leq i \leq n$) and for $L(\text{Spin}(2n+1))$, $X_n = u_n u_0$, $Y_n = u_0 u_{\bar{n}}$ and for $L(\text{Spin}(2n))$, $X_n = u_{n-1} u_n$, $Y_n = u_{\bar{n}} u_{\overline{n-1}}$. Therefore the actions of the above root vectors on V and Δ are given as follows.

Lemma 3.7. *For k ($1 \leq k \leq n-1$) and i ($1 \leq i \leq n$), we have*

$$\begin{aligned} \text{ad}(X_k) &= E_{k,k+1} - E_{\overline{k+1}, \bar{k}}, & d\phi_{\pm}(X_k) &= -1_2 \otimes \dots \otimes 1_2 \otimes \overset{k}{X} \otimes Y \otimes 1_2 \otimes \dots \otimes 1_2, \\ \text{ad}(Y_k) &= E_{k+1,k} - E_{\bar{k}, \overline{k+1}}, & d\phi_{\pm}(Y_k) &= -1_2 \otimes \dots \otimes 1_2 \otimes \overset{k}{Y} \otimes X \otimes 1_2 \otimes \dots \otimes 1_2, \\ \text{ad}(h_i) &= E_{i,i} - E_{\bar{i}, \bar{i}}, & d\phi_{\pm}(h_i) &= \frac{1}{2} 1_2 \otimes \dots \otimes 1_2 \otimes \overset{i}{H} \otimes 1_2 \otimes \dots \otimes 1_2. \end{aligned}$$

And for $L(\text{Spin}(2n+1))$, we have

$$\begin{aligned} \text{ad}(X_n) &= 2E_{n,0} - E_{0,\bar{n}}, & d\phi_{\pm}(X_n) &= \mp 1_2 \otimes 1_2 \otimes \dots \otimes 1_2 \otimes X, \\ \text{ad}(Y_n) &= E_{0,n} - 2E_{\bar{n},0}, & d\phi_{\pm}(Y_n) &= \mp 1_2 \otimes 1_2 \otimes \dots \otimes 1_2 \otimes Y. \end{aligned}$$

And for $L(\text{Spin}(2n))$, we have

$$\begin{aligned} \text{ad}(X_n) &= E_{n-1,\bar{n}} - E_{\bar{n}, \overline{n-1}}, & d\phi(X_n) &= -1_2 \otimes \dots \otimes 1_2 \otimes X \otimes X, \\ \text{ad}(Y_n) &= E_{\bar{n}, n-1} - E_{\overline{n-1}, n}, & d\phi(Y_n) &= -1_2 \otimes \dots \otimes 1_2 \otimes Y \otimes Y. \end{aligned}$$

Then for any $1 \leq k \leq n-1$, we have $[X_k, Y_k] = h_k - h_{k+1} = H_{\alpha_k}$ and for $L(\text{Spin}(2n+1))$, $[X_n, Y_n] = 2h_n = 2H_{\alpha_n}$ and for $L(\text{Spin}(2n))$, $[X_n, Y_n] = h_{n-1} + h_n = H_{\alpha_n}$.

For $\text{Spin}(2n)$, ad is a Lie algebra isomorphism from $L(\text{Spin}(2n))$ to $\mathfrak{so}(V, B)$. Since $B = \frac{1}{2}S$ with respect to the fixed basis (S is the

defining symmetric form of $\mathfrak{so}(N, \mathbb{R})$ in the section 2.) , $\mathfrak{so}(V, B)$ is identified with the Lie algebra $\mathfrak{so}(\mathbb{R}^{2n}, S)$.

So from now on, we consider V has the standard basis $\langle u_i, u_{\bar{i}} \rangle$ such that the defining symmetric bilinear form is given by S . Namely V has the inner product $(\ , \)$ such that $(u_k, u_{\bar{\ell}}) = \delta_{k,\ell}$ and the values of the inner products between the other base elements are all 0.

For $Spin(2n+1)$, we need to change basis. Let us put $u_{0'} = \frac{1}{\sqrt{2}}u_0$.

Then $B(u_{0'}, u_{0'}) = \frac{1}{2}$ and we take a new basis $\langle u_i, u_{0'}, u_{\bar{i}} \rangle$ of V . The representation matrices of X_n and Y_n under 'ad' change into $\text{ad}(u_n u_0) = \text{ad}(X_n) = \sqrt{2}(E_{n,0'} - E_{0',\bar{n}})$ and $\text{ad}(u_0 u_{\bar{n}}) = \text{ad}(Y_n) = \sqrt{2}(E_{0',n} - E_{\bar{n},0'})$. Then ad is a Lie algebra isomorphism from $L(Spin(2n+1))$ to $\mathfrak{so}(V, B)$ and the symmetric matrix corresponding to B on the new base is given by $\frac{1}{2}S$.

So similarly as above, we consider V has the standard basis $\langle u_i, u_{0'}, u_{\bar{i}} \rangle$ such that the defining symmetric bilinear form is given by S . Namely V has the inner product $(\ , \)$ such that $(u_k, u_{\bar{\ell}}) = \delta_{k,\ell}$, $(u_{0'} u_{0'}) = 1$ and the other inner products between these basis elements are all 0.

From Lemma 3.7, We can write down the action of X_i 's and Y_i 's explicitly on the fundamental spin representation space Δ (the dual of Δ) of $\mathfrak{so}(N)$ with respect to the basis $[i_1, i_2, \dots, i_r]$'s.

Lemma 3.8. *In this lemma, \mathbf{I} denotes the index set $\{i_1, i_2, \dots, i_r\}$, where $i_1, i_2, i_3, \dots, i_r$ are in the increasing order. If $1 \leq k \leq n-1$, we can write down the actions of X_k 's and Y_k 's simultaneously for $\mathfrak{so}(N)$ with respect to the basis $[i_1, i_2, \dots, i_r]$'s as follows.*

$$X_k[\mathbf{I}] = \begin{cases} -[i_1, \dots, i_{s-1}, k+1, i_{s+1}, \dots, i_r] & \text{if } k = i_s \text{ and } k+1 < i_{s+1} \\ 0 & \text{otherwise,} \end{cases}$$

and

$$Y_k[\mathbf{I}] = \begin{cases} -[i_1, \dots, i_{s-1}, k, i_{s+1}, \dots, i_r] & \text{if } k+1 = i_s \text{ and } k > i_{s-1} \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{and } h_i[\mathbf{I}] = -\frac{1}{2}[\mathbf{I}] \text{ if } i \in \mathbf{I} \text{ and } h_i[\mathbf{I}] = \frac{1}{2}[\mathbf{I}] \text{ if } i \notin \mathbf{I}.$$

For $L(Spin(2n+1))$, we have

$$d\phi_{\pm}(X_n)[i_1, i_2, \dots, i_r] = \begin{cases} \mp[i_1, i_2, \dots, i_{r-1}] & \text{if } i_r = n \\ 0 & \text{otherwise,} \end{cases}$$

and

$$d\phi_{\pm}(Y_n)[i_1, i_2, \dots, i_r] = \begin{cases} \mp[i_1, i_2, \dots, i_r, n] & \text{if } i_r \neq n \\ 0 & \text{otherwise.} \end{cases}$$

For $\text{Spin}(2n)$, we have

$$X_n[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r] = \begin{cases} -[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_{r-2}] & \text{if } i_{r-1} = n-1 \text{ and } i_r = n \\ 0 & \text{otherwise,} \end{cases}$$

and

$$Y_n[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r] = \begin{cases} -[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r, n-1, n] & \text{if } i_r < n-1 \\ 0 & \text{otherwise} \end{cases}$$

Remark 3.9. As noted before, for $\text{Spin}(2n+1)$, ϕ_\pm are equivalent if we restrict them to the group $\text{Spin}(2n+1)$ and the derivations $d\phi_\pm$ only differ in the sign of the representation matrices of X_n and Y_n , where X_n and Y_n are fixed short root vectors.

As $\mathfrak{so}(2n+1)$ modules, if we take a new base $\{[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]_- = (-1)^r[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]\}$ in the representation space of $d\phi_-$, then the representation matrices of X_i, Y_i, h_i under $d\phi_-$ with respect to this new base coincide with those of them under $d\phi_+$.

So hereafter, in the representation space of ϕ_- for $\text{Pin}(2n+1)$ (and in its dual space), we only consider the elements $[\mathbf{i}_1, \dots, \mathbf{i}_r]_-$ instead of $[\mathbf{i}_1, \dots, \mathbf{i}_r]$.

And from now on, we change the definition of the notation $[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]$ in the representation space ϕ_- and by $[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]$, we denote the element $[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]_-$ in ϕ_- .

For $\dim V = 2n+1$, the center $Z(\text{Cl}(V))$ of the Clifford algebra $\text{Cl}(V)$ is given by $Z(\text{Cl}(V)) = \mathbb{R} \oplus \mathbb{R}z$, where $z = (u_1 - u_{\overline{1}})(u_1 + u_{\overline{1}}) \cdots (u_n - u_{\overline{n}})(u_n + u_{\overline{n}})u_0$. We have $z^2 = 1$ and z acts on V by the scalar -1 and acts on the representation spaces ϕ_\pm by the scalar ± 1 .

The Clifford theorem tells us that there are exactly two non-equivalent irreducible representations of $\text{Pin}(2n+1)$ with the same character and that one is obtained from the other by the tensor product with the linear character \det .

Since $z \notin \text{Spin}(2n+1)$, we can distinguish them by the action of z . By $[\Delta, \delta]_{\text{Pin}(2n+1), \pm}$, we denote the irreducible representation of $\text{Pin}(2n+1)$ with its character $[\Delta, \delta]_{\text{Spin}(2n+1)}$, on which z acts by the scalar ± 1 .

For the representation space of ϕ_\pm , we simply denote them by Δ_\pm instead of $\Delta_{\text{Pin}(2n+1), \pm}$. We have $(\Delta_\pm)^* \cong \Delta_\pm$.

In this paper, for abbreviation, Δ stands for Δ_+ or Δ_- with the base $\{[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]\}$ noted above, if not specified.

For $\mathfrak{so}(2n)$, from Lemma 3.8, the parity of the number of the indices in the base element $[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r]$ under the above action is preserved. We define Δ^+ to be the linear space spanned by all the base elements with even parity of the number of indices and Δ^- to be the linear space spanned by all the base elements with odd parity of indices. Then the highest weight vector with the highest weight

$\frac{1}{2}(\mathbf{e}_1 + \mathbf{e}_2 + \cdots + \mathbf{e}_n)$ of Δ^+ is given by $[\emptyset]$ and the highest weight vector with the highest weight $\frac{1}{2}(\mathbf{e}_1 + \cdots + \mathbf{e}_{n-1} - \mathbf{e}_n)$ of Δ^- is given by $[\mathbf{n}]$. As for the dual representation $(\Delta^+)^*$ and $(\Delta^-)^*$, if $n \equiv 0 \pmod{2}$, we have $(\Delta^+)^* \cong \Delta^+$ and $(\Delta^-)^* \cong \Delta^-$. If $n \equiv 1 \pmod{2}$, we have $(\Delta^+)^* \cong \Delta^-$ and $(\Delta^-)^* \cong \Delta^+$.

As a $Pin(2n)$ module, the action of the element $u_k \pm u_{\bar{k}}$ in $Pin(2n)$ on Δ is given by

$$\phi(u_k \pm u_{\bar{k}}) = H \otimes \cdots H \otimes (X \pm Y) \otimes 1_2 \otimes \cdots \otimes 1_2$$

The adjoint action of the element $u_n - u_{\bar{n}}$ on the Lie algebra $L(Spin(2n))$ in the Clifford algebra $Cl(V)$ gives the standard outer automorphism of $Spin(2n)$ and the sum characters $[\Delta, \delta]_{Spin(2n)}$ are the irreducible characters for $Pin(2n)$.

Also the element $u_n - u_{\bar{n}}$ acts on Δ by $(u_n - u_{\bar{n}})\Delta^\pm = \Delta^\mp$ and $(u_n - u_{\bar{n}})^2 = -\text{id}$. Then the Clifford theorem tells us that as $Pin(2n)$ modules, $(\Delta \otimes \det) \cong \Delta$. The associator A which realizes this isomorphism is, by definition, the linear isomorphism on $\Delta = \Delta^+ \oplus \Delta^-$ given by

$$A = \text{id}_{\Delta^+} \oplus -\text{id}_{\Delta^-}.$$

For, it is easy to see that $(u_n - u_{\bar{n}})A = -A(u_n - u_{\bar{n}})$ and A commutes with the action of $Spin(2n)$. This fact tells us that if two irreducible representations λ and $\tilde{\lambda}$ of $O(2n)$ are associated, i.e., $\lambda \cong \tilde{\lambda} \otimes \det$, we have the equivalent representation $\Delta \otimes \lambda \cong \Delta \otimes \tilde{\lambda}$ of $Pin(2n)$.

A acts on Δ by the degree operator $A[\mathbf{I}] = (-1)^{|\mathbf{I}|}[\mathbf{I}]$.

Also by the same reasoning, all the $[\Delta, \delta]_{Pin(2n)}$ become self-associated since $[\Delta, \delta]_{Pin(2n)}$ is decomposed into two non-equivariant representations under $Spin(2n)$ and the Clifford theorem tells us that it occurs if and only if $[\Delta, \delta]_{Pin(2n)}$ is self-associated. (See p161 in [21].) So we have $[\Delta, \delta]_{Pin(2n)} \cong [\Delta, \delta]_{Pin(2n)} \otimes \det$. This fact means that the characters $[\Delta, \delta]_{Spin(2n)}$ uniquely determine the irreducible representations of $Pin(2n)$.

Since $[\Delta, \delta] = (1/2 + \delta)_{Spin(2n)}^+ \oplus (1/2 + \delta)_{Spin(2n)}^-$ and $(u_n - u_{\bar{n}})(1/2 + \delta)_{Spin(2n)}^\pm = (1/2 + \delta)_{Spin(2n)}^\mp$, the associator A is given by

$$A = \text{id}_{(1/2+\delta)_{Spin(2n)}^+} \oplus -\text{id}_{(1/2+\delta)_{Spin(2n)}^-}.$$

We write down the above as a lemma and use this fact later.

Lemma 3.10. *We have the isomorphism $[\Delta, \delta]_{Pin(2n)} \cong [\Delta, \delta]_{Pin(2n)} \otimes \det$. The associator A which realizes this $Pin(2n)$ isomorphism is given by*

$$A = \text{id}_{(1/2+\delta)_{Spin(2n)}^+} \oplus -\text{id}_{(1/2+\delta)_{Spin(2n)}^-}.$$

Hence the characters $[\Delta, \delta]_{Spin(2n)}$ uniquely determine the irreducible representations of $Pin(2n)$. Since $(\Delta \otimes \det) \cong \Delta$, we have $\Delta \otimes (\lambda \otimes \det) \cong \Delta \otimes \lambda$.

A compact real form $\mathfrak{so}(N)_{cpt}$ of $L(Spin(N)) \otimes \mathbb{C} \cong \mathfrak{so}(N, \mathbb{C})$ is generated by the elements $\sqrt{-1}h_i$ and $\sqrt{-1}(X_i + Y_i)$, $X_i - Y_i$, $(i = 1, 2, \dots, n)$ as a real Lie algebra.

Then we can introduce the invariant hermitian inner products with respect to $\mathfrak{so}(N)_{cpt}$ in the space V and $\Delta^{(\pm)}$ respectively. Such invariant hermitian forms are given explicitly as follows. If we define a hermitian inner product in V such that the basis $\langle u_i, (u_{0'}), u_{\bar{i}} \rangle$ are orthonormal basis, then we can show easily that the hermitian inner product becomes invariant under the action of $\mathfrak{so}(N)_{cpt}$. Also if we define a hermitian inner product on Δ such that the basis $\{[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r]\}$ ($\{i_1, i_2, \dots, i_r\}$ runs over all the subsets of $[1, n] = \{1, 2, \dots, n\}$.) is orthonormal basis, then we can show easily that the hermitian inner product becomes invariant under the action of $\mathfrak{so}(N)_{cpt}$. So we consider V , Δ , Δ^\pm are the hermitian metric spaces with these invariant hermitian metrics.

Let us consider the dual action of Δ and V . Let Δ^* and V^* be the dual vector space of Δ and V respectively and let $\langle u_i^*, (u_{0'}^*), u_{\bar{i}}^* \rangle$ and $\{[\mathbf{i}_1, \dots, \mathbf{i}_r]^*\}$ be the dual basis of $\langle u_i, (u_{0'}), u_{\bar{i}} \rangle$ and $\{[\mathbf{i}_1, \dots, \mathbf{i}_r]\}$ respectively. Since the representations matrices of the dual representations on the dual basis are given by the minus sign of the transposed matrices, we can write down the dual actions explicitly as in Lemma 3.7 and Lemma 3.8.

We fix some notations.

In the exterior product $\bigwedge^r V$, we have the natural basis $\{u_{i_1} \wedge u_{i_2} \wedge \dots \wedge u_{i_r}\}$, where $i_k \in \{1, \dots, n, (0'), \bar{n}, \dots, \bar{1}\}$. We denote these basis elements only by their indices

$$\langle \mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r \rangle = \frac{1}{r!} \sum_{\sigma \in \mathfrak{S}_r} \epsilon(\sigma) u_{i_{\sigma^{-1}(1)}} \otimes u_{i_{\sigma^{-1}(2)}} \otimes \dots \otimes u_{i_{\sigma^{-1}(r)}}.$$

We define an order in the index set by $1 < 2 < \dots < n < (0') < \bar{n} < \dots < \bar{1}$ and introduce the following convention. For any $\mathbf{I}, \mathbf{J}, \mathbf{W} \subseteq [1, n] = \{1, 2, \dots, n\}$, $\langle \mathbf{I}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{J}} \rangle$ denotes the exterior product $u_{\mathbf{I}} \wedge u_{\mathbf{W}} \wedge u_{\bar{\mathbf{W}}} \wedge u_{\bar{\mathbf{J}}}$, where for any subset \mathbf{K} of the index set, $u_{\mathbf{K}}$ denotes the exterior product $\langle \mathbf{k}_1, \mathbf{k}_2, \dots, \mathbf{k}_r \rangle$ for $\mathbf{K} = \{k_1, k_2, \dots, k_r\}$ and the indices are arranged in the increasing order defined above, namely $k_1 < k_2 < \dots < k_r$. Also we put $\langle \mathbf{I}, \mathbf{W}, 0', \bar{\mathbf{W}}, \bar{\mathbf{J}} \rangle = u_{\mathbf{I}} \wedge u_{\mathbf{W}} \wedge u_{0'} \wedge u_{\bar{\mathbf{W}}} \wedge u_{\bar{\mathbf{J}}}$.

For example, for subsets $\mathbf{I} = \{1, 4\}$ and $\mathbf{W} = \{2, 5\}$ and $\mathbf{J} = \{3, 6\}$ of $[1, 7]$, $\langle \mathbf{I}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{J}} \rangle$ denotes the exterior product $u_1 \wedge u_4 \wedge u_2 \wedge u_5 \wedge u_{\bar{3}} \wedge u_{\bar{6}} \wedge u_{\bar{7}}$.

Also for the base $\{[\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r]\}$ of Δ , we introduce a similar notation. Namely for mutually disjoint subsets \mathbf{I} and \mathbf{K} of $[1, n]$, $\{[\mathbf{I}, \mathbf{K}]\}$ denotes the element $[\mathbf{i}_1, \dots, \mathbf{i}_r, \mathbf{k}_1, \dots, \mathbf{k}_s]$, where the indices i_1, i_2, \dots ,

i_r are in the increasing order and the same for K . We note that the indices in the bracket have the alternating property. For the dual space Δ^* , We introduce the same notation $[J, K]^*$, conveying the same meaning.

Then $\{[I, K] \otimes [J, K]^*\}$ becomes a base of $\Delta \otimes \Delta^*$ if I and K and J run over all the mutually disjoint subsets of $[1, n]$.

Since V is a hermitian space, so the exterior products $\bigwedge^k V$ of V are naturally endowed with the hermitian inner product. Namely $\{u_{i_1} \wedge u_{i_2} \wedge \dots \wedge u_{i_r}\}$ becomes an orthonormal basis of $\bigwedge^k V$.

Since V has the $O(N)$ -invariant bilinear form S , we can identify the vector space V and its dual V^* .

Let ι be the \mathbb{R} -linear map from V to V^* defined by $\iota(u_k) = u_{\bar{k}}^*$. Here $k \in \{1, \dots, n, (0'), \bar{n}, \dots, \bar{1}\}$ and we consider $\bar{\bar{k}} = k, \bar{0}' = 0'$. Then $\iota : V \rightarrow V^*$ is an $O(N)$ -equivariant isomorphism. We also denote the $O(N)$ -equivariant isomorphism from V^* to V by the same notation ι , i.e., $\iota(u_k^*) = u_{\bar{k}}$.

The dual base is given by $\langle I, W, (0'), \bar{W}, \bar{J} \rangle^* = k! \langle I^*, W^*, (0')^*, \bar{W}^*, \bar{J}^* \rangle$ (here k is the degree of the exterior product and $*$ denotes the element obtained by replacing each tensor component by its dual base).

Definition 3.11. Let $\iota = (-1)^{\binom{k}{2}} \bigwedge^k \iota : \bigwedge^k V \rightarrow (\bigwedge^k V)^*$ be the $O(N)$ -equivariant isomorphism defined by

$$\iota(\langle I, W, (0'), \bar{W}, \bar{J} \rangle) = \frac{1}{k!} \langle J, W, (0'), \bar{W}, \bar{I} \rangle^*.$$

We also denote the inverse of the above by the same letter ι , that is, ι is also the map $\iota = (-1)^{\binom{k}{2}} \bigwedge^k \iota : (\bigwedge^k V)^* \rightarrow \bigwedge^k V$ defined by $\iota(\langle I, W, (0'), \bar{W}, \bar{J} \rangle^*) = k! \langle J, W, (0'), \bar{W}, \bar{I} \rangle$.

To prove the next lemma, we also introduce the linear map $\iota' : \bigwedge^\ell V \rightarrow \bigwedge^\ell V$.

Definition 3.12. $\iota'(\langle J, W, (0'), \bar{W}, \bar{I} \rangle) = \langle I, W, (0'), \bar{W}, \bar{J} \rangle$.

Then we can check easily that $\iota' X_k = -Y_k \iota'$, $\iota' Y_k = -X_k \iota'$ and $\iota' h_k = -h_k \iota'$ on $\bigwedge^\ell V$.

Lemma 3.13. The $\mathfrak{so}(2n+1)$ -isomorphism $r_\ell : \bigwedge^\ell V \rightarrow \bigwedge^{2n+1-\ell} V$ is given by follows.

$$r_\ell(\langle J, W, \bar{W}, \bar{I} \rangle) = (-1)^{|W|} \langle J, W^c, 0', \bar{W}^c, \bar{I} \rangle$$

and

$$r_\ell(\langle J, W, 0', \bar{W}, \bar{I} \rangle) = (-1)^{|W^c|} \langle J, W^c, \bar{W}^c, \bar{I} \rangle,$$

where $W \sqcup W^c = [1, n] - J - I$.

The $\mathfrak{so}(2n)$ -isomorphism $r_\ell : \bigwedge^\ell V \rightarrow \bigwedge^{2n-\ell} V$ is given by follows.

$$r_\ell(< J, W, \bar{W}, \bar{I} >) = (-1)^{|\bar{I}|} < J, W^c, \bar{W}^c, \bar{I} >,$$

where $W \sqcup W^c = [1, n] - J - I$.

Proof. For $\mathfrak{so}(2n+1)$, we note that we can deduce the above easily from the Theorem 4.1, but we also can check it directly as in the following argument. So we give proof for $\mathfrak{so}(2n)$ here.

We show that r_ℓ is $\mathfrak{so}(2n)$ -equivariant.

Then we have $\iota' \circ r_\ell = (-1)^\ell r_\ell \circ \iota'$, since $i + j + 2w = \ell$.

From the argument just before this lemma, it is enough to show that all the Y_k commute with r_ℓ , since $r_\ell X_k = -r_\ell \iota' Y_k \iota' = (-1)^{\ell+1} \iota' r_\ell Y_k \iota' = (-1)^{\ell+1} \iota' Y_k r_\ell \iota' = -\iota' Y_k \iota' r_\ell = X_k r_\ell$.

Let us prove $r_\ell Y_k(< J, W, \bar{W}, \bar{I} >) = Y_k r_\ell(< J, W, \bar{W}, \bar{I} >)$ for $k = 1, 2, \dots, n-1$.

Here the action of $\mathfrak{so}(2n)$ is the adjoint action and is given by $\text{ad}(Y_k) = E_{k+1,k} - E_{\bar{k}, \bar{k}+1}$.

Case 1. $k, k+1 \in J$ or $k, k+1 \in I$.

In this case, both sides are zero and the equality holds.

Case 2. $k \in J$ and $k+1 \in I$.

Then we have $k, k+1 \notin W$ and

$$\begin{aligned} Y_k(< J, W, \bar{W}, \bar{I} >) &= \epsilon < J - \{k\}, W + \{k+1\}, \overline{W + \{k+1\}}, \overline{I - \{k+1\}} > \\ &\quad - \epsilon < J - \{k\}, W + \{k\}, \overline{W + \{k\}}, \overline{I - \{k+1\}} > \end{aligned}$$

, where $\epsilon = (-1)^{|\{j \in J: j > k+1\}| + |\{i \in I: i > k+1\}|}$.

So we have

$$\begin{aligned} r_\ell Y_k(< J, W, \bar{W}, \bar{I} >) &= \epsilon (-1)^{|\bar{I}|-1} < J - \{k\}, W^c + \{k\}, \overline{W^c + \{k\}}, \overline{I - \{k+1\}} > \\ &\quad - \epsilon (-1)^{|\bar{I}|-1} < J - \{k\}, W^c + \{k+1\}, \overline{W^c + \{k+1\}}, \overline{I - \{k+1\}} >. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} Y_k r_\ell(< J, W, \bar{W}, \bar{I} >) &= \epsilon' < J - \{k\}, W^c + \{k+1\}, \overline{W^c + \{k+1\}}, \overline{I - \{k+1\}} > \\ &\quad - \epsilon' < J - \{k\}, W^c + \{k+1\}, \overline{W^c + \{k\}}, \overline{I - \{k\}} >. \end{aligned}$$

Here $\epsilon' = (-1)^{|\bar{I}| + |\{j \in J: j > k+1\}| + |\{i \in I: i > k+1\}|}$ and the equality holds.

Case 3. $k \in J$ and $k+1 \in W$.

Then we have

$$Y_k(< J, W, \bar{W}, \bar{I} >) = < J - \{k\} + \{k+1\}, W - \{k+1\} + \{k\}, \overline{W - \{k+1\} + \{k\}}, \bar{I} >.$$

So we have

$$r_\ell Y_k(< J, W, \bar{W}, \bar{I} >) = (-1)^{|\bar{I}|} < J - \{k\} + \{k+1\}, W^c, \bar{W}^c, \bar{I} >.$$

On the other hand, since $k, k+1 \notin W^c$, we have

$$Y_k r_\ell(< J, W, \bar{W}, \bar{I} >) = (-1)^{|\bar{I}|} < J - \{k\} + \{k+1\}, W^c, \bar{W}^c, \bar{I} >.$$

So we have the equality.

We omit the proof for the other cases since the proofs are almost similar.

Let us prove $r_\ell Y_n(< J, W, \bar{W}, \bar{I} >) = Y_n r_\ell(< J, W, \bar{W}, \bar{I} >)$, where $\text{ad}(Y_n) = E_{\bar{n}, n-1} - E_{\overline{n-1}, n}$.

Case 1. $n-1, n \in J$ or $n-1, n \in I$.

If $n-1, n \in J$, we have

$$\begin{aligned} Y_n(< J, W, \bar{W}, \bar{I} >) &= - < J - \{n\} - \{n-1\}, W + \{n-1\}, \overline{W + \{n-1\}}, \bar{I} > \\ &- < J - \{n-1\} - \{n\}, W + \{n\}, \overline{W + \{n\}}, \bar{I} > \text{ and} \\ r_\ell Y_n(< J, W, \bar{W}, \bar{I} >) &= (-1)^{|I|+1} (< J - \{n\} - \{n-1\}, W^c + \{n\}, \overline{W^c + \{n\}}, \bar{I} > \\ &+ < J - \{n-1\} - \{n\}, W^c + \{n-1\}, \overline{W^c + \{n-1\}}, \bar{I} >). \end{aligned}$$

The right hand of the above is equal to $Y_n r_\ell(< J, W, \bar{W}, \bar{I} >)$.

So we have the equality. The proof is almost same for the case $n-1, n \in I$.

Case 2. $n-1 \in J$ and $n \in I$.

In this case both sides are zero and the equality holds.

Case 3. $n-1 \in J$ and $n \in W$.

$$\begin{aligned} Y_n(< J, W, \bar{W}, \bar{I} >) &= - < J - \{n-1\}, W - \{n\} + \{n-1\}, \overline{W - \{n\} + \{n-1\}}, \bar{I} + \{n\} > \\ \text{and } r_\ell Y_n(< J, W, \bar{W}, \bar{I} >) &= (-1)^{|I|} (< J - \{n-1\}, W^c, \overline{W^c}, \bar{I} + \{n\} > . \end{aligned}$$

The right hand of the above is equal to $Y_n r_\ell(< J, W, \bar{W}, \bar{I} >)$.

Case 4. $n-1 \in J$ and $n \notin I, W$.

$$\begin{aligned} Y_n(< J, W, \bar{W}, \bar{I} >) &= < J - \{n-1\}, W, \bar{W}, \bar{I} + \{n\} > \text{ and } r_\ell Y_n(< J, W, \bar{W}, \bar{I} > \\ &= (-1)^{|I|+1} (< J - \{n-1\}, W^c - \{n\} + \{n-1\}, \overline{W^c - \{n\} + \{n-1\}}, \bar{I} + \{n\} > \\ & . \end{aligned}$$

Since $n \in W^c$, the right hand of the above is equal to $Y_n r_\ell(< J, W, \bar{W}, \bar{I} >)$.

The rest of the proof follows from similar direct verifications and we omit it. \square

From the above, we can define the explicit base elements of the irreducible constituents e_n^+ and e_n^- of $\bigwedge^n V$ under the action of $Spin(2n)$.

The spaces e_n^+ and e_n^- are given respectively by the ± 1 eigenspaces of the $Spin(2n)$ -equivariant involution r_n on $\bigwedge^n V$ defined in Lemma 3.13.

Definition 3.14. For an exterior product $< J, W, \bar{W}, \bar{I} >$ of degree n , we denote the basis elements of e_n^\pm respectively by

$$< J, W, \bar{W}, \bar{I} >^{(\pm)} = \frac{1}{\sqrt{2}} (< J, W, \bar{W}, \bar{I} > \pm (-1)^{|I|} < J, W^c, \overline{W^c}, \bar{I} >),$$

if the right-hand sides are non-zero.

In the case of $|I| + |J| = n$, we have $< J, \bar{I} >^{(+)} = \sqrt{2} < J, \bar{I} >$, $< J, \bar{I} >^{(-)} = 0$ if $|I| \equiv 0 \pmod{2}$ and $< J, \bar{I} >^{(-)} = \sqrt{2} < J, \bar{I} >$, $< J, \bar{I} >^{(+)} = 0$ if $|I| \equiv 1 \pmod{2}$.

Hence we have

$$\langle J, W, \bar{W}, \bar{I} \rangle = \frac{1}{\sqrt{2}} \left(\langle J, W, \bar{W}, \bar{I} \rangle^{(+)} + \langle J, W, \bar{W}, \bar{I} \rangle^{(-)} \right).$$

We introduce some other duality maps.

For $Spin(2n+1)$, since Δ is self-dual, the isomorphism $\iota_\Delta : \Delta \longrightarrow \Delta^*$ is given by $\iota_\Delta([I]) = (-1)^{(n+1)|I| - \sum I[I^c]^*}$. Here I^c is the complement of I in $\{1, 2, \dots, n\}$ i.e., $I \sqcup I^c = \{1, 2, \dots, n\}$ and $\sum I = \sum_{i \in I} i$ and $|I|$ denotes the number of the elements in I .

Then we can check easily $\iota_\Delta d\phi_+ = d\phi_+^* \iota_\Delta$.

Also for $Spin(2n)$, if n is even positive integer, Δ^\pm is self-dual and the isomorphism $\iota_\Delta : \Delta^\pm \longrightarrow (\Delta^\pm)^*$ is given by $\iota_\Delta([I]) = (-1)^{\sum I[I^c]^*}$.

If n is odd positive integer, then $(\Delta^\pm)^* \cong \Delta^\mp$ and the isomorphism $\iota_\Delta : \Delta^\pm \longrightarrow (\Delta^\mp)^*$ is given by $\iota_\Delta([I]) = \pm(-1)^{\sum I[I^c]^*}$.

Then we can check easily $\iota_\Delta d\phi = d\phi^* \iota_\Delta$.

Also we define an involutive linear automorphism ι' on $\Delta \otimes \Delta^*$ as follows.

Definition 3.15. For any disjoint subsets I, J in $[1, n]$, ι' is defined by

$$\iota'([I] \otimes [J]^*) = [J] \otimes [I]^*.$$

Then for $\mathfrak{so}(N)$, it is easily checked that $\iota'X_k = -Y_k \iota'$, $\iota'Y_k = -X_k \iota'$ and $\iota'h_k = -h_k \iota'$ on $\Delta \otimes \Delta^*$.

If $N = 2n$, ι' is the linear map satisfying $\iota'(\Delta^{\epsilon_1} \otimes (\Delta^{\epsilon_2})^*) = \Delta^{\epsilon_2} \otimes (\Delta^{\epsilon_1})^*$, where $\epsilon_1, \epsilon_2 \in \{\pm 1\}$.

4. THE EQUIVARIANT EMBEDDINGS OF $\bigwedge^k V$ TO $\Delta \otimes \Delta^*$

In this section we give the key theorems, which bridge the gap between the spin representations and the tensor calculus.

Theorem 4.1. For any k ($1 \leq k \leq 2n+1$), there exists an $\mathfrak{so}(2n+1)$ -equivariant embedding ϕ_k from $\bigwedge^k V$ to $\Delta \otimes \Delta^*$ defined as follows.

$$(4.1.1) \quad \phi_k(\langle J, W, \bar{W}, \bar{I} \rangle) = \sum_{[1, n] - J - I \supseteq K} \frac{(-1)^{|W - W \cap K|}}{2^{(n - |J| - |I|)/2}} [I, K] \otimes [J, K]^*$$

$$(4.1.2) \quad \phi_k(\langle J, W, O', \bar{W}, \bar{I} \rangle) = \sum_{[1, n] - J - I \supseteq K} \frac{(-1)^{|K - K \cap W|}}{2^{(n - |J| - |I|)/2}} [I, K] \otimes [J, K]^*$$

Moreover the above ϕ_k is an isometric embedding with respect to the invariant hermitian inner products.

We give proof of the above theorem in the last of this section.

Remark 4.2. Here Δ stands for Δ_+ or Δ_- and in the tensor parts corresponding to Δ_- , as for the bracket notation, we obey the convention of Remark 3.9.

If we compare the actions of the center z (see Remark 3.9.) on both sides, as $Pin(2n+1)$ modules, we have

$$(4.1) \quad \Delta_{\pm} \otimes \Delta_{\pm} \cong \bigoplus_{i=0}^n \bigwedge^{2i} V$$

and

$$(4.2) \quad \Delta_{\pm} \otimes \Delta_{\mp} \cong \bigoplus_{i=0}^n \bigwedge^{2i+1} V$$

and the above ϕ_k 's give $Pin(2n+1)$ -equivariant embeddings.

So for the group $Pin(2n+1)$, we have the following theorem.

Theorem 4.3. *For any k ($1 \leq k \leq 2n+1$), there exists a $Pin(2n+1)$ -equivariant isometric embedding ϕ_k from $\bigwedge^k V$ to $\Delta_{\epsilon_1} \otimes \Delta_{\epsilon_2}^*$ defined by (4.1.1) and (4.1.2) in Theorem 4.1. Here $\epsilon_1, \epsilon_2 \in \{\pm 1\}$ and $(-1)^k = \epsilon_1 \epsilon_2$.*

We move to the case $N = 2n$.

For $\epsilon_1, \epsilon_2 \in \{\pm 1\}$, by $\phi_{\ell}^{\epsilon_1 \epsilon_2}$, we denote an $\mathfrak{so}(2n)$ -equivariant embedding from $\bigwedge^{\ell} V$ to $\Delta^{\epsilon_1} \otimes (\Delta^{\epsilon_2})^*$. For example, $\phi_{\ell}^{\epsilon_1 \epsilon_2} = \phi_{\ell}^{+-}$ if $\epsilon_1 = 1$ and $\epsilon_2 = -1$ and so on.

Theorem 4.4. *If n is even, we have $(\Delta^+)^* \cong \Delta^+$ and $(\Delta^-)^* \cong \Delta^-$ and only the embeddings $\phi_{\ell}^{\epsilon_1 \epsilon_2}$ for any ℓ ($1 \leq \ell \leq 2n$) satisfying the condition $\epsilon_1 = (-1)^{n-\ell} \epsilon_2$ occur.*

If n is odd, we have $(\Delta^+)^ \cong \Delta^-$ and $(\Delta^-)^* \cong \Delta^+$ and only the embeddings $\phi_{\ell}^{\epsilon_1 \epsilon_2}$ for any ℓ ($1 \leq \ell \leq 2n$) satisfying the condition $\epsilon_1 = -(-1)^{n-\ell} \epsilon_2$ occur.*

For $\ell \neq n$, $\phi_{\ell}^{\epsilon_1 \epsilon_2}$ is given by

$$(4.4.1) \quad \phi_{\ell}^{\epsilon_1 \epsilon_2}(< J, W, \bar{W}, \bar{I} >) = \sum_{\substack{[1, n] - J - I \supseteq K \\ (-1)^{i+k} = \epsilon_1 \\ (-1)^{j+k} = \epsilon_2}} \frac{(-1)^{|W-W \cap K|}}{2^{(n-|J|-|I|-1)/2}} [I, K] \otimes [J, K]^*.$$

Here by small letters, we denote the number of the elements in the set indexed by their capital letters.

For e_n^+ and e_n^- , $\phi_n^{\epsilon_1 \epsilon_2}$ is given by

$$(4.4.2) \quad \phi_n^{\epsilon_1 \epsilon_2}(< J, W, \bar{W}, \bar{I} >^{(\epsilon_1)}) = \sum_{\substack{[1, n] - J - I \supseteq K \\ (-1)^{i+k} = \epsilon_1 \\ (-1)^{j+k} = \epsilon_2}} \frac{(-1)^{|W-W \cap K|}}{2^{(n-|J|-|I|-1)/2}} [I, K] \otimes [J, K]^*.$$

Moreover each of the above $\phi_{\ell}^{\epsilon_1 \epsilon_2}$'s is an isometric embedding with respect to the invariant hermitian inner products.

We give proofs of the above theorems for $Spin(2n+1)$ and $Spin(2n)$ in the last of this section simultaneously.

Remark 4.5. For e_n^+ and e_n^- , we note that only the above case occurs from (2.2.1) and (2.2.2).

The last statements in Theorem 4.1 and Theorem 4.4 follow easily from the following fact.

Since ϕ_k and $\phi_k^{\epsilon_1 \epsilon_2}$ become the $\mathfrak{so}(N, \mathbb{C})$ -equivariant map, then of course it becomes $\mathfrak{so}(N, \mathbb{C})_{cpt}$ equivariant map.

$\bigwedge^k V$ and e_n^+ and e_n^- are irreducible under the action of $\mathfrak{so}(N, \mathbb{C})_{cpt}$. Hence the invariant hermitian products only differ in a scalar multiple. If we take the highest weight vector of $\bigwedge^k V$ and compare its value of the hermitian inner product with that of its image, the last statements follow easily.

Before we prove the theorem for $Pin(2n)$, we state several properties, which follow from the theorems easily.

For $Spin(2n+1)$, from Theorem 4.1, we have the isomorphism $\bigoplus_{i=0}^n \bigwedge^{2i} V \cong \Delta \otimes \Delta^*$.

So we give the inverse of the above isomorphism.

Let us compare the same weight spaces in both sides. For simplicity, we omit ϕ_k since the degrees of the exterior products tell us which ϕ_k we are considering in the argument. Let $\mathbf{I} = \{i_1, i_2, \dots, i_r\}$ and $\mathbf{J} = \{j_1, j_2, \dots, j_s\}$ be disjoint subsets of $[1, n]$.

Let $\mathbf{e}_J - \mathbf{e}_I$ denote the weight $\mathbf{e}_J - \mathbf{e}_I = \mathbf{e}_{j_1} + \mathbf{e}_{j_2} + \dots + \mathbf{e}_{j_s} - \mathbf{e}_{i_1} - \mathbf{e}_{i_2} - \dots - \mathbf{e}_{i_r}$. Then a base of the weight space of weight $\mathbf{e}_J - \mathbf{e}_I$ in $\bigoplus_{i=0}^n \bigwedge^{2i} V$ is given by $\{< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >\}$ if $|\mathbf{J}| + |\mathbf{I}| \equiv 0 \pmod{2}$ and $\{< \mathbf{J}, \mathbf{W}, \mathbf{O}', \bar{\mathbf{W}}, \bar{\mathbf{I}} >\}$ if $|\mathbf{J}| + |\mathbf{I}| \equiv 1 \pmod{2}$. Here \mathbf{W} runs over all the subsets in $[1, n] - \mathbf{J} - \mathbf{I}$.

Also a base of the weight space of weight $\mathbf{e}_J - \mathbf{e}_I$ in $\Delta \otimes \Delta^*$ is given by $\{[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*\}$, where \mathbf{K} runs over all the subsets in $[1, n] - \mathbf{J} - \mathbf{I}$.

Since both bases are hermitian orthonormal bases of $\bigoplus_{i=0}^n \bigwedge^{2i} V$ and $\Delta \otimes \Delta^*$ respectively, the transformation matrices $\frac{1}{2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}} ((-1)^{|\mathbf{W}-\mathbf{W} \cap \mathbf{K}|})_{\mathbf{W}, \mathbf{K}}$ are unitary matrices. Moreover all the components in the above are real, so the transformation matrices are orthogonal matrices over \mathbb{R} , hence $H_{\mathbf{J}, \mathbf{I}, n} = ((-1)^{|\mathbf{W}-\mathbf{W} \cap \mathbf{K}|})_{\mathbf{W}, \mathbf{K}}$ (\mathbf{W} and \mathbf{K} run over all the subsets in $[1, n] - \mathbf{J} - \mathbf{I}$.) are Hadamard matrices of size $2^{[1, n] - \mathbf{J} - \mathbf{I}}$. These Hadamard matrices are equivalent to the tensor products of the standard Hadamard matrix of size 2.

So the inverse matrices of these matrices are given by transposing the original matrices.

Therefore if $|\mathbf{J}| + |\mathbf{I}| \equiv 0 \pmod{2}$, we have

$$[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{W}} \frac{(-1)^{|\mathbf{W}-\mathbf{W} \cap \mathbf{K}|}}{2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}} < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >$$

and if $|J| + |I| \equiv 1 \pmod{2}$, we have

$$[I, K] \otimes [J, K]^* = \sum_{[1, n] - J - I \supseteq W} \frac{(-1)^{|K - K \cap W|}}{2^{(n - |J| - |I|)/2}} \langle J, W, O', \bar{W}, \bar{I} \rangle.$$

Similarly for $Spin(2n)$, from Theorem 4.4, we have the isomorphism $e_n^{\epsilon_1} \oplus e_{n-2} \oplus \cdots \oplus e_0 \cong \Delta^{\epsilon_1} \otimes (\Delta^{\epsilon_2})^*$, here $\epsilon_1, \epsilon_2 \in \{\pm 1\}$ and $\epsilon_2 = (-1)^n \epsilon_1$.

We give the inverse of the above isomorphism.

As before, if we compare the same weight spaces in both sides, we obtain the inverse as follows. We also omit $\phi_k^{\epsilon_1 \epsilon_2}$ since the degrees of the exterior products and the parities of the indices in the brackets tell us which $\phi_k^{\epsilon_1 \epsilon_2}$ we are considering in the argument.

For the irreducible decomposition $\Delta^{\epsilon_1} \otimes (\Delta^{\epsilon_2})^*$ with $\epsilon_2 = (-1)^n \epsilon_1$, in the case of $(-1)^{k+i} = \epsilon_1$ and $(-1)^{k+j} = \epsilon_2$, we have

$$\begin{aligned} [I, K] \otimes [J, K]^* &= \sum_{\substack{[1, n] - J - I \supseteq W \\ i+j+2w \leq n-2}} \frac{(-1)^{|W - W \cap K|}}{2^{(n - |J| - |I| - 1)/2}} \langle J, W, \bar{W}, \bar{I} \rangle \\ &+ \sum_{\substack{[1, n] - J - I \supseteq W \\ i+j+2w = n}} \frac{(-1)^{|W - W \cap K|}}{2^{(n - |J| - |I| + 1)/2}} \langle J, W, \bar{W}, \bar{I} \rangle^{(\epsilon_1)}. \end{aligned}$$

We note that in the second sum, if $n - i - j > 0$, $(-1)^{|W - W \cap K|} \langle J, W, \bar{W}, \bar{I} \rangle^{(\epsilon_1)} = (-1)^{|W^c - W^c \cap K|} \langle J, W^c, \bar{W}^c, \bar{I} \rangle^{(\epsilon_1)}$ and we count it twice.

For the irreducible decomposition $\Delta^{\epsilon_1} \otimes (\Delta^{\epsilon_2})^*$ with $\epsilon_2 = -(-1)^n \epsilon_1$, in the case of $(-1)^{k+i} = \epsilon_1$ and $(-1)^{k+j} = \epsilon_2$, we have

$$[I, K] \otimes [J, K]^* = \sum_{\substack{[1, n] - J - I \supseteq W \\ i+j+2w \leq n-1}} \frac{(-1)^{|W - W \cap K|}}{2^{(n - |J| - |I| - 1)/2}} \langle J, W, \bar{W}, \bar{I} \rangle.$$

The matrix $((-1)^{|W - W \cap K|})_{K, W}$ is an Hadamard matrix of size $2^{n - |J| - |I| - 1}$, where W and K run over the subsets in $[1, n] - J - I$ satisfying $|W| \leq (n - 1 - i - j)/2$ and $(-1)^{|K| + i} = \epsilon_1$. These Hadamard matrices are also equivalent to the former ones.¹

Using the above theorem, we obtain the following $Pin(2n)$ ($O(2n)$) equivariant isometric embeddings.

Theorem 4.6. *As $Pin(2n)$ ($O(2n)$) modules, for any even ℓ ($1 \leq \ell \leq 2n$), there exists an equivariant isometric embedding $\phi_\ell : \bigwedge^\ell V \rightarrow \Delta \otimes \Delta^*$ given by*

$$\phi_\ell(\langle J, W, \bar{W}, \bar{I} \rangle) = \sum_{[1, n] - J - I \supseteq K} \frac{(-1)^{|W - W \cap K|}}{2^{(n - |J| - |I|)/2}} [I, K] \otimes [J, K]^*$$

¹We thank Prof. E. Bannai for this comment.

and for any odd ℓ ($1 \leq \ell \leq 2n$), there exists an equivariant isometric embedding $\phi_\ell : \bigwedge^\ell V \longrightarrow \Delta \otimes \Delta^*$ given by

$$\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >) = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}| + |\mathbf{I}| + |\mathbf{K}|}}{2^{(n - |\mathbf{J}| - |\mathbf{I}|)/2}} [\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*.$$

Remark 4.7. This theorem is crucial for us to consider the invariant theory for the group $\text{Pin}(2n)$ ($O(2n)$). The above ϕ_ℓ is defined as the sum $\frac{1}{\sqrt{2}}(\phi_\ell^{++} + \phi_\ell^{--})$ for even $\ell (\neq n)$ and $\frac{1}{\sqrt{2}}(\phi_\ell^{+-} - \phi_\ell^{-+})$ for odd $\ell (\neq n)$ and $\phi_n = \phi_n^{++} \oplus \phi_n^{--}$ for even n and $\phi_n = \phi_n^{+-} \oplus -\phi_n^{-+}$ for odd n .

Proof. From the above remark, ϕ_ℓ is $SO(2n)$ -equivariant. So in order to prove the theorem, it is enough to check that ϕ_ℓ is $(u_n - u_{\bar{n}})$ -equivariant.

We note that the action of $(u_n - u_{\bar{n}})$ on V is given by the transposition of u_n and $u_{\bar{n}}$.

We first prove the case for even $\ell (\neq n)$.

Case 1. $n \notin \mathbf{I}, \mathbf{J}, \mathbf{W}$.

Then we have $(u_n - u_{\bar{n}}) < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > = < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >$ and calculate $(u_n - u_{\bar{n}})\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >)$. We take cases for $n \in \mathbf{K}$ and $n \notin \mathbf{K}$ in the expansion of $\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >)$ in Theorem 4.6.

Case 1-1. $n \in \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{n\}$. From the Lemma 3.6, the action of $(u_n - u_{\bar{n}})$ is given as follows. (We note that in that lemma, the action is described only for the base elements with the increasing indices in the brackets.)

$$\begin{aligned} & (u_n - u_{\bar{n}})[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* \\ &= \epsilon \left(\begin{array}{cc} \mathbf{I}, & \mathbf{K} \\ \mathbf{I} + \mathbf{K}_1, & \{n\} \end{array} \right) \epsilon \left(\begin{array}{cc} \mathbf{J}, & \mathbf{K} \\ \mathbf{J} + \mathbf{K}_1, & \{n\} \end{array} \right) (-1)^{|\mathbf{I}| + |\mathbf{K}_1| + |\mathbf{J}| + |\mathbf{K}_1|} [\mathbf{I} + \mathbf{K}_1] \otimes [\mathbf{J} + \mathbf{K}_1]^* \\ &= (-1)^{|\mathbf{I}| + |\mathbf{J}|} [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1]^*. \end{aligned}$$

Here $\mathbf{I} + \mathbf{K}_1$ denotes the union of the set \mathbf{I} and \mathbf{K}_1 and we have the convention that the indices in the set $\mathbf{I} + \mathbf{K}_1$ are arranged in the increasing order. $\epsilon \left(\begin{array}{cc} \mathbf{I}, & \mathbf{K} \\ \mathbf{I} + \mathbf{K}_1, & \{n\} \end{array} \right)$ denotes the sign of the permutation obtained by arranging \mathbf{I}, \mathbf{K} into $\mathbf{I} + \mathbf{K}_1, \{n\}$ in this order.

Case 1-2. $n \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K}$. The action of $(u_n - u_{\bar{n}})$ is given by

$$\begin{aligned} & (u_n - u_{\bar{n}})[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* \\ &= \epsilon \left(\begin{array}{cc} \mathbf{I}, & \mathbf{K}_1 \\ \mathbf{I} + \mathbf{K}_1, & \end{array} \right) \epsilon \left(\begin{array}{cc} \mathbf{J}, & \mathbf{K}_1 \\ \mathbf{J} + \mathbf{K}_1, & \end{array} \right) (-1)^{|\mathbf{I}| + |\mathbf{K}_1| - 1 + |\mathbf{J}| + |\mathbf{K}_1| - 1} [\mathbf{I} + \mathbf{K}_1, n] \otimes [\mathbf{J} + \mathbf{K}_1, n]^* \\ &= (-1)^{|\mathbf{I}| + |\mathbf{J}|} [\mathbf{I}, \mathbf{K}_1 + \{n\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{n\}]^*. \end{aligned}$$

Since $|\mathbf{I}| + |\mathbf{J}| + 2|\mathbf{W}| = \ell$, we have $(-1)^{|\mathbf{I}| + |\mathbf{J}|} = 1$.

So in this case, the equality holds.

Case 2. $n \in \mathbf{I}$.

Then $n \notin \mathbf{W}, \mathbf{J}, \mathbf{K}$ and $(u_n - u_{\bar{n}}) < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > = < \mathbf{J} + \{\mathbf{n}\}, \mathbf{W}, \bar{\mathbf{W}}, \overline{\mathbf{I} - \{\mathbf{n}\}} >$.

The image of ϕ_ℓ is

$$\phi_\ell(< \mathbf{J} + \{\mathbf{n}\}, \mathbf{W}, \bar{\mathbf{W}}, \overline{\mathbf{I} - \{\mathbf{n}\}} >) = \sum_{[\mathbf{I}, \mathbf{n}] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}|}}{2^{(n - |\mathbf{J}| - |\mathbf{I}|)/2}} [\mathbf{I} - \{\mathbf{n}\}, \mathbf{K}] \otimes [\mathbf{J} + \{\mathbf{n}\}, \mathbf{K}]^*.$$

We calculate $(u_n - u_{\bar{n}})\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >)$. Then we have

$$\begin{aligned} & (u_n - u_{\bar{n}})[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* \\ &= \epsilon \begin{pmatrix} \mathbf{I}, & \mathbf{K} \\ \mathbf{I} - \{\mathbf{n}\} + \mathbf{K}, & \{\mathbf{n}\} \end{pmatrix} \epsilon \begin{pmatrix} \mathbf{J}, & \mathbf{K} \\ \mathbf{J} + \mathbf{K} & \end{pmatrix} (-1)^{|\mathbf{I}| - 1 + |\mathbf{K}| + |\mathbf{J}| + |\mathbf{K}| + 1} [\mathbf{I} - \{\mathbf{n}\} + \mathbf{K}] \otimes [\mathbf{J} + \mathbf{K}, \mathbf{n}]^* \\ &= (-1)^{|\mathbf{I}| + |\mathbf{J}|} [\mathbf{I} - \{\mathbf{n}\}, \mathbf{K}] \otimes [\mathbf{J} + \{\mathbf{n}\}, \mathbf{K}]^*. \end{aligned}$$

Here $\mathbf{I} - \{\mathbf{n}\} + \mathbf{K}$ denotes the set obtained by the set theoretical difference and sum and we consider the indices in it are arranged in the increasing order.

So we have the equality.

The proof for the case $n \in \mathbf{J}$ is almost the same and we omit it.

Case 3. $n \in \mathbf{W}$.

Then $n \notin \mathbf{I}, \mathbf{J}, \mathbf{K}$ and $(u_n - u_{\bar{n}}) < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > = - < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >$.

The image of ϕ_ℓ of this element is $-\sum_{[\mathbf{n}] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} ((-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}|} / 2^{(n - |\mathbf{J}| - |\mathbf{I}|)/2}) [\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*.$

We calculate $(u_n - u_{\bar{n}})\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >)$. We take cases for $n \in \mathbf{K}$ and $n \notin \mathbf{K}$.

Case 3.1 $n \in \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{\mathbf{n}\}$ and the image of the sum over such \mathbf{K} 's is given by $\sum_{[\mathbf{n}] - \mathbf{J} - \mathbf{I} - \{\mathbf{n}\} \supseteq \mathbf{K}_1} ((-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}_1| - 1} / 2^{(n - |\mathbf{J}| - |\mathbf{I}|)/2}) [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1]^*.$

Case 3.2 $n \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K}$ and the image of the sum over such \mathbf{K} 's is given by $\sum_{[\mathbf{n}] - \mathbf{J} - \mathbf{I} - \{\mathbf{n}\} \supseteq \mathbf{K}_1} ((-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}_1|} / 2^{(n - |\mathbf{J}| - |\mathbf{I}|)/2}) [\mathbf{I}, \mathbf{K}_1 + \{\mathbf{n}\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*.$

If we note $|\mathbf{W} - \mathbf{W} \cap \mathbf{K}_1| \equiv |\mathbf{W} - \mathbf{W} \cap (\mathbf{K}_1 + \{\mathbf{n}\})| + 1 \pmod{2}$, we have the equality.

Next we prove the case for odd $\ell (\neq n)$.

Case 1. $n \notin \mathbf{I}, \mathbf{J}, \mathbf{W}$.

Then $(u_n - u_{\bar{n}}) < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > = < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >$.

We calculate $(u_n - u_{\bar{n}})\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >)$.

Case 1.1 $n \in \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{\mathbf{n}\}$, the action of $(u_n - u_{\bar{n}})$ is given by

$$(u_n - u_{\bar{n}})[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* = (-1)^{|\mathbf{I}| + |\mathbf{J}|} [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1]^* = -[\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1]^*,$$

Since $|\mathbf{I}| + |\mathbf{J}| + 2|\mathbf{W}| = \ell$, we have $(-1)^{|\mathbf{I}| + |\mathbf{J}|} = -1$.

Case 1.2 $n \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K}$, the action of $(u_n - u_{\bar{n}})$ is given by

$$(u_n - u_{\bar{n}})[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* = -[\mathbf{I}, \mathbf{K}_1 + \{\mathbf{n}\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*.$$

In both cases the signatures are -1 and the parities of the numbers of the elements in K change. Since the signatures in the definition of ϕ_ℓ are given by $(-1)^{|\mathbf{W}-\mathbf{W}\cap\mathbf{K}|+|\mathbf{I}|+|\mathbf{K}|}$ in this case, we have the equality.

Case 2. $n \in \mathbf{I}$.

Then $n \notin \mathbf{W}, \mathbf{J}, \mathbf{K}$ and $(u_n - u_{\bar{n}}) < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > = < \mathbf{J} + \{\mathbf{n}\}, \mathbf{W}, \bar{\mathbf{W}}, \overline{\mathbf{I} - \{\mathbf{n}\}} >.$

The image of ϕ_ℓ is

$$\phi_\ell(< \mathbf{J} + \{\mathbf{n}\}, \mathbf{W}, \bar{\mathbf{W}}, \overline{\mathbf{I} - \{\mathbf{n}\}} >) = \sum_{[n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W}-\mathbf{W}\cap\mathbf{K}|+|\mathbf{I}|+|\mathbf{K}|}}{2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}} [\mathbf{I} - \{\mathbf{n}\}, \mathbf{K}] \otimes [\mathbf{J} + \{\mathbf{n}\}, \mathbf{K}]^*.$$

We calculate $(u_n - u_{\bar{n}})\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >).$ We have

$$(u_n - u_{\bar{n}})[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* = (-1)^{|\mathbf{I}|+|\mathbf{J}|} [\mathbf{I} - \{\mathbf{n}\}, \mathbf{K}] \otimes [\mathbf{J} + \{\mathbf{n}\}, \mathbf{K}]^* = -[\mathbf{I} - \{\mathbf{n}\}, \mathbf{K}] \otimes [\mathbf{J} + \{\mathbf{n}\}, \mathbf{K}]^*.$$

In this case the parity of \mathbf{I} changes and we have the equality.

The proof for the case $n \in \mathbf{J}$ is almost the same and we omit it.

Case 3. $n \in \mathbf{W}$.

Then $n \notin \mathbf{I}, \mathbf{J}, \mathbf{K}$ and $(u_n - u_{\bar{n}}) < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > = - < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >.$

The image of ϕ_ℓ of this element is $-\sum_{[n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} ((-1)^{|\mathbf{W}-\mathbf{W}\cap\mathbf{K}|+|\mathbf{I}|+|\mathbf{K}|} / 2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}) [\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*.$

We calculate $(u_n - u_{\bar{n}})\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >).$ We take cases for $n \in \mathbf{K}$ and $n \notin \mathbf{K}$.

Case 3.1 $n \in \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{\mathbf{n}\}$ and the image of the sum over such \mathbf{K} 's is given by $\sum_{[n] - \mathbf{J} - \mathbf{I} - \{\mathbf{n}\} \supseteq \mathbf{K}_1} ((-1)^{|\mathbf{W}-\mathbf{W}\cap\mathbf{K}_1|+|\mathbf{I}|+|\mathbf{K}_1|+1} / 2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}) [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1]^*.$

Case 3.2 $n \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K}$ and the image of the sum over such \mathbf{K} 's is given by $\sum_{[n] - \mathbf{J} - \mathbf{I} - \{\mathbf{n}\} \supseteq \mathbf{K}_1} ((-1)^{|\mathbf{W}-\mathbf{W}\cap\mathbf{K}_1|+|\mathbf{I}|+|\mathbf{K}_1|+1} / 2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}) [\mathbf{I}, \mathbf{K}_1 + \{\mathbf{n}\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*.$

If we note $|\mathbf{W}-\mathbf{W}\cap\mathbf{K}_1|+|\mathbf{I}|+|\mathbf{K}_1| \equiv |\mathbf{W}-\mathbf{W}\cap(\mathbf{K}_1 + \{\mathbf{n}\})|+|\mathbf{I}|+|(\mathbf{K}_1 + \{\mathbf{n}\})| \pmod{2}$, we have the equality.

In the case of $\ell = n$, only the difference from the other is a scalar multiple $\frac{1}{\sqrt{2}}$ and the above proof goes well in this case.

The verifications for the other cases are almost similar and we omit it.

□

Remark 4.8. As before, if we compare the same weight spaces in both sides of the above, we obtain the inverse as follows.

If $|\mathbf{J}| + |\mathbf{I}| \equiv 0 \pmod{2}$, we have

$$[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{W}} \frac{(-1)^{|\mathbf{W}-\mathbf{W}\cap\mathbf{K}|}}{2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}} < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >$$

and if $|\mathbf{J}| + |\mathbf{I}| \equiv 1 \pmod{2}$, we have

$$[\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^* = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{W}} \frac{(-1)^{|\mathbf{K}-\mathbf{K}\cap\mathbf{W}|+|\mathbf{I}|+|\mathbf{K}|}}{2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}} < \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} > .$$

As for the composition of r_ℓ in Lemma 3.13 and $\phi_\ell, \phi_\ell^{\epsilon_1\epsilon_2}$, we have the following.

Lemma 4.9. *For $\mathfrak{so}(2n+1)$, as a homomorphism from $\bigwedge^\ell V$ to $\Delta \otimes \Delta^*$, it holds that $\phi_{2n+1-\ell} \circ r_\ell = \phi_\ell$.*

For $\mathfrak{so}(2n)$, as a homomorphism from $\bigwedge^\ell V$ to $\Delta^{\epsilon_1} \otimes (\Delta^{\epsilon_2})^$, it holds that*

$$\phi_{2n-\ell}^{\epsilon_1\epsilon_2} \circ r_\ell = \epsilon_2(-1)^n \phi_\ell^{\epsilon_1\epsilon_2}.$$

Proof. The verification is straightforward. So we omit the proof. \square

Remark 4.10. *If we admit Theorem 4.1 and Theorem 4.4, from this lemma, we obtain the result in Lemma 3.13, which says that r_ℓ is an $\mathfrak{so}(N)$ -equivariant isomorphism. For, ϕ_ℓ and $\phi_{N-\ell}$ are equivariant embedding to the same irreducible component in $\Delta \otimes (\Delta)^*$ respectively. This r_ℓ comes from the determinant of $SO(N)$.*

Finally we give the proofs of Theorem 4.1 and 4.4 simultaneously.

Proof. Since $\bigwedge^\ell V$ ($\ell \neq n$ for $\mathfrak{so}(2n)$), e_n^+ and e_n^- are irreducible, in order to prove the theorem, it is enough to show that ϕ_ℓ and $\phi_\ell^{\epsilon_1\epsilon_2}$ are non-zero equivariant homomorphisms.

First we check the images of the highest weight vectors under the map ϕ_ℓ and $\phi_\ell^{\epsilon_1\epsilon_2}$ are non-zero.

For abbreviation (and to avoid some confusion), we use the following notation hereafter.

For $1 \leq k \leq r \leq n$, by $[k, r]$, we denote the set $\{k, k+1, \dots, r\}$ and by $\overline{[r, k]}$, we denote the set $\{\overline{r}, \overline{r-1}, \dots, \overline{k}\}$.

For $\mathfrak{so}(2n+1)$, the image of the highest weight for $\ell \leq n$ is $\phi_\ell(< [1, \ell] >) = \sum_{[\ell+1, n] \supseteq K} 2^{(n-\ell)/2} [K] \otimes [J, K]^* \neq 0$ and for $\ell \geq n$, the image of the highest weight is $\phi_\ell(< [1, n], 0', \overline{[n, \ell+1]} >) = \sum_{[\ell+1, n] \supseteq K} 2^{(n-\ell)/2} [K] \otimes [J, K]^* \neq 0$, since $W = [\ell+1, n]$.

For $\mathfrak{so}(2n)$, the image of the highest weight for $\ell < n$ is

$$\phi_\ell^{\epsilon_1\epsilon_2}(< [1, \ell] >) = \sum_{[\ell+1, n] \supseteq K} 2^{(n-\ell-1)/2} [K] \otimes [[1, \ell], K]^* \neq 0$$

and if $\ell > n$, the image of the highest weight is

$$\phi_\ell^{\epsilon_1\epsilon_2}(< [1, n], \overline{[n, \ell+1]} >) = \sum_{[\ell+1, n] \supseteq K} 2^{(n-\ell-1)/2} (-1)^{n-\ell-|K|} [K] \otimes [[1, \ell], K]^* \neq 0,$$

since $W = [\ell+1, n]$. For $\ell = n$, if we put $\epsilon = (-1)^n$, we have

$$\phi_n^{+\epsilon}(< [1, n] >) = [\emptyset] \otimes [[1, n]]^* \neq 0$$

$$\phi_n^{-\epsilon}(< [1, n-1], \overline{n} >) = [n] \otimes [[1, n-1]]^* \neq 0.$$

For $\mathfrak{so}(2n)$ and $\ell = n$, if we define the linear homomorphism $\phi_n^{even} = \phi_n^{++} \oplus \phi_n^{--}$ for even $\ell = n$ and $\phi_n^{even} = \phi_n^{+-} \oplus \phi_n^{-+}$ for odd $\ell = n$, then ϕ_n^{even} is of the form

$$\phi_n^{even}(< J, W, \bar{W}, \bar{I} >) = \sum_{[1, n] - J - I \supseteq K} \frac{(-1)^{|W - W \cap K|}}{2^{(n - |J| - |I|)/2}} [I, K] \otimes [J, K]^*.$$

If the above ϕ_n^{even} is $\mathfrak{so}(2n)$ -equivariant isomorphism, then $\phi_n^{\epsilon_1 \epsilon_2}$ are $\mathfrak{so}(2n)$ -equivariant isomorphism. We note that the above ϕ_n^{even} is of the same form to (4.1.1).

For $\mathfrak{so}(N)$, from the definitions (3.12) and (3.15), we can check easily $\phi_\ell \iota' = \iota' \phi_\ell$ and $\phi_\ell^{\epsilon_1 \epsilon_2} \iota' = \iota' \phi_\ell^{\epsilon_2 \epsilon_1}$.

Therefore to prove the theorem, it is enough to show $X_i \phi_\ell = \phi_\ell X_i$ and $X_i \phi_\ell^{\epsilon_1 \epsilon_2} = \phi_\ell^{\epsilon_1 \epsilon_2} X_i$ for any $i \in \{1, 2, \dots, n\}$, since $X_i = -\iota' Y_i \iota'$ in both spaces $\bigwedge^\ell V$ and $\Delta \otimes \Delta^*$. Since X_i 's and Y_i 's generate the Lie algebra $\mathfrak{so}(N)$, we have the theorems.

We note that the definitions of ϕ_ℓ , $\phi_\ell^{\epsilon_1 \epsilon_2}$ and ϕ_n^{even} for the elements $< J, W, \bar{W}, \bar{I} >$ are the same up to the scalar and the action of X_i and Y_i for $i = 1, 2, \dots, n - 1$ are also same for the exterior products of V and Δ and Δ^* . So if we show $X_p \phi_\ell(< J, W, \bar{W}, \bar{I} >) = \phi_\ell X_p(< J, W, \bar{W}, \bar{I} >)$ ($1 \leq p \leq n - 1$) for $\mathfrak{so}(2n + 1)$, the proof holds for the rest.

The proof goes by case-by-case verifications.

Proof of $X_p \phi_\ell(< J, W, \bar{W}, \bar{I} >) = \phi_\ell X_p(< J, W, \bar{W}, \bar{I} >)$ for $1 \leq p < n$

Case 1. $p + 1 \in J$ and $p \in W$.

Then $p, p + 1 \notin I$ and $p \notin J$ and $p + 1 \notin W$. We calculate $\phi_\ell X_p$.

Since $X_p < J, W, \bar{W}, \bar{I} > = - < J, W, \bar{W} - \{p\} + \{p + 1\}, \bar{I} > = < J - \{p + 1\} + \{p\}, W - \{p\} + \{p + 1\}, \bar{W} - \{p\} + \{p + 1\}, \bar{I} >$, we have

$$\begin{aligned} & \phi_\ell(X_p < J, W, \bar{W}, \bar{I} >) \\ &= \sum_{[1, n] - (J - \{p + 1\} + \{p\}) - I \supseteq K} \frac{(-1)^{|(W - \{p\} + \{p + 1\}) - K \cap (W - \{p\} + \{p + 1\})|}}{2^{(n - |J| - |I|)/2}} [I, K] \otimes [J - \{p + 1\} + \{p\}, K]^*. \end{aligned}$$

We take two cases for $p + 1 \in K$ and $p + 1 \notin K$. (We note that $p \notin K$.)

Case 1-1. $p + 1 \in K$.

We put $K_1 = K - \{p + 1\}$. Then if we note that the indices in the bracket have alternating property, we have $[I, K] \otimes [J - \{p + 1\} + \{p\}, K]^* = -[I, K_1 + \{p + 1\}] \otimes [J, K_1 + \{p\}]^*$. So the coefficient of $[I, K_1 + \{p + 1\}] \otimes [J, K_1 + \{p\}]^*$ is $(-1)^{|W - K_1 \cap W|} / 2^{(n - |J| - |I|)/2}$.

Case 1-2. $p + 1 \notin K$.

We put $K_1 = K$. The coefficient of $[I, K_1] \otimes [J - \{p + 1\} + \{p\}, K_1]^*$ is $(-1)^{|W - K_1 \cap W|} / 2^{(n - |J| - |I|)/2}$.

Next we calculate $X_p\phi_\ell$.

$$X_p\phi_\ell(< \mathbf{J}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}} >) = \sum_{[1,n]-\mathbf{J}-\mathbf{I} \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W}-\mathbf{W} \cap \mathbf{K}|}}{2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}} X_p([\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*).$$

We take two cases for $p \in \mathbf{K}$ and $p \notin \mathbf{K}$. (We note that $p+1 \notin \mathbf{K}$ since $p+1 \in \mathbf{J}$.)

Case 1-1'. $p \in \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{p\}$. Then $X_p([\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*) = X_p([\mathbf{I}, \mathbf{K}_1 + \{p\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{p\}]^*) = -[\mathbf{I}, \mathbf{K}_1 + \{p+1\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{p\}]^*$. The coefficient of the base element $[\mathbf{I}, \mathbf{K}_1 + \{p+1\}] \otimes [\mathbf{J}, \mathbf{K}_1 + \{p\}]^*$ is $(-1)^{|\mathbf{W}-\mathbf{W} \cap \mathbf{K}_1|} / 2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}$.

Case 1-2'. $p \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K}$. Then $X_p([\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*) = X_p([\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p+1\}, \mathbf{K}_1]^*) = [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p\}, \mathbf{K}_1]^*$. Here we put $\mathbf{J}_1 = \mathbf{J} - \{p+1\}$.

The coefficient of $[\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p\}, \mathbf{K}_1]^*$ is $(-1)^{|\mathbf{W}-\mathbf{W} \cap \mathbf{K}_1|} / 2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}$.

So comparing the corresponding cases, we obtain $X_p\phi_\ell = \phi_\ell X_p$ for this base element.

Case 2. $p+1 \in \mathbf{J}$ and $p \in \mathbf{I}$.

In this case, $p, p+1 \notin \mathbf{W}$ and we put $\mathbf{J}_1 = \mathbf{J} - \{p+1\}$ and $\mathbf{I}_1 = \mathbf{I} - \{p\}$.

Then $X_p < \mathbf{J}_1 + \{p+1\}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}_1 + \{p\}} > = < \mathbf{J}_1 + \{p\}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}_1 + \{p\}} > - < \mathbf{J}_1 + \{p+1\}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}_1 + \{p+1\}} >$ and $< \mathbf{J}_1 + \{p\}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}_1 + \{p\}} > = \epsilon < \mathbf{J}_1, \mathbf{W} + \{p\}, \overline{\mathbf{W} + \{p\}}, \overline{\mathbf{I}_1} >$ and $< \mathbf{J}_1 + \{p+1\}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}_1 + \{p+1\}} > = \epsilon < \mathbf{J}_1, \mathbf{W} + \{p+1\}, \overline{\mathbf{W} + \{p+1\}}, \overline{\mathbf{I}_1} >$.

Here we put $\epsilon = (-1)^{\#\{j \in \mathbf{J}_1: j > p\} + \#\{i \in \mathbf{I}_1: i > p\}} = (-1)^{\#\{j \in \mathbf{J}_1: j > p+1\} + \#\{i \in \mathbf{I}_1: i > p+1\}}$.

Then we have

$$\begin{aligned} \phi_\ell(X_p < \mathbf{J}, \mathbf{W}, \overline{\mathbf{W}}, \overline{\mathbf{I}} >) &= \epsilon \sum_{[1,n]-\mathbf{J}_1-\mathbf{I}_1 \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W}+\{p\}-\mathbf{K} \cap (\mathbf{W}+\{p\})|}}{2^{(n-|\mathbf{J}_1|-|\mathbf{I}_1|)/2}} [\mathbf{I}_1, \mathbf{K}] \otimes [\mathbf{J}_1, \mathbf{K}]^* \\ &\quad - \epsilon \sum_{[1,n]-\mathbf{J}_1-\mathbf{I}_1 \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W}+\{p+1\}-\mathbf{K} \cap (\mathbf{W}+\{p+1\})|}}{2^{(n-|\mathbf{J}_1|-|\mathbf{I}_1|)/2}} [\mathbf{I}_1, \mathbf{K}] \otimes [\mathbf{J}_1, \mathbf{K}]^*. \end{aligned}$$

We have four cases here, but if $p, p+1 \in \mathbf{K}$, or if $p, p+1 \notin \mathbf{K}$, the two terms in the right-hand side are canceled out. So we take two cases for $p \in \mathbf{K}, p+1 \notin \mathbf{K}$ and $p \notin \mathbf{K}, p+1 \in \mathbf{K}$.

Case 2-1. $p \in \mathbf{K}, p+1 \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{p\}$. Then the base element of the first term in the right-hand side is $[\mathbf{I}_1, \mathbf{K}_1 + \{p\}] \otimes [\mathbf{J}_1, \mathbf{K}_1 + \{p\}]^* = \epsilon [\mathbf{I}_1 + \{p\}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p\}, \mathbf{K}_1]^*$ and the coefficient of $[\mathbf{I}_1 + \{p\}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p\}, \mathbf{K}_1]^*$ is $(-1)^{|\mathbf{W}-\mathbf{K}_1 \cap \mathbf{W}|} / 2^{(n-|\mathbf{J}_1|-|\mathbf{I}_1|)/2}$.

Also the base element of the second term in the right-hand side is $[\mathbf{I}_1 + \{p\}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p\}, \mathbf{K}_1]^*$ and its coefficient is $(-1)^{|\mathbf{W}-\mathbf{K}_1 \cap \mathbf{W}|} / 2^{(n-|\mathbf{J}_1|-|\mathbf{I}_1|)/2}$. So the coefficient are the same and the total coefficient of $[\mathbf{I}_1 + \{p\}, \mathbf{K}_1] \otimes [\mathbf{J}_1 + \{p\}, \mathbf{K}_1]^*$ in this case is given by $(-1)^{|\mathbf{W}-\mathbf{K}_1 \cap \mathbf{W}|} / 2^{(n-2-|\mathbf{J}_1|-|\mathbf{I}_1|)/2} = (-1)^{|\mathbf{W}-\mathbf{K}_1 \cap \mathbf{W}|} / 2^{(n-|\mathbf{J}|-|\mathbf{I}|)/2}$.

Case 2-2. $p \notin \mathbf{K}, p+1 \in \mathbf{K}$.

We put $K_1 = K - \{p+1\}$. By a similar calculation as above, the total coefficient of $[I_1 + \{p+1\}, K_1] \otimes [J_1 + \{p+1\}, K_1]^*$ in this case is given by $(-1)^{1+|W-K_1 \cap W|} / 2^{(n-|J|-|I|)/2}$.

Next we calculate $X_p \phi_\ell$. We have

$$X_p \phi_\ell(< J, W, \bar{W}, \bar{I} >) = \sum_{[1,n]-J-I \supseteq K} \frac{(-1)^{|W-W \cap K|}}{2^{(n-|J|-|I|)/2}} \times \\ (-[I_1 + \{p+1\}, K] \otimes [J_1 + \{p+1\}, K]^* + [I_1 + \{p\}, K] \otimes [J_1 + \{p\}, K]^*).$$

Since $p, p+1 \notin K$, we put $K = K_1$ and compare the coefficients in Case 2-1 and Case 2-2, we obtain $X_p \phi_\ell = \phi_\ell X_p$ for this base element $< J, W, \bar{W}, \bar{I} >$.

Case 3. $p+1 \in W$ and $p \notin I, J, W$.

Then $p+1 \notin I$ and $p+1 \notin J$ and we put $W_1 = W - \{p+1\}$.

Since $X_p < J, W, \bar{W}, \bar{I} > = \epsilon < J + \{p\}, W_1, \bar{W}_1, \bar{I} + \{p+1\} >$, we have

$$\phi_\ell(X_p < J, W, \bar{W}, \bar{I} >) \\ = \epsilon \sum_{[1,n]-(J+\{p\})-(I+\{p+1\}) \supseteq K} \frac{(-1)^{|W_1-W_1 \cap K|}}{2^{(n-|J|-|I|-2)/2}} [I + \{p+1\}, K] \otimes [J + \{p\}, K]^* \\ = 2 \sum_{[1,n]-(J+\{p\})-(I+\{p+1\}) \supseteq K} \frac{(-1)^{|W_1-W_1 \cap K|}}{2^{(n-|J|-|I|)/2}} [I, K + \{p+1\}] \otimes [J, K + \{p\}]^*.$$

Here we put $\epsilon = (-1)^{\#\{j \in J: j > p\} + \#\{i \in I: i > p+1\}}$.

We calculate $X_p \phi_\ell$. Since $W = W_1 + \{p+1\}$, we have

$$\phi_\ell(< J, W, \bar{W}, \bar{I} >) = \sum_{[1,n]-J-I \supseteq K} \frac{(-1)^{|(W_1+\{p+1\})-(W_1+\{p+1\}) \cap K|}}{2^{(n-|J|-|I|)/2}} [I, K] \otimes [J, K]^*.$$

If $p, p+1 \in K$ or if $p, p+1 \notin K$, then the action of X_p on the above element reduces to 0. So we have two cases for $p \in K, p+1 \notin K$ and $p \notin K, p+1 \in K$.

Case 3-1. $p \in K, p+1 \notin K$.

We put $K_1 = K - \{p\}$. Then we have

$$X_p \phi_\ell(< J, W, \bar{W}, \bar{I} >) \\ = \sum_{[1,n]-J-I \supseteq K} \frac{(-1)^{|(W_1+\{p+1\})-(W_1+\{p+1\}) \cap (K_1+\{p\})|}}{2^{(n-|J|-|I|)/2}} X_p([I, K_1 + \{p\}] \otimes [J, K_1 + \{p\}]^*) \\ = \sum_{[1,n]-J-I \supseteq K} \frac{(-1)^{|W_1-W_1 \cap K_1|}}{2^{(n-|J|-|I|)/2}} [I, K_1 + \{p+1\}] \otimes [J, K_1 + \{p\}]^*.$$

Case 3-2. $p \notin K, p+1 \in K$.

We put $K_1 = K - \{p+1\}$.

$$X_p \phi_\ell(< J, W, \bar{W}, \bar{I} >) = \sum_{[1,n]-J-I \supseteq K} \frac{(-1)^{|W_1-W_1 \cap K_1|}}{2^{(n-|J|-|I|)/2}} [I, K_1 + \{p+1\}] \otimes [J, K_1 + \{p\}]^*.$$

So by adding the Case 3-1 and Case 3-2, we have $X_p \phi_\ell = \phi_\ell X_p$ for the base element $< J, W, \bar{W}, \bar{I} >$.

For $\mathfrak{so}(2n+1)$, we must show that $X_p \phi_\ell = \phi_\ell X_p$ ($p = 1, 2, \dots, n-1$) for the base element $< J, W, 0', \bar{W}, \bar{I} >$, but the proof is almost similar as above, so we omit it.

Next, we show that $X_n \phi_\ell = \phi_\ell X_n$ for $\mathfrak{so}(2n+1)$.

Proof of $X_n \phi_\ell = \phi_\ell X_n$.

Case 1. $n \in I$.

We put $I_1 = I - \{n\}$. Then $X_n < J, W, \bar{W}, \bar{I} > = -\sqrt{2} < J, W, \bar{W}, 0', \bar{I}_1 >$ and we have

$$\phi_\ell(X_n < J, W, \bar{W}, \bar{I} >) = \sum_{[1,n]-J-I_1 \supseteq K} \frac{(-1)^{|K-K \cap W|+|W|+1}}{2^{(n-|J|-|I|)/2}} [I_1, K] \otimes [J, K]^*$$

We take the cases for $n \in K$ and $n \notin K$.

Case 1-1. $n \in K$.

We put $K_1 = K - \{n\}$. Then in the right-hand side, the base element is $[I_1, K_1, n] \otimes [J, K_1, n]^*$ and its coefficient is $(-1)^{|K_1-K_1 \cap W|+|W|}/2^{(n-|J|-|I|)/2}$.

Case 1-2. $n \notin K$.

We put $K_1 = K$. Then in the right-hand side, the base element is $[I_1, K_1] \otimes [J, K_1]^*$ and its coefficient is $(-1)^{|K_1-K_1 \cap W|+|W|+1}/2^{(n-|J|-|I|)/2}$.

Next, we calculate $X_n \phi_\ell$.

$$\phi_\ell(< J, W, \bar{W}, \bar{I} >) = \sum_{[1,n]-J-I \supseteq K} \frac{(-1)^{|W-W \cap K|}}{2^{(n-|J|-|I|)/2}} [I, K] \otimes [J, K]^*$$

and since $n \in I, n \notin J, K$. If we write $K = K_1$, we have

$$\begin{aligned} X_n \phi_\ell(< J, W, \bar{W}, \bar{I} >) \\ = \sum_{[1,n]-J-I \supseteq K_1} \frac{(-1)^{|W-W \cap K_1|}}{2^{(n-|J|-|I|)/2}} ((-1)^{|K_1|+1} [I_1, K_1] \otimes [J, K_1]^* + [I, K_1] \otimes [J, K_1, n]^*). \end{aligned}$$

Here we note that $X_n [I, K_1] = X_n [I_1, n, K_1] = (-1)^{|K_1|} X_n [I_1, K_1, n] = (-1)^{|K_1|+1} [I_1, K_1]$.

So the coefficient of $[I_1, K_1] \otimes [J, K_1]^*$ is equal to $\frac{(-1)^{|W-W \cap K_1|+|K_1|+1}}{2^{(n-|J|-|I|)/2}}$, which is equal to that of Case 1-2. Since $[I, K_1] \otimes [J, K_1, n]^* = [I_1, n, K_1] \otimes [J, K_1, n]^* = (-1)^{|K_1|} [I_1, K_1, n] \otimes [J, K_1, n]^*$, the coefficient of $[I_1, K_1, n] \otimes$

$[J, K_1, \mathbf{n}]^*$ is given by $\frac{(-1)^{|\mathbf{W}-\mathbf{W}\cap K_1|+|K_1|}}{2^{(n-|J|-|I|)/2}}$, which is equal to that of Case 1-1. So we have done the verification in this case.

Next we check the equation for the base element $\langle J, \mathbf{W}, O', \bar{\mathbf{W}}, \bar{I} \rangle$. (We are still in the Case 1 and assume $I \ni n$.)

Since $X_n \langle J, \mathbf{W}, O', \bar{\mathbf{W}}, \bar{I} \rangle = \sqrt{2} \langle J, \mathbf{W}, \mathbf{n}, \bar{\mathbf{W}}, \bar{I} \rangle = \sqrt{2}(-1)^{|\mathbf{W}|} \langle J, \mathbf{W} + \{\mathbf{n}\}, \bar{\mathbf{W}} + \{\mathbf{n}\}, \bar{I}_1 \rangle$, we have

$$\phi_\ell(X_n \langle J, \mathbf{W}, O', \bar{\mathbf{W}}, \bar{I} \rangle) = \sqrt{2}(-1)^{|\mathbf{W}|} \sum_{[1, n] - J - I_1 \supseteq K} \frac{(-1)^{|\mathbf{W} + \{\mathbf{n}\} - (\mathbf{W} + \{\mathbf{n}\}) \cap K|}}{2^{(n-|J|-|I|+1)/2}} [I_1, K] \otimes [J, K]^*.$$

We take the cases for $n \in K$ and $n \notin K$.

Case 1-3. $n \in K$.

We put $K_1 = K - \{n\}$. Then in the right-hand side, the base element is $[I_1, K_1, \mathbf{n}] \otimes [J, K_1, \mathbf{n}]^*$ and its coefficient is $(-1)^{|\mathbf{W}-K_1 \cap \mathbf{W}|+|\mathbf{W}|} / 2^{(n-|J|-|I|)/2}$.

Case 1-4. $n \notin K$.

We put $K_1 = K$. Then in the right-hand side, the base element is $[I_1, K_1] \otimes [J, K_1]^*$ and its coefficient is $(-1)^{|\mathbf{W}-K_1 \cap \mathbf{W}|+|\mathbf{W}|+1} / 2^{(n-|J|-|I|)/2}$.

Next we calculate $X_n \phi_\ell$. Since $n \in I$, $n \notin J, K$, we write $K = K_1$ and we have

$$\begin{aligned} X_n \phi_\ell(\langle J, \mathbf{W}, O', \bar{\mathbf{W}}, \bar{I} \rangle) &= X_n \sum_{[1, n] - J - I \supseteq K} \frac{(-1)^{|K_1 - K_1 \cap \mathbf{W}|}}{2^{(n-|J|-|I|)/2}} [I, K_1] \otimes [J, K_1]^* \\ &= \sum_{[1, n] - J - I \supseteq K_1} \frac{(-1)^{|K_1 - K_1 \cap \mathbf{W}|}}{2^{(n-|J|-|I|)/2}} ((-1)^{|K_1|+1} [I_1, K_1] \otimes [J, K_1]^* + [I, K_1] \otimes [J, K_1, \mathbf{n}]^*). \end{aligned}$$

So the coefficient of $[I_1, K_1] \otimes [J, K_1]^*$ is equal to $(-1)^{|K_1 - \mathbf{W} \cap K_1| + |K_1| + 1} / 2^{(n-|J|-|I|)/2}$, which is equal to that of Case 1-4.

As before, $[I, K_1] \otimes [J, K_1, \mathbf{n}]^* = [I_1, \mathbf{n}, K_1] \otimes [J, K_1, \mathbf{n}]^* = (-1)^{|K_1|} [I_1, K_1, \mathbf{n}] \otimes [J, K_1, \mathbf{n}]^*$, the coefficient of $[I_1, K_1, \mathbf{n}] \otimes [J, K_1, \mathbf{n}]^*$ is given by $(-1)^{|K_1 - \mathbf{W} \cap K_1| + |K_1|} / 2^{(n-|J|-|I|)/2}$, which is equal to that of Case 1-3.

Case 2. $n \in \mathbf{W}$.

Then $n \notin I$ and $n \notin J$ and we put $\mathbf{W}_1 = \mathbf{W} - \{n\}$.

We calculate $\phi_\ell X_n$. Since $X_n \langle J, \mathbf{W}_1 + \{\mathbf{n}\}, \{\mathbf{n}\} + \bar{\mathbf{W}}, \bar{I} \rangle = -\sqrt{2} \langle J, \mathbf{W}_1 + \{\mathbf{n}\}, O', \bar{\mathbf{W}}_1, \bar{I} \rangle = (-1)^{|\mathbf{W}|} \sqrt{2} \langle J + \{\mathbf{n}\}, \mathbf{W}_1, O', \bar{\mathbf{W}}_1, \bar{I} \rangle$, we have

$$\phi_\ell(X_n \langle J, \mathbf{W}, \bar{\mathbf{W}}, \bar{I} \rangle) = 2 \sum_{[1, n] - J_1 - I \supseteq K} \frac{(-1)^{|K - K \cap \mathbf{W}_1| + |\mathbf{W}|}}{2^{(n-|J|-|I|)/2}} [I, K] \otimes [J_1, K]^*.$$

Here we put $J_1 = J + \{n\}$. Since $n \in J_1$, $n \notin K$ and we put $K_1 = K$.

Then in the right-hand side, the base element is $[\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}_1, \mathbf{K}_1]^* = [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J} + \{\mathbf{n}\}, \mathbf{K}_1]^* = (-1)^{|\mathbf{K}_1|} [\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*$ and the coefficient of $[\mathbf{I}, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*$ is $(-1)^{|\mathbf{K}_1 - \mathbf{K}_1 \cap \mathbf{W}_1| + |\mathbf{W}| + |\mathbf{K}_1|} / 2^{(n-2-|\mathbf{J}| - |\mathbf{I}|)/2} = (-1)^{|\mathbf{W} - \mathbf{K}_1 \cap \mathbf{W}|} / 2^{(n-2-|\mathbf{J}| - |\mathbf{I}|)/2}$.

We calculate $X_n \phi_\ell$. Since

$$\phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >) = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}|}}{2^{(n-|\mathbf{J}| - |\mathbf{I}|)/2}} [\mathbf{I}, \mathbf{K}] \otimes [\mathbf{J}, \mathbf{K}]^*$$

and $n \in \mathbf{I}$, $n \notin \mathbf{J}$, we take two cases for $n \in \mathbf{K}$ and $n \notin \mathbf{K}$.

Case 2-1. $n \in \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K} - \{n\}$ and we have

$$X_n \phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >) = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}} \frac{(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}|}}{2^{(n-|\mathbf{J}| - |\mathbf{I}|)/2}} (-[\mathbf{I}_1, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*).$$

So the coefficient of the base element $[\mathbf{I}_1, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*$ is $-(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}|} / 2^{(n-|\mathbf{J}| - |\mathbf{I}|)/2} = (-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}_1|} / 2^{(n-|\mathbf{J}| - |\mathbf{I}|)/2}$.

Case 2-2. $n \notin \mathbf{K}$.

We put $\mathbf{K}_1 = \mathbf{K}$ and we have

$$X_n \phi_\ell(< \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} >) = \sum_{[1, n] - \mathbf{J} - \mathbf{I} \supseteq \mathbf{K}_1} \frac{(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}_1|}}{2^{(n-|\mathbf{J}| - |\mathbf{I}|)/2}} ([\mathbf{I}_1, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*).$$

So the coefficient of the base element $[\mathbf{I}_1, \mathbf{K}_1] \otimes [\mathbf{J}, \mathbf{K}_1 + \{\mathbf{n}\}]^*$ is $(-1)^{|\mathbf{W} - \mathbf{W} \cap \mathbf{K}_1|} / 2^{(n-|\mathbf{J}| - |\mathbf{I}|)/2}$.

By adding the results of Case 2-1 and Case 2-2, we have $\phi_\ell X_n = X_n \phi_\ell$ in this case.

There are several remaining cases, in which we must show $\phi_\ell X_n = X_n \phi_\ell$, but the proof is almost similar, so we omit it.

We must show that $\phi_\ell^{\epsilon_1 \epsilon_2} X_n = X_n \phi_\ell^{\epsilon_1 \epsilon_2}$ and $\phi_n^{\text{even}} X_n = X_n \phi_n^{\text{even}}$, but the proofs go in a similar way by (lengthy) case-by-case verifications and so we omit it.

□

5. INVARIANT THEORY AND THE CENTRALIZER ALGEBRAS

In this section we define a set of elements based on the invariant theory in the centralizer algebra $\mathbf{CP}_\mathbf{k} = \text{Hom}_{\text{Pin}(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^k V)$. They are parameterized by the generalized Brauer diagrams. In the classical situation for the orthogonal group $O(N)$, R. Brauer ([3]) considered the centralizer algebra $\text{Hom}_{O(N)}(\otimes^k V, \otimes^k V)$ and he determined the special basis (parameterized by the Brauer diagrams) and the multiplication rules of the basis elements by using the first and the second fundamental theorem of the invariant theory of $O(N)$.

Since we have already the explicit isomorphisms between the exterior algebras $\bigwedge^i V$ and the irreducible components of $\Delta \otimes \Delta^*$ as $\text{Pin}(N)$ ($O(N)$) modules in the previous section, we can follow the above way.

More generally, we define the spaces of the equivariant homomorphisms.

Definition 5.1.

$$\mathbf{CP}_1^k = \text{Hom}_{Pin(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V).$$

and

$$\mathbf{CS}_1^k = \text{Hom}_{Spin(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V).$$

We write \mathbf{CP}_k and \mathbf{CS}_k for \mathbf{CP}_1^k and \mathbf{CS}_1^k respectively.

Remark 5.2. We note that for $Pin(2n+1)$, Δ stands for Δ_\pm , so there are four cases for \mathbf{CP}_1^k . (But two of them are trivial. See Lemma 5.4.) For $Spin(2n)$, we can change Δ into Δ^\pm in the above and so there are also four cases for \mathbf{CS}_1^k .

We note that from the formulas (2.2.3), (2.2.4), for $\epsilon_1, \epsilon_2 \in \{\pm 1\}$, the irreducible representation $(1/2 + \delta)_{Spin(2n)}^{\epsilon_1}$ of $Spin(2n)$ occurs in $\Delta^{\epsilon_2} \otimes \otimes^k V$ if and only if $(-1)^{k-|\delta|} = \epsilon_1 \epsilon_2$.

So we have $\text{Hom}_{Spin(2n)}(\Delta^{\epsilon_1} \otimes \otimes^k V, \Delta^{\epsilon_2} \otimes \otimes^l V) = 0$ if $\epsilon_1 \epsilon_2 \neq (-1)^{(k-l)}$.

We interpret them as the elements of the invariant subspace of the tensor spaces under the action of $Pin(N)$.

From Theorem 4.3 and Theorem 4.6 and $V \cong V^*$, we have the following lemma.

Lemma 5.3. For $Pin(2n)$, we have

$$(5.3.1) \quad \text{Hom}_{Pin(2n)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V) \cong \bigoplus_{i=0}^{2n} ((\bigwedge^i V) \otimes \otimes^{k+l} V)^{O(2n)}.$$

For $Pin(2n+1)$, we have

$$(5.3.2) \quad \text{Hom}_{Pin(2n+1)}(\Delta_\pm \otimes \otimes^k V, \Delta_\pm \otimes \otimes^l V) \cong \bigoplus_{i=0}^n ((\bigwedge^{2i} V) \otimes \otimes^{k+l} V)^{O(2n+1)}.$$

and

$$(5.3.3) \quad \text{Hom}_{Pin(2n+1)}(\Delta_\pm \otimes \otimes^k V, \Delta_\mp \otimes \otimes^l V) \cong \bigoplus_{i=0}^n ((\bigwedge^{2i+1} V) \otimes \otimes^{k+l} V)^{O(2n+1)}.$$

Here the superscript $O(N)$ denotes the subspace obtained by taking the $O(N)$ -invariants.

First let us recall the first main theorem and the second main theorem for the polynomial invariants under the action of the orthogonal groups.

Let $V = K^N$ be a vector space over a field K of characteristic 0 and let $P(\bigoplus^k V)$ be the polynomial ring over the vector space $\bigoplus^k V$. Let $(\ , \)$ be a non-degenerate symmetric bilinear form which defines

the orthogonal group $O(N)$. Then ‘The First Main Theorem’ (see [21] p53 Theorem 2.9A, p64 Theorem 2.11A.) tells us that the polynomial invariants $P(\bigoplus^k V)^{O(N)}$ are generated by the scalar products $(\mathbf{v}_i, \mathbf{v}_j)$ ($1 \leq i, j \leq k$), where \mathbf{v}_i denotes the vector in the i -th direct summand in $\bigoplus^k V$. ‘The Second Main Theorem’ (see [21] p75 Theorem 2.17A.) tells us that the relations of the polynomial invariants $(\mathbf{v}_i, \mathbf{v}_j)$ ’s in $P(\bigoplus^k V)^{O(N)}$ are generated by the determinants $\det((\mathbf{v}_{i_s}, \mathbf{v}_{j_t}))$ ($0 \leq s, t \leq N$) of size $N + 1$. We note that the minimum (total) degree of the relations is $2N + 2$.

We consider the invariant tensors $(\bigwedge^r V^* \otimes^s V^*)^{O(N)}$. This space is included in the tensor space $(\bigotimes^{r+s} V^*)^{O(N)}$ and we regard this space as the subspace of $P(\bigoplus^{r+s} V)^{O(N)}$. The elements in this space are multi-linear (i.e., degree 1) in each of the tensor components and alternating concerning the first r components. Since the invariants are generated by the scalar products of degree 2, the multi-linear property tells us that $(\bigotimes^{r+s} V^*)^{O(N)} = 0$ if $r + s \equiv 1 \pmod{2}$.

So we have the following.

Lemma 5.4. *If $(-1)^{k+l}\epsilon_1\epsilon_2 = -1$, then*

$$\mathbf{CP}_1^k = \text{Hom}_{\text{Pin}(2n+1)}(\Delta_{\epsilon_1} \otimes^k V, \Delta_{\epsilon_2} \otimes^l V) = 0.$$

Remark 5.5. *From the above lemma, we only consider the case $(-1)^{k+l}\epsilon_1\epsilon_2 = 1$ for the centralizer algebras of $\text{Pin}(2n + 1)$. If we write simply Δ in the formulas for $\text{Pin}(2n + 1)$, we always assume this condition without mentioning it.*

We can write down the elements in $(\bigwedge^r V^* \otimes^s V^*)^{O(N)}$ as follows. If $s < r$, $\bigotimes^s V$ does not contain the representations $\bigwedge^r V$, so this space must be 0.

Let $s \geq r$. For the subsets $\mathbf{t} = \{t_1, t_2, \dots, t_r\}$ ($t_1 < t_2 < \dots < t_r$ and $r \leq N$) of $[1, s]$ and the sets of $u = \frac{s-r}{2}$ pairs of indices $\{\mathbf{m}, \mathbf{l}\} = \{\{m_1, l_1\}, \{m_2, l_2\}, \dots, \{m_u, l_u\}\}$ such that $[1, s] = \mathbf{t} \sqcup \mathbf{m} \sqcup \mathbf{l}$ (disjoint unions), where we put $\mathbf{m} = \{m_1, \dots, m_u\}$ and $\mathbf{l} = \{l_1, \dots, l_u\}$), we define the invariant elements $\mathbf{T}_{\mathbf{t}, \{\mathbf{m}, \mathbf{l}\}}$ by

$$\mathbf{T}_{\mathbf{t}, \{\mathbf{m}, \mathbf{l}\}} = \frac{1}{r!} \det((\mathbf{x}_i, \mathbf{y}_{t_j})) \times \prod_{j=1}^u (\mathbf{y}_{m_j}, \mathbf{y}_{l_j}).$$

Here \mathbf{x}_j ($j = 1, 2, \dots, r$) are the components of the first r tensors in $\bigotimes^{r+s} V$ and \mathbf{y}_j ($j = 1, 2, \dots, s$) are the components of the latter s tensors in $\bigotimes^{r+s} V$.

Moreover $\frac{1}{r!} \det((\mathbf{x}_i, \mathbf{y}_{t_j}))$ is the invariant element in $((\bigwedge^r V)^* \otimes (\bigwedge^r V)^*)^{O(N)}$.

Since $\bigwedge^r V$ is an irreducible $O(N)$ module, the trivial representation occurs in $(\bigwedge^r V)^* \otimes \bigwedge^r V$ with multiplicity one. The invariant element is given by

$$(5.1) \quad \mathfrak{I}_r = \sum_{\substack{[1,n] \supseteq J \sqcup I \sqcup \bar{W} \text{ (disjoint sets)} \\ |J|+|I|+2|\bar{W}|=r}} \langle J, \bar{W}, \bar{I} \rangle^* \otimes \langle J, \bar{W}, \bar{I} \rangle \\ (+ \sum_{\substack{[1,n] \supseteq J \sqcup I \sqcup \bar{W} \text{ (disjoint sets)} \\ |J|+|I|+2|\bar{W}|=r-1}} \langle J, \bar{W}, 0', \bar{W}, \bar{I} \rangle^* \otimes \langle J, \bar{W}, 0', \bar{W}, \bar{I} \rangle).$$

Here the second term in the parentheses are considered only for $O(2n+1)$.

So $\frac{1}{r!} \det((\mathbf{x}_i, \mathbf{y}_{t_j}))$ must be a scalar multiple of $(\text{id} \otimes \iota)(\mathfrak{I}_r)$ (see Definition 3.11.) and the scalar is given by $(-1)^{\binom{r}{2}}$. (To obtain the scalar, take cases of the parity of r and compare the values of both sides at the special elements $\mathbf{x}_1 = u_1, \mathbf{x}_2 = u_2$, etc..)

Lemma 5.6. *For $\text{Pin}(2n)$, if we take only \mathbf{t} 's such that $|\mathbf{t}| \equiv k+l \pmod{2}$ and $|\mathbf{t}| \leq 2n$, then the above $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$'s span linearly the whole space of the $\mathbf{CP}_1^k = \text{Hom}_{\text{Pin}(2n)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V)$.*

Moreover if $k+l < 2n+2$, $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ ($\mathbf{t} \sqcup \mathbf{m} \sqcup \mathbf{l} = [1, s]$) are linearly independent.

For $\text{Pin}(2n+1)$, if we take only \mathbf{t} 's such that $|\mathbf{t}| \leq 2n+1$ and $|\mathbf{t}| \equiv k+l \equiv 0 \pmod{2}$, then the above $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$'s span linearly the whole space of the $\mathbf{CP}_1^k = \text{Hom}_{\text{Pin}(2n+1)}(\Delta_{\pm} \otimes \otimes^k V, \Delta_{\pm} \otimes \otimes^l V)$.

For $\text{Pin}(2n+1)$, if we take only \mathbf{t} 's such that $|\mathbf{t}| \leq 2n+1$ and $|\mathbf{t}| \equiv k+l \equiv 1 \pmod{2}$, then the above $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$'s span linearly the whole space of the $\mathbf{CP}_1^k = \text{Hom}_{\text{Pin}(2n+1)}(\Delta_{\pm} \otimes \otimes^k V, \Delta_{\mp} \otimes \otimes^l V)$.

In both cases, if $k+l < 2n+3$, $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ ($\mathbf{t} \sqcup \mathbf{m} \sqcup \mathbf{l} = [1, s]$) are linearly independent.

Proof. The statements follow from the argument developed above and the Second Main Theorem. \square

From Lemma 5.3, we consider this invariant element $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ as a homomorphism from $\Delta \otimes \otimes^k V$ to $\Delta \otimes \otimes^l V$.

By the original definition, $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ are the elements in $(\bigwedge^r V^*) \otimes \otimes^k V^* \otimes \otimes^l V^*$ for $r \equiv k+l \pmod{2}$. We denote the degree 2 tensor corresponding to the invariant symmetric bilinear form defined by S on $V \times V$ by id_V .

Then as the homomorphism in $\text{Hom}_{\text{Pin}(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V)$, $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ is given as follows.

The pair $(\mathbf{y}_{m_j}, \mathbf{y}_{l_j})$ with $m_j, l_j \leq k$ corresponds to the contraction of the specified tensor components (i.e., m_j -th and l_j -th components) in $\Delta \otimes \otimes^k V$. We denote this contraction by $C_{\{m_j, l_j\}}$.

The pair $(\mathbf{y}_{m_j}, \mathbf{y}_{l_j})$ with $m_j, l_j > k$ corresponds to the insertion of the invariant tensor (id_V) into the specified tensor components (i.e., $m_j - k$ -th and $l_j - k$ -th components) in $\Delta \otimes \otimes^l V$. We denote this insertion by $\text{id}_{V_{m_j-k, l_j-k}}$. (We introduce the numbering such that the first k tensors correspond to the domain and the last l tensors correspond to the range in $\text{Hom}_{\text{Pin}(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V)$.)

The pair $(\mathbf{y}_{m_j}, \mathbf{y}_{l_j})$ with $m_j \leq k$ and $l_j > k$ (resp. with $l_j \leq k$ and $m_j > k$) corresponds to the map which transfers the m_j -th (resp. l_j -th) component of the domain tensor $\Delta \otimes \otimes^k V$ into $l_j - k$ -th (resp. $m_j - k$ -th) component of the image tensor $\Delta \otimes \otimes^l V$. Namely this corresponds to the partial permutation of the tensor components.

The above three operators are the same in the usual Brauer centralizer algebras. For the remaining part $\det((\mathbf{x}_i, \mathbf{y}_{t_j}))$ ($1 \leq i, j \leq r$) in $\mathbf{T}_{\mathbf{t}, \{\mathbf{m}, \mathbf{l}\}}$, we will give the explicit description as the homomorphism. (See Theorem 7.1.)

Let us use a diagrammatic parametrization of the elements $\{\mathbf{T}_{\mathbf{t}, \{\mathbf{m}, \mathbf{l}\}}\}$ of $\text{Hom}_{\text{Pin}(N)}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^l V)$ as follows. We consider the diagrams of two lines vertices, with k vertices in the upper row and l vertices in the lower row, in which vertices are connected with each other as in the usual Brauer diagrams except for admitting isolated vertices. Namely they are graphs with no loops, in which the number of edges connected to each vertex is either 0 or 1.

We call such diagrams the generalized Brauer diagrams and denote the set of the generalized Brauer diagrams by \mathbf{GB}_1^k . (Later we use another parametrization using the same diagrams \mathbf{GB}_1^k from the representation theoretical viewpoint.)

The explicit bijection is given as follows. Let us number the vertices from 1 to k from left to right in the upper row and from 1 to l in the lower row.




FIGURE 2. An example of the generalized Brauer diagrams with $k = 6$ and $l = 5$

We correspond each of $\mathbf{T}_{\mathbf{t}, \{\mathbf{m}, \mathbf{l}\}}$ to a generalized Brauer diagram as follows. We consider the tensor components in the range $\Delta \otimes \otimes^l V$ are indexed by $k+1, k+2, \dots, k+l$ in this argument. For the vertices

with indices in \mathbf{t} , we leave them the isolated vertices and for the other vertices (namely the vertices with indices in \mathbf{l} and \mathbf{m}), we connect l_j -th vertex and m_j -th vertex. Except for the isolated vertices, the edges in the diagrams convey the same meaning as in the usual Brauer diagrams.

So we obtain the bijective correspondence between the elements $\{\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}\}$ of \mathbf{CP}_1^k and the diagrams with the isolated vertices of at most N in \mathbf{GB}_1^k .

From the formula (2.4.2) in Theorem 2.4 and (2.2.3) and (2.2.4) in Theorem 2.2, the multiplicity of $[\Delta, \lambda]_{Pin(N)}$ in $\Delta \otimes \otimes^k V$ is equal to the number of the ‘up-down-stable (n -) tableaux’ of degree k starting with the empty set \emptyset and ending with the Young diagram λ and consisting of the diagrams with their lengths at most n . We show an example of up-down-stable tableaux.




FIGURE 3. An example of ‘up-down-stable tableaux’ of $\lambda = (2, 2)$ and $k = 8$

Namely by definition, the ‘up-down-stable (n -)tableaux’ of degree k are the $k + 1$ sequences of Young diagrams, in which the adjacent diagrams are obtained by attaching a cell to or removing a cell from the previous one or are the same (if its length is less than n for $N = 2n$).

So figure 3 is not an ‘up-down-stable 2-tableau’ for $Pin(4)$ because there exist adjacent diagrams which are the same and its length is 2.

Since \mathbf{CP}_k and $Pin(N)$ are the dual pair on the space $\Delta \otimes \otimes^k V$, the irreducible representations of \mathbf{CP}_k are in bijective correspondence with the irreducible representations of $Pin(N)$ on this space.

So we can expect the base of the irreducible representation λ of \mathbf{CP}_k (corresponding to $[\Delta, \lambda]_{Pin(N)}$) are parameterized by the ‘up-down-stable (n -) tableaux’ of degree k , starting with \emptyset and ending with λ and consisting of the diagrams with their lengths at most n .

From the above, the number of the ‘up-down-stable (n -) tableaux’ of degree $2k + 1$, starting and ending with \emptyset and consisting of the diagrams with their lengths at most n , is equal to the dimension \mathbf{CP}_k . (We note that $\dim \mathbf{CP}_k$ also depends on n if $n \leq k$.)

It seems to be interesting problem to determine the explicit action of \mathbf{CP}_k on the ‘up-down-stable n -tableaux’ base.

We can identify the elements in \mathbf{GB}_k^k with the involutions in the symmetric group of degree $2k$. So we can expect a bijective correspondence between the ‘up-down-stable tableaux’ of degree $2k + 1$ defined

above and the involutions in \mathfrak{S}_{2k} . The author heard from I. Terada that Tom Roby privately communicated to him that the method used in his Thesis [19] also gives such a bijection. Namely the pictorial rules of the Robinson-Schensted correspondence initiated by Sergey Fomin [7] give a bijective correspondence between the ‘up-down-stable tableaux’ of degree k and the involutions in \mathfrak{S}_k .

Finally we give the dimension of \mathbf{CP}_k .

Lemma 5.7. *If $k \leq n$, we have*

$$\dim \mathbf{CP}_k = (2k-1)!! + \binom{2k}{2}(2k-3)!! + \binom{2k}{4}(2k-5)!! + \dots + \binom{2k}{2k-2}(1)!! + \binom{2k}{2k},$$

where $(2k-1)!! = \prod_{i=0}^{k-1} (2k-1-2i)$.

6. REPRESENTATION THEORETIC PARAMETRIZATION

In this section we first give explicit formulas for the projection from $\Delta \otimes \bigwedge^k V$ to Δ and the injection from Δ to $\Delta \otimes \bigwedge^k V$ to Δ .

Then we give another kind of parametrization of the elements in \mathbf{CP}_1^k by the generalized Brauer diagrams, using the above operators.

From Theorem 2.4 and Theorem 2.2, we have $\Delta \otimes \bigwedge^k V = \sum_{i=0}^k [\Delta, (1^i)]_{Spin(N)}$

for any $k \leq n$.

For $Pin(2n)$, the representations $[\Delta, (1^i)]_{Pin(2n)}$ are determined by their characters and the above decomposition literally holds for $Pin(2n)$. But for $Pin(2n+1)$, if we consider the action of the center z , we have the following. For $\epsilon_1 \in \{\pm 1\}$, $\Delta_{\epsilon_1} \otimes \bigwedge^k V = \sum_{i=0}^k [\Delta, (1^i)]_{Pin(2n+1), \epsilon_2}$, where $\epsilon_2 = (-1)^k \epsilon_1$.

We note that the above decomposition is multiplicity free. So the $Pin(N)$ -equivariant projection from $\Delta \otimes \bigwedge^k V$ to Δ and the $Pin(N)$ -equivariant embedding from Δ to $\Delta \otimes \bigwedge^k V$ are determined uniquely up to scalars. We write down those maps explicitly.

The $Pin(N)$ -equivariant projection in $\text{Hom}_{Pin(N)}(\Delta \otimes \bigwedge^k V, \Delta)$ ($\cong (\Delta^* \otimes \Delta \otimes (\bigwedge^k V)^*)^{Pin(N)}$) is given by the image of the following element under the map $\phi_k \otimes \text{id}$. (See Theorem 4.3 and 4.6.)

$$\begin{aligned} & \sum_{\substack{[1,n] \supseteq J \sqcup I \sqcup W \text{ (disjoint sets)} \\ |J|+|I|+2|W|=k}} \langle J, W, \bar{W}, \bar{I} \rangle \otimes \langle J, W, \bar{W}, \bar{I} \rangle^* \\ & (+ \sum_{\substack{[1,n] \supseteq J \sqcup I \sqcup W \text{ (disjoint sets)} \\ |J|+|I|+2|W|=k-1}} \langle J, W, O', \bar{W}, \bar{I} \rangle \otimes \langle J, W, O', \bar{W}, \bar{I} \rangle^*). \end{aligned}$$

For convenience sake, we multiply $2^{n/2}$ to the above and denote the corresponding homomorphism by pr_k . We note that for $\text{Pin}(2n+1)$ and odd k , pr_k is the projection from $\Delta_{\pm} \otimes \bigwedge^k V$ to Δ_{\mp} .

Definition 6.1. For $\text{Pin}(2n+1)$, the projection pr_k is given by

$$\text{pr}_k([T] \otimes \langle I, W, \bar{W}, \bar{J} \rangle) = \begin{cases} 0 & \text{if } I \not\subseteq T, \\ \epsilon \begin{pmatrix} T & \\ I & K \end{pmatrix} (-1)^{|W-W \cap K|} 2^{(|I|+|J|)/2} [J, K] & \text{if } I \subseteq T, \end{cases}$$

and

$$\text{pr}_k([T] \otimes \langle I, W, O', \bar{W}, \bar{J} \rangle) = \begin{cases} 0 & \text{if } I \not\subseteq T, \\ \epsilon \begin{pmatrix} T & \\ I & K \end{pmatrix} (-1)^{|K-W \cap K|} 2^{(|I|+|J|)/2} [J, K] & \text{if } I \subseteq T. \end{cases}$$

Here we put $K = T - I$ and $\epsilon \begin{pmatrix} T & \\ I & K \end{pmatrix}$ denotes the sign of the permutation obtained by arranging T into I, K in this order.

For $\text{Pin}(2n)$, the projection pr_k is given as follows. If k is even, we define

$$\text{pr}_k([T] \otimes \langle I, W, \bar{W}, \bar{J} \rangle) = \begin{cases} 0 & \text{if } I \not\subseteq T, \\ \epsilon \begin{pmatrix} T & \\ I & K \end{pmatrix} (-1)^{|W-W \cap K|} 2^{(|I|+|J|)/2} [J, K] & \text{if } I \subseteq T. \end{cases}$$

If k is odd, we define

$$\text{pr}_k([T] \otimes \langle I, W, \bar{W}, \bar{J} \rangle) = \begin{cases} 0 & \text{if } I \not\subseteq T, \\ \epsilon \begin{pmatrix} T & \\ I & K \end{pmatrix} (-1)^{|W-W \cap K|+|K|+|J|} 2^{(|I|+|J|)/2} [J, K] & \text{if } I \subseteq T. \end{cases}$$

Similarly we have $\dim(\text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)}, \Delta_{(\epsilon_2)} \otimes \bigwedge^k V)) = 1$. (If $N = 2n$, ignore the subscripts (ϵ_1) and (ϵ_2) of Δ . The subscripts (ϵ_1) and (ϵ_2) of Δ are only considered for the case $N = 2n+1$. From now on, we assume this convention. Here we also assume that $\epsilon_2 = (-1)^k \epsilon_1$ is satisfied for $N = 2n+1$.)

$$\text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)}, \Delta_{(\epsilon_2)} \otimes \bigwedge^k V) \cong (\Delta_{(\epsilon_1)}^* \otimes \Delta_{(\epsilon_2)} \otimes \bigwedge^k V)^{\text{Pin}(N)}$$

and the corresponding invariant element is given by \mathfrak{I}_k (ref. (5.1)). So as in the case of pr_k , we obtain the corresponding homomorphism inj_k (after a scalar multiplication by $2^{n/2}$) as follows. We note that for $\text{Pin}(2n+1)$ and odd k , inj_k is the immersion of Δ_{\pm} into $\Delta_{\mp} \otimes \bigwedge^k V$.

Definition 6.2. For $Pin(2n+1)$, the immersion inj_k is given by

$$\begin{aligned} \text{inj}_k([T]) = & \sum_{\substack{T \supseteq I \\ K = T - I}} \epsilon \begin{pmatrix} T \\ I, K \end{pmatrix} \left(\sum_{\substack{J \subseteq ([1,n] - T) \\ W \subseteq ([1,n] - I - J) \\ |J| + |I| + 2|W| = k}} (-1)^{|W - W \cap K|} 2^{(|J| + |I|)/2} [J, K] \otimes k! \langle J, W, \bar{W}, \bar{I} \rangle \right. \\ & + \sum_{\substack{J \subseteq ([1,n] - T) \\ W \subseteq ([1,n] - I - J) \\ |J| + |I| + 2|W| + 1 = k}} (-1)^{|K - W \cap K|} 2^{(|J| + |I|)/2} [J, K] \otimes k! \langle J, W, O', \bar{W}, \bar{I} \rangle \Big). \end{aligned}$$

For $Pin(2n)$, the immersion inj_k is given as follows. For even k , we define

$$\text{inj}_k([T]) = \sum_{\substack{T \supseteq I \\ K = T - I}} \epsilon \begin{pmatrix} T \\ I, K \end{pmatrix} \left(\sum_{\substack{J \subseteq ([1,n] - T) \\ W \subseteq ([1,n] - I - J) \\ |J| + |I| + 2|W| = k}} (-1)^{|W - W \cap K|} 2^{(|J| + |I|)/2} [J, K] \otimes k! \langle J, W, \bar{W}, \bar{I} \rangle \right)$$

For odd k , we define

$$\text{inj}_k([T]) = \sum_{\substack{T \supseteq I \\ K = T - I}} \epsilon \begin{pmatrix} T \\ I, K \end{pmatrix} \left(\sum_{\substack{J \subseteq ([1,n] - T) \\ W \subseteq ([1,n] - I - J) \\ |J| + |I| + 2|W| = k}} (-1)^{|W - W \cap K| + |J| + |K|} 2^{(|J| + |I|)/2} [J, K] \otimes k! \langle J, W, \bar{W}, \bar{I} \rangle \right).$$

Using the above projection pr_p , we introduce a $Pin(N)$ -equivariant homomorphism pr_T from $\Delta_{(\epsilon_1)} \otimes \otimes^k V$ to $\Delta_{(\epsilon_2)} \otimes \otimes^{k-p} V$ as follows. (If $N = 2n + 1$, we assume $\epsilon_2 = (-1)^p \epsilon_1$ is satisfied.)

Let $T = \{t_1, t_2, \dots, t_p\}$ ($t_1 < t_2 < \dots < t_p$) be a subset of $[1, k]$ and denote the positions in the tensor product $\otimes^k V$.

First we prepare a general notation.

Definition 6.3. Let Alt_T be the alternating operator of the tensor components of $\otimes^k V$ whose positions are in T . Namely

$$\begin{aligned} \text{Alt}_T(v_1 \otimes v_2 \otimes \dots \otimes v_k) = & \frac{1}{p!} \sum_{\sigma \in \mathfrak{S}_p} \epsilon(\sigma) v_1 \otimes \dots \otimes v_{t_{\sigma^{-1}(1)}} \otimes \dots \otimes v_{t_{\sigma^{-1}(2)}} \otimes \dots \otimes v_{t_{\sigma^{-1}(p)}} \otimes \dots \otimes v_k. \end{aligned}$$

Then Alt_T acts naturally on $\Delta \otimes \otimes^k V$ and we consider the operator Alt_T itself has the alternating property with the indices in T .

Definition 6.4. $\text{pr}_{\mathbf{T}} : \Delta_{(\epsilon_1)} \otimes \otimes^k V \longrightarrow \Delta_{(\epsilon_2)} \otimes \otimes^{k-p} V$ is defined by the composition of $\text{Alt}_{\mathbf{T}}$ and pr_p , namely $\text{pr}_{\mathbf{T}} = \text{pr}_p \circ \text{Alt}_{\mathbf{T}}$. Here pr_p acts on the alternating tensors of the position \mathbf{T} in $\Delta \otimes \otimes^k V$.

From the definition, $\text{pr}_{\mathbf{T}}$ is the element in $\text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)} \otimes \otimes^k V, \Delta_{(\epsilon_2)} \otimes \otimes^{k-p} V)$ and has the alternating property with the indices in \mathbf{T} .

Next we define the element $\text{inj}_{\mathbf{T}}$ in $\text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)} \otimes \otimes^{k-p} V, \Delta_{(\epsilon_2)} \otimes \otimes^k V)$ as follows.

Definition 6.5. $\text{inj}_{\mathbf{T}} : \Delta_{(\epsilon_1)} \otimes \otimes^{k-p} V \longrightarrow \Delta_{(\epsilon_2)} \otimes \otimes^k V$ is defined by the composition of $\text{inj}_p : \Delta \longrightarrow \Delta \otimes \wedge^p V$ and the linear immersion of the alternating tensor to the positions indexed by \mathbf{T} . Namely the image of this linear immersion of the exterior product $\langle \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} \rangle$ with the (ordered) index set \mathbf{T} is the tensor obtained by inserting the first component of this alternating tensor in the t_1 -th position of $\Delta \otimes \otimes^{k-p} V$ and inserting the second component in the t_2 -th position of the resulting tensor in $\Delta \otimes \otimes^{k-p+1} V$ and so on. We write this linear immersion by the exterior bracket with the subscript \mathbf{T} e.g., $\langle \mathbf{J}, \mathbf{W}, \bar{\mathbf{W}}, \bar{\mathbf{I}} \rangle_{\mathbf{T}}$.

From the definition, $\text{inj}_{\mathbf{T}}$ has also the alternating property with the indices in \mathbf{T} .

So we define another parametrization of the elements in \mathbf{CP}_1^k by the generalized Brauer diagrams \mathbf{GB}_1^k .

To a given generalized Brauer diagram, let us denote the indices of the isolated vertices in the upper row by \mathbf{T}_u and in the lower row by \mathbf{T}_ℓ . We correspond the following $\text{Pin}(N)$ -equivariant homomorphism to it.

To the isolated vertices in the upper row, we correspond $\text{pr}_{\mathbf{T}_u}$ and the rest is determined by the same rule as in the usual Brauer diagrams. To the isolated vertices in the lower row, we correspond $\text{inj}_{\mathbf{T}_\ell}$ and the rest is determined by the same rule as in the usual Brauer diagrams.

We call this parametrization the representation theoretical parametrization by the generalized Brauer diagrams and to distinguish the invariant theoretical ones, we add the subscript *rt* for the representation theoretical ones and the subscript *inv* for the invariant theoretical ones to the generalized Brauer diagrams. At this moment, we have not yet known that these representation theoretical operators span the whole space of the centralizer spaces linearly. If $k \leq n$ and $l \leq n$, we show this in the section 7, namely they become linearly independent and a base of \mathbf{CP}_1^k .

7. DESCRIPTION OF THE INVARIANT THEORETICAL EQUIVARIANT HOMOMORPHISMS

In this section, we give the description of the invariant theoretical equivariant homomorphisms. First in the invariant theoretical parametrization, we interpret the invariant element \mathfrak{J}_r (ref. (5.1))

corresponding to the isolated vertices in a generalized Brauer diagram as the equivariant homomorphism. The corresponding element in $\Delta_{(\epsilon_1)} \otimes \Delta_{(\epsilon_2)}^* \otimes \bigwedge^r V$ is given by $((\phi_r \circ \iota) \otimes \text{id})(\mathfrak{I}_r)$. (If $N = 2n + 1$, we assume $\epsilon_2 = (-1)^r \epsilon_1$ is satisfied.)

Let us recall that the index set \mathbf{t} which parameterizes $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ corresponds to the isolated vertices in the generalized Brauer diagrams. (We note that we numbered the vertices from 1 to $k + l$ when we defined $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$.) We put $\mathbf{T}_u = \mathbf{t} \cap [1, k]$ and define $\mathbf{T}_\ell = \{s_1, s_2, \dots, s_q\} \subseteq [1, l]$ by $\{k + s_1, k + s_2, \dots, k + s_p\} = \mathbf{t} \cap [k + 1, k + l]$. We put $p = |\mathbf{T}_u|$ and $q = |\mathbf{T}_\ell|$ and then $p + q = r$. We consider the homomorphism caused by the isolated vertices in $\mathbf{T}_{\mathbf{t},\{\mathbf{m},\mathbf{l}\}}$ and for convenience sake, we multiply $2^{n/2}$ to it and we denote the resulting homomorphism by $\psi_{\mathbf{T}_\ell}^{\mathbf{T}_u}$. When we need not to show the index set explicitly, we just write ψ_q^p . As a homomorphism from $\Delta \otimes \bigotimes^p V$ to $\Delta \otimes \bigotimes^q V$ (here we pick up only the \mathbf{T}_u -th components in the upper row $\Delta \otimes \bigotimes^k V$ and \mathbf{T}_ℓ -th components in the lower row $\Delta \otimes \bigotimes^l V$), $\psi_{\mathbf{T}_\ell}^{\mathbf{T}_u}$ is the composition of three homomorphisms, namely $\text{Alt}_{\mathbf{T}_u} : \Delta \otimes \bigotimes^p V \rightarrow \Delta \otimes \bigwedge^p V$ and the homomorphism caused by the \mathfrak{I}_r from $\Delta \otimes \bigwedge^p V$ to $\Delta \otimes \bigwedge^q V$ and finally the linear immersion of the exterior products of degree q to the prescribed positions in the tensor products. We note that in the space $\bigwedge^p V \otimes \bigwedge^q V$, $\bigwedge^r V$ ($r = p + q$) occurs exactly once for any $r \leq N$.

We divide the exterior product $\bigwedge^r V$ occurring in \mathfrak{I}_r into the first p tensor product and the remaining q tensor product. We also consider $\psi_{\mathbf{T}_\ell}^{\mathbf{T}_u}$ has the alternating property both in the superscripts and the subscripts respectively.

Direct computation tells us that the equivariant homomorphism ψ_q^p is given as follows.

Theorem 7.1. *For $\text{Pin}(2n + 1)$, the homomorphism ψ_q^p is given as follows.*

$$\begin{aligned} \psi_q^p([\mathbf{T}] \otimes \langle \mathbf{I}, \mathbf{S}_2, \overline{\mathbf{S}}_2, \overline{\mathbf{J}} \rangle) &= \sum_{i_2+j_2+s_1+s_3+2s_4=q} (-1)^{\sum_{i=1}^4 |\mathbf{S}_i - \mathbf{S}_i \cap \mathbf{K}|} 2^{(i_1+j_1+i_2+j_2)/2} (-1)^{j_2 s_1 + s_3 j_2 + s_3 + i_1 q} \\ &\times \epsilon \left(\begin{matrix} \mathbf{T} \\ \mathbf{I}_1, & \mathbf{I}_2, & \mathbf{K} \end{matrix} \right) \epsilon \left(\begin{matrix} \mathbf{I} \\ \mathbf{I}_1, & \mathbf{S}_3 \end{matrix} \right) \epsilon \left(\begin{matrix} \overline{\mathbf{J}} \\ \overline{\mathbf{S}}_1, & \overline{\mathbf{J}}_1 \end{matrix} \right) [\mathbf{J}_1, \mathbf{J}_2, \mathbf{K}] \otimes q! \langle \mathbf{J}_2, \mathbf{S}_3, \mathbf{S}_4, \overline{\mathbf{S}}_4, \overline{\mathbf{S}}_1, \overline{\mathbf{I}}_2 \rangle \\ &+ \sum_{i_2+j_2+s_1+s_3+2s_4=q-1} (-1)^{|\mathbf{K} - \mathbf{K} \cap (\cup_{i=1}^4 \mathbf{S}_i)|} 2^{(i_1+j_1+i_2+j_2)/2} (-1)^{j_2(s_1+s_3)+s_2+i_1 q} \\ &\times \epsilon \left(\begin{matrix} \mathbf{T} \\ \mathbf{I}_1, & \mathbf{I}_2, & \mathbf{K} \end{matrix} \right) \epsilon \left(\begin{matrix} \mathbf{I} \\ \mathbf{I}_1, & \mathbf{S}_3 \end{matrix} \right) \epsilon \left(\begin{matrix} \overline{\mathbf{J}} \\ \overline{\mathbf{S}}_1, & \overline{\mathbf{J}}_1 \end{matrix} \right) [\mathbf{J}_1, \mathbf{J}_2, \mathbf{K}] \otimes q! \langle \mathbf{J}_2, \mathbf{S}_3, \mathbf{S}_4, \mathbf{O}', \overline{\mathbf{S}}_4, \overline{\mathbf{S}}_1, \overline{\mathbf{I}}_2 \rangle. \end{aligned}$$

In the above, the sums run over all the \mathbf{I}_1 ($\mathbf{I}_1 \subseteq \mathbf{T} \cap \mathbf{I}$) and \mathbf{J}_1 ($\mathbf{J}_1 \subseteq ([1, n] - \mathbf{T}) \cap \mathbf{J}$) and \mathbf{I}_2 ($\mathbf{I}_2 \subseteq \mathbf{T} - (\mathbf{T} \cap (\mathbf{I} \cup \mathbf{S}_2 \cup \mathbf{J}))$) and \mathbf{J}_2 ($\mathbf{J}_2 \subseteq ([1, n] - \mathbf{T}) - (([1, n] - \mathbf{T}) \cap (\mathbf{I} \cup \mathbf{S}_2 \cup \mathbf{J}))$) and \mathbf{S}_4 ($\mathbf{S}_4 \subseteq [1, n] - \mathbf{I} - \mathbf{J} - \mathbf{S}_2 - \mathbf{J}_2 - \mathbf{I}_2$). We also put $\mathbf{I} - \mathbf{I}_1 = \mathbf{S}_3$ and $\mathbf{J} - \mathbf{J}_1 = \mathbf{S}_1$.

and $\mathbf{T} - \mathbf{I}_1 - \mathbf{I}_2 = \mathbf{K}$ and we denote the sizes of the sets by their small letters. As before, we assume that the indices in the sets \mathbf{I} , $\bar{\mathbf{J}}$ and the rest are in the increasing order.

For the element $\langle \mathbf{I}, \mathbf{S}_2, \mathbf{O}', \bar{\mathbf{S}}_2, \bar{\mathbf{J}} \rangle$, we have

$$\begin{aligned} \psi_q^p([\mathbf{T}] \otimes \langle \mathbf{I}, \mathbf{S}_2, \mathbf{O}', \bar{\mathbf{S}}_2, \bar{\mathbf{J}} \rangle) = \\ \sum_{i_2+j_2+s_1+s_3+2s_4=q} (-1)^{|\mathbf{K}-\mathbf{K} \cap (\cup_{i=1}^4 \mathbf{S}_i)|} 2^{(i_1+j_1+i_2+j_2)/2} (-1)^{j_2(s_1+s_3+1)+s_4+i_1q} \times \\ \epsilon \begin{pmatrix} \mathbf{T} \\ \mathbf{I}_1, \mathbf{I}_2, \mathbf{K} \end{pmatrix} \epsilon \begin{pmatrix} \mathbf{I} \\ \mathbf{I}_1, \mathbf{S}_3 \end{pmatrix} \epsilon \begin{pmatrix} \bar{\mathbf{J}} \\ \bar{\mathbf{S}}_1, \bar{\mathbf{J}}_1 \end{pmatrix} [\mathbf{J}_1, \mathbf{J}_2, \mathbf{K}] \otimes q! \langle \mathbf{J}_2, \mathbf{S}_3, \mathbf{S}_4, \bar{\mathbf{S}}_4, \bar{\mathbf{S}}_1, \bar{\mathbf{I}}_2 \rangle \end{aligned}$$

and the notations are the same as above.

For $\text{Pin}(2n)$, the homomorphism ψ_q^p is given as follows.

For even $r = p + q$, we have

$$\begin{aligned} \psi_q^p([\mathbf{T}] \otimes \langle \mathbf{I}, \mathbf{S}_2, \bar{\mathbf{S}}_2, \bar{\mathbf{J}} \rangle) = \\ \sum_{i_2+j_2+s_1+s_3+2s_4=q} (-1)^{\sum_{i=1}^4 |\mathbf{S}_i - \mathbf{S}_i \cap \mathbf{K}|} 2^{(i_1+j_1+i_2+j_2)/2} (-1)^{j_2s_1+s_3j_2+s_3+i_1q} \times \\ \epsilon \begin{pmatrix} \mathbf{T} \\ \mathbf{I}_1, \mathbf{I}_2, \mathbf{K} \end{pmatrix} \epsilon \begin{pmatrix} \mathbf{I} \\ \mathbf{I}_1, \mathbf{S}_3 \end{pmatrix} \epsilon \begin{pmatrix} \bar{\mathbf{J}} \\ \bar{\mathbf{S}}_1, \bar{\mathbf{J}}_1 \end{pmatrix} [\mathbf{J}_1, \mathbf{J}_2, \mathbf{K}] \otimes q! \langle \mathbf{J}_2, \mathbf{S}_3, \mathbf{S}_4, \bar{\mathbf{S}}_4, \bar{\mathbf{S}}_1, \bar{\mathbf{I}}_2 \rangle, \end{aligned}$$

and the notations are the same as above.

For odd $r = p + q$, we have

$$\begin{aligned} \psi_q^p([\mathbf{T}] \otimes \langle \mathbf{I}, \mathbf{S}_2, \bar{\mathbf{S}}_2, \bar{\mathbf{J}} \rangle) = \\ \sum_{i_2+j_2+s_1+s_3+2s_4=q} (-1)^{\sum_{i=1}^4 |\mathbf{S}_i - \mathbf{S}_i \cap \mathbf{K}| + |\mathbf{J}_1| + |\mathbf{J}_2| + |\mathbf{K}|} 2^{(i_1+j_1+i_2+j_2)/2} (-1)^{j_2s_1+s_3j_2+s_3+i_1q} \times \\ \epsilon \begin{pmatrix} \mathbf{T} \\ \mathbf{I}_1, \mathbf{I}_2, \mathbf{K} \end{pmatrix} \epsilon \begin{pmatrix} \mathbf{I} \\ \mathbf{I}_1, \mathbf{S}_3 \end{pmatrix} \epsilon \begin{pmatrix} \bar{\mathbf{J}} \\ \bar{\mathbf{S}}_1, \bar{\mathbf{J}}_1 \end{pmatrix} [\mathbf{J}_1, \mathbf{J}_2, \mathbf{K}] \otimes q! \langle \mathbf{J}_2, \mathbf{S}_3, \mathbf{S}_4, \bar{\mathbf{S}}_4, \bar{\mathbf{S}}_1, \bar{\mathbf{I}}_2 \rangle, \end{aligned}$$

and the notations are the same as above.

For any $\sigma \in \mathfrak{S}_k$ and $\tau \in \mathfrak{S}_l$, we have $\tau \circ \psi_{\mathbf{T}_\ell}^{\mathbf{T}_u} \circ \sigma = \psi_{\tau(\mathbf{T}_\ell)}^{\sigma^{-1}(\mathbf{T}_u)}$. So it is enough to give the explicit description for $\psi_{[1, \mathbf{q}]}^{[1, \mathbf{p}]}$ in terms of the representation theoretical operators.

Theorem 7.2. Let $p \leq n$ and $q \leq n$.

Then we have

(7.2.1)

$$\psi_{[1, \mathbf{q}]}^{[1, \mathbf{p}]} = \sum_{i=0}^{\min(p, q)} \epsilon(i) \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{inj}_{\{\sigma([1, \mathbf{q}])\}}}{(q-i)!} \frac{\left(\tau([1, \mathbf{i}]) \right)}{i!} \frac{\text{pr}_{\{\tau([1, \mathbf{p}])\}}}{(p-i)!}.$$

Here $\epsilon(i) = (-1)^{(p-i)(q-i)}$ for $\text{Pin}(2n+1)$ and $\epsilon(i) = 1$ for $\text{Pin}(2n)$ ($0 \leq i \leq \min(p, q)$).

Also we have

(7.2.2)

$$\text{inj}_{[1, \mathbf{q}]} \circ \text{pr}_{[1, \mathbf{p}]} = \sum_{i=0}^{\min(p, q)} \epsilon(i) \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p}} \epsilon(\sigma) \epsilon(\tau) \frac{\psi_{\{\sigma([1, \mathbf{q}])\}}^{\{\tau([1, \mathbf{p}])\}}}{(q-i)!(p-i)!} \otimes \frac{\binom{\tau([1, \mathbf{i}])}{\sigma([1, \mathbf{i}])}}{i!}.$$

Here $\epsilon(i) = (-1)^{i+pq}$ for $\text{Pin}(2n+1)$ and $\epsilon(i) = (-1)^i$ for $\text{Pin}(2n)$ ($0 \leq i \leq \min(p, q)$).

We denote $\sigma([1, \mathbf{q}]) = \{\sigma(1), \sigma(2), \dots, \sigma(q)\}$ and $\tau([1, \mathbf{p}]) = \{\tau(1), \tau(2), \dots, \tau(p)\}$ and $\binom{\tau([1, \mathbf{i}])}{\sigma([1, \mathbf{i}])}$ stands for the partial permutation of the tensor components obtained by sending the $\tau(1)$ -th component in $\Delta \otimes \otimes^p V$ in the upper row to the $\sigma(1)$ -th component in $\Delta \otimes \otimes^q V$ in the lower row and so on.

So if $k \leq n$ and $l \leq n$, the generalized Brauer diagrams under the representation theoretical parametrization also span the whole space \mathbf{CP}_1^k and become a linear base of \mathbf{CP}_1^k .

Remark 7.3. The right-hand side of (7.2.1) can be considered as the composition of the homomorphisms in this order, but the formula (7.2.2) cannot. So we put \otimes in the right-hand side to show that each summand becomes a homomorphism as a whole.

Proof. Both sides of the formula (7.2.1) are the homomorphisms from $\Delta \otimes \otimes^p V$ to $\Delta \otimes \otimes^q V$ and moreover both of them are the compositions of the alternating operators $\text{Alt}_{[1, \mathbf{p}]}$ and the homomorphisms from $\Delta \otimes \wedge^p V$ to $\Delta \otimes \wedge^q V$ and the linear immersion from $\wedge^q V$ to $\otimes^q V$.

As we noted before, we have $\dim \text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)} \otimes \wedge^p V, \Delta_{(\epsilon_2)} \otimes \wedge^q V) = \min(p, q) + 1$. Here we assume that $\epsilon_1 \epsilon_2 = (-1)^{p+q}$ for $\text{Pin}(2n+1)$. (ref. Lemma 5.4.)

On the way to proving the formula (7.2.1), we show that the terms occurring in the right-hand side of the formula are linearly independent.

The proof goes as follows.

For $0 \leq i \leq \min(p, q)$, let us put

$$i\phi_{[1, \mathbf{q}]}^{[1, \mathbf{p}]} = \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{inj}_{\{\sigma([1, \mathbf{q}])\}}}{(q-i)!} \frac{\binom{\tau([1, \mathbf{i}])}{\sigma([1, \mathbf{i}])}}{i!} \frac{\text{pr}_{\{\tau([1, \mathbf{p}])\}}}{(p-i)!}.$$

We find a sequence of pairs of elements $\{x_i, y_i\}$ (here $x_i \in \Delta \otimes \otimes^p V$, $y_i \in \Delta \otimes \otimes^q V$ and $0 \leq i \leq \min(p, q)$) satisfying the following condition.

For $\text{Pin}(2n+1)$, the coefficients of y_i in the image $j\phi_{[1, \mathbf{q}]}^{[1, \mathbf{p}]}(x_i)$ is 0 if $j \neq i$ and non-zero if $j = i$.

For $\text{Pin}(2n)$, the coefficients of y_{2i} in the image $j\phi_{[1,q]}^{[1,p]}(x_{2i})$ is 0 if $j \neq 2i$ and non-zero if $j = 2i$ and the coefficients of y_{2i+1} in the image $j\phi_{[1,q]}^{[1,p]}(x_{2i+1})$ is 0 if $j \neq 2i, 2i+1$ and non-zero if $j = 2i, 2i+1$.

If we find such a sequence, $i\phi_{[1,q]}^{[1,p]}$'s ($0 \leq i \leq \min(p, q)$) are linearly independent and span the intertwining space $\text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)} \otimes \bigwedge^p V, \Delta_{(\epsilon_2)} \otimes \bigwedge^q V)$.

We put

$$\psi_{[1,q]}^{[1,p]} = \sum_{i=0}^{\min(p,q)} d_i i\phi_{[1,q]}^{[1,p]}$$

and give the above sequence $\{x_i, y_i\}$ and apply both sides to the elements x_i and compare the coefficients of y_i . Then we determine the coefficients d_i inductively.

We first prove the formula (7.2.1) for $\text{Pin}(2n+1)$.

For abbreviation, we write $\otimes\{\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_r\}$ for the tensor product of the base elements $u_{i_1} \otimes u_{i_2} \otimes \dots \otimes u_{i_r}$ and denote $\otimes\{\mathbf{k}, \mathbf{k}+1, \dots, \mathbf{k}+r\}$ by $\otimes\{\overline{\mathbf{k}}, \mathbf{k}+r\}$ and $\otimes\{\overline{\mathbf{k}+r}, \overline{\mathbf{k}+r-1}, \dots, \overline{\mathbf{k}}\}$ by $\otimes\{\overline{\mathbf{k}+r}, \mathbf{k}\}$ and we use the juxtaposition of the above notations.

We take cases for the parities of p and q .

Case 1 $p = 2r$ and $q = 2s$

We note $r + s \leq n$.

Case 1-0. The determination of d_0 .

We take $y_0 = [\emptyset] \otimes \{\overline{\mathbf{r}+1}, \mathbf{r}+s\}, \overline{\mathbf{r}+s}, \mathbf{r}+1\}$, $x_0 = [\emptyset] \otimes \{\overline{\mathbf{1}}, \mathbf{r}\}, \overline{\mathbf{r}}, \mathbf{1}\}$.

To calculate the image of $\psi_{[1,q]}^{[1,p]}(x_0)$, we put $\mathbf{S}_2 = \{\overline{\mathbf{1}}, \mathbf{r}\}$ and $\mathbf{T} = \mathbf{J} = \mathbf{I} = \emptyset$. So $\mathbf{I}_1 = \mathbf{J}_1 = \mathbf{I}_2 = \mathbf{S}_3 = \mathbf{S}_1 = \mathbf{K} = \emptyset$ and $\mathbf{J}_2 \subseteq [1, n] - \mathbf{S}_2$ and $\mathbf{S}_4 \subseteq [1, n] - \mathbf{J}_2 - \mathbf{S}_2$ and apply Theorem 7.1. Since we consider the coefficients of y_0 , $\mathbf{J}_2 = \emptyset$ and $\mathbf{S}_4 = \{\overline{\mathbf{r}+1}, \mathbf{r}+s\}$ and the coefficient is given by $(-1)^{r+s}$. On the other hand, since there are no common numbers in the sets $\{\overline{\mathbf{1}}, \mathbf{r}\}$ and $\{\overline{\mathbf{r}+1}, \mathbf{r}+s\}$, only the term coming from $i = 0$ contributes to the coefficient.

We calculate $\text{inj}_{\{\overline{\mathbf{1}}, \mathbf{q}\}} \circ \text{pr}_{\{\overline{\mathbf{1}}, \mathbf{p}\}}(x_0)$. Since $\mathbf{I} = \mathbf{J} = \mathbf{T} = \emptyset$ and $\mathbf{W} = \{\overline{\mathbf{1}}, \mathbf{r}\}$ in the definition 6.1, $\text{pr}_{\{\overline{\mathbf{1}}, \mathbf{p}\}}(x_0) = (-1)^r [\emptyset]$. We calculate the coefficient of y_0 in the image $\text{inj}_{\{\overline{\mathbf{1}}, \mathbf{q}\}}((-1)^r [\emptyset])$. Since $\mathbf{T} = \emptyset$, $\mathbf{I} = \mathbf{K} = \emptyset$ and we only consider the case $\mathbf{J} = \emptyset$ and $\mathbf{W} = \{\overline{\mathbf{r}+1}, \mathbf{r}+s\}$ in the definition 6.2. So the coefficient is given by $(-1)^{r+s}$. Therefore we have $d_0 = 1 = (-1)^{pq}$ in this case.

Case 1-1. The determination of d_1 .

We take $y_1 = [\mathbf{r}+1] \otimes \{\overline{\mathbf{1}}, \overline{\mathbf{r}+1}, \mathbf{r}+s\}, \overline{\mathbf{r}+s}, \mathbf{r}+2\}$, $x_1 = [\emptyset] \otimes \{\overline{\mathbf{1}}, \mathbf{r}\}, \mathbf{0}', \overline{\mathbf{r}}, \mathbf{2}\}$.

In the left-hand side, since $\mathbf{I} = \{\mathbf{1}\}$, $\mathbf{S}_2 = \{\overline{\mathbf{2}}, \mathbf{r}\}$, $\mathbf{J} = \emptyset$, $\mathbf{T} = \emptyset$, we have $\mathbf{J}_1 = \emptyset$, $\mathbf{S}_1 = \emptyset$, $\mathbf{K} = \emptyset$, $\mathbf{I}_1 = \emptyset$, $\mathbf{I}_2 = \emptyset$. Only the case of $\mathbf{S}_3 = \{\mathbf{1}\}$, $\mathbf{J}_2 = \{\mathbf{r}+1\}$ and $\mathbf{S}_4 = \{\overline{\mathbf{r}+2}, \mathbf{r}+s\}$ contributes to the coefficient of $[\mathbf{r}+1] \otimes \{\overline{\mathbf{1}}, \overline{\mathbf{r}+1}, \mathbf{r}+s\}, \overline{\mathbf{r}+s}, \mathbf{r}+2\}$. Namely we have $\psi_{[1,q]}^{[1,p]}([\emptyset] \otimes \{\overline{\mathbf{1}}, \mathbf{r}\}, \mathbf{0}', \overline{\mathbf{r}}, \mathbf{2}\}) = \sqrt{2}(-1)^{s-1+1+1}[\mathbf{r}+1] \otimes q! <$

$r + 1, 1, [r + 2, r + s], \overline{[r + s, r + 2]} >$ + the other terms. So the coefficient is $\sqrt{2}(-1)^s$ in the left-hand side.

On the other hand since $\{1, [r + 1, r + s], \overline{[r + s, r + 2]}\} \cap \{[1, r], 0', \overline{[r, 2]}\} = \{1\}$, the possible terms to contribute to the coefficient are only $i = 0$ or $i = 1$. Moreover if we consider the weight of $[\emptyset] \otimes \{[1, r], 0', \overline{[r, 2]}\}$, this is not in the weight set of Δ , so the image of pr must be 0 and $i = 0$ term also does not contribute to the coefficient.

In the case of $i = 1$, to obtain the coefficient, $\sigma(1)$ and $\tau(1)$ must be 1 and in the $i = 1$ term, we have only to consider the term $d_1 \text{inj}_{\{[2, q]\}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{pr}_{\{[2, p]\}} \cdot$

Since $\text{pr}_{\{[2, p]\}}([\emptyset] \otimes < [2, r], 0', \overline{[r, 2]} >) = [\emptyset]$ and $\text{inj}_{\{[2, q]\}}([\emptyset]) = \sqrt{2}(-1)^{s-1} [r + 1] \otimes (q - 1)! < [r + 1, r + s], \overline{[r + s, r + 2]} >$, the coefficient of the right-hand side is $d_1 \sqrt{2}(-1)^{s-1}$. So we have $d_1 = -1$.

The similar argument goes on for the general case and we only give the elements x_i and y_i and d_i for each of the cases.

Case 1-2i The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, 2i], [r + i + 1, r + s], \overline{[r + s, r + i + 1]}\}$, $x_{2i} = [\emptyset] \otimes \{[1, r + i], \overline{[r + i, 2i + 1]}\}$. In this case $d_{2i} = 1$.

Case 1-2i + 1 The determination of d_{2i+1} .

We take $y_{2i+1} = [r + i + 1] \otimes \{[1, 2i + 1], [r + i + 1, r + s], \overline{[r + s, r + i + 2]}\}$, $x_{2i+1} = [\emptyset] \otimes \{[1, r + i], 0', \overline{[r + i, 2i + 2]}\}$. In this case $d_{2i+1} = -1$.

Case 2. $p = 2r + 1$ and $q = 2s$.

In this case, we have $r + s < n$.

Case 2-2i. The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, 2i], [r + i + 1, r + s], \overline{[r + s, r + i + 1]}\}$, $x_{2i} = [\emptyset] \otimes \{[1, r + i], 0', \overline{[r + i, 2i + 1]}\}$. In this case $d_{2i} = 1$.

Case 2-2i + 1. The determination of d_{2i+1} .

We take $y_{2i+1} = [\emptyset] \otimes \{[1, 2i + 1], [r + i + 2, r + s], 0', \overline{[r + s, r + i + 2]}\}$, $x_{2i+1} = [\emptyset] \otimes \{[1, r + i + 1], \overline{[r + i + 1, 2i + 2]}\}$.

In this case $d_{2i+1} = 1$.

Case 3. $p = 2r$ and $q = 2s + 1$.

In this case, we have $r + s < n$.

Case 3-2i. The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, 2i], [r + i + 1, r + s], \overline{[r + s, r + i + 1]}\}$, $x_{2i} = [2i + 1] \otimes \{[1, r + i], 0', \overline{[r + i, 2i + 2]}\}$.

In this case $d_{2i} = 1$.

Case 3-2i + 1. The determination of d_{2i+1} .

We take $y_{2i+1} = [\emptyset] \otimes \{[1, 2i + 1], [r + i + 1, r + s], \overline{[r + s, r + i + 1]}\}$, $x_{2i+1} = [\emptyset] \otimes \{[1, r + i], 0', \overline{[r + i, 2i + 2]}\}$.

In this case $d_{2i+1} = 1$.

Case 4. $p = 2r + 1$ and $q = 2s + 1$.

In this case, we note that $r + s \leq n - 1$.

Case 4-2i. The determination of d_{2i} .

We take $y_{2i} = [\mathbf{r} + \mathbf{i} + 1] \otimes \{[1, 2\mathbf{i}], [\mathbf{r} + \mathbf{i} + 1, \mathbf{r} + \mathbf{s} + 1], \overline{[\mathbf{r} + \mathbf{s} + 1, \mathbf{r} + \mathbf{i} + 2]}\}$,
 $x_{2i} = [\emptyset] \otimes \{[1, \mathbf{r} + \mathbf{i}], \mathbf{0}', \overline{[\mathbf{r} + \mathbf{i}, 2\mathbf{i} + 1]}\}$.

In this case $d_{2i} = -1$.

Case 4-2i + 1. The determination of d_{2i+1} .

We take $y_{2i+1} = [\emptyset] \otimes \{[1, 2\mathbf{i} + 1], [\mathbf{r} + \mathbf{i} + 2, \mathbf{r} + \mathbf{s} + 1], \overline{[\mathbf{r} + \mathbf{s} + 1, \mathbf{r} + \mathbf{i} + 2]}\}$,
 $x_{2i+1} = [\emptyset] \otimes \{[1, \mathbf{r} + \mathbf{i} + 1], \overline{[\mathbf{r} + \mathbf{i} + 1, 2\mathbf{i} + 2]}\}$.

In this case $d_{2i+1} = 1$.

We verify all the cases and we have proved the formula (7.2.1) for $Pin(2n + 1)$.

We will prove the formula (7.2.1) for $Pin(2n)$.

Case 1. $p = 2r$ and $q = 2s$.

We note $r + s \leq n$.

Case 1-0. The determination of d_0 .

We take $y_0 = [\emptyset] \otimes \{[\mathbf{r} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + 1]}\}$, $x_0 = [\emptyset] \otimes \{[1, \mathbf{r}], \overline{[\mathbf{r}, 1]}\}$.
In this case $d_0 = 1$.

Case 1-1. The determination of d_1 .

We take $y_1 = [1, \mathbf{r} + 1] \otimes \{1, [\mathbf{r} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + 2]}\}$, $x_1 = [\emptyset] \otimes \{[1, \mathbf{r}], \overline{[\mathbf{r}, 1]}\}$.

In this case, the coefficient of the left-hand side is 0.

On the right-hand side, since $\{1, [\mathbf{r} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + 2]}\} \cap \{[1, \mathbf{r}], \overline{[\mathbf{r}, 1]}\} = \{1\}$, the possible terms to contribute to the coefficient are only $i = 0$ or $i = 1$.

We first calculate the contribution coming from the term $i = 1$.
The image of $\text{pr}_{2r-1}([\emptyset] \otimes \{[2, \mathbf{r}], \overline{[\mathbf{r}, 1]}\})$ is $(-1)^{r-1+1}\sqrt{2}[1]$. The image $\text{inj}_{2s-1}([1])$ is given by $(-1)^{s-1+1+1}\sqrt{2}[\mathbf{r} + 1, 1] \otimes (2s-1)! < [\mathbf{r} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + 2]} > +$ the other terms. So the contribution from this term to the coefficient is $2(-1)^{r+s}$.

We calculate the contribution coming from the term $i = 0$. The image of $\text{pr}_{2r}([\emptyset] \otimes \{[1, \mathbf{r}], \overline{[\mathbf{r}, 1]}\})$ is $(-1)^r[\emptyset]$. The image $\text{inj}_{2s}([\emptyset])$ is given by $(-1)^{s-1}2[1, \mathbf{r} + 1] \otimes (2s)! < 1, [\mathbf{r} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + 2]} > +$ the other terms. So the contribution from this term to the coefficient is $2(-1)^{r+s-1}$. Hence we have $d_1 = 1$.

Case 1-2i. The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, 2\mathbf{i}], [\mathbf{r} + \mathbf{i} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + \mathbf{i} + 1]}\}$, $x_{2i} = [\emptyset] \otimes \{[1, \mathbf{r} + \mathbf{i}], \overline{[\mathbf{r} + \mathbf{i}, 2\mathbf{i} + 1]}\}$.

In this case $d_{2i} = 1$.

Case 1-2i + 1. The determination of d_{2i+1} .

We take $y_{2i+1} = [2\mathbf{i} + 1, \mathbf{r} + \mathbf{i} + 1] \otimes \{[1, 2\mathbf{i} + 1], [\mathbf{r} + \mathbf{i} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + \mathbf{i} + 2]}\}$,
 $x_{2i+1} = [\emptyset] \otimes \{[1, \mathbf{r} + \mathbf{i}], \overline{[\mathbf{r} + \mathbf{i}, 2\mathbf{i} + 1]}\}$. In this case $d_{2i+1} = 1$.

For the remaining cases of parities of p and q , the similar argument goes on and so we omit it.

Next we show the formula (7.2.2).

For $0 \leq i \leq \min(p, q)$, let us put

$$i\psi_{[1,q]}^{[1,p]} = \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p}} \epsilon(\sigma) \epsilon(\tau) \frac{\psi_{\{\sigma([i+1,q])\}}^{\{\tau([i+1,p])\}}}{(q-i)!(p-i)!} \otimes \frac{\left(\begin{smallmatrix} \tau([1,i]) \\ \sigma([1,i]) \end{smallmatrix} \right)}{i!}$$

We find a sequence of pairs of elements $\{x_i, y_i\}$ (here $x_i \in \Delta \otimes \otimes^p V$, $y_i \in \Delta \otimes \otimes^q V$ and $0 \leq i \leq \min(p, q)$) such that the coefficients of y_i in the image $i\psi_{[1,q]}^{[1,p]}(x_i)$ is 0 if $j > i$ and non-zero if $j = i$.

In this case $i\psi_{[1,q]}^{[1,p]}$ is equal to $\text{Alt}_{[1,q]} \circ (\phi_{[i+1,q]}^{[i+1,p]} \otimes \left(\begin{smallmatrix} [1,i] \\ [1,i] \end{smallmatrix} \right)) \circ \text{Alt}_{[1,p]}$ up to scalar and so $i\psi_{[1,q]}^{[1,p]}$'s ($0 \leq i \leq \min(p, q)$) are linearly independent. (See Lemma 5.6.)

We first put

$$\text{inj}_{[1,q]} \circ \text{pr}_{[1,p]} = \sum_{u=0}^{\min(p,q)} d_u' u\psi_{[1,q]}^{[1,p]}$$

and give the above sequence $\{x_u, y_u\}$ and apply both sides to the elements x_u and compare the coefficients of y_u as before. Then we can determine the coefficients d_u' inductively.

We take cases for the parities of p and q .

Case 1. $p = 2r$ and $q = 2s$.

We note $r + s \leq n$ because of $p \leq n$ and $q \leq n$.

Case 1-0 The determination of d_0' .

We take $y_0 = [\emptyset] \otimes \{[r+1, r+s], [\overline{r+s, r+1}]\}$, $x_0 = [\emptyset] \otimes \{[1, r], [\overline{r, 1}]\}$.

In this case $d_0' = 1$.

Case 1-1. The determination of d_1' .

We take $y_1 = [1, r+1] \otimes \{1, [r+1, r+s], [\overline{r+s, r+2}]\}$, $x_1 = [1] \otimes \{[1, r], 0', [\overline{r, 2}]\}$. The coefficient of the left-hand side is $2\sqrt{2}(-1)^{s-1}$.

In the right-hand side, by considering the number of the common letters, the u term with $u > 1$ never contributes to the coefficient. We calculate the contribution coming from the $u = 0$ term. Then we have $\mathbf{I} = \{1\}$, $\mathbf{S}_2 = \{[2, r]\}$, $\mathbf{J} = \emptyset$ and $\mathbf{T} = \{1\}$. Only the case of $\mathbf{J}_2 = \{r+1\}$, $\mathbf{K} = \{1\}$, $\mathbf{S}_4 = \{[r+2, r+s]\}$, $\mathbf{S}_3 = \{1\}$, $\mathbf{I}_1 = \emptyset$, $\mathbf{J}_1 = \emptyset$, $\mathbf{I}_2 = \emptyset$ and $\mathbf{S}_1 = \emptyset$ contributes to the coefficient. The image of the $u = 0$ term is given by $\sqrt{2}(-1)^{s-1}[r+1, 1] \otimes q! < r+1, 1, \mathbf{S}_4, \overline{\mathbf{S}_4} > +$ the other terms and the coefficient is given by $\sqrt{2}(-1)^{s-1}d_0'$. Next we consider the $u = 1$ term. Then only the case $\tau(1) = \sigma(1) = 1$ contributes to the coefficient and we calculate $d_1' \psi_{[1,q]}^{[1,p]}$. At this time only the case $\mathbf{I} = \emptyset$, $\mathbf{S}_2 = \{[2, r]\}$, $\mathbf{T} = \{1\}$, $\mathbf{J}_2 = \{r+1\}$, $\mathbf{J} = \mathbf{J}_1 = \emptyset$, $\mathbf{S}_1 = \emptyset$, $\mathbf{S}_4 = \{[r+2, r+s]\}$, $\mathbf{K} = \{1\}$, $\mathbf{I}_1 = \emptyset$, $\mathbf{I}_2 = \emptyset$ and $\mathbf{S}_3 = \emptyset$ contributes to the coefficient. The image of the $u = 1$ term is given by $\psi_{[1,q]}^{[1,p]}([1] \otimes < \mathbf{S}_2, 0', \overline{\mathbf{S}_2} >) = (-1)^{1+1+s-1}\sqrt{2}[r+1, 1] \otimes (q-1)! < r+1, \mathbf{S}_4, \overline{\mathbf{S}_4} > +$ the other terms

and the coefficient is $\sqrt{2}(-1)^s d_1'$ because of $[\mathbf{r} + 1, 1] = -[1, \mathbf{r} + 1]$ in Δ .

So we have $d_1' = -1$.

We prove $d_i' = (-1)^i$ by induction on i in this case.

We assume that the above holds for $i = 1, 2, \dots, 2u - 1$ and prove the case of $i = 2u$.

We take $y_{2u} = [[1, 2u]] \otimes \{[1, 2u], [\mathbf{r} + \mathbf{u} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + \mathbf{u} + 1]}\},$
 $x_{2u} = [[1, 2u]] \otimes \{[1, \mathbf{r} + \mathbf{u}], \overline{[\mathbf{r} + \mathbf{u}, 2u + 1]}\}.$

The coefficient of the left-hand side is $2^{2u}(-1)^{r+s}$.

We calculate the right-hand side.

The contribution to the coefficient under consideration from the $i = 0$ term comes from $\mathbf{I} = \{[1, 2u]\}, \mathbf{S}_2 = \{[2u + 1, r + u]\}, \mathbf{T} = \{[1, 2u]\},$
 $\mathbf{J} = \mathbf{J}_1 = \emptyset, \mathbf{S}_1 = \emptyset, \mathbf{S}_4 = \{[r + u + 1, r + s]\}, \mathbf{K} = \{[1, 2u]\}, \mathbf{I}_1 = \emptyset, \mathbf{I}_2 = \emptyset,$
 $\mathbf{J}_2 = \emptyset$ and $\mathbf{S}_3 = \{[1, 2u]\}.$ The coefficient is given by $(-1)^{s-u+r-u+2s} d_0'.$

We consider the contribution from the $i = 1$ term. Then we must have $1 \leq \tau(1) = \sigma(1) = j \leq 2u$ and for each j , the right-hand side of the $i = 1$ term is reduced to $\psi_{\{[1, \mathbf{p}] - \{j\}\} \{[1, \mathbf{q}] - \{j\}\}}^{\{[1, \mathbf{p}] - \{j\}\} \{[1, \mathbf{q}] - \{j\}\}}$ from the alternating property with the indices.

The contribution to the coefficient from this term comes from $\mathbf{I} = \{[1, 2u] - \{j\}\}, \mathbf{S}_2 = \{[2u + 1, r + u]\}, \mathbf{T} = \{[1, 2u]\}, \mathbf{J}_2 = \emptyset, \mathbf{J} = \mathbf{J}_1 = \emptyset,$
 $\mathbf{S}_1 = \emptyset, \mathbf{S}_4 = \{[r + u + 1, r + s]\}, \mathbf{K} = \mathbf{T}, \mathbf{I}_1 = \emptyset, \mathbf{I}_2 = \emptyset$ and $\mathbf{S}_3 = \mathbf{I}.$
Then we have $\psi_{\{[1, \mathbf{p}] - \{j\}\} \{[1, \mathbf{q}] - \{j\}\}}^{\{[1, \mathbf{p}] - \{j\}\} \{[1, \mathbf{q}] - \{j\}\}}([1, 2u] \otimes \{[1, \mathbf{r} + \mathbf{u}] - \{j\}, \overline{[\mathbf{r} + \mathbf{u}, 2u + 1]}\}) =$
 $(-1)^{s-u+r-u+2u-1} [\mathbf{T}] \otimes (q-1)! < \mathbf{S}_3, \mathbf{S}_4, \overline{\mathbf{S}_4} >_{\{[1, \mathbf{q}] - \{j\}\}} + \text{the other terms}.$

So the total contribution to the coefficient from the $i = 1$ term is given by $\binom{2u}{1}(-1)^{s+r-1} d_1'.$

We consider the contribution from the $i = 2$ term. Then we must have $\tau(1) = \sigma(1)$ and $\tau(2) = \sigma(2)$ and the right-hand side of the $i = 2$ term is given by $\sum_{1 \leq j_1 < j_2 \leq 2u} \psi_{\{[1, \mathbf{p}] - \{j_1\} - \{j_2\}\} \{[1, \mathbf{q}] - \{j_1\} - \{j_2\}\}}^{\{[1, \mathbf{p}] - \{j_1\} - \{j_2\}\} \{[1, \mathbf{q}] - \{j_1\} - \{j_2\}\}} \otimes \begin{pmatrix} j_1 & j_2 \\ j_1 & j_2 \end{pmatrix}.$

The contribution to the coefficient from this term comes from $\mathbf{I} = \{[1, 2u] - \{j_1\} - \{j_2\}\} = \mathbf{S}_3, \mathbf{S}_2 = \{[2u + 1, r + u]\}, \mathbf{T} = \{[1, 2u]\} = \mathbf{K},$
 $\mathbf{J}_2 = \emptyset, \mathbf{J} = \mathbf{J}_1 = \emptyset, \mathbf{S}_1 = \emptyset, \mathbf{S}_4 = \{[r + u + 1, r + s]\}$ and $\mathbf{I}_1 = \mathbf{I}_2 = \emptyset.$
The coefficient of the image under consideration by this map is given by $(-1)^{s-u+r-u+2u-2}$ and the total contribution coming from the $i = 2$ term is given by $\binom{2u}{2}(-1)^{s+r} d_2'.$

So from the same argument, we can deduce that the total contribution to the coefficient from the i term is given by $\binom{2u}{i}(-1)^{s+r-i} d_i'.$

From the induction hypothesis, we have $d_{2u}' = 1$ since $2^{2u} = \sum_{i=0}^{2u} \binom{2u}{i}.$

Next we assume that the $d_i' = (-1)^i$ holds for $i = 1, 2, \dots, 2u$ and prove the case of $i = 2u + 1$.

We take $y_{2u+1} = [[1, 2u + 1], \mathbf{r} + \mathbf{u} + 1] \otimes \{[1, 2u + 1], [\mathbf{r} + \mathbf{u} + 1, \mathbf{r} + \mathbf{s}], \overline{[\mathbf{r} + \mathbf{s}, \mathbf{r} + \mathbf{u} + 2]}\},$
 $x_{2u+1} = [[1, 2u + 1]] \otimes \{[1, \mathbf{r} + \mathbf{u}], 0', \overline{[\mathbf{r} + \mathbf{u}, 2u + 2]}\}.$

The coefficient of the left-hand side is $2^{2u+1+1/2}(-1)^{s-u-1}.$

We calculate the right-hand side.

The contribution to the coefficient under consideration from the $i = 0$ term comes from $\mathbf{I} = \{[1, 2u + 1]\} = \mathbf{S}_3$, $\mathbf{S}_2 = \{[2u + 2, r + u]\}$, $\mathbf{T} = \{[1, 2u + 1]\} = \mathbf{K}$, $\mathbf{J}_2 = \{r + u + 1\}$, $\mathbf{J} = \mathbf{J}_1 = \emptyset$, $\mathbf{S}_1 = \emptyset$, $\mathbf{S}_4 = \{[r + u + 2, r + s]\}$, $\mathbf{I}_1 = \emptyset$ and $\mathbf{I}_2 = \emptyset$. The image of ψ is $\sqrt{2}(-1)^{1(2u+1+1)+s-u-1}[\mathbf{r} + \mathbf{u} + 1, \mathbf{T}] \otimes q! < \mathbf{r} + \mathbf{u} + 1, \mathbf{S}_3, \mathbf{S}_4, \overline{\mathbf{S}_4} > + \text{the other terms and the coefficient is given by } \sqrt{2}(-1)^{s-u-1}d_0'$.

We consider the contribution from the $i = 1$ term. As before, we must have $1 \leq \tau(1) = \sigma(1) = j \leq 2u$ and for each j , the contribution to the coefficient from this term comes from $\mathbf{I} = \{[1, 2u + 1] - \{j\}\} = \mathbf{S}_3$, $\mathbf{S}_2 = \{[2u + 2, r + u]\}$, $\mathbf{T} = \{[1, 2u + 1]\} = \mathbf{K}$, $\mathbf{J}_2 = \{r + u + 1\}$, $\mathbf{J} = \mathbf{J}_1 = \emptyset$, $\mathbf{S}_1 = \emptyset$, $\mathbf{S}_4 = \{[r + u + 2, r + s]\}$, $\mathbf{I}_1 = \emptyset$ and $\mathbf{I}_2 = \emptyset$. Then we have $\psi_{\{\{1, \mathbf{p}\} - \{j\}\} \atop \{\{1, \mathbf{q}\} - \{j\}\}}([1, 2u + 1] \otimes \{[1, \mathbf{r} + \mathbf{u}] - \{j\}, 0', [\mathbf{r} + \mathbf{u}, 2u + 2]\}) = \sqrt{2}(-1)^{1+1(2u+1)+s-u-1}[\mathbf{r} + \mathbf{u} + 1, \mathbf{T}] \otimes (q-1)! < \mathbf{r} + \mathbf{u} + 1, \mathbf{S}_3, \mathbf{S}_4, \overline{\mathbf{S}_4} >_{\{\{1, \mathbf{q}\} - \{j\}\}} + \text{the other terms. From the alternating property, we have } [\mathbf{r} + \mathbf{u} + 1, \mathbf{T}] \otimes (q-1)! < \mathbf{r} + \mathbf{u} + 1, \mathbf{S}_3, \mathbf{S}_4, \overline{\mathbf{S}_4} > = -[\mathbf{T}, \mathbf{r} + \mathbf{u} + 1] \otimes (q-1)! < \mathbf{S}_3, \mathbf{r} + \mathbf{u} + 1, \mathbf{S}_4, \overline{\mathbf{S}_4} > \text{ and the total contribution to the coefficient from the } i = 1 \text{ term is given by } \binom{2u+1}{1} \sqrt{2}(-1)^{s-u}d_1'$.

Similarly we can calculate the contribution coming from the $i = 2$ term. As before, we must have $\tau(1) = \sigma(1)$ and $\tau(2) = \sigma(2)$ and the right-hand side of the $i = 2$ term is given by $\sum_{1 \leq j_1 < j_2 \leq 2u} \psi_{\{\{1, \mathbf{p}\} - \{j_1\} - \{j_2\}\} \atop \{\{1, \mathbf{q}\} - \{j_1\} - \{j_2\}\}} \otimes \begin{pmatrix} j_1 & j_2 \\ j_1 & j_2 \end{pmatrix}$.

The contribution to the coefficient from this term comes from $\mathbf{I} = \{[1, 2u + 1] - \{j_1\} - \{j_2\}\} = \mathbf{S}_3$, $\mathbf{S}_2 = \{[2u + 2, r + u]\}$, $\mathbf{T} = \{[1, 2u + 1]\} = \mathbf{K}$, $\mathbf{J}_2 = \{r + u + 1\}$, $\mathbf{J} = \mathbf{J}_1 = \emptyset$, $\mathbf{S}_1 = \emptyset$, $\mathbf{S}_4 = \{[r + u + 2, r + s]\}$ and $\mathbf{I}_1 = \mathbf{I}_2 = \emptyset$. So from the same argument, the total contribution to the coefficient from the $i = 2$ term is given by $\binom{2u+1}{2} \sqrt{2}(-1)^{s-u-1}d_2'$.

Similarly we can calculate the contribution to the coefficient from the i term and it is given by $\binom{2u+1}{i} \sqrt{2}(-1)^{s-u-1+i}d_i'$.

So from the induction hypothesis, we obtain $d_i' = (-1)^i$ in this case.

For the remaining cases of parities of p and q , the proof goes almost in a similar way. So we omit it and summarizing all the cases, we have $d_i' = (-1)^{pq+i}$.

For $\text{Pin}(2n)$, the proof goes almost same and we omit it.

From the above two formulas, if $n \geq k$, the generalized Brauer diagrams under the representation theoretical parametrization also span the full centralizer algebra \mathbf{CP}_k and they become a linear base of \mathbf{CP}_k since $\dim \mathbf{CP}_k$ is equal to the number of the generalized Brauer diagrams.

□

Corollary 7.4. *Let $p \leq n$ and $q \leq n$. For $0 \leq i \leq \min(p, q)$, if we put*

$$\phi_{[1,q]}^{[1,p]} = \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{inj}_{\{\sigma([i+1,q])\}}}{(q-i)!} \frac{\binom{\tau([1,i])}}{\sigma([1,i])!} \frac{\text{pr}_{\{\tau([i+1,p])\}}}{(p-i)!},$$

then $\phi_{[1,q]}^{[1,p]}$'s ($0 \leq i \leq \min(p, q)$) are linearly independent and span the intertwining space $\text{Hom}_{\text{Pin}(N)}(\Delta_{(\epsilon_1)} \otimes \bigwedge^p V, \Delta_{(\epsilon_2)} \otimes \bigwedge^q V)$.

Example 7.5. *We show a few example of the relations of two parametrization in figure 4 for $n \geq 2$.*

$$\begin{array}{c} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} = \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} - \begin{array}{c} | \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} + \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} + \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \text{inv} \qquad \qquad \text{rt} \qquad \qquad \text{rt} \qquad \qquad \text{rt} \qquad \qquad \text{rt} \\ \\ \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} | \\ \cdot \\ \cdot \end{array} + \begin{array}{c} | \\ \cdot \\ \cdot \end{array} \begin{array}{c} | \\ \cdot \\ \cdot \end{array} - \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} \\ \text{rt} \qquad \qquad \text{rt} \qquad \qquad \text{rt} \\ \\ \begin{array}{c} | \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} = - \begin{array}{c} | \\ \cdot \\ \cdot \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array} + \begin{array}{c} | \\ \cdot \\ \cdot \end{array} \begin{array}{c} | \\ \cdot \\ \cdot \end{array} \\ \text{inv} \qquad \qquad \text{rt} \qquad \qquad \text{rt} \end{array}$$

FIGURE 4. The relations of two parametrization for $\text{Pin}(2n+1)$

In figure 4, if we change simultaneously the signatures of the diagrams which have exactly one vertical edge (the edge from the upper row to the lower row), we have the result for $\text{Pin}(2n)$.

If $p > n$, we have $\Delta_{(\epsilon)} \otimes \bigwedge^p V \cong \Delta_{(\epsilon)} \otimes \bigwedge^{N-p} V \otimes \det \cong \Delta_{(-\epsilon)} \otimes \bigwedge^{N-p} V$.

For $\text{Pin}(2n+1)$, the isomorphism $A : \Delta_{\epsilon} \otimes \det \cong \Delta_{-\epsilon}$ is given by the map $A[\mathbf{I}] = [\mathbf{I}]$ for any \mathbf{I} . Here we obey the convention in Remark 3.9. Then we have the $\text{Pin}(2n+1)$ isomorphism $A \otimes r_p : \Delta_{(\epsilon)} \otimes \bigwedge^p V \longrightarrow \Delta_{(\epsilon)} \otimes \bigwedge^{2n+1-p} V$. Here r_p is defined in Lemma 3.13.

For $\text{Pin}(2n)$, let A be the associator in Lemma 3.10, the isomorphism from $\Delta \otimes \det$ to Δ . Then we have the isomorphism $A \otimes r_p : \Delta \otimes \bigwedge^p V \longrightarrow \Delta \otimes \bigwedge^{2n-p} V$.

As before, we define a homomorphism $r_{[1,2n-p]}^{[1,p]}$ by the composition of the alternating operator $\text{Alt}_{[1,p]}$ and r_p and the immersion of the alternating tensor to the positions $[1, 2n-p]$.

Theorem 7.6. *Let $p > n$ and $p + q \leq N$. As a homomorphism from $\Delta_{(\epsilon_1)} \otimes \otimes^p V$ to $\Delta_{(\epsilon_2)} \otimes \otimes^q V$ (we assume $(-1)^{p+q} = \epsilon_1 \epsilon_2$ for $Pin(2n+1)$.), we have*

(7.6.1)

$$\psi_{[1,q]}^{[1,p]} = \epsilon \sum_{\substack{\sigma \in \mathfrak{S}_{N-p} \\ \tau \in \mathfrak{S}_q}} \epsilon(\sigma) \epsilon(\tau) \frac{\binom{\sigma([1,q])}{\tau([1,q])}}{q!} \frac{\text{pr}_{\{\sigma([q+1, N-p])\}}}{(N-p-q)!} (A \otimes r_{[1, N-p]}^{[1,p]}).$$

Here $\epsilon = 1$ for $Pin(2n+1)$ and $\epsilon = (-1)^{q+n}$ for $Pin(2n)$.

Let $q > n$ and $p+q \leq N$. As homomorphisms from $\Delta_{(\epsilon_1)} \otimes \otimes^p V$ to $\Delta_{(\epsilon_2)} \otimes \otimes^q V$ (we assume $(-1)^{p+q} = \epsilon_1 \epsilon_2$ for $Pin(2n+1)$.) is given by

(7.6.2)

$$\psi_{[1,q]}^{[1,p]} = \epsilon \frac{q!}{(N-q)!} (A \otimes r_{[1,q]}^{[1, N-q]}) \sum_{\substack{\sigma \in \mathfrak{S}_p \\ \tau \in \mathfrak{S}_{N-q}}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{inj}_{\{\tau([p+1, N-q])\}}}{(N-p-q)!} \frac{\binom{\sigma([1,p])}{\tau([1,p])}}{p!}.$$

Here $\epsilon = 1$ for $Pin(2n+1)$ and $\epsilon = (-1)^{p+n}$ for $Pin(2n)$.

Proof. We prove the formula (7.6.1) for $Pin(2n+1)$. Let $p > n$ and $q \leq n$. Since $q \leq 2n+1-p$ and Corollary 7.4, we can put

$$\psi_{[1,q]}^{[1,p]} = \sum_{i=0}^q d_i i \phi_{[1,q]}^{[1, 2n+1-p]} (A \otimes r_{[1, 2n-p]}^{[1,p]}).$$

As before, we find a sequence of pairs $\{x_i, y_i\}$ and apply both sides of the above to x_i 's and compare the coefficients of y_i 's and determine the coefficients d_i from $i = q$ to $i = 0$ inductively.

We take cases for the parities of p and q .

Case 1 $p \equiv 1 \pmod{2}$ and $q \equiv 0 \pmod{2}$.

We put $2n+1-p = 2r$ and $q = 2s$. We have $r \geq s$.

We determine the coefficients d_q first.

Case 1- $2s$. $d_{2s} = 1$.

We take $y_{2s} = [\emptyset] \otimes \{[1, 2s]\}$, $x_{2s} = [\emptyset] \otimes \{[1, 2s], [r+s+1, n], 0', \overline{[n, r+s+1]}\}$.

If $i < 2s$, the term i never contributes to the coefficient since the image of pr is 0 because of the weight argument. And the direct calculation tells us that $d_{2s} = 1$.

Case 1- $2s-1$. $d_{2s-1} = 0$.

We take $y_{2s-1} = [\emptyset] \otimes \{[1, 2s-1], 0'\}$, $x_{2s-1} = [\emptyset] \otimes \{[1, 2s-1], [r+s, n], \overline{[n, r+s]}\}$.

Only the terms $2s$ and $2s-1$ contribute to the coefficient because of the weight argument. And the direct calculation tells us that $d_{2s-1} = 0$.

We assume that $d_{2s} = 1$, $d_{2s-1} = \dots = d_{2s-2i+1} = 0$ and we show that $d_{2s-2i} = 0$.

Case 1- $2s - 2i$. $d_{2s-2i} = 0$.

We take $y_{2s-2i} = [\emptyset] \otimes \{[1, 2s - i], \overline{[2s - i, 2s - 2i + 1]}\}, x_{2s-2i} = [\emptyset] \otimes \{[1, 2s - 2i], [r + s + 1 - i, n], 0', \overline{[n, r + s + 1 - i]}\}$. From the induction hypothesis and the weight argument, only the terms $2s$ and $2s - 2i$ contribute to the coefficient and the direct calculation tells us that $d_{2s-2i} = 0$.

We assume that $d_{2s} = 1, d_{2s-1} = \dots = d_{2s-2i} = 0$ and we show that $d_{2s-2i-1} = 0$.

Case 1- $2s - 2i - 1$. $d_{2s-2i-1} = 0$ ($i \geq 1$).

We take $y_{2s-2i-1} = [\emptyset] \otimes \{[1, 2s - i - 1], 0', \overline{[2s - i - 1, 2s - 2i]}\}, x_{2s-2i-1} = [\emptyset] \otimes \{[1, 2s - 2i - 1], [r + s - i, n], \overline{[n, r + s - i]}\}$.

For the remaining cases of parities of p and q , the proof goes almost in a similar way. So we omit it. Summarizing all the cases, we have $d_q = 1$ and $d_i = 0$ if $i < q$ in the formula (7.6).

Next we prove the formula (7.6) for $Pin(2n + 1)$.

Let $p \leq n$ and $q > n$.

We can put

$$\psi_{[1,q]}^{[1,p]} = \sum_{i=0}^p d_i \frac{q!}{(2n + 1 - q)!} (A \otimes r_{[1,q]}^{[1,2n+1-q]}) \phi_{[1,2n+1-q]}^{[1,p]}$$

We determine the coefficients d_i just as in the above.

We take cases for the parities of p and q .

Case 1 $p \equiv 0 \pmod{2}$ and $q \equiv 1 \pmod{2}$.

We put $p = 2r$ and $2n + 1 - q = 2s$. We have $s \geq r$.

We determine the coefficients d_p first.

Case 1- $2r$. $d_{2r} = 1$.

We take $y_{2r} = [\emptyset] \otimes \{[1, 2r], [r + s + 1, n], 0', \overline{[n, r + s + 1]}\}, x_{2r} = [\emptyset] \otimes \{[1, 2r]\}$. We have $d_{2r} = 1$.

Case 1- $2r - 1$. $d_{2r-1} = 0$.

We take $y_{2r-1} = [\emptyset] \otimes \{[1, 2r - 1], [r + s, n], \overline{[n, r + s]}\}, x_{2r-1} = [\emptyset] \otimes \{[1, 2r - 1], 0'\}$. We have $d_{2r-1} = 0$.

We assume that $d_{2r} = 1, d_{2r-1} = \dots = d_{2r-2i+1} = 0$ and we show that $d_{2r-2i} = 0$.

Case 1- $2r - 2i$. $d_{2r-2i} = 0$. ($i \geq 1$).

We take $y_{2r-2i} = [\emptyset] \otimes \{[1, 2r - 2i], [r + s - i + 1, n], 0', \overline{[n, r + s - i + 1]}\}, x_{2r-2i} = [\emptyset] \otimes \{[1, 2r - i], \overline{[2r - i, 2r - 2i + 1]}\}$. We have $d_{2r-2i} = 0$.

We assume that $d_{2r} = 1, d_{2r-1} = \dots = d_{2r-2i} = 0$ and we show that $d_{2r-2i-1} = 0$.

Case 1- $2r - 2i - 1$. $d_{2r-2i-1} = 0$. ($i \geq 1$).

We take $y_{2r-2i-1} = [\emptyset] \otimes \{[1, 2r - 2i - 1], [r + s - i, n], \overline{[n, r + s - i]}\}, x_{2r-2i-1} = [\emptyset] \otimes \{[1, 2r - i - 1], 0', \overline{[2r - i - 1, 2r - 2i]}\}$. We have $d_{2r-2i-1} = 0$.

For the remaining cases, proof goes almost in a similar way. So we omit it and summarizing all the cases, we have $d_p = 1$ and $d_i = 0$ if $i < p$ in the formula (7.6).

For $Pin(2n)$, the proof is almost similar to the above and we omit it. \square

8. RELATIONS BETWEEN $Pin(N)$ -EQUIVARIANT HOMOMORPHISMS

In this section we give the relations between the compositions of pr and inj and the contractions and the immersion of the invariant forms, from which we can deduce the multiplication rules of the generalized Brauer diagrams.

We recall some notations. Let $C_{\{i,j\}}$ be the contraction operator of the i -th and j -th tensor components (with respect to S) and let $\text{id}_V^{\{i,j\}}$ be the linear immersion of the invariant element $\text{id}_V = \sum_{i=1}^n (u_i \otimes u_{\bar{i}} + u_{\bar{i}} \otimes u_i)(+u_{0'} \otimes u_{0'})$ to the i -th and the j -th position.

As in Corollary 7.4, we define the equivariant homomorphism from $\Delta \otimes \otimes^p$ (here we consider the tensor $\otimes^p V$ sits in the positions $\{[r+1, r+p]\}$) to $\Delta \otimes \otimes^q$ (here we consider the tensor $\otimes^q V$ sits in the positions $\{[s+1, s+q]\}$) by

$$\phi_{[s+1, s+q]}^{[r+1, p+r]} = \sum_{\substack{\sigma \in \mathfrak{S}_q[s+1, s+q] \\ \tau \in \mathfrak{S}_p[r+1, r+p]}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{inj}_{\{\sigma([s+1+1, s+q])\}}}{(q-i)!} \frac{\left(\tau([r+1, r+i]) \right)}{\sigma([s+1, s+i])} \frac{\text{pr}_{\{\tau([r+i+1, r+p])\}}}{(p-i)!}.$$

Here $\mathfrak{S}_q[s+1, s+q]$ and $\mathfrak{S}_p[r+1, r+p]$ denote the symmetric groups of degree q and p acting on the sets $\{s+1, s+2, \dots, s+q\}$ and $\{r+1, r+2, \dots, r+p\}$ respectively.

Then from Corollary 7.4, $\phi_{[s+1, s+q]}^{[r+1, p+r]}$'s ($0 \leq i \leq \min(p, q)$) become a base of $\text{Hom}_{Pin(N)}(\Delta \otimes \wedge^p V, \Delta \otimes \wedge^q V)$.

Theorem 8.1. *The following formulas hold for pr , inj , $C_{\{i,j\}}$ and $\text{id}_V^{\{i,j\}}$.*

- (i) *If $p \leq n$, as the homomorphisms from $\Delta \otimes \otimes^p V$ to Δ , (here we consider the tensor $\otimes^p V$ sits in the positions $\{[q+1, p+q]\}$), we have*

$$(8.1.1) \quad \text{pr}_{\{[1, q+p]\}} \circ \text{inj}_{\{[1, q]\}} = (N-p)_q \in \text{pr}_{\{[q+1, q+p]\}}.$$

Here $\epsilon = 1$ for $Pin(2n+1)$ and $\epsilon = (-1)^{(p+q)q}$ for $Pin(2n)$.

$(N-p)_q$ denotes the lower factorial, that is, for any x and non-negative integer i , we define $(x)_i = x(x-1)(x-2) \cdots (x-(i-1))$.

The above also holds for $p = 0$ and in this case we regard pr in the right-hand side as the identity map of Δ .

- (ii) If $p \leq n$, as the homomorphisms from Δ to $\Delta \otimes \otimes^p V$ (here we consider the tensor $\otimes^p V$ sits in the positions $\{[q+1, p+q]\}$), we have

$$(8.1.2) \quad \text{pr}_{\{[1, q]\}} \circ \text{inj}_{\{[1, q+p]\}} = (N-p)_q \in \text{inj}_{\{[q+1, q+p]\}}.$$

Here $\epsilon = 1$ for $\text{Pin}(2n+1)$ and $\epsilon = (-1)^{(p+q)q}$ for $\text{Pin}(2n)$.

- (iii) If $p \leq n$ and $q \leq n$, as the homomorphisms from Δ to $\Delta \otimes \otimes^{p+q} V$, we have

$$(8.1.3)$$

$$\text{inj}_{\{[1, q]\}} \circ \text{inj}_{\{[q+1, q+p]\}} = \sum_{i=0}^{\min(p, q)} \epsilon(i) \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p[q+1, q+p]}} \epsilon(\sigma) \epsilon(\tau) \frac{\prod_{u=1}^i \text{id}_V \{\sigma(u), \tau(q+u)\}}{i!} \frac{\text{inj}_{\{\sigma([i+1, q]), \tau([q+i+1, q+p])\}}}{(q-i)!(p-i)!}.$$

Here $\epsilon(i) = (-1)^{qi + \binom{i+1}{2}}$ for $\text{Pin}(2n+1)$ and $\epsilon(i) = (-1)^{qi + \binom{i+1}{2} + pq}$ for $\text{Pin}(2n)$.

- (iv) If $p \leq n$ and $q \leq n$, as the homomorphisms from $\Delta \otimes \otimes^{p+q} V$ to Δ , we have

$$(8.1.4)$$

$$\text{pr}_{\{[1, q]\}} \circ \text{pr}_{\{[q+1, q+p]\}} = \sum_{i=0}^{\min(p, q)} \epsilon(i) \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p[q+1, q+p]}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{pr}_{\{\sigma([i+1, q]), \tau([q+i+1, q+p])\}}}{(q-i)!(p-i)!} \frac{\prod_{u=1}^i C_{\{\sigma(u), \tau(q+u)\}}}{i!}.$$

Here $\epsilon(i) = (-1)^{qp + qi + \binom{i}{2}}$ for $\text{Pin}(2n+1)$ and $\epsilon(i) = (-1)^{qi + \binom{i}{2}}$ for $\text{Pin}(2n)$.

- (v) If $p \geq t$ and $p-t \leq n$ and $q \leq n$, as the homomorphisms from $\Delta \otimes \otimes^{p-t} V$ to $\Delta \otimes \otimes^q V$, (here we consider the tensor $\otimes^{p-t} V$ sits in the positions $\{[q+t+1, q+p]\}$.) we have

$$(8.1.5)$$

$$\text{pr}_{\{[q+1, q+p]\}} \circ \text{inj}_{\{[1, q+t]\}} = \sum_{i=0}^{\min(p-t, q)} \epsilon(i) \left(\sum_{u=0}^i \binom{i}{u} (N-p-q+t+i-u)_t \right) i \phi_{[1, q]}^{[q+t+1, q+p]}$$

Here $\epsilon(i) = (-1)^{(q-i)(p-i)+it}$ for $\text{Pin}(2n+1)$ and $\epsilon(i) = (-1)^{pq+t(p+q)}$ for $\text{Pin}(2n)$ and $\binom{i}{u}$ in the parentheses denotes the ordinary binomial coefficient. If $t = 0$, $(N-p-q+0+i-u)_0 = 1$ and the sum in the parentheses is equal to 2^i .

- (vi) If $p \leq n$ and $q \leq n$, as the homomorphisms from $\Delta \otimes \otimes^q V$ to $\Delta \otimes \otimes^p V$, (here we consider the tensor $\otimes^q V$ sits in the positions $\{[p+q+1, p+2q]\}$.) we have

$$(8.1.6) \quad \prod_{i=1}^q C_{\{i, p+q+i\}} \text{inj}_{\{[1, q+p]\}} = \sum_{i=0}^{\min(p, q)} \epsilon(i) i \phi_{[q+1, q+p]}^{[p+q+1, p+2q]}.$$

Here $\epsilon(i) = (-1)^{pq + \binom{q}{2} + i(p+q-1)}$ for $Pin(2n+1)$ and $\epsilon(i) = (-1)^{\binom{q}{2}}$ for $Pin(2n)$.

- (vii) If $p \leq n$ and $q \leq n$, as the homomorphisms from $\Delta \otimes \otimes^q V$ to $\Delta \otimes \otimes^p V$, (here we consider the tensor $\otimes^q V$ sits in the positions $[1, q]$ and $\otimes^p V$ sits in the positions $\{[p+q+1, 2p+q]\}$.) we have

$$(8.1.7) \quad \text{pr}_{\{[1, q+p]\}} \prod_{i=1}^p \text{id}_{V_{\{q+i, p+q+i\}}} = \sum_{i=0}^{\min(p, q)} \epsilon(i) i \phi_{[p+q+1, q+2p]}^{[1, q]}.$$

Here $\epsilon(i) = (-1)^{\binom{p}{2} + i(p+q+1)}$ for $Pin(2n+1)$ and $\epsilon(i) = (-1)^{\binom{p}{2} + pq}$ for $Pin(2n)$.

Remark 8.2. If we change N in the above formulas into an indeterminate X simultaneously, we can define the ‘generic’ centralizer algebra of \mathbf{CP}_k just as in the ordinary Brauer algebras.

Proof. We prove the formula (8.1.1). If we note that $\text{Alt}_{[1, p+q]}(\text{Alt}_{[1, q]} \times \text{Alt}_{[q+1, p+q]}) = \text{Alt}_{[1, p+q]}(\text{Alt}_{[1, q]}) = \text{Alt}_{[1, p+q]}$, both sides factor through the homomorphisms from $\Delta \otimes \wedge^p V$ to Δ . Because of $\dim \text{Hom}_{Pin(N)}(\Delta \otimes \wedge^p V, \Delta) = 1$, they only differ in some scalar if they are non-zero. We compare the coefficients of $[\emptyset]$ of the images of $[[1, p]] \otimes \{[1, p]\}$ under both sides of the formula (8.1.1). Direct calculation tells us that the scalar is given by $(2n+1-p)_q$ for $Pin(2n+1)$ and the scalar is given by $(-1)^{(p+q)q}(2n-p)_q$ for $Pin(2n)$.

We prove the formula (8.1.2).

Both sides of the above can be considered as the homomorphisms from Δ to $\Delta \otimes \wedge^p V$. As before, they only differ in some scalar. We compare the coefficients of $[[1, p]] \otimes \{[1, p]\}$ of the images of $[\emptyset]$ under both sides of the formula (8.1.2).

Direct calculation tells us that the scalar is given by $(2n+1-p)_q$ for $Pin(2n+1)$ and $(-1)^{(p+q)q}(2n+1-p)_q$ for $Pin(2n)$.

Next we prove the formula (8.1.3).

From Theorem 2.4 and Theorem 2.2, Δ occurs in the representation $\Delta \otimes \mu_{O(N)}$ if and only if $\mu_{O(N)} = (1^i)$ and then the multiplicity is one. Also the character theory tells us that in the representation $\wedge^q V \otimes \wedge^p V$, $\wedge^{p+q-2i} V$ ($0 \leq i \leq \min(p, q)$) occurs exactly once and the other exterior products never appears.

So the left-hand side in (8.1.3) can be considered as the homomorphism in $\text{Hom}_{Pin(N)}(\Delta, \Delta \otimes \wedge^q V \otimes \wedge^p V) \cong \bigoplus_{i=0}^{\min(p, q)} \text{Hom}_{Pin(2n)}(\Delta, \Delta \otimes \wedge^{p+q-2i} V)$.

A $Pin(2n)$ equivariant embedding from $\wedge^{p+q-2i} V$ to $\wedge^q V \otimes \wedge^p V$ is given as follows. For any subset $U \subseteq \{[1, n], (0'), \overline{[n, 1]}\}$ of size $p+q-2i$,

we define

$$< \mathbf{U} > \longrightarrow \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p[q]}} \epsilon(\sigma) \epsilon(\tau) \prod_{u=1}^i \text{id}_V \{ \sigma(\mathbf{u}), \tau(\mathbf{q}+\mathbf{u}) \} \times < \mathbf{U} >_{\{ \sigma([\mathbf{i}+1, \mathbf{q}]), \tau([\mathbf{q}+\mathbf{i}+1, \mathbf{q}+\mathbf{p}]) \}}.$$

Here the indices in \mathbf{U} are in the increasing order.

So in the formula (8.1.3), the i term in the right-hand side is the homomorphism in the direct summand $\text{Hom}_{\text{Pin}(N)}(\Delta, \Delta \otimes \bigwedge^{p+q-2i} V)$.

So we can put

$$\begin{aligned} & \text{inj}_{\{[1, \mathbf{q}]\}} \circ \text{inj}_{\{[\mathbf{q}+1, \mathbf{q}+\mathbf{p}]\}} = \\ & \sum_{i=0}^{\min(p, q)} d_i \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p[q+1, q+p]}} \epsilon(\sigma) \epsilon(\tau) \frac{\prod_{u=1}^i \text{id}_V \{ \sigma(\mathbf{u}), \tau(\mathbf{q}+\mathbf{u}) \}}{i!} \frac{\text{inj}_{\{ \sigma([\mathbf{i}+1, \mathbf{q}]), \tau([\mathbf{q}+\mathbf{i}+1, \mathbf{q}+\mathbf{p}]) \}}}{(q-i)!(p-i)!}. \end{aligned}$$

As before, we find a sequence of pairs $\{x_i, y_i\}$ and apply both sides of the above to x_i 's and compare the coefficients of y_i 's and determine the coefficients d_i inductively.

We prove the formula (8.1.3) for $\text{Pin}(2n+1)$.

We take the cases for the parities of p and q .

Case 1. $p = 2r$ and $q = 2s$.

We note $r + s \leq n$ because of $p \leq n$ and $q \leq n$.

Case 1-0. The determination of d_0 .

We take $y_0 = [\emptyset] \otimes \{ [1, \mathbf{s}], [\mathbf{s}, 1], [\mathbf{s} + 1, \mathbf{r} + \mathbf{s}], [\mathbf{r} + \mathbf{s}, \mathbf{s} + 1] \}$, $x_0 = [\emptyset]$.

In this case $d_0 = 1$.

Case 1-1. The determination of d_1 .

We take $y_1 = [\mathbf{s} + 1] \otimes \{ [\overline{1}, [2, \mathbf{s}], 0', [\mathbf{s}, 2], 1, [\mathbf{s} + 1, \mathbf{r} + \mathbf{s}], [\mathbf{r} + \mathbf{s}, \mathbf{s} + 2] \}$, $x_1 = [\emptyset]$. In this case $d_1 = -1$.

We prove $d_i = (-1)^{\binom{i+1}{2}}$ by induction on i in this case.

Case 1-2u. The determination of d_{2u} .

We assume that the above holds for $i = 1, 2, \dots, 2u - 1$ and prove the case of $i = 2u$. We take

$$\begin{aligned} y_{2u} &= [\emptyset] \otimes \\ & \{ [\overline{2u, 1}], [2u + 1, \mathbf{s} + \mathbf{u}], [\overline{\mathbf{s} + \mathbf{u}, 2u + 1}], [1, 2u], [\mathbf{s} + \mathbf{u} + 1, \mathbf{r} + \mathbf{s}], [\overline{\mathbf{r} + \mathbf{s}, \mathbf{s} + \mathbf{u} + 1}] \} \end{aligned}$$

and $x_{2u} = [\emptyset]$. Then in the right-hand side, only the terms from 0 to $2u$ contribute to the coefficient and we have $d_{2u} = (-1)^{\binom{2u+1}{2}}$.

Case 1-2u+1. The determination of d_{2u+1} .

We take

$$\begin{aligned} y_{2u+1} &= [\mathbf{s} + \mathbf{u} + 1] \otimes \\ & \{ [\overline{1, 2u + 1}], [2u + 2, \mathbf{s} + \mathbf{u}], 0', [\overline{\mathbf{s} + \mathbf{u}, 2u + 2}], [1, 2u + 1], [\mathbf{s} + \mathbf{u} + 1, \mathbf{r} + \mathbf{s}], [\overline{\mathbf{r} + \mathbf{s}, \mathbf{s} + \mathbf{u} + 2}] \} \end{aligned}$$

and $x_{2u+1} = [\emptyset]$.

For the remaining cases for $Pin(2n+1)$ and for $Pin(2n+1)$, the proof goes in a similar way and so we omit it.

Next we prove the formula (8.1.4). As in the proof of the formula (8.1.3), the left-hand side can be considered as the homomorphism in $\text{Hom}_{Pin(N)}(\Delta \otimes \wedge^q V \otimes \wedge^p V, \Delta) \cong \bigoplus_{i=0}^{\min(p,q)} \text{Hom}_{Pin(N)}(\Delta \otimes \wedge^{p+q-2i} V, \Delta)$. A $Pin(N)$ -equivariant embedding from $\wedge^q V \otimes \wedge^p V$ to $\wedge^{p+q-2i} V$ is given as follows. For any subsets $Q, P \subseteq \{[1, n], (0'), \overline{[n, 1]}\}$ of size q, p respectively, we define

$$\begin{aligned} \langle Q \rangle \otimes \langle P \rangle &\longrightarrow \text{Alt}_{\{[i+1, q], [q+i+1, q+p]\}} \prod_{u=1}^i C_{\{u, q+u\}} \text{Alt}_{\{[1, q]\}}(\otimes Q) \otimes \text{Alt}_{\{[q+1, q+p]\}}(\otimes P) \\ &= \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p[q+1, q+p]}} \frac{1}{q!p!} \epsilon(\sigma) \epsilon(\tau) \text{Alt}_{\{\sigma([i+1, q]), \tau([q+i+1, q+p])\}} \prod_{u=1}^i C_{\{\sigma(u), \tau(q+u)\}}(\otimes Q) \otimes (\otimes P) \end{aligned}$$

Here if $Q = \{\ell_1 \ell_2, \dots, \ell_q\}$, $(\otimes Q)$ denotes the ordinary tensor product (not exterior one) $\otimes \{\ell_1 \ell_2, \dots, \ell_q\}$.

So in the formula (8.1.4), the i term in the right-hand side corresponds to the homomorphism in the direct summand $\text{Hom}_{Pin(N)}(\Delta \otimes \wedge^{p+q-2i} V, \Delta)$.

So we can put

$$\begin{aligned} \text{pr}_{\{[1, q]\}} \circ \text{pr}_{\{[q+1, q+p]\}} &= \\ \sum_{i=0}^{\min(p,q)} d_i \sum_{\substack{\sigma \in \mathfrak{S}_q \\ \tau \in \mathfrak{S}_p[q+1, q+p]}} \epsilon(\sigma) \epsilon(\tau) \frac{\text{pr}_{\{\sigma([i+1, q]), \tau([q+i+1, q+p])\}}}{(q-i)!(p-i)!} \frac{\prod_{u=1}^i C_{\{\sigma(u), \tau(q+u)\}}}{i!}. \end{aligned}$$

The rest of the proof is almost similar to (8.1.3) and we omit it.

Next we prove the formula (8.1.5).

Both sides of the above can be considered as the homomorphisms from $\Delta \otimes \wedge^{p-t} V$ to $\Delta \otimes \wedge^q V$.

So we can put (see Corollary 7.4.)

$$\text{pr}_{\{[q+1, q+p]\}} \circ \text{inj}_{\{[1, q+t]\}} = \sum_{i=0}^{\min(p-t, q)} d_i {}^i\phi_{[1, q]}^{[q+t+1, q+p]}.$$

We show $d_i = (-1)^{(q-i)(p-i)+it} (\sum_{u=0}^i \binom{i}{u} (2n+1-p-q+t+i-u)_t)$ for $Pin(2n+1)$ and $d_i = (-1)^{pq+t(p+q)} (\sum_{u=0}^i \binom{i}{u} (2n+1-p-q+t+i-u)_t)$ for $Pin(2n)$.

As before, we find a sequence of pairs $\{x_i, y_i\}$ and apply both sides of the above to x_i 's and compare the coefficients of y_i 's and determine the coefficients d_i inductively.

We prove the formula (8.1.5) for $Pin(2n+1)$.

We take the cases for the parities of $p-t$ and q .

Case 1. $p - t = 2r$ and $q = 2s$.

We note $r + s \leq n$.

Case 1-0. The determination of d_0 .

We take $y_0 = [\emptyset] \otimes \{[1, \mathbf{s}], [\overline{\mathbf{s}}, 1]\}$, $x_0 = [\emptyset] \otimes \{[\mathbf{n} - \mathbf{r} + 1, \mathbf{n}], [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + 1]\}$.

In the left-hand side, from the definition 6.2, we have

$$(8.1) \quad \begin{aligned} \text{inj}_{q+t}([\emptyset]) &= \sum_{\substack{J \subseteq [1, n] \\ W \subseteq ([1, n] - J) \\ j+2w=q+t}} (-1)^w 2^{j/2} [J] \otimes (q+t)! \langle J, W, \overline{W} \rangle \\ &+ \sum_{\substack{J \subseteq [1, n] \\ W \subseteq ([1, n] - J) \\ j+2w+1=q+t}} 2^{j/2} [J] \otimes (q+t)! \langle J, W, O', \overline{W} \rangle. \end{aligned}$$

In the above only the W 's which include the set $\{[1, s]\}$ contribute to the coefficient and for them, we put $W_1 = W - \{[1, s]\}$. The exterior products can be rewritten as $(q+t)! \langle J, W, \overline{W} \rangle = \otimes \{[1, \mathbf{s}], [\overline{\mathbf{s}}, 1]\} \otimes t! \langle J, W_1, \overline{W_1} \rangle + \text{the other terms}$ and $(q+t)! \langle J, W, O', \overline{W} \rangle = (-1)^s \otimes \{[1, \mathbf{s}], [\overline{\mathbf{s}}, 1]\} \otimes t! \langle J, W_1, O', \overline{W_1} \rangle + \text{the other terms}$.

We tensor $\otimes \{[\mathbf{n} - \mathbf{r} + 1, \mathbf{n}], [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + 1]\}$ after the exterior products in the above and alternate them. If $(J \cup W_1) \cap \{[n-r+1, n]\} \neq \emptyset$, then the resulting exterior products are 0. So we can assume $(J \cup W_1) \cap \{[n-r+1, n]\} = \emptyset$. Then we have $t! \langle J, W_1, [\mathbf{n} - \mathbf{r} + 1, \mathbf{n}], [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + 1], \overline{W_1} \rangle$ and $t!(-1)^r \langle J, W_1, [\mathbf{n} - \mathbf{r} + 1, \mathbf{n}], O', [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + 1], \overline{W_1} \rangle$.

So the coefficient in the left-hand side is equal to

$$\begin{aligned} &\sum_{\substack{\{[s+1, n-r]\} \supseteq W_1 \sqcup J \text{ (disjoint)} \\ j+2w_1=t}} (-1)^{w_1+s} 2^{j/2} t! 2^{j/2} (-1)^{w_1+r} + \sum_{\substack{\{[s+1, n-r]\} \supseteq W_1 \sqcup J \text{ (disjoint)} \\ j+2w_1+1=t}} 2^{j/2} t! (-1)^{s+r} 2^{j/2} \\ &= (-1)^{r+s} t! ((1 + 2x + x^2)^{n-r-s} [t] + (1 + 2x + x^2)^{n-r-s} [t-1]) \\ &= (-1)^{r+s} t! (1+x)^{2n+1-p+t-q} [t] = (-1)^{r+s} (2n+1-p+t-q)_t. \end{aligned}$$

Here for the formal power series $f(x)$, by $f(x)[t]$ we denote the coefficient of the power x^t in $f(x)$.

In the right-hand side, only the $i = 0$ term contributes to the coefficient and it is given by $(-1)^{r+s} d_0$ and we have $d_0 = (2n+1-p+t-q)_t$.

Case 1-1. The determination of d_1 .

We take $y_1 = [\mathbf{s} + 1] \otimes \{[1, \mathbf{s} + 1], [\overline{\mathbf{s}}, 2]\}$, $x_1 = [\emptyset] \otimes \{1, [\mathbf{n} - \mathbf{r} + 2, \mathbf{n}], O', [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + 2]\}$.

We calculate the left-hand side.

In the equation (8.1), only the first sum contributes to the coefficient and moreover in the first sum the W 's and J 's such that $W \supseteq \{2, \dots, s\}$ and $J \cup W \supseteq \{1, s+1\}$ contribute to the coefficient. There are four subcases, 1) $W \supseteq \{1, s+1\}$ (if $t \geq 2$) 2) $J \supseteq \{1\}$, $W \supseteq \{s+1\}$ 3) $J \supseteq \{s+1\}$, $W \supseteq \{1\}$ 4) $J \supseteq \{1, s+1\}$. We put $J_1 = J - J \cap \{1, s+1\}$ and $W_1 = W - W \cap [s+1]$.

In the subcase 1), we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2+2w_1=t}} -2^{j_1+1/2}t! = -\sqrt{2}t!(1+x)^{2n-2r-2s}[t-2].$$

In the subcase 2) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+1+2w_1=t}} -2^{j_1+1+1/2}t! = -2\sqrt{2}t!(1+x)^{2n-2r-2s}[t-1].$$

In the subcase 3) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+1+2w_1=t}} -2^{j_1+1/2}t! = -\sqrt{2}t!(1+x)^{2n-2r-2s}[t-1].$$

In the subcase 4) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2w_1=t}} -2^{j_1+1+1/2}t! = -2\sqrt{2}t!(1+x)^{2n-2r-2s}[t].$$

So the total coefficient of the left-hand side is $-\sqrt{2}t!((1+x)^{2n+1-2r-2s}[t] + (1+x)^{2n+1-2r-2s+1}[t] = -\sqrt{2}((2n+1-p-q+t)_t + (2n+1-p-q+t+1)_t)$. If $t = 1$, the subcase 1) does not occur, but the result also holds in this case since $(1+x)^{2n-2r-2s}[-1] = 0$.

In the right-hand side, only the $i = 1$ term contributes to the coefficient and it is given by $\sqrt{2}d_1$ and we have $d_1 = -((2n+1-p-q+t)_t + (2n+1-p-q+t+1)_t)$.

Case 1-2. The determination of d_2 .

We take $y_2 = [\emptyset] \otimes \{[1, s+1], [s+1, 3]\}$, $x_2 = [\emptyset] \otimes \{1, 2, [n-r+2, n], [n, n-r+2]\}$.

We calculate the left-hand side.

In the equation (8.1), only the W 's and J 's such that $W \supseteq \{[3, s+1]\}$ and $J \cup W \supseteq \{1, 2\}$ contribute to the coefficient. There are four subcases, 1) $W \supseteq \{1, 2\}$ (if $t \geq 2$) 2) $J \supseteq \{1\}$, $W \supseteq \{2\}$ 3) $J \supseteq \{2\}$, $W \supseteq \{1\}$ 4) $J \supseteq \{1, 2\}$. We put $J_1 = J - J \cap \{1, 2\}$ and $W_1 = W - W \cap [s+1]$.

In the subcase 1) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2+2w_1=t}} (-1)^{s+r}2^{j_1}t! + \sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2+2w_1+1=t}} (-1)^{s+r}2^{j_1}t!.$$

In the subcase 2) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+1+2w_1=t}} (-1)^{s+r}2^{j_1+1}t! + \sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2+2w_1=t}} (-1)^{s+r}2^{j_1+1}t!.$$

In the subcase 3) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+1+2w_1=t}} (-1)^{s+r} 2^{j_1+1} t! + \sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2+2w_1=t}} (-1)^{s+r} 2^{j_1+1} t!.$$

In the subcase 4) we have the coefficient

$$\sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2w_1=t}} (-1)^{s+r} 2^{j_1+2} t! + \sum_{\substack{\{[s+2, n-r+1]\} \supseteq J_1 \sqcup W_1 \text{ (disjoint)} \\ j_1+2w_1+1=t}} (-1)^{s+r} 2^{j_1+2} t!.$$

So the total coefficient in the left-hand side is given as follows.

$$(-1)^{s+r} t! \{ (1+x)^{2n-p-q+t} [t-2] + 2(1+x)^{2n-p-q+t} [t-1] + 2(1+x)^{2n-p-q+t} [t-1] + 4(1+x)^{2n-p-q+t} [t] + (1+x)^{2n-p-q+t} [t-3] + 2(1+x)^{2n-p-q+t} [t-2] + 2(1+x)^{2n-p-q+t} [t-2] + 4(1+x)^{2n-p-q+t} [t-1] \}.$$

Here we prove the following lemma.

Lemma 8.3. *For any indeterminate X and for any non-negative integers t and i , we have*

$$\sum_{u=0}^i \binom{i}{u} 2^u \binom{X}{t-i+u} = \sum_{u=0}^i \binom{i}{u} \binom{X+i-u}{t}.$$

Proof. This follows easily from the formula

$$\binom{X+i}{t} = \sum_{u=0}^i \binom{X}{t-u} \binom{i}{u}.$$

□

From this lemma, the coefficient of the left-hand side is $(-1)^{s+r} \{ (2n+3-p-q+t)_t + 2(2n+2-p-q+t)_t + (2n+1-p-q+t)_t \}$.

In the right-hand side, only the $i=2$ term contributes to the coefficient and it is given by $(-1)^{s-1+r-1} d_2$. So $d_2 = (2n+3-p-q+t)_t + 2(2n+2-p-q+t)_t + (2n+1-p-q+t)_t$.

Case 1-2*i* + 1. The determination of d_{2i+1} .

We take $y_{2i+1} = [\mathbf{s} + \mathbf{i} + 1] \otimes \{ [1, \mathbf{s} + \mathbf{i} + 1], [\overline{\mathbf{s} + \mathbf{i}}, 2\mathbf{i} + 2] \}$, $x_{2i+1} = [\emptyset] \otimes \{ [1, 2\mathbf{i} + 1], [\mathbf{n} - \mathbf{r} + \mathbf{i} + 2, \mathbf{n}], \mathbf{0}', [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + \mathbf{i} + 2] \}$.

In the equation (8.1), only the W 's and J 's such that $W \supseteq \{ [2i+2, s+i] \}$ and $J \cup W \supseteq \{ [1, 2i+1], s+i+1 \}$ contribute to the coefficient.

We put $J_2 = J \cap [1, 2i+1]$ and $J_1 = J \cap \{ [s+i+2, n-r+i+1] \}$ and $W_2 = W \cap [1, 2i+1]$ and $W_1 = W \cap \{ [s+i+2, n-r+i+1] \}$. We fix a subset $J_2 \subseteq [1, 2i+1]$ with $|J_2| = v$ and take cases for $s+i+1 \in J$ and $s+i+1 \notin J$.

Subcase 1-2*i* + 1-1. $s+i+1 \notin J$.

In this subcase we have the coefficient

$$\begin{aligned}
& -t! \sum_{\substack{\{[s+i+2, n-r+i+1]\} \supseteq W_1 \sqcup J_1 \text{ (disjoint)} \\ j_1+2w_1+2i+2-v=t}} 2^{j_1+v+1/2} \\
& = -t! 2^{v+1/2} \sum_{j_1+2w_1+2i+2-v=t} \binom{n-r-s}{j_1} \binom{n-r-s-j_1}{w_1} 2^{j_1} \\
& = -t! 2^{v+1/2} (1+2x+x^2)^{n-r-s} [t-2i-2+v].
\end{aligned}$$

Subcase 1-2*i* + 1-2. $s+i+1 \in J$.

In this subcase we have the coefficient

$$\begin{aligned}
& -t! \sum_{\substack{\{[s+i+2, n-r+i+1]\} \supseteq W_1 \sqcup J_1 \text{ (disjoint)} \\ j_1+2w_1+2i+1-v=t}} 2^{j_1+v+1/2} \\
& = -t! 2^{v+1/2} (1+2x+x^2)^{n-r-s} [t-2i-1+v].
\end{aligned}$$

Therefore for a fixed $J_2 \subseteq [2i+1]$, we have the coefficient $-t! 2^{v+1/2} (1+x)^{2n+1-p-q+t} [t-2i-1+v]$ and the total coefficient is

$$\begin{aligned}
& -t! \sqrt{2} \sum_{v=0}^{2i+1} \binom{2n+1-p-q+t}{t-2i-1+v} 2^v \binom{2i+1}{v} \\
& = -t! \sqrt{2} \sum_{v=0}^{2i+1} \binom{2i+1}{v} \binom{2n+1-p-q+t+2i+1-v}{t}.
\end{aligned}$$

In the right-hand side, only the $2i+1$ term contributes to the coefficient and it is given by $\sqrt{2} d_{2i+1}$. So $d_{2i+1} = -\sum_{v=0}^{2i+1} \binom{2i+1}{v} (2n+1-p-q+t+2i+1-v)_t$.

Case 1-2*i*. The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, \mathbf{s} + \mathbf{i}], [\overline{\mathbf{s} + \mathbf{i}}, 2\mathbf{i} + 1]\}$,
 $x_{2i} = [\emptyset] \otimes \{[1, 2\mathbf{i}], [\mathbf{n} - \mathbf{r} + \mathbf{i} + 1, \mathbf{n}], [\overline{\mathbf{n}}, \mathbf{n} - \mathbf{r} + \mathbf{i} + 1]\}$.

In the equation (8.1), only the W 's and J 's such that $W \supseteq \{2i+1, \dots, s+i\}$ and $J \cup W \supseteq \{[1, 2i]\}$ contribute to the coefficient.

We put $J_2 = J \cap [1, 2i]$ and $J_1 = J \cap \{[s+i+1, n-r+i]\}$ and $W_2 = W \cap [1, 2i]$ and $W_1 = W \cap \{[s+i+1, n-r+i]\}$. We fix a subset $J_2 \subseteq [1, 2i]$ with $|J_2| = v$. We take the cases.

Subcase 1-2*i*-1. The terms do not contain $0'$.

In this subcase we have the coefficient

$$\begin{aligned}
& t! \sum_{\substack{\{[s+i+1, n-r+i]\} \supseteq W_1 \sqcup J_1 \text{ (disjoint)} \\ j_1 + 2w_1 + 2i - v = t}} (-1)^{s-i+r-i} 2^{j_1+v} \\
&= (-1)^{s+r} t! 2^v \sum_{j_1 + 2w_1 + 2i - v = t} \binom{n-r-s}{j_1} \binom{n-r-s-j_1}{w_1} 2^{j_1} \\
&= (-1)^{s+r} t! 2^v (1+x)^{2n-p-q+t} [t-2i+v].
\end{aligned}$$

Subcase 1-2i-2. The terms contain $0'$.

In this subcase we have the coefficient

$$\begin{aligned}
& t! 2^v (-1)^{s+r} \sum_{\substack{\{[s+i+1, n-r+i]\} \supseteq W_1 \sqcup J_1 \text{ (disjoint)} \\ j_1 + 2w_1 + 2i + 1 - v = t}} 2^{j_1} \\
&= t! 2^v (-1)^{s+r} (1+2x+x^2)^{n-r-s} [t-2i-1+v].
\end{aligned}$$

Therefore for a fixed $J_2 \subseteq [1, 2i]$, we have the coefficient $t! 2^v (-1)^{s+r} (1+x)^{2n+1-p-q+t} [t-2i+v]$ and the total coefficient is

$$\begin{aligned}
& t! (-1)^{s+r} \sum_{v=0}^{2i} \binom{2n+1-p-q+t}{t-2i+v} 2^v \binom{2i}{v} \\
&= t! (-1)^{s+r} \sum_{v=0}^{2i} \binom{2i}{v} \binom{2n+1-p-q+t+2i-v}{t}.
\end{aligned}$$

In the right-hand side, only the $2i$ term contributes to the coefficient and it is given by $(-1)^{s+r} d_{2i}$. So $d_{2i} = \sum_{v=0}^{2i} \binom{2i}{v} (2n+1-p-q+t+2i-v)_t$.

Therefore in this case we have $d_i = (-1)^i \sum_{v=0}^i \binom{i}{v} (2n+1-p-q+t+i-v)_t$.

The proof goes in a similar way for the remaining cases and so we omit it. d_i is given by $(-1)^{(q-i)(p-i)+it} \sum_{u=0}^i \binom{i}{u} (2n+1-p-q+t+i-u)_t$ for $Pin(2n+1)$ and d_i is given by $(-1)^{qp+t(p+q)} \sum_{u=0}^i \binom{i}{u} (2n-p-q+t+i-u)_t$ for $Pin(2n)$.

Next we prove the formula (8.1.6).

Both sides of the formula can be considered as the homomorphisms from $\Delta \otimes \otimes^q V$ to $\Delta \otimes \otimes^p V$.

So we can put

$$\prod_{i=1}^q C_{\{i, p+q+i\}} \text{inj}_{\{[1, q+p]\}} = \sum_{i=0}^{\min(p, q)} d_i {}^i \phi_{[q+1, q+p]}^{[p+q+1, p+2q]}$$

As before, we find a sequence of pairs $\{x_i, y_i\}$ and apply both sides of the above to x_i 's and compare the coefficients of y_i 's and determine the coefficients d_i inductively.

We determine d_i for $Pin(2n+1)$.

Case 1 $p = 2r$ and $q = 2s$.

We note $r + s \leq n$.

Case 1-2i. The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, 2i], [\mathbf{s} + \mathbf{i} + 1, \mathbf{s} + \mathbf{r}], \overline{[\mathbf{s} + \mathbf{r}, \mathbf{s} + \mathbf{i} + 1]}\}, x_{2i} = [\emptyset] \otimes \{[1, \mathbf{s} + \mathbf{i}], \overline{[\mathbf{s} + \mathbf{i}, 2i + 1]}\}$. In this case $d_{2i} = (-1)^{\binom{q}{2}}$.

Case 1-2i + 1 The determination of d_{2i+1} .

We take $y_{2i+1} = [\mathbf{s} + \mathbf{i} + 1] \otimes \{[1, 2i + 1], [\mathbf{s} + \mathbf{i} + 1, \mathbf{s} + \mathbf{r}], \overline{[\mathbf{s} + \mathbf{r}, \mathbf{s} + \mathbf{i} + 2]}\}, x_{2i+1} = [\emptyset] \otimes \{[1, \mathbf{s} + \mathbf{i}], 0', \overline{[\mathbf{s} + \mathbf{i}, 2i + 2]}\}$. In this case $d_{2i+1} = (-1)^{s+1}$.

The proof goes in a similar way for the remaining cases and so we omit it. Summarizing all the calculation, we have $d_i = (-1)^{pq + \binom{q}{2} + i(p+q-1)}$ for $Pin(2n+1)$.

For $Pin(2n)$, a similar calculation gives us $d_i = (-1)^{\binom{q}{2}}$.

Finally we prove the formula (8.1.7).

Both sides of the formula can be considered as the homomorphisms from $\Delta \otimes \otimes^q V$ to $\Delta \otimes \otimes^p V$.

So we can put

$$\text{pr}_{\{[1, \mathbf{q} + \mathbf{p}]\}} \prod_{i=1}^p \text{id}_{V_{\{\mathbf{q} + \mathbf{i}, \mathbf{p} + \mathbf{q} + \mathbf{i}\}}} = \sum_{i=0}^{\min(p, q)} d_i \phi_{[\mathbf{p} + \mathbf{q} + 1, \mathbf{q} + 2\mathbf{p}]}^{i, [1, \mathbf{q}]}$$

As before, we find a sequence of pairs $\{x_i, y_i\}$ and apply both sides of the above to x_i 's and compare the coefficients of y_i 's and determine the coefficients d_i inductively.

We determine d_i for $Pin(2n+1)$.

Case 1. $p = 2r$ and $q = 2s$.

We note $r + s \leq n$.

Case 1-2i. The determination of d_{2i} .

We take $y_{2i} = [\emptyset] \otimes \{[1, 2i], [\mathbf{s} + \mathbf{i} + 1, \mathbf{s} + \mathbf{r}], \overline{[\mathbf{s} + \mathbf{r}, \mathbf{s} + \mathbf{i} + 1]}\}, x_{2i} = [\emptyset] \otimes \{[1, \mathbf{s} + \mathbf{i}], \overline{[\mathbf{s} + \mathbf{i}, 2i + 1]}\}$.

In this case $d_{2i} = (-1)^r$.

Case 1-2i + 1. The determination of d_{2i+1} .

We take $y_{2i+1} = [\mathbf{s} + \mathbf{i} + 1] \otimes \{[1, 2i + 1], [\mathbf{s} + \mathbf{i} + 1, \mathbf{s} + \mathbf{r}], \overline{[\mathbf{s} + \mathbf{r}, \mathbf{s} + \mathbf{i} + 2]}\}, x_{2i+1} = [\emptyset] \otimes \{[1, \mathbf{s} + \mathbf{i}], 0', \overline{[\mathbf{s} + \mathbf{i}, 2i + 2]}\}$.

In this case $d_{2i+1} = (-1)^{r-1}$.

The proof goes in a similar way for the remaining cases and so we omit it. Summarizing all the calculation, we have $d_i = (-1)^{\binom{p}{2} + i(p+q+1)}$ for $Pin(2n+1)$.

For $Pin(2n)$, a similar calculation gives us $d_i = (-1)^{\binom{p}{2} + pq}$.

□

9. EXAMPLES OF PRODUCTS OF THE GENERALIZED BRAUER DIAGRAMS

From the result of the previous section, we can calculate the product of the generalized Brauer diagrams.

As we remarked after the statement of Theorem 8.1, we change N in the formulas of the theorem into an indeterminate X simultaneously and we write down the relations of the ‘generic’ centralizer algebra of \mathbf{CP}_k .

We summarize other relations between the contractions and the immersions and pr and inj which follow easily from the definitions. We fix the index set of the tensor positions of $\bigotimes^k V$ from 1 to k and after the contraction $C_{\{i,j\}}$ we consider the i -th and the j -th components are occupied by the empty set and $\text{id}_{V_{\{i,j\}}}$ is allowed if the positions i and j are occupied by the empty set.

Then we have the followings.

Lemma 9.1. (i)

$$C_{\{s,t\}} \text{id}_{V_{\{i,j\}}} = \text{id}_{V_{\{i,j\}}} C_{\{s,t\}} \quad \text{if } \{s,t\} \cap \{i,j\} = \emptyset.$$

(ii)

$$C_{\{s,t\}} \text{id}_{V_{\{t,j\}}} = \begin{pmatrix} s \\ j \end{pmatrix} \quad \text{if } s \neq j.$$

Here $\begin{pmatrix} s \\ j \end{pmatrix}$ denotes the (partial) permutation which sends s -th component to the j -th position.

(iii)

$$C_{\{s,t\}} \text{id}_{V_{\{s,t\}}} = N \text{id}.$$

(iv) If $s, t \in T$, we have

$$C_{\{s,t\}} \text{inj}_T = 0, \quad \text{pr}_T \text{id}_{V_{\{s,t\}}} = 0.$$

(v) If $s, t_1, t_2, \dots, t_r \in [1, n]$ are different from each other, we have

$$\begin{pmatrix} t_i \\ s \end{pmatrix} \text{inj}_{\{t_1, \dots, t_i, \dots, t_r\}} = \text{inj}_{\{t_1, \dots, \overset{i}{s}, \dots, t_r\}}, \quad \begin{pmatrix} t_1 \\ s \end{pmatrix} \text{id}_{V_{\{t_1, t_2\}}} = \text{id}_{V_{\{s, t_2\}}}.$$

(vi) If $s, t_1, t_2, \dots, t_r \in [1, n]$ are different from each other, we have

$$\text{pr}_{\{t_1, \dots, t_i, \dots, t_r\}} \begin{pmatrix} s \\ t_i \end{pmatrix} = \text{pr}_{\{t_1, \dots, \overset{i}{s}, \dots, t_r\}}, \quad C_{\{t_1, t_2\}} \begin{pmatrix} s \\ t_1 \end{pmatrix} = C_{\{s, t_2\}}.$$

(vii) If $s, t, t_1, t_2, \dots, t_r \in [1, n]$ are different from each other, $\text{id}_{V_{\{s,t\}}}$ commutes with inj_T and pr_T and $C_{\{s,t\}}$ commutes with inj_T and pr_T , where $T = \{t_1, t_2, \dots, t_r\}$.

Proof. The first three formulas are easy and the fourth formula follows directly from the alternating property with the indices in T of pr and inj . The rest are obvious from the definitions. \square

Remark 9.2. If we put $e_i = \text{id}_V \{i, i+1\} C_{\{i, i+1\}}$, we can deduce easily $e_i e_{i+1} e_i = e_i$ from the above lemma. These relations were used to define the Birman-Wenzl algebras (q -analogs of the Brauer centralizer algebras). ([2].)

Remark 9.3. We note that

$$\text{inj}_{\{[1, q]\}} \circ \text{inj}_{\{[q+1, q+p]\}} \neq \text{inj}_{\{[q+1, q+p]\}} \circ \text{inj}_{\{[1, q]\}}$$

and

$$\text{pr}_{\{[1, q]\}} \circ \text{pr}_{\{[q+1, q+p]\}} \neq \text{pr}_{\{[q+1, q+p]\}} \circ \text{pr}_{\{[1, q]\}}.$$

Let us show some examples.

Example 9.4. In this example we always assume $n \geq k$ and use the representation theoretical parametrization of the generalized Brauer diagrams, so we omit the subscript rt here. We calculate the product of $y_5 y_8$ in figure 1.

$$y_5 y_8 = \begin{array}{c} \text{---} \\ \vdots \quad \vdots \\ \vdots \end{array} = (X-1) \begin{array}{c} \text{---} \\ \cdot \quad \cdot \end{array} + (X-1) \begin{array}{c} \text{---} \\ \text{---} \end{array}$$

FIGURE 5. The result of the product of $y_5 y_8$ for $\text{Pin}(2n+1)$

Here $y_8 = \text{inj}_{\{1,2\}} C_{\{1,2\}}$ and $y_5 = \text{inj}_{\{1\}} \binom{2}{2} \text{pr}_{\{1\}}$. From the formula (8.1.2), we have $\text{pr}_{\{1\}} \text{inj}_{\{1,2\}} = (X-1)_1 \text{inj}_{\{2\}}$ (we put X for $2n+1$.) and the targeting homomorphism is $\text{inj}_{\{1\}} \binom{2}{2} (X-1) \text{inj}_{\{2\}} C_{\{1,2\}} = (X-1) \text{inj}_{\{1\}} \text{inj}_{\{2\}} C_{\{1,2\}}$. From the formula (8.1.3), we have $\text{inj}_{\{1\}} \text{inj}_{\{2\}} = \text{inj}_{\{1,2\}} + \text{id}_V \{1,2\}$ and the final formula is given in figure 5.

We calculate a more complicated example of figure 6.

In this case we must calculate

$$\text{id}_V \{3,5\} \text{inj}_{\{1,2,6,7\}} \binom{5}{4} \text{pr}_{\{1,2,4,7\}} C_{\{3,6\}} \text{id}_V \{5,7\} \text{inj}_{\{1,2,4,6\}} \binom{1}{3} \text{pr}_{\{2,3,5,6\}} C_{\{4,7\}}.$$

The inside homomorphism of the above, i.e., the homomorphism obtained by taking the middle part between the contractions and pr of the first element and the immersion and inj of the second element, is given by $\binom{5}{4} \text{pr}_{\{1,2,4,7\}} C_{\{3,6\}} \text{id}_V \{5,7\} \text{inj}_{\{1,2,4,6\}} \binom{1}{3}$. This element is in $\mathbf{CP}_1^1 = \text{Hom}_{\text{Pin}(2n)}(\Delta \otimes V, \Delta \otimes V)$. (V sits in the first place in the upper row and sits in the forth place in the lower row.) So it can be written as a sum of the basis of \mathbf{CP}_1^1 i.e., \mathbf{GB}_1^1 . In general, this fact tells us that the linear spaces spanned by all the diagrams in \mathbf{GB}_k^k with at most s ($0 \leq s \leq k$) vertical edges (the edge from the upper row to the lower row), become two-sided ideals in \mathbf{CP}_k .

$$\begin{array}{c}
\begin{array}{ccc}
\begin{array}{c} \cdot \cdot \cdot \cdot \cdot \\ \cdot \cdot \cdot \cdot \cdot \end{array} & \begin{array}{c} \cdot \cdot \cdot \cdot \cdot \\ \cdot \cdot \cdot \cdot \cdot \end{array} & \begin{array}{c} \cdot \cdot \cdot \cdot \cdot \\ \cdot \cdot \cdot \cdot \cdot \end{array} \\
\text{diagram 1} & \text{diagram 2} & \text{diagram 3}
\end{array} = 3(X-2)(X-3) \begin{array}{ccc}
\begin{array}{c} \cdot \cdot \cdot \cdot \cdot \\ \cdot \cdot \cdot \cdot \cdot \end{array} & \begin{array}{c} \cdot \cdot \cdot \cdot \cdot \\ \cdot \cdot \cdot \cdot \cdot \end{array} & \begin{array}{c} \cdot \cdot \cdot \cdot \cdot \\ \cdot \cdot \cdot \cdot \cdot \end{array} \\
\text{diagram 4} & \text{diagram 5} & \text{diagram 6}
\end{array} \\
& & - (X-2)(X-3)(X-4) \sum_{i=1}^5 \sum_{j=1}^5 (-1)^{i+j} z_i y_j
\end{array}$$

Here y_j denotes the upper row and z_i denotes the lower row given as follows.




FIGURE 6. The result of the product of a more complicated example for $Pin(2n)$

From the formula (8.1.7), we have

$$\text{pr}_{\{1,2,3,4\}} \text{id}_V \{4,5\} = -\text{inj}_{\{5\}} \text{pr}_{\{1,2,3\}} - \binom{1}{5} \text{pr}_{\{2,3\}} + \binom{2}{5} \text{pr}_{\{1,3\}} - \binom{3}{5} \text{pr}_{\{1,2\}}.$$

If we apply the conjugation of the permutation $\eta = \begin{pmatrix} 3 & 4 \\ 4 & 7 \end{pmatrix}$, we have

$$\text{pr}_{\{1,2,4,7\}} \text{id}_V \{5,7\} = -\text{inj}_{\{5\}} \text{pr}_{\{1,2,4\}} - \binom{1}{5} \text{pr}_{\{2,4\}} + \binom{2}{5} \text{pr}_{\{1,4\}} - \binom{4}{5} \text{pr}_{\{1,2\}}.$$

From the formula (8.1.6), we have

$$C_{\{1,5\}} \text{inj}_{\{1,2,3,4\}} = \text{inj}_{\{2,3,4\}} \text{pr}_{\{5\}} + \text{inj}_{\{3,4\}} \binom{5}{2} - \text{inj}_{\{2,4\}} \binom{5}{3} + \text{inj}_{\{2,3\}} \binom{5}{4}.$$

If we apply the conjugation of the permutation $\eta = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 6 & 2 & 4 & 1 & 3 \end{pmatrix}$, we have

$$C_{\{3,6\}} \text{inj}_{\{1,2,4,6\}} = -\text{inj}_{\{1,2,4\}} \text{pr}_{\{3\}} + \text{inj}_{\{1,4\}} \binom{3}{2} - \text{inj}_{\{1,2\}} \binom{3}{4} - \text{inj}_{\{2,4\}} \binom{3}{1}.$$

Since $\text{pr}_{\{2,3\}} \text{inj}_{\{1,2\}} = -(X-2) \text{inj}_{\{1\}} \text{pr}_{\{3\}} - \{(X-1-2+1+1-0) + (X-1-2+1+1-1)\} \binom{3}{1}$ from the formula (8.1.5), we have

$\text{pr}_{\{2,4\}} \text{inj}_{\{1,4\}} = (X-2) \text{inj}_{\{1\}} \text{pr}_{\{2\}} + (2X-3) \binom{2}{1}$. The calculation goes on in almost similar way and the final result consists of 26 terms in which only one term contains a vertical line. It is given in figure 6.

10. DUAL PAIR AND SPIN REPRESENTATIONS

In this section we define the subspace in the tensor space $T_k = \Delta \otimes \otimes^k V$, on which the symmetric group of degree k and $Pin(N)$ (or $Spin(N)$) act as the dual pair.

By \mathbf{GB}_k , we denote \mathbf{GB}_k^k .

From Lemma 5.6, the elements of \mathbf{GB}_k under the invariant theoretical parametrization with at most N isolated vertices (we denote a set of these elements by $\mathbf{GB}_{k,\text{inv}}$.) span linearly \mathbf{CP}_k . If $n \geq k$, they become a base of \mathbf{CP}_k and in this case, the elements of \mathbf{GB}_k under the representation theoretical parametrization (we denote a set of these elements by $\mathbf{GB}_{k,\text{rt}}$.) also become a base of \mathbf{CP}_k .

Since \mathbf{CP}_k contains $\mathbb{R}[\mathfrak{S}_k]$ naturally, For $0 \leq s \leq \min(k, n)$, let us define the subspace $T_{k,s}^0$ of $\Delta \otimes \otimes^k V$ as follows.

Definition 10.1. Let $T_{k,s}^0$ denote the intersection of the all the kernels of the contractions $C_{\{i,j\}}$ ($1 \leq i < j \leq k$) and the projections $\text{pr}_{\{i_1, i_2, \dots, i_r\}}$ ($r > 0$ and $1 \leq i_1 < i_2 < \dots < i_r \leq k$) and the alternating operators $\text{Alt}_{\{i_1, i_2, \dots, i_r\}}$ (see Definition 6.3 and $r > s$) of degree greater than s .

If $n \geq k$, then we ignore the alternating operators in the above and put $T_{k,k}^0 = T_k^0$.

Lemma 10.2. The space $T_{k,s}^0$ becomes a $\mathbf{CP}_k \times Pin(N)$ module.

Proof. It is obvious that $T_{k,s}^0$ is a $Pin(N)$ module from the definition.

Let $y_{\text{inv}} \in \mathbf{GB}_{k,\text{inv}}$ and let $\psi_{T_\ell}^{\text{T}_u}$ be the homomorphism corresponding to the isolated vertices in y .

Then if $|T_u| > s$, $\psi_{T_\ell}^{\text{T}_u}$ acts on the space $T_{k,s}^0$ by 0 and so $y_{\text{inv}} T_{k,s}^0 = 0$.

We can assume $|T_u| \leq s = \min(k, n)$. If $|T_\ell| \neq |T_u|$, the contractions or pr must appear in the upper row, so $y_{\text{inv}} T_{k,s}^0 = 0$. (See (7.2.1).)

We can assume $|T_u| = |T_\ell| \leq s = \min(k, n)$. Then from Theorem 7.2, $\psi_{T_\ell}^{\text{T}_u}$ acts on this space by the alternating operators of degree $|T_u|$. Since the space $T_{k,s}^0$ is invariant under the action of \mathfrak{S}_k , the lemma holds. \square

Let us denote the set of the annihilators in \mathbf{CP}_k of the module $T_{k,s}^0$ by $\mathfrak{J}_{k,s}$. Then from the above, $\mathfrak{J}_{k,s}$ becomes a two sided ideal in \mathbf{CP}_k and $\mathbf{CP}_k/\mathfrak{J}_{k,s}$ is the surjective image of the subalgebra $\mathbb{R}[\mathfrak{S}_k]$. Since $Pin(N)$ and \mathbf{CP}_k are dual pair on T_k , on this direct summand $T_{k,s}^0$, the images of $\mathbb{R}[\mathfrak{S}_k]$ and $Pin(2n)$ become a dual pair.

So we have the following theorem.

Theorem 10.3. The symmetric group of degree k and $Pin(N)$ become a dual pair on the subspace $T_{k,s}^0$ of $\Delta_{(\epsilon_1)} \otimes \otimes^k V$. Namely the space $T_{k,s}^0$ is decomposed into the direct sum of the tensor products of the

irreducible representations with multiplicity free as follows.

$$(10.3.1) \quad T_{k,s}^0 = \sum_{\substack{\lambda: \text{partitions of size } k \\ \ell(\lambda) \leq s}} \lambda_{\mathfrak{S}_k} \otimes [\Delta, \lambda]_{Pin(N), (\epsilon_2)}.$$

Here for $Pin(2n+1)$, $\epsilon_1(-1)^k = \epsilon_2$

Remark 10.4. For $Spin(2n+1)$, on the above $T_{k,s}^0$, \mathfrak{S}_k and $Spin(2n+1)$ also act as a dual pair, since the center z acts on it by the scalar $\epsilon_1(-1)^k$. The decomposition is the same if we ignore ϵ_1 and ϵ_2 .

Remark 10.5. From the above, if $n \geq k$, $\mathfrak{J}_{k,k}$ is the linear space spanned by $y_{rt} \in \mathbf{GB}_{k,rt}$ with at most $k-1$ vertical edges and in this case, we have $\mathbf{CP}_k = \mathbb{R}[\mathfrak{S}_k] \oplus \mathfrak{J}_{k,k}$.

Proof. From the argument just before this theorem, $\mathbb{R}[\mathfrak{S}_k]$ and $Pin(N)$ become a dual pair on the space $T_{k,s}^0$.

The Schur-Weyl reciprocity tells us that, as $\mathfrak{S}_k \times GL(N)$ module, $\bigotimes^k V$ is decomposed into

$$(10.1) \quad \bigotimes^k V = \sum_{\lambda: \text{partitions of size } k} \lambda_{\mathfrak{S}_k} \otimes \lambda_{GL(N)}.$$

For λ with $\ell(\lambda) \leq s \leq n$, the restriction rules (See the proposition 1.5.3 in [16].) tells us that we have

$$\lambda_{GL(N)} \downarrow_{O(N)}^{GL(N)} = \sum_{\kappa, \mu} LR_{2\kappa, \mu}^\lambda \mu_{O(N)}.$$

Here 2κ denotes the even partition and $LR_{2\kappa, \mu}^\lambda$ denotes the Littlewood-Richardson coefficient. So $\lambda_{GL(2n)} \downarrow_{O(2n)}^{GL(2n)}$ contains $\lambda_{O(2n)}$ with multiplicity one and $\lambda_{O(2n)}$ never appears in the other components of the right-hand sides of (10.1), since the size $|\lambda|$ of λ is equal to k .

From (2.4.3) in Theorem 2.4 and (2.2.6) in Theorem 2.2, $[\Delta, \lambda]_{Pin(N)}$ occurs exactly once in $\Delta \otimes \lambda_{O(2n)}$.

Since $|\lambda| = k$, from (2.4.2) and (2.2.3) and (2.2.4), $[\Delta, \lambda]_{Pin(N)}$ never appears in $\Delta \otimes \bigotimes^u V$ for $u < k$. So this representation must be in the kernels of the contractions and pr's.

From the property of the Young symmetrizers, all the alternating operators $\text{Alt}_{\{i_1, i_2, \dots, i_r\}}$ ($r > s$) of degree greater than s annihilate the space $\sum_{\ell(\lambda) \leq s} \lambda_{\mathfrak{S}_k} \otimes \lambda_{GL(2n)}$.

So $\lambda_{\mathfrak{S}_k} \otimes [\Delta, \lambda]_{Pin(N)}$ for partitions λ of size k and of $\ell(\lambda) \leq s$ must be in the space $T_{k,s}^0$.

Since we already know that on the space $T_{k,s}^0$, \mathfrak{S}_k and $Pin(N)$ act as a dual pair and all the irreducible representations $\lambda_{\mathfrak{S}_k}$ of \mathfrak{S}_k with $\ell(\lambda) \leq s$ already appear, so we have the theorem. \square

Next we move to the case for $Spin(2n)$.

For $Spin(2n)$ representations, we consider the endomorphism $(A \otimes \text{id}) \in \text{End}(\Delta \otimes \otimes^k V)$. Here A is the associator for Δ and is given by the degree operator $A[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r] = (-1)^r[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r]$.

Lemma 10.6. *The endomorphism $A \otimes \text{id}$ commutes with both of the actions of \mathbf{CP}_k and $Spin(2n)$ on the space $\Delta \otimes \otimes^k V$.*

Moreover we have

$$\text{pr}_p \circ (A \otimes \text{id}) = (-1)^p A \circ \text{pr}_p$$

and

$$\text{inj}_q \circ A = (-1)^q (A \otimes \text{id}) \text{inj}_q.$$

Proof. It is obvious that $(A \otimes \text{id})$ commutes with the action of the Brauer algebra $\text{End}_{O(2n)}(\otimes^k V)$ and the action of $Spin(2n)$, since the action of $Spin(2n)$ preserves the parity r , where r is the number of the indices in $[\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \dots, \mathbf{i}_r]$.

We note that from the parity argument, ψ_q^p corresponding to the isolated points in any generalized Brauer diagrams $y_{inv} \in \mathbf{GB}_{k,inv}$ must satisfy $p + q \equiv 0 \pmod{2}$.

From the formula for $Pin(2n)$ in Theorem 7.1, if $p + q \equiv 0 \pmod{2}$, we can easily check $\psi_q^p \circ (A \otimes \text{id}) = (A \otimes \text{id}) \circ \psi_q^p$.

The last two formulas are straightforward. \square

So we first decompose the space $T_k = \Delta \otimes \otimes^k V$ into the irreducible constituents under the action of $Pin(2n) \times \mathbf{CP}_k$. Since $Pin(2n)$ and \mathbf{CP}_k are semisimple and act on the space as the dual pair, from the Wedderburn's theorem, we have

$$\Delta \otimes \otimes^k V = \bigoplus_{\substack{\ell(\lambda) \leq n \\ |\lambda| \leq k}} [\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}.$$

Here $\lambda_{\mathbf{CP}_k}$ is the irreducible representation of \mathbf{CP}_k . We consider the subspace $[\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}$ as $Spin(2n) \times \mathbf{CP}_k$ module. Then we have

$$[\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k} = (1/2 + \lambda)_{Spin(2n)}^+ \otimes \lambda_{\mathbf{CP}_k} \oplus (1/2 + \delta)_{Spin(2n)}^- \otimes \lambda_{\mathbf{CP}_k}.$$

Since $A \otimes \text{id}$ commutes with the action of \mathbf{CP}_k and the space $[\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}$ is the homogeneous component of the irreducible representation $\lambda_{\mathbf{CP}_k}$ of \mathbf{CP}_k , we have

$$(A \otimes \text{id})([\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}) = [\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}.$$

Since $(A \otimes \text{id})^2 = \text{id}$, ± 1 eigenspaces of $(A \otimes \text{id})$ in $[\Delta, \lambda] \otimes \lambda_{\mathbf{CP}_k}$ become non-equivalent $Spin(2n) \times \mathbf{CP}_k$ modules.

Here $\Delta^+ \otimes \otimes^k V$ is $+1$ eigenspace and $\Delta^- \otimes \otimes^k V$ is -1 eigenspace of $A \otimes \text{id}$ in $\Delta \otimes \otimes^k V$.

From Theorem 2.2, $(1/2 + \lambda)_{Spin(2n)}^{\pm}$ appears in $\Delta^{\pm} \otimes \otimes^k V$ if and only if $k - |\lambda| \equiv 0 \pmod{2}$. Also $(1/2 + \lambda)_{Spin(2n)}^{\mp}$ appears in $\Delta^{\pm} \otimes \otimes^k V$ if and only if $k - |\lambda| \equiv 1 \pmod{2}$.

So $(-1)^{k-|\lambda|}$ eigenspace of $(A \otimes \text{id})$ in $[\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}$ is the irreducible representation $(1/2 + \lambda)_{Spin(2n)}^+ \otimes \lambda_{\mathbf{CP}_k}$ and $(-1)^{k-|\lambda|+1}$ eigenspace of $(A \otimes \text{id})$ in $[\Delta, \lambda]_{Pin(2n)} \otimes \lambda_{\mathbf{CP}_k}$ is the irreducible representation $(1/2 + \lambda)_{Spin(2n)}^- \otimes \lambda_{\mathbf{CP}_k}$.

We define the extension of the algebra \mathbf{CP}_k by $(A \otimes \text{id})$.

Definition 10.7. Let \mathbf{CS}_k be the subalgebra of $\text{Hom}(\Delta \otimes \otimes^k V, \Delta \otimes \otimes^k V)$ generated by \mathbf{CP}_k and $(A \otimes \text{id})$.

Then \mathbf{CS}_k and $Spin(2n)$ act on the space $\Delta \otimes \otimes^k V$ as the dual pair.

Then Theorem 10.3 for $Spin(2n)$ is given as follows. Since we have Lemma 10.6, $T_{k,s}^0$ in Definition 10.1 is $(A \otimes \text{id})$ stable and We decompose $T_{k,s}^0$ into the ± 1 eigenspaces of $(A \otimes \text{id})$ and denote them by $T_{k,s}^{0,\pm}$. Then from Theorem 10.3, if we note that only λ 's considering here satisfy $|\lambda| = k$, we have the following theorem.

Theorem 10.8. Then the symmetric group of degree k and $Spin(2n)$ becomes the dual pair on the subspaces $T_{k,s}^{0,+}$ and $T_{k,s}^{0,-}$ respectively. Namely, $T_{k,s}^{0,\pm}$ are decomposed into the direct sum of the tensor products of the irreducible representations with multiplicity free as follows.

$$(10.8.1) \quad T_{k,s}^{0,\pm} = \sum_{\substack{\lambda: \text{partitions of size } k \\ \ell(\lambda) \leq s}} \lambda_{\mathfrak{S}_k} \otimes (1/2 + \lambda)_{Spin(2n)}^{\pm}.$$

REFERENCES

- [1] M. F. Atiyah, R. Bott and A. Shapiro Clifford Modules, *Topology* **3**, (1964) pp. 3–38.
- [2] J. Birman and H. Wenzl, Braids, link polynomials and a new algebra, *Trans. Amer. Math. Soc.* **313**, (1989) pp. 249–273
- [3] R. Brauer, On algebras which are connected with the semisimple continuous groups, *Ann. of Math.* **38**, No 4, (1937) pp. 857–872.
- [4] R. Brauer and H. Weyl, Spinors in n dimensions, *Amer. J. Math.* **57**, (1935) pp. 425–449
- [5] C. Chevalley, The Algebraic Theory of Spinors and Clifford Algebras, *Collected Works*, **2**, Springer, (1997)
- [6] W. F. Doran IV and D. B. Wales and P. J. Hanlon, On the semisimplicity of the Brauer Centralizer Algebras, *J. Algebra*, **211**, (1999) pp. 647–685
- [7] S. V. Fomin, Generalized Robinson-Schensted-Knuth correspondence, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **155**, (1986) pp. 156–175. [in Russian]
- [8] W. Fulton and J. Harris, Representation Theory – A First Course, *Graduate Texts in Mathematics* **129**, Springer-Verlag, (1991)

- [9] R. Goodman and N. Wallach, Representations and Invariants of the Classical Groups, *Encyclopedia of Mathematics and its Applications* **68**, Cambridge Univ. Press, (1998)
- [10] R. Howe, Remarks on classical invariant theory, *Trans. Amer. Math. Soc.* **313**, (1989) pp. 539–570.
- [11] J. E. Humphreys, Introduction to Lie Algebras and Representation Theory, *Graduate Texts in Mathematics*, **9**, Springer-Verlag, (1972), (1980)
- [12] K. Koike, On the decomposition of tensor products of the representations of the classical groups, *Adv. Math.* **74**, (1989) pp. 57–86.
- [13] K. Koike, Principal specializations of the classical groups and q-analogs of the dimension formulas, *Adv. Math.* **125**, (1997) pp. 236–274.
- [14] K. Koike, Representations of Spinor Groups and the Difference Characters of $SO(2n)$, *Adv. Math.* **128**, (1997) pp. 40–81.
- [15] K. Koike, On Representation of the Classical Groups, *Amer. Math. Soc. Trans.* **183**, (1998) pp. 79–100.
- [16] K. Koike and I. Terada, Young-diagrammatic methods for the representation theory of the classical groups of type B_n , C_n , D_n , *J. Algebra* **107**, (1987) pp. 466–511.
- [17] K. Koike and I. Terada, Young-diagrammatic methods for the restriction of representations of complex classical Lie groups to reductive subgroups of maximal rank, *Adv. Math.* **79**, (1990) pp. 104–135.
- [18] D. E. Littlewood, The Theory of Group Characters, *2nd ed.*, Oxford University Press, Oxford, (1950)
- [19] T. Roby, Applications and Extensions of Fomin’s Generalization of the Robinson-Schensted Correspondence to Differential Posets, *Ph. D Thesis M.I.T.*, (1991)
- [20] G. Takeuchi, *Lie Algebra and Particle Theory (in Japanese)*, Shoukabou, (in Japanese), (1983)
- [21] H. Weyl, The Classical Groups, their Invariants and Representations, *2nd edition*, Princeton Univ. Press, Princeton, N.J., (1946)

DEPARTMENT OF MATHEMATICS, AOYAMA GAKUIN UNIVERSITY, 5-10-1, SAGAMIHARA-SHI, FUCHINOBE, 229-8558, JAPAN

E-mail address: koike@gem.aoyama.ac.jp