

On Evaluation of Nonlinear Exponential Sums

N. A. Carella, June 2002

Abstract: This paper provides a technique for evaluating the nonlinear exponential sums $T_d(\chi, s) = \sum_{0 \leq x < p} \chi(x) \exp(i2\pi s x^d / p)$ in closed forms. The evaluation is obtained from the known values of simpler exponential sums.

Keywords: Character sums, Gaussian sums.

Mathematics Subject Classification: Primary 11L05, Secondary 11L03.

1 Introduction

This paper provides a technique for evaluating the nonlinear exponential sums $T_d(\chi, s) = \sum_{0 \leq x < p} \chi(x) \exp(i2\pi s x^d / p)$ in closed forms. The evaluations are obtained from the known values of simpler exponential sums. Investigation of the properties of the related gaussian sums $S_k(s) = \sum_{0 \leq x < p} \exp(i2\pi s x^k / p)$ and $G_n(\chi, s) = \sum_{0 \leq x < p} \chi(x) \exp(i2\pi s x / p)$ has a long history, it extends back to at least two centuries. Very elaborate algebraic and analytical techniques have been developed to deal with this problem. Nonetheless, many questions about these sums are still unresolved, even for the simplest cases of the cubic, quartic and quintic gaussian sums.

2 Closed Forms Evaluations of $T_d(\chi, s)$

Let χ be a multiplicative character on \mathbf{F}_p of order $\text{ord}(\chi) = n$. The exact value of the nonlinear exponential sum

$$(1) \quad T_d(\chi, s) = \sum_{x=0}^{p-1} \chi(x) e^{i2\pi s x^d / p}$$

of degree d will be determined in terms of the values of sums S_2, S_3, \dots, S_k , or equivalently as a polynomial $T_d(\chi, s) \in \mathbb{Z}[S_2, S_3, \dots, S_k]$. The evaluations already known are the followings:

- (1) $T_1(\chi, s) = G_n(\chi, s)$, with characters of order $n = \text{ord}(\chi) = 2, 3$, and 4. The analysis of the cases $n = 3, 4$ appear in [6], [7], see also [3], [5], [9], and [10], etc.
- (2) $T_d(\chi, s) = 0$, with $\text{ord}(\chi) = 2$, d even and $p \equiv 3 \pmod{4}$.

Case (2) above is easy. The remaining sums $T_d(\chi, s)$ with $p - 1 \equiv 0 \pmod{d}$ are not necessarily trivial. The closed form evaluation of $T_2(\chi)$, $T_3(\chi)$, and $T_4(\chi)$, $\text{ord}(\chi) = 2$, up to a sign will be determined next. Some of the sign ambiguity has not been resolved yet.

Theorem 1. Let $p = a_4^2 + b_4^2$ be a prime for which $a_4 \equiv -\mu \pmod{4}$, $\mu \equiv 2^{(p-1)/2} \pmod{p}$, and let $\text{ord}(\chi) = 2$. Then

$$(2) \quad \sum_{x=0}^{p-1} \left(\frac{x}{p} \right) e^{i2\pi x^2/p} = \pm \sqrt{2\mu(p + a_4\sqrt{p})}.$$

Proof: It is simple matter to show that $S_4 = S_2 + T_2(\chi)$. Substituting the value of S_4 , [2, p. 362], completes the claim. (Note that $T_2(\chi) = 0$ for $p = 4m + 3$). ■

The symbol $i^* = (-1)^{(p-1)^2/8}$ used below is equal to either 1 or i depending on whether $p \equiv 1$ or $3 \pmod{4}$.

Theorem 2. Let $p \equiv 1 \pmod{6}$, and let $\text{ord}(\chi) = 2$. Then

$$(3) \quad T_3(\chi) = \begin{cases} i^*(S_3^2 - p)p^{-1/2} & \text{if } 2 \text{ is a cubic residue,} \\ i^*(4p - S_3^2 \pm S_3(12p - 3S_3^2)^{1/2})p^{-1/2}/2 & \text{if } 2 \text{ is a cubic nonresidue.} \end{cases}$$

Proof: Consider the equation $S_6 = S_3 + T_3(\chi)$, and substitute the value of S_6 , [2, p. 360]. ■

The case of $p \equiv 5 \pmod{6}$ reduces to $S_6 = S_2$, this follows from $S_k = S_d$, for $d = \gcd(k, p - 1)$. A theorem of Gauss asserts that $2 \in \mathbf{F}_p$ is a cubic residue if and only if the prime $p = 3n + 1$ has the quadratic form representation $p = x^2 + 27y^2$, for some integers $x, y \in \mathbb{Z}$. So the complex number $T_3(\chi)$ assumes the first entry in the formula if p is a prime of this form. The sign in the second part of the formula is not difficult to determine, for details, see [2, p. 360].

Corollary 3. If χ is the quadratic symbol and $p \equiv 1 \pmod{6}$, then

$$(4i) \quad \sqrt{p} \leq |T_3(\chi, s)| \leq 3\sqrt{p} \quad \Leftrightarrow \quad p = x^2 + 3y^2,$$

$$(4ii) \quad 0 \leq |T_3(\chi, s)| \leq 3\sqrt{p} \quad \Leftrightarrow \quad p \neq x^2 + 3y^2.$$

Theorem 4. Let $p = a_8^2 + 2b_8^2 = 8m + 1$ be prime, $a_8 \equiv -1 \pmod{4}$, and $\eta \equiv 2^{(p-1)/4} \pmod{p}$. Then

$$(5) \quad \sum_{x=0}^{p-1} \chi(x) e^{i2\pi x^4/p} = \pm \sqrt{(a_8 + \sqrt{p})(2\eta \sqrt{2(p + a_4 \sqrt{p})} + (-1)^m \sqrt{p})}.$$

The proof/evaluation of the sum $T_d(\chi)$, $\text{ord}(\chi) = 2$, has the same format: $T_d(\chi, s) = S_{2d}(s) - S_d(s)$, with $\text{gcd}(2d, p-1) = 2d$. Currently it can also be evaluated for $d = 2, 3, 4, 6, 8$ and 12 in terms of the known values $S_2, S_4, S_6, S_8, S_{12}, S_{16}$, and S_{24} , up to signs see [2], [1]. In addition, the exact values of S_k over \mathbf{F}_q for the parameter $k = p^e + 1, p \mid q$, are given in [4].

The last sum considered is evaluated in terms of the well known Kloostermann sum and Salie sum:

$$(6) \quad K(a, b) = \sum_{x=0}^{p-1} e^{i2\pi(ax+bx^{-1})/p} \quad \text{and} \quad S(a, b) = \sum_{x=0}^{p-1} \left(\frac{x}{p} \right) e^{i2\pi(ax+bx^{-1})/p},$$

respectively

Theorem 5. Let p be a prime, and let $0 \neq a, b \in \mathbb{Z}$. Then

$$(7) \quad \sum_{x=0}^{p-1} e^{i2\pi(ax^2+bx^{-2})/p} = 2\sqrt{p} \left(\left(\frac{a}{p} \right) (-1)^{(p-1)^2/8} \cos(4\pi a/p) + \cos(2\pi \theta/p) \right),$$

where the $p-1$ discrete angles $\theta \in [0, \pi]$ are distributed in accordance to the Sato-Tate measure $2\pi^{-1} \sin(\theta) d\theta$.

On Evaluation of Nonlinear Exponential Sum

References:

- [1] B.C. Berndt, R.J. Evans, *The Determination of Gaussian Sums*, Bulletin Amer. Math. Soc., Vol. 5, No. 2, Sept. 1981, p.107-129.
- [2] _____, *The Determination of Gauss and Jacobi Sums* Journal of Number Theory Vol. No. 11, 1979, p.349-398.
- [3] J.W.S. Cassels, *On Kummer sums*, Proc. London Math. Soc., (3) 21 (1970), 19 - 27.
- [4] RS Coulter, *Evaluation of Weil sums*, Acta Arith. Vol. 86, No. 3 (1998) 217-226.
- [5] J.H. Loxton, *Some conjectures concerning Gaussian sums*, J. Reine Angew. Math. 297 (1978), 153 - 158.
- [6] C.R. Matthews, *Gaussian sums and elliptic functions I*, Invent. Math. 52 (1978), 2, 163 - 185.
- [7] _____, *Gaussian sums and elliptic functions II*, Invent. Math. 54 (1979), 1, 23 - 52.
- [8] A.D. McGettrick, *On the biquadratic gaussian sums*, Proc. Cambridge, Phil. Soc., 71 (1972), 79 - 83.
- [9] Yamamoto, *On gaussian sums with biquadratic residue characters*, J. Reine Angew. Math. 219 (1965), 200-213.
- [10] K.S. Williams, *On Salie's Sum*, J. Number Theory, 3, p.311-317, 1971.