

OPEN CONDITIONS FOR INFINITE MULTIPLICITY EIGENVALUES ON ELLIPTIC CURVES

BO-HAE IM AND MICHAEL LARSEN

ABSTRACT. Let E be an elliptic curve defined over a number field K . We show that for each root of unity ζ , the set Σ_ζ of $\sigma \in \text{Gal}(\overline{K}/K)$ such that ζ is an eigenvalue of infinite multiplicity for σ acting on $E(\overline{K}) \otimes \mathbb{C}$ has non-empty interior.

For the eigenvalue -1 , we can show more: for any σ in $\text{Gal}(\overline{K}/K)$, the multiplicity of the eigenvalue -1 is either 0 or ∞ . It follows that Σ_{-1} is open.

1. INTRODUCTION

Let K be a number field, $\overline{K}$ an algebraic closure of K , and $G_K := \text{Gal}(\overline{K}/K)$ the absolute Galois group of $\overline{K}$ over K . Let E be an elliptic curve defined over K . There is a natural continuous action of G_K on the countably infinite-dimensional complex vector space $V_E := E(\overline{K}) \otimes \mathbb{C}$. The resulting representation decomposes as a direct sum of finite-dimensional irreducible representations in each of which G_K acts through a finite quotient group.

In particular, the action of every $\sigma \in G_K$ on V_E is diagonalizable, with all eigenvalues roots of unity. In [3], the first-named author showed that for *generic* σ , every root of unity appears as an eigenvalue of countably infinite multiplicity. This is true both in terms of measure and of Baire category. However, there exist σ for which the spectrum is quite different: trivially, the identity and complex conjugation elements; less trivially, examples which can be constructed for an arbitrary set S of primes, such that ζ is an eigenvalue if and only if every prime factor of its order lies in S .

Throughout this paper, we will write Σ_ζ for the subset of G_K consisting of elements σ acting as ζ on an infinite-dimensional subspace of V_E (E and K being fixed). For $\zeta = 1$, a good deal is known. In [2], it is proved that whenever 1 appears as an eigenvalue of σ at all, we have $\sigma \in \Sigma_1$. It follows that Σ_1 is open. By [4], when $K = \mathbb{Q}$, Σ_1 is all of G_K , and

Date: December 8, 2018.

2000 *Mathematics Subject Classification.* Primary 11G05.

quite possibly this may be true without restriction on K . We have already observed that $\Sigma_\zeta \neq G_K$ for $\zeta \neq 1$. We can still hope for positive answers to the following progression of increasingly optimistic questions:

Question 1.1. *Does Σ_ζ have non-empty interior for all ζ ?*

Question 1.2. *Is Σ_ζ open for all ζ ?*

Question 1.3. *Do all eigenvalues of σ acting on V_E appear with infinite multiplicity?*

In this paper, we give an affirmative answer to Question 1.1 for all ζ and an affirmative answer to all three questions for $\zeta = -1$.

The difficulty in proving such theorems is that placing σ in a basic open subset U of G_K amounts to specifying the action of σ on a finite Galois extension L of K . By the Mordell-Weil theorem, $E(L) \otimes \mathbb{C}$ cannot provide an infinite eigenspace for ζ . Thus, the intersection of eigenspaces

$$\bigcap_{\sigma \in U} V_E^{\sigma - \zeta}$$

is finite-dimensional. Thus, the behavior of a finite collection of rational points must be enough to guarantee the existence of infinitely many linearly independent points on the curve with specified σ -action.

We would like to thank L. Moret-Bailly for correcting an error in the proof of Proposition 3.1 in an earlier draft of this manuscript.

2. MULTIPLICITY OF THE EIGENVALUE -1

In this section, we answer Questions 1.2 and 1.3 for $\zeta = -1$.

Proposition 2.1. *Let E/K be an elliptic curve over K . Suppose -1 is an eigenvalue of the action of $\sigma \in G_K$ on V_E . Then the -1 -eigenspace of σ is infinite-dimensional.*

Proof. As -1 is an eigenvalue of σ acting on V_E , it is an eigenvalue of σ acting on $E(\overline{K}) \otimes \mathbb{Q}$. Clearing denominators, there exists a non-torsion $P \in E(\overline{K})$ such that $\sigma(P) + P \in E(\overline{K})_{\text{tor}}$. Replacing P by a suitable positive integral multiple, $\sigma(P) = -P$.

Let $y^2 = f(x)$ be a fixed Weierstrass equation of E/K . Let $P = (\alpha, \sqrt{f(\alpha)})$. As $\sigma(P) = -P$, we have $\alpha \in \overline{K}^\sigma$ but $\sigma(\sqrt{f(\alpha)}) = -\sqrt{f(\alpha)}$ so $\sqrt{f(\alpha)} \notin \overline{K}^\sigma$. Then, $\sqrt{f(\alpha)} \notin K(\alpha)$, since $K(\alpha) \subseteq \overline{K}^\sigma$

Note that $f(\alpha) \in K(\alpha) \subseteq \overline{K}^\sigma$. Let $c = f(\alpha) \in K(\alpha)$. We still have $\sigma \in \text{Gal}(\overline{K}/K(\alpha))$ and $\sigma(\sqrt{c}) = -\sqrt{c}$.

Let $E'/K(\alpha)$ denote the twist $y^2 = cf(x)$. Then, E' has a rational point $P' = (\alpha, f(\alpha))$ over $K(\alpha)$. The $\overline{K}$ -isomorphism $\phi: E \rightarrow E'$ mapping $(x, y) \mapsto (x, \sqrt{f(\alpha)y})$ sends P to P' , so P' is of infinite order on E' . By ([2, Theorem 5.3], $E'(\overline{K}^\sigma)$ has infinite rank. Let $\{P'_i = (x_i, \sqrt{cf(x_i)})\}_{i=1}^\infty$ be an infinite sequence of linearly independent points of E' generating the infinite dimensional eigenspace of 1 of σ in $E'(\overline{K}) \otimes \mathbb{C}$. Then, $\sigma(x_i) = x_i$ and $\sigma(\sqrt{f(x_i)}) = -\sqrt{f(x_i)}$ for all i , since $\sigma(\sqrt{c}) = -\sqrt{c}$.

Let $P_i = \phi^{-1}(P'_i) = (x_i, \sqrt{f(x_i)})$. These are points of the given elliptic curve E such that $\sigma(P_i) = -P_i$ for all i , since $\sigma(x_i) = x_i$ and $\sigma(\sqrt{f(x_i)}) = -\sqrt{f(x_i)}$.

The points P_i are linearly independent because the P'_i are so. Therefore, $\{P_i \otimes 1\}_{i=1}^\infty$ generates an infinite dimensional subspace of the -1 -eigenspace of σ on V_E . This completes the proof. $\square$

Theorem 2.2. *Let E/K be an elliptic curve over K . Then, Σ_{-1} is open.*

Proof. We have already seen that if $\sigma \in \Sigma_{-1}$, we can choose a point $P \in E(\overline{K})$ of infinite order such that $\sigma(P) = -P$. By Proposition 2.1, $\tau(P) = -P$ implies $\tau \in \Sigma_{-1}$. It follows that Σ_{-1} contains the open neighborhood $\{\tau \in G_K \mid \tau(P) = \sigma(P)\}$ of σ . $\square$

Remark 2.3. The same argument shows that Questions 1.2 and 1.3 have an affirmative answer for $\zeta = \omega$ (resp. $\zeta = i$) when E has complex multiplication by $\mathbb{Z}[\omega]$ (resp. $\mathbb{Z}[i]$).

3. INTERIOR POINTS

In this section, we show that for every root of unity ζ , the set Σ_ζ contains a non-empty open subset. We assume that the order of ζ is $n \geq 3$, the case $n = 1$ having been treated in [2], and the case $n = 2$ in Theorem 2.2.

Our strategy will be to find points $Q_i \in E(\overline{K})$ such that the σ -orbit of Q_i has length n . For each such point Q_i , we set

$$(1) \quad R_i := \sum_{i=0}^{n-1} \sigma^i(Q_i) \otimes \zeta^{-i}$$

and observe that R_i is a ζ -eigenvector of σ provided that it is non-zero.

We therefore begin with the following proposition:

Proposition 3.1. *Let X be a Riemann surface of genus $g > 1$ with an automorphism σ of order n . Then X contains a non-empty open set U such that $x \in U$ implies that*

$$\sum_{i=0}^{n-1} [\sigma^i x] \otimes \zeta^{-i} \neq 0$$

in $\text{Pic } X \otimes \mathbb{C}$.

Proof. We can regard X as the group of complex points of a non-singular projective curve whose Picard scheme has complex locus $\text{Pic } X$. Then $\text{Pic } X \otimes \mathbb{Z}[\zeta]$ is the group of complex points of a group scheme whose identity component $\text{Pic}^0 X \otimes \mathbb{Z}[\zeta]$ is isomorphic to the $\phi(n)$ th power of the Jacobian variety of this curve. The action of σ on X defines an action on $\text{Pic } X$, and the map $\psi: \text{Pic } X \rightarrow \text{Pic } X \otimes \mathbb{Z}[\zeta]$ given by

$$\psi(y) = \sum_{i=0}^{n-1} \sigma^i y \otimes \zeta^{-i}$$

then comes from a morphism of group schemes. The image of ψ actually lies in $\text{Pic}^0 X \otimes \mathbb{Z}[\zeta]$, and its kernel P_ζ^0 is Zariski-closed in $\text{Pic } X$.

The set P_ζ of y such that $\psi(y)$ maps to 0 in $\text{Pic } X \otimes \mathbb{C}$ is the union of all translates of P_ζ^0 by torsion points of $\text{Pic } X$. Applying Raynaud's theorem [7] (i.e., the proof of the Manin-Mumford conjecture) to the image of X in $\text{Pic } X / P_\zeta^0$, the intersection $X \cap P_\zeta$ is finite whenever P_ζ^0 does not contain the identity component of $\text{Pic } X$. It therefore suffices to prove that the Lie algebra of P_ζ^0 is a proper subspace of the Lie algebra of $\text{Pic } X$. We identify the latter Lie algebra in the usual way [1, Ch. 2, §6] with $H^1(X, \mathcal{O}_X) = H^{0,1}(X)$. As $V_E = (E(\overline{K}) \otimes \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C}$, the eigenvalues of σ form a Galois-invariant set. Therefore, it suffices to show *some* primitive n th root of unity is an eigenvalue of σ acting on $H^{0,1}$. The Hodge decomposition

$$H^1(X(\mathbb{C}), \mathbb{C}) = H^{1,0} \oplus \overline{H^{1,0}}$$

implies that it suffices to show that ζ is an eigenvalue of σ acting on $H^1(X(\mathbb{C}), \mathbb{C})$.

Let $R_{\mathbb{C}}(G)$ denote the ring of complex (virtual) representations of G . For any subgroup H of $G := \langle \sigma \rangle$, let $R_{G/H}$ denote the regular representation of G/H regarded as an element of $R_{\mathbb{C}}(G)$, and let $I_H := R_G - R_{G/H}$. In particular, $I_{\{1\}} = 0$. Regarded as an element of

$R_{\mathbb{C}}(G)$, the G -representation $H^1(X(\mathbb{C}), \mathbb{C})$ is

$$2g + (2h - 2)I_G + \sum_{[x] \in X/G} I_{\text{Stab}_G(x)},$$

where h is the genus of X/G , and $\text{Stab}_G(x)$ is the stabilizer of any element of X representing the G -orbit $[x]$. This is worked out in the case that $h = 0$ in [6, Prop. 2.2]), but the method (in which the character of $H^1(X(\mathbb{C}), \mathbb{C})$ as a representation of G is deduced from the Hurwitz formula and the Lefschetz trace formula) works in general.

The multiplicity of ζ as an eigenvalue of σ on $H^1(X(\mathbb{C}), \mathbb{C})$ is therefore $2g - 2$ plus the number of ramification points of the cover $X \rightarrow X/G$. This is positive except in two cases: the cyclic cover $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ of degree n (necessarily ramified over two points) and a degree n isogeny of elliptic of elliptic curves. As $g \geq 2$, neither case applies.

□

Theorem 3.2. *Let E/K be an elliptic curve over a number field K . For each root of unity ζ , there exists a nonempty open subset Σ_{ζ} of $\text{Gal}(\overline{K})$ such that the multiplicity of the eigenvalue ζ for $\sigma \in \Sigma_{\zeta}$ acting on $E(\overline{K}) \otimes \mathbb{C}$ is infinite.*

Proof. Let ζ be an n th root of unity. Let $\lambda_1, \lambda_2, \lambda_3, \infty$ be the ramification points of a double cover $E \rightarrow \mathbb{P}^1$, and let λ denote the cross-ratio of $(\lambda_1, \lambda_2, \lambda_3, \infty)$. Choose $a, b \in \overline{K}$ such that the ordered quadruple $(a, b, \zeta a, \zeta b)$ satisfies

$$\frac{(\zeta a - a)(\zeta b - b)}{(\zeta b - a)(\zeta a - b)} = \lambda$$

This is always possible; for instance, setting $a = 1$, we get a non-trivial quadratic equation for b , and since λ is not 1 or ∞ , we have $b, \zeta b \notin \{a, \zeta a\}$. Thus the elliptic curves

$$X_i : y^2 = (x - \zeta^{i-1}a)(x - \zeta^{i-1}b)(x - \zeta^i a)(x - \zeta^i b), \text{ for } i = 1, \dots, n.$$

all have the same j -invariant as E .

Let $L = K(a, b, \zeta)$. Fix $q \in K$ such that $L(\sqrt[n]{q})$ is a Galois $\mathbb{Z}/n\mathbb{Z}$ -extension of L . We claim that Σ_{ζ} contains the open set

$$U_{\zeta} := \{\sigma \in \text{Gal}(\overline{K}/L) \mid \sigma(\sqrt[n]{q}) = \zeta \sqrt[n]{q}\}.$$

Let $M = L(\sqrt[n]{q})$. For N any number field containing M , let C_N denote the affine curve over N

$$\text{Spec } N[x, y_1, \dots, y_n] / (P_1(x, y_1), \dots, P_n(x, y_n), y_1 \cdots y_n - (x^n - a^n)(x^n - b^n))$$

where

$$P_i(x, y) = y^2 - (x - \zeta^{i-1}a)(x - \zeta^i a)(x - \zeta^{i-1}b)(x - \zeta^i b).$$

Note that the equation $y_1 \cdots y_n - (x^n - a^n)(x^n - b^n) = 0$ merely selects one of the two irreducible components of the 1-dimensional affine scheme cut out by the other equations.

Let X denote the compact Riemann surface which is the compactification of $C_N(\mathbb{C})$. By the Hurwitz genus formula, the genus of X is $(n-2)2^{n-2} + 1$, which is greater than 1 since $n \geq 3$. For any n -tuple $(k_1, \dots, k_n) \in \{0, 1\}^n$ with even sum, the map

$$(2) \quad (x, y_1, \dots, y_n) \mapsto (\zeta x, (-1)^{k_1} \zeta^2 y_n, (-1)^{k_2} \zeta^2 y_1, (-1)^{k_3} \zeta^2 y_2, \dots, (-1)^{k_n} \zeta^2 y_{n-1})$$

defines an automorphism σ of C_N and therefore of X . As the k_i have even sum, σ is of order n . If $x \in \sqrt[n]{q}L^*$ and $\sigma \in U_\zeta$, then $\sigma(x) = \zeta x$, so

$$\sigma(y_i)^2 = \zeta^4 y_{i-1}^2,$$

and so there exists an n -tuple $(k_1, \dots, k_n)$ with even coordinate sum such that σ acts on $Q := (x, y_1, \dots, y_n)$ by (2). By Proposition 3.1, for all but finitely many values of x ,

$$R := \sum_{i=0}^{n-1} \sigma^i(Q) \otimes \zeta^{-i}$$

is a non-zero eigenvector of σ with eigenvalue ζ .

Assume now that N is a finite Galois extension of M . Consider the morphism from C_N to the affine line over M given by $(x, y_1, \dots, y_n) \mapsto x$. This is a branched Galois cover with Galois group $\text{Gal}(N/M) \times (\mathbb{Z}/2\mathbb{Z})^{n-1}$. There exists a Hilbert set of values $t \in M$ such that the geometric points lying over $x = \sqrt[n]{qt}$ in C_M consists of a single $\text{Gal}(\overline{K}/M)$ -orbit or, equivalently, $\text{Gal}(M(y_1, \dots, y_n)/M) \cong (\mathbb{Z}/2\mathbb{Z})^{n-1}$ and $M(y_1, \dots, y_n)$ is linearly disjoint from N over M . As a Hilbert set of a finite extension of L always contains some Hilbert set of L ([5, Ch. 9, Prop. 3.3]), it follows that there exists $t \in L$ such that setting $x = \sqrt[n]{qt}$, relative to M , the extension $M(y_1, \dots, y_n)$ is linearly disjoint from N and has Galois group $(\mathbb{Z}/2\mathbb{Z})^{n-1}$.

We can therefore iteratively construct a sequence $t_1, t_2, \dots \in L^*$ such that the extensions

$$M_i := M \left(\sqrt{(\sqrt[n]{qt_i} - a)(\sqrt[n]{qt_i} - b)(\sqrt[n]{qt_i} - \zeta a)(\sqrt[n]{qt_i} - \zeta b), \dots, \sqrt{(\sqrt[n]{qt_i} - \zeta^{n-1}a)(\sqrt[n]{qt_i} - \zeta^{n-1}b)(\sqrt[n]{qt_i} - a)(\sqrt[n]{qt_i} - b)}} \right)$$

are all linearly disjoint over M . Let Q_i be a point with x -coordinate $\sqrt[n]{qt_i}$, and R_i the corresponding ζ -eigenvector of σ given by (1). We claim that the R_i span a space of infinite dimension. The Q_i do so by [2, Lemma 3.12], and as the ζ^{-j} are linearly independent over $\mathbb{Q}$, it follows that the R_i do so as well. □

We conclude with a question that does not seem to be directly amenable to the methods of this paper:

Question 3.3. *Does the set $\bigcap_{\zeta \in \mathbb{C}_{\text{tor}}^*} \Sigma_{\zeta}$ of elements of G_K having generic spectrum on V_E always have an interior point?*

REFERENCES

- [1] P. Griffiths and J. Harris: *Principles of Algebraic Geometry*, John Wiley & Sons, New York, 1978.
- [2] B. Im: Mordell-Weil groups and the rank over large fields of elliptic curves over large fields, arXiv: math.NT/0411533, to appear in *Canadian J. Math.*
- [3] B. Im: Infinite multiplicity of roots of unity of the Galois group in the representation on elliptic curves, preprint, 2003.
- [4] B. Im: Heegner points and Mordell-Weil groups of elliptic curves over large fields, preprint, arXiv: math.NT/0411534.
- [5] S. Lang: *Fundamentals of Diophantine Geometry*, Springer-Verlag, New York, 1983.
- [6] M. Larsen: *Rigid lattices are Mordell-Weil*, preprint, arXiv: math.NT/0401209.
- [7] M. Raynaud: Courbes sur une variété abélienne et points de torsion. *Invent. Math.* **71** (1983), no. 1, 207–233.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH 84112, USA

E-mail address: im@math.utah.edu

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405, USA

E-mail address: `larsen@math.indiana.edu`