# UNLINKING NUMBER AND UNLINKING GAP

SLAVIK JABLAN, RADMILA SAZDANOVIĆ<sup>†</sup>

The Mathematical Institute Knez Mihailova 35 P.O. Box 367 11001 Belgrade Serbia

jablans@yahoo.com

The George Washington University<sup>†</sup> Department of Mathematics Monroe Hall of Government 2115 G Street, NW Washington, D.C. 20052 USA

radmila@gwu.edu

"It is very easy to define a number of knot invariants so long as one is not concerned with giving algorithms for their computation. For instance,... One can change each knot projection into the projection of a circle by reversing the overcrossing and undercrossing at, say k double points of the projection. The minimum number m(k) of these operations, that is, the minimal number of self-piercings, by which a knot is transformed into a circle, is a natural measure of knottedness." K. Reidemeister [Re]

### ABSTRACT

Computing unlinking number is usually very difficult and complex problem, therefore we define BJ-unlinking number and recall Bernhard-Jablan conjecture stating that the classical unknotting/unlinking number is equal to the BJ-unlinking number. We compute BJ-unlinking number for various families of knots and links for which the unlinking number is unknown. Furthermore, we define BJ-unlinking gap and construct examples of links with arbitrarily large BJ-unlinking gap. Experimental results for BJunlinking gap of rational links up to 16 crossings, and all alternating links up to 12 crossings are obtained using programs LinKnot and K2K. Moreover, we propose families of rational links with arbitrarily large BJ-unlinking gap and polyhedral links with constant non-trivial BJ-unlinking gap. Computational results suggest existence of families of non-alternating links with arbitrarily large BJ-unlinking gap.

Keywords: Unknotting number, unlinking number, unknotting gap, unlinking gap, rational knot, pretzel knot, Conway notation, *n*-move.

Mathematics Subject Classification 2000: 57M25, 57M27

### 1. Introduction

The main topic of this paper is Bernhard-Jablan unlinking number and BJunlinking gap. Computing unlinking number is usually very difficult and complex problem, therefore we define BJ-unlinking number which will be computable due to the algorithmic nature of its definition.

In order to make precise statements we first need to introduce basic notation. The term "link" will be used for both knots and links. Accordingly, under the terms containing the word "link" we consider both knot and link properties/invariants.

We use Conway notation [Co,KL], and the following related symbols denoting rational and pretzel links. Rational link diagrams given by Conway symbols  $a_1 a_2 \ldots a_n$  are denoted by  $C_{[a_1,a_2,\ldots,a_n]}$ .  $C_{[a_1,a_2,\ldots,a_n]}$  is a standard diagram of a rational link denoted by  $R_{[a_1,a_2,\ldots,a_n]}$  which corresponds to the continued fraction

 $a_n + \frac{1}{a_{n-1} + \frac{1}{a_{n-2} + \ldots + \frac{1}{a_2 + \frac{1}{a_1}}}}.$  For a detailed explanation see [KL]. The pretzel

link with n columns of  $a_i$  half-twists each is denoted by  $P_{(a_1,a_2,\ldots,a_n)}$  (Fig. 1).



Fig. 1. Pretzel link  $P_{(a_1,a_2,\ldots,a_n)}$ .

The *BJ*-unlinking gap, (Def. 1.2), is motivated by the following example given by Y. Nakanishi [Na1] and S. Bleiler [Bl]. They have noticed that the rational knot 10<sub>8</sub>, that is  $R_{[5,1,4]}$ , has the unlinking number 2 but its unique minimal diagram  $C_{[5,1,4]}$  (Fig. 2a) has diagram unlinking number 3. Unlinking number 2 can be achieved using non-minimal diagram illustrated in Fig. 2(b) (crossings needed to be changed are denoted by circles). The goal of this paper is to determine how much

#### Unlinking Number and BJUnlinking Gap 3

the unlinking number of any minimal diagram differs from the unlinking number of a link they represent.



Fig. 2. The Nakanishi-Bleiler example: (a) the minimal diagram of the knot 514 that requires at least three simultaneous crossing changes to be unknotted. Switching a particular crossing results in rational knot whose Conway symbol is given next to the crossing. Notice that all of the diagrams obtained are unknotting number 2; (b) non-minimal diagram of the knot 514 from which we obtain the correct unknotting number u(514) = 2.

In this setting we have the following definition:

**Definition 1.1.** For a given crossing v of a diagram D representing link L let  $D_v$  denote the link diagram obtained from D by switching crossing v.

- a) The unlinking number u(D) of a link diagram D is the minimal number of crossing changes on the diagram required to obtain an unlink.
- b) The classical unlinking number of a link L, denoted by u(L) can be defined by  $u(L) = \min_{D} u(D)$  where the minimum is taken over all diagrams Drepresenting L.
- c) The *BJ*-unlinking number  $u_{BJ}(D)$  of a diagram *D* is defined recursively in the following manner:
  - (1)  $u_{BJ}(D) = 0$  iff D represents an unlink.
  - (2) Assume that the sets of diagrams  $\mathcal{D}_k$  with  $u_{BJ}(D) \leq k$  are already defined. A diagram D has  $u_{BJ}(D) = k + 1$  if  $D \notin \mathcal{D}_k$  and there exists a crossing v on the diagram D such that  $u_{BJ}(D'_v) = k$  where  $D'_v$  is a minimal link diagram representing the same link as diagram  $D_v$  obtained from D by a crossing change at v. Notice that  $u_{BJ}(D)$  is well defined for every diagram D as  $D \in \mathcal{D}_n!$  where n is the number of crossings in D.

- 4 Slavik Jablan and Radmila Sazdanović
  - d)  $u_M(L) = \min_D u(D)$  where the minimum is taken over all minimal diagrams D representing L.
  - e) The *BJ*-unlinking number  $u_{BJ}(L)$  of a link *L* is defined by  $u_{BJ}(L) = \min_{D} u_{BJ}(D)$  where the minimum is taken over all minimal diagrams *D* representing *L*.

J.A. Bernhard [Be] in 1994 and independently S. Jablan [Ja] in 1995, conjectured that for every link L we have that  $u(L) = u_{BJ}(L)$ . In the next section we discuss BJ-conjecture and illustrate the importance of the conjecture on the example of pretzel knots whose unlinking number is unknown, except for some small values.

### Definition 1.2.

- a) The *BJ*-unlinking gap of a diagram *D* denoted by  $\delta_{BJ}(D)$  is the difference  $\delta_{BJ}(D) = u(D) u_{BJ}(D)$ .
- b) The *BJ*-unlinking gap of a link *L* denoted by  $\delta_{BJ}(L)$  is defined by  $\delta_{BJ}(L) = u_M(L) u_{BJ}(L)$ .

It is natural to consider  $\delta'_{BJ}(L) = \min_{D} \delta(D)$  where *D* denotes minimal diagram of *L*. For alternating links  $\delta'_{BJ}(L) = \delta_{BJ}(L)$  and it would be interesting to check whether this equality holds in greater generality.

Section 3 contains experimental results– lists of rational knots and links up to 16 crossings that have a non-trivial BJ-unlinking gap. Experimental results imply that knots and links with this property are not so exceptional. In fact, they represent a considerable portion of rational knots and links, e.g., about 4% for n = 15 or n = 16.

In the Section 4 we give explicit formulas for BJ unlinking number,  $u_M$  and BJ unlinking gap for several families of knots (two-bridge and pretzel knots with up to 3 parameters). Moreover, we provide formulas for BJ-unlinking number and BJ-unlinking gap of the family  $R_{[2k,2m,1,2n]}$  and conclude it has an arbitrarily large unlinking gap. Both Sections 4 and 5 contain experimental results: multi-parameter families of alternating rational and polyhedral knots and links with positive BJ-unlinking gap. In the Section 5, based on experimental results, we propose families of non-alternating minimal knot and link diagrams with possibly arbitrarily large BJ-unlinking gap.

## 2. Bernhard-Jablan Conjecture

J. Bernhard [Be] in 1994 and independently S. Jablan [Ja] in 1995, conjectured:

Conjecture 2.1 (Bernhard-Jablan Conjecture). For every link L we have that  $u(L) = u_{BJ}(L)$ .

#### Unlinking Number and BJ Unlinking Gap 5

*BJ*-conjecture holds for all links for which unlinking number is computed. In particular, it holds for all knots up to 11 crossings [Liv] and 2-component links up to 9 crossings [Ko2]. Furthermore, T. Kanenobu, H. Murakami and P. Kohn proved that for unknotting number 1 rational links unknotting crossing appears in the minimal diagram [KM,Ko1].

Notice that, even if the Bernhard-Jablan Conjecture does not hold for all links, BJ-unlinking number is an upper bound for the unlinking number.

We illustrate the importance of the conjecture by the following example. Consider alternating pretzel knots  $P_{(a,b,c)}$  where  $0 < a \leq b \leq c$  and a, b, c are all odd numbers. We show in the Proposition 2.2(b), that  $u_{BJ}(P_{(a,b,c)}) = \frac{a+b}{2}$ . However, the unknotting number of these knots is still unknown, except for the smallest knots, e.g.,  $P_{(1,3,3)}$  (with unknotting number 2 computed by W.B.R. Lickorish [Lic]) and  $P_{(3,3,3)}$  (with unknotting number 3 computed recently by B. Owens [Ow]).



Fig. 3. (a) Rational knot  $R_{[2p-1,1,2q-1]}$ ; (b) Pretzel knot  $P_{(a,b,c)}$ .

## Proposition 2.2.

- a) For 2-bridge knots with Conway type 2p 1 1 2q 1 denoted by  $R_{[2p-1,1,2q-1]}$ (in the fraction form  $\frac{4pq-1}{2q}$ ), p,q > 0 BJ-unknotting number satisfies  $u_{BJ}(R_{[2p-1,1,2q-1]}) = min(p,q)$ .
- b) For pretzel knots  $P_{(a,b,c)}$  with  $0 < a \le b \le c$  and a, b, c are all odd numbers

<sup>a</sup>Pretzel knots  $P_{(a,b,c)}$  are preserved by permutations of symbols a,b,c.

we have that:

$$u_{BJ}(P_{(a,b,c)}) = \frac{a+b}{2}$$

## Proof.

a) We proceed by induction on min(p,q). For min(p,q) = 1 that is, p = 1 or q = 1 our knot is a twist knot  $(R_{[1,1,q]} \text{ or } R_{[p,1,1]})$ . Therefore, it has unknotting number 1, so the proposition holds. Assume that proposition holds for min(p,q) < n for n > 1. Since our link is alternating we can work with the specific minimal diagram (Fig. 3a). We consider every crossing of the diagram. If the crossing is chosen among those representing 2p - 1 or 2q - 1 we obtain either  $R_{[2p-3,1,2q-1]}$  or  $R_{[2p-1,1,2q-3]}$ , then the inductive step is immediate. If we consider the remaining crossing v, after switching at this crossing and using ambient isotopy we get rational knot  $R_{[2(p-1),1,2(q-1)]}$ . In each step of the inductive construction we decrease either p or q or both, therefore we have:

$$u_{BJ}(R_{[2p-1,1,2q-1]}) =$$

$$= 1 + min(u_{BJ}(R_{[2(p-1),1,2q-1]}), u_{BJ}(R_{[2(p-1),1,2(q-1)]}), u_{BJ}(R_{[2p-1,1,2(q-1)]}))$$

$$= 1 + min(p-1,q-1)$$

$$= min(p,q)$$

- b) We proceed by induction on a+b. For a+b=2 our knot  $P_{[1,1,c]}$  is a twist knot. Such a knot has unknotting number 1, so the proposition holds. Assume that proposition holds for a+b < n, n > 2. Since our link is alternating we can work with the specific minimal diagram (Fig. 3b). Assume a = 1 (a+b > 2 implies b, c > 1). We should consider every crossing, so we have the following cases:
  - If the crossing is chosen among those representing c we obtain  $P_{[1,b,c-2]} = P_{[1,c-2,b]}$ , therefore by inductive hypothesis

$$u_{BJ}(P_{[1,b,c-2]}) = \frac{1+b}{2}$$

- If the crossing is chosen among those representing b we obtain  $P_{[1,b-2,c]}$ , so by inductive hypothesis  $u_{BJ}(P_{[1,b-2,c]}) = \frac{b-1}{2}$
- Crossing change at the remaining crossing corresponding to 1 in Conway symbol gives the rational knot  $R_{[b-2,1,c-2]}$ . According to the part a) we have that:

$$u_{BJ}(R_{[1,b-2,c-2]}) = min(\frac{b-1}{2}, \frac{c-1}{2}) = \frac{b-1}{2}$$

To summarize:

$$u_{BJ}(P_{[1,b,c]}) = 1 + \min(u_{BJ}(P_{[1,b,c-2]}), u_{BJ}(P_{[1,b-2,c]}), u_{BJ}(R_{[1,b-2,c-2]}))$$
$$= 1 + \min(\frac{b-1}{2}, \frac{1+b}{2})$$
$$= 1 + \frac{b-1}{2} = \frac{b+1}{2}$$

Unlinking Number and BJUnlinking Gap 7

which is exactly  $\frac{a+b}{2}$  for a = 1.

If a, b > 1 then we have immediate induction, which completes the proof. Since,  $u_M(R_{[2p-1,1,2q-1]}) = u_{BJ}(R_{[2p-1,1,2q-1]})$  and  $u_M(P_{[a,b,c]}) = u_{BJ}(P_{[1,b,c]})$  the BJ-unlinking gap is equal to zero in both cases.

Waldhausen [Wa] has proven Smith's conjecture for double branch covers and we will use it in the following form:

**Theorem 2.3 (Waldhausen).** Double branched covering  $M_K^{(2)}$  of  $S^3$  along a knot K is  $S^3$  if and only if K is a trivial knot (unknot).

**Lemma 2.4 (Montesinos).** If L and m(L) are mutant pairs of links then  $M_L^{(2)}$ and  $M_{m(L)}^{(2)}$  are homeomorphic [Mo, Vi].

**Corollary 2.5.** *Knot* K *is trivial if and only if* m(K) *is trivial.* 

**Lemma 2.6.** If m(D) is a mutation of a diagram D then u(D) = u(m(D)).

**Proof.** Consider an arbitrary 2-tangle T inside a diagram D and a diagram m(D) obtained from D by mutation of T (the remaining part of a diagram is intact). If we make u(D) crossing changes necessary to unknot diagram D and corresponding crossing changes on a diagram m(D), then, by Corollary 2.5, a diagram obtained from m(D) also represents an unknot. Therefore,  $u(D) \leq u(m(D))$ . Since D can be obtained by mutation on m(D) we have that u(D) = u(m(D)).

In the following sections we use Tait's first and third Conjectures. Term minimum diagram stands for minimum crossing number diagram of a link and reduced diagram is a diagram with no nugatory crossings.

## Theorem 2.7.

- a) (Tait's First Conjecture) A reduced alternating diagram is the minimum diagram of its alternating link. Moreover, the minimum diagram of a prime alternating link can only be an alternating diagram. In other words, a non-alternating diagram can never be the minimum diagram of a prime alternating link.
- b) (**Tait's flyping Conjecture**) Two reduced alternating diagrams of the same link, are related by a finite series of flypes.

The first Tait's conjecture [MT1] was proven independently in 1986 by L. Kauffman, K. Murasugi and M. Thistlethwaite [Kau,Mu,Th]. The third Tait's conjecture (Tait's flyping conjecture) was proven by W. Menasco and M. Thistlethwaite [MT1,MT2] and we use it to prove the following Corollary:

**Corollary 2.8.** For every prime alternating link L and its minimal diagrams D and D' the following holds:  $u(D) = u(D') = u_M(L)$ ,  $u_{BJ}(D) = u_{BJ}(D') = u_{BJ}(L)$ ,  $\delta_{BJ}(D) = \delta_{BJ}(D') = \delta_{BJ}(L)$ .

**Proof.** The proof follows from Lemma 2.6 and Tait flyping theorem since flype can be viewed as a special case od mutation.

Corollary 2.8. enables us to compute BJ-unlinking number or BJ-unlinking gap using arbitrary minimal diagram of an alternating link. On the other hand, minimal diagrams of non-alternating links can have different unlinking numbers, for example knot  $14n_{36750}$  discovered by A. Stoimenow [St1].

## 3. Computations of BJ-unlinking gap for knots and links

Experimental results presented in this section are obtained using *Mathematica* based knot theory program *LinKnot* [JS1,JS2,JS3]. For a link given in Conway notation functions **UnKnotLink** and **fGap** compute *BJ*-unlinking number, the unlinking number of its fixed minimal diagram, and *BJ*-unlinking gap  $\delta_{BJ}(L)$ . Unfortunately, these functions are dependent on the function **ReductionKnotLink** [Oc] which sometimes fails in simplifying links. Therefore, for rational links, we use the *LinKnot* function **fGapRat** which is based on the following Theorem of Schubert [Sch].

**Theorem 3.1 (Schubert).** Unoriented rational links  $K(\frac{p}{q})$  and  $K(\frac{p'}{q'})$  are ambient isotopic if and only if:

(1) p = p' and (2) either  $q \equiv q' \pmod{p}$  or  $qq' \equiv 1 \pmod{p}$ 

The following tables contain Conway symbols of rational knots and links up to 16 crossings with a non-trivial BJ-unlinking gap, given according to the number of crossings and whether they are knots or links. Symbols given in bold denote the links with BJ-unlinking gap 2 (others have BJ-unlinking gap 1). The first column in each table gives the number of crossings, second the number of knots or links with non-trivial BJ-unlinking gap, and third column their list.

n	No. of KL's	List of all KL's							
9	1 Link	414							
10	1 Knot				5	14			
	1 Knot				4	142			
11	4 Links		4 3	34 (	314	$5 \ 1 \ 3 \ 2$	5 1	1 1 3	
12	5 Knots		714	534	414	3 6 1	32	61113	3
	7 Knots	4 4	414	6 3	1 4 2	4 1 3	14	51323	2
		2 3 1	1412	51	$1 \ 1 \ 3 \ 2$	513	$1 \ 1 \ 2$		
13		6	16	6	34	8 1	4	515	2
13	16 Links	5 3	332	6	13 3	7 1	32	3413	3 2
		4 1	$4\ 2\ 2$	5 1	$1 \ 1 \ 5$	611	2 3	711	1 3
		2 4 1	1312	41	$1\ 1\ 4\ 2$	611	1 2 2	4211	1 1 3
		7 1 6	73	4	9	14	4	163	4343
		$5 \ 4 \ 1 \ 4$	6 1			5 2		332	7133
		8 1 3 2	331			132		1423	$5\ 1\ 3\ 1\ 4$
14	31 Knot	51323	5 1 4			124		1322	7 1 1 2 3
14	5 Links	81113	313	$1 \ 4 \ 2$	3 1 4	$1 \ 3 \ 2$	3 4	$1 \ 3 \ 1 \ 2$	351113
		$5\ 1\ 1\ 1\ 4\ 2$	$7\ 1\ 1$	1 2 2	2 1 4	$1 \ 3 \ 1 \ 2$	24	11132	4 2 1 1 1 1 4
		$5\ 2\ 1\ 1\ 1\ 1\ 3$							
		$4\ 1\ 4\ 3\ 2$	4 1 6	1 2	4 1 3	$1 \ 3 \ 2$	5 1	$3\ 2\ 1\ 2$	$5\ 1\ 3\ 1\ 1\ 1\ 2$

Among the links with n = 13 crossings we find the first link 616 with the *BJ*-unlinking gap  $\delta_{BJ} = 2$ .

	n =	15	
	43 Kı	nots	
$2\ 1\ 3\ 1\ 4\ 1\ 1\ 2$	$2\ 2\ 4\ 1\ 3\ 1\ 2$	$2\ 4\ 1\ 1\ 1\ 3\ 1\ 2$	$2\ 4\ 1\ 1\ 1\ 4\ 2$
$2\ 4\ 1\ 4\ 2\ 2$	$3\ 2\ 1\ 1\ 6\ 2$	$3\ 3\ 1\ 5\ 1\ 2$	$3\ 3\ 1\ 6\ 2$
$3\ 4\ 1\ 3\ 2\ 2$	$4\ 1\ 1\ 1\ 4\ 2\ 2$	$4\ 1\ 4\ 1\ 1\ 1\ 3$	$4\ 1\ 4\ 1\ 3\ 2$
4 1 4 2 2 2	$4\ 1\ 4\ 3\ 3$	$4\ 1\ 5\ 1\ 4$	4 1 6 1 3
$4\ 1\ 7\ 1\ 2$	$4\ 2\ 1\ 1\ 1\ 1\ 3\ 2$	$4\ 4\ 3\ 4$	$4\ 5\ 1\ 1\ 1\ 3$
$4\ 5\ 1\ 3\ 2$	$4\ 6\ 1\ 4$	$5\ 1\ 1\ 1\ 2\ 1\ 4$	$5\ 1\ 1\ 1\ 3\ 4$
$5\ 1\ 3\ 1\ 1\ 4$	$5\ 1\ 3\ 2\ 1\ 3$	$5\ 1\ 3\ 2\ 4$	$5\ 1\ 4\ 3\ 2$
$5\ 1\ 5\ 2\ 2$	$5\ 1\ 6\ 1\ 2$	$5\ 3\ 3\ 2\ 2$	$6\ 1\ 1\ 1\ 2\ 1\ 1\ 2$
$6\ 1\ 1\ 2\ 3\ 2$	$6\ 1\ 3\ 1\ 1\ 1\ 2$	61314	$6\ 1\ 3\ 2\ 1\ 2$
61332	6144	6162	$6\ 4\ 1\ 4$
7 1 1 1 3 2	713112	8142	
10.1.4	63 Li		0.01.4.1.0.0
10 1 4		$2\ 3\ 1\ 1\ 1\ 4\ 1\ 2$	$2\ 3\ 1\ 4\ 1\ 2\ 2$
2 4 1 5 1 2	25111212	251322	3111522
341313	3 4 1 5 2	3 4 3 3 2	3511122
35133	36132	4111232	41211113
4 1 4 1 2 1 2	41442	4 1 6 2 2	4211313
$\begin{array}{r} 4 & 2 & 1 & 1 & 5 & 2 \\ \hline 4 & 3 & 1 & 1 & 2 & 1 & 3 \end{array}$	$\begin{array}{r} 4 & 2 & 2 & 1 & 1 & 2 & 3 \\ \hline & 4 & 3 & 4 & 2 & 2 \end{array}$	4231113	$\frac{4\ 2\ 4\ 1\ 4}{5\ 1\ 1\ 1\ 2\ 4}$
$     \begin{array}{r}             4 3 1 1 2 1 3 \\             5 1 1 1 2 2 3         \end{array}     $	43422 5111322	441312	
5111223 51513	5111322 52111122	$\frac{5\ 1\ 1\ 1\ 4\ 3}{5\ 2\ 1\ 1\ 1\ 2\ 3}$	$\frac{5\ 1\ 3\ 1\ 5}{5\ 3\ 5\ 2}$
51513 54132	5211122 5532	611142	$\begin{array}{r} 5 \ 5 \ 5 \ 2 \\ \hline 6 \ 1 \ 1 \ 2 \ 5 \end{array}$
61224	$\begin{array}{r} 5 \ 5 \ 5 \ 2 \\ \hline 6 \ 1 \ 3 \ 2 \ 3 \end{array}$	611142 61422	61123 6153
621224 6211113	63123	6333	636
711123	71115	711222	71133
71313	7134	7152	7332
811122	81123	8133	816
834	91113	9132	010
	n =		
	138 K		
10 1 1 1 3	10 1 3 2	11 1 4	$2\ 1\ 4\ 1\ 1\ 1\ 3\ 1\ 2$
$\begin{array}{r} 101113\\214131112\end{array}$	21511212		2314111312 231622
2321112 232111132	241432	2 4 1 6 1 2	251022 2511232
2611222	31113142	31211152	3124132
31311142	3131422	3121102 31411132	31413112
314332	31511122	3214132	3 2 3 1 5 2
3241312	3 2 5 1 1 1 3	33111412	$3\ 3\ 1\ 5\ 1\ 3$
3 3 3 5 2	334132	3 4 1 1 1 4 2	3 4 1 4 2 2
3 4 1 5 1 2	$3\ 4\ 2\ 1\ 1\ 1\ 1\ 3$	$3\ 5\ 1\ 1\ 1\ 2\ 1\ 2$	$3\ 5\ 1\ 3\ 2\ 2$
$3\ 5\ 1\ 5\ 2$	$3\ 5\ 3\ 3\ 2$	$3\ 6\ 1\ 1\ 1\ 2\ 2$	$3\ 6\ 1\ 1\ 2\ 3$
36133	371113	37132	4 1 1 1 4 2 3
4 1 2 1 1 1 1 1 4	$4\ 1\ 3\ 1\ 4\ 3$	$4\ 1\ 4\ 1\ 2\ 1\ 3$	$4\ 1\ 4\ 1\ 4\ 2$
$4\ 1\ 4\ 2\ 1\ 2\ 2$	$4\ 1\ 4\ 2\ 2\ 3$	$4\ 1\ 4\ 2\ 3\ 2$	$4\ 1\ 4\ 3\ 2\ 2$
$4\ 1\ 4\ 4\ 1\ 2$	$4\ 1\ 4\ 4\ 3$	$4\ 1\ 6\ 1\ 2\ 2$	$4\ 1\ 6\ 2\ 3$
4183	$4\ 2\ 1\ 1\ 1\ 1\ 3\ 3$	$4\ 2\ 1\ 1\ 3\ 1\ 4$	$4\ 2\ 2\ 1\ 1\ 2\ 4$
4311214	$4\ 3\ 1\ 6\ 2$	$4\ 3\ 4\ 2\ 3$	$4\ 3\ 6\ 3$
4 4 1 3 1 3	$5\ 1\ 1\ 1\ 1\ 2\ 3\ 2$	$5\ 1\ 1\ 1\ 2\ 2\ 4$	$5\ 1\ 1\ 1\ 4\ 2\ 2$
$5\ 1\ 2\ 1\ 1\ 1\ 1\ 3$	$5\ 1\ 2\ 1\ 1\ 1\ 5$	$5\ 1\ 4\ 1\ 1\ 1\ 3$	$5\ 1\ 4\ 1\ 2\ 1\ 2$
$5\ 1\ 4\ 1\ 3\ 2$	$5\ 1\ 4\ 4\ 2$	$5\ 1\ 5\ 1\ 4$	$5\ 1\ 5\ 2\ 3$
$5\ 2\ 1\ 1\ 3\ 1\ 3$	$5\ 2\ 1\ 1\ 5\ 2$	5221123	$5\ 2\ 3\ 1\ 1\ 1\ 3$
$5\ 2\ 3\ 1\ 5$	$5\ 2\ 4\ 1\ 4$	$5\ 3\ 1\ 1\ 1\ 5$	$5\ 3\ 1\ 1\ 2\ 1\ 3$

# Unlinking Number and BJUnlinking Gap 9

$5\ 3\ 3\ 2\ 3$	$5\ 3\ 4\ 2\ 2$	$5\ 4\ 1\ 3\ 1\ 2$	$5\ 4\ 3\ 4$
$5\ 5\ 1\ 1\ 1\ 3$	$5\ 5\ 1\ 3\ 2$	$5\ 6\ 1\ 4$	$6\ 1\ 1\ 1\ 1\ 2\ 4$
$6\ 1\ 1\ 1\ 2\ 2\ 3$	$6\ 1\ 1\ 1\ 3\ 2\ 2$	$6\ 1\ 1\ 1\ 4\ 3$	$6\ 1\ 1\ 2\ 3\ 3$
$6\ 1\ 2\ 2\ 5$	$6\ 1\ 3\ 1\ 5$	$6\ 1\ 3\ 2\ 1\ 3$	$6\ 1\ 3\ 2\ 4$
$6\ 1\ 3\ 3\ 3$	$6\ 1\ 4\ 2\ 3$	$6\ 1\ 4\ 5$	$6\ 1\ 5\ 1\ 3$
$6\ 1\ 5\ 2\ 2$	$6\ 1\ 6\ 3$	$6\ 2\ 1\ 1\ 1\ 1\ 2\ 2$	$6\ 2\ 1\ 1\ 1\ 2\ 3$
$6\ 2\ 1\ 1\ 6$	$6\ 3\ 1\ 2\ 4$	$6\ 3\ 3\ 2\ 2$	$6\ 3\ 4\ 3$
$7\ 1\ 1\ 1\ 4\ 2$	$7\ 1\ 1\ 3\ 4$	$7\ 1\ 2\ 2\ 4$	$7\ 1\ 3\ 1\ 4$
$7\ 1\ 3\ 2\ 3$	$7\ 1\ 3\ 3\ 2$	$7\ 1\ 4\ 2\ 2$	$7\ 1\ 5\ 3$
$7\ 2\ 1\ 1\ 1\ 3$	$7\ 3\ 1\ 2\ 3$	$7\ 3\ 3\ 3$	$7\ 3\ 6$
$7\ 4\ 1\ 4$	$8\ 1\ 1\ 1\ 2\ 3$	$8\ 1\ 1\ 1\ 5$	8 1 1 2 2 2
8 1 1 3 3	$8\ 1\ 3\ 1\ 3$	$8\ 1\ 3\ 4$	$8\ 1\ 5\ 2$
8332	$9\ 1\ 1\ 1\ 2\ 2$	$9\ 1\ 1\ 2\ 3$	$9\ 1\ 3\ 3$
916	934		
	42 Lin	ks	
$2\ 3\ 1\ 4\ 1\ 3\ 2$	$2\ 3\ 1\ 4\ 3\ 1\ 2$	$2\;3\;1\;7\;1\;2$	$2\ 3\ 5\ 1\ 3\ 2$
$2\ 4\ 1\ 3\ 1\ 1\ 1\ 1\ 2$	$2\ 4\ 1\ 3\ 1\ 2\ 1\ 2$	$3\ 1\ 1\ 1\ 5\ 3\ 2$	$3\ 1\ 1\ 1\ 7\ 1\ 2$
$3\ 4\ 1\ 3\ 1\ 1\ 1\ 2$	$3\ 4\ 1\ 3\ 2\ 1\ 2$	$4\ 1\ 3\ 1\ 5\ 2$	$4\ 1\ 4\ 2\ 2\ 1\ 2$
$4\ 1\ 4\ 5\ 2$	$4\ 1\ 5\ 1\ 1\ 1\ 3$	$4\ 1\ 5\ 2\ 1\ 1\ 2$	41614
$4\ 1\ 6\ 3\ 2$	$4\ 1\ 7\ 1\ 1\ 2$	$4\ 1\ 8\ 1\ 2$	$4\ 3\ 4\ 1\ 4$
$4\ 3\ 4\ 3\ 2$	$4\ 3\ 6\ 1\ 2$	$5\ 1\ 1\ 1\ 2\ 1\ 3\ 2$	$5\ 1\ 1\ 1\ 3\ 3\ 2$
$5\ 1\ 1\ 1\ 5\ 1\ 2$	$5\ 1\ 3\ 1\ 1\ 1\ 4$	$5\ 1\ 3\ 1\ 1\ 3\ 2$	$5\ 1\ 3\ 1\ 2\ 1\ 1\ 2$
$5\ 1\ 3\ 1\ 3\ 1\ 2$	$5\ 1\ 3\ 2\ 1\ 4$	$5\ 1\ 3\ 2\ 3\ 2$	$5\ 3\ 3\ 1\ 1\ 1\ 2$
$5\ 3\ 3\ 2\ 1\ 2$	$6\ 1\ 1\ 1\ 2\ 1\ 1\ 1\ 2$	$6\ 1\ 1\ 1\ 2\ 2\ 1\ 2$	$6\ 1\ 3\ 1\ 3\ 2$
$6\ 1\ 3\ 2\ 1\ 1\ 2$	$6\ 1\ 3\ 3\ 1\ 2$	$6\ 1\ 4\ 3\ 2$	$6\ 1\ 6\ 1\ 2$
$7\ 1\ 3\ 1\ 1\ 2$	$7\ 1\ 3\ 2\ 1\ 2$		
1101112	115212		l.

First rational knots with the non-trivial unknotting gap  $\delta_{BJ} = 2$  are 6163 and 8152 with n = 16 crossings. First non-rational alternating knots with BJunlinking gap  $\delta_{BJ} = 1$  appear for  $n \geq 12$  crossings: the pretzel knot  $P_{(5,4,3)}$  $(12a_{1242})$  and polyhedral knots  $6^*2.40 : 30 (12a_{970}), 6^*2.210 : 40 (12a_{76}), and$  $<math>6^*2.2.2.40 (12a_{1153})$ . In the next section they will be extended to families (see Def. 4.1) with BJ-unlinking gap  $\delta_{BJ} = 1$ .

### 4. BJ-unlinking gap for Some Families of Alternating Links

In this section we explore the effect 2n-moves [Pr] have on the BJ-unlinking number,  $u_M$  and BJ-unlinking gap. Applying 2n-move on an integer tangle decreases or increases its Conway symbol by 2n. If we allow applying 2n-moves on an arbitrary subset of integer tangles of a given link we get its infinite families defined below:

**Definition 4.1.** For a link or knot L given in an unreduced <sup>b</sup> Conway notation C(L) denote by S a set of numbers in the Conway symbol excluding numbers denoting basic polyhedron and zeros (determining the position of tangles in the vertices of polyhedron). For C(L) and an arbitrary (non-empty) subset  $\tilde{S}$  of S the

 $^{\rm b}{\rm The}$  Conway notation is called unreduced if in symbols of polyhedral knots or links elementary tangles 1 in single vertices are included.

#### Unlinking Number and BJUnlinking Gap 11

family  $F_{\tilde{S}}(L)$  of knots or links derived from L is constructed by substituting each  $a \in S_f$  by  $sgn(a)(|a| + 2k_a)$  for  $k_a \in N$ .

J. Bernhard [Be] and D. Garity [Ga] used this approach to obtain general formulas for unlinking numbers of the following families of diagrams of rational knots:  $C_{[(2k+1),1,(2k)]}$   $(k \ge 2)$  and  $C_{[(2k+1),(2l+1),(2k)]}$   $(k \ge 2, l \ge 0, k > l)$  whose unknotting gap is  $\delta(C_{[(2k+1),(2l+1),(2k)]}) = k + l + 1 - (k + l) = 1$ . Moreover, the two-parameter family of rational link diagrams  $C_{[2k,1,2l]}$   $(k \ge 2, l \ge 2)$  [Ga] has  $u_M(C_{[2k,1,2l]}) = k + l - 1$  and  $u(C_{[2k,1,2l]}) \le l$ , so the unlinking gap of a given diagram is at least k - 1 and can be made arbitrarily large for a sufficiently large k.

In the similar manner, we try to obtain explicit formulas for BJ-unlinking gap of the infinite family (with up to k parameters) obtained from a link denoted by its Conway symbol containing k integer tangles. First, we consider rational links containing only 2 or 3 parameters.



Fig. 4. (a) Rational link  $R_{[2m+1,2n+1]}$ ; (b) rational knot  $R_{[2m+1,2n]}$ ; (c) rational knot  $R_{[2m,2n]}$ .

**Lemma 4.2.** Let  $R_{[a,b]}$  denote a 2-bridge link with the Conway symbol a b. Then the following holds:

- (a) If a, b are both odd then for a link  $R_{[a,b]} = R_{[2m+1,2n+1]}$  we have  $u_{BJ}(R_{[2m+1,2n+1]}) = u_M(R_{[2m+1,2n+1]}) = u(R_{[2m+1,2n+1]}) = \frac{a+b}{2} = m+n+1.$
- (b) If a is odd and b is even then for a knot  $R_{[a,b]} = R_{[2m+1,2n]}$  we have  $u_{BJ}(R_{[2m+1,2n]}) = u_M(R_{[2m+1,2n]}) = u(R_{[2m+1,2n+1]}) = n$ .
- (c) If a, b are both even then for a knot  $R_{[a,b]} = R_{[2m,2n]}$  we have  $u_{BJ}(R_{[2m,2n]}) = u_M(R_{[2m,2n]}) = min(m,n)$ .<sup>c</sup>

<sup>c</sup>Unknotting number  $u(R_{[2m,2n]})$  is an open question for the most of knots with m, n > 1.

### Proof.

(a) Notice that switching m + 1 crossings corresponding to 2m + 1 and n crossings corresponding to 2n + 1, as shown in the picture below, leaves us with the unlink of 2 components. Therefore we have:

$$u_M(R_{[2m+1,2n+1]}) \le m+n+1$$

Since every crossing on the diagram is between different components with sign equal to 1 belongs to different components and has the sign equal to 1, then:  $lk(R_{[2m+1,2n+1]}) = m + n + 1$ . Combining these results we have:

$$m + n + 1 = lk(R_{[2m+1,2n+1]}) \le u(R_{[2m+1,2n+1]})$$
$$\le u_{BJ}(R_{[2m+1,2n+1]}) \le u_M(R_{[2m+1,2n+1]})$$
$$\le m + n + 1$$

Therefore  $u(R_{[a,b]}) = u_{BJ}(R_{[a,b]}) = u_M(R_{[a,b]}) = \frac{a+b}{2}$ .

(b) In the same manner as in the proof of part a), using signature (which is equal to b in this case) instead of linking number we get:

$$n = \frac{\sigma(R_{[2m+1,2n]}]}{2} \le u(R_{[2m+1,2n]})$$
  
$$\le u_{BJ}(R_{[2m+1,2n]}) \le u_M(R_{[2m+1,2n]})$$
  
$$\le n.$$

The last inequality is obtained directly from the diagram (switching *n* crossings corresponding to 2n gives the unknot  $R_{[2m+1,0]}$ ) so we have:  $u(R_{[a,b]}) = u_{BJ}(R_{[a,b]}) = u_M(R_{[a,b]}) = \frac{b}{2}$ .

(c) We proceed by induction on min(m, n). If min(m, n) = 0, then we have an unknot, so the proposition holds. Assume that the proposition holds for min(m, n) smaller than fixed positive number p. Now consider min(m, n) = p. We have a choice of switching crossing corresponding to 2m or 2n. As a result we get either  $R_{[2(m-1),2n]}$  or  $R_{[2m,2(n-1)]}$  with min(m-1,n) = min(m,n-1) = p-1 so according to the induction hypothesis  $u_{BJ}(R_{[2m,2n]}) = 1+p-1 = p = min(m,n)$ . Therefore:

$$u(R_{[2m,2n]}) \le u_{BJ}(R_{[2m,2n]}) = u_M(R_{[2m,2n]}) = \min(m,n).$$

**Lemma 4.3.** Let  $R_{[a,b,c]}$  denote a 2-bridge link with the Conway symbol  $a \ b \ c$ . Then the following holds:

- (1) If  $R_{[a,b,c]} = R_{[2k,2l,2m]}$  we have a 2-component link with  $u_{BJ}(R_{[2k,2l,2m]}) = u_M(R_{[2k,2l,2m]}) = u(R_{[2k,2l,2m]}) = k + m.$
- (2) If  $R_{[a,b,c]} = R_{[2k+1,2l+1,2m+1]}$  then for  $(k, m \ge 1)$  has  $u_{BJ}(R_{[2k+1,2l+1,2m+1]}) = u_M(R_{[2k+1,2l+1,2m+1]}) = l + \min(m,k) + 1.$
- (3) If  $R_{[a,b,c]} = R_{[2k+1,2l,2m+1]}$   $(k,m \ge 1)$  then  $u_{BJ}(R_{[2k,2l+1,2m]}) = u_M(R_{[2k,2l+1,2m]}) = u(R_{[2k,2l+1,2m]}) = k + l + m + 1.$

July 16, 2018 7:2 WSPC/INSTRUCTION FILE UnlinkningNumberUnlinkingGap

> Unlinking Number and BJ Unlinking Gap 13

- (4) If  $R_{[a,b,c]} = R_{[2k+1,2l,2m]}$   $(k,m \ge 1)$  then  $u_{BJ}(R_{[2k,2l,2m]}) = u_M(R_{[2k,2l,2m]}) =$  $u(R_{[2k,2l,2m]}) = k + m.$
- $\begin{aligned} &(5) \quad If \ R_{[a,b,c]} = R_{[2k,2l+1,2m]} \ (k,m \ge 1) \ then \\ & u_{BJ}(R_{[2k,2l+1,2m]}) = \begin{cases} l + max(k,m), \ l \le k,m; \\ k+m, & k,m \le l, \end{cases} \\ &i.e., \ u_{BJ}(R_{[2k,2l+1,2m]}) = \min(k+m,\max(k,m)+l) \ and \\ & u_M(R_{[2k,2l+1,2m]}) = \begin{cases} k+m, & 1 \le l \le k,m; \\ k+m-1, \ 0 = l \le k,m; \\ k+m, & k,m \le l. \end{cases} \end{aligned}$

$$\begin{array}{ll} \text{(6)} \ If \ R_{[a,b,c]} = R_{[2k+1,2l+1,2m]} \ (k,m \ge 1) \ then \\ u_{BJ}(R_{[2k+1,2l+1,2m]}) & = & \begin{cases} k+min(l,m), \ l,m \le k \ or \ k,m < l; \\ k+l+1, \ k,l \le m; \end{cases} \\ u_M(R_{[2k+1,2l+1,2m]}) = & \begin{cases} k+min(l,m), \ k,m < l; \\ k+min(l+1,m), \ l,m \le k; \\ k+l+1, \ k,l \le m. \end{cases} \end{array}$$

**Proof.** Cases 1, 3 and 4 are resolved using of linking number or signature while the rest require a detailed analysis (similar to that in the proof of Lemma 4.2) of all possible cases, and will be omitted. 

## Corollary 4.4.

- (a) Links  $R_{[2k,2l+1,2m]}$  have non-trivial gap if  $k, m \geq 2$  and  $m \geq l+1$ . Then  $\delta_{BJ}(R_{[2k,2l+1,2m]}) = \begin{cases} \min(m,k) - 1, \ l = 0; \\ \min(m,k) - l, \ l \ge 1. \end{cases}$ (b) Family  $R_{[2k+1,2l+1,2m]}$  has non-trivial gap  $\delta_{BJ}(R_{[2k+1,2l+1,2m]}) = 1$  if  $m \ge 2$
- and  $l + 1 < m < k^{d}$ .

 $R_{[a]}$  is a torus knot or link of type [2, a] and therefore  $u_{BJ} = u_M = u$ , so both gap and BJ-unlinking gap are trivial. From Lemma 4.2 it follows that all rational links 2 parameters and  $R_{[a,b]}$  have trivial BJ-unlinking gap. The same holds for all 3-parameter families  $R_{[a,b,c]}$  except two families listed in Corollary 4.4. One can try to extend this classification to rational links with more parameters and more complicated generating links but the computations based on parity of parameters and symmetries of the links are very long and tedious so we give only experimental results.

All rational links up to 14 crossings with positive BJ-unlinking gap  $^{\rm e}$  are described by 68 one-parameter<sup>f</sup> families. For all families we predict values of BJunlinking number and BJ-unlinking gap based on computations for links with less

 $<sup>\</sup>overset{\mathrm{d}}{\overset{\mathrm{e}}} u_{BJ}(R_{[2k+1,2l+1,2m]}) = k+l \\ \overset{\mathrm{e}}{\operatorname{Compare}} \text{ with the first table in the Section 3.}$ 

<sup>&</sup>lt;sup>f</sup>One parameter family is obtained by applying the same 2n-move to all chosen integral tangles.

than 48 crossings. Each family in the following table is given by its Conway symbol; the next entry is the number of components followed by experimental results for BJ-unlinking number  $u_{BJ}$  and BJ-unlinking gap  $\delta_{BJ}$ .

	Family	Comp. No.	21 D I	$\delta_{BJ}$
(1)	(2k+2) 1 (2k+2)	2	$\frac{u_{BJ}}{k+1}$	$\frac{0BJ}{k}$
(1) (2)	$\frac{(2k+2)1(2k+2)}{(2k+3)1(2k+2)}$	1	$\frac{k+1}{k+1}$	<u>κ</u> 1
(2) (3)	$\frac{(2k+3)1(2k+2)}{(2k+2)(2k)}$	1	$\frac{k+1}{k+1}$	$\frac{1}{k}$
(4)	$\frac{(2k+2)(2k+2)(2k)}{(2k+2)(2k+1)(2k+2)}$	2	$\frac{n+1}{2k+1}$	1
(5)	$\frac{(2k+2)(2k+2)(2k+2)}{(2k+4)1(2k+2)}$	2	k+2	k
(6)	(2k+3) 1 (2k+1) (2k)	2	k+1	1
(7)	(2k+3) 1 1 1 $(2k+1)$	2	k+1	k
(8)	(2k+5) 1 (2k+2)	1	k+2	1
(9)	(2k+3)(2k+1)(2k+2)	1	2k + 1	1
(10)	(2k+2) 1 $(2k+2)$ $(2k+1)$	1	k+1	k
(11)	(2k+4) 1 (2k+1) (2k)	1	k+1	1
(12)	(2k+4) 1 1 1 $(2k+1)$	1	k+1	1
(13)	(2k+2)(2k+2)1(2k+2)	1	k+1	1
(14)	(2k+4) 1 (2k+2) (2k)	1	k+2	k
(15)	(2k+2) 1 $(2k+1)$ 1 $(2k+2)$	1	k+1	k
(16)	(2k+3) 1 (2k+1) (2k) (2k)	1	k+1	1
(17)	(2k)(2k+1)1(2k+2)1(2k)	1	k+1	1
(18)	(2k+3) 1 1 1 $(2k+1)$ $(2k)$	1	k+1	k
(19)	(2k+4)(2k+1)(2k+2)	2	2k + 2	1
(20)	(2k+6) 1 (2k+2)	2	k+3	k
(21)	$\frac{(2k+3)1(2k+3)(2k)}{(2k+3)(2k+1)(2k+1)(2k)}$	2	k+3	1
(22)	$\frac{(2k+3)(2k+1)(2k+1)(2k)}{(2k+1)(2k+1)(2k+1)}$	2	2k+1	1
(23)	$\frac{(2k+4) 1 (2k+1) (2k+1)}{(2k+5) 1 (2k+1) (2k)}$	2	$\frac{k+1}{3}$	$\frac{1}{1 \text{ if } k = 1}$
(24)	$(2\kappa + 3) = (2\kappa + 1) (2\kappa)$	2	$\frac{3}{k+1}$	$\begin{array}{c}1 \ \text{if } k \equiv 1\\2 \ \text{if } k \geq 2\end{array}$
(25)	(2k+1)(2k+2)1(2k+1)(2k)	2	2k + 1	1
(26)	(2k+2) 1 (2k+2) (2k) (2k)	2	2k + 1	k
(27)	(2k+3) 1 1 1 $(2k+3)$	2	k+2	k
(28)	(2k+4) 1 1 $(2k)$ $(2k+1)$	2	k+1	1
(29)	(2k+5) 1 1 1 $(2k+1)$	2	k+1	k
(30)	(2k)(2k+2)1(2k+1)1(2k)	2	k+1	k
(31)	(2k+2) 1 1 1 $(2k+2)$ $(2k)$	2	k+1	1
(32)	(2k+4) 1 1 1 $(2k)$ $(2k)$	2	k+1 k	1 if $k = 1, 2$ 2 if $k \ge 3$
(33)	(2k+2)(2k) 1 1 1 1 $(2k+1)$	2	2k	1
(34)	(2k+5)(2k+1)(2k+2)	1	2k + 2	1
(35)	(2k+7) 1 (2k+2)	1	k+3	1
(36)	(2k+2) 1 (2k+4) (2k+1)	1	k+2	k
(37)	(2k+2)(2k+1)(2k+2)(2k+1)	1	k+2	1
(38)	(2k+3)(2k+2)1(2k+2)	1	k+1	k
(39)	(2k+4) 1 (2k+2) (2k+1)	1	k+2	k
(40)	(2k+4) 1 $(2k+3)$ $(2k)$	1	k+2	1

(41)	(2k+4)(2k+1)(2k+1)(2k)	1	2k + 1	1
(42)	(2k+5) 1 $(2k+1)$ $(2k+1)$	1	k+1	1
(43)	(2k+6) 1 $(2k+1)$ $(2k)$	1	3	1 if $k = 1$
			k+1	2 if $k \ge 2$
(44)	(2k+1)(2k+1)1(2k+3)(2k)	1	2	1 for $k = 1$
			k+2	0 if $k \ge 2$
(45)	(2k+1)(2k+3)1(2k+1)(2k)	1	k+1	1
(46)	(2k+2) 1 $(2k+2)$ $(2k)$ $(2k+1)$	1	2k + 1	k
(47)	(2k+1) 1 $(2k+1)$ 1 $(2k+2)$	1	k+1	k
(48)	(2k+3) 1 $(2k+1)$ $(2k)$ $(2k+1)$	1	k+1	1
(49)	(2k+3) 1 $(2k+2)$ $(2k)$ $(2k)$	1	2k + 1	1
(50)	(2k+4) 1 1 $(2k)$ $(2k+2)$	1	k+1	1
(51)	(2k+4) 1 $(2k+1)$ $(2k)$ $(2k)$	1	2k	1
(52)	(2k+5) 1 1 $(2k)$ $(2k+1)$	1	2	1 if $k = 1$
			3	0 if $k \ge 2$
(53)	(2k+6) 1 1 1 $(2k+1)$	1	k+2	1
(54)	(2k+1) 1 $(2k+1)$ 1 $(2k+2)$ $(2k)$	1	k+1	k
(55)	(2k+1) 1 $(2k+2)$ 1 $(2k+1)$ $(2k)$	1	k+1	1
(56)	(2k+1)(2k+2)1(2k+1)1(2k)	1	k+1	k
(57)	(2k+1)(2k+3)111(2k+1)	1	k+1	k
(58)	(2k+3) 1 1 1 $(2k+2)$ $(2k)$	1	k+1	1
(59)	(2k+5) 1 1 1 $(2k)$ $(2k)$	1	k+1	1 if $k = 1, 2$
			k	2 if $k \ge 3$
(60)	(2k) 1 (2k+2) 1 (2k+1) 1 (2k)	1	2k	1 if $k = 1, 2$
			2k	$k \text{ if } k \geq 3$
(61)	(2k) (2k+2) 1 1 1 (2k+1) (2k)	1	2k	1
(62)	(2k+2)(2k) 1 1 1 1 1 (2k+2)	1	2k	1
(63)	(2k+3)(2k) 1 1 1 1 $(2k+1)$	1	2k	1
(64)	(2k+2) 1 $(2k+2)$ $(2k+1)$ $(2k)$	2	2k + 1	k
(65)	(2k+2) 1 $(2k+4)$ 1 $(2k)$	2	2k + 1	k
(66)	(2k+2) 1 $(2k+1)$ 1 $(2k+1)$ $(2k)$	2	2k + 1	1
(67)	(2k+3) 1 $(2k+1)$ $(2k)$ 1 $(2k)$	2	2k + 1	1
(68)	(2k+3) 1 $(2k+1)$ 1 1 1 $(2k)$	2	2k + 1	1

#### Unlinking Number and BJUnlinking Gap 15

The following results (unless explicitly stated otherwise) are based on the properties of the generating links and experimental results for rational, pretzel and polyhedral links up to 16 crossings. First, we present several multi-parameter families of rational links with an arbitrarily large BJ-unlinking gap.

- (1) The family  $R_{[(2k),1,(2m),(2n)]}$  has an arbitrarily large BJ-unlinking gap (see Theorem 4.5).
- (2) The family  $R_{[(2k+2),1,(2k+2),(2k-1)]}(k \ge 1)$ , starting with the knot  $R_{[6,1,6,3]}$ , has *BJ*-unlinking number k + 1 and  $\delta_{BJ}(R_{[(2k+2),1,(2k+2),(2k-1)]}) = k$ .
- (3) The family  $R_{[(4(k+1),1,(2k+3),(2k)]}(k \ge 1)$ , starting with the knot  $R_{[8,1,5,2]}$ , has  $u_{BJ}(R_{[(4(k+1),1,(2k+3),(2k)]}) = k+2$  and BJ-unlinking gap is k+1.
- (4) The family  $R_{[(2k+1),1,1,1,(2l+1)]}$ , starting with link  $R_{[5,1,1,1,3]}$ , has the unlinking

gap 
$$\delta_{BJ}(R_{[(2k+1),1,1,1,(2l+1)]}) = \begin{cases} 0, & k = l = 0; \\ l-1, & k = l > 0; \\ l, & k > l > 1. \end{cases}$$
 for  $k \ge l \ge 0$ 

(5) Knots in the family  $R_{[(2k),...,1,...(2k)]}$   $(k \ge 2)$  and every link in the family  $R_{[(2k),...,(2k),1,(2m),...,(2m)]}$   $(k, m \ge 2)$  have members with arbitrarily large BJ-unlinking gaps. If symbol k occurs j times BJ-unlinking number is  $[\frac{j}{2}]k$  and  $\delta_{BJ} = k - 1$ .

We use Lemma 4.2 to prove the following theorem about an example of a family of rational knots with an arbitrarily large BJ-unlinking gap <sup>g</sup>

**Theorem 4.5.** Let  $R_{[2k,2m,1,2n]}$  be a rational knot with diagram  $C_{[2k,2m,1,2n]}$  $(m,k,n \ge 0)$ . Then the following holds:

- a) Diagram unlinking number is  $u_M(R_{[2k,2m,1,2n]}) = n + min(k,m-1)$
- b) BJ-unlinking number is

$$u_{BJ}(R_{[2k,2m,1,2n]}) = \begin{cases} n, & \text{if } m \le n; \\ n + \min(k,m-n), & \text{if } m > n. \end{cases}$$

c) BJ-unlinking gap is

$$\delta_{BJ}(R_{[2k,2m,1,2n]}) = \begin{cases} \min(k,m-1), & \text{if } m \le n; \\ \min(k,m-1) - \min(k,m-n), & \text{if } m > n. \end{cases}$$

Proof.



Fig. 5. Family of rational links  $R_{[2k,2m,1,2n]}$  with arbitrarily large BJ-unlinking gap.

<sup>g</sup>Bleiler asked if  $\delta(L) = u_M(L) - u(L)$  has an upper bound [Bl]. Since  $\delta_{BJ}(L) \leq \delta(L)$ , Theorem 4.5 provides more examples of links with unbounded  $\delta(L)$  (compare [St2]).

Unlinking Number and BJ Unlinking Gap 17

a) The diagram of  $R_{[2k,2m,1,2n]}$  (see Fig. 5) is the minimal as it is reduced alternating [Kau,Mu,Th]. From Corollary 2.8 it follows that it is sufficient to consider only one minimal diagram, so the proof of a) follows from the next lemma:

**Lemma 4.6.** For  $k \ge 0, m > 0, n > 0$ 

(i)  $u(C_{[2k,2m,1,2n]}) = n + min(k,m-1);$ (ii)  $u(C_{[2k,2m,-1,2n]}) = n - 1 + min(k,m-1).$ 

**Proof.** We prove (i) and (ii) simultaneously by induction on k + m + n. For k + m + n=2 and k + m + n = 3 we get:  $u(C_{[0,2,1,2]}) = u(C_{[2,1]}) = 1$ , and  $u(C_{[0,2,-1,2]}) = 0$ , and  $u(C_{[0,4,1,2]}) = u(C_{[2,1]}) = 1$ ;  $u(C_{[0,4,-1,2]}) = 1$  is an unknot,  $u(C_{[0,2,1,4]}) = u(C_{[4,1]}) = 2$ ,  $u(C_{[0,2,-1,4]}) = u(C_{[3]}) = 1$ ,  $u(C_{[2,2,1,2]}) = 1$ , and  $u(C_{[2,2,-1,2]}) = 2$ ,  $u(C_{[0,2,-1,4]}) = u(C_{[3]}) = 1$ ,  $u(C_{[2,2,1,2]}) = 1$ , and  $u(C_{[2,2,-1,2]}) = 0$ , and the lemma holds for k + m + n < p for p > 3. Before we proceed, notice that:  $u(R_{[2k,2,1,2]}) = u_M(R_{[2k,2,-1,2]}) = 1$ ,  $u(R_{[2k,2,-1,2]}) = 0$ , and that signature of  $\sigma(R_{[0,2m,1,2n]}) = 2n$ . Therefore,  $u(R_{[0,2m,1,2n]}) = u_M(R_{[0,2m,1,2n]}) = u(C_{[0,2m,1,2n]}) = u(C_{[0,2m,1,2n]}) = n$  (see Fig. 6). Hence, when k = 0, m = n = 2 lemma holds.



Fig. 6. Family of rational links  $R_{[0,2m,1,2n]}$  with unlinking number n which can be obtained from a minimal projection.

$$\begin{aligned} u(C_{[2k,2m,1,2n]}) &= 1 + \min(\frac{u(C_{[2(k-1),2m,1,2n]}), u(C_{[2k,2(m-1),1,2n]})}{u(C_{[2k,2m,-1,2n]}), u(C_{[2k,2m,1,2(n-1)]}))} \\ &= 1 + \min(\frac{n + \min(k - 1, m - 1), n + \min(k, m - 1), n}{u(C_{[2k,2m,-1,2n]}), n - 1 + \min(k, m - 1))} \\ &= 1 + \min(n - 1 + \min(k, m - 1), u(C_{[2k,2m,-1,2n]})) \end{aligned}$$

$$\begin{split} u(C_{[2k,2m,-1,2n]}) &= 1 + \min(\frac{u(C_{[2(k-1),2m,-1,2n]}), u(C_{[2k,2(m-1),-1,2n]})}{u(C_{[2k,2m,1,2n]}), u(C_{[2k,2m,-1,2(n-1)]}))} \\ &= 1 + \min(\frac{n-1 + \min(k-1,m-1), n-1 + \min(k,m-2),}{u(C_{[2k,2m,1,2n]}), n-2 + \min(k,m-1))} \\ &= 1 + \min(n-2 + \min(k,m-1), u(C_{[2k,2m,1,2n]})) \end{split}$$

From the equations above we get:

$$\begin{aligned} u(C_{[2k,2m,1,2n]}) &= 1 + \min(n-1 + \min(k,m-1), u(C_{[2k,2m,-1,2n]})) \\ &= \min(n + \min(k,m-1), \min(n + \min(k,m-1), 2 + u(C_{[2k,2m,1,2n]}))) \\ &= \min(n + \min(k,m-1), 2 + (C_{[2k,2m,1,2n]})) \\ &= n + \min(k,m-1) \end{aligned}$$

Furthermore:

$$u(C_{[2k,2m,-1,2n]}) = 1 + \min(n-2 + \min(k,m-1), u(C_{[2k,2m,1,2n]}))$$
  
= 1 + min(n - 2 + min(k,m - 1), min(n - 2 + min(k,m - 1)))  
= 1 + n - 2 + min(k,m - 1) = n - 1 + min(k,m - 1)

which completes the proof of the Lemma 4.4 and part a) of the theorem, i.e.,  $u_M(R_{[2k,2m,1,2n]}) = n + \min(k,m-1).$ 



Fig. 7. Family of rational links  $R_{[2k,2m,1,0]} = R_{[2k,2m]}$  with BJ-unlinking number min(m,k).

Unlinking Number and BJ Unlinking Gap 19

- b) We proceed by induction on m + n + k. Using the similar arguments as in Proposition 2.2 and results of the previous lemma we get that the proposition holds for:
  - $u_{BJ}(R_{[0,2,1,2n]}) = u_{BJ}(R_{[1,2n]}) = n$   $- u_{BJ}(R_{[2k,2m,1,0]}) = u_{BJ}(R_{[2k,2m]}) = min(k,m) \text{ (see Fig.7)}$   $- u_{BJ}(R_{[2k,2,1,0]}) = u_{BJ}(R_{[2k,2]}) = 1$  $- u_{BJ}(R_{[2k,0,1,2]}) = u_{BJ}(R_{[2k+1,2]}) = 1$

In particular, the proposition holds for  $m + n + k \leq 3$ . Assume that proposition holds for m + n + k < p for  $p \geq 3$ . Since our link is alternating we can work with the specific minimal diagram (Fig. 3). In the unlinking process we can distinguish 4 cases based on where we perform the crossing change:

- Switch at one of the crossings representing k gives:

$$u_{BJ}(R_{[2(k-1),2m,1,2n]}) = \begin{cases} n, & m \le n; \\ n+min(k-1,m-n), & m > n. \end{cases}$$

- Switch at one of the crossings representing m gives:

$$u_{BJ}(R_{[2k,2(m-1),1,2n]}) = \begin{cases} n, & m \le n+1; \\ n+min(k,m-1-n), & m > n+1. \end{cases}$$

- Switch at one of the crossings representing n gives:

$$u_{BJ}(R_{[2k,2m,1,2(n-1)]}) = \begin{cases} n-1, & m \le n; \\ n-1+\min(k,m-n), & m > n. \end{cases}$$

– Switching the crossing representing 1 to -1 results in  $R_{[2k,2m,-1,2n]} = R_{[2k,2(m-1),1,2(n-1)]}$ , therefore:

$$u_{BJ}(R_{[2k,2(m-1),1,2(n-1)]}) = \begin{cases} n-1, & m \le n-1; \\ n-1+\min(k,m-n+1), & m > n-1. \end{cases}$$

To find  $u_{BJ}(R_{[2k,2m,1,2n]})$  we need to take the minimum over all 4 cases:

$$\begin{split} u_{BJ}(R_{[2k,2m,1,2n]}) &= \\ &= 1 + \min(u_{BJ}(R_{[2k,2(m-1),1,2n]}), u_{BJ}(R_{[2k,2(m-1),1,2(n-1)]}), \\ &u_{BJ}(R_{[2k,2m,1,2(n-1)]}), u_{BJ}(R_{[2k,2(m-1),1,2n]})) \\ &= 1 + \begin{cases} \min(n,n,n-1,n-1), & m < n; \\ \min(n,n,n-1,n-1+\min(k,m-n-1)), & m = n; \\ \min(n+\min(k-1,m-n),n+\min(k,m-n-1)), & m \ge n. \end{cases} \\ &= 1 + \begin{cases} n-1, & m < n; \\ n-1+\min(k,m-n), m > n. \\ n-1+\min(k,m-n), m > n. \end{cases} \\ &= \begin{cases} n, & m \le n; \\ n+\min(k,m-n), m \ge n. \end{cases} \end{split}$$

c) Follows from parts a) and b).

Next we consider the family of pretzel knots  $P_{(a,b,c)}$ . For the pretzel knots  $P_{(2k+1,2l+1,2m+1)}$   $(k \ge l \ge m \ge 1)$  we proved (Proposition 2.2b) that  $u_{BJ}(P_{(2k+1,2l+1,2m+1)}) = l + m$  and  $\delta_{BJ}(P_{(2k+1,2l+1,2m+1)}) = 0$ . For the families of pretzel KLs with three columns we have the following:

## Theorem 4.7.

- (1)  $P_{(2k+1,2l+1,2m+1)}$  has  $u_{BJ}(P_{(2k+1,2l+1,2m+1)}) = u_M(P_{(2k+1,2l+1,2m+1)}) = l + m$ and  $\delta_{BJ}(P_{(2k+1,2l+1,2m+1)}) = 0.$
- (2)  $P_{(2k,2l,2m)}$  has  $u_{BJ}(P_{(2k,2l,2m)}) = u_M(P_{(2k,2l,2m)}) = u(P_{(2k,2l,2m)}) = k + l + m$ , and therefore  $\delta_{BJ}(P_{(2k,2l,2m)}) = 0^{h}$ ;
- (3) For pretzel knots  $P_{(2k+1,2l,2m+1)}$  with  $(k \ge m \ge 1)$  we have i:

$P_{(2k+1,2l,2m+1)}$	$u_{BJ}$	$u_M$	$\delta_{BJ}$
l = 1	m+k	m+k	0
$k \ge l > 1$	m+k	m + k + 1	1
$l > k \ge 1$	m + k + 1	m + k + 1	0

(4)  $P_{(2k,2l+1,2m)}$   $(k \ge m)$  has  $u_{BJ}(P_{(2k,2l+1,2m)}) = u(P_{(2k,2l+1,2m)}) = k+l$ , and gap  $\delta_{BJ}(P_{(2k,2l+1,2m)}) = m-1$ <sup>j</sup>.

Pretzel links  $P_{(2k,2l+1,2m)}$   $(k \ge m)$  are the example of links where BJ-unlinking number and BJ-unlinking gap coincide with unlinking number and unlinking gap (since half signature equals  $u_{BJ}$ ), but not with  $u_M$ , so the gap is non-trivial and grows as we increase parameter m  $(u_M(P_{(2k,2l+1,2m)}) = k + l + m - 1)$ .

# 5. Experimental results and speculations about BJ-unlinking gap for polyhedral and non-alternating links

In this section we give experimental results, which (combined with results from Section 3 for rational links) make computations of the BJ-unlinking gap complete for alternating links up to 12 crossings. Furthermore, we propose the family of non-alternating pretzel link diagrams with an arbitrarily large BJ-unlinking gap.

The first alternating algebraic non-rational link with positive BJ-unlinking gap is the pretzel link 4, 4, 3 with 11-crossings and  $\delta_{BJ} = 1$ , and the remaining seven links with 12-crossings are given in the following table:

<sup>&</sup>lt;sup>h</sup>Notice that the linking number guarantees that  $u_{BJ} = u_M = u$  and  $\delta = \delta_{BJ}$ .

<sup>&</sup>lt;sup>i</sup>Notice that in first two cases, l = 1 and  $k \ge l > 1$ , the signature guarantees that  $u_{BJ} = u$  and  $\delta = \delta_{BJ}$ .

<sup>&</sup>lt;sup>j</sup>Notice that the signature guarantees that  $u_{BJ} = u$  and  $\delta = \delta_{BJ}$ .

Unlinking Number and BJUnlinking Gap 21

22112,2,2	411,3,3	211,31,31	5, 4, 3
31, 31, 21 +	(21, 2111)(2, 2)	(21, 22)1(2, 2)	

Polyhedral knots, defined by Conway [Co], with n = 12 crossings and positive *BJ*-unlinking gap are given in the table below. Second column contains the one-parameter families derived from these knots, followed by the first step of the unknotting process which reduces them to families of rational, pretzel, or polyhedral knots <sup>k</sup>:

No.	Knot	Family	Reduction	$\delta_{BJ}$
1	6*2.31:30	$6^*(2k).31:30$	$6^*(2k).31 1.30$	<u>зы</u>
1	$u_{B,I} = 2$	$u_{B,I} = k$	$k > 1: \approx (2k-1)  1  1  1  2  2$	1
	$u_{BJ} = 2$	$u_{BJ} = \kappa$	$k > 1. \approx (2k - 1) 11122$ $k = 1: \approx 21122$	
	0+0.010.10			
2	$6^*2.210:40$	$6^*(2k).210:40$	$6^*(2k).210:40:-1$	1
		$u_{BJ} = k + 1$	$\approx 31(2k-1)22$	
3	$6^*2.220:30$	$6^*(2k).220:30$	$6^*(2k).220:30:-1$	1
		$u_{BJ} = k + 1$	$\approx 21(2k-1), 21, 2$	
4	$6^*2.40:30$	$6^*2.(2k)0:30 \ (k \ge 2)$	$6^*2.(2k)0:30:-1$	1
	$u_{BJ} = 2$	$u_{BJ} = k$	$\approx (2k)  1  1  2$	
5	$6^{*}2.2.3.30$	$6^*(2k).2.3.30$	$6^*(2k).2.3.30 1$	1
		$u_{BJ} = k + 1$	$\approx 21(2k-1), 21, 2$	
6	$6^{*}2.2.2.40$	$6^*2.2.2.(2k) 0 \ (k \ge 2)$	$6^{*}2.2.2.(2k)01$	1
	$u_{BJ} = 2$	$u_{BJ} = k$	$\approx (2k-1)11112$	
7	$6^{*}2.20.3.30$	$6^*(2k).20.3.30$	$6^*(2k).2.3.30 1$	1
		$u_{BJ} = k + 1$	$\approx 41(2k-1)12$ or	
		-	$6^*(2k).2.3.30:-1$	
			$\approx 21(2k-1), 3, 2$	
8	$6^{*}2.(3,3)$	$6^*(2k).(3,3)$	$6^*(2k).(3,3) :: -1$	1
		$u_{BJ} = k + 1$	$k \ge 2$ : $\approx 6^* 2.(2k - 2) : 20$	
			$k = 1: \approx 21112$	
9	$6^*2.(3,2).2$	$6^*2.(2k+1,2).2$	$6^*2.(2k+1,2).21$	1
		$u_{BJ} = k + 1$	$\approx 221(2k)$	
10	$8^{*}2:2:.30$	$8^*(2k):2:.30$	$8^{*}2:2130$	1
	$u_{BJ} = 2$		$\approx (2k+1) 3 2$ (see Lemma 4.3.)	

Moreover, the following n = 12-crossing links have BJ-unlinking gap  $\delta_{BJ} = 1^{l}$ :

$6^*2.2.2:2110$	$6^*2.2.2:211$	$6^*(21,22)$	$6^*2.(2,2)11$	$6^*2.(2,2), 20$
$6^{*}2.2, (2, 2)0$	$6^{*}2.(2,2).210$	$6^*(2,2).21:2$	$6^{*}(2,2)  1.2:20$	
$6^*211:.(2,2)0$	$6^*211:.(2,2)$	$6^*2.21.2.20:20$	$8^*21:.20:20$	

The question of finding BJ-unlinking gap of non-alternating links is much more difficult because of the lack of classification of their minimal diagrams. For a few classes of non-alternating links partial results can be obtained using the work of W.B.R. Lickorish and M.B. Thistlethwaite [LT]. Unfortunately, this is not sufficient to find all minimal diagrams corresponding to non-alternating link families and compute BJ-unlinking gap for non-alternating links.

<sup>1</sup>In the first 2 rows we give 2-component links and the third row contains 3-component links.

<sup>&</sup>lt;sup>k</sup> The symbol  $\approx$  is used to denote ambient isotopy between two links; for example, in the first row symbol  $\approx$  means that  $6^*(2k).31. - 1.30$  is ambient isotopic to (2k - 1)11122 if k > 1 and 21122 if k = 1.

The following table contains non-alternating KL diagrams with n = 11 and n = 12 crossings. In all cases  $\delta_{BJ} = 1$ :

n = 11	411, 3, -2	32, 3, -3	4, 4, -3	.(3, -2).2	.2.(3,-2)
n = 12	5, -31, 21	-5, 31, 21	(-4, 21)(3, 2)	(-4, -21)(3, 2)	5, -31, 21
	-5, 31, 21	(-3, -3)(3, 21)	(3,3)(-3,21)	3:2:-40	-30.2.20.30

Non-alternating minimal diagrams  $4 \ 1 \ 1, 3, -2 \ \text{and} \ 3 \ 2, 3, -3$  (Fig. 8) of the nonalternating knots  $11n_{64}$  and  $11n_{122}$  [Ho,Liv] have the unknotting gap  $\delta_M = 1$ . These diagrams can be extended to two-parameter families of minimal diagrams  $(2k+2) \ 1 \ 1, (2l+1), (-2m)$  and  $(2k+1) \ 2, (2r+1), -3$  representing Montesionos knots with the diagram unlinking gap  $\delta_M = 1$ .



Fig. 8. The diagrams (a) 411, 3, -2; (b) 32, 3, -3.

As we described before, even minimal diagrams can have a non-trivial unlinking gap. Hence, it is not surprising that some non-minimal diagrams can have a nontrivial unlinking gap.

For example, the 11-crossing non-alternating knot  $11n_{138}$  [Ho,Liv] has the nonminimal diagram 311,3,3- with the unknotting gap  $\delta_M = 1$ , while the (fixed) minimal diagram 311,3,-21 gives the unknotting number u = 2 (Fig. 9).

The family of non-alternating pretzel links  $P_{(2k,-3,2k)}$   $(k \ge 2)$  is the candidate for non-alternating link family with an arbitrarily large unlinking gap (Fig. 10).

This family is obtained from the family of rational links  $R_{[2k,1,2k]} = P_{(2k,1,2k)}$  $(k \ge 2)$  which is a special case of the family  $R_{[2k,2l+1,2m]}$  from the Corollary 4.4a for l = 0, k = m with arbitrarily large BJ-unlinking gap  $\delta_{BJ} = k - 1$ . In the similar manner as in Section 4, we may obtain that the family of standard diagrams of  $P_{(2k,-3,2k)}$  has BJ-unlinking number k. Furthermore, the unlinking number of the standard diagram of  $P_{(2k,-3,2k)}$  is equal to 2k - 1, hence the diagram BJ-unlinking gap is k - 1.



Unlinking Number and BJUnlinking Gap 23

Fig. 9. (a) Non-minimal diagram 311, 3, 3-; (b) the minimal diagram 311, 3, -21 of the non-alternating knot  $11n_{138}$ .



Fig. 10. The family  $P_{(2k,-3,2k)}$  of non alternating minimal diagrams with an arbitrarily large BJ-unlinking gap.

Since the classification of all minimal diagrams of the link family  $P_{(2k,-3,2k)}$  is, up to our knowledge, not yet achieved we are not able to show that the link family  $P_{(2k,-3,2k)}$  has an arbitrarily large unlinking gap.

# Acknowledgments

We would like to express our gratitude to Józef Przytycki for his critical reading of the manuscript, corrections, advice and suggestions.

### References

- [Be] J. A. Bernhard, Unknotting numbers and their minimal knot diagrams, J. Knot Theory Ramifications, 3, 1 (1994) 1–5.
- [Bl] S. A. Bleiler, A note on unknotting number, Math. Proc. Camb. Phil. Soc., 96 (1984) 469–471.
- [Co] J. Conway, An enumeration of knots and links and some of their related properties, in Computational Problems in Abstract Algebra, Proc. Conf. Oxford 1967 (Ed. J. Leech), 329–358, Pergamon Press, New York (1970).
- [Ga] D. Garity, Unknotting Numbers are not Realized in Minimal Projections for a Class of Rational Knots, Proceedings of the "II Italian-Spanish Congress on General Topology and its Applications" (Italian) (Trieste, 1999). Rend. Istit. Mat. Univ. Trieste, **32** (2001), suppl. 2, 59–72 (2002).
- [Ho] J. Hoste and M. Thistlethwaite, *Knotscape*, http://www.math.utk.edu/ ~morwen/
- [Ja] S. Jablan, Unknotting number and ∞-unknotting number of a knot, Filomat, 12, 1, (1998) 113–120.
- [JS1] S. Jablan and R. Sazdanović, LinKnot, http://www.mi.sanu.ac.yu/vismath/linknot/ (2003).
- [JS2] S. Jablan, R. Sazdanovic, LinKnot- Knot Theory by Computer, Series on Knots and Everything, Volume 21, World Scientific Publishing Co., to appear.
- [JS3] S. Jablan and R. Sazdanović, LinKnot, http://math.ict.edu.yu/ (2007).
- [KM] T. Kanenobu, H. Murakami Two-bridge knots with Unknotting Number One, Proceedings of the American Mathematical Society, 98, 3 (1986) 499–502.
- [Kau] L.H. Kauffman, State Models and the Jones polynomial, Topology, 26 (1987) 395– 407.
- [KL] L.H. Kauffman and S. Lambropoulou On the classification of rational tangles Advances in Applied Mathematics, 33, 2 (2004) 199–237 (see arXiv: math.GT/0212011).
- [Ko1] P. Kohn, Two Bridge Links with Unlinking Number One, Proceedings of the American Mathematical Society, 98, 4 (1991) 1135–1147.
- [Ko2] P. Kohn, Unlinking two component links, Osaka J. Math., **30** (1993) 741–752.
- [Lic] W.B.R. Lickorish, The unknotting number of a classical knot, in Combinatorial methods in topology and algebraic geometry (Rocherster, N.Y., 1982), Vol. 44 of Cont. Math., 117–121.
- [LT] W.B.R. Lickorish and M. B.Thislethwaite, Some links with non-trivial polynomials and their crossing-numbers, Comment. Math. Helvetici, 63 (1988) 527–539.
- [Liv] C. Livingston, Knot Tables, http://www.indiana.edu/~knotinfo/ accessed on June 6, 2007.
- [Mo] J.M. Montesions, Surgery on links and doublebranch covers of S<sup>3</sup>, in Knots, groups and 3-manifolds, (Ed. L.P. Neuwrith), Ann. Math. Studies 84, Princeton Univ. Press (1975) 227–259.
- [MT1] W. W. Menasco and M. B. Thistlethwaite, The Tait flyping conjecture, Bull. Amer. Math. Soc., 25, 2 (1991) 403–412.
- [MT2] W. W. Menasco and M. B. Thistlethwaite, The classification of alternating links, Annals of Math., 138 (1993) 113–171.
- [Mu] K. Murasugi, Jones Polynomials and Classical Conjectures in Knot Theory II. Math. Proc. Cambridge Philos. Soc., 102 (1987) 317–318.
- [Na1] Y. Nakanishi, Unknotting numbers and knot diagrams with the minimum crossings, Math. Sem. Notes Kobe Univ., 11 (1983) 257–258.
- [Na2] Y. Nakanishi, Unknotting number and knot diagram, Rev. Mat. Univ. Complut., Madrid, 9 (1996), 2, 359–366.

Unlinking Number and BJUnlinking Gap 25

- [Oc] N. Imafuji and M. Ochiai, Computed aided knot theory using Mathematica and Math-Link, J. Knot Theory Ramifications, 11, 6 (2002) 945–954.
- [Ow] B. Owens, Unknotting information from Heegaard Floer homology, Accepted for publication in Advances in Mathematics; arxiv.org/math.GT/0506485.
- [Pr] J. Przytycki, t<sub>k</sub> moves on links, Contemporary Math., Vol. 78, Braids Proceedings of the Santa Cruz conference on Artin's braid groups (July 1986), 1988, 615-656 (arxiv.org/math.GT/0606633)
- [Re] K. Reidemeister, Knotentheorie. Ergebn. Math. Grenzgeb., Bd.1; Berlin: Springer-Verlag (1932) pg. 25. (English translation: Knot theory, BSC Associates, Moscow, Idaho, USA, 1983).
- [Ro] D. Rolfsen, Knots and Links, Publish or Perish, 1976 (second edition, 1990; third edition, AMS Chelsea Publishing, 2003).
- [Sch] H. Schubert, Knoten mit zwei Brücken, Math. Zeit., 65 (1956) 133–170.
- [St1] A. Stoimenow, On unknotting numbers and knot triviadjacency, Mathematica Scandinavica, 94, 2 (2004) 227–248.
- [St2] A. Stoimenow, On the unknotting number of minimal diagrams, Mathematics of Computation, 72, 244 (2003) 2043–2057.
- [Th] M. B.Thislethwaite A spanning tree expansion for the Jones polynomial, Topology, 26 (1987), 297–309.
- [Vi] O. Ya.Viro Nonprojecting isotopies and knots with homeomorphic coverings, Zap. Nauchn. Semin. LOMI, 66, (1976) 133–147, Russian; English transl. in Journal of Mathematical Sciences, 12, 1, (1979), 86–96.
- [Wa] F. Waldhausen Über Involutionen der 3-Sphäre, Topology, 8 (1969) 81–91.