#### 2000]Primary 32D15, 32D10

# GENERALIZATION OF DRUŻKOWSKI'S AND GONCHAR'S "EDGE-OF-THE-WEDGE" THEOREMS

PETER PFLUG AND VIÊT-ANH NGUYÊN

ABSTRACT. Let  $D, G \subset \mathbb{C}$  be two open sets, let A (resp. B) be a subset of  $\partial D$  (resp.  $\partial G$ ), and let X be the 2-fold cross  $((D \cup A) \times B) \cup (A \times (B \cup G))$ . Suppose in addition that D (resp. G) is *locally rectifiable on* A (resp. B) and that A and B are of positive one-dimensional Hausdorff measure. We determine the "envelope of holomorphy"  $\hat{X}$  of X in the sense that any function locally bounded on X, measurable on  $A \times B$ , and separately holomorphic on  $(A \times G) \cup (D \times B)$  "extends" to a function holomorphic on the interior of  $\hat{X}$ . Generalizations of this result for an N-fold cross are also given.

# 1. INTRODUCTION

The "Edge-of-the-Wedge" type theorems deal with the continuation of holomorphic functions of several complex variables. The first theorem was discovered and proved by N. N. Bogolyubov in 1956 in connection with quantum field theory and dispersion relations. Since then, there is a long list of papers dealing with this theorem and its generalizations under various assumptions (see [16], [20] and the references therein).

Here we consider a one-sided version of the "Edge-of-the-Wedge" type theorem in the spirit of the pioneer work of Malgrange–Zerner [21]. Epstein's survey article [3] gives a historical discussion and motivation for this version of an "Edge-ofthe-Wedge" theorem, as well as its natural connections with theorems on separate analyticity.

The first results in this direction are obtained by Komatsu [10] and Drużkowski [2], but only for some special cases. Recently, Gonchar [5, 6] has proved a remarkable more general result for the one-dimensional case. In a recent work [14], the authors are able to generalize Gonchar's result to the higher dimensional case.

However, in all these cases the hypotheses on the function to extend and its domain of definition are, in some sense, rather restrictive and strong. Therefore, the main goal of our work is to establish one-sided "Edge-of-the-Wedge" type theorems in some more general one-dimensional cases with more optimal hypotheses. Perhaps, this will be the first step towards understanding the higher dimensional case and the manifold settings in its full generality.

This paper is organized as follows.

Date: 1 December 2004.

<sup>1991</sup> Mathematics Subject Classification. [.

Key words and phrases. "Edge-of-the-Wedge" Theorem, Carleman formula, Gonchar–Carleman operator, holomorphic extension, harmonic measure.

In Section 2 we gather some necessary notions and auxiliary results. This preparation will enable us, at the end of this section, to formulate the above mentioned results of Drużkowski and Gonchar and to discuss in more details the motivation for our work.

The statements of our main results as well as an outline of their proofs are given in Section 3.

The tools which are needed for the proof of the main results are developed in Sections 4, 5, and 8.

The proofs of the main results are given in Sections 6, 7, 9, and 10.

Section 11 gives examples showing the optimality of our results. Finally, we conclude the article with some remarks and open questions.

Our approach is based on our previous work [14], the *Gonchar–Carleman operator* developed in [5, 6], conformal mapping theory, and a thorough geometric study of harmonic measures.

Acknowledgment. The paper was written while the second author was visiting the Carl von Ossietzky Universität Oldenburg being supported by The Alexander von Humboldt Foundation. He wishes to express his gratitude to these organizations.

### 2. Preliminaries

In order to recall the classical one-sided versions of the "Edge-of-the-Wedge" theorem and to discuss in more detail our motivation, we need to introduce some notation and terminology. In fact, we keep the main notation from the previous work [14].

2.1. Planar domains with partly rectifiable boundary. We collect here some classical facts from the books by Goluzin, Koosis and Pommerenke ([4], [11], [13]).

A Jordan curve is the image  $\mathcal{C} := \{\gamma(t), t \in [a, b]\}$  of a continuous one-toone map  $\gamma : [a, b] \longrightarrow \mathbb{C}$ , where  $a, b \in \mathbb{R}$ , a < b. The set  $\{\gamma(t), t \in (a, b)\}$  is said to be the *interior* of the Jordan curve. A closed Jordan curve is the image  $\mathcal{C} := \{\gamma(t), t \in [a, b]\}$  of a continuous map  $\gamma : [a, b] \longrightarrow \mathbb{C}$ , which is one-to-one in [a, b) and which satisfies  $\gamma(a) = \gamma(b)$ . The map  $\gamma$  is called a *parametrization* of  $\mathcal{C}$ . Moreover,  $\mathcal{C}$  is said to be *rectifiable* if

$$\sup\left\{\sum_{k=0}^{n-1}|\gamma(t_{k+1})-\gamma(t_k)|\right\}<\infty,$$

where the supremum is taken over all possible positive integers n and sequences of values  $t_0, \ldots, t_n \in [a, b]$  such that  $t_0 < t_1 < \cdots < t_n$ . It is easy to see that this supremum is independent of the choice of a parametrization. It is called the *length* of C. A *(rectifiable) Jordan domain* is a bounded domain in  $\mathbb{C}$  whose boundary is a (rectifiable) closed Jordan curve.

Consider an open set  $D \subset \mathbb{C}$ . Then D is said to be *locally rectifiable at a point*  $\zeta \in \partial D$  if there is a neighborhood U of  $\zeta$  in  $\mathbb{C}$  such that  $U \cap \partial D$  is the interior of a rectifiable Jordan curve. Moreover, D is said to be *locally rectifiable on a subset* A of  $\partial D$  if D is locally rectifiable at all points of A.

 $\mathbf{2}$ 

Now let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a set  $A \subset \partial D$ . We consider the induced topology on  $\partial D$  with respect to the Euclidean topology of  $\mathbb{C}$ . Therefore, there is an open set V in  $\partial D$  such that  $A \subset V$  and D is locally rectifiable on V. Denote by mes the linear measure (i.e. the one-dimensional Hausdorff measure) in  $\mathbb{C}$ . Notice that when A is a Jordan curve, then the length of A coincides with mes(A). We fix the following terminology: A set  $A \subset \mathbb{C}$  is said to be *linearly measurable* if it is measurable with respect to the linear measure.

Let  $\zeta \in \partial D$  be a point with the following property: There is a closed neighborhood V of  $\zeta$  such that  $V \cap \partial D$  is a Jordan curve whose interior contains  $\zeta$ . Then  $\zeta$  is said to be of type I if there is a neighborhood U of  $\zeta$  such that  $U \cap D$  is a Jordan domain. Otherwise,  $\zeta$  is said to be of type II. We see easily that if  $\zeta$  is of type II, then there are an open neighborhood U of  $\zeta$  and two Jordan domains  $U_1$ ,  $U_2$  such that  $U \cap D = U_1 \cup U_2$ .

Let  $\zeta \in \partial D$  be a point of type I or II.  $\zeta$  is said to be *a point where*  $\partial D$  *admits a tangent* if there are a continuous one-to-one map  $\gamma : [a, b] \longrightarrow \partial D$  and a point  $t_0 \in (a, b)$  such that  $\gamma(t_0) = \zeta$  and the following limit exists

$$\lim_{t \to t_0} \frac{\gamma(t) - \gamma(t_0)}{t - t_0} \cdot \frac{|t - t_0|}{|\gamma(t) - \gamma(t_0)|} = \lambda.$$

The tangent line of D at  $\zeta$  is, by definition, the real line  $L := \{t\lambda, t \in \mathbb{R}\}$ . In Proposition 5.6 below we shall see that all the above notions are independent of the choice of a parametrization  $\gamma$ . Let  $\partial^* D$  denote the set of all points  $\zeta \in \partial D$ where  $\partial D$  admits a tangent at  $\zeta$ . Given a linearly measurable set  $A \subset \partial D$  such that D is locally rectifiable on A, then it is well-known (see [11, p. 68–69]) that  $\operatorname{mes}(A \cap \partial^* D) = \operatorname{mes}(A)$ .

We define the concept of angular approach regions at every point of  $\zeta \in \partial^* D$  as follows. Let  $L_{\zeta}$  be the tangent line of D at  $\zeta$  and let  $n_{\zeta}$  be any element of  $L_{\zeta} \setminus \{0\}$ . Then, for any  $0 < \alpha < \frac{\pi}{2}$ , the Stolz region or angular approach region  $\mathcal{A}_{\alpha}(\zeta)$  is given by

$$\mathcal{A}_{\alpha}(\zeta) := \left\{ z \in D : \left| \frac{\pi}{2} - \alpha < \left| \arg\left(\frac{z-\zeta}{n_{\zeta}}\right) \right| < \frac{\pi}{2} + \alpha \right\},\right.$$

where arg :  $\mathbb{C} \longrightarrow (-\pi, \pi]$  is as usual the argument function. Geometrically,  $\mathcal{A}_{\alpha}(\zeta)$  is the intersection of D with two cones of aperture  $2\alpha$  and vertex  $\zeta$ .

Let  $\zeta \in \partial^* D$  and let U be an open neighborhood of  $\zeta$ . We say that a function f defined on  $U \cap D$  admits the angular limit  $\lambda$  at  $\zeta$  if

$$\lim_{z \in \mathcal{A}_{\alpha}(\zeta), \ z \to \zeta} f(z) = \lambda,$$

for all  $0 < \alpha < \frac{\pi}{2}$ .

We conclude this subsection with a simple example which may clarify the above definitions. Let G be the open square whose four vertex are 1 + i, -1 + i, -1 - i, and 1 - i. Define the domain

$$D := G \setminus \left[ -\frac{1}{2}, \frac{1}{2} \right].$$

Then *D* is locally rectifiable on  $\partial G \cup \left(-\frac{1}{2}, \frac{1}{2}\right)$ . Every point of  $\partial G$  is of type I and every point of  $\left(-\frac{1}{2}, \frac{1}{2}\right)$  is of type II. Moreover,

$$\partial^* D = \left(\partial G \setminus \left\{1 + i, -1 + i, -1 - i, 1 - i\right\}\right) \cup \left(-\frac{1}{2}, \frac{1}{2}\right).$$

2.2. Harmonic measure for an open set of  $\mathbb{C}$ . Let D be a proper open subset of  $\mathbb{C} \cup \{\infty\}$  such that the boundary  $\partial D$  (with respect to  $\mathbb{C} \cup \{\infty\}$ ) is non-polar (and A a subset of  $\partial D$ .) Let A be a subset of  $\partial D$ .

Consider the characteristic function

$$1_{\partial D \setminus A}(\zeta) := \begin{cases} 1, & \zeta \in \partial D \setminus A, \\ 0, & \zeta \in A. \end{cases}$$

Then the harmonic measure of the set  $\partial D \setminus A$  (denoted by  $\omega(\cdot, A, D)$ ) is the Perron solution of the generalized Dirichlet problem with boundary data  $1_{\partial D \setminus A}$ . In other words, one has

$$\omega(\cdot, A, D) := \sup_{u \in \widehat{\mathcal{U}}} u,$$

where  $\mathcal{U} = \mathcal{U}(A, D)$  denotes the family of all subharmonic functions u on D such that  $\limsup_{D \ni z \to \zeta} u(z) \leq 1_{\partial D \setminus A}(\zeta)$  for each  $\zeta \in \partial D$ .

It is well-known (see, for example, the book of Ransford [15]) that  $\omega(\cdot, A, D)$  is harmonic on D.

Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a subset A of  $\partial D$ . We say that a point  $\zeta \in \partial^* D$  is a *locally regular point relative to* A if

$$\lim_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} \omega(z, A \cap U, D \cap U) = 0$$

for any  $0 < \alpha < \frac{\pi}{2}$  and any open neighborhood U of  $\zeta$ . Obviously,  $\zeta \in \overline{A}$ . If, moreover,  $\zeta \in A$ , then  $\zeta$  is said to be a *locally regular point of* A. The set of all locally regular points relative to A is denoted by  $A^*$ . Observe that in general  $A^* \not\subset A$ ,  $A \not\subset A^*$ . However, if A is open, then  $A \cap \partial^* D \subset A^*$ .

As an immediate consequence of the Subordination Principle for the harmonic measure (see Corollary 4.3.9 in [15]), one gets

(2.1) 
$$\lim_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} \omega(z, A, D) = 0, \qquad \zeta \in A^*, \ 0 < \alpha < \frac{\pi}{2}.$$

We extend the function  $\omega(\cdot, A, D)$  to  $D \cup A^*$  by simply setting

$$\omega(z, A, D) := 0, \qquad z \in A^*.$$

Geometric properties of the harmonic measure will be discussed in Section 5 below.

2.3. Cross and separate holomorphicity. Let  $N \in \mathbb{N}$ ,  $N \geq 2$ , and let  $D_j$  be a planar domain which is locally rectifiable on a linearly measurable subset  $A_j$  of  $\partial D_j$ ,

j = 1, ..., N. We define an N-fold cross X, its regular part X<sup>\*</sup>, its interior X<sup>o</sup>, its edge A and its regular edge A<sup>\*</sup> as

$$X := \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$$
  

$$:= \bigcup_{j=1}^N A_1 \times \dots \times A_{j-1} \times (D_j \cup A_j) \times A_{j+1} \times \dots \times A_N \subset \mathbb{C}^N$$
  

$$X^* := \mathbb{X}(A_1^*, \dots, A_N^*; D_1, \dots, D_N),$$
  

$$X^\circ = \mathbb{X}^\circ(A_1, \dots, A_N; D_1, \dots, D_N)$$
  

$$:= \bigcup_{j=1}^N A_1 \times \dots \times A_{j-1} \times D_j \times A_{j+1} \times \dots \times A_N,$$
  

$$A := A_1 \times \dots \times A_N, \qquad A^* := A_1^* \times \dots \times A_N^*.$$

Moreover, put

$$\omega(z) := \sum_{j=1}^N \omega(z_j, A_j, D_j), \qquad z = (z_1, \dots, z_N) \in (D_1 \cup A_1^*) \times \dots \times (D_N \cup A_N^*).$$

It is clear that  $\omega|_{D_1 \times \cdots \times D_N}$  is harmonic.

For an N-fold cross  $X := \mathbb{X}(A_1, \ldots, A_N; D_1, \ldots, D_N)$  define its wedge

$$\widehat{X} := \widehat{X}(A_1, \dots, A_N; D_1, \dots, D_N) := \{ z = (z_1, \dots, z_N) \in (D_1 \cup A_1^*) \times \dots \times (D_N \cup A_N^*) : \omega(z) < 1 \}.$$

Then the set of all interior points of the wedge  $\hat{X}$  is given by

$$X^{\circ} := \mathbb{X}^{\circ}(A_1, \dots, A_N; D_1, \dots, D_N)$$
$$:= \{ z = (z_1, \dots, z_N) \in D_1 \times \dots \times D_N : \omega(z) < 1 \}.$$

In particular, if  $A_j$  is an open set of  $\partial D_j$  and  $A_j \subset \partial^* D_j$ ,  $j = 1, \ldots, N$ , one has  $A \subset A^*$  and  $X \subset X^* \subset \widehat{X}$ .

We say that a function  $f: X \longrightarrow \mathbb{C}$  is separately holomorphic on  $X^o$  and write  $f \in \mathcal{O}_s(X^o)$ , if for any  $j \in \{1, \ldots, N\}$  and  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N)$  the function  $f(a', \cdot, a'')|_{D_j}$  is holomorphic on  $D_j$ .

We say that a function  $f: X \longrightarrow \mathbb{C}$  (resp.  $f: A \longrightarrow \mathbb{C}$ ) is separately continuous on X (resp. on A) and write  $f \in \mathcal{C}_s(X)$  (resp.  $f \in \mathcal{C}_s(A)$ ), if for any  $j \in \{1, \ldots, N\}$ and  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N)$  the function  $f(a', \cdot, a'')|_{(D_j \cup A_j)}$ (resp.  $f(a', \cdot, a'')|_{A_j}$ ) is continuous.

In the sequel, for a subset J' of  $\{1, \ldots, N\}$ , we write  $J'' := \{1, \ldots, N\} \setminus J'$ . Moreover, one often identifies  $z \in \mathbb{C}^n$  with (z', z''), where  $z' := (z_j)_{j \in J'}$  and  $z'' := (z_j)_{j \in J''}$ .

In this paragraph, suppose that  $A_j \subset A_j^*$ , j = 1, ..., N. We say that a function  $f: \widehat{X} \setminus (A_1 \times \cdots \times A_N) \longrightarrow \mathbb{C}$  is superholomorphic on  $\widehat{X}$  and write  $f \in \mathcal{O}(\widehat{X})$ , if for

every proper subset J' of  $\{1, \ldots, N\}$  (including  $\emptyset$ ) and any  $a' := (a_j)_{j \in J'} \in \prod_{j \in J'} A_j$ , the restricted function  $f_{a'} : \widehat{\mathbb{X}}^o(A_j, j \in J''; D_j, j \in J'') \longrightarrow \mathbb{C}$ , given by  $f_{a'}(z'') := f(a', z'')$ , where  $J'' := \{1, \ldots, N\} \setminus J'$ , is holomorphic.

For any  $\lambda \in \mathbb{C}$  and r > 0, let  $\mathbb{B}(\lambda, r)$  denote the ball with center  $\lambda$  with radius r. Using identity (2.1) we make the following observation: For any  $a = (a', a'') \in \widehat{X}$  with  $a' = (a_j)_{j \in J'} \in \prod_{j \in J'} A_j^*$  and  $a'' = (a_j)_{j \in J''} \in \prod_{j \in J''} D_j$  and any  $0 < \alpha < \frac{\pi}{2}$ , there is a neighborhood U of a and an  $\epsilon > 0$  such that

$$U \cap \left(\prod_{j \in J'} \mathcal{A}_{\alpha}(a_j) \times \prod_{j \in J''} \mathbb{B}(a_j, \epsilon)\right) \subset \widehat{X}^{\circ}.$$

We say that a function  $f: \widehat{X}^{\circ} \longrightarrow \mathbb{C}$  admits an *angular limit*  $\lambda$  at  $a \in \widehat{X}$  if

$$\lim_{z \to a, \ z \in \prod_{j \in J'} \mathcal{A}_{\alpha}(a_j) \times \prod_{j \in J''} \mathbb{B}(a_j, \epsilon)} f(z) = \lambda, \qquad 0 < \alpha < \frac{\pi}{2}.$$

Throughout the paper, for a subset M of an Euclidean space,  $\mathcal{C}(M)$  denotes the space of all continuous functions  $f : M \longrightarrow \mathbb{C}$  equipped with the sup-norm  $|f|_M := \sup_M |f|$ . Moreover, a function  $f : M \longrightarrow \mathbb{C}$  is said to be *locally bounded* on M if, for any point  $z \in M$ , there are an open neighborhood U of z (with respect to the induced topology on M) and a positive number  $K = K_z$  such that  $|f|_U < K$ . Finally, for an open set  $\Omega \subset \mathbb{C}^n$ ,  $\mathcal{SH}(\Omega)$  (resp.  $\mathcal{PSH}(\Omega)$ ,  $\mathcal{O}(\Omega)$ ) denotes the set of all subharmonic (resp. plurisubharmonic, holomorphic) functions on  $\Omega$ .

2.4. Motivations for our work. We are now able to formulate what, in the sequel, we quote as the *classical one-sided version of the "Edge-of-the-Wedge" theorem*.

**Theorem 1.** (Gonchar [5, 6]) Let  $D_j \subset \mathbb{C}$  be a rectifiable Jordan domain and  $\emptyset \neq A_j$  an open set of the boundary  $\partial D_j$ , j = 1, ..., N. Then, for any function  $f \in \mathcal{C}(X) \cap \mathcal{O}_s(X^o)$ , there is a unique function  $\hat{f} \in \mathcal{C}(\hat{X}) \cap \mathcal{O}(\hat{X})$  such that  $\hat{f} = f$  on X. Moreover, if  $|f|_X < \infty$  then

$$|\widehat{f}(z)| \le |f|_A^{1-\omega(z)} |f|_X^{\omega(z)}, \qquad z \in \widehat{X},$$

where X, X<sup>o</sup>, and  $\widehat{X}$  denote the N-fold cross, its interior and its wedge, respectively, associated to the  $A_j$ ,  $D_j$ .

Theorem 1 admits various generalizations. The following theorem is announced by Gonchar in [5].

**Theorem 2.** Let  $D_j \subset \mathbb{C}$  be a Jordan domain and let  $\emptyset \neq A_j$  be an open set of the boundary  $\partial D_j$  such that  $D_j$  is locally rectifiable on  $A_j$ , j = 1, ..., N. Let f be a function defined on the N-fold cross X with the following properties:

(i)  $f|_{X^o} \in \mathcal{C}(X^o) \cap \mathcal{O}_s(X^o);$ 

(ii) f is locally bounded on X;

- (iii) for any  $j \in \{1, ..., N\}$  there is a function  $f_j$  defined on A such that for any  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N)$ , the holomorphic function  $f(a', \cdot, a'')|_{D_j}$  has the angular limit  $f_j(a', a_j, a'')$  at  $a_j$  for a.e.  $a_j \in A_j$  and  $f_1 = \cdots = f_N = f$  a.e. on A.
- 1) Then there is a unique function  $\hat{f} \in \mathcal{O}(\widehat{X}^o)$  such that

z

$$\lim_{\in \widehat{X}^o, \ z \to \zeta} \widehat{f}(z) = f(\zeta), \qquad \zeta \in X^o.$$

2) If, moreover,  $|f|_X < \infty$ , then

$$|\widehat{f}(z)| \le |f|_A^{1-\omega(z)} |f|_X^{\omega(z)}, \qquad z \in \widehat{X}^o.$$

3) If, moreover, f is continuous at a point  $a \in A$ , then

$$\lim_{z \in \hat{X}^o, \ z \to a} \hat{f}(z) = f(a).$$

On the other hand, the following result due to Drużkowski [2] gives a different flavor.

**Theorem 3.** Let  $D_j \subset \mathbb{C}$  be a rectifiable Jordan domain and let  $\emptyset \neq A_j$  be an open connected set of the boundary  $\partial D_j$ , j = 1, ..., N. Let f be a function defined on X with the following properties:

- (i)  $f \in \mathcal{C}_s(X) \cap \mathcal{O}_s(X^o);$
- (ii) f is locally bounded on X;
- (iii)  $f|_A$  is continuous on A.

Then all conclusions of Theorem 1 still hold.

Observe that all these theorems require the following very strong hypothesis:  $D_1, \ldots, D_N$  are rectifiable Jordan domains and the edge A is an open set of  $\partial D_1 \times \cdots \times \partial D_N$ . Moreover, the assumptions on the boundedness and continuity of f are rather restrictive.

The question naturally arises whether Theorems 1–3 are still true if  $D_1, \ldots, D_N$ are open sets in  $\mathbb{C}$  and the edge A is not necessarily an open set of  $\partial D_1 \times \cdots \times \partial D_N$ . In addition, if one drops the hypothesis on the local boundedness and the continuity of f, can one obtain a holomorphic extension of f and what are its properties? These matters seem to be of interest especially when one seeks to generalize Theorems 1–3 to higher dimensions.

The present paper is motivated by these questions. Our first purpose is to generalize Gonchar's theorems to a very general situation, where  $D_1, \ldots, D_N$  are, in some sense, almost general open subsets of  $\mathbb{C}$  and where the boundary sets  $A_1, \ldots, A_N$ are almost general subsets of  $\partial D_1, \ldots, \partial D_N$ . Our second goal is to establish, in this general context, an extension theorem analogous to Drużkowski's theorem with a minimum of hypothesis on f.

3. Statement of the main results and outline of the proofs

We are now ready to state the first main result.

**Theorem A.** Let  $D_j \subset \mathbb{C}$  be an open set and  $A_j$  a linearly measurable subset of  $\partial D_j$  such that  $D_j$  is locally rectifiable on  $A_j$ ,  $\operatorname{mes}(A_j) > 0$ ,  $j = 1, \ldots, N$ . Let  $f : X \longrightarrow \mathbb{C}$  be such that:

- (i) f is locally bounded on X and  $f \in \mathcal{O}_s(X^o)$ ;
- (ii)  $f|_A$  is measurable with respect to the N-dimensional Hausdorff measure on A;
- (iii) for any  $j \in \{1, ..., N\}$ , there is a function  $f_j : A \longrightarrow \mathbb{C}$  such that, for any  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N)$ , the holomorphic function  $f(a', \cdot, a'')|_{D_j}$  (see (i)) has the angular limit  $f_j(a', a_j, a'')$  at  $a_j$  for a.e.  $a_j \in A_j$  and  $f_1 = \cdots = f_N = f$  a.e. on A.

Then there exists a unique function  $\hat{f} \in \mathcal{O}(\widehat{X}^o)$  with the following property:

- 1) There are subsets  $\tilde{A}_1 \subset A_1 \cap A_1^*, \ldots, \tilde{A}_N \subset A_N \cap A_N^*$  such that
- 1a)  $mes(A_j \setminus \tilde{A}_j) = 0, \ j = 1, \dots, N;^1$
- 1b)  $\hat{f}$  can be extended to a function (still denoted by)

$$\hat{f} \in \mathcal{O}\Big(\widehat{\mathbb{X}}(\tilde{A}_1,\ldots,\tilde{A}_N;D_1,\ldots,D_N)\Big)$$

which admits the angular limit  $\hat{f}(a)$  at every point

$$a \in \widehat{\mathbb{X}}(\widetilde{A}_1, \ldots, \widetilde{A}_N; D_1, \ldots, D_N) \setminus A;$$

1c)  $\hat{f} = f$  on  $\mathbb{X}(\tilde{A}_1, \ldots, \tilde{A}_N; D_1, \ldots, D_N) \setminus A$ .

In addition,  $\hat{f}$  enjoys the following properties: 2) If  $|f|_X < \infty$ , then

$$|\widehat{f}(z)| \le |f|_A^{1-\omega(z)} |f|_X^{\omega(z)}, \qquad z \in \widehat{X}^o.$$

3) For any  $j \in \{1, \ldots, N\}$  and  $a^0 = (a', z_j^0, a'') \in A_1^* \times \cdots \times A_{j-1}^* \times D_j \times A_{j+1}^* \times \cdots \times A_N^*$ , if  $\lim_{a \to a^0, a \in X} f(a)$  exists, then  $\hat{f}$  admits the angular limit  $\lim_{a \to a^0, a \in X} f(a)$  at  $a^0$ .

4) For any  $a^0 \in A^*$ , if  $\lim_{a \to a^0, a \in A} f(a)$  exists, then  $\hat{f}$  admits the angular limit  $\lim_{a \to a^0, a \in A} f(a)$  at  $a^0$ .

5) If  $f|_A$  can be extended to a continuous function defined on  $A^*$ , then f can be extended to a unique continuous function (still denoted by) f defined on  $X^* := \mathbb{X}(A_1^*, \ldots, A_N^*; D_1, \ldots, D_N)$  and  $\hat{f}$  admits the angular limit f(a) at every  $a \in X^*$  and  $f_1 = \cdots = f_N = f$  on  $A \cap A^*$ .

Theorem A has an immediate consequence.

**Corollary A'.** We keep the hypotheses and the notation of Theorem A. Suppose in addition that  $f \in \mathcal{C}(X^o)$ . Then there exists a unique function  $\hat{f} \in \mathcal{O}(\hat{X}^o)$  with the following property:

<sup>&</sup>lt;sup>1</sup> Under this condition it follows from Lemma 5.7 and Part 1) of Proposition 5.9 below that  $\tilde{A}_j \subset \tilde{A}_i^*, j = 1, \ldots, N.$ 

1a')  $\hat{f}$  can be extended to a function (still denoted by)

$$\hat{f} \in \mathcal{O}\Big(\widehat{\mathbb{X}}(A_1 \cap A_1^*, \dots, A_N \cap A_N^*; D_1, \dots, D_N)\Big),$$

which admits the angular limit f(a) at every point

$$a \in \widetilde{\mathbb{X}} (A_1 \cap A_1^*, \dots, A_N \cap A_N^*; D_1, \dots, D_N) \setminus A;$$

1b')  $\hat{f} = f \text{ on } (X \cap X^*) \setminus A.$ 

It is worthy to note that Theorem A and Corollary A' generalize, in some sense, Theorems 1–3.

Now we drop the hypothesis on local boundedness and continuity of f. Then the examples of Drużkowski in [2] (see Section 11 below) show that, without these conditions, the extended function  $\hat{f}$  (if it does exist) is, in general, not continuous on  $\hat{X}$ . However, our second main result gives a partially positive answer to this question.

**Theorem B.** Let  $D_j \subset \mathbb{C}$  be an open set and  $A_j$  a linearly measurable subset of  $\partial D_j$  such that  $D_j$  is locally rectifiable on  $A_j$ ,  $\operatorname{mes}(A_j) > 0$ , and  $A_j \subset \partial^* D_j$ ,  $j = 1, \ldots, N$ . Let  $f \colon X \longrightarrow \mathbb{C}$  satisfy the following properties:

- (i)  $f|_A \in \mathcal{C}_s(A)$  and  $f \in \mathcal{O}_s(X^o)$ ;
- (ii) for any  $j \in \{1, \ldots, N\}$  and for any  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N)$ , the function  $f(a', \cdot, a'')$  is locally bounded on  $D_j \cup A_j$  and the (holomorphic) restriction function  $f(a', \cdot, a'')|_{D_j}$  has the angular limit  $f(a', a_j, a'')$  at  $a_j$  for every  $a_j \in A_j$ .

Then there are subsets  $\tilde{A}_1 \subset A_1 \cap A_1^*, \ldots, \tilde{A}_N \subset A_N \cap A_N^*$ , and a unique function  $\hat{f} \in \mathcal{O}(\hat{X}^o)$  with the following properties:

- 1) mes $(A_j \setminus \tilde{A}_j) = 0, j = 1, \dots, N;$
- 2)  $\hat{f}$  can be extended to a function (still denoted by)

$$\hat{f} \in \mathcal{O}\left(\widehat{\mathbb{X}}\left(\tilde{A}_1, \ldots, \tilde{A}_N; D_1, \ldots, D_N\right)\right)$$

which admits the angular limit  $\hat{f}(a)$  at every point

$$a \in \widehat{\mathbb{X}}\left(\widetilde{A}_1, \ldots, \widetilde{A}_N; D_1, \ldots, D_N\right);$$

3)  $\hat{f} = f \text{ on } \mathbb{X}\left(\tilde{A}_1, \ldots, \tilde{A}_N; D_1, \ldots, D_N\right).$ 

Observe that if  $f \in \mathcal{C}_s(X) \cap \mathcal{O}_s(X^\circ)$ , then conditions (i)–(ii) above are fulfilled. Below we give some ideas how to prove Theorems A and B.

Our method consists of two steps. In the first step we suppose that each  $D_j$  is a Jordan domain, j = 1, ..., N. In the second one we treat the general case. The key technique here is to use *level sets* of the harmonic measure. More precisely, we exhaust each  $D_j$  by the level sets of the harmonic measure  $\omega(\cdot, A_j, D_j)$ , i.e. by  $D_{j,\delta} := \{z_j \in D_j : \omega(z_j, A_j, D_j) < 1 - \delta\} \ (0 < \delta < 1).$ 

#### PETER PFLUG AND VIỆT-ANH NGUYÊN

In order to carry out the first step, we improve Gonchar's method [5, 6] and make intensive use of Carleman's formula and of geometric properties of the level sets of harmonic measures.

In the second step we apply some mixed cross type theorems (see [14]) in order to prove Theorems A and B with  $D_j$  replaced by  $D_{j,\delta}$ . Then we construct the solution for the original domains  $D_j$  by means of a gluing procedure.

Although our results have been stated for the general case  $N \ge 2$ , the proofs will be presented only in the case N = 2. At the end of Section 9 we will give the ideas how to get the general results.

#### 4. GONCHAR-CARLEMAN OPERATOR

In this section we reformulate a result due to Gonchar [5, 6] to our context. This result will play an important role for the proof of Theorems A and B.

Let  $D_j$  be a rectifiable Jordan domain and let  $A_j$  be a linearly measurable subset of  $\partial D_j$  such that  $\operatorname{mes}(A_j) > 0$ ,  $j = 1, \ldots, N$ . Let f be a function defined on  $X := \mathbb{X}(A_1, \ldots, A_N; D_1, \ldots, D_N)$  with the following properties:

- (i)  $f|_A$  is measurable;
- (ii)  $f \in \mathcal{O}_s(X^{\mathrm{o}});$
- (iii) there exist a constant C > 0 and N functions  $f_j : A \longrightarrow \mathbb{C}, j = 1, ..., N$ , such that for any  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N), |f(a', \cdot, a'')|_{D_j} < C$ , and  $f(a', \cdot, a'')$  has the angular limit  $f_j(a', a_j, a'')$  at  $a_j$  for a.e.  $a_j \in A_j$ , and  $f_1 = \cdots = f_N = f$  a.e. on A.

For j = 1, ..., N, let  $\tilde{\omega}_j$  be the conjugate function of  $\omega(\cdot, A_j, D_j)$  such that  $\tilde{\omega}_j(z_j^0) = 0$  for a certain fixed point  $z_j^0 \in D_j$ . Therefore, we can define the holomorphic functions  $g_j(z_j) := \omega(z_j, A_j, D_j) + i\tilde{\omega}_j(z_j), j = 1, ..., N$ , and

$$g(z) := \sum_{j=1}^{N} g_j(z_j), \qquad z = (z_1, \dots, z_N) \in D_1 \times \dots \times D_N.$$

Each function  $e^{-g_j}$  is bounded on  $D_j$ , j = 1, ..., N. Therefore, in virtue of [4, p. 439], we may define  $e^{-g_j(a_j)}$  for a.e.  $a_j \in A_j$  to be the angular boundary limit of  $e^{-g_j}$  at  $a_j$ .

In virtue of (i), for each positive integer M, we define the Gonchar-Carleman operator as follows

(4.1) 
$$K_M(z) = K_M[f](z) := \frac{1}{(2\pi i)^N} \int_A e^{-M(g(a) - g(z))} \frac{f(a)da}{a - z}, \ z \in D_1 \times \dots \times D_N,$$

where  $da := da_1 \dots da_N$ ,  $a - z := (a_1 - z_1) \cdots (a_N - z_N)$ .

Moreover, we can extend  $K_M$  to  $(D_1 \cup A_1) \times \cdots \times (D_N \cup A_N)$  in the following way. Let  $z = (z_1, \ldots, z_N) \in (D_1 \cup A_1) \times \cdots \times (D_N \cup A_N)$  and J' the set of  $\{ j : 1 \le j \le N, z_j \in A_j \}$ . Set  $J'' := \{1, \ldots, N\} \setminus J'$ . In the sequel one often identifies z

with (z', z''), where  $z' := (z_j)_{j \in J'}$  and  $z'' := (z_j)_{j \in J''}$ . Then we define

(4.2) 
$$K_{M_1,\dots,M_N}(z) = K_{M_1,\dots,M_N}[f](z) := \begin{cases} f(z), & J'' = \emptyset, \\ \frac{1}{(2\pi i)^N} \int_{A''} e^{-\sum_{j \in J''} M_j(g_j(a_j) - g(z_j))} \frac{f(z',a'')da''}{a'' - z''}, & J'' \neq \emptyset, \end{cases}$$

where  $M_1, \ldots, M_N \in \mathbb{N}, A'' := \prod_{j \in J''} A_j, da'' := \prod_{j \in J''} da_j$ , and  $a'' - z'' := \prod_{j \in J''} (a_j - z_j)$ . Moreover, we will write simply  $K_M(z)$  in place of  $K_{M,\dots,M}(z)$  for any  $M \in \mathbb{N}$ . This notation is in accordance with (4.1).

The following Carleman Theorem due to Goluzin and Krylov (see, for example, [1, p. 2]) will be very useful.

**Theorem 4.1.** Let D be a rectifiable Jordan domain and A a linearly measurable subset of  $\partial D$  such that  $\operatorname{mes}(A) > 0$ . Then for any function  $f \in H^{\infty}(D)$ , any relatively compact subset  $V \subseteq D$ , and any  $\epsilon > 0$ , there exists a positive integer  $M_0$ (depending only on the sets D, A, V and the number  $||f||_{H^{\infty}(D)}$ ) such that

$$|f(z) - K_M[f|_A](z)| < \epsilon, \qquad z \in V, \ M \ge M_0,$$

where  $K_M$  is given by (4.1) for N = 1 and  $f|_A$  is the angular boundary limit of f on A. (Notice that by [4, p. 439],  $f|_A$  is linearly measurable).

*Proof.* For every  $z \in D$ , applying the Cauchy formula to the function  $w \mapsto$  $e^{-M(g(w)-g(z))}f(w) \in H^{\infty}(D), M \in \mathbb{N}$ , we obtain

$$f(z) = \frac{1}{2\pi i} \int_{A} e^{-M(g(a) - g(z))} \frac{f(a)da}{a - z} + \frac{1}{2\pi i} \int_{\partial D \setminus A} e^{-M(g(a) - g(z))} \frac{f(a)da}{a - z}.$$

Since  $|e^{-(g(a)-g(z))}| < 1$  for a.e.  $a \in \partial D \setminus A$ , the theorem follows.

**Theorem 4.2.** Let  $D_i$  be a rectifiable Jordan domain and  $A_i$  a linearly measurable subset of  $\partial D_j$  such that  $\operatorname{mes}(A_j) > 0, \ j = 1, \ldots, N$ . Let f be a function defined on X which satisfies (i)-(iii) above. Then 1) the following limit

$$K(z) = K[f](z) := \lim_{M \to \infty} K_M(z)$$

exists for all  $z \in \widehat{X} \cap (D_1 \cup A_1) \times \cdots \times (D_N \cup A_N)$ , and its limit is uniform on compact subsets of  $\widehat{X}^{o}$ ; 2) there is a finite constant  $C_0$  such that

$$|K(z)| \leq \frac{C_0 C}{\prod_{j \in J''} \operatorname{dist}(z_j, \partial D_j)(1 - e^{-(1 - \omega(z))})}, \qquad z \in \widehat{X} \cap (D_1 \cup A_1) \times \dots \times (D_N \cup A_N),$$

where  $\operatorname{dist}(z_j, \partial D_j) := \inf_{\zeta_j \in \partial D_j} |z_j - \zeta_j|$  and C is the constant given in (iii) above.

*Proof.* We briefly recall the argument of Gonchar in [6]. Write the difference  $K_{M+1} - K_M$  as follows:

(4.3) 
$$K_{M+1} - K_M = \sum_{j=1}^N K_M^j,$$

where

$$K_M^j := K_{M+1,\dots,M+1,\underbrace{M,\dots,M}_{j-1}} - K_{M+1,\dots,M+1,\underbrace{M,\dots,M}_{j}}, \qquad j = 1,\dots,N.$$

Reasoning as in formulas (6)–(9) in [6] and taking (iii) into account, we see that there is a constant  $C_0$  such that

(4.4) 
$$|K_{M+1}(z) - K_M(z)| \le \sum_{j=1}^N |K_M^j(z)| \le \frac{CC_0}{\prod_{j \in J''} \operatorname{dist}(z_j, \partial D_j)} e^{-M(1-\omega(z))}$$

for  $z \in \widehat{X} \cap (D_1 \cup A_1) \times \cdots \times (D_N \cup A_N)$ . Since  $|e^{-(1-\omega(z))}| < 1$ , the desired conclusion of Part 1) follows immediately from this estimate. Finally, using (4.4), Part 2) follows.

The following version of Privalov's Uniqueness Theorem will be also needed.

**Theorem 4.3.** Let  $D \subset \mathbb{C}$  be a domain which is locally rectifiable on a linearly measurable subset A of  $\partial D$  with  $\operatorname{mes}(A) > 0$ . Let f be a holomorphic function on D such that the angular limits of f on the set  $A \cap \partial^* D$  are equal to 0. Then  $f \equiv 0$ .

*Proof.* Using the hypothesis, one may find a rectifiable Jordan domain  $U \subset D$  such that  $\operatorname{mes}(A \cap \partial U) > 0$  and  $f|_U$  admits the angular limit 0 at almost every point of  $A \cap \partial U$ . By Privalov's Uniqueness Theorem (see [11]),  $f|_U \equiv 0$ . Hence,  $f \equiv 0$ .

Finally, the following result will play a key role in the proof of Theorems A and B.

**Lemma 4.4.** Let D be a rectifiable Jordan domain,  $(f_n)_{n=0}^{\infty}$  a sequence of holomorphic functions on D, and  $\Delta$  a linearly measurable subset of  $\partial D$  with  $\operatorname{mes}(\Delta) > 0$ . Assume that  $f_n$  admits the angular limit (denoted by)  $f_n(t)$  at every point  $t \in \Delta$ ,  $n \in \mathbb{N}$ , <sup>2</sup> that  $\lim_{n \to \infty} f_n(t) = f_0(t)$ ,  $t \in \Delta$ , and that  $\sup_{n\geq 0} |f_n|_D < \infty$ . Then  $(f_n)_{n=1}^{\infty}$  converges uniformly on every compact subset of D to  $f_0$ .

*Proof.* Let

$$g(t,\Delta,D):=\omega(t,\Delta,D)+i\widetilde{\omega}(t,\Delta,D),\quad t\in D,$$

where  $\widetilde{\omega}(\cdot, \Delta, D)$  is the conjugate harmonic function of  $\omega(\cdot, \Delta, D)$ . Since  $g \in \mathcal{O}(D)$  is bounded, it follows from [4, p. 439] that for a.e.  $t \in \partial D$ , g admits an angular limit (denoted by g(t)) at t. Fix an arbitrary compact subset H of D. Recall that

<sup>&</sup>lt;sup>2</sup> In our notation  $\mathbb{N} = \{0, 1, 2, ...\}$ .

 $|f_n|_D < M < \infty, n \in \mathbb{N}$ . Consequently, applying Theorem 4.1 we see that, for any  $\epsilon > 0$ , there is an  $M_{\epsilon}$  such that, for any  $M \ge M_{\epsilon}$ ,

(4.5) 
$$\left| f_n(\tau) - \frac{1}{2\pi i} \int_{\Delta} e^{-M(g(t,\Delta,D) - g(\tau,\Delta,D))} \frac{f_n(t)dt}{t - \tau} \right| < \epsilon, \ \tau \in H, \ n \in \mathbb{N}$$

On the other hand, using that  $\lim_{n\to\infty} f_n(t) = f_0(t), t \in \Delta$ , and applying Lebesgue's Dominated Convergence Theorem, we see that

$$\lim_{n \to \infty} \frac{1}{2\pi i} \int\limits_{\Delta} e^{-M(g(t,\Delta,D) - g(\tau,\Delta,D))} \frac{f_n(t)dt}{t - \tau} = \frac{1}{2\pi i} \int\limits_{\Delta} e^{-M(g(t,\Delta,D) - g(\tau,\Delta,D))} \frac{f_0(t)dt}{t - \tau}$$

for all  $\tau \in H$  and  $M > M_{\epsilon}$ . This, combined with (4.5), implies that  $\lim_{n \to \infty} f_n(\tau) = f_0(\tau), \tau \in H$ . Hence, by the Montel Theorem the sequence  $(f_n)_{n=1}^{\infty}$  converges uniformly on compact subsets of D to  $f_0$ . This completes the proof.

# 5. Properties of the harmonic measure and some elements of Conformal mapping theory

In this section we develop the tools needed for the proofs of Theorems A and B. In the sequel,  $D \subset \mathbb{C}$  is an open set and A is a linearly measurable nonempty subset of  $\partial D$  such that D is locally rectifiable on A. Observe that under the above assumption  $\partial D$  is non-polar. Let  $\partial^* D$  be the set of all points  $\zeta \in \partial D$  where  $\partial D$ admits a tangent. Recall from Subsection 2.1 that  $\operatorname{mes}(A \cap \partial^* D) = \operatorname{mes}(A)$ .

Let  $\mathcal{P}_D$  be the generalized Poisson integral of D. If, in addition, A is a Borel set, then, by Theorem 4.3.3 of [15], the harmonic measure of  $\partial D \setminus A$  is given by

(5.1) 
$$\omega(\cdot, A, D) = \mathcal{P}_D[1_{\partial D \setminus A}].$$

Next, let E denote the unit disc of C. Then, for an  $f \in L^1(\partial E)$ , a point  $\zeta \in \partial E$  is said to be a *Lebesgue point* of f if

$$\lim_{r \to 0} \frac{1}{\max\left(\partial E \cap \mathbb{B}(\zeta, r)\right)} \int_{\partial E \cap \mathbb{B}(\zeta, r)} |f(\theta) - f(\zeta)| d\theta = 0,$$

where  $d\theta$  is the linear measure defined on  $\partial E$ . If A is a linearly measurable subset of  $\partial E$  and  $f := 1_A$ , then every point of A that is a Lebesgue point of f is called a *density point* of A.

**Proposition 5.1.** 1) Let  $f \in L^1(\partial E)$ . Then a.e. points of  $\partial E$  are Lebesgue points of f and

$$\lim_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} \mathcal{P}_{E}[f] = f(\zeta), \qquad 0 < \alpha < \frac{\pi}{2}.$$

for every Lebesgue point  $\zeta$  of f.

2) Let A be a subset of positive linear measure of  $\partial E$  and A' the set of all density points of A. Then  $mes(A \setminus A') = 0$  and

$$\lim_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} \mathcal{P}_E[1_{\partial E \setminus A}] = 0, \qquad 0 < \alpha < \frac{\pi}{2},$$

for every  $\zeta \in A'$ . Moreover, if  $\mathcal{N}$  is a subset of  $\partial E$  with  $\operatorname{mes}(\mathcal{N}) = 0$ , then

$$\lim_{z \to \zeta, \ z \in E} \mathcal{P}_E[1_{\partial E \setminus A}] = 0, \qquad 0 < \alpha < \frac{\pi}{2},$$

for all interior points  $\zeta$  of  $A \cup \mathcal{N}$ .

*Proof.* Part 1) is classical (see, for example, Theorem 5.4.8 in [17]). The first assertion of Part 2) is also classical. Applying Part 1) to the function  $f := 1_{\partial E \setminus A}$ , the second assertion of Part 2) follows. The last assertion is almost trivial.

We recall the following well-known result due to Carathéodory and F. and M. Riesz (see [4, p. 44, p. 420]).

**Theorem 5.2.** Let  $D \subset \mathbb{C}$  be a Jordan domain and let  $\Phi$  be a conformal mapping from D onto the unit disc E.

1) Then  $\Phi$  extends to a (unique) homeomorphic map (still denoted by)  $\Phi$  from D onto  $\overline{E}$ . If, in addition, D is a rectifiable Jordan domain, then, for any linearly measurable subset A of  $\partial D$ ,  $\operatorname{mes}(A) > 0$  if and only if  $\operatorname{mes}(\Phi(A)) > 0$ .

2) Suppose that  $\gamma : [0,1] \longrightarrow \mathbb{C}$  is a one-to-one real analytic map and U is an open set such that  $\gamma([0,1]) \subset U \cap \partial D$ , and  $U \cap D$  is a Jordan domain. Then  $\Phi$  extends to a conformal mapping from  $D \cup \gamma([0,1])$  onto  $E \cup \Phi(\gamma([0,1]))$  (i.e.  $\Phi'(z)$  exists and is nonzero for  $z \in \gamma([0,1])$ ).

Now we are ready to formulate the following

**Definition 5.3.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a linearly measurable subset A of  $\partial D$ . A point  $\zeta \in A$  is said to be a density point of  $A^3$  if one of the following cases happens:

**Case 1:**  $\zeta$  is of type I. There are an open neighborhood U of  $\zeta$  such that  $U \cap D$  is a Jordan domain and a conformal mapping  $\Phi$  from  $U \cap D$  onto the unit disc E which extends homeomorphically from  $\overline{U \cap D}$  onto  $\overline{E}$  such that  $\Phi(\zeta)$  is a density point of the set  $\Phi(\partial(U \cap D) \cap A)$ . In other words,

$$\lim_{r \to 0} \frac{\operatorname{mes}(\Phi(\partial(U \cap D) \cap A) \cap \mathbb{B}(\Phi(\zeta), r))}{\operatorname{mes}(\partial E \cap \mathbb{B}(\Phi(\zeta), r))} = 1.$$

**Case 2:**  $\zeta$  is of type II. There are an open neighborhood U of  $\zeta$  such that  $U \cap D = U_1 \cup U_2$  with Jordan domains  $U_1$ ,  $U_2$ , and conformal mappings  $\Phi_j$  (j = 1, 2) from  $U_j$  onto E which extends homeomorphically from  $\overline{U_j}$  onto  $\overline{E}$  such that  $\Phi_j(\zeta)$  is a density point of the set  $\Phi_j(\partial U_j \cap A)$ . In other words,

$$\lim_{r \to 0} \frac{\operatorname{mes}(\Phi_j(\partial U_j \cap A) \cap \mathbb{B}(\Phi_j(\zeta), r))}{\operatorname{mes}(\partial E \cap \mathbb{B}(\Phi_j(\zeta), r))} = 1, \qquad j = 1, 2$$

<sup>&</sup>lt;sup>3</sup> In the case when D = E, Definition 5.3 coincides with the classical definition of density points of a subset of  $\partial E$ .

**Proposition 5.4.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a linearly measurable subset A of  $\partial D$ .

1) Then Definition 5.3 is independent of the choice of U and  $\Phi$  in Case 1 (resp. U,  $\Phi_1, \Phi_2$  in Case 2).

2) Let A' denote the set of all density points of A. Then  $mes(A \setminus A') = 0$ .

*Proof.* To prove Part 1), let  $\zeta_0$  be a density point of A. We consider two cases. **Case 1:**  $\zeta_0$  is of type I.

In virtue of Definition 5.3 and the assumption that  $\zeta_0$  is of type I, for each  $j \in \{1,2\}$  let  $U_j$  be an open neighborhood of  $\zeta_0$  such that  $U_j \cap D$  is a Jordan domain and let  $\Phi_j$  be a conformal mapping from  $U_j \cap D_j$  onto E which extends homeomorphically from  $\overline{U_j} \cap \overline{D}$  onto  $\overline{E}$ . Suppose that  $\Phi_1(\zeta_0)$  is a density point of  $\Phi_1(\partial(U_1 \cap D) \cap A) \subset \partial E$ . We would like to show that  $\Phi_2(\zeta_0)$  is a density point of  $\Phi_2(\partial(U_2 \cap D) \cap A) \subset \partial E$ .

Let V be an open neighborhood of  $\zeta_0$  such that  $V \subset U_1 \cap U_2$ ,  $V \cap D$  is a rectifiable Jordan domain and  $V \cap \partial D$  is connected. We deduce that  $\Phi_j(V \cap \partial D)$  is an open arc of  $\partial E$  (which is obviously real analytic) and  $\Phi_j(V \cap D)$  is a Jordan subdomain of E for  $j \in \{1, 2\}$ .

Let  $\Psi_1$  be a conformal mapping from the Jordan domain  $\Phi_1(V \cap D)$  onto E which extends homeomorphically from  $\overline{\Phi_1(V \cap D)}$  onto  $\overline{E}$ . By Part 2) of Theorem 5.2 and the fact that  $\Phi_1(V \cap \partial D)$  is an open real analytic arc, we see that  $\Psi_1$  extends conformally to the arc  $\Phi_1(V \cap \partial D) \subset \partial E$ , and  $(\Psi_1 \circ \Phi_1)(V \cap \partial D)$  is an arc of  $\partial E$ . Next, consider the conformal mapping  $\Psi_2$ :  $\Phi_2(V \cap D) \longrightarrow E$  given by

$$\Psi_2(z) := (\Psi_1 \circ \Phi_1 \circ \Phi_2^{-1})(z), \qquad z \in \Phi_2(V \cap D).$$

Since by Part 1) of Theorem 5.2  $\Phi_2(V \cap D)$  is a Jordan domain,  $\Psi_2$  extends homeomorphically from  $\overline{\Phi_2(V \cap D)}$  onto  $\overline{E}$  and satisfies the equation  $\Phi_2 \circ \Phi_1^{-1} = \Psi_2^{-1} \circ \Psi_1$ . Since  $\Phi_2(V \cap \partial D) \subset \partial E$  is an open real analytic arc, applying again Part 2) of Theorem 5.2, we deduce that  $\Psi_2$  extends conformally to the arc  $\Phi_2(V \cap \partial D) \subset \partial E$ , and  $(\Psi_2 \circ \Phi_2)(V \cap \partial D)$  is an arc of  $\partial E$ .

We summarize what has been done so far: We have shown that  $\Psi_2^{-1} \circ \Psi_1$  is conformal from the arc  $\Phi_1(V \cap \partial D)$  onto the arc  $\Phi_2(V \cap \partial D)$  and  $\Phi_2 \circ \Phi_1^{-1} = \Psi_2^{-1} \circ \Psi_1$ . Hence,  $|(\Phi_2 \circ \Phi_1^{-1})'(\eta)| > 0, \eta \in \Phi_1(V \cap \partial D)$ . Choosing  $\eta_0 := \Phi_1(\zeta_0)$  and noting that  $(\Phi_2 \circ \Phi_1^{-1})(\eta_0) = \Phi_2(\zeta_0)$ , we deduce that

$$\lim_{r \to 0} \frac{\operatorname{mes} \left( \Phi_2(\partial(U_2 \cap D) \cap A) \cap \mathbb{B}(\Phi_2(\zeta_0), r) \right)}{\operatorname{mes} \left( \partial E \cap \mathbb{B}(\Phi_2(\zeta_0), r) \right)}$$

$$= \lim_{r \to 0} \frac{\operatorname{mes} \left( \Phi_2(\partial(V \cap D) \cap A) \cap \mathbb{B}(\Phi_2(\zeta_0), r) \right)}{\operatorname{mes} \left( \partial E \cap \mathbb{B}(\Phi_2(\zeta_0), r) \right)}$$

$$= \lim_{r \to 0} \frac{\operatorname{mes} \left( \Phi_2(\partial(V \cap D) \cap A) \cap \mathbb{B}(\Phi_2(\zeta_0), |(\Phi_2 \circ \Phi_1^{-1})'(\eta_0)|r) \right)}{\operatorname{mes} \left( \partial E \cap \mathbb{B}(\Phi_2(\zeta_0), |(\Phi_2 \circ \Phi_1^{-1})'(\eta_0)|r) \right)}$$

$$= \lim_{r \to 0} \frac{\operatorname{mes}\left(\Phi_1(\partial(V \cap D) \cap A) \cap \mathbb{B}(\Phi_1(\zeta_0), r)\right)}{\operatorname{mes}\left(\partial E \cap \mathbb{B}(\Phi_1(\zeta_0), r)\right)}$$
  
= 1,

where the last identity follows from the assumption that  $\Phi_1(\zeta_0)$  is a density point of  $\Phi_1(\partial(V \cap D) \cap A) \subset \partial E$ .

Consequently,  $\Phi_2(\zeta_0)$  is a density point of  $\Phi_2(\partial(V \cap D) \cap A) \subset \partial E$ . Hence the proof of Part 1) is complete in this first case. **Case 2:**  $\zeta_0$  is of type II.

Then there is an open neighborhood U of  $\zeta_0$  such that  $U \cap D = U' \cup U''$ , where U', U'' are some Jordan domains. We apply the result of Case 1 to each Jordan domain U', U''. Hence the proof of Part 1) is finished.

It remains to prove Part 2). We may find a sequence  $(U_k)_{k=1}^{\infty}$  of open sets of  $\mathbb{C}$  such that  $U_k \cap D$  is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains and  $A \subset \bigcup_{k=1}^{\infty} \partial(U_k \cap D)$ . Using Part 1) and Proposition 5.1, we see that almost every point in  $(A \cap \partial(D \cap U_k))$  is a density point of this set, i.e. belongs to  $(A \cap \partial(D \cap U_k))'$ . On the other hand, clearly  $(A \cap \partial(D \cap U_k))' \subset A'$ . Consequently,

$$\operatorname{mes}(A \setminus A') \le \sum_{k=1}^{\infty} \operatorname{mes}\left( (A \cap \partial (D \cap U_k)) \setminus (A \cap \partial (D \cap U_k))' \right) = \sum_{k=1}^{\infty} 0 = 0.$$

Hence, A' is linearly measurable and  $mes(A \setminus A') = 0$ . This completes the proof of Part 2).

The following classical result will be needed to prove Proposition 5.6 below (see [11, p. 60] or [13, p. 51]).

**Theorem 5.5.** (A theorem of Lindelöf) Let D be a Jordan domain and let  $\Phi$  be a conformal mapping from E onto D. By Part 1) of Theorem 5.2 we still denote by  $\Phi$  its homeomorphic extension from  $\overline{E}$  onto  $\overline{D}$ . Let  $\zeta = e^{i\theta} \in \partial E$  and let  $\beta :=$  $\lim_{t\to\theta+} \arg\left(\Phi(e^{it}) - \Phi(e^{i\theta})\right)$ . Then  $\partial D$  has a corner <sup>4</sup> of opening  $\pi\alpha$  ( $0 \le \alpha \le 2$ ) at  $\Phi(\zeta)$  if and only if

$$\arg \frac{\Phi(z) - \Phi(\zeta)}{z - \zeta} \longrightarrow \beta - \alpha \left(\theta + \frac{\pi}{2}\right) \qquad \text{as } z \to \zeta, \qquad z \in E.$$

The next proposition justifies the coherence of the definitions of the tangent line and the angular approach regions given in Subsection 2.1.

**Proposition 5.6.** Let  $D \subset \mathbb{C}$  be an open set and let  $\zeta \in \partial^* D$ . 1) Suppose that  $\zeta$  is either of type I or of type II. For  $j \in \{1, 2\}$ , let  $\gamma_j : [a_j, b_j] \longrightarrow$ 

<sup>&</sup>lt;sup>4</sup> For the notion of a corner, which is more general than the notion of a point admitting a tangent, see [13].

 $\partial D$  be a parametrization of one Jordan curve contained in  $\partial D$  such that  $\gamma_j(c_j) = \zeta$ ,  $c_j \in (a_j, b_j)$ . Suppose that the following limit exists

$$\lim_{t \to c_1} \frac{\gamma_1(t) - \zeta}{t - c_1} \cdot \frac{|t - c_1|}{|\gamma_1(t) - \zeta|} = \lambda_1$$

Then the following limit also exists

$$\lim_{t \to c_2} \frac{\gamma_2(t) - \zeta}{t - c_2} \cdot \frac{|t - c_2|}{|\gamma_2(t) - \zeta|} = \lambda_2.$$

and  $\lambda_2 = \pm \lambda_1$ . In other words, the tangent line at  $\zeta$  is independent of the choice of parametrizations.

2) Suppose that  $\zeta$  is of type I. Suppose that U is a neighborhoods of  $\zeta$  such that  $D \cap U$  is a Jordan domain. Let  $\Phi$  be a conformal mapping from  $D \cap U$  onto E such that  $\Phi(\zeta) = 1$ , where  $\Phi(\zeta)$  is defined as in Part 1) of Theorem 5.2. Then for any  $0 < \alpha < \beta < \gamma < \frac{\pi}{2}$ , and any neighborhood V of  $\zeta$ , there are neighborhoods  $R \subset V$  of  $\zeta$  and S of 1 such that

$$\Phi\left(\mathcal{A}_{\alpha}(\zeta)\cap R\right)\subset\mathcal{A}_{\beta}(1)\cap S\subset\Phi\left(\mathcal{A}_{\gamma}(\zeta)\cap R\right).$$

3) Suppose that  $\zeta$  is of type II. Suppose that U is a neighborhood of  $\zeta$  such that  $D \cap U = U_1 \cup U_2$  with some disjoint Jordan domains  $U_1$ ,  $U_2$  satisfying  $\zeta \in \partial^* U_1 \cap \partial^* U_2$ . For  $j \in \{1, 2\}$ , let  $\Phi_j$  be a conformal mapping from  $D \cap U_j$  onto E such that  $\Phi_j(\zeta) = 1$ . Then for any  $0 < \alpha < \beta < \gamma < \frac{\pi}{2}$ ,  $j \in \{1, 2\}$ , and any neighborhood V of  $\zeta$ , there are neighborhoods  $R \subset V$  of  $\zeta$  and S of 1 such that

$$\Phi_j \left( \mathcal{A}_{\alpha}(\zeta) \cap U_j \cap R \right) \subset \mathcal{A}_{\beta}(1) \cap S \subset \Phi_j \left( \mathcal{A}_{\gamma}(\zeta) \cap U_j \cap R \right).$$

*Proof.* First one proves Part 1). Without loss of generality suppose that  $\gamma_1([a_1, b_1]) = \gamma_2([a_2, b_2]) \subset \partial D$ . Then the function  $\gamma := \gamma_1^{-1} \circ \gamma_2 : [a_2, b_2] \longrightarrow [a_1, b_1]$  is a bijective continuous map and  $\gamma(c_2) = c_1$ . Therefore, it is either monotone increasing or monotone decreasing. Observe that

$$\lim_{t \to c_2} \left( \frac{\gamma_2(t) - \zeta}{t - c_2} \cdot \frac{|t - c_2|}{|\gamma_2(t) - \zeta|} \right) = \lim_{t \to c_2} \left( \frac{(\gamma_1 \circ \gamma)(t) - \zeta}{t - c_2} \cdot \frac{|t - c_2|}{|(\gamma_1 \circ \gamma)(t) - \zeta|} \right)$$
$$= \lim_{t \to c_2} \left( \frac{(\gamma_1 \circ \gamma)(t) - \zeta}{\gamma(t) - c_1} \cdot \frac{|\gamma(t) - c_1|}{|(\gamma_1 \circ \gamma)(t) - \zeta|} \right) \cdot \lim_{t \to c_2} \left( \frac{\gamma(t) - c_1}{t - c_2} \cdot \frac{|t - c_2|}{|\gamma(t) - c_1|} \right)$$
$$= \lambda_1 \cdot \epsilon(\gamma),$$

where

$$\epsilon(\gamma) := \begin{cases} 1, & \text{if } \gamma \text{ is monotone increasing} \\ -1, & \text{if } \gamma \text{ is monotone decreasing} \end{cases}.$$

Hence,  $\lambda_2 = \pm \lambda_1$  and the proof of Part 1) is complete.

To prove Part 2), let R' be an open neighborhood of  $\zeta$  such that  $R' \subset U \cap V$  and  $R' \cap D$  is a Jordan domain. Then  $\Phi(R' \cap D)$  is a Jordan subdomain of E.

Let  $\Psi$  be a conformal mapping from E onto the Jordan domain  $R' \cap D$  such that  $\Psi(1) = \zeta$ . Then  $\Phi \circ \Psi$  is a conformal mapping from E onto the Jordan domain  $\Phi(R' \cap D)$  and  $(\Phi \circ \Psi)(1) = \Phi(\zeta)$ .

By Part 1) of Theorem 5.2, we have that

(5.2) 
$$\lim_{z \to 1, z \in E} \Psi(z) = \Psi(1) \quad \text{and} \quad \lim_{z \to 1, z \in E} (\Phi \circ \Psi)(z) = (\Phi \circ \Psi)(1) = \Phi(\zeta).$$

On the other hand, observe that  $\zeta \in \partial^*(R' \cap D)$  as  $\zeta \in \partial^*D$ . Moreover,  $\Phi(\zeta) \in \partial^*(\Phi(R' \cap D))$  because  $\partial(\Phi(R' \cap D))$  contains a neighborhood of  $\Phi(\zeta)$  in  $\partial E$ . Consequently, applying Theorem 5.5 to the two conformal mappings  $\Psi$  and  $\Phi \circ \Psi$ , we obtain the existence of the following two limits

$$\lim_{z \to 1, \ z \in E} \arg\left(\frac{\Psi(z) - \Psi(1)}{z - 1}\right) \quad \text{and} \quad \lim_{z \to 1, \ z \in E} \arg\left(\frac{(\Phi \circ \Psi)(z) - (\Phi \circ \Psi)(1)}{z - 1}\right),$$

where  $\arg z$  denotes the argument of a complex number z.

Choose an  $\alpha'$  such that  $\alpha < \alpha' < \beta$ . Then using (5.2) and (5.3), it follows that there is an open neighborhood T of 1 such that

(5.4) 
$$\begin{aligned} \mathcal{A}_{\alpha}(\zeta) \cap \Psi(T \cap E) \subset \Psi\left(\mathcal{A}_{\alpha'}(1) \cap T\right), \\ \left(\Phi \circ \Psi\right)\left(\mathcal{A}_{\alpha'}(1) \cap T\right) \subset \mathcal{A}_{\beta}(1) \cap \left(\Phi \circ \Psi\right)\left(T \cap E\right) \end{aligned}$$

Let R be a neighborhood of  $\zeta$  such that  $R \cap D = \Psi(T \cap E)$  and S a neighborhood of 1 such that  $S \cap E = (\Phi \circ \Psi) (T \cap E)$ . Then we deduce immediately from (5.4) that

$$\Phi_j\left(\mathcal{A}_\alpha(\zeta)\cap U_j\cap R\right)\subset \mathcal{A}_\beta(1)\cap S.$$

The remaining inclusion of Part 2) can be proved in exactly the same manner.

For Part 3) one argues as in the proof of Part 2). Hence the proof of the proposition is complete.  $\hfill \Box$ 

The following two lemmas will be very useful.

**Lemma 5.7.** Let  $D \subset \mathbb{C}$  be a bounded domain with  $\mathcal{C}^2$  smooth boundary and let A be a linearly measurable subset of  $\partial D$ . Then

$$\omega(z, A, D) = \mathcal{P}_D[1_{\partial D \setminus A}](z), \qquad z \in D.$$

*Proof.* Since D is bounded with  $C^2$  boundary, using (5.1) it is a classical fact (see, for example, [19]) that

(5.5) 
$$\omega(z, B, D) = \mathcal{P}_D[1_{\partial D \setminus B}](z) := \int_{\partial D} P(z, \zeta) 1_{\partial D \setminus B}(\zeta) d\sigma(\zeta), \qquad z \in D,$$

for any Borel subset B of  $\partial D$ . Here  $P(\cdot, \cdot)$  is the Poisson kernel of D and  $d\sigma$  is the Lebesgue boundary measure of  $\partial D$ . Since A is linearly measurable, there are two Borel sets  $A_1$ ,  $A_2$  such that  $A_1 \subset A \subset A_2$  and  $mes(A_2 \setminus A_1) = 0$ . It follows from definition that

(5.6) 
$$\omega(z, A_1, D) \ge \omega(z, A, D) \ge \omega(z, A_2, D), \qquad z \in D$$

On the other hand, (5.5) and the assumption on  $A_1$  and  $A_2$  imply that

$$\omega(z, A_1, D) = \omega(z, A_2, D) = \mathcal{P}_D[1_{\partial D \setminus A}](z), \qquad z \in D.$$

This, combined with (5.6), gives the desired conclusion.

**Lemma 5.8.** Let *E* be the unit disc, *A* a linearly measurable subset of  $\partial E$ , and *u* a subharmonic function defined on *E* with  $u \leq 1$ . Let  $\alpha \in (0, \frac{\pi}{2})$  be such that

$$\limsup_{\substack{\to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)}} u(z) \leq 0 \qquad for \ a.e. \ \zeta \in A.$$

Then  $u \leq \omega(\cdot, A, E)$  on E.

*Proof.* Fix a point 
$$z_0 \in E$$
. We wish to show that

(5.7) 
$$u(z_0) \le \omega(z_0, A, E).$$

As in the proof of Lemma 5.7, we may assume (without loss of generality) that A is a Borel set. Moreover, we may suppose that u is continuous at  $z_0$ . Otherwise, one considers a *Poisson Modification* v of u given by

$$v(w) := \begin{cases} u(w), & w \in E \setminus \mathbb{B}(z_0, r) \\ \mathcal{P}_{\mathbb{B}(z_0, r)}[u|_{\partial \mathbb{B}(z_0, r)}], & w \in \mathbb{B}(z_0, r) \end{cases}$$

where r > 0 is chosen so that  $B(z_0, r) \Subset E$ . Then by the hypothesis on u and by Lemma 4.1.3 in [15], we deduce that v is subharmonic on E,  $u \le v \le 1$  on E, and  $v|_{\mathbb{B}(z_0,r)}$  is harmonic.

For 
$$0 < r < 1$$
 put  $u_r(z) := u(rz), z \in E$ . We obtain  
 $u(z_0) = \lim_{r \to 1} u(rz_0) \leq \limsup_{r \to 1} \mathcal{P}_E[u_r](z_0) \leq \mathcal{P}_E[\limsup_{r \to 1} u_r](z_0)$   
 $\leq \mathcal{P}_E[1_{\partial E \setminus A}](z_0) = \omega(z_0, A, E),$ 

where the second estimate holds because of Fatou's Lemma, the third one follows from the hypothesis on u, and the last equality is a consequence of (5.1). This proves (5.7). Hence, the proof is complete.

The following estimate will be crucial for the future development.

**Proposition 5.9.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on linearly measurable subsets A,  $\mathcal{N}$  of  $\partial D$  with  $\operatorname{mes}(\mathcal{N}) = 0$ . Then

1) all density points of A which are elements of  $\partial^* D$  are locally regular points of A and  $\operatorname{mes}(A \setminus A^*) = 0$ ;

2) for every  $\zeta \in A^*$  and every open neighborhood U of  $\zeta$ , it holds

$$\sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} \omega(z, A, D) = 0$$

and  $\operatorname{mes}(A \cap U) > 0;$ 

3) for any interior point  $\zeta$  of  $A \cup \mathcal{N}$ , we have

$$\lim_{z \to \zeta, \ z \in D} \omega(z, A, D) = 0.$$

*Proof.* To prove Part 1), let  $\zeta_0$  be a density point of A. There are two cases to consider according to the type of  $\zeta_0$ . We only treat the case when  $\zeta_0$  is of type I. The second case where  $\zeta_0$  is of type II is almost analogous, and therefore, left to the interested reader.

Since  $\zeta_0$  is of type I and D is rectifiable at  $\zeta_0$ , there is an open neighborhood U of  $\zeta_0$  such that  $U \cap D$  is a rectifiable Jordan domain. Fix a conformal mapping  $\Phi$ 

from  $U \cap D$  onto E which extends to a homeomorphism from  $\overline{U \cap D}$  onto  $\overline{E}$ . By Definition 5.3 and Proposition 5.4,  $\eta_0 := \Phi(\zeta_0)$  is a density point of the linearly measurable subset  $\Phi(\partial(U \cap D) \cap A)$  of  $\partial E$ . Consequently, applying identity (5.1) and Part 1) of Proposition 5.1 yields that

(5.8) 
$$\sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{w \in \mathcal{A}_{\alpha}(\eta_0)} \omega \left( w, \Phi(\partial(U \cap D) \cap A), E \right) = 0.$$

On the other hand, one has

(5.9)  $\omega(z,\partial(U\cap D)\cap A), U\cap D) = \omega(\Phi(z), \Phi(\partial(U\cap D)\cap A), E), \qquad z \in U \cap D.$ 

Applying Theorem 5.5 to  $\Phi^{-1}$  at  $\eta_0$  and combining (5.8) and (5.9), it follows that

$$\sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{z \in \mathcal{A}_{\alpha}(\zeta_0)} \omega\left(z, \partial(U \cap D) \cap A, U \cap D\right) = 0.$$

Hence,  $\zeta_0 \in A^*$ . This proves the first assertion of Part 1).

Let A' denote the set of all density points of A. By Part 2) of Proposition 5.4, mes $(A \setminus A') = 0$ . On the other hand, we have already shown that  $A' \cap \partial^* D \subset A^*$ . Consequently,

$$\operatorname{mes}(A \setminus A^*) \le \operatorname{mes}\left(A \setminus (A' \cap \partial^* D)\right) = \operatorname{mes}(A \setminus A') + \operatorname{mes}(A \setminus \partial^* D) = 0.$$

which completes the proof of the last assertion of Part 1).

Part 2) follows from the definition of regular points and the Subordination Principle (see Corollary 4.3.9 in [15]).

In virtue of Lemma 5.8 and the second assertion of Part 2) of Proposition 5.1, the proof of Part 3) follows along the same lines as that of Part 2).  $\Box$ 

In the sequel we formulate some important stability property of the angular harmonic measure. Recall that D is a proper open subset of  $\mathbb{C} \cup \{\infty\}$  such that the boundary  $\partial D$  (with respect to  $\mathbb{C} \cup \{\infty\}$ ) is non-polar. Let A be a linearly measurable subset of  $\partial^* D$ . Let  $\phi : \partial D \longrightarrow \mathbb{R}$  be a bounded function. The associated *Perron* function  $H_{D,A}: D \longrightarrow \mathbb{R}$  is defined by

(5.10) 
$$H_{D,A}\phi := \sup_{u \in \widehat{\mathcal{U}}} u,$$

where  $\widehat{\mathcal{U}} = \widehat{\mathcal{U}}(\phi, A, D)$  denotes the family of all subharmonic functions u on D such that

$$\limsup_{z \to \zeta} u(z) \leq \phi(\zeta), \qquad \zeta \in \partial D \setminus A,$$
$$\limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} u(z) \leq \phi(\zeta), \qquad \zeta \in A, \ 0 < \alpha < \frac{\pi}{2}.$$

In the sequel,  $\widehat{\mathcal{U}}(A, D)$  will stand for  $\widehat{\mathcal{U}}(1_{\partial D \setminus A}, A, D)$ .

**Proposition 5.10.** (Maximum Principle) Let  $u \in SH(D)$  be such that u is bounded from the above and

$$\limsup_{z \to \zeta} u(z) \leq 0, \qquad \zeta \in \partial D \setminus A,$$
$$\limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} u(z) \leq 0, \qquad \zeta \in A, \ 0 < \alpha < \frac{\pi}{2}.$$

Then  $u \leq 0$  on D.

*Proof.* Suppose that u < M for some M. Let  $\zeta_0$  be an arbitrary point of A. Fix a rectifiable Jordan domain U such that  $U \subset D$  and  $\partial U \cap \partial D$  is a neighborhood of  $\zeta_0$  in  $\partial D$ . Applying Lemma 5.8 and Part 3) of Proposition 5.9 to  $u|_U$  yields that

$$\limsup_{z \to \zeta, \ z \in U.} u(z) \le 0.$$

Since  $\zeta_0$  is an arbitrary point of A, the desired conclusion follows from the classical Maximum Principle (see Theorem 2.3.2 in [15]).

Using the above proposition, the corresponding results in Sections 4.1 and 4.2 of [15] with respect to  $H_{D,A}$  (instead of  $H_D$ ) are still valid making the obviously necessary changes. In particular, we have the following (see Corollary 4.2.6 in [15]):

**Proposition 5.11.** Let D be a proper open subset of  $\mathbb{C} \cup \{\infty\}$  such that the boundary  $\partial D$  (with respect to  $\mathbb{C} \cup \{\infty\}$ ) is non-polar. Let A be a linearly measurable subset of  $\partial^* D$  and  $\phi : \partial D \longrightarrow \mathbb{R}$  a bounded function which is continuous n.e. on  $\partial D$ . Then there exists a unique bounded harmonic function h on D such that  $\lim_{z \to \zeta} h(z) = \phi(\zeta)$ 

for n.e.  $\zeta \in \partial D$ . Moreover,  $h = H_D \phi = H_{D,A} \phi$ .

In virtue of this result, Theorem 4.3.3 in [15] is still valid in the context of  $H_{D,A}$ . More precisely,

**Proposition 5.12.** Let D be a proper open subset of  $\mathbb{C} \cup \{\infty\}$  such that  $\partial D$  (with respect to  $\mathbb{C} \cup \{\infty\}$ ) is non-polar. Let A be a linearly measurable subset of  $\partial^* D$  and  $\phi: \partial D \longrightarrow \mathbb{R}$  a bounded Borel function. Then  $H_D \phi = H_{D,A} \phi = \mathcal{P}_D[\phi]$ .

We also need the following

**Proposition 5.13.** Let D be a proper open subset of  $\mathbb{C} \cup \{\infty\}$  such that  $\partial D$  (with respect to  $\mathbb{C} \cup \{\infty\}$ ) is non-polar. Let A be a Borel subset of  $\partial D$  such that  $A \subset \partial^* D$  and  $\operatorname{mes}(A) = 0$ . Then  $\mathcal{P}_D[1_A] \equiv 0$  on D.

*Proof.* Suppose without loss of generality that D is locally rectifiable on the interval  $[0,1] \subset \partial D$  and that A is a Borel subset of [0,1] with  $\operatorname{mes}(A) = 0$ . Since  $D \subset \mathbb{C} \setminus [0,1]$ , it follows from the Subordination Principle that

$$\mathcal{P}_D[1_A] \le \mathcal{P}_{\mathbb{C}\setminus[0,1]}[1_A] \quad \text{on } D.$$

Therefore, it suffices to show that  $\mathcal{P}_{\mathbb{C}\setminus[0,1]}[1_A] \equiv 0$  on  $\mathbb{C}\setminus[0,1]$ . To this end consider the conformal mapping  $\Phi(z) := \sqrt{\frac{1}{z} - 1}$  which maps  $\mathbb{C} \cup \{\infty\} \setminus [0,1]$  onto  $\mathbb{H} :=$   $\{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ . It is not difficult to show that

$$\mathcal{P}_{\mathbb{C}\setminus[0,1]}[1_A] = \mathcal{P}_{\mathbb{H}}[1_{\Phi(A)}] \circ \Phi^{-1} \equiv 0.$$

This concludes the proof.

Now we arrive at one of the main results of the section

**Theorem 5.14.** Let  $D \subset \mathbb{C}$  be a proper open subset of  $\mathbb{C} \cup \{\infty\}$  such that  $\partial D$  (with respect to  $\mathbb{C} \cup \{\infty\}$ ) is non-polar. Suppose that D is locally rectifiable on a linearly measurable subset A of  $\partial D$ . Let B be a linearly measurable subset of  $\partial D$  with  $B \subset A \cap A^*$  and  $\operatorname{mes}(A \setminus B) = 0$ . Then  $\omega(z, A, D) = H_{D,B} \mathbb{1}_{\partial D \setminus B}$  for  $z \in D$ . In particular, for every subset  $\mathcal{N}$  of  $\partial D$  with  $\operatorname{mes}(\mathcal{N}) = 0$ , we have  $\omega(z, A \setminus \mathcal{N}, D) = \omega(z, A, D)$ for  $z \in D$ .

*Proof.* Replacing A by two Borel sets  $A_1$ ,  $A_2$  such that  $A_1 \subset B$  and  $A \subset A_2 \subset \partial D$ and  $\operatorname{mes}(B \setminus A_1) = \operatorname{mes}(A_2 \setminus A) = 0$ , one gets that  $\operatorname{mes}(A_2 \setminus A_1) = 0$ . Then we conclude by the Subordination Principle and Proposition 5.13 that

$$\omega(z, A_2, D) \le \omega(z, A, D) \le \omega(z, B, D) \le \omega(z, A_1, D) = \omega(z, A_2, D), \qquad z \in D.$$

In virtue of Proposition 5.12, we have that

$$\omega(z, A_1, D) = H_{D, A_1} \mathbf{1}_{\partial D \setminus A_1}.$$

On the other hand, it follows from the definition that

$$H_{D,A_1} \mathbb{1}_{\partial D \setminus A_1} \ge H_{D,B} \mathbb{1}_{\partial D \setminus B} \ge \omega(\cdot, A, D)$$
 on  $D$ .

Combining the above three estimates, the proof of the first assertion of the theorem follows.

The second assertion is a direct consequence of the first one.

**Proposition 5.15.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a linearly measurable subset A of  $\partial D$ . Let  $(D_k)_{k=1}^{\infty}$  be a sequence of open subsets  $D_k$  of D and  $(A_k)_{k=1}^{\infty}$  a sequence of measurable subsets of A such that

- (i)  $D_k \subset D_{k+1}$  and  $\bigcup_{k=1}^{\infty} D_k = D;$
- (ii)  $A_k \subset A_{k+1}$  and  $A_k \subset \partial D \cap \partial D_k$  and  $D_k$  is locally rectifiable on  $A_k$  and  $\bigcup_{k=1}^{\infty} A_k = A;$
- (iii) for any point  $\zeta \in A$  there is an open neighborhood  $V = V_{\zeta}$  of  $\zeta$  in  $\mathbb{C}$  such that  $V \cap D = V \cap D_k$  for some k.

Then

$$\omega(z, A, D) = \lim_{k \to \infty} \omega(z, A_k, D_k), \qquad z \in D.$$

**Remark 5.16.** We may always choose sequences  $(D_k)_{k=1}^{\infty}$  and  $(A_k)_{k=1}^{\infty}$  such that (i)-(iii) are satisfied and the open sets  $D_k$  are bounded. For example, take  $D_k := D \cap \mathbb{B}(0, k)$  and  $A_k := A \cap \mathbb{B}(0, k), k \ge 1, k \in \mathbb{N}$ .

22

	-	-	
L			

*Proof.* Using the Subordination Principle it is easy to see that the sequence  $(\omega(\cdot, A_k, D_k))_{k=1}^{\infty}$  is decreasing and the following limit

$$u := \lim_{k \to \infty} \omega(\cdot, A_k, D_k)$$

exists and defines a subharmonic function in D. By the Subordination Principle again, we have  $u \ge \omega(\cdot, A, D)$ . Therefore, it remains to establish the converse inequality. In virtue of (iii) and of Part 2) of Proposition 5.9, we conclude that

(5.11) 
$$\sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} u = 0, \qquad \zeta \in B,$$

where  $B := \bigcup_{k=1}^{\infty} A_k^*$ .

On the other hand, in virtue of Part 1) of Proposition 5.9, we have that

$$\operatorname{mes}((A \cap A^*) \setminus B) \le \sum_{k=1}^{\infty} \operatorname{mes}(A_k \setminus A_k^*) = 0.$$

Consequently, applying Theorem 5.14, we deduce from (5.11) that  $u(z) \leq \omega(z, A, D)$ ,  $z \in D$ . This completes the proof.

Next, we introduce a notion which will be relevant for our further study.

**Definition 5.17.** Let  $D, G \subset \mathbb{C}$  be two open sets such that  $G \subset D$ . A point  $\zeta \in \partial^* D$ is said to be an end-point of G in D if, for every  $0 < \alpha < \frac{\pi}{2}$ , there is an open neighborhood  $U = U_{\alpha}$  of  $\zeta$  such that  $U \cap \mathcal{A}_{\alpha}(\zeta) \subset G$ . The set of all end-points of Gin D is denoted by  $G^D$ .

The remaining part of this section is devoted to the study of level sets of the harmonic measure. We begin with the following important properties of these sets.

**Theorem 5.18.** Let  $D \subset \mathbb{C}$  be an open set and A a linearly measurable subset of  $\partial D$  such that D is locally rectifiable on A and mes(A) > 0. Then, for any  $0 < \epsilon < 1$ , the " $\epsilon$ -level set"

$$D_{\epsilon} := \{ z \in D : \ \omega(z, A, D) < 1 - \epsilon \}$$

enjoys the following properties:

- (i) Let  $G_1, G_2$  be arbitrary distinct connected components of  $D_{\epsilon}$ , then  $G_1^D \cap G_2^D = \emptyset$ .
- (ii) For any point  $\zeta \in A^*$ , there is exactly one connected component G of  $D_{\epsilon}$  such that  $\zeta \in G^D$ .
- (iii)  $G^D \cap A$  is linearly measurable and  $\operatorname{mes}(G^D \cap A) > 0$  for every connected component G of  $D_{\epsilon}$ .

*Proof.* To prove (i), suppose, in order to reach a contradiction, that  $G_1^D \cap G_2^D \neq \emptyset$ . Fix a point  $\zeta_0 \in G_1^D \cap G_2^D$ . Then, for any  $0 < \alpha < \frac{\pi}{2}$ , there is an open neighborhood  $U_{\alpha}$  of  $\zeta_0$  such that  $\mathcal{A}_{\alpha}(\zeta) \cap U_{\alpha} \subset G_1 \cap G_2$ . This implies that  $G_1 \cap G_2 \neq \emptyset$ . Hence,  $G_1 = G_2$ , which contradicts the hypothesis that  $G_1 \neq G_2$ . The proof of (i) is complete.

Next, we turn to the proof of (ii). Fix a  $\zeta_0 \in A^*$ . In virtue of assertion (i), it suffices to show the existence of a connected component G of  $D_{\epsilon}$  such that  $\zeta_0 \in G^D$ .

Applying Part 2) of Proposition 5.9, we see that, for every  $0 < \alpha < \frac{\pi}{2}$ , there is an open neighborhood  $U_{\alpha}$  of  $\zeta_0$  such that

(5.12) 
$$\mathcal{A}_{\alpha}(\zeta_0) \cap U_{\alpha} \subset D_{\epsilon}$$

Fix an arbitrary  $0 < \alpha_0 < \frac{\pi}{2}$ , and let G be the connected component of  $D_{\epsilon}$  containing  $\mathcal{A}_{\alpha_0}(\zeta_0) \cap U_{\alpha_0}$ . Since

$$\left(\mathcal{A}_{\alpha_0}(\zeta_0)\cap U_{\alpha_0}\right)\cap \left(\mathcal{A}_{\alpha}(\zeta_0)\cap U_{\alpha}\right)\neq \emptyset, \quad 0<\alpha<\frac{\pi}{2},$$

we deduce from (5.12) that G also contains  $\mathcal{A}_{\alpha}(\zeta_0) \cap U_{\alpha}$  for every  $0 < \alpha < \frac{\pi}{2}$ . Hence  $\zeta_0 \in G^D$ . The proof of (ii) is finished.

Finally, we prove (iii). First, we may find a sequence  $(U_k)_{k=1}^{\infty}$  of open sets of  $\mathbb{C}$  such that  $U_k \cap D$  is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains and  $A \subset \bigcup_{k=1}^{\infty} \partial(U_k \cap D)$ . Since A is linearly measurable, we see that in order to prove the measurability of  $G^D \cap A$ , it is sufficient to check that  $G^D \cap \partial(D \cap U_k)$  is linearly measurable for every  $k \geq 1$ . To prove the latter assertion, fix an  $k_0 \geq 1$  and let  $U := U_{k_0}$ . Let  $\Phi$  be a conformal mapping from  $D \cap U$  onto E which extends to a homeomorphic mapping (still denoted by)  $\Phi$  from  $\overline{D \cap U}$  onto  $\overline{E}$ . Applying Part 2) of Proposition 5.6 and using Definition 5.17, we see that, for any  $\zeta \in \partial(D \cap U), \zeta \in G^D$  if and only if  $\Phi(\zeta) \in [\Phi(G \cap U)]^E$ . We shall prove, in the sequel, that  $[\Phi(G \cap U)]^E$  is a Borel subset of  $\partial E$ . Taking this for granted, then  $G^D \cap \partial(D \cap U)$  is also a Borel set. Consequently,  $G^D \cap A$  is linearly measurable. To check that  $[\Phi(G \cap U)]^E$  is a Borel set, put

To eneck that  $[\Psi(G + U)]$  is a Dorel set, put

(5.13) 
$$\mathcal{A}_{n,m}(\eta) := \left\{ w \in E \cap \mathcal{A}_{\left(1-\frac{1}{n}\right) \cdot \frac{\pi}{2}}(\eta) : |w-\eta| < \frac{1}{m} \right\}, \ n, m \ge 1, \ \eta \in \partial E.$$

For any  $n, m, p \ge 1$ , let

(5.14) 
$$T_{nmp} := \left\{ \eta \in \partial E : \mathcal{A}_{n,m}(\eta) \subset \Phi(G \cap U) \text{ and} \\ \omega(\Phi^{-1}(w), A, D) \leq 1 - \epsilon - \frac{1}{p}, \forall w \in \mathcal{A}_{n,m}(\eta) \right\}.$$

We observe the following:

**Geometric fact.** Let  $\eta_0 \in \partial E$  and  $(\eta_q)_{q=1}^{\infty} \subset \partial E$  such that  $\lim_{q \to \infty} \eta_q = \eta_0$ . Then

$$\mathcal{A}_{n,m}(\eta_0) \subset \bigcup_{q=1}^{\infty} \mathcal{A}_{n,m}(\eta_q)$$

The proof of this fact follows immediately from the geometric shape of the cone  $\mathcal{A}_{n,m}(\eta)$  given in (5.13).

Let  $(\eta_q)_{q=1}^{\infty} \subset T_{nmp}$  such that  $\lim_{q \to \infty} \eta_q = \eta_0 \in \partial E$ . Using the above geometric fact, we see that  $\mathcal{A}_{n,m}(\eta_0) \subset \Phi(G \cap U)$ . This, combined with (5.14) and the continuity of  $\omega(\Phi^{-1}(\cdot), A, D)|_E$ , implies that  $\eta_0 \in T_{nmp}$ . Hence, the set  $T_{nmp}$  is closed. Clearly, we have

$$[\Phi(G \cap U)]^E = \bigcap_{n=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcup_{p=1}^{\infty} T_{nmp}$$

It follows immediately from this identity that  $[\Phi(G \cap U)]^E$  is a Borel set. Consequently, as was already discussed before,  $G^D \cap A$  is linearly measurable.

To finish assertion (iii), it remains to prove that mes  $(G^E \cap A) > 0$ . Suppose, in order to reach a contradiction, that mes  $(G^E \cap A) = 0$ . Consider the following function

$$u(z) := \begin{cases} \omega(z, A, D), & z \in D \setminus G\\ 1 - \epsilon, & z \in G \end{cases}$$

Then clearly  $u \in SH(D)$  and  $u \leq 1$ . In virtue of assertions (i) and (ii) and Part 2) of Proposition 5.9, we have that

 $\sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} u(z) = \sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} \omega(z, A, E) = 0, \qquad \zeta \in (A \cap A^*) \setminus (G^D \cap A).$ 

Consequently, using the notation in (5.10), we conclude that

$$u \in \mathcal{U}\left((A \cap A^*) \setminus \mathcal{N}, D\right),$$

where  $\mathcal{N} := G^D \cap A$ . Since, by our above assumption,  $\operatorname{mes}(\mathcal{N}) = 0$ , it follows from Theorem 5.14 that  $u \leq \omega(\cdot, A, D)$ . But on the other hand, one has  $\omega(z, A, D) < 1 - \epsilon = u(z)$  for  $z \in G$ . This leads to the desired contradiction. Hence, the proof of (iii) is finished.

**Theorem 5.19.** Let  $D \subset \mathbb{C}$  be an open set and A a linearly measurable subset of  $\partial D$  such that D is locally rectifiable on A and  $\operatorname{mes}(A) > 0$ . For any  $0 \leq \epsilon < 1$ , let  $D_{\epsilon} := \{z \in D : \omega(z, A, D) < 1 - \epsilon\}$ . 1) For any subset  $\mathcal{N}$  of  $\partial D$  such that  $\operatorname{mes}(\mathcal{N}) = 0$ , let

$$\mathcal{U}_{\epsilon}(A, \mathcal{N}, D) := \left\{ u \in \mathcal{SH}(D_{\epsilon}) : \ u \leq 1 \ and \ \sup_{0 < \alpha < \frac{\pi}{2}} \limsup_{z \to \zeta, \ z \in \mathcal{A}_{\alpha}(\zeta)} u(z) \leq 0, \ \zeta \in (A \cap A^{*}) \setminus \mathcal{N} \right\}$$

Then  $\mathcal{U}_{\epsilon}(A, \mathcal{N}, D) = \mathcal{U}_{\epsilon}(A, \emptyset, D).$ 2) Define the "harmonic measure of the  $\epsilon$ -level set"  $\omega_{\epsilon}(\cdot, A, D)$  as

$$\omega_{\epsilon}(z, A, D) := \begin{cases} \sup_{u \in \mathcal{U}_{\epsilon}(A, \emptyset, D)} u(z), & z \in D_{\epsilon} \\ 0, & z \in A^* \end{cases}$$

Then

$$\omega_{\epsilon}(z, A, D) = \frac{\omega(z, A, D)}{1 - \epsilon}, \qquad z \in D_{\epsilon} \cup A^*.$$

*Proof.* Clearly, by definition,  $\mathcal{U}_{\epsilon}(A, \emptyset, D) \subset \mathcal{U}_{\epsilon}(A, \mathcal{N}, D)$ . To prove the converse inclusion, fix an arbitrary  $u \in \mathcal{U}_{\epsilon}(A, \mathcal{N}, D)$ . Consider the following function

$$\hat{u}(z) := \begin{cases} \max\left\{(1-\epsilon)u(z), \omega(z, A, D)\right\}, & z \in D_{\epsilon} \\ \omega(z, A, D), & z \in D \setminus D_{\epsilon} \end{cases}$$

Then  $\hat{u} \in \mathcal{SH}(D)$  and  $\hat{u} \leq 1$ . Moreover, in virtue of Part 2) of Proposition 5.9, we have that  $A^* \subset (D_{\epsilon})^D$ . Consequently, for every  $\zeta \in (A \cap A^*) \setminus \mathcal{N}$ ,

(5.15) 
$$\sup_{0<\alpha<\frac{\pi}{2}} \limsup_{z\to\zeta,\ z\in\mathcal{A}_{\alpha}(\zeta)} \hat{u}(z)$$
$$\leq \max\left\{\sup_{0<\alpha<\frac{\pi}{2}} \limsup_{z\to\zeta,\ z\in\mathcal{A}_{\alpha}(\zeta)} u(z), \sup_{0<\alpha<\frac{\pi}{2}} \limsup_{z\to\zeta,\ z\in\mathcal{A}_{\alpha}(\zeta)} \omega(z,A,D)\right\}.$$

Observe that the first term in the latter line of (5.15) is equal to 0 because  $u \in \mathcal{U}_{\epsilon}(A, \mathcal{N}, D)$ . In addition, in virtue of Part 2) of Proposition 5.9, the second term in the latter line of (5.15) is also equal to 0. Hence,  $\hat{u} \in \mathcal{U}((A \cap A^*) \setminus \mathcal{N}, D)$ . Consequently, by Theorem 5.14,  $\hat{u} \leq \omega(\cdot, A, D)$ . In particular, one has

(5.16) 
$$u(z) \le \frac{\omega(z, A, D)}{1 - \epsilon}, \qquad z \in D, \ u \in \mathcal{U}_{\epsilon}(A, \mathcal{N}, D).$$

On the other hand, as an immediate consequence of Part 2) of Proposition 5.9, we get that  $\frac{\omega(\cdot,A,D)}{1-\epsilon} \in \mathcal{U}_{\epsilon}(A, \emptyset, D) \subset \mathcal{U}_{\epsilon}(A, \mathcal{N}, D)$ . This, combined with (5.16), implies the desired conclusions of Part 1) and Part 2).

An immediate consequence of Theorem 5.19 is the following Two-Constant Theorem for level sets.

**Corollary 5.20.** Let  $D \subset \mathbb{C}$  be an open set and let A and  $\mathcal{N}$  be two linearly measurable subsets of  $\partial D$  such that D is locally rectifiable on A,  $\operatorname{mes}(A) > 0$  and  $\operatorname{mes}(\mathcal{N}) = 0$ . Let  $0 \leq \epsilon < 1$  and put  $D_{\epsilon} := \{z \in D : \omega(z, A, D) < 1 - \epsilon\}$ . If  $u \in \mathcal{SH}(D_{\epsilon})$  satisfies  $u \leq M$  on  $D_{\epsilon}$  and  $\sup_{0 \leq \alpha \leq \frac{\pi}{2}} \lim_{z \to \zeta, z \in \mathcal{A}_{\alpha}(\zeta)} u(z) \leq m, \zeta \in (A \cap A^*) \setminus \mathcal{N}$ ,

then

$$u(z) \le m(1 - \omega_{\epsilon}(z, A, D)) + M \cdot \omega_{\epsilon}(z, A, D).$$

### 6. Boundary behaviour of the Gonchar-Carleman operator

Before investigating the boundary behavior of the Gonchar–Carleman operator, we first introduce the following notion and study its properties.

## 6.1. Angular Jordan domains. Let E be the unit disc. We begin with the

**Definition 6.1.** For every closed subset F of  $\partial E$  and any real number h such that  $\operatorname{mes}(F) > 0$  and  $\sup_{x,y \in F} |x-y| < h < 1 - \frac{\sqrt{2}}{2}$ , the open set

$$\Omega = \Omega(F, h) := \bigcup_{\zeta \in F} \left\{ z \in \mathcal{A}_{\frac{\pi}{4}}(\zeta) : |z| > 1 - h \right\}$$

is called the angular Jordan domain with base F and height h.

We give below a list of properties of such angular Jordan domains.

**Proposition 6.2.** Let  $\Omega = \Omega(F, h)$  be an angular Jordan domain. 1) Then there exist exactly two points  $\zeta_1, \zeta_2 \in F$  such that  $|\zeta_1 - \zeta_2| = \sup_{x,y \in F} |x-y|$ and  $F \subset [\zeta_1, \zeta_2]$ , where  $[\zeta_1, \zeta_2]$  is the (small) closed arc of  $\partial E$  which is oriented in the positive sense and which starts from  $\zeta_1$  and ends at  $\zeta_2$ .

2) Write the open set  $[\zeta_1, \zeta_2] \setminus F$  as the union of disjoint open arcs

$$[\zeta_1, \zeta_2] \setminus F = \bigcup_{j \in J} (a_j, b_j),$$

where  $(a_j, b_j)$  is the (small) open arc of  $\partial E$  which goes from  $a_j$  to  $b_j$  and which is oriented in the positive sense, and the index set J is finite or countable.

For  $j \in J$ , we construct the isosceles triangle with the three vertices  $a_j$ ,  $b_j$  and  $c_j$  such that the base of the isosceles triangle is the segment connecting  $a_j$  to  $b_j$ , and  $c_j$  satisfies

$$\operatorname{arg}\left(\frac{c_j-a_j}{a_j}\right) = \frac{3\pi}{4} \quad and \quad \operatorname{arg}\left(\frac{c_j-b_j}{b_j}\right) = \frac{-3\pi}{4}.$$

Let  $[a_j, c_j]$  (resp.  $[c_j, b_j]$ ) denote the segment connecting  $a_j$  to  $c_j$  (resp. the segment connecting  $c_j$  to  $b_j$ ). Put

$$F_0 := F \cup \bigcup_{j \in J} ([a_j c_j] \cup [c_j b_j])$$

Then  $F_0$  is a rectifiable Jordan curve starting from  $\zeta_1$  and ending at  $\zeta_2$ . 3) Let  $\eta_1$  (resp.  $\eta_2$ ) be the unique point in the circle  $\partial \mathbb{B}(0, 1-h)$  such that

$$\arg\left(\frac{\eta_1-\zeta_1}{\zeta_1}\right) = \frac{-3\pi}{4} \quad \left(resp. \quad \arg\left(\frac{\eta_2-\zeta_2}{\zeta_2}\right) = \frac{3\pi}{4}\right)$$

and that  $|\eta_1 - \zeta_1|$  (resp.  $|\eta_2 - \zeta_2|$ ) is minimal. Let  $F_1$  (resp.  $F_2$ ) denote the segment connecting  $\eta_1$  to  $\zeta_1$  (resp. the segment connecting  $\zeta_2$  to  $\eta_2$ ). Let  $F_3$  be the (small) closed arc of the circle  $\partial \mathbb{B}(0, 1 - h)$  which starts from  $\eta_2$  and ends at  $\eta_1$  and which is oriented in the negative sense.

Then  $\Omega$  is a rectifiable Jordan domain and its boundary  $\Gamma$  consists of the rectifiable Jordan curve  $F_0$ , two segments  $F_1$ ,  $F_2$  and the closed arc  $F_3$ .

4) For every  $\epsilon \in (0, \frac{h}{4})$  define the dilatation  $\tau_{\epsilon} : E \longrightarrow E$  as follows

$$\tau_{\epsilon}(z) := (1 - \epsilon)z, \qquad z \in E$$

Put

$$\Omega_{\epsilon} := \tau_{\epsilon}(\Omega) \setminus \overline{\mathbb{B}(0, (1+\epsilon)(1-h))}.$$

Then  $\Omega_{\epsilon}$  is a rectifiable Jordan domain and its boundary  $\Gamma_{\epsilon}$  consists of the rectifiable Jordan curve  $F_{0\epsilon} := \tau_{\epsilon}(F_0)$ , a sub-segment  $F_{1\epsilon}$  of  $\tau_{\epsilon}(F_1)$ , a sub-segment  $F_{2\epsilon}$  of  $\tau_{\epsilon}(F_2)$ , and a closed arc  $F_{3\epsilon}$  of  $\partial \mathbb{B}(0, (1 + \epsilon)(1 - h))$ .

5) Consider the projection  $\tau : E \setminus \{0\} \longrightarrow \partial E$  given by  $\tau(z) := \frac{z}{|z|}, z \in E \setminus \{0\}.$ 

For every  $\epsilon \in (0, \frac{h}{4})$  notice that  $F_{0\epsilon} \cup F_{1\epsilon} \cup F_{2\epsilon} = \Gamma_{\epsilon} \setminus \partial \mathbb{B}(0, (1+\epsilon)(1-h))$ . Then the two maps

$$F_{0\epsilon} \cup F_{1\epsilon} \cup F_{2\epsilon} \ni \zeta \quad \mapsto \quad \tau(\zeta) \in \partial E,$$
  
$$F_{3\epsilon} \ni \zeta \quad \mapsto \quad \tau(\zeta) \in \partial E,$$

are one-to-one. In addition, for any linearly measurable subset A of  $\Gamma_{\epsilon}$ ,

$$\operatorname{mes}(A) \le 10 \cdot \operatorname{mes}(\tau(A)).$$

6)  $\Omega_{\epsilon} \nearrow \Omega$  as  $\epsilon \searrow 0$ . 7) For any closed Jordan curve  $\mathcal{C}$  contained in  $\Omega$  there is an  $\epsilon > 0$  such that  $\mathcal{C} \subset \Omega_{\epsilon}$ . 8)  $\operatorname{mes}(F \setminus \Omega^E) = 0.$ 

*Proof.* All assertions are quite simple using an elementary geometric argument. Therefore, we leave the details of their proofs to the reader. However, we will give the proof of the fact that  $\Omega$  is a domain. This proof will clarify Definition 6.1.

In virtue of the condition on F and h given in Definition 6.1, we see that  $\left\{z \in \mathcal{A}_{\frac{\pi}{4}}(\zeta) : |z| > 1 - h\right\}, \zeta \in \partial E$ , is connected, and that

$$\{z \in \mathcal{A}_{\frac{\pi}{4}}(\zeta) : |z| > 1 - h\} \cap \{z \in \mathcal{A}_{\frac{\pi}{4}}(\eta) : |z| > 1 - h\} \neq \emptyset,$$
  
 
$$\forall \zeta, \eta \in \partial E : |\zeta - \eta| < h < 1 - \frac{\sqrt{2}}{2}.$$
  
ence,  $\Omega$  is a domain.

Hence,  $\Omega$  is a domain.

**Theorem 6.3.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a linearly measurable subset A of  $\partial D$  with mes(A) > 0. Then, for any  $0 \le \epsilon < 1$  and any connected component G of  $D_{\epsilon} := \{z \in D : \omega(z, A, D) < 1 - \epsilon\}$ , there are an open set  $U \subset \mathbb{C}$ , a conformal mapping  $\Phi$ , and an angular Jordan domain  $\Omega = \Omega(F, h)$ such that

- (i)  $U \cap D$  is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains;
- (ii)  $\Phi$  maps E conformally onto one connected component of  $U \cap D$  (notice that, in virtue of (i),  $U \cap D$  has at most two connected components);
- (iii)  $\Phi(F) \subset A \cap A^* \cap G^D$  and  $\Phi(\Omega) \subset G$ .

*Proof.* We have already shown in the proof of (iii) of Theorem 5.18 that there is a sequence  $(U_k)_{k=1}^{\infty}$  of open sets of  $\mathbb{C}$  such that  $U_k \cap D$  is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains, and  $A \subset \bigcup_{k=1}^{\infty} \partial(U_k \cap U_k)$ D), and  $\operatorname{mes}(A \cap A^* \cap G^D) > 0$ . Consequently, there is an index  $k_0$  such that

(6.1) 
$$\operatorname{mes}\left(A \cap A^* \cap G^D \cap \partial(D \cap U)\right) > 0,$$

where  $U := U_{k_0}$ . Suppose without loss of generality that  $U \cap D$  is a rectifiable Jordan domain. The remaining case where  $U \cap D$  is the disjoint union of two rectifiable Jordan domains may be proved in the same way. Let  $\Phi$  be a conformal mapping

from E onto  $D \cap U$ . By Theorem 5.2,  $\Phi$  extends to a homeomorphic map (still denoted by)  $\Phi$  from  $\overline{E}$  onto  $\overline{D \cap U}$ . Hence, (i) and (ii) are satisfied.

In virtue of Theorem 5.2, it follows from (6.1) that

(6.2) 
$$\operatorname{mes}\left(\Phi^{-1}\left(A \cap A^* \cap G^D \cap \partial(D \cap U)\right)\right) > 0.$$

For any  $m \geq 1$ , let

(6.3) 
$$A_m := \left\{ \eta \in \partial E : \mathcal{A}_{2,m}(\eta) \subset \Phi^{-1}(G) \right\},$$

where  $\mathcal{A}_{2,m}(\eta)$  is given by formula (5.13).

Using the Geometric fact just after (5.14), we see that  $A_m$  is closed. On the other hand, applying Proposition 5.6 leads to

$$\Phi^{-1}\left(A \cap A^* \cap G^D \cap \partial(D \cap U)\right) \subset \bigcup_{m=1}^{\infty} A_m.$$

Therefore, in virtue of (6.2), there is an index  $m_0$  such that

$$\operatorname{mes}\left(A_{m_0} \cap \Phi^{-1}\left(A \cap A^* \cap G^D \cap \partial(D \cap U)\right)\right) > 0.$$

Put  $h := \frac{1}{2m_0}$ . By the latter estimate one may find a closed set F contained in  $A_{m_0} \cap \Phi^{-1} \left( A \cap A^* \cap G^D \cap \partial(D \cap U) \right)$  such that  $\operatorname{mes}(F) > 0$  and  $\sup_{x,y \in F} |x - y| < h$ . Since  $h = \frac{1}{2m_0}$ , a geometric argument shows that

$$\left\{z \in \mathcal{A}_{\frac{\pi}{4}}(\zeta) : |z| > 1 - h\right\} \subset \mathcal{A}_{2,m_0}(\zeta), \qquad \zeta \in \partial E.$$

This together with (6.3) implies that  $\Omega = \Omega(F, h) \subset \Phi^{-1}(G)$ . Hence, (iii) is verified. This completes the proof.

In the sequel, the following uniqueness theorem will play a vital role.

**Theorem 6.4.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a linearly measurable subset A of  $\partial D$  with  $\operatorname{mes}(A) > 0$ . Let  $\mathcal{N} \subset \partial D$  with  $\operatorname{mes}(\mathcal{N}) = 0$ . Let  $0 \leq \epsilon < 1$  and G a connected component of  $D_{\epsilon} := \{z \in D : \omega(z, A, D) < 1 - \epsilon\}$ . If  $f \in \mathcal{O}(G)$  admits the angular limit 0 at every point of  $(A \cap A^* \cap G^D) \setminus \mathcal{N}$ , then  $f \equiv 0$ .

*Proof.* Applying Theorem 6.3 we obtain an open set U in  $\mathbb{C}$ , a conformal mapping  $\Phi$  from E onto  $D \cap U$  which extends homeomorphically to  $\overline{E}$ , and an angular Jordan domain  $\Omega := \Omega(F, h)$  satisfying assertions (i)–(iii) listed in that theorem.

Consider the function  $f \circ \Phi : \Omega \longrightarrow \mathbb{C}$ . By the hypothesis and by Proposition 5.6,  $f \circ \Phi \in \mathcal{O}(\Omega)$  admits the angular limit 0 at a.e point in F. Since  $\operatorname{mes}(F) > 0$ , Theorem 4.3 gives that  $f \circ \Phi \equiv 0$  on  $\Omega$ . Hence,  $f \equiv 0$  on the subdomain  $\Phi(\Omega)$  of G. This proves  $f \equiv 0$ .

6.2. Main result of the section. The boundary behavior of Gonchar–Carleman operator is described below.

**Theorem 6.5.** Let  $D_j$  be the unit disc E and  $A_j$  a linearly measurable subset of  $\partial D_j$  such that  $\operatorname{mes}(A_j) > 0$ ,  $j = 1, \ldots, N$ . Let f be a function defined on  $X := \mathbb{X}(A_1, \ldots, A_N; D_1, \ldots, D_N)$  with the following properties:

- (i)  $f|_A$  is measurable;
- (ii)  $f \in \mathcal{O}_s(X^o);$
- (iii) there exist a constant C > 0 and N functions  $f_j : A \longrightarrow \mathbb{C}, j = 1, ..., N$ , such that for any  $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N), |f(a', \cdot, a'')|_{D_j} < C$ and  $f(a', \cdot, a'')$  has the angular limit  $f_j(a', a_j, a'')$  at  $a_j$  for a.e.  $a_j \in A_j$ , and  $f_1 = \cdots = f_N = f$  a.e. on A.

Let  $0 < \delta_1 < 1, z' = (z_2, \dots, z_N) \in (D_2 \cup (A_2 \cap A_2^*)) \times \dots \times (D_N \cup (A_N \cap A_N^*))$  such that

$$\sum_{j=2}^{N} \omega(z_j, A_j, D_j) < \delta_1,$$

and let G be any connected component of

$$D_{1,\delta_1} := \{ z_1 \in D_1 : \ \omega(z_1, A_1, D_1) < 1 - \delta_1 \}.$$

Then there is an angular Jordan domain  $\Omega = \Omega(F,h)$  such that  $\Omega \subset G$ ,  $F \subset A_1 \cap A_1^* \cap G^{D_1}$ , and the Gonchar-Carleman operator K[f] (see formulas (4.1)-(4.2)) satisfies

$$K[f](a_{1}, z') = \lim_{z_{1} \to a_{1}, z_{1} \in \mathcal{A}_{\alpha}(a_{1})} K[f](z_{1}, z'), \qquad 0 < \alpha < \frac{\pi}{2},$$

for a.e.  $a_1 \in F$ .

6.3. **Preparatory results.** For the proof of Theorem 6.5 we need the following results.

In the sequel, for every function  $f \in L^1(\partial E, |d\zeta|)$ , let  $\mathcal{C}[f]$  denote the Cauchy integral

$$\mathcal{C}[f](z) := \frac{1}{2\pi i} \int_{\partial E} \frac{f(\zeta)d\zeta}{z-\zeta}, \qquad z \in E.$$

For a function  $F: E \longrightarrow \mathbb{C}$ , the radial maximal function  $M_{\text{rad}}F: \partial E \rightarrow [0, \infty]$  is defined by

$$(M_{\mathrm{rad}}F)(\zeta) := \sup_{0 \le r < 1} |F(r\zeta)|, \qquad \zeta \in \partial E.$$

Now we are able to state the following classical result (see Theorem 6.3.1 in Rudin's book [16])

**Theorem 6.6.** (Korányi-Vági type theorem) There is a constant C > 0 such that (changes)

$$\int_{\partial E} |M_{\rm rad} \mathcal{C}[f](\zeta)|^2 |d\zeta| \le C \int_{\partial E} |f(\zeta)|^2 |d\zeta|$$

for every  $f \in L^2(\partial E, |d\zeta|)$ .

We recall the definition of the Smirnov class  $E^p$ , p > 0, on rectifiable Jordan domains.

**Definition 6.7.** Let p > 0 and  $\Omega$  a rectifiable Jordan domain. A function  $f \in \mathcal{O}(\Omega)$ is said to belong to the Smirnov class  $E^p(\Omega)$  if there exists a sequence of rectifiable closed Jordan curves  $(\mathcal{C}_n)_{n=1}^{\infty}$  in  $\Omega$ , tending to the boundary in the sense that  $\mathcal{C}_n$ eventually surrounds each compact subdomain of  $\Omega$ , such that

$$\int_{\mathcal{C}_n} |f(z)|^p |dz| \le M < \infty, \qquad n \ge 1.$$

Next, we rephrase some facts concerning the Smirnov class  $E^p$ , p > 0 on rectifiable Jordan domains in the context of angular Jordan domains  $\Omega(F, h)$ .

**Theorem 6.8.** 1) Let  $\Omega$  be a rectifiable Jordan domain. Then, for every  $f \in E^p(\Omega)$ , p > 0, f admits the angular limit  $f^*$  a.e. on  $\partial\Omega$ .

2) Let  $\Omega := \Omega(F, h)$  be an angular Jordan domain and let  $\Gamma := \partial \Omega$ . For any  $0 < \epsilon < \frac{h}{4}$ , let  $\Gamma_{\epsilon}$  be the rectifiable closed Jordan curve defined in Part 4) of Proposition 6.2. Then  $f \in E^p(\Omega)$  if  $\sup_{0 < \epsilon < \frac{h}{4}\Gamma_{\epsilon}} \int |f(z)|^p |dz| < \infty$ . In addition, for an  $f \in E^p(\Omega)$ , p > 0,

it holds that

$$\int_{\Gamma} |f^*(z)|^p |dz| \le \sup_{0 < \epsilon < \frac{h}{4}} \int_{\Gamma_{\epsilon}} |f(z)|^p |dz|.$$

3) Every  $f \in E^1(E)$  has a Cauchy representation  $f := \mathcal{C}[f^*]$ . Conversely, if  $g \in L^1(\partial E, |dz|)$  and

$$\int_{\partial E} z^n g(z) dz = 0, \qquad n = 0, 1, 2, \dots,$$

then  $f := \mathcal{C}[g] \in E^1(E)$  and g coincides with  $f^*$  a.e. on  $\partial E$ .

*Proof.* For the proof of Parts 1) and 3), see [4, p. 438–441]. Taking into account Parts 6) and 7) of Proposition 6.2, Part 2) also follows from the results in [4, p. 438–441]. Hence, the proof is complete.  $\Box$ 

6.4. **Proof of Theorem 6.5.** We only give the proof of this theorem in the case N = 2. The proof for the general case is quite similar, therefore, it is left to the interested reader.

We fix an arbitrary  $z_2^0 \in D_2 \cup (A_2 \cap A_2^*)$ ,  $0 < \delta_2 < \delta_1$  such that  $\omega(z_2^0, A_2, D_2) < \delta_2$ , and an arbitrary connected component G of  $D_{1,\delta_1} :=$ 

 $\{z_1 \in D_1 : \omega(z_1, A_1, D_1) < 1 - \delta_1\}$ . Applying Theorem 6.3, we may find an angular Jordan domain  $\Omega := \Omega(F, h) \subset G$  such that  $F \subset A \cap A^* \cap G^{D_1}$ . Without loss of generality we may assume that  $z_2^0 \in D_2$ , since the case where  $z_2^0 \in A_2 \cap A_2^*$  is trivial because of (iii). In the course of the proof, the letter C will denote a positive constant that is not necessarily the same at each step.

Applying Theorem 4.1, we have

$$f(z_1, a_2) = \lim_{M \to \infty} K_M[f|_{A_1 \times \{a_2\}}](z_1), \qquad z_1 \in D_1, \ a_2 \in A_2,$$
  
$$f(a_1, a_2) = \lim_{r \to 1^-} f(ra_1, a_2), \qquad a_1 \in \partial D_1, \ a_2 \in A_2.$$

Consequently,  $f|_{\partial D_1 \times A_2}$  is measurable. In addition, by (iii) this function is bounded. Therefore, for every  $M \in \mathbb{N}$  we are able to define the function  $K_{\infty,M}(\cdot, z_2^0) : \partial D_1 \longrightarrow \mathbb{C}$ ,

(6.4) 
$$K_{\infty,M}(a_1, z_2^0) := \frac{1}{2\pi i} \int_{A_2} e^{M(g_2(z_2^0) - g_2(a_2))} \frac{f(a_1, a_2) da_2}{a_2 - z_2^0}, \qquad a_1 \in \partial D_1.$$

Since, in virtue of (ii)–(iii),  $f(a_1, \cdot) \in \mathcal{O}(D_2)$  and  $|f(a_1, \cdot)|_{D_2} < C$  for  $a_1 \in A_1$ , it follows from Theorem 4.1 that

(6.5) 
$$\lim_{M \to \infty} K_{\infty,M}(a_1, z_2^0) = f(a_1, z_2^0), \qquad a_1 \in A_1,$$

and the above convergence is uniform with respect to  $a_1 \in A_1$ .

On the other hand, by (6.4) we see that  $K_{\infty,M}(\cdot, z_2^0)$  is measurable and bounded. In addition, for any  $n = 0, 1, 2, \ldots$ , taking (ii) into account, we have that

$$\int_{\partial D_1} K_{\infty,M}(a_1, z_2^0) a_1^n da_1 = \frac{1}{2\pi i} \int_{A_2} \left( \int_{\partial D_1} f(a_1, a_2) a_1^n da_1 \right) \frac{e^{M(g_2(z_2^0) - g_2(a_2))} da_2}{a_2 - z_2^0} = 0,$$

where the first equality follows from an application of Fubini's Theorem and the second one from an application of Part 3) of Theorem 6.8 to  $f(\cdot, a_2)$ ,  $a_2 \in A_2$ . Consequently, in virtue of Part 3) of Theorem 6.8, we can extend  $K_{\infty,M}(\cdot, z_2^0)$  to  $\overline{D}_1$  by setting

(6.6) 
$$K_{\infty,M}(z_1, z_2^0) := \mathcal{C}[K_{\infty,M}(\cdot, z_2^0)](z_1) = \frac{1}{2\pi i} \int_{\partial D_1} \frac{K_{\infty,M}(a_1, z_2^0) da_1}{a_1 - z_1}, \qquad z_1 \in D_1.$$

Then the following identity holds

(6.7) 
$$\lim_{z_1 \to a_1, \ z_1 \in \mathcal{A}_{\alpha}(a_1)} K_{\infty,M}(z_1, z_2^0) = K_{\infty,M}(a_1, z_2^0), \qquad 0 < \alpha < \frac{\pi}{2},$$

for a.e.  $a_1 \in \partial D_1$ .

Now we come back to the angular Jordan domain  $\Omega$ . We keep the notation introduced in Proposition 6.2. Put K := K[f] and  $K_{M,M} := K_{M,M}[f|_A]$  (see formulas (4.1)–(4.2)). For any  $0 < \epsilon < \frac{h}{4}$  and any  $z_1 \in \Gamma_{\epsilon}$ , applying the Cauchy integral formula, we obtain

(6.8) 
$$K_{\infty,M}(z_1, z_2^0) - K_{M,M}(z_1, z_2^0) = \frac{1}{(2\pi i)^2} \int_{\partial D_1 \setminus A_1} \int_{A_2} e^{M(g_1(z_1) - g_1(t_1)) + M(g_2(z_2^0) - g_2(t_2))} \frac{f(t_1, t_2) dt_1 dt_2}{(t_1 - z_1)(t_2 - z_2^0)} = e^{M(g_1(z_1) - (1 - \delta_1))} \int_{\partial D_1} \frac{p_N(t_1) dt_1}{t_1 - z_1}$$

Using the choice of G and the hypothesis on  $\delta_1$  and  $\delta_2$ , it can be checked that

(6.9)  $|e^{M(g_1(\cdot)-(1-\delta_1))}|_G \le 1, \qquad |p_N|_{\partial D_1} \le Ce^{-M(\delta_1-\delta_2)}.$ 

Therefore, recalling the projection  $\tau : E \setminus \{0\} \longrightarrow \partial E$  is defined in Part 5) of Proposition 6.2, we estimate

$$(6.10) \int_{\Gamma_{\epsilon}} |K_{\infty,M}(z_{1}, z_{2}^{0}) - K_{M,M}(z_{1}, z_{2}^{0})|^{2} |dz_{1}| \leq C \int_{\Gamma_{\epsilon}} |M_{\mathrm{rad}} \mathcal{C}[p_{N}](\tau(z_{1}))|^{2} |dz_{1}|$$

$$\leq 10C \int_{\tau(F_{0\epsilon} \cup F_{1\epsilon} \cup F_{2\epsilon})} |M_{\mathrm{rad}} \mathcal{C}[p_{N}](a_{1})|^{2} |da_{1}| + 10C \int_{\tau(F_{3\epsilon})} |M_{\mathrm{rad}} \mathcal{C}[p_{N}](a_{1})|^{2} |da_{1}|$$

$$\leq 20C \int_{\partial E} |M_{\mathrm{rad}} \mathcal{C}[p_{N}](a_{1})|^{2} |da_{1}| \leq C \int_{\partial E} |p_{N}(a_{1})|^{2} |da_{1}| \leq Ce^{-M(\delta_{1} - \delta_{2})}.$$

where the first estimate follows from (6.8)–(6.9) and the definition of the radial maximal function, the second and the third one from Part 5) of Proposition 6.2, the fourth estimate holds by an application of Theorem 6.6, and the last one follows from (6.9).

On the other hand, for any  $0 < \epsilon < \frac{h}{4}$ ,

(6.11) 
$$\int_{\Gamma_{\epsilon}} |K_{M+1,M+1}(z_1, z_2^0) - K_{M,M}(z_1, z_2^0)|^2 |dz_1| \\ \leq 2 \int_{\Gamma_{\epsilon}} |K_M^1(z_1, z_2^0)|^2 |dz_1| + 2 \int_{\Gamma_{\epsilon}} |K_M^2(z_1, z_2^0)|^2 |dz_1| \leq Ce^{-M\delta_0},$$

where the latter estimate follows from the same argument as in the proof of (6.8)–(6.10). We recall from Part 1) of Theorem 4.2 that

$$\lim_{M \to \infty} K_{M,M}(z_1, z_2^0) = K(z_1, z_2^0), \qquad z_1 \in \Gamma_{\epsilon},$$

and that the convergence is uniform with respect to  $z_1 \in \Gamma_{\epsilon}$ . This, combined with (6.10)–(6.11), implies that

(6.12) 
$$\int_{\Gamma_{\epsilon}} |K_{\infty,M}(z_1, z_2^0) - K(z_1, z_2^0)|^2 |dz_1| \le C \cdot e^{-M(\delta_1 - \delta_2)}, \quad 0 < \epsilon < \frac{h}{4}.$$

Since we have already shown that  $|K_{\infty,M}(\cdot, z_2^0)|_{D_1} < \infty$ , in virtue of Part 2) of Theorem 6.8, we deduce from (6.12) that  $K(\cdot, z_2^0)|_{\Omega} \in E^2(\Omega)$ . For every  $a_1 \in \partial D_1$ , let  $K(a_1, z_2^0)$  denote the angular limit of  $K(\cdot, z_2^0)|_{\Omega}$  at  $a_1$  (if the limit exists). It follows from (6.12) and Part 2) of Theorem 6.8 that

$$\lim_{M \to \infty} \int_{\Gamma} |K_{\infty,M}(a_1, z_2^0) - K(a_1, z_2^0)|^2 |da_1|$$
  
$$\leq \sup_{0 < \epsilon < \frac{h}{4}} \int_{\Gamma_{\epsilon}} |K_{\infty,M}(z_1, z_2^0) - K(z_1, z_2^0)|^2 |dz_1| \leq \lim_{M \to \infty} C \cdot e^{-M(\delta_1 - \delta_2)} = 0.$$

This, combined with (6.5) and Part 8) of Proposition 6.2, implies finally that

$$K(a_1, z_2^0) = f(a_1, z_2^0),$$
 for a.e.  $a_1 \in F.$ 

Hence, Theorem 6.5 has been proved.

### 7. Proof of Theorem A in a special case

In this section, we prove Theorem A under the following hypotheses:

 $D_1, \ldots, D_N$  are rectifiable Jordan domains and  $|f|_X < \infty$ . (\*)

We first prove that Theorem A under the above hypotheses may be reduced to the case when  $D_1 = \cdots = D_N = E$ . Indeed, let  $\Phi_j$  be a conformal mapping from  $D_j$ onto E which extends to a homeomorphic map (still denoted by)  $\Phi$  from  $\overline{D_j}$  onto  $\overline{E}$  $j = 1, \ldots, N$ . Consider the function  $\tilde{f}$  given by

$$\tilde{f}(w) := f\left(\Phi_1^{-1}(w_1), \dots, \Phi_N^{-1}(w_N)\right), w = (w_1, \dots, w_N) \in \mathbb{X}\left(\Phi_1(A_1), \dots, \Phi_N(A_N); E, \dots, E\right).$$

Then we deduce easily from the hypothesis on f that  $\tilde{f}$  satisfies the hypotheses (i)–(iii) on the cross  $\mathbb{X}(\Phi_1(A_1),\ldots,\Phi_N(A_N);E,\ldots,E)$ . Let  $\hat{f} \in \mathcal{O}(\widehat{\mathbb{X}}^{\circ}(\Phi_1(A_1),\ldots,\Phi_N(A_N);E,\ldots,E))$  denote the extension function of  $\tilde{f}$  provided by Theorem A in the case where  $D_1 = \cdots = D_N = E$ . Using Proposition 5.6 and Part 1) of Theorem 5.2, we see that the function  $\hat{f}$  defined by

$$\hat{f}(z) := \hat{f}(\Phi_1(z_1), \dots, \Phi_N(z_N)), z = (z_1, \dots, z_N) \in \hat{X}^{\circ},$$

satisfies the conclusions of Parts 1)-4). This proves the above reduction.

We give here only the proof of Theorem A for the case N = 2. Since the general case is quite similar and does not require any new ideas, it is therefore left to the interested reader. In summary, from now on

we assume that  $N = 2, D_1 = D_2 = E, and |f|_X < \infty.$  (\*)

Using hypotheses (i)–(iii) and  $\binom{*}{*}$ , we are able to apply Theorem 6.5 and obtain a function  $K[f] \in \mathcal{O}(\widehat{X})$  (see the notation in Subsection 2.3). We define the desired extension function  $\hat{f}$  as follows

(7.1) 
$$\hat{f} := K[f] \quad \text{on } \hat{X} \setminus (A_1 \times A_2)$$

The remaining part of the proof is divided into several steps.

**Step 1:** *Proof of the estimate* 

$$|\hat{f}|_{\hat{X}^o} \le |f|_X.$$

Proof of Step 1. Let  $z^0 = (z_1^0, z_2^0)$  be an arbitrary point of  $\widehat{X}^\circ$ . Then we may find an  $\delta_1 \in (0, 1)$  such that  $0 < \omega(z_2^0, A_2, D_2) < \delta_1 < 1 - \omega(z_1^0, A_1, D_1)$ . Let G be the connected component of  $D_{1,\delta_1} := \{z_1 \in D_1 : \omega(z_1, A_1, D_1) < 1 - \delta_1\}$  that contains  $z_1^0$ . By Theorem 6.3 we may find an angular Jordan domain  $\Omega := \Omega(F, h)$  contained in G such that  $F \subset A_1 \cap A_1^* \cap G^{D_1}$ . In addition, for every  $M \in \mathbb{N}$ , applying Theorem 6.5 to the function  $f^M$ , we obtain the function  $K[f^M] \in \mathcal{O}(\widehat{X}^\circ)$  with the following property

$$\lim_{z_1 \to a_1, z_1 \in \mathcal{A}_{\alpha}(a_1)} K[f^M](z_1, z_2^0) = f(a_1, z_2^0)^M$$
$$= \lim_{z_1 \to a_1, z_1 \in \mathcal{A}_{\alpha}(a_1)} \left( K[f](z_1, z_2^0) \right)^M, \ 0 < \alpha < \frac{\pi}{2},$$

for a.e.  $a_1 \in F$ .

Consequently, arguing as in the proof of Theorem 6.4, we get

$$K[f^{M}](z^{0}) = \left(K[f](z^{0})\right)^{M}, \ M \in \mathbb{N},$$

Since  $z_0 \in \widehat{X}^{\circ}$  is arbitrarily chosen, it follows from the latter identity that

(7.2) 
$$K[f^M](z) = \left(K[f](z)\right)^M, \ M \in \mathbb{N}, \ z \in \widehat{X}^{\circ}.$$

Now we are able to conclude the proof in the same way as in [6, p. 23]. More precisely, taking into account (7.1)–(7.2) and Part 2) of Theorem 4.2, one gets that

$$|\hat{f}^M(z)| \le |K[f^M](z)| \le \frac{C_0|f|_X^M}{D_1)\operatorname{dist}(z_2,\partial D_2)(1-e^{-(1-\omega(z))})}, \ z=(z_1,z_2)\in\widehat{X}^\circ.$$

• •

Extracting the *M*th roots of both sides and letting *M* tend to  $\infty$ , the desired estimate of Step 1 follows.

**Step 2:** We shall prove that  $\hat{f}$  is the unique function  $\mathcal{O}(\hat{X}^o)$  which verifies Property 1).

Proof of Step 2. First we show that the function  $\hat{f}$  given by (7.1) satisfies Property 1). Without loss of generality, it suffices to prove that there is a subset  $\tilde{A}_2$  of  $A_2 \cap A_2^*$ such that  $\operatorname{mes}(\tilde{A}_2) = \operatorname{mes}(A_2)$  and  $\hat{f}$  admits the angular limit f at every point of  $D_1 \times \tilde{A}_2$ .

For any  $a_1 \in A_1$  put

$$A_{a_1}' := \{a_2 \in A_2 : f(a_1, \cdot) \text{ has an angular limit at } a_2\}.$$

By hypothesis (iii), we have  $mes(A'_{a_1}) = mes(A_2), a_1 \in A_1$ . Consequently, applying Fubini's Theorem, we obtain that

$$\int_{A_1} \operatorname{mes}(A'_{a_1}) |da_1| = \operatorname{mes}(A_1) \operatorname{mes}(A_2) = \int_{A_2} \operatorname{mes}\left(\left\{a_1 \in A_1 : a_2 \in A'_{a_1}\right\}\right) |da_2|$$

Hence,

(7.3) 
$$\max\left(\left\{a_1 \in A_1 : a_2 \in A'_{a_1}\right\}\right) = \max(A_1) \quad \text{for a.e. } a_2 \in A_2.$$

The same reasoning also gives that

(7.4) 
$$\operatorname{mes}\left(\{a_1 \in A_1 : f(a_1, a_2) = f_1(a_1, a_2)\}\right) = \operatorname{mes}(A_1)$$
 for a.e.  $a_2 \in A_2$ .  
Set

(7.5) 
$$\tilde{A}_2 := \left\{ a_2 \in A_2 \cap A_2^* : \operatorname{mes} \left( \left\{ a_1 \in A_1 : a_2 \in A_{a_1}' \right\} \right) = \operatorname{mes}(A_1) \\ \operatorname{and} \operatorname{mes} \left( \left\{ a_1 \in A_1 : f(a_1, a_2) = f_1(a_1, a_2) \right\} \right) = \operatorname{mes}(A_1) \right\}$$

We deduce from (7.3)–(7.5) that

(7.6) 
$$\operatorname{mes}(\tilde{A}_2) = \operatorname{mes}(A_2).$$

Fix an arbitrary point  $a_2^0 \in \tilde{A}_2$  and let  $(z_2^n)_{n=1}^\infty$  be an arbitrary sequence of  $D_2$  such that  $\lim_{n\to\infty} z_2^n = a_2^0$  and  $z_2^n \in \mathcal{A}_\alpha(a_2^0)$  for some fixed number  $0 < \alpha < \frac{\pi}{2}$ . Fix an arbitrary point  $z_1^0$  of  $D_1$  and let  $(z_1^n)_{n=1}^\infty$  be an arbitrary sequence of  $D_1$  such that  $\lim_{n \to \infty} z_1^n = z_1^0.$ Clearly, we may find  $0 < \delta_1 < 1$  such that

(7.7) 
$$\sup_{n \in \mathbb{N}} \omega(z_1^n, A_1, D_1) < 1 - \delta_1.$$

Fix an  $\delta_2$  such that  $0 < \delta_2 < \delta_1$ . Since  $a_2^0$  is locally regular relative to  $A_2$  and  $\lim_{n \to \infty} z_2^n = a_2^0$  and  $z_2^n \in \mathcal{A}_{\alpha}(a_2^0)$ , there is a sufficiently large number  $N_0$  with

(7.8) 
$$\omega(z_2^n, A_2, D_2) < \delta_2, \qquad n > N_0.$$

Let G be that connected component of the following open set

$$D_{1,\delta_1} := \{ z_1 \in D_1 : \ \omega(z_1, A_1, D_1) < 1 - \delta_1 \}$$

which contains  $z_1^0$  (see (7.7)). Applying Theorem 6.3, we may find an angular Jordan domain  $\Omega := \Omega(F, h)$  contained in G such that  $F \subset A_1 \cap A_1^* \cap G^{D_1}$ . Let V be a rectifiable Jordan domain with  $\Omega \subset V \subset G$ ,  $z_2^0 \in V$ , and U some neighborhood of the base F of  $\Omega$  with  $V \cap U = \Omega \cap U$ .

In virtue of (7.8) and of the fact that  $V \subset G \subset D_{1,\delta_1}$ , we obtain that

(7.9) 
$$V \times \{z_2^n\} \subset \widehat{X}^{\mathbf{o}}, \qquad n > N_0$$

Consequently, Theorem 6.5 yields that for any  $n > N_0$ ,

(7.10) 
$$f(a_1, z_2^n) = \lim_{z_1 \to a_1, \ z_1 \in \mathcal{A}_{\alpha}(a_1)} \hat{f}(z_1, z_2^n), \qquad 0 < \alpha < \frac{\pi}{2},$$

for a.e.  $a_1 \in F$ .

Next, for any  $n > N_0$  let

$$F_{n} := \left\{ a_{1} \in F : a_{2}^{0} \in A_{a_{1}}^{'} \text{ and } f(a_{1}, z_{2}^{n}) = \lim_{z_{1} \to a_{1}, z_{1} \in \mathcal{A}_{\alpha}(a_{1})} \hat{f}(z_{1}, z_{2}^{n}) \right\},$$
  
$$F_{0} := \bigcap_{n=N_{0}+1}^{\infty} F_{n}.$$

It follows from (7.5), (7.10) and the fact that  $a_2^0 \in \tilde{A}_2$  that  $\operatorname{mes}(F_n) = \operatorname{mes}(F)$ ,  $n > N_0$ . Hence

(7.11) 
$$mes(F_0) = mes(F) > 0.$$

In virtue of (7.9), consider the following holomorphic functions on V

(7.12) 
$$h_n(t) := \hat{f}(t, z_2^n) \text{ and } h_0(t) := f(t, a_2^0), \quad t \in V, \ n > N_0.$$

Since we have already shown in Step I that  $|h_n|_V \leq |f|_X < \infty$ ,  $n > N_0$  or n = 0, applying Part 1) of Theorem 6.8, we may find a subset  $\Delta$  of  $F_0$  with  $\operatorname{mes}(\Delta) = \operatorname{mes}(F_0) > 0$  such that  $h_n$ ,  $n > N_0$  (resp.  $h_0$ ) admits the angular limit  $f_1(t, z_2^n)$ (resp.  $f_1(t, a_2^0)$ ) at  $t \in \Delta$ . Observe that by (7.5) and the fact that  $a_2^0 \in \tilde{A}_2$  we have that

$$\lim_{n \to \infty} f_1(t, z_2^n) = f_1(t, a_2^0) = f(t, a_2^0) \quad \text{for a.e. } t \in \Delta.$$

Using this and (7.12), we are able to apply Lemma 4.4 to the sequence  $(h_n)_{n=0}^{\infty}$ . Consequently, one gets

$$\lim_{n \to \infty} \hat{f}(z_1^n, z_2^n) = f(z_1^0, a_2^0)$$

This shows that  $\hat{f}$  admits the angular limit f at every point of  $D_1 \times \tilde{A}_2$ . Hence,  $\hat{f}$  satisfies Property 1).

In order to complete Step 2 we need to show the uniqueness of  $\hat{f}$ . To do this, let  $\hat{f} \in \mathcal{O}(\hat{X}^{\circ})$  be a function with the following property: There is a subset  $\tilde{A}_j$  of  $A_j \cap A_j^*$  (j = 1, 2) such that  $\operatorname{mes}(A_j \setminus \tilde{A}_j) = 0$  and  $\hat{f}$  admits the angular limit f at every point of  $(\tilde{A}_1 \times D_2) \cup (D_1 \times \tilde{A}_2)$ . Fix an arbitrary point  $z^0 = (z_1^0, z_2^0) \in \hat{X}^{\circ}$ . Let G be the connected component containing  $z_1^0$  of the following open set

$$\{z_1 \in D_1 : \omega(z_1, A_1, D_1) < 1 - \omega(z_2^0, A_2, D_2)\}$$

We deduce from the property of  $\hat{f}$  and  $\hat{f}$  that both holomorphic functions  $\hat{f}(\cdot, z_2^0)|_G$ and  $\hat{f}(\cdot, z_2^0)|_G$  admit the angular limit  $f(\cdot, z_2^0)$  at every point of  $\tilde{A}_1 \cap \tilde{A}_1 \cap G^{D_1}$ . Consequently, applying Theorem 6.4 yields that  $\hat{f}(\cdot, z_2^0) = \hat{f}(\cdot, z_2^0)$  on G. Hence,  $\hat{f}(z^0) = \hat{f}(z^0)$ . Since  $z^0 \in \hat{X}^\circ$  is arbitrary, the uniqueness of  $\hat{f}$  is established. This completes Step 2.

Step 3: Proof of Part 2).

Proof of Step 3. Fix  $(z_1^0, z_2^0) \in \widehat{X}^{\circ}$ . For every  $a_2 \in A_2$  we have

$$|f(a_1, a_2)| \le |f|_{A_1 \times A_2}, a_1 \in A_1,$$
 and  $|f(z_1, a_2)| \le |f|_X, z_1 \in D_1.$ 

Therefore, the Two-Constant Theorem (see Theorem 2.2 in [14]) implies that

(7.13) 
$$|f(z_1, a_2)| \le |f|_{A_1 \times A_2}^{1-\omega(z_1, A_1, D_1)} |f|_X^{\omega(z_1, A_1, D_1)}, \quad z_1 \in D_1, \ a_2 \in A_2$$

Let  $\delta := \omega(z_1^0, A_1, D_1)$  and consider the  $\delta$ -level set

$$D_{2,\delta} := \{ z_2 \in D_2 : \omega(z_2, A_2, D_2) < 1 - \delta \}.$$

Clearly,  $z_2^0 \in D_{2,\delta}$ .

Recall from Step 2 that  $\tilde{A}_2 \subset A_2 \cap A_2^*$ , mes  $\left( (A_2 \cap A_2^*) \setminus \tilde{A}_2 \right) = 0$ , and

(7.14) 
$$f(z_1^0, a_2) = \lim_{z_2 \to a_2, \ z_2 \in \mathcal{A}_\alpha(a_2)} \hat{f}(z_1^0, z_2), \qquad 0 < \alpha < \frac{\pi}{2}, \ a_2 \in \tilde{\mathcal{A}}_2.$$

Consider the following function  $h: D_{2,\delta} \cup \tilde{A}_2 \longrightarrow \mathbb{C}$  defined by

(7.15) 
$$h(t) := \begin{cases} \hat{f}(z_1^0, t), & t \in D_{2,\delta} \\ f(z_1^0, t), & t \in \tilde{A}_2 \end{cases}.$$

Clearly,  $h|_{D_{2,\delta}} \in \mathcal{O}(D_{2,\delta})$ .

On the other hand, in virtue of (7.15) and the result of Step 1, we have

(7.16) 
$$|h|_{D_{2,\delta}} \le |\hat{f}|_{\hat{X}^0} \le |f|_X < \infty.$$

In addition, applying Corollary 5.20 and taking (7.14)-(7.15) into account yields

$$|h(t)| \le |h|_{\tilde{A}_2}^{1-\omega_{\delta}(t,A_1,D_1)} |h|_{D_{2,\delta}}^{\omega_{\delta}(t,A_1,D_1)}, \quad t \in D_{2,\delta},$$

where, by Theorem 5.19,

$$\omega_{\delta}(t, A_2, D_2) = \frac{\omega(t, A_2, D_2)}{1 - \omega(z_1^0, A_1, D_1)}.$$

This, combined with (7.13)-(7.16), implies that

$$|\hat{f}(z_1^0, z_2^0)| = |h(z_2^0)| \le |f|_{A_1 \times A_2}^{1-\omega(z_1^0, A_1, D_1) - \omega(z_2^0, A_2, D_2)} |f|_X^{\omega(z_1^0, A_1, D_1) + \omega(z_2^0, D_2, A_2)}.$$

Hence Part 2) for the point  $(z_1^0, z_2^0)$  is proved.

Step 4: Proof of Part 3).

Proof of Step 4. Let  $(a_1^0, z_2^0) \in A_1^* \times D_2$  be such that the following limit exists

$$\lambda := \lim_{(a_1, z_2) \to (a_1^0, z_2^0), \ (a_1, z_2) \in A_1 \times D_2} f(a_1, z_2).$$

We like to show that  $\hat{f}$  admits the angular limit  $\lambda$  at  $(a_1^0, z_2^0)$ .

In virtue of assumption  $\binom{*}{*}$ , we may suppose (without loss of generality) that  $|f|_X < 1$ . For any  $0 < \epsilon < \frac{1}{2}$ , we may find an open neighborhood  $A_{a_1^0}$  of  $a_1^0$  in  $A_1$  and a positive number r > 0 such that  $\mathbb{B}(z_2^0, r) \in D_2$  and

(7.17) 
$$|f(a_1, z_2) - \lambda| < \epsilon^2, \quad a_1 \in A_{a_1^0}, \ |z_2 - z_2^0| \le r.$$

38

Put

(7.18) 
$$\delta := \sup_{z_2 \in \overline{\mathbb{B}(z_2^0, r)}} \omega(z_2, A_2, D_2).$$

Since  $a_1^0 \in A_1^*$ , it follows from Part 2) of Proposition 5.9 that  $\operatorname{mes}(A_{a_1^0}) > 0$ . Next, consider the level set

$$D_{1,\delta} := \left\{ z_1 \in D_1 : \ \omega(z_1, A_{a_1^0}, D_1) < 1 - \delta \right\}.$$

In virtue of (7.18), we can define

(7.19) 
$$h(t, z_2) := \hat{f}(t, z_2) - \lambda, \quad t \in D_{1,\delta}, \ z_2 \in \overline{\mathbb{B}(z_2^0, r)}.$$

Clearly,

(7.20) 
$$|h|_{D_{1,\delta}} \le 2|\hat{f}|_{\hat{X}^0} = 2|f|_X = 2.$$

By (7.19) and using the result of Step 2, we know that for every  $z_2 \in \mathbb{B}(z_2^0, r)$  the holomorphic function  $h(\cdot, z_2)|_{D_{1,\delta}}$  admits the angular limit  $f(a_1, z_2) - \lambda$  at  $a_1$  for  $a_1 \in \tilde{A}_1 \cap A_{a_1^0}$ , where  $\tilde{A}_1$  is given in Step 2. Consequently, applying Corollary 5.20 and taking (7.17) and (7.20) into account, we see that

$$|h(t, z_2)| < \epsilon^{2(1-\omega_{\delta}(t, A_{a_1^0}, D_1))} 2^{\omega_{\delta}(t, A_{a_1^0}, D_1)}, \qquad t \in D_{1, \delta}.$$

Let  $0 < \alpha < \frac{\pi}{2}$ . In virtue of Theorem 5.19 and the hypothesis that  $a_1^0 \in A_1^*$ , we deduce that  $\lim_{t \to a_1^0, t \in \mathcal{A}_{\alpha}(a_1^0)} \omega_{\delta}(t, A_{a_1^0}, D_1) = 0$ . Consequently, there is an  $r_{\alpha} > 0$  such that

$$|f(z_1, z_2) - \lambda| = |h(z_1, z_2)| < \epsilon, \qquad z_1 \in \mathcal{A}_{\alpha}(a_1^0) \cap \{|z_1 - a_1^0| < r_{\alpha}\}, \ z_2 \in \mathbb{B}(z_2^0, r).$$

This completes the above assertion.

Similarly, we can prove that f admits the angular limit

$$\lim_{(z_1,a_2)\to(z_1^0,a_2^0),\ (z_1,a_2)\in D_1\times A_2} f(z_1,a_2)$$

at any point  $(z_1^0, a_2^0)$ , if the latter limit exists. Hence the proof of Step 4 (i.e. Part 3)) is finished.

Step 5: Proof of Part 4).

Proof of Step 5. Let  $(a_1^0, a_2^0) \in A_1^* \times A_2^*$  be such that the following limit exists

$$\lambda := \lim_{(a_1, a_2) \to (a_1^0, a_2^0), \ (a_1, a_2) \in A_1 \times A_2} f(a_1, a_2).$$

We like to show that  $\hat{f}$  admits the angular limit  $\lambda$  at  $(a_1^0, a_2^0)$ .

Suppose without loss of generality that  $|f|_X < 1$ , and fix an arbitrary  $0 < \epsilon < \frac{1}{2}$ . Since  $(a_1^0, a_2^0) \in A_1^* \times A_2^*$ , we may find an open neighborhood  $A_{a_1^0}$  of  $a_1^0$  in  $A_1$  (resp. an open neighborhood  $A_{a_2^0}$  of  $a_2^0$  in  $A_2$ ) such that

(7.21) 
$$|f(a_1, a_2) - \lambda| < \epsilon^2, \qquad a_1 \in A_{a_1^0}, \ a_2 \in A_{a_2^0}.$$

By Part 2) of Proposition 5.9, one gets  $\operatorname{mes}(A_{a_1^0}) > 0$  and  $\operatorname{mes}(A_{a_2^0}) > 0$ .

Consider the function

(7.22) 
$$h(z_1, z_2) := f(z_1, z_2) - \lambda, \quad (z_1, z_2) \in \mathbb{X}(A_{a_1^0}, A_{a_2^0}; D_1, D_2).$$

Clearly,

(7.23) 
$$|h(z_1, z_2)| \le 2, \quad (z_1, z_2) \in \mathbb{X}(A_{a_1^0}, A_{a_2^0}; D_1, D_2).$$

Applying the results of Steps 1-3 to h, we obtain the function

(7.24) 
$$\hat{h} := K[h]$$
 on  $\hat{\mathbb{X}}^0(A_{a_1^0}, A_{a_2^0}; D_1, D_2).$ 

so that  $\hat{h}$  admits the angular limit h on  $(\tilde{A}_{a_1^0} \times D_2) \cup (D_1 \times \tilde{A}_{a_1^0})$ , where  $\tilde{A}_{a_1^0}$ ,  $\tilde{A}_{a_1^0}$  are given by Step 2. Clearly,

$$\widehat{\mathbb{X}}(A_{a_1^0}, A_{a_2^0}; D_1, D_2) \subset \widehat{\mathbb{X}}(A_1, A_2; D_1, D_2).$$

Consequently, arguing as in Step 1 and taking into account the above mentioned angular limit of  $\hat{h}$ , we conclude that

$$\hat{h} = \hat{f} - \lambda \qquad \text{on } \widehat{\mathbb{X}}(A_{a_1^0}, A_{a_2^0}; D_1, D_2).$$

Consequently, applying Step 3 and taking into account (7.21)–(7.24) and the inequality  $|f|_X < 1$ , we see that

$$\begin{split} \left| \hat{f}(z_1, z_2) - \lambda \right| &= \left| \hat{h}(z_1, z_2) \right| \le \left| h \right|_{A_{a_1^0} \times A_{a_2^0}}^{1 - \omega(z_1, A_{a_1^0}, D_1) - \omega(z_2, A_{a_2^0}, D_2)} (2|f|_X)^{\omega(z_1, A_{a_1^0}, D_1) + \omega(z_2, A_{a_2^0}, D_2)} \\ &< \epsilon^{2 \left( 1 - \omega(z_1, A_{a_1^0}, D_1) - \omega(z_2, A_{a_2^0}, D_2) \right)} 2^{\omega(z_1, A_{a_1^0}, D_1) + \omega(z_2, A_{a_2^0}, D_2)}. \end{split}$$

Therefore, for all  $z = (z_1, z_2) \in \widehat{\mathbb{X}}(A_{a_1^0}, A_{a_2^0}; D_1, D_2)$  satisfying

(7.25) 
$$\omega(z_1, A_{a_1^0}, D_1) + \omega(z_2, A_{a_2^0}, D_2) < \frac{1}{3},$$

we deduce from the latter estimate that

(7.26) 
$$\left| \hat{f}(z) - \lambda \right| < \epsilon.$$

Since  $a_1^0$  (resp.  $a_2^0$ ) is locally regular relative to  $A_{a_1^0}$  (resp.  $A_{a_2^0}$ ), there is an  $r_{\alpha} > 0$  such that (7.25) is fulfilled for

$$z = (z_1, z_2) \in \left( \mathcal{A}_{\alpha}(a_1^0) \cap \{ |z_1 - a_1^0| < r_{\alpha} \} \right) \times \left( \mathcal{A}_{\alpha}(a_2^0) \cap \{ |z_2 - a_2^0| < r_{\alpha} \} \right).$$

This, combined with (7.26), completes the proof. Hence Step 5 (i.e. Part 4)) is finished.  $\hfill \Box$ 

# Step 6: Proof of Part 5).

Proof of Step 6. In virtue of Step 5, we only need to show that  $\hat{f}$  admits the angular limit f on  $(A_1^* \times D_2) \cup (D_1 \times A_2^*)$ . To do this let  $(a_1^0, z_2^0) \in A_1^* \times D_2$  and choose an arbitrary  $0 < \epsilon < 1$ . Fix a compact subset K of  $A_2 \cap A_2^*$  such that  $\operatorname{mes}(K) > 0$  and a sufficiently large N such that

(7.27) 
$$\epsilon^{N(1-\omega(z_2^0,A_2,D_2))}(2|f|_X)^{\omega(z_2^0,A_2,D_2)} < \frac{\epsilon}{2}.$$

Using the hypothesis that f can be extended to a continuous function on  $A_1^* \times A_2^*$ , we may find an open neighborhood  $A_{a_1^0}$  of  $a_1^0$  in  $A_1^*$  such that

(7.28) 
$$|f(a_1, a_2) - f(a_1^0, a_2)| \le \epsilon^N, \quad a_1 \in A_{a_1^0} \cap A_{a_1^0}^*, \ a_2 \in K.$$

On the other hand,

(7.29) 
$$\left| f(a_1, z_2) - f(a_1^0, z_2^0) \right| \le 2|f|_X < \infty, \quad a_1 \in A_{a_1^0} \cap A_{a_1^0}^*, \ z_2 \in D_2.$$

For  $a_1 \in A_{a_1^0} \cap A_{a_1^0}^*$ , applying the Two-Constant Theorem to the function  $f(a_1, \cdot) - f(a_1^0, \cdot) \in \mathcal{O}(D_2)$  and taking (7.27)–(7.29) into account, we deduce that

(7.30) 
$$\left| f(a_1, z_2^0) - f(a_1^0, z_2^0) \right| \le \epsilon^{N(1 - \omega(z_2^0, K, D_2))} (2|f|_X)^{\omega(z_2^0, K, D_2)} < \frac{\epsilon}{2}.$$

Since  $f(a_1, \cdot)|_{D_2}$  is a bounded holomorphic function for  $a_1 \in A_1$ , there is an open neighborhood V of  $z_2^0$  such that

$$\left|f(a_1, z_2) - f(a_1, z_2^0)\right| < \frac{\epsilon}{2}, \qquad a_1 \in A_1, \ z_2 \in V.$$

This, combined with (7.30), implies that

$$\begin{aligned} \left| f(a_1, z_2) - f(a_1^0, z_2^0) \right| &\leq \left| f(a_1, z_2^0) - f(a_1^0, z_2^0) \right| + \left| f(a_1, z_2) - f(a_1, z_2^0) \right| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon, \qquad a_1 \in A_{a_1^0}, \ z_2 \in V. \end{aligned}$$

Therefore, f is continuous at  $(a_1^0, z_2^0)$ . Consequently, we conclude, by Step 5, that  $\hat{f}$  admits the angular limit  $f(a_1^0, z_2^0)$  at  $(a_1^0, z_2^0)$ . Similarly, we may also show that  $\hat{f}$  admits the angular limit  $f(z_1^0, a_2^0)$  at every point  $(z_1^0, a_2^0) \in D_1 \times A_2^*$ . This completes the proof of the last step.

# 8. Preparatory results

We first develop some auxiliary results. This preparation will enable us to generalize the results of section 7 to the general case considered in Theorem A.

**Definition 8.1.** Let  $\Omega \subset \mathbb{C}^n$  be an open set and let  $A \subset \Omega$ . Define

$$h_{A,\Omega} := \sup \left\{ u : u \in \mathcal{PSH}(\Omega), u \leq 1 \text{ on } \Omega, u \leq 0 \text{ on } A \right\},$$
  
$$h_{A,\Omega}^*(z) := \limsup_{w \to z} h_{A,\Omega}(w), \qquad z \in \Omega.$$

The function  $h_{A,\Omega}^*$  is called the plurisubharmonic measure of A relative to  $\Omega$ .

**Proposition 8.2.** Let  $D \subset \mathbb{C}$  be an open set which is locally rectifiable on a linearly measurable subset A of  $\partial D$  with  $\operatorname{mes}(A) > 0$ . Let  $\{a_j\}_{j \in J}$  be a finite or countable subset of A with the following properties:

(i) For any  $j \in J$ , there is an open neighborhood  $U_j$  of  $a_j$  such that  $D \cap U_j$  is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains;

(ii) 
$$A \subset \bigcup_{j \in J} U_j$$
.

For any  $0 < \delta < \frac{1}{2}$ , define

$$U_{j,\delta} := \{ z \in D \cap U_j : \ \omega(z, A \cap U_j, D \cap U_j) < \delta \}, \qquad j \in J,$$
$$A_{\delta} := \bigcup_{j \in J} U_{j,\delta},$$
$$D_{\delta} := \{ z \in D : \ \omega(z, A, D) < 1 - \delta \}.$$

Then:

1)  $A \cap A^* \subset A^D_{\delta}$  and  $A_{\delta} \subset D_{1-\delta} \subset D_{\delta};$ 2)  $\omega(z, A, D) - \delta \leq h^*_{A_{\delta}, D}(z) \leq \omega(z, A, D), \ z \in D.$ 

*Proof.* To prove Part 1), let  $a \in A \cap A^*$  and fix an  $j \in J$  such that  $a \in U_j$ . Then

$$\lim_{z \to a, \ z \in \mathcal{A}_{\alpha}(a)} \omega(z, A \cap U_j, D \cap U_j) = 0, \qquad 0 < \alpha < \frac{\pi}{2}.$$

Consequently, for every  $0 < \alpha < \frac{\pi}{2}$ , there is an open neighborhood  $V_{\alpha} \subset U_j$  of a such that

$$\omega(z, A \cap U_j, D \cap U_j) < \delta, \qquad z \in \mathcal{A}_{\alpha}(a) \cap V_{\alpha}.$$

This proves  $A \cap A^* \subset A^D_{\delta}$ .

To prove the second assertion of Part 1), one invokes the Subordination Principle and obtains for  $z \in U_{i,\delta}$ ,

(8.1) 
$$\omega(z, A, D) \le \omega(z, A \cap U_j, D \cap U_j) < \delta < 1 - \delta.$$

Hence,  $z \in D_{1-\delta}$ . This implies that  $A_{\delta} \subset D_{1-\delta}$ . In addition, since  $0 < \delta < \frac{1}{2}$ , it follows that  $D_{1-\delta} \subset D_{\delta}$ . Hence, Part 1) is proved.

We turn to Part 2). Since  $A_{\delta}$  is an open set and, by Part 1),  $A \cap A^* \subset A_{\delta}^D$ , it follows from Definitions 5.17, 8.1 that

$$h^*_{A_{s,D}}(z) \le \omega(z, A \cap A^*, D), \qquad z \in D.$$

Hence, in virtue of Proposition 5.9 and Theorem 5.14, it follows that

$$h_{A_{\delta},D}^*(z) \le \omega(z,A,D), \qquad z \in D,$$

which proves the second estimate of Part 2).

To complete Part 2), let  $z \in A_{\delta}$ . Choose  $j \in J$  such that  $z \in U_{j,\delta}$ . We deduce from (8.1) that  $\omega(z, A, D) - \delta \leq 0$ . Hence,

$$\omega(z, A, D) - \delta \le 0, \qquad z \in A_{\delta}.$$

On the other hand,  $\omega(z, A, D) - \delta < 1$ ,  $z \in D$ . Consequently, the first estimate of Part 2) follows. The proof of the lemma is finished.

The next result gives a nice geometric property of the interior of a wedge.

**Proposition 8.3.** Let  $N \in \mathbb{N}$ ,  $N \geq 2$ ,  $D_j \subset \mathbb{C}$  a domain and  $A_j \subset \partial D_j$  such that  $D_j$  is locally rectifiable on  $A_j$ ,  $\operatorname{mes}(A_j) > 0$ ,  $j = 1, \ldots, N$ . Put  $\widehat{X}^o := \widehat{X}^o(A_1, \ldots, A_N; D_1, \ldots, D_N)$ . Then  $\widehat{X}^o$  is a domain.

*Proof.* We only give the proof for the case N = 2. The general case when N > 2 can be proved in exactly the same way. Fix arbitrary points  $z^0 = (z_1^0, z_2^0)$  and  $w^0 = (w_1^0, w_2^0)$  of  $\widehat{X}^{\circ}$ . Let

(8.2)  

$$\epsilon := \frac{1}{2} \Big( \min \left\{ 1 - \omega(z_1^0, A_1, D_1) - \omega(z_2^0, A_2, D_2), 1 - \omega(w_1^0, A_1, D_1) - \omega(w_2^0, A_2, D_2) \right\} \Big).$$

For  $j, k \in \{1, 2\}, j \neq k$ , let  $D_{j, z_j^0}$  (resp.  $D_{j, w_j^0}$ ) be the connected component containing  $z_j^0$  (resp.  $w_j^0$ ) of the following open set

(8.3) 
$$\left\{ z_j \in D_j : \ \omega(z_j, A_j, D_j) < 1 - \omega(z_k^0, A_k, D_k) - \epsilon \right\}, \\ \left( \operatorname{resp} \left\{ z_j \in D_j : \ \omega(z_j, A_j, D_j) < 1 - \omega(w_k^0, A_k, D_k) - \epsilon \right\} \right),$$

where  $\epsilon$  is given in (8.2).

By Theorem 5.18 (iii), for  $j \in \{1, 2\}$ , we may find a point  $a_j \in A_j \cap A_j^* \cap D_{j, z_j^0}^{D_j}$ (resp.  $b_j \in A_j \cap A_j^* \cap D_{j, w_i^0}^{D_j}$ ).

For  $j \in \{1, 2\}$ , fix a connected open neighborhood  $V_{z_j^0} \subset D_j$  of  $z_j^0$  and an open neighborhood  $V_{a_j} \subset D_j$  of  $a_j$  such that

(8.4)  

$$V_{a_j} \cap \mathcal{A}_{\frac{\pi}{4}}(a_j) \quad \text{is a domain,}$$

$$\omega(z_j, A_j, D_j) < \omega(z_j^0, A_k, D_j) + \epsilon, \quad z_j \in V_{z_j^0},$$

$$\omega(z_j, A_j, D_j) < \epsilon, \quad z_j \in V_{a_j} \cap \mathcal{A}_{\frac{\pi}{4}}(a_j).$$

Combining (8.2)–(8.4), we see that  $V_{z_1^0} \times V_{z_2^0}$  and  $V_{z_1^0} \times \left(V_{a_2} \cap \mathcal{A}_{\frac{\pi}{4}}(a_2)\right)$  are in the same connected component of  $\widehat{X}^{\circ}$ . The same argument also shows that  $V_{z_1^0} \times \left(V_{a_2} \cap \mathcal{A}_{\frac{\pi}{4}}(a_2)\right)$  and  $\left(V_{a_1} \cap \mathcal{A}_{\frac{\pi}{4}}(a_1)\right) \times \left(V_{a_2} \cap \mathcal{A}_{\frac{\pi}{4}}(a_2)\right)$  are in the same connected component of  $\widehat{X}^{\circ}$ . Hence,  $V_{z_1^0} \times V_{z_2^0}$  and  $\left(V_{a_1} \cap \mathcal{A}_{\frac{\pi}{4}}(a_1)\right) \times \left(V_{a_2} \cap \mathcal{A}_{\frac{\pi}{4}}(a_2)\right)$  are in the same connected component of  $\widehat{X}^{\circ}$ .

We apply the same argument as above to  $w^0$ . Consequently, one may find, for  $j \in \{1, 2\}$ , a connected open neighborhood  $V_{w_j^0} \subset D_j$  of  $w_j^0$  and an open neighborhood  $V_{b_j} \subset D_j$  of  $b_j$  such that

(8.5) 
$$V_{b_j} \cap \mathcal{A}_{\frac{\pi}{4}}(b_j) \quad \text{is a domain,} \\ \omega(z_j, A_j, D_j) < \omega(w_j^0, A_k, D_j) + \epsilon, \qquad z_j \in V_{w_j^0}, \\ \omega(z_j, A_j, D_j) < \epsilon, \qquad z_j \in V_{b_j} \cap \mathcal{A}_{\frac{\pi}{4}}(b_j), \end{cases}$$

and  $V_{w_1^0} \times V_{w_2^0}$  and  $\left(V_{b_1} \cap \mathcal{A}_{\frac{\pi}{4}}(b_1)\right) \times \left(V_{b_2} \cap \mathcal{A}_{\frac{\pi}{4}}(b_2)\right)$  are in the same connected component of  $\widehat{X}^{\circ}$ .

Recall that  $D_1$  and  $D_2$  are domains. Then in virtue of (8.4) and (8.5), one may find, for  $j \in \{1, 2\}$ , a Jordan curve  $\gamma_j : [0, 1] \longrightarrow D_j$  such that

(8.6) 
$$\gamma_j(0) \in V_{a_j} \cap \mathcal{A}_{\frac{\pi}{4}}(a_j) \text{ and } \gamma_j(1) \in V_{b_j} \cap \mathcal{A}_{\frac{\pi}{4}}(b_j).$$

Let

(8.7) 
$$\delta := \frac{1}{2} \left( \min \left\{ 1 - \sup_{t \in [0,1]} \omega(\gamma_1(t), A_1, D_1), 1 - \sup_{t \in [0,1]} \omega(\gamma_2(t), A_2, D_2) \right\} \right).$$

Next, we shrink  $V_{b_2}$  such that

$$\omega(z_2, A_2, D_2) < \delta, \qquad z_2 \in V_{b_2}$$

This, combined with (8.4)–(8.7), implies that  $(V_{a_1} \cap \mathcal{A}_{\frac{\pi}{4}}(a_1)) \times (V_{b_2} \cap \mathcal{A}_{\frac{\pi}{4}}(b_2))$ and  $(V_{b_1} \cap \mathcal{A}_{\frac{\pi}{4}}(b_1)) \times (V_{b_2} \cap \mathcal{A}_{\frac{\pi}{4}}(b_2))$  are in the same connected component of  $\widehat{X}^{\circ}$ . Similarly, by shrinking  $V_{a_1}$  if necessary, we see that  $(V_{a_1} \cap \mathcal{A}_{\frac{\pi}{4}}(a_1)) \times (V_{a_2} \cap \mathcal{A}_{\frac{\pi}{4}}(a_2))$ and  $(V_{a_1} \cap \mathcal{A}_{\frac{\pi}{4}}(a_1)) \times (V_{b_2} \cap \mathcal{A}_{\frac{\pi}{4}}(b_2))$  are in the same connected component of  $\widehat{X}^{\circ}$ . Hence,  $(V_{a_1} \cap \mathcal{A}_{\frac{\pi}{4}}(a_1)) \times (V_{a_2} \cap \mathcal{A}_{\frac{\pi}{4}}(a_2))$  and  $(V_{b_1} \cap \mathcal{A}_{\frac{\pi}{4}}(b_1)) \times (V_{b_2} \cap \mathcal{A}_{\frac{\pi}{4}}(b_2))$  are in the same connected component of  $\widehat{X}^{\circ}$ .

In summary, combining the above fact, (8.5) and the similar conclusion for  $z^0$ , we have shown that  $V_{z_1^0} \times V_{z_2^0}$  and  $V_{w_1^0} \times V_{w_2^0}$  are in the same connected component of  $\hat{X}^{\circ}$ . Consequently,  $z^0$  and  $w^0$  are in the same connected component of  $\hat{X}^{\circ}$ , which completes the proof.

The main ingredient in the proof of Theorem A is the following mixed cross theorem.

**Theorem 8.4.** Let  $D \subset \mathbb{C}^n$  be a domain of holomorphy,  $\Omega \subset \mathbb{C}$  an open set,  $A \subset D$ , and B a linearly measurable subset of  $\partial\Omega$ . Assume that  $A = \bigcup_{k=1}^{\infty} A_k$  with  $A_k$  locally pluriregular<sup>5</sup> compact subsets of D,  $A_k \subset A_{k+1}$ ,  $k \geq 1$ . In addition,  $\Omega$  is locally rectifiable on B with mes(B) > 0, and  $B \subset B^*$ . For  $0 \leq \delta < 1$  put G :=  $\{w \in \Omega : \omega(w, B, \Omega) < 1 - \delta\}$ . Let  $X := \mathbb{X}(A, B; D, G), X^o := \mathbb{X}^o(A, B; D, G)$ , and (using the notation of Theorem 5.19)

$$\widehat{X}^o = \widehat{\mathbb{X}}^o(A, B; D, G) := \left\{ (z, w) \in D \times G : h^*_{A, D}(z) + \omega_\delta(w, B, \Omega) < 1 \right\}.$$

Let  $f: X \longrightarrow \mathbb{C}$  be such that

(i)  $f \in \mathcal{O}_s(X^o);$ 

- (ii) f is measurable and bounded on X;
- (iii) for any  $z \in A$ ,

$$\lim_{w \to \eta, \ w \in \mathcal{A}_{\alpha}(\eta)} f(z, w) = f(z, \eta), \qquad \eta \in B, \ 0 < \alpha < \frac{\pi}{2}$$

<sup>&</sup>lt;sup>5</sup>See Sections 5.1-5.3 in the book by Klimek [9] for the definition of the notion *local* pluriregularity.

Then there is a unique function  $\hat{f} \in \mathcal{O}(\widehat{X}^o)$  such that  $\hat{f} = f$  on  $A \times G$  and

$$\lim_{z \to z_0, \ w \to \eta_0, \ w \in \mathcal{A}_{\alpha}(\eta_0)} \hat{f}(z, w) = f(z_0, \eta_0), \qquad 0 < \alpha < \frac{\pi}{2}$$

for every  $z_0 \in D$  and  $\eta_0 \in B$ . Moreover,  $|\hat{f}|_{\hat{X}^o} \leq |f|_X$ .

*Proof.* First one proves the existence and uniqueness of f. To do this we argue as in the proof of Theorem 3.5.1 in [7]. For the sake of completeness, we give here a sketchy proof. Fix an  $f: X \longrightarrow \mathbb{C}$  which satisfies (i)–(iii) above.

**Step I:** Reduction to the case where D is strongly pseudoconvex and A is a locally pluriregular compact subset of D.

One proceeds as in the first and second step in that proof. More precisely, since D is a domain of holomorphy, we may find an exhaustion sequence  $(D_k)_{k=1}^{\infty}$  of relatively compact, strongly pseudoconvex subdomains  $D_k$  of D with  $A_k \subset D_k \nearrow D$ .

By reduction assumption, for each k there exists an  $\hat{f}_k \in \mathcal{O}\left(\widehat{\mathbb{X}}^{\circ}(A_k, B; D_k, G)\right)$ such that  $\hat{f}_k$  admit the angular limit  $f|_{\mathbb{X}(A_k, B; D_k, G)}$  on  $\mathbb{X}(A_k, B; D_k, G)$ .

We would like to show that  $\hat{f}_{k+1} = \hat{f}_k$  on  $\widehat{\mathbb{X}}^{\circ}(A_k, B; D_k, G)$ . Indeed, fix an arbitrary  $k_0 \geq 1$  and an arbitrary point  $(z_0, w_0) \in \widehat{\mathbb{X}}^{\circ}(A_{k_0}, B; D_{k_0}, G)$ . Let  $k \in \mathbb{N}$  such that  $k \geq k_0$ . Let  $\mathcal{D}$  be the connected component containing  $z_0$  of the following open set

$$\left\{z \in D: h^*_{A_{k_0}, D_{k_0}}(z) < 1 - \omega_{\delta}(w_0, B, \Omega)\right\}.$$

Observe that both functions  $\hat{f}_{k_0}(\cdot, w_0)|_{\mathcal{D}}$  and  $\hat{f}_k(\cdot, w_0)|_{\mathcal{D}}$  are holomorphic and

$$\hat{f}_k(z, w_0) = f_k(z, w_0) = \hat{f}_{k_0}(z, w_0), \qquad z \in A_k \cap \mathcal{D}.$$

Since  $A_k \cap \mathcal{D}$  is non-pluripolar, we deduce that  $\hat{f}_{k_0}(\cdot, w_0)|_{\mathcal{D}} = \hat{f}_k(\cdot, w_0)|_{\mathcal{D}}$ . Hence,  $\hat{f}_{k_0}(z_0, w_0) = \hat{f}_k(z_0, w_0)$ , which proves the above assertion.

On the other hand, by Proposition 5.15 one gets  $\widehat{\mathbb{X}}^{\circ}(A_k, B; D_k, G) \nearrow \widehat{X}^{\circ}$  as  $k \nearrow \infty$ . Therefore, we may glue  $\widehat{f}_k$  together to obtain a function  $\widehat{f} \in \mathcal{O}(\widehat{X}^{\circ})$  such that  $\widehat{f}$  admits the angular limit f on X and  $\widehat{f} = f$  on  $A \times G$ . The uniqueness of such an extension  $\widehat{f}$  can be proved using the argument given in the previous paragraph.

This completes Step I.

**Step II:** The case where D is strongly pseudoconvex and A is a locally pluriregular compact subset of D.

Suppose without loss of generality that  $|f|_X < 1$ . The key observation is that we are still able to apply the classical method of doubly orthogonal bases of Bergman type.

Next one observes that Lemma 3.5.10 in [7] is still valid in the present context. Look at Step 3 in that proof. In the sequel, we will use the notation from [7].

Let  $\mu := \mu_{A,D}$ ,  $H_0 := L_h^2(D)$ ,  $H_1 :=$  the closure of  $H_0|_A$  in  $L^2(A,\mu)$  and let  $(b_k)_{k=1}^{\infty} \subset H_0$  be the basis from Lemma 3.5.10 in [7],  $\nu_k := \|b_k\|_{H_0}$ ,  $k \in \mathbb{N}$ , with the

following property:

(8.8) 
$$\sum_{k=1}^{\infty} \nu_k^{-\epsilon} < \infty, \qquad \epsilon > 0.$$

For any  $w \in B$ , we have  $f(\cdot, w) \in H_0$  and  $f(\cdot, w)|_A \in H_1$ . Hence

(8.9) 
$$f(\cdot, w) = \sum_{k=1}^{\infty} c_k(w) b_k,$$

where

(8.10) 
$$c_k(w) = \frac{1}{\nu_k^2} \int_D f(z, w) \overline{b_k(z)} d\Lambda_{2n}(z) = \int_A f(z, w) \overline{b_k(z)} d\mu(z), \qquad k \in \mathbb{N}.$$

Taking the hypotheses (i)–(iii) into account and applying Lebesgue's Dominated Convergence Theorem, we see that the formula

(8.11) 
$$\widehat{c}_k(w) := \int_A f(z, w) \overline{b_k(z)} d\mu(z), \qquad w \in G \cup B, \ k \in \mathbb{N};$$

defines a bounded function which is holomorphic in G. Moreover, by (iii) and (8.10)–(8.11) it follows that

(8.12) 
$$\lim_{w \to \eta, \ w \in \mathcal{A}_{\alpha}(\eta)} \widehat{c_k}(w) = \widehat{c_k}(\eta) = c_k(\eta), \qquad \eta \in B, \ 0 < \alpha < \frac{\pi}{2}$$

Observe, as in [7] and using (8.10)–(8.12), that we obtain the following estimates

$$\begin{aligned} \frac{\log |\widehat{c}_k(w)|}{\log \nu_k} &\leq \frac{\log \sqrt{\mu(A)}}{\log \nu_k}, \qquad w \in G, \ k \in \mathbb{N}, \\ \limsup_{w \to \eta, \ w \in \mathcal{A}_\alpha(\eta)} \frac{\log |\widehat{c}_k(\eta)|}{\log \nu_k} &\leq \frac{\log \sqrt{\Lambda_{2n}(D)}}{\log \nu_k} - 1, \qquad \eta \in B, \ 0 < \alpha < \frac{\pi}{2}, \ k \in \mathbb{N}, \end{aligned}$$

where  $\Lambda_{2n}(D)$  is the volume of D with respect to the Lebesgue measure of  $\mathbb{C}^n$ .

This shows that for any  $\epsilon > 0$ , there is a sufficiently large N such that for all  $k \ge N$ ,

(8.13) 
$$\frac{\log |\widehat{c}_k|}{\log \nu_k} \le \omega_{\delta}(\cdot, B, \Omega) + \epsilon - 1 \quad \text{on } G.$$

Take a compact set  $K \Subset D$  and let  $1 > \alpha > \max_{K} h_{A,D}^*$ . Choose an  $\epsilon > 0$  so small that  $\alpha + 2\epsilon < 1$ . Consider the open set

$$G_K := \{ w \in G : \omega_{\delta}(\cdot, B, \Omega) < 1 - \alpha - 2\epsilon \}.$$

By (8.13) there is a constant C'(K) such that

(8.14) 
$$|\widehat{c}_k|_{G_k} \le C'(K)\nu_k^{\omega_\delta(\cdot,B,\Omega)+\epsilon-1} \le C'(K)\nu_k^{-\alpha-\epsilon}, \qquad k \ge 1.$$

Now we wish to show that

(8.15) 
$$\sum_{k=1}^{\infty} \widehat{c}_k(w) b_k(z)$$

converges locally uniformly in  $\widehat{X}^{\text{o}}$ . Indeed, by (8.8), (8.14), and Lemma 3.5.10 in [7],

$$(8.16)$$

$$\sum_{k=1}^{\infty} |\widehat{c_k}|_{G_K} |b_k|_K \leq \sum_{k=1}^{\infty} C'(K) \nu_k^{-\alpha-\epsilon} C(K,\alpha) \nu_k^{\alpha} \leq C'(K) C(K,\alpha) \sum_{k=1}^{\infty} \nu_k^{-\epsilon} < \infty,$$

which gives the normal convergence on  $K \times G_K$ . Since the compact set K and  $\epsilon > 0$  are arbitrary, the series in (8.15) converges uniformly on compact subsets of  $\widehat{X}^{\circ}$ . Let  $\widehat{f}$  denote this limit function in (8.15).

Fix  $z_0 \in D$  and  $\eta_0 \in B$ . We choose a compact K of D as above. Moreover, we may suppose without loss of generality that K contains a neighborhood of  $z_0$ . Let  $\epsilon_0 > 0$ .

In virtue of (8.16), there is an  $N_0$  such that

(8.17) 
$$\sum_{k=N_0+1}^{\infty} |\widehat{c}_k|_{G_K} |b_k|_K < \frac{\epsilon_0}{2}.$$

On the other hand, in virtue of (8.9)–(8.12), we may find, for any  $0 < \alpha < \frac{\pi}{2}$ , an open neighborhood  $V_{\alpha}$  of  $\eta_0$  such that

$$\left|\sum_{k=1}^{N_0} \widehat{c_k}(w) b_k(z) - \sum_{k=1}^{N_0} c_k(\eta_0) b_k(z)\right| < \frac{\epsilon_0}{2}, \qquad z \in K, \ w \in \mathcal{A}_\alpha(\eta_0) \cap V_\alpha.$$

This, combined with (8.15) and (8.17), implies that

$$\lim_{x \to z_0, w \to \eta_0, w \in \mathcal{A}_{\alpha}(\eta_0)} \left| \hat{f}(z, w) - f(z_0, \eta_0) \right| < \epsilon_0, \qquad 0 < \alpha < \frac{\pi}{2}.$$

Since  $\epsilon_0 > 0$  and  $(z_0, \eta_0) \in D \times B$  can be arbitrarily chosen, we conclude that

$$\lim_{z \to z_0, w \to \eta_0, w \in \mathcal{A}_{\alpha}(\eta_0)} \hat{f}(z, w) = f(z_0, \eta_0), \qquad (z_0, \eta_0) \in D \times B, \ 0 < \alpha < \frac{\pi}{2}.$$

To complete Step II, it remains to show that  $\hat{f} = f$  on  $A \times G$ . To do this, fix an arbitrary  $(z_0, w_0) \in A \times G$ . Let  $\mathcal{G}$  be the connected component of G containing  $w_0$ . Recall that  $G = \{w \in \Omega : \ \omega(w, B, \Omega) < 1 - \delta\}$ . Then observe that both functions  $\hat{f}(z_0, \cdot)|_{\mathcal{G}}$  and  $f(z_0, \cdot)|_{\mathcal{G}}$  admit the same angular limit f on  $B \cap \mathcal{G}^{\Omega}$ . Consequently, applying Theorem 6.4 yields that  $\hat{f}(z_0, \cdot)|_{\mathcal{G}} = f(z_0, \cdot)|_{\mathcal{G}}$ . Hence,  $\hat{f}(z_0, w_0) = f(z_0, w_0)$ , which proves the above assertion.

This completes the proof of Step II.

It remains to prove the estimate  $|\hat{f}|_{\hat{X}} \leq |f|_X$ . In order to reach a contradiction assume that there is a point  $z^0 \in \hat{X}^\circ$  such that  $|\hat{f}(z^0)| > |f|_X$ . Put  $\alpha := \hat{f}(z^0)$  and consider the function

(8.18) 
$$g(z) := \frac{1}{f(z) - \alpha}, \qquad z \in X.$$

Using the above assumption, it can be checked that g satisfies hypotheses (i)–(iii) of Theorem 8.4. Hence applying the first assertion of the theorem, there is exactly one

function  $\hat{g} \in \mathcal{O}(\hat{X}^{\circ})$  with  $\hat{g} = g$  on  $A \times G$ . Therefore, by (8.18) we have on  $A \times G$ :  $g(f - \alpha) \equiv 1$ . Thus  $\hat{g}(\hat{f} - \alpha) \equiv 1$  on  $\hat{X}^{o}$ . In particular,

$$0 = \hat{g}(z^0)(\hat{f}(z^0) - \alpha) = 1,$$

a contradiction. Hence the inequality  $|\hat{f}|_{\hat{X}} \leq |f|_X$  is proved.

Finally, we conclude this section with two uniqueness results.

**Proposition 8.5.** Let  $D_i \subset \mathbb{C}$  be an open set and  $A_i$  a linearly measurable subset of  $\partial D_j$  such that  $D_j$  is locally rectifiable on  $A_j$ ,  $A_j \subset A_j^*$ ,  $mes(A_j) > 0$ , j = 1, 2. Let  $\tilde{D}_1 \subset \mathbb{C}$  be a domain,  $D_1 \cap \tilde{D}_1 \neq \emptyset$ , and let  $\tilde{A}_1$  be a linearly measurable subset of  $\partial \widetilde{D}_1$  such that  $\widetilde{D}_1$  is locally rectifiable on  $\widetilde{A}_1$ , and  $\operatorname{mes}(\widetilde{A}_1) > 0$ . Put

$$\widehat{X}^{o} := \widehat{\mathbb{X}}^{o} \Big( A_{1}, A_{2}; D_{1}, D_{2} \Big),$$
$$\widehat{\widetilde{X}}^{o} := \widehat{\mathbb{X}}^{o} \Big( \widetilde{A}_{1}, A_{2}; \widetilde{D}_{1}, D_{2}, \Big).$$

Let  $\hat{f} \in \mathcal{O}(\hat{X}^o)$ ,  $\hat{\tilde{f}} \in \mathcal{O}(\hat{\tilde{X}}^o)$ , and  $z_1^0 \in D_1 \cap \tilde{D}_1$  be such that both  $\hat{f}$  and  $\hat{\tilde{f}}$  admit the same angular limit at  $(z_1^0, a_2)$  for a.e.  $a_2 \in A_2$ . Then  $\hat{f}(z) = \hat{f}(z)$  for every  $z = (z_1^0, z_2) \in \widehat{X}^o \cap \widehat{\widetilde{X}}^o.$ 

*Proof.* Fix an arbitrary  $z_2^0 \in D_2$  such that  $z^0 := (z_1^0, z_2^0) \in \widehat{X}^\circ \cap$  $\widehat{\widetilde{X}}^{\circ}$ . Let  $\epsilon_0 := \frac{1}{4} \Big( 1 - \max \Big\{ \omega(z_1^0, A_1, D_1), \omega(z_1^0, \widetilde{A}_1, \widetilde{D}_1) \Big\} \Big)$ , and put  $D_{2,\epsilon_0} :=$  $\{z_2 \in D_2: \ \omega(z_2, A_2, D_2) < 1 - \epsilon_0\}$ . Applying Theorem 6.4 to  $\hat{f}|_{D_{2,\epsilon_0}}$  and  $\hat{\tilde{f}}|_{D_{2,\epsilon_0}}$ , it follows that  $\widehat{\widetilde{f}}(z_1^0, z_2^0) = \widehat{f}(z_1^0, z_2^0)$ . Hence, the proof is finished. 

Now we are able to prove the uniqueness stated in Theorem A.

**Corollary 8.6.** We keep the hypotheses and the notation of Theorem A. Then there is at most one function  $\hat{f} \in \mathcal{O}(\hat{X}^o)$  which satisfies Property 1) of Theorem A.

*Proof.* It follows immediately from Proposition 8.5.

#### 9. Proof of Theorem A

We mainly consider the case N = 2. In this case we proceed by four steps. Recall that by Corollary 8.6, the function  $\hat{f}$  satisfying Part 1) is uniquely determined (if exists).

In the first two steps we mainly prove the following assertion:

There are a function  $\hat{f} \in \mathcal{O}(\widehat{X}^o)$  and a subset  $\tilde{A}_j$  of  $A_j \cap A_j^*$  (j = 1, 2) such that  $\operatorname{mes}(A_j \setminus \tilde{A}_j) = 0$  and  $\hat{f}$  admits the angular limit f at every point of  $(\tilde{A}_1 \times D_2) \cup \tilde{A}_j$ (\*)  $(D_1 \times A_2).$ 

In fact, in the first two steps we always assume that  $|f|_X < \infty$ . Using this and the previous observation and taking  $\binom{*}{**}$  for granted, we conclude the proof of Steps

1 and 2 below in exactly the same way as we did in Section 7 starting from Step 2 of that section.

**Step 1:** Proof of Theorem A for the case where N = 2,  $|f|_X < \infty$ , and  $D_2$  is a rectifiable Jordan domain.

*Proof of Step 1.* In virtue of Proposition 8.2, let  $\{a_{1j}\}_{j\in J_1}$  be a finite or countable subset of  $A_1$  with the following properties:

• For any  $j \in J_1$ , there is an open neighborhood  $U_{1j}$  of  $a_{1j}$  such that  $D_1 \cap U_{1j}$  is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains (according to the type of  $a_{1j}$ );

• 
$$A_1 \subset \bigcup_{j \in J_1} U_{1j}$$
.

For any  $0 < \delta < \frac{1}{2}$ , define

$$U_{1j,\delta} := \{ z_1 \in D_1 \cap U_{1j} : \ \omega(z_1, A_1 \cap U_{1j}, D_1 \cap U_{1j}) < \delta \}, \qquad j \in J_1,$$
$$A_{1,\delta} := \bigcup_{j \in J_1} U_{1j,\delta},$$
$$D_{2,\delta} := \{ z_2 \in D_2 : \ \omega(z_2, A_2, D_2) < 1 - \delta \}.$$

Moreover, for every  $j \in J_1$  let

$$X_j := \mathbb{X} \left( \partial (D_1 \cap U_{1j}) \cap A_1, A_2; D_1 \cap U_{1j}, D_2) \right),$$

(9.1) 
$$\widehat{X_j}^{\circ} := \widehat{\mathbb{X}}^{\circ} \left( \partial (D_1 \cap U_{1j}) \cap A_1, A_2; D_1 \cap U_{1j}, D_2) \right),$$
$$\widetilde{f_j} := f|_{X_j}.$$

Using the hypotheses on f, we conclude that  $f_j$ ,  $j \in J_1$ , satisfies (i)–(iii) of Theorem A. Moreover,  $|\tilde{f}_j|_{X_j} \leq |f|_X < \infty$ . Since  $D_2$  is a rectifiable Jordan domain and  $D_1 \cap U_{1j}$ ,  $j \in J_1$ , is either a rectifiable Jordan domain or the disjoint union of two rectifiable Jordan domains, we are able to apply the result of Section 7 to  $\tilde{f}_j$ . Consequently, we obtain, for  $j \in J_1$ , a unique function  $\hat{f}_{1j} \in \mathcal{O}\left(\widehat{X_j}^\circ\right)$ , a subset  $A_{1j}$  of  $A_1$ , a subset  $A_{2j}$  of  $A_2$  such that

 $\operatorname{mes}(A_1 \setminus A_{1j}) = \operatorname{mes}(A_2 \setminus A_{2j}) = 0,$ 

 $\hat{f}_{1j}$  admits the angular limit f on  $((\partial (D_1 \cap U_{1j}) \cap A_{1j}) \times D_2) \cup (D_1 \times A_{2j})$ . Put

(9.3)  

$$\tilde{A}_{1} := \bigcap_{j \in J_{1}} A_{1j} \quad \text{and} \quad \tilde{A}_{2} := \bigcap_{j \in J_{2}} A_{2j},$$

$$X_{\delta} := \mathbb{X} \left( A_{1,\delta}, \tilde{A}_{2}; D_{1}, D_{2,\delta} \right),$$

$$\widehat{X}_{\delta}^{o} := \widehat{\mathbb{X}}^{o} \left( A_{1,\delta}, \tilde{A}_{2}; D_{1}, D_{2,\delta} \right).$$

In virtue of Proposition 8.5, we are able to collect the family  $\left(\hat{f}_{1j}|_{U_{1j,\delta}\times D_{2,\delta}}\right)_{j\in J_1}$  in order to obtain a function  $\tilde{\tilde{f}}_{\delta} \in \mathcal{O}(A_{1,\delta}\times D_{2,\delta})$ .

Next, consider the function  $\tilde{f}_{\delta}$ :  $X_{\delta} \longrightarrow \mathbb{C}$  given by

(9.4) 
$$\tilde{f}_{\delta} := \begin{cases} \tilde{f}_{\delta}, & \text{on } A_{1,\delta} \times D_{2,\delta} \\ f, & \text{on } D_1 \times \tilde{A}_2 \end{cases}$$

In virtue of (9.1)-(9.4), we deduce that

(9.5) 
$$\operatorname{mes}(A_1 \setminus A_1) = \operatorname{mes}(A_2 \setminus A_2) = 0,$$

and

(9.6) 
$$\lim_{\substack{z_1 \to z_1^0, \ z_2 \to a_2^0, \ z_2 \in \mathcal{A}_{\alpha}(a_2^0)}} \tilde{f}_{\delta}(z) = f(z_1^0, a_2^0), \qquad 0 < \alpha < \frac{\pi}{2}, \ z_1^0 \in D_1, \ a_2^0 \in \tilde{A}_2, \lim_{\substack{z_1 \to a_1^0, \ z_1 \in \mathcal{A}_{\alpha}(a_1^0), \ z_2 \to a_2^0}} \tilde{f}_{\delta}(z) = f(a_1^0, z_2^0), \qquad 0 < \alpha < \frac{\pi}{2}, \ a_1^0 \in \tilde{A}_1, \ z_2^0 \in D_{2,\delta}.$$

In virtue of (9.4)–(9.6),  $\tilde{f}_{\delta}$  satisfies the hypotheses (i)–(iii) of Theorem 8.4. Applying this theorem to  $\tilde{f}_{\delta}$ , we obtain, for every  $0 < \delta < \frac{1}{2}$ , a function  $\hat{f}_{\delta} \in \mathcal{O}\left(\widehat{X}_{\delta}^{\circ}\right)$ . In virtue of (9.6), we see that

$$\hat{f}_{\delta} = \tilde{f}_{\delta} \quad \text{on } A_{1,\delta} \times D_{2,\delta},$$

$$(9.7) \quad \lim_{z_1 \to z_1^0, \ z_2 \to a_2^0, \ z_2 \in \mathcal{A}_{\alpha}(a_2^0)} \hat{f}_{\delta}(z) = f(z_1^0, a_2^0), \qquad 0 < \alpha < \frac{\pi}{2}, \ z_1^0 \in D_1, \ a_2^0 \in \tilde{A}_2,$$

$$\lim_{z_1 \to a_1^0, \ z_1 \in \mathcal{A}_{\alpha}(a_1^0), \ z_2 \to a_2^0} \hat{f}_{\delta}(z) = f(a_1^0, z_2^0), \qquad 0 < \alpha < \frac{\pi}{2}, \ a_1^0 \in \tilde{A}_1, \ z_2^0 \in D_{2,\delta}$$

We are now in a position to define the desired extension function  $\hat{f}$ . Indeed, one glues  $(\hat{f}_{\delta})_{0<\delta<\frac{1}{2}}$  together to obtain  $\hat{f}$  in the following way

(9.8) 
$$\hat{f} := \lim_{\delta \to 0} \hat{f}_{\delta} \quad \text{on } \widehat{X}^{\circ} = \widehat{\mathbb{X}}^{\circ} \left( A_1, A_2; D_1, D_2 \right)$$

Now one has to check that the limit (9.8) exists and possesses all the required properties. This will be an immediate consequence of the following

Lemma 9.1. For any point 
$$z \in \widehat{X}^o$$
 put  
(9.9)  $\delta_z := \frac{1 - \omega(z_1, A_1, D_1) - \omega(z_2, A_2, D_2)}{2}.$ 

Then  $\hat{f}(z) = \hat{f}_{\delta}(z)$  for all  $0 < \delta \leq \delta_z$ .

Proof of Lemma 9.1. Fix an arbitrary point  $z^0 = (z_1^0, z_2^0) \in \widehat{\mathbb{X}}^{\circ}(A_1, A_2; D_1, D_2)$  and let  $\delta_0 := \delta_{z^0}$ . Let  $0 < \delta \leq \delta_0$ . Then, by Proposition 8.2, we see that  $\omega(z_2^0, \tilde{A}_2, D_2) < 1 - \delta_0$  and

$$\begin{aligned} h^*_{A_{1,\delta},D_1}(z_1^0) + \omega_{\delta_0}(z_2^0,\tilde{A}_2,D_2) &\leq \omega(z_1^0,A_1,D_1) + \frac{\omega(z_2^0,\tilde{A}_2,D_2)}{1-\delta_0} \\ &\leq \frac{\omega(z_1^0,A_1,D_1) + \omega(z_2^0,A_2,D_2)}{1-\delta_0} < 1, \end{aligned}$$

where the latter estimate follows from formula (9.9). Consequently,

(9.10) 
$$z^{0} \in \widehat{\mathbb{X}}^{o}\left(A_{1,\delta}, \widetilde{A}_{2}; D_{1}, D_{2,\delta_{0}}\right)$$

On the other hand, using Part 1) of Proposition 8.2, it is clear that (9.11)

$$\widehat{\mathbb{X}}^{\mathrm{o}}\left(A_{1,\delta}, \tilde{A}_{2}; D_{1}, D_{2,\delta_{0}}\right) \subset \widehat{\mathbb{X}}^{\mathrm{o}}\left(A_{1,\delta}, \tilde{A}_{2}; D_{1}, D_{2,\delta}\right) \cap \widehat{\mathbb{X}}^{\mathrm{o}}\left(A_{1,\delta_{0}}, \tilde{A}_{2}; D_{1}, D_{2,\delta_{0}}\right).$$

Moreover, in virtue of (9.4) and (9.7), we have

(9.12) 
$$\hat{f}_{\delta} = \tilde{f}_{\delta} = \hat{f}_{\delta_0} \quad \text{on } A_{1,\delta} \times D_{2,\delta_0}.$$

Next, let  $\mathcal{G}$  be the connected component containing  $z_1^0$  of the following open set

$$\left\{z_1 \in D_1: h^*_{A_{1,\delta},D_1}(z_1) < 1 - \omega_{\delta_0}(z_2^0, A_2, D_2)\right\}$$

Observe that, in virtue of (9.10)–(9.11), both functions  $\hat{f}_{\delta}|_{\mathcal{G}}$  and  $\hat{f}_{\delta_0}|_{\mathcal{G}}$  are holomorphic and  $\mathcal{G} \cap A_{1,\delta}$  is a nonempty open set. Therefore, we deduce from (9.12) that  $\hat{f}_{\delta} = \hat{f}_{\delta_0}$  on  $\mathcal{G}$ . Hence,  $\hat{f}_{\delta}(z^0) = \hat{f}_{\delta_0}(z^0)$ , which completes the proof of the lemma.  $\Box$ 

We complete the proof as follows. An immediate consequence of Lemma 9.1 is that  $\hat{f} \in \mathcal{O}(\widehat{X}^{\circ})$ . Next, we apply Lemma 9.1 and make use of (9.4)–(9.9) and of the fact that  $\widehat{X}_{\delta}^{\circ} \to \widehat{X}^{\circ}$  as  $\delta \searrow 0$ . Consequently, we conclude that  $\hat{f}$  satisfies  $\binom{*}{**}$ .

Taking into account the remark made at the beginning of this section, the proof of the theorem in Step 1 is finished.  $\Box$ 

**Step 2:** Proof of Theorem A for the case N = 2 and  $|f|_X < \infty$ .

*Proof of Step 2.* We proceed using Step 1 in exactly the same way as we proved Step 1 using the result of Section 7. Hence, Step 2 is finished.  $\Box$ 

**Step 3:** Proof of Theorem A for the case when N = 2 and  $A_1$ ,  $A_2$  are compacts.

Proof of Step 3. Recall from the hypothesis (i) of Theorem A that f is locally bounded. Since  $A_1$ ,  $A_2$  are compact, a compactness argument shows that there are a positive number  $M_0$  and an open neighborhood  $U_1$  (resp.  $U_2$ ) of  $A_1$  (resp.  $A_2$ ) such that

(9.13) 
$$|f| < M_0$$
 on  $([U_1 \cap (D_1 \cup A_1)] \times A_2) \bigcup (A_1 \times [U_2 \cap (D_2 \cup A_2)]).$ 

For  $j \in \{1, 2\}$ , fix a sequence  $\left(\tilde{D}_{j,k}\right)_{k=1}^{\infty}$  of subdomains of  $D_j$  such that

(9.14) 
$$\tilde{D}_{j,k} \in D_j \text{ and } \tilde{D}_{j,k} \nearrow D_j \text{ as } k \nearrow \infty.$$

Consider the sequence  $(D_{j,k})_{k=1}^{\infty}$  of open sets of  $D_j$ , the sequence of crosses  $(X_k)_{k=1}^{\infty}$ , and the sequence of the interior of their wedges  $(\widehat{X}_k^{\text{o}})_{k=1}^{\infty}$  defined as

(9.15)  
$$D_{j,k} := \tilde{D}_{j,k} \cup (U_j \cap D_j), \qquad k \ge 1$$
$$X_k := \mathbb{X} (A_1, A_2; D_{1,k}, D_{2,k}), \\\widehat{X}_k^{o} := \widehat{\mathbb{X}}^{o} (A_1, A_2; D_{1,k}, D_{2,k}).$$

Since f is locally bounded,  $A_1, A_2$  are compact and, by (9.14),  $\tilde{D}_{1,k} \subseteq D_1, \tilde{D}_{2,k} \subseteq D_2$ , a compactness argument shows that for every  $k \ge 1$ , there is a number  $M_k$  such that  $M_k \ge M_0$  and

$$|f| < M_k$$
 on  $\left(\tilde{D}_{1,k} \times A_2\right) \cup \left(A_1 \times \tilde{D}_{2,k}\right)$ 

This, combined with (9.13) and (9.15), implies that

(9.16) 
$$|f|_{X_k} < M_k, \quad k \ge 1.$$

On the other hand, in virtue of (9.14) and (9.15), we see that the sequences  $(D_{j,k})_{k=1}^{\infty}$ and  $(A_{j,k} := A_j)_{k=1}^{\infty}$  satisfy the hypotheses of Proposition 5.15. Consequently, applying this proposition and taking (9.15) into account, we obtain

(9.17) 
$$\widehat{X}_k^{\mathrm{o}} \nearrow \widehat{X}^{\mathrm{o}} = \widehat{\mathbb{X}}^{\mathrm{o}} (A_1, A_2; D_1, D_2) \text{ as } k \nearrow \infty.$$

In virtue of (9.16) and the hypothesis on f, we see that  $f|_{X_k}$  satisfies the hypotheses of Step 2. Consequently, applying the result of Step 2 to  $f|_{X_k}$ ,  $k \ge 1$ , we obtain a unique function  $\hat{f}_k \in \mathcal{O}(\hat{X}_k^{\circ})$  and a subset  $\tilde{A}_{1,k}$  (resp.  $\tilde{A}_{2,k}$ ) of  $A_1$  (resp.  $A_2$ ) such that

(9.18) 
$$\begin{aligned} \max(A_1 \setminus A_{1,k}) &= \max(A_2 \setminus A_{2,k}) = 0, \\ \hat{f}_k \text{ admits the angular limit } f \text{ on } \left(D_{1,k} \times \tilde{A}_{2,k}\right) \cup \left(\tilde{A}_{1,k} \times D_{2,k}\right). \end{aligned}$$

Put

(9.19) 
$$\tilde{A}_1 := \bigcap_{k=1}^{\infty} \tilde{A}_{1,k} \quad \text{and} \quad \tilde{A}_2 := \bigcap_{k=1}^{\infty} \tilde{A}_{2,k}.$$

Using (9.18), we deduce from (9.19) that

(9.20) 
$$\begin{aligned} \max(A_1 \setminus A_1) &= \max(A_2 \setminus A_2) = 0, \\ \hat{f}_k & \text{admits the angular limit } f \text{ on } \left(D_{1,k} \times \tilde{A}_2\right) \cup \left(\tilde{A}_1 \times D_{2,k}\right). \end{aligned}$$

Using (9.17), (9.20) and applying Corollary 8.6, we obtain

(9.21) 
$$\hat{f}_k = \hat{f}_{k+1}$$
 on  $\hat{X}_k^{\text{o}}, \ k \ge 1.$ 

Therefore, we may glue  $\hat{f}_k$  together in order to obtain the desired extension function  $\hat{f}$  as

(9.22) 
$$\hat{f} = \lim_{k \to \infty} \hat{f}_k \quad \text{on } \hat{X}^{\circ}.$$

In virtue of this formula and (9.17), (9.20)–(9.21), we conclude that  $\hat{f}$  satisfies  $\binom{*}{**}$ . Hence, the proof of Part 1) of Theorem A is complete.

Using  $\binom{*}{**}$  and arguing as in Step 3 of Section 7, Part 2) of Theorem A follows.

Now we turn to Part 3) of Theorem A. Let  $(a_1^0, z_2^0) \in A_1^* \times D_2$  be such that the following limit exists

$$\lambda := \lim_{(a_1, z_2) \to (a_1^0, z_2^0), \ (a_1, z_2) \in A_1 \times D_2} f(a_1, z_2).$$

We would like to show that  $\hat{f}$  admits the angular limit  $\lambda$  at  $(a_1^0, z_2^0)$ .

Let  $k_0$  be a sufficiently large integer such that  $z_2^0 \in D_{2,k_0}$ . Then in virtue of (9.17) and (9.20)–(9.21),

(9.23) 
$$\hat{f} = \hat{f}_{k_0} \qquad \text{on } \hat{X}_{k_0}^{\text{o}}$$

In addition, applying Part 2) of Theorem A established in Step 2 and taking (9.16) into account, we see that

(9.24) 
$$|\hat{f}_{k_0}|_{\hat{X}_{k_0}^{o}} \le |f|_{X_{k_0}} < M_{k_0} < \infty.$$

Consequently, the proof of Step 4 in Section 7 applied to  $\hat{f}_{k_0}$  still works in this context making the obviously necessary changes. Hence,  $\hat{f}_{k_0}$  and then  $\hat{f}$  by (9.23) admits the angular limit  $\lambda$  at  $(a_1^0, z_2^0)$ . This completes the proof of Part 3) of Theorem A.

Using  $\binom{*}{**}$  and (9.23)-(9.24) (for a sufficiently large  $k_0$ ), the proof of Parts 4) and 5) of Theorem A given in Steps 5 and 6 of Section 7 still works in this context making the obviously necessary changes. This completes Step 3.

**Step 4:** Proof of Theorem A for the case N = 2.

*Proof of Step 4.* For each  $j \in \{1, 2\}$ , since  $A_j \subset \partial D_j$  is linearly measurable, one may find a sequence  $(A_{j,k})_{k=1}^{\infty}$  of subsets of  $A_j$  such that

- (a)  $A_{j,k}$  is a compact subset of  $\partial D_j$  and  $\operatorname{mes}(A_{j,k}) > 0, k \ge 1$ ;
- (b)  $A_{j,k} \subset A_{j,k+1}, k \ge 1;$ (c)  $\operatorname{mes}\left(A_j \setminus \bigcup_{k=1}^{\infty} A_{j,k}\right) = 0.$

For  $k \geq 1$ , let

(9.25) 
$$X_k := \mathbb{X} (A_{1,k}, A_{2,k}; D_1, D_2), \\ \widehat{X}_k^{\circ} := \widehat{\mathbb{X}}^{\circ} (A_{1,k}, A_{2,k}; D_1, D_2).$$

On the other hand, in virtue of (a)–(c), we see that the sequences  $(D_{j,k} := D_j)_{k=1}^{\infty}$ and  $(A_{j,k})_{k=1}^{\infty}$  satisfy the hypotheses of Proposition 5.15. Consequently, applying this proposition and taking (9.25) into account, we obtain

(9.26) 
$$\widehat{X}_k^{\mathrm{o}} \nearrow \widehat{X}^{\mathrm{o}} = \widehat{\mathbb{X}}^{\mathrm{o}} (A_1, A_2; D_1, D_2) \text{ as } k \nearrow \infty.$$

In virtue of (a) and the hypothesis on f, we see that  $f|_{X_k}$  satisfies the hypotheses of Step 3. Consequently, applying the result of Step 3 to  $f|_{X_k}$ ,  $k \ge 1$ , we obtain a unique function  $\hat{f}_k \in \mathcal{O}(\hat{X}_k^{\circ})$  and a subset  $\tilde{A}_{1,k}$  (resp.  $\tilde{A}_{2,k}$ ) of  $A_1$  (resp.  $A_2$ ) such that

(9.27) 
$$\begin{aligned} \max(A_1 \setminus \tilde{A}_{1,k}) &= \max(A_2 \setminus \tilde{A}_{2,k}) = 0, \\ \hat{f}_k \quad \text{admits the angular limit } f \text{ on } \left(D_1 \times \tilde{A}_{2,k}\right) \cup \left(\tilde{A}_{1,k} \times D_2\right). \end{aligned}$$

Put

(9.28) 
$$\tilde{A}_1 := \bigcup_{k=1}^{\infty} \tilde{A}_{1,k}$$
 and  $\tilde{A}_2 := \bigcup_{k=1}^{\infty} \tilde{A}_{2,k}$ .

Using (9.27), we deduce from (9.28) that

(9.29) 
$$\begin{array}{l} \operatorname{mes}(A_1 \setminus \tilde{A}_1) = \operatorname{mes}(A_2 \setminus \tilde{A}_2) = 0, \\ \hat{f}_k \quad \text{admits the angular limit } f \text{ on } \left(D_{1,k} \times \tilde{A}_2\right) \cup \left(\tilde{A}_1 \times D_{2,k}\right). \end{array}$$

Applying (9.26), (9.29), and Corollary 8.6, it follows that

(9.30) 
$$\hat{f}_k = \hat{f}_{k+1} \quad \text{on } \widehat{\mathbb{X}}_k^{\text{o}}, \ k \ge 1.$$

Therefore, we may glue  $\hat{f}_k$  together in order to obtain the desired extension function  $\hat{f}$  as

(9.31) 
$$\hat{f} = \lim_{k \to \infty} \hat{f}_k \quad \text{on } \hat{X}^{\circ}.$$

In virtue of this formula and (9.26), (9.29)–(9.30), we conclude that f satisfies  $\binom{*}{**}$ . Hence, the proof of Part 1) of Theorem A is complete.

Using  $\binom{*}{**}$  and arguing as in Step 3 of Section 7, Part 2) of Theorem A follows. The remaining parts of Theorem A can also be proved using Step 5 and 6 of Section 7 and making the obviously necessary changes.

This completes the last step of Theorem A in the case N = 2.

Finally, we present a sketch of the argument for the general case N > 2. In order to prove Theorem A in its full generality, we use induction twice. More precisely, we proceed by induction (I) on  $N \ge 2$ . Suppose the theorem is true for  $N - 1 \ge 2$ . We have to discuss the case of an N-fold cross  $X := \mathbb{X}(A_1, \ldots, A_N; D_1, \ldots, D_N)$ , where  $D_1, \ldots, D_N \subset \mathbb{C}$  and  $A_1, \ldots, A_N$  are linearly measurable subsets of  $\partial D_1, \ldots, \partial D_N$ such that  $D_j$  is locally rectifiable on  $A_j$   $(1 \le j \le N)$ .

We proceed again by induction (II) on the positive integer j ( $0 \le j \le N$ ) such that there are at least j rectifiable Jordan domains among the open sets  $D_1, \ldots, D_N$ . For j = N we are reduced to Section 7.

In fact, our proof follows essentially the scheme of the works in [14], [12], and that of Sections 7 and the previous proof for N = 2.

### 10. Proof of Theorem B

We will only give the proof of Theorem B for the case when N = 2 and  $D_1$ ,  $D_2$  are rectifiable Jordan domains. Since the general case can be proved using the scheme of Section 7 and 9, it is left to the interested reader. The proof is divided into two steps.

**Step 1:** Proof of Theorem B for the case when the slice functions  $f(a_1, \cdot)|_{D_2}$  and  $f(\cdot, a_2)|_{D_1}$  are bounded for every  $a_1 \in A_1$  and  $a_2 \in A_2$ .

Proof of Step 1. For any  $M \in \mathbb{N}$  let

(10.1)

$$A_{1M} := \{a_1 \in A_1 : |f(a_1, \cdot)|_{D_2} \le M\} \text{ and } A_{2M} := \{a_2 \in A_2 : |f(\cdot, a_2)|_{D_1} \le M\}.$$

Using the assumption of Step 1 and (10.1), we obtain

(10.2)  $A_{jM} \nearrow A_j$  as  $M \nearrow \infty$ ,  $j \in \{1, 2\}$ .

Now we would like to show that for  $j \in \{1, 2\}, M \in \mathbb{N}$ ,

(10.3)

 $A_{jM}$  is a closed subset of  $A_j$  and  $f|_{A_{jM} \times D_k} \in \mathcal{C}(A_{jM} \times D_k), \ k \neq j, \ k \in \{1, 2\}.$ 

To do this, suppose without loss of generality that j = 1, k = 2, and fix an arbitrary  $M \in \mathbb{N}$  and a point  $a_1^0 \in A_1$ . Let  $(a_{1n})_{n=1}^{\infty}$  be a sequence in  $A_{1M}$  such that  $\lim_{n \to \infty} a_{1n} = a_1^0$ . Consequently, using the hypothesis (i), we see that

(10.4) 
$$\lim_{n \to \infty} f(a_{1n}, t) = f(a_1^0, t), \qquad t \in A_2.$$

On the other hand, it follows from the assumption  $(a_{1n})_{n=1}^{\infty} \subset A_{1M}$  and the hypothesis of Step 1 that

$$|f(a_{1n}, \cdot)|_{D_2} \le M$$
 and  $|f(a_1^0, \cdot)|_{D_2} < \infty$ .

Combining this and (10.4), we are able to apply Lemma 4.4 to the sequence  $(f(a_{1n}, \cdot)|_{D_2})_{n=0}^{\infty} \subset \mathcal{O}(D_2)$ , where  $a_{10} := a_1^0$ . Consequently, the sequence  $(f(a_{1n}, \cdot))_{n=1}^{\infty}$  converges uniformly on compact subsets of  $D_2$  to  $f(a_1^0, \cdot)$ . This completes the proof of (10.3).

On the other hand, by hypothesis (ii), the holomorphic function  $f(a_1, \cdot)$  admits the angular limit  $f(a_1, a_2)$  at  $a_2 \in A_2$ . Hence, it follows that  $f|_{A_{1M} \times A_{2M}}$  is measurable. Moreover, by (10.1),  $|f|_{\mathbb{X}(A_{1M}, A_{2M}; D_1, D_2)} \leq M$  for every  $M \in \mathbb{N}$ . In addition, in virtue of (10.2), there exists a sufficiently large integer  $M_0$  such that  $\operatorname{mes}(A_{jM}) > 0$  for  $j \in \{1, 2\}$  and  $M \geq M_0$ . Consequently, we are in a position to apply Theorem A to the function f restricted to the cross  $\mathbb{X}(A_{1M}, A_{2M}; D_1, D_2)$  for  $M \geq M_0$ . Therefore, we obtain a function  $\hat{f}_M \in \mathcal{O}\left(\widehat{\mathbb{X}}^o(A_{1M}, A_{2M}; D_1, D_2)\right)$  and a subset  $\widetilde{A}_{1M}$  (resp.  $\widetilde{A}_{2M}$ ) of  $A_{1M}$  (resp.  $A_{2M}$ ) for  $M \geq M_0$ , such that

(10.5) 
$$\operatorname{mes}(A_{1M} \setminus \tilde{A}_{1M}) = \operatorname{mes}(A_{2M} \setminus \tilde{A}_{2M}) = 0,$$
$$\hat{f}_M \text{ admits the angular limit } f \text{ on } \left(\tilde{A}_{1M} \times D_2\right) \bigcup \left(D_1 \times \tilde{A}_{2M}\right).$$

Put

(10.6) 
$$\tilde{A}_1 := \bigcup_{M=M_0}^{\infty} \tilde{A}_{1M}$$
 and  $\tilde{A}_2 := \bigcup_{M=M_0}^{\infty} \tilde{A}_{2M}$ .

Using (10.5), we deduce from (10.6) that

(10.7) 
$$\operatorname{mes}(A_1 \setminus \tilde{A}_1) = \operatorname{mes}(A_2 \setminus \tilde{A}_2) = 0,$$
$$\hat{f}_M \text{ admits the angular limit } f \text{ on } \left(\tilde{A}_1 \times D_2\right) \bigcup \left(D_1 \times \tilde{A}_2\right).$$

Applying (10.2), (10.7), and Corollary 8.6, we obtain

(10.8) 
$$\hat{f}_M = \hat{f}_{M+1}$$
 on  $\widehat{\mathbb{X}}^o\left(\tilde{A}_{1M}, \tilde{A}_{1M}; D_1, D_2\right), \ M \ge M_0.$ 

Therefore, we may glue the  $\hat{f}_M$  together to obtain the desired extension function  $\hat{f}$  as

(10.9) 
$$\hat{f} = \lim_{M \to \infty} \hat{f}_M \quad \text{on } \hat{X}^\circ = \widehat{\mathbb{X}}^\circ \left( A_1, A_2; D_1, D_2 \right).$$

Next, for every  $j \in \{1, 2\}$  and  $M \ge M_0$ , in virtue of (10.2)–(10.3) and (10.5), one may find a sequence  $(F_{j,M,n})_{n=1}^{\infty}$  of compact subsets of  $\partial D_j$  such that

(10.10)  

$$F_{j,M,n} \subset F_{j,M,n+1} \subset A_j,$$

$$\operatorname{mes}(F_{j,M,n}) > 0,$$

$$\operatorname{mes}\left(\tilde{A}_{jM} \setminus \bigcup_{n=1}^{\infty} F_{j,M,n}\right) = 0$$

Moreover, for any  $k \in \mathbb{N}$ ,  $k \ge 1$ , and for any  $m \in \mathbb{N}$ , put (10.11)

$$A_{1Mnmk} := \left\{ a_1 \in A_{1M} : |f(a_1, \zeta) - f(a_1, \eta)| \le \frac{1}{2k^2}, \quad \forall \zeta, \eta \in F_{2,M,n} : |\zeta - \eta| < \frac{1}{m} \right\},$$
$$A_{2Mnmk} := \left\{ a_2 \in A_{2M} : |f(\zeta, a_2) - f(\eta, a_2)| \le \frac{1}{2k^2}, \quad \forall \zeta, \eta \in F_{1,M,n} : |\zeta - \eta| < \frac{1}{m} \right\}.$$

Since, by hypothesis (i),  $f \in C_s(A_1 \times A_2)$ , we deduce from (10.10) and (10.11) that  $A_{jMnmk}$  is a closed subset of  $A_{jM}$  and

(10.12) 
$$A_{jMnmk} \nearrow A_{jM} \quad \text{as } m \nearrow \infty, \ j \in \{1, 2\}, \ k \ge 1.$$

Consequently, there is an  $m_0 := m_0(M, n, k)$  such that  $\operatorname{mes}(A_{jMnmk} \cap F_{1,M,n}) > 0$ for any  $m > m_0$ . Now we are in a position to apply Theorem A to the function f restricted on the cross  $\mathbb{X}(A_{1Mnmk} \cap F_{1,M,n}, A_{2Mnmk} \cap F_{2,M,n}; D_1, D_2)$ . Using (10.7)–(10.9) and Corollary 8.6, we obtain exactly the function  $\hat{f}$  restricted to  $\widehat{\mathbb{X}}^{\circ}(A_{1Mnmk} \cap F_{1,M,n}, A_{2Mnmk} \cap F_{2,M,n}; D_1, D_2)$ . Let

(10.13) 
$$\tilde{A}_{jMnmk} := (A_{jMnmk} \cap F_{j,M,n}) \cap (A_{jMnmk} \cap F_{j,M,n})^*,$$

where  $T^*$  denotes as usual the set of locally regular points relative to T.

Taking (10.11)-(10.13) into account and arguing as in Step 5 of Section 7, we may show that

(10.14) 
$$\operatorname{mes}\left(\tilde{A}_{jMnmk} \setminus F_{j,n}\right) = 0,$$
$$\lim_{z \to a, \ z_1 \in \mathcal{A}_{\alpha}(a_1), \ z_2 \in \mathcal{A}_{\alpha}(a_2), \ z \in \hat{X}^{\circ}} |\hat{f}(z) - f(a)| < \frac{1}{k}, \qquad 0 < \alpha < \frac{\pi}{2}$$

for every  $a = (a_1, a_2) \in \tilde{A}_{1Mnmk} \times \tilde{A}_{2Mnmk}$ . Now it suffices to put

$$\tilde{A}_j := \bigcap_{k=1}^{\infty} \bigcup_{M=M_0}^{\infty} \bigcup_{n=1}^{\infty} \bigcup_{m=m_0(M,n,k)}^{\infty} \tilde{A}_{jMnmk}, \qquad j \in \{1,2\}.$$

Combining this and (10.14), (10.12), (10.2), we may check that all the conclusions 1)-3 of Theorem B are satisfied. Hence the proof is complete in this first step.  $\Box$ 

Step 2: The general case.

*Proof of Step 2.* Arguing as in Section 7, it suffices to treat the case when  $D_1 = D_2 =$  the unit disc *E*. We begin with the following

**Definition 10.1.** For every closed subset F of  $\partial E$  and any  $n \in \mathbb{N}$ , n > 1, define the following open set

$$\Delta = \Delta(F, n) := \bigcup_{\zeta \in F} \left\{ z \in \mathcal{A}_{\frac{\pi}{4}}(\zeta) : |z| \ge 1 - \frac{1}{n} \right\} \cup \mathbb{B}\left(0, 1 - \frac{1}{n}\right).$$

The reader should compare this definition with Definition 6.1. Below we give a list of properties of such open sets.

### **Proposition 10.2.** Let F be a closed subset of $\partial E$ .

1) Let  $\Delta(F,n)$  be as in Definition 10.1, then  $\Delta(F,n)$  is a rectifiable Jordan domain and  $F \subset \partial \Delta(F,n)$ .

2)  $\Delta(F,n) \nearrow E$  as  $n \nearrow \infty$ .

3) Consider a locally bounded function  $f : E \cup F \longrightarrow \mathbb{C}$ . Then  $|f|_{\Delta(F,n)} < \infty$  for every  $n \in \mathbb{N}, n > 1$ .

4) There holds the following equality

$$\omega(z, F, E) = \lim_{n \to \infty} \omega(z, F, \Delta(F, n)), \qquad z \in E.$$

*Proof of Proposition 10.2.* Part 1) may be done as in the proof of Proposition 6.2.

Part 2) is an immediate consequence of Definition 10.1.

Part 3) follows immediately from the compactness of F.

The proof of Proposition 5.15 still works in the context of Part 4) making the obviously necessary changes. This completes Part 4).  $\Box$ 

Now we are in a position to complete Step 2. Indeed, first suppose that both  $A_1$  and  $A_2$  are closed. Then for  $j \in \{1, 2\}$ , consider the sequence of rectifiable Jordan domain  $(D_{jn})_{n=2}^{\infty}$  given by

$$D_{jn} := \Delta(A_j, n), \qquad n \in \mathbb{N}, \ n > 1.$$

For  $n \in \mathbb{N}$ , n > 1, let  $f_n := f|_{\mathbb{X}(A_1,A_2;D_{1n},D_{2n})}$ . In virtue of Proposition 10.2, we are able to apply the result of Step 1 to  $f_n$ . Consequently, we obtain a function  $\hat{f}_n \in \widehat{\mathbb{X}}^o(A_1, A_2; D_{1n}, D_{2n})$ . Therefore, we may glue  $\hat{f}_n$  together in order to obtain the desired extension function  $\hat{f}$  as

$$\hat{f} = \lim_{n \to \infty} \hat{f}_n$$
 on  $\widehat{X}^{\circ} = \widehat{\mathbb{X}}^{\circ} (A_1, A_2; D_1, D_2)$ .

Because of Proposition 10.2, we can show that  $\hat{f}$  possesses all the assertions of Theorem B.

The case when  $A_1$  and  $A_2$  are only measurable is similar. It suffices to find a sequence  $(A_{jn})_{n=1}^{\infty}$  of subsets of  $A_j$  such that  $A_{jn}$  is compact and mes  $\left(A_j \setminus \bigcup_{n=1}^{\infty} A_{jn}\right) =$ 0. Then we may apply the previous discussion to  $f|_{\mathbb{X}(A_{1n},A_{2n};D_1,D_2)}$  in order to obtain a function  $\hat{f}_n \in \widehat{\mathbb{X}}^{o}(A_{1n}, A_{2n}; D_1, D_2)$ . Finally, the desired extension function  $\hat{f}$  is defined by

$$\hat{f} = \lim_{n \to \infty} \hat{f}_n$$
 on  $\hat{X}^{\circ} = \hat{\mathbb{X}}^{\circ} (A_1, A_2; D_1, D_2)$ .

This completes the proof in this last step.

#### 11. Examples and Concluding Remarks

The following examples of Drużkowski [2] show the optimality of Theorem A and B.

Consider N = 2,  $D_1 = D_2 = E$ ,  $A_1 = A_2 = \{t \in \partial E : \text{Re } t > 0\}$ ,  $A := A_1 \times A_2$ ,  $X := \mathbb{X}(A_1, A_2; D_1, D_2)$ , and  $Y := (D_1 \cup A_1) \times (D_2 \cup A_2)$ . **Example 1.** Define a function  $h: Y \longrightarrow \mathbb{C}$  as follows

$$h(z) := \begin{cases} \exp\left(-\left[\log\left(1-z_1\right) + \log\left(1-z_2\right)\right] \log\frac{2+z_1z_2}{3}\right), & z_1 \neq 1, \ z_2 \neq 1\\ 0, & z_1 = 1 \text{ or } z_2 = 1 \end{cases}$$

where Log is the principal branch of logarithm.

Put  $f := h|_X$ . As in [2] observe that f is measurable,  $f \in \mathcal{C}_s(X) \cap \mathcal{O}_s(X^\circ)$ ,  $|f|_X < \infty$ , but  $f|_A$  is not continuous at (1, 1). Since  $h|_{\widehat{X}^\circ} \in \mathcal{O}(\widehat{X}^\circ)$ , using the uniqueness established in Theorem A, we conclude that the solution  $\widehat{f}$  provided by Theorem A and B satisfies  $\widehat{f} = h|_{\widehat{X}^\circ}$ . In addition, we see that, for  $0 < \alpha < \frac{\pi}{2}$ , the angular limit of  $\widehat{f}$  at (1, 1) does not exist. Thus the condition in assertion 3) of Theorem A is necessary. Moreover, the sets  $\widetilde{A}_1$ ,  $\widetilde{A}_2$  given by Theorem B do depend on f.

**Example 2.** Define a function  $h: Y \longrightarrow \mathbb{C}$  as follows

$$h(z) := \begin{cases} \exp\left(-(z_1 - \lambda) \operatorname{Log}^2 \frac{3+z_2}{1-z_2}\right), & z_2 \neq 1\\ 0, & z_2 = 1 \end{cases}.$$

where  $z \in Y$ ,  $0 < \lambda \leq \frac{\sqrt{2}}{2}$ .

Define  $f := h|_X$ . Then  $\hat{f} = h|_{\hat{X}^o}$ . As in [2] observe that  $f|_A$  is continuous,  $f \in \mathcal{C}_s(X) \cap \mathcal{O}_s(X^o)$ , but f is not locally bounded on X.

In addition, for  $\frac{\pi}{3} < \alpha < \frac{\pi}{2}$ , consider the functions  $z_1^{\alpha,\lambda}, z_2^{\alpha} : [0,1] \to \mathbb{C}$  given by

$$z_{2}^{\alpha}(t) := 1 + te^{i\left(\pi - \frac{9\alpha}{10}\right)},$$
  
$$z_{1}^{\alpha,\lambda}(t) := \lambda + \left(\operatorname{Re}\operatorname{Log}^{2}\frac{3 + z_{2}^{\alpha}(t)}{1 - z_{2}^{\alpha}(t)}\right)^{-1} + i\lambda, \qquad t \in [0, 1].$$

We may prove that there is an  $t_{\alpha,\lambda} > 0$  and a neighborhood  $U_{\alpha,\lambda}$  of  $\lambda + i\lambda$  in  $\mathbb{C}$  such that

$$\left( z_1^{\alpha,\lambda}(t), z_2^{\alpha}(t) \right) \in \begin{cases} \left( \left( \mathcal{A}_{\alpha}(\lambda + i\lambda) \cap U_{\alpha,\lambda} \right) \times \mathcal{A}_{\alpha}(1) \right) \cap \widehat{X}^{\circ}, & 0 < t < t_{\alpha,\lambda}, \ \lambda = \frac{\sqrt{2}}{2} \\ \left( U_{\alpha,\lambda} \times \mathcal{A}_{\alpha}(1) \right) \cap \widehat{X}^{\circ}, & 0 < t < t_{\alpha,\lambda}, \ 0 < \lambda < \frac{\sqrt{2}}{2} \end{cases} \end{cases}$$

58

In addition, it can be checked that

$$\lim_{t \to 0} \left( z_1^{\alpha,\lambda}(t), z_2^{\alpha}(t) \right) = (\lambda + i\lambda, 1) \quad \text{and} \quad \lim_{t \to 0} \left| \hat{f} \left( z_1^{\alpha,\lambda}(t), z_2^{\alpha}(t) \right) \right| = \infty.$$

This shows that the assumption of the local boundedness on f is necessary in Theorem A.

Finally, we conclude the article by some remarks and open questions.

1. We may generalize Theorems A and B in the following directions: Let  $D \subset \mathbb{C}$  be an open set.

- 1a. We extend  $\partial^* D$  to the set of all points  $\zeta \in \partial D$  where  $\partial D$  admits a corner at  $\zeta$  (see [13]). Note that there are at most countably many points where  $\partial D$  admits a corner but not a tangent (see Exercises 3.4 in [13]).
- 1b. We extend the notion type of a point in  $\partial D$  as follows. A point  $\zeta \in \partial D$  is said to be of type  $m \ (m \ge 1)$  if there are an open neighborhood U of  $\zeta$  and mdisjoint rectifiable Jordan domains  $U_1, \ldots, U_m$  such that  $U \cap D = \bigcup_{i=1}^m U_i$ .
- 1c. Instead of the notion *local rectifiability*, we use the weaker notion *local Jordan* curve. More precisely, D is said to be *locally Jordan curve like* at a point  $\zeta \in \partial D$  if there is a neighborhood U of  $\zeta$  in  $\mathbb{C}$  such that  $U \cap \partial D$  is the union of m disjoint Jordan curves and the interior of at least one of them contains  $\zeta$ .

Since the corresponding statements of Theorems A and B under these generalizations are somewhat complicated and their proofs do not require any new method, we leave the interested reader to develop these ideas.

2. It may be proved that  $\widehat{X}^{\circ}$  provided by Theorem A is the maximal domain of holomorphic extension of the function f. We postpone the proof of this result to an ongoing work.

3. It seems to be of interest to consider Theorem A and B under the following general settings: Let  $G_j$  be a complex manifold of dimension  $d_j$  and  $D_j \Subset G_j$  an open set,  $j = 1, \ldots, N$ . Let  $A_j$  be a subset of positive  $d_j$ -dimensional Hausdorff measure of  $M_j$ , where  $M_j$  is a real  $d_j$ -dimensional generating submanifold contained in  $\partial D_j$ ,  $j = 1, \ldots, N$ , etc. We postpone this issue to an ongoing work.

4. Does Theorem A still hold if we omit the assumption (ii) " $f|_A$  is measurable"?

5. Does Theorem B still hold if we omit the assumption that  $f|_A \in \mathcal{C}_s(A)$ ?

### References

- L. Aizenberg, Carleman's formulas in complex analysis. Theory and applications, Transl. from the Russian, updated, enlarged and revised. (English) Mathematics and its Applications (Dordrecht). 244. Dordrecht: Kluwer Academic Publishers. xx, 299 p. (1993).
- [2] L. M. Drużkowski, A generalization of the Malgrange-Zerner theorem, Ann. Polon. Math., 38, (1980), 181–186.
- [3] H. Epstein, Generalization of the "Edge of the Wedge" Theorem, J. Math. Phys., 1, (1960), 524-531.
- [4] G. M. Goluzin, Geometric theory of functions of a complex variable, (English), Providence, R. I.:American Mathematical Society (AMS). VI, (1969), 676 p.

- [5] A. A. Gonchar, On analytic continuation from the "edge of the wedge" theorem, Ann. Acad. Sci. Fenn. Ser. A.I: Mathematica, 10, (1985), 221–225.
- [6] A. A. Gonchar, On Bogolyubov's "edge of the wedge" theorem, Proc. Steklov Inst. Math., 228, (2000), 18–24.
- [7] M. Jarnicki, P. Pflug, *Extension of Holomorphic Functions*, de Gruyter Expositions in Mathematics 34, Walter de Gruyter, 2000.
- [8] M. Jarnicki, P. Pflug, An extension theorem for separately holomorphic functions with analytic singularities, Ann. Pol. Math., 80, (2003), 143–161.
- [9] M. Klimek, *Pluripotential theory*, London Mathematical society monographs, Oxford Univ. Press., 6, (1991).
- [10] H. Komatsu, A local version of Bochner's tube theorem, J. Fac. Sci., Univ. Tokyo, Sect. I A 19, (1972), 201–214.
- [11] P. Koosis, Introduction to H<sub>p</sub> spaces. With an appendix on Wolff's proof of the corona theorem, London Mathematical Society Lecture Note Series. 40. Cambridge etc.: Cambridge University Press. XV, 376 p. (1980). (1991), 183–194.
- [12] V.-A. Nguyên, A general version of the Hartogs extension theorem for separately holomorphic mappings between complex analytic spaces, preprint of the University of Oldenburg (2004), 33 p.
- [13] Ch. Pommerenke, Boundary behaviour of conformal maps, Grundlehren der Mathematischen Wissenschaften, 299, Berlin: Springer-Verlag, 300 p. (1992).
- [14] P. Pflug and V.-A. Nguyên, A boundary cross theorem for separately holomorphic functions, Ann. Polon. Math., 84, (2004), no. 3, 237–271.
- [15] T. Ransford, Potential theory in the complex plane, London Mathematical Society Student Texts, 28, Cambridge: Univ. Press., (1995).
- [16] W. Rudin, Lectures on the edge-of-the-wedge theorem, (English) Conference Board of the Mathematical Sciences. Regional Conference Series in Mathematics. No.6. Providence, R.I.:American Mathematical Society (1971).
- [17] W. Rudin, Function theory in the unit ball of  $\mathbb{C}^n$ , (English) Grundlehren der Mathematischen Wissenschaften **241**, Springer-Verlag, New York-Berlin, 1980.
- [18] W. Rudin, Function theory in polydiscs (English) Mathematics Lecture Note Series. New York-Amsterdam: W.A. Benjamin, Inc., 188 p. (1969).
- [19] E. M. Stein, Boundary behavior of holomorphic functions of several complex variables, (English) Mathematical Notes. Princeton, N. J.: Princeton University Press. IX, 72 p. (1972).
- [20] V. S. Vladimirov, Methods of the Theory of Functions of Several Complex Variables, (Russian), Moscow: 'Nauka' Publisher, 411 p. (1964).
- [21] M. Zerner, Quelques résultats sur le prolongement analytique des fonctions de variables complexes, Séminaire de Physique Mathématique.

Peter Pflug, Carl von Ossietzky Universität Oldenburg, Fachbereich Mathe-Matik, Postfach 2503, D–26111, Oldenburg, Germany

E-mail address: pflug@mathematik.uni-oldenburg.de

VIỆT-ANH NGUYÊN, CARL VON OSSIETZKY UNIVERSITÄT OLDENBURG, FACHBEREICH MATHEMATIK, POSTFACH 2503, D-26111, OLDENBURG, GERMANY *E-mail address*: nguyen@mathematik.uni-oldenburg.de