

ON NORMAL SUBGROUPS OF PRODUCT OF GROUPS

ASHISH KUMAR DAS AND DIPANKAR DEB

Abstract. The object of this paper is to find a necessary and sufficient condition for a pair of groups G_1 and G_2 so that every normal subgroup of $G_1 \times G_2$ is of the type $N_1 \times N_2$ where $N_i \in G_i$, $i = 1, 2$.

1. Introduction

Let G be a finite group. Then G has a composition series given by $\text{feg} = H_0 \subset H_1 \subset \dots \subset H_n = G$. The set with multiplicities of composition factors associated to this composition series is given by $C(G) = \{H_i/H_{i-1} : i = 1, 2, \dots, n\}$; noting that any two composition series of G give rise to the same set with multiplicities of composition factors, upto isomorphism of factors (see [2], Jordan-Hölder Theorem). It is a standard fact (see [1]) that $\bigcup_{K \subset G} C(K)$ is the disjoint union (i.e. union counting multiplicities) of the sets $C(G=K)$ and $C(K)$. By convention $C(\text{feg}) = \emptyset$.

In [3], Leinster has proved that if G_1 and G_2 are any two finite groups such that the sets $C(G_1)$ and $C(G_2)$ have no member in common then every normal subgroup of $G_1 \times G_2$ is of the type $N_1 \times N_2$ where $N_i \in G_i$, $i = 1, 2$. However, considering $G_1 = G_2 =$ the alternating group A_5 , one can see that the converse is not true.

In this paper we derive a condition which is necessary as well as sufficient for any pair of groups G_1 and G_2 (finite or infinite, abelian or non-abelian) so that all normal subgroups of $G_1 \times G_2$ are of the above mentioned type. This condition seems to provide more options for finding examples of non-abelian perfect groups; a perfect group being one in which the sum of the orders of all normal subgroups equals twice the order of the group (see [3]).

2. The NS-condition

We begin this section with a few definitions.

1991 Mathematics Subject Classification. 20D 06, 20D 40.

Key words and phrases. normal subgroups, composition series, simple groups.

Definition 2.1. Let G_1 and G_2 be any two groups. A normal subgroup N of $G_1 \times G_2$ is said to be of standard type if $N = N_1 \times N_2$ where $N_i \in G_i, i = 1, 2$:

Definition 2.2. Two groups G_1 and G_2 are said to have a subgroup in common if there exist non-trivial subgroups H_i of $G_i, i = 1, 2$; such that $H_1 = H_2$.

Definition 2.3. Two groups G_1 and G_2 are said to satisfy NS-condition if $H_i \in G_i, i = 1, 2$; the centres $Z(G_1/H_1)$ and $Z(G_2/H_2)$ of the quotient groups G_1/H_1 and G_2/H_2 have no subgroup in common.

Remark 2.4. If $H \in G$ then the subgroups of $Z(G/H)$ are of the type K/H where $H \in K \leq G$ and $gkg^{-1}k^{-1} \in H, \forall g \in G$ and $k \in K$. In particular $K \in G$:

The following proposition shows that NS-condition is strictly weaker than Leinster's hypothesis (see Proposition 3.1 of [3]).

Proposition 2.5. If G_1 and G_2 are two finite groups such that the sets $C(G_1)$ and $C(G_2)$ have no member in common then G_1 and G_2 satisfy NS-condition, but not conversely.

Proof. Suppose $H_i \in G_i, i = 1, 2$; are such that the centres $Z(G_1/H_1)$ and $Z(G_2/H_2)$ have a subgroup in common. So, there are non-trivial subgroups K_i/H_i of $Z(G_i/H_i), i = 1, 2$; such that $K_1/H_1 = K_2/H_2$. Then $C(K_1/H_1) = C(K_2/H_2)$. Since $H_i \in K_i \in G_i$; we have $C(K_i/H_i) \leq C(K_i) \leq C(G_i), i = 1, 2$. Thus, $C(G_1)$ and $C(G_2)$ have a member in common.

That the converse is not true can be easily seen by considering $G_1 = G_2 =$ a finite simple non-abelian group or a product of such groups.

Proposition 2.6. Two finite groups G_1 and G_2 satisfy NS-condition if and only if $\gcd(n_1, n_2) = 1$ where

$$n_i = \prod_{H_i \in G_i} |Z(G_i/H_i)|, i = 1, 2;$$

Proof. If each n_i is divisible by a prime p , then p divides $|Z(G_i/H_i)|$ for some $H_i \in G_i, i = 1, 2$. So, each of $Z(G_i/H_i), i = 1, 2$, has a subgroup of order p (see [1]). Obviously such a subgroup is isomorphic to Z_p .

Conversely, if $\gcd(n_1, n_2) = 1$ then $H_i \in G_i, i = 1, 2$, the centres $Z(G_1/H_1)$ and $Z(G_2/H_2)$ have coprime orders and so they can not have any subgroup in common.

The following corollary is immediate.

Corollary 2.7. If two finite groups G_1 and G_2 have coprime orders then they satisfy NS-condition. Converse also holds if G_1 and G_2 are finite abelian.

3. Main Theorem

In this section we show that the NS-condition is indeed necessary as well as sufficient.

Theorem 3.1. Let G_1 and G_2 be any two groups (finite or infinite, abelian or non-abelian). Then all normal subgroups of $G_1 \times G_2$ are of standard type if and only if G_1 and G_2 satisfy NS-condition.

Proof. Suppose G_1 and G_2 satisfy NS-condition. Let $N \in \mathcal{N}(G_1 \times G_2)$. Set $H_1 = \pi_1((G_1 \times e_2g) \setminus N)$ and $H_2 = \pi_2((e_1g \times G_2) \setminus N)$ where e_i are identities of G_i and $\pi_i : G_1 \times G_2 \rightarrow G_i$ are projections, $i = 1, 2$. Then $H_1 \times e_2g, e_1g \times H_2 \subseteq N \subseteq \pi_1(N) \times \pi_2(N)$ and so

$$H_1 \times H_2 = (H_1 \times e_2g)(e_1g \times H_2) \subseteq \pi_1(N) \times \pi_2(N) \quad (3.2)$$

It may be noted here that $H_i \in \mathcal{N}(G_i)$ and $H_i \in \mathcal{N}(G_i)$, $i = 1, 2$. Now, suppose $a_1 \in \pi_1(N)$. Then $(a_1, a_2) \in N$ for some $a_2 \in G_2$; in fact $a_2 \in \pi_2(N)$. Therefore, $g_1 a_1 g_1^{-1} \in G_1$; we have

$$\begin{aligned} (g_1 a_1 g_1^{-1}, a_2) &= (g_1, e_2)(a_1, a_2)(g_1^{-1}, e_2) \in N \\ \Rightarrow (g_1 a_1 g_1^{-1}, a_1^{-1}, e_2) &\in N \\ \Rightarrow g_1 a_1 g_1^{-1} a_1^{-1} &\in H_1 \\ \Rightarrow g_1 H_1 a_1 H_1 &= a_1 H_1 g_1 H_1 \subseteq G_1 : \end{aligned}$$

Thus, $a_1 H_1 \subseteq Z(G_1 = H_1)$. So, we have $\pi_1(N) = H_1 \subseteq Z(G_1 = H_1)$. Similarly, $\pi_2(N) = H_2 \subseteq Z(G_2 = H_2)$. Note that if $a_1, b_1 \in \pi_1(N)$ then $(a_1, a_2), (b_1, b_2) \in N$ for some $a_2, b_2 \in \pi_2(N)$; and so $(a_1 b_1^{-1}, a_2 b_2^{-1}), (a_1 b_1, a_2 b_2) \in N$. Therefore,

$$\begin{aligned} a_1 H_1 &= b_1 H_1 \quad (\quad) \quad a_1 b_1^{-1} \in H_1 \quad (\quad) \quad (a_1 b_1^{-1}, e_2) \in N \\ (\quad) \quad (e_1, a_2 b_2^{-1}) &\in N \quad (\quad) \quad a_2 b_2^{-1} \in H_2 \quad (\quad) \quad a_2 H_2 = b_2 H_2 : \end{aligned}$$

This means that we have a well-defined injective map $f : \pi_1(N) = H_1 \rightarrow \pi_2(N) = H_2$ given by $f(a_1 H_1) = a_2 H_2$ where $(a_1, a_2) \in N$. Also, $f(a_1 H_1 b_1 H_1) = f(a_1 b_1 H_1) = a_2 b_2 H_2 = a_2 H_2 b_2 H_2 = f(a_1 H_1) f(b_1 H_1)$, showing that f is a homomorphism. Finally, if $b \in \pi_2(N)$ then $(a, b) \in N$ for some $a \in \pi_1(N)$ and so $f(a H_1) = b H_2$, which implies that f is surjective. Thus f is an isomorphism. Hence it follows from the hypothesis that $\pi_i(N) = H_i$ are trivial subgroups of $Z(G_i = H_i)$, $i = 1, 2$. Therefore, $H_i = \pi_i(N)$; $i = 1, 2$, and so $N = H_1 \times H_2$; by (3.2).

Conversely, suppose G_1 and G_2 do not satisfy the NS-condition. So, there exist $H_i \in G_i, i = 1, 2$, such that $Z(G_1 = H_1)$ and $Z(G_2 = H_2)$ have a subgroup in common. Let $K_i = H_i$ be non-trivial subgroups of $Z(G_i = H_i), i = 1, 2$, such that there is an isomorphism $F: K_1 = H_1 \rightarrow K_2 = H_2$. Put $N = \{f(a_1; a_2) \in K_1 \times K_2 : F(a_1 H_1) = a_2 H_2\}$. Let $(a_1; a_2); (b_1; b_2) \in N$ then $F(a_1 H_1) = a_2 H_2$ and $F(b_1 H_1) = b_2 H_2$. So, $F(a_1 b_1^{-1} H_1) = a_2 b_2^{-1} H_2$. Thus $(a_1; a_2)(b_1; b_2)^{-1} = (a_1 b_1^{-1}; a_2 b_2^{-1}) \in N$, showing that N is a subgroup of $G_1 \times G_2$. Again let $(a_1; a_2) \in N$ and $(g_1; g_2) \in G_1 \times G_2$. Then, $(g_1; g_2)(a_1; a_2)(g_1; g_2)^{-1} = (g_1 a_1 g_1^{-1}; g_2 a_2 g_2^{-1}) \in K_1 \times K_2$, since, by Remark 2.4, $K_i \in G_i, i = 1, 2$. Also, since $a_i H_i \in K_i = H_i \subseteq Z(G_i = H_i), i = 1, 2$, we have

$$F(g_1 a_1 g_1^{-1} H_1) = F(a_1 H_1) = a_2 H_2 = g_2 a_2 g_2^{-1} H_2.$$

Thus $(g_1; g_2)(a_1; a_2)(g_1; g_2)^{-1} \in N$, and so $N \in G_1 \times G_2$. On the other hand, suppose N is of standard form $N_1 \times N_2$ where $N_i \in G_i, i = 1, 2$. Then, $N_i \cap (N) = N_i, i = 1, 2$. But since F is bijective, we have $N_i \cap (N) = K_i, i = 1, 2$. Therefore, $N = K_1 \times K_2$. Since $K_1 = H_1$ is non-trivial, there is some $a_1 \in K_1$ such that $a_1 H_1 \notin H_1$. But $(a_1; e_2) \in K_1 \times K_2 = N$. So, $F(a_1 H_1) = e_2 H_2 = H_2$, the zero element of $K_2 = H_2$. Therefore, since F is injective, we have $a_1 H_1 = H_1$, the zero element of $K_1 = H_1$. This contradiction shows that N is not of standard type.

As an immediate consequence, using induction and the fact that the centre of a product of groups equals the product of the centres of the constituent groups, we have

Corollary 3.3. Let $G = \prod_{i=1}^n G_i$ where all or all but one of the groups G_i are simple non-abelian. Then every normal subgroup of G is of the standard type $\prod_{i=1}^n N_i$ where $N_i \in G_i, i = 1, 2$.

References

- [1] J. B. Fraleigh, A first course in abstract algebra, Addison-Wesley Publishing Company, Inc. Philippines, 1982.
- [2] N. Jacobson, Basic algebra - vol I, W. H. Freeman and Company, U.S.A., 1974
- [3] T. Leinster, Perfect number and groups, arXiv math.GR/0104012 v1 Apr 2001.

Department of Mathematics, North Eastern Hill University, Permanent Campus, Shillong-793022, Meghalaya, India.

E-mail address: akdas@nehu.ac.in