

# CONJUGACY CLASSES AND FINITE $p$ -GROUPS

EDITH ADAN-BANTE

**ABSTRACT.** Let  $G$  be a finite  $p$ -group, where  $p$  is a prime number and  $a \in G$ . Denote by  $\text{Cl}(a) = \{gag^{-1} \mid g \in G\}$  the conjugacy class of  $a$  in  $G$ . Assume that  $|\text{Cl}(a)| = p^n$ . Then  $\text{Cl}(a)\text{Cl}(a^{-1}) = \{xy \mid x \in \text{Cl}(a), y \in \text{Cl}(a^{-1})\}$  is the union of at least  $n(p-1) + 1$  distinct conjugacy classes of  $G$ .

## 1. INTRODUCTION

Let  $G$  be a finite group. Denote by  $\text{Cl}(a) = \{gag^{-1} \mid g \in G\}$  the conjugacy class of  $a$  in  $G$ , and by  $|\text{Cl}(a)|$  the size of  $\text{Cl}(a)$ . If the subset  $X$  of  $G$  is  $G$ -invariant, i.e.  $X^g = \{x^g \mid x \in X\} = X$  for all  $g \in G$ , then  $X$  is the union of  $m$  distinct conjugacy classes of  $G$ , for some integer  $m$ . Set  $\eta(X) = m$ .

Given any conjugacy classes  $\text{Cl}(a)$  and  $\text{Cl}(b)$ , we can check that the product  $\text{Cl}(a)\text{Cl}(b) = \{xy \mid x \in \text{Cl}(a), y \in \text{Cl}(b)\}$  is a  $G$ -invariant set. In this note, we will explore the relation between  $|\text{Cl}(a)|$  and  $\eta(\text{Cl}(a)\text{Cl}(a^{-1}))$ . Those results are the equivalent in conjugacy classes as some of the ones in irreducible characters in [1] and [2].

In Theorem A of [2], it is proved that if  $G$  is a  $p$ -group,  $\chi$  is an irreducible character with degree  $p^n$ , then the product  $\chi\bar{\chi}$  of  $\chi$  with its complex conjugate  $\bar{\chi}$  is the sum of at least  $2n(p-1) + 1$  distinct irreducible characters. The following is the equivalent for conjugacy classes

**Theorem A.** *Let  $G$  be a finite  $p$ -group and  $a \in G$ . Assume that  $|\text{Cl}(a)| = p^n$ . Then the product  $\text{Cl}(a)\text{Cl}(a^{-1})$  of the conjugacy class of  $a$  in  $G$  and the conjugacy class of the inverse of  $a$  in  $G$ , is the union of at least  $n(p-1) + 1$  distinct conjugacy classes of  $G$ , i.e.  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) \geq n(p-1) + 1$ .*

In Proposition 5.3, it is shown that for every prime  $p$  and every integer  $n \geq 0$ , there exist a  $p$ -group  $G$  and a conjugacy class  $\text{Cl}(a)$  of  $G$  such that  $|\text{Cl}(a)| = p^n$  and  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) = n(p-1) + 1$ . Thus the bound in Theorem A is optimal.

An application of Theorem A is the following

**Theorem B.** *Let  $n$  be a positive integer. Then there exists a finite set  $S_n$  of positive integers such that for any nilpotent group  $G$  and any conjugacy class  $\text{Cl}(a)$  of  $G$  with  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) \leq n$ , we have that*

$$|\text{Cl}(a)| \in S_n.$$

In Proposition 5.5, we prove that given any prime  $p$ , there exist a supersolvable group and a conjugacy class  $\text{Cl}(a)$  of  $G$  with  $|\text{Cl}(a)| = p$  and  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) = 2$ . Thus the previous result does not remain true assuming the weaker hypothesis that

*Date:* 2004.

*1991 Mathematics Subject Classification.* 20D15.

*Key words and phrases.* Conjugacy class,  $p$ -groups, products.

the groups are supersolvable. Theorem A is the equivalent in conjugacy classes of Theorem B of [1].

**Theorem C.** *Let  $p$  be a prime number. Let  $G$  be a finite  $p$ -group and  $\text{Cl}(a)$  be a conjugacy class of  $G$ . Then one of the following holds:*

- i)  $|\text{Cl}(a)| = 1$  and  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = 1$ .
- ii)  $|\text{Cl}(a)| = p$  and  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = p$ .
- iii)  $|\text{Cl}(a)| \geq p^2$  and  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) \geq 2p - 1$ .

Given a fix prime  $p > 2$ , observe that Theorem C implies that there are “gaps” among the possible values that  $\eta(\text{Cl}(a) \text{Cl}(a^{-1}))$  can take for any finite  $p$ -group and any conjugacy class  $\text{Cl}(a)$  in  $G$ . The previous result is the equivalent in conjugacy classes of Theorem B of [2].

**Acknowledgment.** I would like to thank Manoj Kumar for bringing to my attention products of conjugacy classes. I also want to thank Professor Everett C. Dade for useful advise and corrections.

## 2. PROOF OF THEOREM A

**Notation.** Let  $G$  be a finite  $p$ -group and  $N$  be a normal subgroup of  $G$ . Denote by  $\bar{a}$  the element in  $G/N$  that contains  $a$ . Thus  $\text{Cl}(\bar{a})$  is the conjugacy class of  $\bar{a}$  in  $G/N$ .

**Lemma 2.1.** *Let  $G$  be a finite  $p$ -group and  $N$  be a normal subgroup of  $G$ . Let  $a$  and  $b$  be elements of  $G$ . Then*

- i)  $\text{Cl}(\bar{a}) \text{Cl}(\bar{b})$  is a  $G$ -invariant set. If  $\text{Cl}(\bar{a}) \cap \text{Cl}(\bar{b}) = \emptyset$  then  $\text{Cl}(a) \cap \text{Cl}(b) = \emptyset$ . Thus  $\eta(\text{Cl}(\bar{a}) \text{Cl}(\bar{b})) \leq \eta(\text{Cl}(a) \text{Cl}(b))$ .
- ii) If, in addition,  $|N| = p$ , then either  $|\text{Cl}(\bar{a})| = |\text{Cl}(a)|$  or  $|\text{Cl}(\bar{a})| = \frac{|\text{Cl}(a)|}{p}$ . Furthermore, if  $|\text{Cl}(\bar{a})| = \frac{|\text{Cl}(a)|}{p}$ , then  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) \geq \eta(\text{Cl}(\bar{a}) \text{Cl}(\bar{a}^{-1})) + (p-1)$ .

*Proof.* i) Clearly if  $a = gbg^{-1}$ , then  $\bar{a} = \bar{g}\bar{b}(\bar{g})^{-1}$ . Thus if  $\text{Cl}(\bar{a}) \cap \text{Cl}(\bar{b}) = \emptyset$  then  $\text{Cl}(a) \cap \text{Cl}(b) = \emptyset$ . Therefore  $\eta(\text{Cl}(\bar{a}) \text{Cl}(\bar{b})) \leq \eta(\text{Cl}(a) \text{Cl}(b))$ .

ii) Since  $N$  is normal,  $|N| = p$  and  $G$  is a  $p$ -group, then  $N$  is contained in the center  $\mathbf{Z}(G)$  of  $G$ . Thus given any  $n \in N$ ,  $\text{Cl}(n) = \{n\}$ .

Suppose that  $|\text{Cl}(\bar{a})| \neq |\text{Cl}(a)|$ . Since  $|N| = p$ , we have that  $|\text{Cl}(a)| \leq p|\text{Cl}(\bar{a})|$ . Therefore  $\frac{|\text{Cl}(a)|}{p} \leq |\text{Cl}(\bar{a})| \leq |\text{Cl}(a)|$ . Thus  $|\text{Cl}(\bar{a})| = \frac{|\text{Cl}(a)|}{p}$  since  $G$  is a  $p$ -group and  $|\text{Cl}(\bar{a})|$  divides  $|G/N|$ .

If  $|\text{Cl}(\bar{a})| = \frac{|\text{Cl}(a)|}{p}$ , then given any  $x \in \text{Cl}(a)$  and any  $n \in N$ , we have that  $nx \in \text{Cl}(a)$ . Thus  $n = nx(x^{-1}) \in \text{Cl}(a) \text{Cl}(a^{-1})$  for any  $n \in N$ . Therefore  $N \leq \text{Cl}(a) \text{Cl}(a^{-1})$  and ii) follows.  $\square$

*Proof of Theorem A.* We are going to use induction on the order of  $G$ . Let  $N$  be a normal subgroup of  $G$  of order  $p$ . Observe such group exists since  $G$  is a  $p$ -group. Let  $|\text{Cl}(\bar{a})| = p^m$ . Since  $|G/N| < |G|$ , by induction we have that  $\eta(\text{Cl}(\bar{a}) \text{Cl}(\bar{a}^{-1})) \geq m(p-1) + 1$ . If  $|\text{Cl}(\bar{a})| = |\text{Cl}(a)|$ , i.e if  $m = n$ , then by Lemma 2.1 i) we have that

$$\eta(\text{Cl}(a) \text{Cl}(a^{-1})) \geq \eta(\text{Cl}(\bar{a}) \text{Cl}(\bar{a}^{-1})) \geq m(p-1) + 1 = n(p-1) + 1.$$

We may assume then that  $|\text{Cl}(\bar{a})| \neq |\text{Cl}(a)|$ . By Lemma 2.1 ii), we have that  $m = n - 1$  and

$$\begin{aligned} \eta(\text{Cl}(a) \text{Cl}(a^{-1})) &\geq \eta(\text{Cl}(\bar{a}) \text{Cl}(\bar{a}^{-1})) + (p-1) \\ &= (n-1)(p-1) + 1 + (p-1) = n(p-1) + 1. \end{aligned}$$

□

## 3. PROOF OF THEOREM B

*Proof of Theorem B.* Let

$$S_n = \{\Pi(p_i)^{t_i} \mid p_i \text{ is a prime number for all } i, t_i \geq 0 \text{ and } t_i(p_i - 1) + 1 \leq n\}.$$

Observe that the set  $S_n$  is a finite set of positive integers since  $0 \leq t_i \leq n$  and if  $t_i > 0$  then  $p_i \leq n$ .

Let  $\{p_1, \dots, p_r\}$  be the set of distinct prime divisors of  $|G|$ . For  $i = 1, \dots, r$ , let  $P_i$  be the Sylow  $p_i$ -subgroup of  $G$ . Observe that  $a = \prod_{i=1}^r a_i$ , for some  $a_i \in P_i$  for  $i = 1, \dots, r$ . Since  $G$  is nilpotent, we have that  $\text{Cl}(a) = \prod_{i=1}^r \text{Cl}(a_i)$ , where  $\text{Cl}(a_i)$  is the conjugacy class of  $a_i$  in  $P_i$ , for  $i = 1, \dots, r$ . Let  $m_i = \eta(\text{Cl}(a_i) \text{Cl}(a_i^{-1}))$ . Observe that  $m_i \leq n$  and

$$|\text{Cl}(a)| = \prod_{i=1}^r |\text{Cl}(a_i)|.$$

We can check that

$$\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = \prod_{i=1}^r \eta(\text{Cl}(a_i) \text{Cl}(a_i^{-1})).$$

For each  $i$ , let  $|\text{Cl}(a_i)| = p_i^{t_i}$ . Since  $\text{Cl}(a_i)$  is the conjugacy class of  $a_i$  in the  $p_i$ -group  $P_i$ , by Theorem A we have that  $m_i = \eta(\text{Cl}(a_i) \text{Cl}(a_i^{-1})) \geq t_i(p_i - 1) + 1$ . Thus  $n \geq t_i(p_i - 1) + 1$ . Therefore  $|\text{Cl}(a)| = \prod_{i=1}^r |\text{Cl}(a_i)| = \prod_{i=1}^r p_i^{t_i} \in S_n$ . □

## 4. PROOF OF THEOREM C

**Lemma 4.1.** *Let  $G$  be a finite  $p$ -group and  $\text{Cl}(a)$  be a conjugacy class of  $G$  with  $|\text{Cl}(a)| = p$ . Then one of the following holds:*

- i)  $\text{Cl}(a) = \{az \mid z \in Z\}$  for some subgroup  $Z$  of the center  $\mathbf{Z}(G)$  of  $G$ . Therefore  $\text{Cl}(a) \text{Cl}(a^{-1}) = Z$  and  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = p$ .
- ii)  $\text{Cl}(a) \text{Cl}(a^{-1})$  is the union of  $p - 1$  distinct conjugacy classes of size  $p$  and the class  $\text{Cl}(e) = \{e\}$ . Therefore  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = p$ .

*Proof.* Observe that if  $z \in \text{Cl}(a) \text{Cl}(a^{-1})$  and  $|\text{Cl}(z)| = 1$ , then  $z$  is in the center  $\mathbf{Z}(G)$  of  $G$ . Since  $z = a^g a^{-1}$  for some  $g \in G$  and  $z \in \mathbf{Z}(G)$ ,  $z^i \in \text{Cl}(a) \text{Cl}(a^{-1})$  and  $a^{g^i} = az^i$  for all integer  $i$ . Thus  $\langle z \rangle \leq \text{Cl}(a) \text{Cl}(a^{-1})$ . Set  $Z = \langle z \rangle$ .

i) If  $z \neq e$ , it follows that  $|Z| \geq p$ . Since  $|\text{Cl}(a)| = p$  and  $a^{g^i} = az^i$  for all integer  $i$ , we have that  $\text{Cl}(a) = \{az \mid z \in Z\}$  and  $|Z| = p$ . Since  $Z$  is contained in  $\mathbf{Z}(G)$ , then  $\text{Cl}(a) \text{Cl}(a^{-1}) = Z$  and  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = p$ .

ii) We may assume now that if  $z \in \text{Cl}(a) \text{Cl}(a^{-1})$  and  $|\text{Cl}(z)| = 1$ , then  $z = e$ . Thus all the conjugacy classes different from  $\text{Cl}(e)$  are of size  $p$ . Observe that  $a^{g(a^{-1})^g} = e$  for all  $g \in G$ . Thus  $|\text{Cl}(a) \text{Cl}(a^{-1})| \leq p^2 - p + 1 = (p - 1)p + 1$ . Therefore by Theorem A it follows that  $\text{Cl}(a) \text{Cl}(a^{-1})$  is the union of  $p - 1$  distinct conjugacy classes of size  $p$  and  $\text{Cl}(e)$ . □

**Remark.** Let  $p$  be a prime number.

a) Let  $G$  be an extra special group of order  $p^3$  and exponent  $p$ . We can check that given any  $a \in G$ , where  $a$  is not in the center of  $G$ , then  $\text{Cl}(a) \text{Cl}(a^{-1}) = \mathbf{Z}(G)$  and thus Lemma 4.1 i) occurs.

b) Let  $G$  be the wreath product of a cyclic group  $C_{p^2}$  of order  $p^2$  by a cyclic group  $C_p$  of order  $p$ . Thus  $|G| = p^{2p+1}$ . Let  $a = (c, e, \dots, e)$  in  $G$ , where  $c \in C_{p^2}$  has order  $p^2$ . Observe that  $a^{-1} \neq a$ . Observe also that

$$\text{Cl}(a) = \{(c, e, \dots, e), (e, c, \dots, e), \dots, (e, e, \dots, e, c)\}.$$

Thus  $|\text{Cl}(a)| = p$ . Let  $b_i = (c, e, \dots, c^{-1}, e, \dots, e)$ , where  $c^{-1}$  is in the  $i$ -position for  $i = 0, \dots, p-1$ , i.e.  $b_0 = (cc^{-1}, \dots, e) = (e, e, \dots, e)$ ,  $b_1 = (c, c^{-1}, e, \dots, e)$  and so for. Observe that  $\text{Cl}(b_0) = \text{Cl}((e, e, \dots, e))$  has class size 1. We can check that  $\text{Cl}(b_i)$  has size  $p$  for  $i = 1, \dots, p-1$ . Since  $c \neq c^{-1}$ , then  $\text{Cl}(b_i) \cap \text{Cl}(b_j) = \emptyset$  if  $i \neq j$  and  $i, j = 0, \dots, p-1$ . Observe that  $\text{Cl}(a)\text{Cl}(a^{-1}) = \cup_{i=0}^{p-1} \text{Cl}(b_i)$ . Thus  $\text{Cl}(a)\text{Cl}(a^{-1})$  is the union of a conjugacy class of size 1, namely  $\text{Cl}(b_0)$  and  $p-1$  distinct conjugacy classes of size  $p$ , namely  $\text{Cl}(b_i)$  for  $i = 1, \dots, p-1$ . We conclude that given any prime  $p$ , there exist some group  $G$  and some conjugacy class  $\text{Cl}(a)$  of  $G$  satisfying the condition in case ii) of Lemma 4.1.

*Proof of Theorem C.* If  $|\text{Cl}(a)| = \{a\}$ , then  $\text{Cl}(a)\text{Cl}(a^{-1}) = \{e\}$  and so i) holds. Lemma 4.1 implies ii) and iii) follows from Theorem A.  $\square$

## 5. EXAMPLES

**Lemma 5.1.** *Let  $G_0$  be a  $p$ -group and  $\text{Cl}(g_0)$  be the conjugacy class containing  $g_0 \in G_0$ . Assume that  $\text{Cl}(g_0) \neq \text{Cl}(g_0^{-1})$ . Let  $N = G_0 \times G_0 \times \dots \times G_0$  be the direct product of  $p$ -copies of  $G_0$ . Let  $C = \langle c \rangle$  be a cyclic group of order  $p$ . Observe that  $C$  acts on  $N$  by*

$$(5.2) \quad c : (n_0, n_1, \dots, n_{p-1}) \mapsto (n_{p-1}, n_0, \dots, n_{p-2})$$

for any  $(n_0, n_1, \dots, n_{p-1}) \in N$ .

Let  $G$  be the semidirect product of  $N$  and  $C$ , i.e.  $G$  is the wreath product of  $G_0$  and  $C$ . Set  $a = (g_0, e, \dots, e)$  in  $N$ , where  $e$  is the identity of  $G_0$ . Then  $|\text{Cl}(a)| = p|\text{Cl}(g_0)|$ ,  $\text{Cl}(a) \neq \text{Cl}(a^{-1})$  and  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) = \eta(\text{Cl}(g_0)\text{Cl}(g_0^{-1})) + (p-1)$ .

*Proof.* Observe that  $\text{Cl}(a) \neq \text{Cl}(a^{-1})$  since  $\text{Cl}(g_0) \neq \text{Cl}(g_0^{-1})$ .

Let  $\text{Cl}(g_0)\text{Cl}(g_0^{-1}) = C_1 \cup C_2 \dots \cup C_m$ , where  $C_1, \dots, C_m$  are distinct conjugacy classes of  $G_0$ . Thus  $m = \eta(\text{Cl}(g_0)\text{Cl}(g_0^{-1}))$ . We can check that the distinct conjugacy classes of  $\text{Cl}(a)\text{Cl}(a^{-1})$  are of the following two types:

- i)  $\{(x, e, \dots, e)^c \mid x \in C_i, c \in C\}$  for  $i = 1, \dots, m$ .
- ii)  $\{(x, y, \dots, e, e)^c \mid x \in \text{Cl}(g_0), y \in \text{Cl}(g_0^{-1}), c \in C\}$ ,  $\{(x, e, y, \dots, e)^c \mid x \in \text{Cl}(g_0), y \in \text{Cl}(g_0^{-1}), c \in C\}$  and  $\{(x, e, \dots, e, y)^c \mid x \in \text{Cl}(g_0), y \in \text{Cl}(g_0^{-1}), c \in C\}$ .

Observe that there are  $\eta(\text{Cl}(g_0)\text{Cl}(g_0^{-1}))$  distinct conjugacy classes of type i) and exactly  $p-1$  distinct conjugacy classes of type ii). Thus  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) = \eta(\text{Cl}(g_0)\text{Cl}(g_0^{-1})) + (p-1)$ .  $\square$

**Proposition 5.3.** *Given any prime  $p$ , and any integer  $n \geq 0$ , there exist a finite  $p$ -group  $G$  and a conjugacy class  $\text{Cl}(a)$  of  $G$  with  $|\text{Cl}(a)| = p^n$ ,  $\text{Cl}(a) \neq \text{Cl}(a^{-1})$  and  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) = n(p-1) + 1$ .*

*Proof.* Observe that if  $G$  is an abelian group and  $a \in G$  has order  $p^2$ , then  $|\text{Cl}(a)| = 1$ ,  $\text{Cl}(a) \neq \text{Cl}(a^{-1})$  and  $\eta(\text{Cl}(a)\text{Cl}(a^{-1})) = 1 = 0(p-1) + 1$ . Thus the statement is true for  $n = 0$ . Assume by induction that the statement is true for  $n-1$ , i.e. there exist a finite  $p$ -group  $G_0$  and a conjugacy class  $\text{Cl}(g_0)$  of  $G_0$  with  $|\text{Cl}(g_0)| = p^{n-1}$ ,

$\text{Cl}(g_0) \neq \text{Cl}(g_0^{-1})$  and  $\eta(\text{Cl}(g_0) \text{Cl}(g_0^{-1})) = (n-1)(p-1) + 1$ . Using the notation of Lemma 5.1, we have that

$$\begin{aligned} \eta(\text{Cl}(a) \text{Cl}(a^{-1})) &= \eta(\text{Cl}(g_0) \text{Cl}(g_0^{-1})) + (p-1) \\ &= (n-1)(p-1) + 1 + (p-1) = n(p-1) + 1. \end{aligned}$$

Since  $|\text{Cl}(a)| = p |\text{Cl}(g_0)| = p \times p^{n-1} = p^n$ , the proof is complete.  $\square$

**Hypothesis 5.4.** Fix a prime  $p$  and let  $F = \{0, 1, \dots, p-1\}$  be the finite field with  $p$  elements. Observe that  $F$  is also a vector space of dimension 1 over itself. Let  $A = \text{Aff}(F)$  be the affine group of  $F$ . Observe that the group  $A$  is a cyclic by cyclic group and thus it is supersolvable.

Let  $C$  be a cyclic group of order  $p$ . Set  $X = F$  and  $K = C^X = \{f : X \rightarrow C\}$ . Observe that  $K$  is a group via pointwise multiplication, and clearly  $A$  acts on this group (via its action on  $X$ ).

Let  $G$  be the wreath product of  $C$  and  $A$  relative to  $X$ , i.e.  $G = K \rtimes A$ . We can check that  $G$  is a supersolvable group.

**Proposition 5.5.** Assume Hypotheses 5.4. Set  $a = (c, e, e, \dots, e)$  in  $K$ . Then  $a \in G$ , the conjugacy class  $\text{Cl}(a)$  of  $G$  has size  $p$  and  $\text{Cl}(a) \text{Cl}(a^{-1}) = \text{Cl}((e, e, \dots, e)) \cup \text{Cl}((c, c^{-1}, e, \dots, e))$ . Thus  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = 2$ .

Therefore, given any prime  $p$ , there exist a supersolvable group  $G$  and a conjugacy class  $\text{Cl}(a)$  of  $G$  with  $|\text{Cl}(a)| = p$  and  $\eta(\text{Cl}(a) \text{Cl}(a^{-1})) = 2$ .

*Proof.* Observe that  $\text{Cl}(a) = \{(c, e, \dots, e), (e, c, e, \dots, e), \dots, (e, e, \dots, e, c)\}$ . Thus  $\text{Cl}(a)$  has  $p$ -elements. Observe that

$$\{(c, c^{-1}, \dots, e)^y \mid y \in F \setminus \{0\}\} = \{(c, c^{-1}, e, \dots, e), (c, e, c^{-1}, \dots, e), \dots, (c, e, \dots, e, c^{-1})\}.$$

Observe also that

$$\{(c, c^{-1}, e, \dots, e)^x \mid x \in F\} = \{(c, c^{-1}, e, \dots, e), \dots, (e, e, \dots, c, c^{-1}), (c^{-1}, e, \dots, e, c)\}.$$

Thus

$$\begin{aligned} \text{Cl}((c, c^{-1}, e, \dots, e)) &= \{(c, c^{-1}, e, \dots, e)^x, (c, e, c^{-1}, \dots, e)^x, \dots, \\ &\quad (c, e, \dots, c^{-1}, e), (c, e, \dots, e, c^{-1})^x \mid x \in F\} \end{aligned}$$

has  $(p-1)p = p^2 - p$  elements. Since  $a^g(a^{-1})^g = (e, \dots, e)$ , then  $\text{Cl}(a) \text{Cl}(a^{-1})$  has at most  $p^2 - p + 1$  elements. We conclude that

$$\text{Cl}(a) \text{Cl}(a^{-1}) = \text{Cl}((e, e, \dots, e)) \cup \text{Cl}((c, c^{-1}, e, \dots, e)).$$

$\square$

## REFERENCES

- [1] E. Adan-Bante, Products of characters and finite  $p$ -groups, J. Algebra 277 (1), 236-255.
- [2] E. Adan-Bante, Products of characters and finite  $p$ -groups II, Arch. Math. (82) (2004), 289-297.

UNIVERSITY OF SOUTHERN MISSISSIPPI GULF COAST, 730 EAST BEACH BOULEVARD, LONG BEACH MS 39560

E-mail address: Edith.Bante@usm.edu