

ON THE HOMOLOGY OF THE SPACE OF KNOTS

R. BUDNEY AND F. R. COHEN*

ABSTRACT. Consider the space of ‘long knots’ in $\mathbb{R}^3$ $\mathcal{K}_{3,1}$. This is the space of knots as studied by V. Vassiliev. Based on work in [5, 9], it follows that the rational homology of $\mathcal{K}_{3,1}$ is a free Poisson algebra. A rough, as yet incomplete, ‘approximation’ for a basis is given. In addition, the mod- p homology of this space is a ‘free, restricted Poisson algebra’. The main new results here is that there is p -torsion of all orders in the integer homology $\mathcal{K}_{3,1}$.

This leads to some natural questions about the homotopy type of $\mathcal{K}_{n,1}$, the space of long knots in $\mathbb{R}^n$ for $n > 3$, as well as consequences for the space of smooth embeddings of S^1 in S^3 .

1. INTRODUCTION

The purpose of this paper is to give homological properties of the classical spaces of smooth ‘long’ embeddings $\mathcal{K}_{3,1} = \text{Emb}(\mathbb{R}, \mathbb{R}^3)$ and smooth embeddings $\text{Emb}(S^1, S^3)$. Some results here also apply to the embedding spaces $\text{Emb}(S^k, S^n)$, and ‘long’ embedding spaces $\mathcal{K}_{n,k} = \text{Emb}(\mathbb{R}^k, \mathbb{R}^n)$, with the main results focused on the 3-dimensional case $k = 1$, $n = 3$.

The approach here to these homological problems follows recent work of Hatcher [18] and Budney [5]. The homotopy type of ‘most’ of the components of $\text{Emb}(S^1, S^3)$ and $\mathcal{K}_{3,1}$ are understood completely in terms of configuration spaces in the plane, Stiefel manifolds and various natural iterated bundle operations. Many of the homological properties of both $\mathcal{K}_{3,1}$, and $\text{Emb}(S^1, S^3)$ follow from combining this information together with earlier work of Cohen [9] on configuration spaces.

In addition, the space of long knots was shown to be a free $\mathcal{C}_2$ -space in the sense of May, with “generating set” given by the space of prime long knots [5]. One consequence is that the homotopy type of the space of long knots is determined completely by the homotopy type of the prime long knots. Information concerning spaces of prime long knots is combined with bundle theoretic constructions to give a large contribution to the homology groups for spaces of long knots, as well as $\text{Emb}(S^1, S^3)$.

The structure of a Poisson algebra arises in the work here. An introduction to Poisson algebras is given in [8], pages 177-182 while some applications are given in [9, 32]. A Poisson algebra A is a graded commutative algebra over $\mathbb{Q}$ given by A_n in gradation n together with a graded skew symmetric bilinear map

$$\{-, -\} : A_s \otimes A_t \rightarrow A_{s+t+1}$$

which satisfies

- (1) the Jacobi identity

$$\{\{a, \{b, c\}\} = \{\{a, b\}, c\} + (\pm 1)\{\{a, c\}, b\},$$

and

1991 *Mathematics Subject Classification*. Primary: 58D10, 57T25; Secondary: 57M25, 57Q45.

Key words and phrases. knots, embeddings, spaces, cubes, homology.

*Partially supported by the NSF grant number 9704410.

(2) the Leibniz formula

$$\{a \cdot b, c\} = a \cdot \{b, c\} + (\pm 1)b\{a, c\}.$$

A standard example of such a Poisson algebra is given by the rational homology algebra of $\Omega^2(X)$ for X a bouquet of spheres of dimension at least 3 [9].

The mod- p homology of $\mathcal{K}_{3,1}$ has more detailed structure, and is a ‘free restricted Poisson algebra’ with additional structure satisfied by free $\mathcal{C}_2$ -spaces [9]. In addition, these homology classes provide invariants of long knots; remarks concerning this point are given below.

There are certain components of $\text{Emb}(S^1, S^3)$ whose homotopy types are not yet completely understood which arise from certain ‘symmetric’ hyperbolic satellite operations: Whitehead doubles and ‘borromean’ hyperbolic satellite operations are some of the most elementary examples of this. Thus, the homological results given here for $\mathcal{K}_{3,1}$, and $\text{Emb}(S^1, S^3)$ are not yet in a closed form as they depend on the homology of components of knots given in terms of hyperbolic satellite operations.

These results lead to some natural questions about the structure of the homology of the higher-dimensional embedding spaces ($n > 3$) recently studied by Sinha [29], Volic [34] and Lambrechts [21]. Constructions related to these questions are also addressed here.

The main results of the current article are Theorem 1.3 on the structure of the homology of $\mathcal{K}_{3,1}$, Proposition 1.4 concerning implications for the space of smooth embeddings of S^1 in S^3 , and Corollary 1.5, a homological characterization of the unknot.

Definition 1.1. $\mathcal{K}_{n,j} := \{f : \mathbb{R}^j \rightarrow \mathbb{R}^n : f \text{ is an embedding and } f(x_1, x_2, \dots, x_j) = (x_1, x_2, \dots, x_j, 0, \dots, 0) \text{ for } |x| \geq 1\}$. $\mathcal{K}_{n,1}$ is traditionally called the space of long knots in $\mathbb{R}^n$, and $\mathcal{K}_{n,j}$ the space of long j -knots in $\mathbb{R}^n$. Given an element $f \in \mathcal{K}_{n,j}$ the connected-component of $\mathcal{K}_{n,j}$ containing f is denoted $\mathcal{K}_{n,j}(f)$. Two knots are considered equivalent if they are in the same connected component of $\mathcal{K}_{n,j}$ (That is the knots are isotopic).

The homotopy type of the space of long knots in $\mathbb{R}^3$, $\mathcal{K}_{3,1}$, was analyzed in work of Hatcher, and Budney [17, 18, 5, 6]. One result of Budney is recorded next [5]. Let $X_{\mathcal{K}} = \{f \in \mathcal{K}_{3,1} : f \text{ is prime}\}$, where the word ‘prime’ is used in the traditional sense of Schubert [28], that is $X_{\mathcal{K}}$ is the union of the connected components of $\mathcal{K}_{3,1}$ which contain prime knots. Alternatively, a knot can only fail to be prime in only two ways: 1) The unknot is not prime. 2) A connected sum of 2 or more non-trivial knots is not prime.

Theorem 1.2. *The space $\mathcal{K}_{3,1}$ is homotopy equivalent to*

$$C(\mathbb{R}^2, X_{\mathcal{K}} \amalg \{*\})$$

that is the labelled configuration space of points in the plane with labels in $X_{\mathcal{K}} \amalg \{\}$. Furthermore, the following hold:*

- (1) *Each path-component of $\mathcal{K}_{3,1}$ is a $K(\pi, 1)$.*
- (2) *The path-components of $\mathcal{C}_2(n) \times_{\Sigma_n} (X_{\mathcal{K}})^n$ for all n , and thus the path-components of $\mathcal{K}_{3,1}$ are given by*

$$\mathcal{C}_2(n) \times_{\Sigma_f} \prod_{i=1}^n \mathcal{K}_{3,1}(f_i)$$

for certain choices of $f_1, \dots, f_n \in X_{\mathcal{K}}$, and Young subgroups Σ_f .

The above theorem can be thought of as a generalization of Schubert’s Theorem that $\pi_0 \mathcal{K}_{3,1}$ is a free commutative monoid on countably-infinite many generators [28] to a space-level theorem about $\mathcal{K}_{3,1}$.

In general, $\mathcal{K}_{n,1}$ is a homotopy-associative H -space with multiplication induced by concatenation. Since $\mathcal{K}_{3,1}$ admits the action of the operad of little 2-cubes, there is an induced map

$$\Theta : S^1 \times \mathcal{K}_{3,1} \times \mathcal{K}_{3,1} \rightarrow \mathcal{K}_{3,1}$$


together with an operation in homology with any coefficients

$$H_s(\mathcal{K}_{3,1}) \otimes H_t(\mathcal{K}_{3,1}) \rightarrow H_{1+s+t}(\mathcal{K}_{3,1})$$

which is denoted, up to sign, by

$$\lambda_1(\alpha, \beta)$$

for α in $H_s(\mathcal{K}_{3,1})$, and β in $H_t(\mathcal{K}_{3,1})$. These operations satisfy the structure of a graded Poisson algebra [9] for which the bracket operation is called the Browder operation.



In what follows below, these bracket operations will be denoted

$$\{\alpha, \beta\} = \lambda_1(\alpha, \beta).$$

There is also a product operation

$$H_s(\mathcal{K}_{n,1}) \otimes H_t(\mathcal{K}_{n,1}) \rightarrow H_{s+t}(\mathcal{K}_{n,1})$$

which, together with the bracket operations, gives $H_*\mathcal{K}_{3,1}$ the structure of a graded Poisson algebra.

The next result uses the product operation above as well as the bracket operation $\{\alpha, \beta\} = \lambda_1(\alpha, \beta)$, and follows by interweaving the results of 1.2, and [9]. To state these results, additional information given by three functors from graded vector spaces V over a field $\mathbb{F}$ are described next with complete details given in section 6:

- (1) If the characteristic of $\mathbb{F}$ is 0, then the value of the functor on objects V is the symmetric algebra generated by an algebraically ‘desuspended’ free Lie algebra generated by the suspension of V , and denoted

$$S[\sigma^{-1}L[\sigma(V)]].$$

This last algebra is a free Poisson algebra.

- (2) If $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$, then the value of the functor on objects V is the symmetric algebra generated by an algebraically ‘desuspended’ free, mod-2 restricted Lie algebra generated by the suspension of V , and denoted

$$S[\sigma^{-1}L^{(2)}[\sigma(V)]].$$

This last algebra is a free restricted Poisson algebra.

- (3) If $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ for odd primes p , then the value of the functor on objects V is the symmetric algebra generated by an algebraically “desuspended” mod- p free restricted Lie algebra generated by the suspension of V plus an additional summand as described in section 6, and denoted

$$S[\sigma^{-1}L^{(p)}[\sigma(V)] \oplus \sigma^{-2}W^p[\sigma(V)]].$$

Theorem 1.3. *The homology of $\mathcal{K}_{3,1}$ satisfies the following properties.*

- (1) *The rational homology of $\mathcal{K}_{3,1}$ is a free Poisson algebra generated by $V = H_*(X_{\mathcal{K}}; \mathbb{Q})$.*
- (2) *The homology of $\mathcal{K}_{3,1}$ with $\mathbb{F}_p$ coefficients is a free Poisson algebra generated by $V = H_*(X_{\mathcal{K}}; \mathbb{F}_p)$ as described in [9].*
- (3) *There are isomorphisms of Hopf algebras*
 - (a) $S[\sigma^{-1}L[\sigma(V)]] \rightarrow H_*(\mathcal{K}_{3,1}; \mathbb{Q})$ for $V = H_*(X_{\mathcal{K}}; \mathbb{Q})$,
 - (b) $S[\sigma^{-1}L^{(2)}[\sigma(V)]] \rightarrow H_*(\mathcal{K}_{3,1}; \mathbb{Z}/2\mathbb{Z})$ for $V = H_*(X_{\mathcal{K}}; \mathbb{Z}/2\mathbb{Z})$, and
 - (c) $S[\sigma^{-1}L^{(p)}[\sigma(V)] \oplus \sigma^{-2}W^p[\sigma(V)]] \rightarrow H_*(\mathcal{K}_{3,1}; \mathbb{Z}/p\mathbb{Z})$ for $V = H_*(X_{\mathcal{K}}; \mathbb{Z}/p\mathbb{Z})$ in case p is an odd prime.

These isomorphisms specialize to an identification of the homology of each path-component of $\mathcal{K}_{3,1}$ with the one ambiguity that the homology of the components of knots arising from hyperbolic satellite operations is not given in a closed form here. More information is described in section 3.

- (4) *The integer homology of $\mathcal{K}_{3,1}$ has p -torsion of arbitrarily large order (with examples listed in section 6, and 7).*

By Theorem 1.3, there is arbitrarily large p -torsion in the homology of $\mathcal{K}_{3,1}$. Examples of knots whose path-components have higher 2-torsion in their integer homology is given next. This higher torsion can be regarded as a ‘measure’ of the ‘complexity’ of a knot, or more precisely, the ‘depth’ of its JSJ-tree.

- (1) Let $\mathcal{K}_{3,1}(f)$ denote the path-component of a non-trivial torus knot. Thus $\mathcal{K}_{3,1}(f)$ has the homotopy type of a circle [18].
- (2) Given any space X , and a strictly positive integer q , define

$$E(q, X) = \text{Conf}(\mathbb{R}^2, q) \times_{\Sigma_q} (X)^q.$$

Then $E(4, \mathcal{K}_{3,1}(f))$ has the property that $H_2(E(4, \mathcal{K}_{3,1}(f))) = \mathbb{Z}/2\mathbb{Z}$, and $H_3(E(4, \mathcal{K}_{3,1}(f))) = 0$ by [9].

- (3) Then $H_3(S^1 \times E(4, \mathcal{K}_{3,1}(f)))$ is isomorphic to $\mathbb{Z}/2\mathbb{Z}$.
- (4) Furthermore, $E(2^s, S^1 \times E(4, \mathcal{K}_{3,1}(f)))$ has the homotopy type of a path-component $\mathcal{K}_{3,1}(g)$ for a long knot g as given in [5], and has torsion of order 2^{s+1} in its integer homology by section 6, and 7 here. In the language of Section 2, $g = \#_{2^s}(c_{p/q}(\#_4(f)))$ ie, g is a connected-sum of 2^s copies of the same knot, which is a p/q -cable of a connected sum of four copies of the same torus knot.
- (5) A second example is $E(2^s, \mathcal{K}_{3,1}(h))$ where h is a hyperbolic satellite knot with $H_1(\mathcal{K}_{3,1}(h)) = \mathbb{Z}/2\mathbb{Z} \oplus A$ for some abelian group A . Such knots can be shown to exist – for example the Whitehead double of a trefoil knot.

A more complete description of the homology of $\mathcal{K}_{3,1}$ is given in sections 6, and 7. The homology of each path-component is given in terms of Theorem 1.3 as well as filtrations of the values of the functors listed in that theorem.

Consider the subspace $\mathcal{TK}_{3,1}$ of $\mathcal{K}_{3,1}$ consisting of the union of the components of $\mathcal{K}_{3,1}$ corresponding to knots which can be obtained from torus knots via iterated the operations: cabling and connected-sum. These are the knots whose complements have JSJ-decompositions built up entirely of Seifert-fibered manifolds. The structure of the

homology of the subspace $\mathcal{TK}_{3,1}$ of $\mathcal{K}_{3,1}$ is described in section 9. For example, some components of $\mathcal{TK}_{3,1}$, $\mathcal{K}_{3,1}(g)$, have the property that there is p^s -torsion for a fixed prime p , and a fixed positive integer s in the integer homology of $\mathcal{K}_{3,1}(g)$.

A further consequence of Theorem 1.2 is the next result which follows directly, and is proven in section 5. Let $\text{Emb}_*(S^1, S^n)$ denote the space of smooth pointed embeddings.

Proposition 1.4. *The group $SO(n-1)$ acts naturally on $\mathcal{K}_{n,1}$ (rotations perpendicular to the ‘long axis’), and there are morphisms of bundles for which each vertical map is a homotopy equivalence:*

$$\begin{array}{ccccc} SO(n) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{i} & SO(n+1) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{p} & SO(n+1)/SO(n) \\ \downarrow \Theta_n & & \downarrow \Theta_n & & \downarrow 1 \\ \text{Emb}_*(S^1, S^n) & \xrightarrow{j} & \text{Emb}(S^1, S^n) & \longrightarrow & S^n \end{array}$$

Thus, there is a bundle

$$SO(n+1) \times_{SO(n-1)} \mathcal{K}_{n,1} \rightarrow SO(n+1)/SO(n-1)$$

with fibre $\mathcal{K}_{n,1}$. Furthermore, there is a homeomorphism

$$\text{Emb}(S^1, S^3) \rightarrow S^3 \times \text{Emb}_*(S^1, S^3)$$

for which $\text{Emb}_*(S^1, S^3)$ denotes the space of smooth pointed embeddings, and the bundle

$$\mathcal{K}_{3,1} \rightarrow \text{Emb}_*(S^1, S^3) \rightarrow S^2$$

is the induced bundle with fiber $\mathcal{K}_{3,1}$ from the bundle

$$SO(2) \rightarrow SO(3) \rightarrow S^2.$$

Notice that the homological properties of each path component thus give knot invariants. These invariants are illustrated by the next direct corollary.

Corollary 1.5. (1) *A knot $f : S^1 \rightarrow S^3$ is the unknot if and only if its component in $\text{Emb}(S^1, S^3)$ contains no 2-torsion in its homology.*
 (2) *A long knot $f : \mathbb{R}^1 \rightarrow \mathbb{R}^3$ in $\mathcal{K}_{3,1}$ is the ‘long unknot’ if and only if its component has trivial homology.*

One question which arises here is whether Vassiliev invariants of long knots detect whether the path component of a long knot has non-trivial homology. The methods here in conjunction with T. Kohno’s analysis of Vassiliev invariants of braids [19] might be useful in addressing this question.

A classical theorem of R. Thom concerning the classical Steenrod Realization Problem gives that any mod-2 homology class for a space of the homotopy type of a CW-complex is represented by the fundamental cycle of a closed manifold. The next theorem addresses the question of representing mod-2 homology classes for $\mathcal{K}_{3,1}$ in the classical sense of Steenrod, and Thom. In particular, the mod-2 homology classes for $\mathcal{K}_{3,1}$ are represented by a restricted family of manifolds which is described next.

Consider manifolds M which represent the mod-2 homology of the spaces $\mathcal{K}_{3,1}(\mathcal{P})$ for prime knots $\mathcal{P}$. Consider the manifolds obtained by the following constructions.

- (1) finite products of M with finite products of circles,
- (2) the “wreath product construction” $S^1 \times_{\mathbb{Z}/2\mathbb{Z}} X^2$ for X one of the manifolds listed in 1 above, and
- (3) finite iterations of the constructions in 1, and 2 above.

Theorem 1.6. *Any class in the mod-2 homology of $\mathcal{K}_{3,1}$ is represented by the manifolds in 1-3 above.*

This last result provides information concerning the unoriented bordism ring of the space of long knots. In addition, the ‘group completion’ $\Omega B(\mathcal{K}_{3,1})$ of $\mathcal{K}_{3,1}$ exists. Furthermore, there is a natural bundle over the completion $\Omega B(\mathcal{K}_{3,1})$ whose Thom space has the property that the mod-2 cohomology is a free module over the mod-2 Steenrod algebra. These remarks are discussed in section 12.

One of the main results here is that the homology of the path-components of $\mathcal{K}_{3,1}$ is given iteratively in terms of the homology of ‘simpler’ components in Section 7. There exists torsion of all orders in the homology of $\mathcal{K}_{3,1}$. This is shown in section 8 using the results of section 7. The focus of section 8 restricts attention to knots whose JSJ-decompositions consist of Seifert Fibred manifolds (ie: knots whose exteriors are graph manifolds). Another way to say this is that the knots that can be obtained from the unknot by iterating cabling and connected-sum operations.

Much recent progress on the structure of spaces of embeddings via finite-dimensional model spaces and approximations. Some of this was first given by Vassiliev [33] and has been the subject of further study via the Goodwillie Calculus of Embeddings [29] [34] [21], or cohomological techniques such as Bott-Taubes integrals [4, 7].

The Poisson algebra considered above was first considered on the E^2 -level of the Vassiliev spectral sequence by Tourtchine [32]. Related progress is given in work of Altshuler-Freidel [2], Bar-Natan [3], Cattaneo, Cotta-Ramusino-Longoni [7], Kontsevich [20], Lescop [22], Polyak-Viro [27], as well as others.

This paper takes the path of using Allen Hatcher’s techniques for understanding the homotopy type of $\mathcal{K}_{3,1}$, one component at a time. The central construction of Hatcher [18] is to consider the components of the knot space as the classifying space of the mapping class group of the knot complement. One then studies how such a mapping class group acts on the JSJ-tree of the knot complement as in [5], using Hatcher’s results on the homotopy type of diffeomorphism groups of Haken manifolds [17] to assemble an answer. Thus most of the results here are complementary to the results of Vassiliev, Goodwillie, Sinha, Volic and Lambrechts.

TABLE OF CONTENTS

- 1: Introduction
- 2: Notation, labeling components
- 3: The homotopy type of $\mathcal{K}_{3,1}$
- 4: Actions on $\mathcal{K}_{3,1}$
- 5: Relations among various spaces
- 6: On the homology of $C_2(X \amalg \{*\})$
- 7: On the homology of $\mathcal{K}_{3,1}$
- 8: Higher p -torsion in the integer homology of $\mathcal{K}_{3,1}$
- 10: On the subspace generated by torus knots
- 11: Steenrod-Thom representability
- 12: Closed knots, and homology
- 13: Problems

2. NOTATION, LABELING COMPONENTS

Whitney [38] showed that the embedding space $\mathcal{K}_{n,j}$ is connected for $n > 2j + 1$. By work of Wu [35], $\mathcal{K}_{n,j}$ is also connected provided both inequalities:

$$n > 2j, \text{ and } j > 1$$

are satisfied. That $\mathcal{K}_{n,1}$ is also connected for $n = 1$ is elementary. The analogous result for $n = 2$ is the Alexander/Schoenflies theorem.

In case $n \leq 2j + 1$, the space $\mathcal{K}_{n,j}$ could potentially have many connected components. A primary example of this is the case $\mathcal{K}_{3,1}$ which has countably infinite many components (this is for example, an elementary consequence of Reidemeister’s theorem

and the Alexander polynomial). Given $f \in \mathcal{K}_{n,j}$ let $\mathcal{K}_{n,j}(f)$ denote the path-component of $\mathcal{K}_{n,j}$ containing f .

This section collects information in the indexing of the components of $\mathcal{K}_{3,1}$ which is given in terms of the Jaco-Shalen-Johannson (JSJ) decomposition of knot complements. Similar constructions have appeared in the works [5] and [10]. In the book by Eisenbud-Neumann [10] this method of indexing is called the splice decomposition of links.

The description given in [5] and [10] is expanded below to give a more complete treatment of the consequences of the JSJ-decomposition for knots in S^3 from the point of view of ‘satellite operations.’

An alternative long knot space is the space $\text{EC}(1, D^{n-1})$ of ‘framed’ long knots from [5]. Given a compact manifold M ,

$$\text{EC}(k, M) = \{f \in \text{Emb}(\mathbb{R}^k \times M, \mathbb{R}^k \times M), \text{supp}(f) \subset I^k \times M\}$$

Here the support of f , $\text{supp}(f)$ is defined by $\text{supp}(f) = \{x \in \mathbb{R}^k \times M : f(x) \neq x\}$ and $I = [-1, 1]$. $\text{EC}(1, D^{n-1})$ is not homotopy equivalent to $\mathcal{K}_{n,1}$ but as described in [5] there is a fibration

$$\Omega SO(n-1) \rightarrow \text{EC}(1, D^{n-1}) \rightarrow \mathcal{K}_{n,1}$$


which is trivial for $n \in \{1, 2, 3\}$. This allows us to think of $\mathcal{K}_{3,1}$ as a subspace of $\text{EC}(1, D^2)$. When we think of $\mathcal{K}_{3,1}$ as a subspace of $\text{EC}(1, D^2)$ we will use the notation $\hat{\mathcal{K}}$ for the space of long knots in $\mathbb{R}^3$, as in [5]. This is the space of ‘trivially framed’ long knots in $\mathbb{R}^3$, which has the same homotopy type as $\mathcal{K}_{3,1}$. The spaces $\text{EC}(1, D^{n-1})$ and $\hat{\mathcal{K}}$ admit an action of the operad of little 2-cubes [5]. In addition, using the connectedness of $\mathcal{K}_{n,1}$ for $n \geq 4$ together with the cubes action one can prove that $\text{EC}(1, D^{n-1})$ has the homotopy type of a 2-fold loop space for $n \geq 4$.

The previously described fibre bundle

$$\text{Emb}(S^1, S^3) \rightarrow SO(4)/SO(2)$$

whose fibre inclusion $i : \mathcal{K}_{3,1} \rightarrow \text{Emb}(S^1, S^3)$ is induced by the one-point compactification is explored more deeply in Section 5. For the purpose of this section and the study of the components of $\mathcal{K}_{3,1}$ and $\text{Emb}(S^1, S^3)$ respectively we simply treat the above result as saying there is a canonical bijective correspondence between $\pi_0 \mathcal{K}_{3,1}$, $\pi_0 \hat{\mathcal{K}}$ and $\pi_0 \text{Emb}(S^1, S^3)$ thus we move freely between the models when discussing our notation to index components.


Definition 2.1. Let $\nu f : S^1 \times D^2 \rightarrow S^3$ be a closed tubular neighborhood of the knot $f : S^1 \rightarrow S^3$. Let T be the boundary of the image of νf . Up to isotopy there are two distinguished oriented closed curves in T : l and m . l , the longitude of T , is the intersection of a Seifert surface for f with T , with orientation parallel to f (we demand that the Seifert surface intersects νf in an annulus). m is called the meridian of T and is any curve in T which has linking number 1 with f and which bounds a disc in $\text{img}(\nu f)$.



Definition 2.2. Given n long knots $f_i \in \hat{\mathcal{K}}$ and disjoint embeddings $g_i : S^1 \times D^2 \rightarrow I \times D^2$ for $i \in \{1, 2, \dots, n\}$ such that $g = \sqcup_{i=1}^n g_i : \sqcup_{i=1}^n S^1 \times D^2 \rightarrow I \times D^2$ is the n -component unlink, let $J_i = \text{int}(\text{img}(g_i))$, $J = \cup_{i=1}^n J_i$. Let $h_i : \mathbb{R} \times D^2 \rightarrow (I \times D^2) - J$ be an embedding such that $h_i(\{0\} \times S^1)$ is the (oriented) longitude of g_i and $h_i(\mathbb{R} \times S^1) \subset \partial J_i$. Let $h = \sqcup_{i=1}^n h_i$. Define an embedding $R[g, f, h] : \mathbb{R} \times D^2 - J \rightarrow \mathbb{R} \times D^2 - J$ with support contained in $\text{img}(h)$ so that on the image of h_i it is defined to be the function $h_i \circ f_i \circ h_i^{-1}$, for all $i \in \{1, 2, \dots, n\}$. We call $R[g, f, h]$ the re-embedding map associated to (g, f, h) .

Proposition 2.3. [6] Up to isotopy, $R[g, f, h]$ does not depend on the choice of the embeddings $h = \sqcup_{i=1}^n h_i$.


Since the isotopy class of $R[g, f, h]$ does not depend on h we sometimes abuse notation and refer to $R[g, f]$.



Definition 2.4. The standard 2-component p/q -Seifert link is defined to be the pair $(t_{p/q}, w)$ where $t_{p/q} \in \hat{\mathcal{K}}$ is a p/q -torus knot, $w : S^1 \times D^2 \rightarrow I \times D^2$ is an unknotted embedding disjoint from the image of $t_{p/q}$ making the link complement a Seifert-fibered manifold, fibered over a twice-punctured disc with one singular fiber. Thus we demand that $\gcd(p, q) = 1$ and $|q| \geq 1$.

Thus, for example, $(t_{0/1}, w)$ is the Hopf link.

Definition 2.5. Let $(t_{p/q}, w)$ be the p/q -Seifert link. Let $R[w, f]$ be the re-embedding function with respect to w and f , then the p/q -cabling of f is defined to be the composite $R[w, f] \circ t_{p/q}$. Denote the p/q -cabling of f by $c_{p/q}f \in \mathcal{K}_3$.



Observe that if g is the p/q -cabling of a knot f then if $q = 0$, g is the unknot. If $q = \pm 1$ then g and f are either isotopic or inverses of each other. A p/q -cabling $g = c_{p/q}f$ is said to be non-trivial if one of the following is true:

- $|p| \geq 2, |q| \geq 2$ and f is the unknot. In this case g is a torus knot.
- $|q| \geq 2$ and f is a non-trivial knot.


Proposition 2.6. [6] For two non-trivial cable knots, $\mathcal{K}_{c_{p/q}f_1} = \mathcal{K}_{c_{p'/q'}f_2}$ if and only if $\mathcal{K}_{f_1} = \mathcal{K}_{f_2}$ and one of the following are true

- $p/q = p'/q'$.
- $p/q = q'/p'$ and $\mathcal{K}_{f_1} = \mathcal{K}_{f_2}$ is the unknot component.

Definition 2.7. (in the terminology of [5]) A knot $f \in \hat{\mathcal{K}}$ is a connected-sum $f = f_1 \# \cdots \# f_n$ of n knots for $n \geq 2$ if f is isotopic to a knot in the image of $\kappa_n|_{\mathcal{C}_2(n) \times \prod_{i=1}^n \hat{\mathcal{K}}_{f_i}}$ where $f_i \in \hat{\mathcal{K}}$ are non-trivial knots. If a knot is not a connected-sum, and not the unknot, then we call it prime.

Proposition 2.8. (Schubert) [28] If the knots $f_1, f_2, \dots, f_n$ and $g_1, g_2, \dots, g_m$ are prime then $\mathcal{K}_{f_1 \# \cdots \# f_n} = \mathcal{K}_{g_1 \# \cdots \# g_m}$ if and only if $n = m$ and for some permutation σ of $\{1, \dots, n\}$ and all $i \in \{1, 2, \dots, n\}$, $\mathcal{K}_{f_i} = \mathcal{K}_{g_{\sigma(i)}}$.

Schubert's result also follows from basic results on the JSJ-decomposition of the knot complement (see for example [5]).



Definition 2.9. Given $g_0 \in \hat{\mathcal{K}}$, an embedding $g : \sqcup_{i=1}^n S^1 \times D^2 \rightarrow I \times D^2$ disjoint from g_0 , and n non-trivial long knots $f = (f_1, \dots, f_n) \in \hat{\mathcal{K}}^n$ such that

- $g = \sqcup_{i=1}^n g_i$ is the unlink
- $\tilde{g} = \sqcup_{i=0}^n g_i$ has a hyperbolic complement in $\mathbb{R}^3$


define the hyperbolic satellite knot $(\tilde{g}, f)$ to be the composite $R[g, f] \circ g_0$ where $R[g, f]$ is the re-embedding function.


This operation is a special case of what Eisenbud and Neumann call splicing in their book [10].

Proposition 2.10. [6] $\mathcal{K}_{3,1}(\tilde{g}, f) = \mathcal{K}_{3,1}(\tilde{g}', f')$ if and only if:

- $\tilde{g}$ and $\tilde{g}'$ have isotopic images where the isotopy is required to send the g_0 component to the g'_0 component, preserving the orientations of those components.
- Let σ_i be the permutation of $\{1, 2, \dots, n\}$ so that the above isotopy sends g_i to $g'_{\sigma(i)}$, and let $\epsilon_i \in \{0, 1\}$ describe whether or not the isotopy preserves or reverses the orientation of g_i to $g'_{\sigma(i)}$, ie: $\epsilon_i = 0$ if g_i to $g'_{\sigma(i)}$ preserves orientation and $\epsilon_i = 1$ otherwise. Then we require that $\hat{\mathcal{K}}(N^{\epsilon_i} f_i) = \hat{\mathcal{K}}(f'_{\sigma(i)})$. Here $N : \hat{\mathcal{K}} \rightarrow \hat{\mathcal{K}}$ is the involution of $\hat{\mathcal{K}}$ given by reversing a long knot (rotation by π about any axis perpendicular to the long axis).

Example 2.11. Examples of hyperbolic satellite knots and their JSJ-decompositions.





The purpose of Propositions 2.6 2.8 and 2.10 is that it gives a canonical description of $\pi_0 \mathcal{K}_{3,1}$. A non-trivial knot is either a cable, a connect-sum or a hyperbolic satellite knot [5], and these three classes of knots are disjoint. Thus the above three propositions give a bijective correspondence between $\pi_0 \mathcal{K}_{3,1}$ and certain finite rooted trees (JSJ-trees) decorated by various Seifert and hyperbolic links in S^3 , as described above.

3. THE HOMOTOPY TYPE OF $\mathcal{K}_{3,1}$

A detailed description of the homotopy type of $\mathcal{K}_{3,1}$ is given in this section. This description is given in terms of the satellite decomposition of knots as described in Section 2.

As mentioned in the previous section, the spaces $EC(1, D^n)$ admit an action of the operad of little 2-cubes, moreover the subspace $\hat{\mathcal{K}} \subset EC(1, D^2)$ also admits an action of the 2-cubes operad. $\hat{\mathcal{K}}$ is homotopy equivalent to $\mathcal{K}_{3,1}$, thus there is a homotopy equivalence

$$\mathcal{C}_2(X_{\mathcal{K}} \sqcup \{*\}) \rightarrow \mathcal{K}_{3,1}$$

We list the current understanding of the homotopy type of $\mathcal{K}_{3,1}$.

- If f is the unknot then $\mathcal{K}_{3,1}(f)$ is contractible by work of Hatcher [18].
- If f is a p/q -cabling of g then by work of Hatcher [18], there is a homotopy equivalence

$$S^1 \times \mathcal{K}_{3,1}(g) \rightarrow \mathcal{K}_{3,1}(f).$$

- If $f = f_1 \# \cdots \# f_n$ where $\{f_i : i \in \{1, 2, \dots, n\}\}$ are the prime summands of f , then there is a homotopy equivalence

$$\mathcal{C}_2(n) \times_{\Sigma_f} (\mathcal{K}_{3,1}(f_1) \times \cdots \times \mathcal{K}_{3,1}(f_n)) \rightarrow \mathcal{K}_{3,1}(f)$$

as described in [5].

- If a knot f is obtained from knots $\{f_i : i \in \{1, 2, \dots, n\}\}$ by a hyperbolic satellite operation then there is a fibration

$$\prod_{i=1}^n \mathcal{K}_{3,1}(f_i) \rightarrow \mathcal{K}_{3,1}(f) \rightarrow (S^1)^2.$$

Moreover, this fibration can be non-trivial and there exists a finite-sheeted cover of $(S^1)^2$ so that the pull-back of the bundle $\mathcal{K}_{3,1}(f) \rightarrow (S^1)^2$ is the product

$$\left(\prod_{i=1}^n \mathcal{K}_{3,1}(f_i) \right) \times (S^1)^2.$$

Proofs of these statements (will) appear in [17], [18], [5], and [6]. The goal of [6] is to work out the monodromy of the above bundle in the hyperbolic satellite case and is currently incomplete. Since every knot is either trivial, a cable, a hyperbolic satellite or a connected-sum of 2 or more knots, the above gives a description of $\mathcal{K}_{3,1}$ up to a finite-sheeted cover.

Lemma 3.1. *Let f denote a non-trivial long knot. Then there is a surjection*

$$\pi_1(\mathcal{K}_{3,1}(f)) \rightarrow \mathbb{Z}.$$

Proof. A case-by-case proof is given next.

- (1) If the long knot f is a p/q -cabling of a (perhaps trivial) knot g then there is a homotopy equivalence $S^1 \times \mathcal{K}_{3,1}(g) \rightarrow \mathcal{K}_{3,1}(f)$, thus the map $\mathcal{K}_{3,1}(f) \rightarrow S^1$ is projection onto the first factor.
- (2) If $f = f_1 \# \cdots \# f_n$ where $\{f_i : i \in \{1, 2, \dots, n\}\}$ are the prime summands of f , then there is a homotopy equivalence $\mathcal{C}_2(n) \times_{\Sigma_f} (\mathcal{K}_{3,1}(f_1) \times \cdots \times \mathcal{K}_{3,1}(f_n)) \rightarrow \mathcal{K}_{3,1}(f)$. Consider the projection map $\mathcal{K}_{3,1}(f) \rightarrow \mathcal{C}_2(n)/\Sigma_n$. $\pi_1 \mathcal{C}_2(n)/\Sigma_n$ is the braid group on n strands. Since $n \geq 2$, $\pi_1 \mathcal{C}_2(n)/\Sigma_n$ has a rank 1 abelianization, so take the composite $\mathcal{K}_{3,1}(f) \rightarrow \mathcal{C}_2(n)/\Sigma_n \rightarrow S^1$ where $\mathcal{C}_2(n)/\Sigma_n \rightarrow S^1$ is the abelianization map.
- (3) If a knot f is obtained from knots $\{f_i : i \in \{1, 2, \dots, n\}\}$ by a hyperbolic satellite operation then there is a fibration

$$\prod_{i=1}^n \mathcal{K}_{3,1}(f_i) \rightarrow \mathcal{K}_{3,1}(f) \rightarrow (S^1)^2$$

Since the fiber is connected, the map $\pi_1 \mathcal{K}_{3,1}(f) \rightarrow \pi_1 (S^1)^2$ is epic.

Thus in all three cases, there is a surjection of the fundamental group $\mathcal{K}_{3,1}(f)$ to the integers. $\square$

The second statement in Corollary 1.5 follows from Lemma 3.1.

4. ACTIONS ON $\mathcal{K}_{3,1}$

Consider the subgroup of $SO(3)$ which preserves the x -axis $\{(x, 0, 0) \in \mathbb{R}^3\}$ denoted here by $O(2)$. Then $O(2)$ acts on $\mathcal{K}_{3,1}$ by conjugation. The purpose of this section is to describe the $O(2)$ -action on $\mathcal{K}_{3,1}$ for connected-sums and cable knots.

Fix once and for all an element N in $O(2) - SO(2)$. Observe that if $A \in SO(2)$ then $A(\mathcal{K}_{3,1}(f)) = \mathcal{K}_{3,1}(f)$ since SO_2 is connected. However, $N(\mathcal{K}_{3,1}(f))$ could potentially be a different component of $\mathcal{K}_{3,1}$. This happens when the knot f is not invertible (or equivalently, ‘not reversible’). Non-invertible knots were shown to exist by Trotter [31].

Proposition 4.1. ($O(2)$ -action on a cabling) *If $f = c_{p/q}g$ then*

- *By [6], there is an $SO(2)$ -equivariant homotopy equivalence*

$$S^1 \times \mathcal{K}_{3,1}(g) \rightarrow \mathcal{K}_{3,1}(f)$$

where the $SO(2)$ action on $S^1 \times \mathcal{K}_{3,1}(g)$ is given by $A \cdot (v, x) = (A \cdot v, x)$.

- *$N(\mathcal{K}_{3,1}(f)) = \mathcal{K}_{3,1}(f)$ if and only if $N(\mathcal{K}_{3,1}(g)) = \mathcal{K}_{3,1}(g)$ and in this case the homotopy equivalence $S^1 \times \mathcal{K}_{3,1}(g) \rightarrow \mathcal{K}_{3,1}(f)$ is $O(2)$ -equivariant where the $O(2)$ action on $S^1 \times \mathcal{K}_{3,1}(g)$ is given by $A \cdot (v, g) = (A \cdot v, N^{\epsilon(A)} \cdot g)$ where $\epsilon : O(2) \rightarrow \mathbb{Z}_2$ is the epimorphism.*

Proposition 4.2. ($O(2)$ -action on a connected-sum)

- *By [5], if $f = f_1 \# \cdots \# f_n$ where $\{f_i : i \in \{1, 2, \dots, n\}\}$ are the prime summands of f , then there is an $SO(2)$ -equivariant homotopy equivalence*

$$\mathcal{C}_2(n) \times_{\Sigma_f} \prod_{i=1}^n \mathcal{K}_{3,1}(f_i) \rightarrow \mathcal{K}_{3,1}(f)$$

where the $SO(2)$ action on $\mathcal{C}_2(n) \times_{\Sigma_f} \prod_{i=1}^n \mathcal{K}_{3,1}(f_i)$ is given by

$$A \cdot ((F_1, \dots, F_n), (f_1, \dots, f_n)) = ((F_1, \dots, F_n), (A \cdot f_i, \dots, A \cdot f_n))$$

- The $O(2)$ action on $\mathcal{K}_{3,1}$ restricts to $\mathcal{K}_{3,1}(f)$ if and only if there is some involution $\sigma \in S_n$ so that $\mathcal{K}_{3,1}(Rf_i) = \mathcal{K}_{3,1}(f_{\sigma(i)})$ for all $i \in \{1, 2, \dots, n\}$. In this case the map $\mathcal{C}_2(n) \times_{\Sigma_f} \prod_{i=1}^n \mathcal{K}_{3,1}(f_i) \rightarrow \mathcal{K}_{3,1}(f)$ is $O(2)$ -equivariant where the $O(2)$ action on $\mathcal{C}_2(n) \times_{\Sigma_f} \prod_{i=1}^n \mathcal{K}_{3,1}(f_i)$ is given by

$$A \cdot ((F_1, \dots, F_n), (f_1, \dots, f_n)) = ((A \cdot F_{\sigma(1)}, \dots, A \cdot F_{\sigma(n)}), (A \cdot f_{\sigma(1)}, \dots, A \cdot f_{\sigma(n)}))$$

where $A \in O(2) - SO(2)$ and A acts on $\mathcal{C}_2(n)$ by mirror reflection about the y -axis, and acts as usual on the $\mathcal{K}_{3,1}(f_i)$. As before, if $A \in SO(2)$ then

$$A \cdot ((F_1, \dots, F_n), (f_1, \dots, f_n)) = ((F_1, \dots, F_n), (A \cdot f_1, \dots, A \cdot f_n))$$

These propositions are elementary consequences of the works [5] and [18] and appear in [6]. Since the precise homotopy type of a hyperbolic satellite knot is not known, little can be said about the $O(2)$ action on these components. This issue will be addressed in [6].

5. RELATIONS AMONG VARIOUS SPACES

The goal of this section is to compare the homotopy types of the spaces $\mathcal{K}_{n,1}$ and the space of smooth embeddings of S^1 in S^n , $\text{Emb}(S^1, S^n)$. The space of pointed, smooth embeddings $\text{Emb}_*(S^1, S^n)$ will be a useful auxiliary space.

Proposition 5.1. *If $n \geq 3$, there are morphisms of fibrations for which each vertical map is a homotopy equivalence:*

$$\begin{array}{ccccc} SO(n) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{i} & SO(n+1) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{p} & SO(n+1)/SO(n) \\ \downarrow \Theta_n & & \downarrow \Theta_n & & \downarrow 1 \\ \text{Emb}_*(S^1, S^n) & \xrightarrow{j} & \text{Emb}(S^1, S^n) & \longrightarrow & S^n, \end{array}$$

and

$$\begin{array}{ccccc} \mathcal{K}_{n,1} & \xrightarrow{i} & SO(n) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{p} & SO(n)/SO(n-1) \\ \downarrow 1 & & \downarrow \Theta_n & & \downarrow 1 \\ \mathcal{K}_{n,1} & \xrightarrow{j} & \text{Emb}_*(S^1, S^n) & \longrightarrow & S^{n-1}. \end{array}$$

Proof. Consider the maps $\Theta_n : SO(n+1) \times \mathcal{K}_{n,1} \rightarrow \text{Emb}(S^1, S^n)$ obtained from the natural $SO(n+1)$ -action on $\text{Emb}(S^1, S^n)$ together with the natural inclusion $\mathcal{K}_{n,1} \rightarrow \text{Emb}(S^1, S^n)$. Notice that the map Θ_n is $SO(n-1)$ -equivariant, and thus there is an induced map

$$\Theta_n : SO(n+1) \times_{SO(n-1)} \mathcal{K}_{n,1} \rightarrow \text{Emb}(S^1, S^n).$$

Consider the natural fibrations

$$\text{Emb}_*(S^1, S^n) \rightarrow \text{Emb}(S^1, S^n) \rightarrow S^n,$$

and

$$\mathcal{K}_{n,1} \rightarrow \text{Emb}_*(S^1, S^n) \rightarrow S^{n-1}$$

The first map is a fibration by the isotopy extension theorem. Indeed, Palais proved that in general ‘restriction maps’ are locally trivial fiber bundles [26]. The map $\text{Emb}_*(S^1, S^n) \rightarrow S^{n-1}$ is the composition of the restriction map $\text{Emb}_*(S^1, S^n) \rightarrow \text{Emb}_*(U, S^n)$ with the homotopy equivalence $\text{Emb}_*(U, S^n) \rightarrow S^{n-1}$ given by the derivative at $*$ where U is some closed interval neighbourhood of $*$ in S^1 . Thus there is a map of fibrations:

$$\begin{array}{ccccc}
SO(n) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{i} & SO(n+1) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{p} & SO(n+1)/SO(n) \\
\downarrow \Theta_n & & \downarrow \Theta_n & & \downarrow 1 \\
Emb_*(S^1, S^n) & \xrightarrow{j} & Emb(S^1, S^n) & \longrightarrow & S^n
\end{array}$$

as well as

$$\begin{array}{ccccc}
\mathcal{K}_{n,1} & \xrightarrow{i} & SO(n) \times_{SO(n-1)} \mathcal{K}_{n,1} & \xrightarrow{p} & SO(n)/SO(n-1) \\
\downarrow 1 & & \downarrow \Theta_n & & \downarrow 1 \\
\mathcal{K}_{n,1} & \xrightarrow{j} & Emb_*(S^1, S^n) & \longrightarrow & S^{n-1}.
\end{array}$$

The map $\Theta_n : SO(n) \times_{SO(n-1)} \mathcal{K}_{n,1} \rightarrow Emb_*(S^1, S^n)$ is thus a homotopy equivalence. Hence the map

$$\Theta_n : SO(n+1) \times_{SO(n-1)} \mathcal{K}_{n,1} \rightarrow Emb(S^1, S^n)$$

is also a homotopy equivalence. $\square$

Notice that Proposition 5.1 provides a method for comparing the homology of $Emb(S^1, S^n)$ with the homology of $\mathcal{K}_{n,1}$. Next, restrict attention to the special case given by $n = 3$.

Corollary 5.2. *There is a homeomorphism*

$$S^3 \times Emb_*(S^1, S^3) \rightarrow Emb(S^1, S^3).$$

Furthermore, the bundle

$$\mathcal{K}_{3,1} \rightarrow Emb_*(S^1, S^3) \rightarrow S^2$$

is the induced bundle with fiber $\mathcal{K}_{3,1}$ from the bundle

$$SO(2) \rightarrow SO(3) \rightarrow S^2$$

where $SO(2)$ acts on $\mathcal{K}_{3,1}$ in the standard way.

There is a morphism of fibrations which is a fibrewise homotopy equivalence

$$\begin{array}{ccccc}
\mathcal{K}_{3,1} & \xrightarrow{i} & SO(3) \times_{SO(n-1)} \mathcal{K}_{3,1} & \xrightarrow{p} & SO(3)/SO(2) \\
\downarrow 1 & & \downarrow \Theta_n & & \downarrow 1 \\
\mathcal{K}_{3,1} & \xrightarrow{j} & Emb_*(S^1, S^3) & \longrightarrow & S^2.
\end{array}$$

Proof. Let A be a pointed space with $map_*(A, G)$ the space of pointed maps from A to a topological group G . Then the space of continuous functions $map(A, G)$ is homeomorphic to a product $G \times map_*(A, G)$ for any topological group. This homeomorphism specializes to a homeomorphism

$$Emb_*(S^1, S^3) \rightarrow Emb(S^1, S^3) \rightarrow S^3.$$

The remainder of this corollary is given in Proposition 5.1. $\square$

Notice that Proposition 1.4 follows at once from 5.2, and 5.1. In addition, there is an analogous decomposition for $Emb(S^1, S^7)$, but that will not be addressed.

Observe that Proposition 5.1 immediately generalizes to a proposition about the embedding spaces $Emb(S^k, S^n)$. Let $\mathcal{K}_{n,k}$ be the space of embeddings of $\mathbb{R}^k$ in $\mathbb{R}^n$ which agree with the standard inclusion $(x_1, \dots, x_k) \mapsto (x_1, \dots, x_k, 0, \dots, 0)$ outside of the unit ball. Let $EC(k, D^{n-k})$ be as in [5]. Thus there is a fibration $\Omega^k SO(n-k) \rightarrow EC(k, D^{n-k}) \rightarrow \mathcal{K}_{n,k}$.

Proposition 5.3. *There is a commutative diagram of 6 fibrations:*

$$\begin{array}{ccccc}
 \mathcal{K}_{n,k} & \longrightarrow & \text{Emb}(S^k, S^n) & \longrightarrow & V_{n+1,k+1} \\
 \uparrow & & \uparrow & & \uparrow \\
 \text{EC}(k, D^{n-k}) & \longrightarrow & \text{Emb}(S^k \times D^{n-k}, S^n) & \longrightarrow & SO_{n+1} \\
 \uparrow & & \uparrow & & \uparrow \\
 \Omega^k SO(n-k) & \longrightarrow & SO(n-k) \times \Omega^k SO(n-k) & \longrightarrow & SO(n-k)
 \end{array}$$

Moreover, the top horizontal fibration $\mathcal{K}_{n,k} \rightarrow \text{Emb}(S^k, S^n) \rightarrow V_{n+1,k+1}$ is fiber-homotopy equivalent to the fibration $\mathcal{K}_{n,k} \rightarrow SO(n+1) \times_{SO(n-k)} \mathcal{K}_{n,k} \rightarrow V_{n+1,k+1} \equiv SO(n+1)/SO(n-k)$. Both of the other two horizontal fibrations are trivial product bundles, the rightmost fibration is the canonical quotient $V_{n+1,k+1} \equiv SO(n+1)/SO(n-k)$ while the other two are ‘Palais’ restriction fibrations.

Proof. Consider the inclusion $\mathcal{K}_{n,k} \rightarrow \text{Emb}(S^k, S^n)$ given by taking the 1-point compactification of an element in $\mathcal{K}_{n,k}$ and identifying the one point compactification of $\mathbb{R}^n$ with S^n . Since $SO(n+1)$ acts on S^n , $SO(n+1)$ there is a map $\mathcal{K}_{n,k} \times SO(n+1) \rightarrow \text{Emb}(S^k, S^n)$. Observe this map factors to a map $\mathcal{K}_{n,k} \times_{SO(n-1)} SO(n+1) \rightarrow \text{Emb}(S^k, S^n)$. Thus there is a commuting diagram

$$\begin{array}{ccccc}
 \mathcal{K}_{n,k} & \longrightarrow & \text{Emb}(S^k, S^n) & \longrightarrow & \text{Emb}(D^k, S^n) \\
 \uparrow & & \uparrow & & \uparrow \\
 \mathcal{K}_{n,k} & \longrightarrow & \mathcal{K}_{n,k} \times_{SO(n-1)} SO(n+1) & \longrightarrow & V_{n+1,k+1}
 \end{array}$$

The two vertical maps on the extreme ends of the diagram are homotopy equivalences by design, thus the center map is a homotopy equivalence by the 5-lemma. $\square$

6. ON THE HOMOLOGY OF $C_2(X \amalg \{*\})$

The purpose of this section is to recall the homology of

$$C_2(X \amalg \{*\})$$

for X not necessarily path-connected. These results will then be combined with Theorem 1.2 to obtain Theorem 1.3. The space X is assumed to be compactly generated, and weak Hausdorff as a topological space [9]; the base-point $\{*\}$ is non-degenerate by construction.

Formal constructions are given next for which $\mathbb{F}$ is a field, and all modules are assumed to be vector spaces over $\mathbb{F}$. Let V denote a graded vector space which splits as a direct sum

$$V = V_+ \oplus V_-$$

for which V_+ consists of the elements concentrated in even degrees, and V_- consists of the elements concentrated in odd degrees. Let

$$\sigma(V)$$

denote the ‘‘algebraic suspension’’ of V . That is $\sigma(V)$ is the module V with all degrees raised by one. In addition,

$$\sigma^n(V) = \sigma(\sigma^{n-1}(V)).$$

The ‘‘algebraic desuspension’’ of V denoted

$$\sigma^{-1}(V)$$

is defined by $\sigma(\sigma^{-1}(V)) = V$.

Next consider the free Lie algebra

$$L[\sigma(V)]$$

and, if $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$, the free restricted Lie algebra over $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ denoted

$$L^{(p)}[\sigma(V)].$$

In this last case, consider the natural inclusion

$$j : L[\sigma(V)] \rightarrow L^{(p)}[\sigma(V)]$$

with co-kernel denoted $W^p[\sigma(V)]$ (for which $L[\sigma(V)]$ is the free Lie algebra defined over the field $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$). The definition of a restricted Lie algebra is given in Jacobson's book "Lie Algebras", and may be regarded as the module of primitive elements in the tensor algebra $T[\sigma(V)]$ defined over the field $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$. Notice that $\sigma^{-1}(\sigma V) = V$, but that $\sigma^{-1}L[\sigma(V)]$ is not isomorphic to $L[V]$ in general.

Consider the symmetric algebra $S[V]$ defined as follows:

- (1) If $\mathbb{F} = \mathbb{Q}$, then $S[V] = E[V_-] \otimes \mathbb{F}[V_+]$ where $E[\cdot]$ denotes an exterior algebra, and $\mathbb{F}[\cdot]$ denotes a polynomial algebra.
- (2) If $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$, then $S[V] = \mathbb{F}[V]$, a polynomial algebra.
- (3) If $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ for an odd prime p , then $S[V] = E[V_-] \otimes \mathbb{F}[V_+]$ where $E[\cdot]$ denotes an exterior algebra, and $\mathbb{F}[\cdot]$ denotes a polynomial algebra.

The homology of $C_2(X \amalg \{*\})$ is described below in three distinct cases

- (1) $\mathbb{F} = \mathbb{Q}$,
- (2) $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$, and
- (3) $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ for an odd prime p .

Consider the first case with $\mathbb{F} = \mathbb{Q}$. Let $V = H_*(X; \mathbb{Q})$, and form the symmetric algebra $S[\sigma^{-1}L[\sigma(H_*(X; \mathbb{Q}))]]$. By [9], there is an isomorphism of Hopf algebras

$$S[\sigma^{-1}L[\sigma(H_*(X; \mathbb{Q}))]] \rightarrow H_*(C_2(X \amalg \{*\}); \mathbb{Q})$$

with co-product determined by that of $H_*(X; \mathbb{Q})$.

The analogous theorem for $\mathbb{F}_2 = \mathbb{Z}/2\mathbb{Z}$ is given as follows. Let $V = H_*(X; \mathbb{F}_2)$, and form the symmetric algebra

$$S[\sigma^{-1}L^{(2)}[\sigma(H_*(X; \mathbb{F}_2))]].$$

By [9], there is an isomorphism of Hopf algebras

$$S[\sigma^{-1}L^{(2)}[\sigma(H_*(X; \mathbb{F}_2))]] \rightarrow H_*(C_2(X \amalg \{*\}); \mathbb{F}_2)$$

with co-product determined by that of $H_*(X; \mathbb{F}_2)$. The role of the restriction in a restricted Lie algebra over $\mathbb{F}_2$ is to create the Araki-Kudo-Dyer-Lashof operation, the operation which sends an element $\sigma(v)$ to $Q_1(\sigma(v))$.

The result for odd primes p with $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ is given as follows. Let $V = H_*(X; \mathbb{F}_p)$, and form the symmetric algebra

$$S[\sigma^{-1}L^{(p)}[\sigma(V)] \oplus \sigma^{-2}W^p[\sigma(V)]].$$

By [9], there is an isomorphism of Hopf algebras

$$S[\sigma^{-1}L^{(p)}[\sigma(V)] \oplus \sigma^{-2}W^p[\sigma(V)]] \rightarrow H_*(C_2(X \amalg \{*\}); \mathbb{F}_p)$$

with coproduct determined by that of $H_*(X; \mathbb{F}_p)$.

Assuming familiarity with the Araki-Kudo-Dyer-Lashof operations, some remarks are given next. The role of the restriction in a restricted Lie algebra over $\mathbb{F}_p$ is to create the Araki-Kudo-Dyer-Lashof operation, an operation which sends an element $\sigma(v)$ to $Q_{p-1}(\sigma(v))$ precisely in case $\sigma(v)$ for which $\sigma(v)$ is of odd degree in the case

that p is an odd prime. The role of $\sigma^{-2}W^p[\sigma(V)]$ is the contribution of the first Bockstein of $Q_{p-1}(\sigma(v))$ in case $\sigma(v)$ is of odd degree, and p is an odd prime.

7. ON THE HOMOLOGY OF $\mathcal{K}_{3,1}$

Recall the homotopy equivalence of Theorem 1.2,

$$C(\mathbb{R}^2, X_{\mathcal{K}} \amalg \{*\}) \rightarrow \mathcal{K}_{3,1}$$

Thus there are isomorphisms of Hopf algebras

- (1) $S[\sigma^{-1}L[\sigma(V)]] \rightarrow H_*(\mathcal{K}_{3,1}; \mathbb{Q})$ for $V = H_*(X_{\mathcal{K}}; \mathbb{Q})$,
- (2) $S[\sigma^{-1}L^2[\sigma(V)]] \rightarrow H_*(\mathcal{K}_{3,1}; \mathbb{Z}/2\mathbb{Z})$ for $V = H_*(X_{\mathcal{K}}; \mathbb{Z}/2\mathbb{Z})$, and
- (3) $S[\sigma^{-1}L^{(p)}[\sigma(V)] \oplus \sigma^{-2}W^p[\sigma(V_-)]] \rightarrow H_*(\mathcal{K}_{3,1}; \mathbb{Z}/p\mathbb{Z})$ for odd primes p .

These isomorphisms, given in terms of the homology of $X_{\mathcal{K}}$, are also filtered [9] with these filtrations specializing to an identification of the homology of each path-component $\mathcal{K}_{3,1}(f)$. There is one ambiguity arising from the fact that the homology of the components corresponding to hyperbolic satellite operations is not given in a closed form. Furthermore, these isomorphisms indicate how the homology of $\mathcal{K}_{3,1}$ is determined by the homology of $X_{\mathcal{K}}$, and are listed in detail in [9].

More precisely, the homology of the space $C_2(X \amalg \{*\})$ is filtered in [9] with the associated filtration quotients in homology given by $H_*(\mathcal{C}_2(n) \times_{\Sigma_n} (X)^n)$. This suffices.

The homology of the components of knots arising from hyperbolic satellite operations is not given in a closed form here as described in section 3. Thus, the homology of $X_{\mathcal{K}}$ has not been determined completely. Further information concerning the homology of $X_{\mathcal{K}}$ is given in section 9.

Thus the above isomorphisms give the homology of $\mathcal{K}_{3,1}$ with field coefficients. The first, and second parts of Theorem 1.2 follow. The proof of Theorem 1.2 will be concluded in section 8 in which higher torsion is constructed.

Notice that the space $C_2(X \amalg \{*\})$ is naturally a disjoint union of X with another space. Thus, there is a natural direct sum decomposition of graded vector spaces

$$\bar{H}_*(X; \mathbb{F}) \oplus \Gamma(X; \mathbb{F}) \rightarrow H_*(C_2(X \amalg \{*\}); \mathbb{F})$$

for a choice of graded vector space

$$\Gamma(X; \mathbb{F})$$

which is functor of $H_*(X; \mathbb{F})$.

The construction $\Gamma(X; \mathbb{F})$ is used in section 9 to describe the homology of the subspace of $\mathcal{K}_{3,1}$ generated by torus knots, as well as the operations of connected sums, cablings, and the action of the little two-cubes.

The structure considered here on the homology of the space of long knots in $\mathbb{R}^3$ also appears in several other contexts. In characteristic zero, the homology of the space of long knots also satisfies structures known as a Batalin-Vilkovisky algebra, or Gerstenhaber algebra, and appears within the context of vertex operator algebras, as well as some physically inspired mathematics. Thus the rational homology of the space of long knots in $\mathbb{R}^3$ is a free Batalin-Vilkovisky algebra.

8. HIGHER p -TORSION IN THE INTEGER HOMOLOGY OF $\mathcal{K}_{3,1}$

One way in which higher order p -torsion in the homology of $\mathcal{K}_{3,1}$ arises is summarized next. The way in which little cubes $\mathcal{C}_2(n)$ are related to configuration spaces $Conf(\mathbb{R}^2, n)$ is as follows. There are maps $\mathcal{C}_2(n) \rightarrow Conf(\mathbb{R}^2, n)$ which are both homotopy equivalences, and equivariant with respect to the action of the symmetric group Σ_n [25]. Thus it suffices to exhibit higher torsion in the integer homology of

$Conf(\mathbb{R}^2, n) \times_{\Sigma_n} X^n$ for certain choices of spaces $X = \mathcal{K}_{3,1}(f)$. Since the construction $Conf(\mathbb{R}^2, n) \times_{\Sigma_n} X^n$ occurs numerous times below, it is convenient to define

$$E_n(X) = Conf(\mathbb{R}^2, n) \times_{\Sigma_n} X^n.$$

Given a prime long knot f , consider the path-component in $\mathcal{K}_{3,1}$ with

$$g = \#_L f$$

where $\#_L f$ denotes the connected-sum of L copies of the same knot f . Thus there are homotopy equivalences

$$K_{3,1}(g) = K_{3,1}(\#_L f) \rightarrow Conf(\mathbb{R}^2, L) \times_{\Sigma_L} K_{3,1}(f)^L = E_L(f).$$

First consider p -torsion of order exactly p obtained from the equivariant cohomology of $Conf(\mathbb{R}^2, p)$ as constructed in [9].

Proposition 8.1. *Let Y denote any connected CW-complex.*

- (1) *If $H_{2t-1}(Y; \mathbb{F}_p)$ is non-zero, then $\mathbb{Z}/p\mathbb{Z}$ is a direct summand of*

$$H_{2pt-2}(Conf(\mathbb{R}^2, p) \times_{\Sigma_p} Y^p; \mathbb{Z}).$$

- (2) *If $\mathbb{Z}/p^s\mathbb{Z}$ is a direct summand of $H_{2t-1}(Y; \mathbb{Z})$, then*

$$H_{2tp^r-1}(E_{p^r}(Y); \mathbb{Z}) = H_{2tp^r-1}(Conf(\mathbb{R}^2, p^r) \times_{\Sigma_{p^r}} Y^{p^r}; \mathbb{Z})$$

has a $\mathbb{Z}/p^{s+q}\mathbb{Z}$ -summand.

- (3) *There is a homotopy equivalence*

$$E_L(K_{3,1}(f)) \rightarrow K_{3,1}(\#_L f).$$

- (4) *Thus if $\mathcal{K}_{3,1}(f)$ has any non-trivial mod- p homology in degree $2t-1$, then $\mathbb{Z}/p\mathbb{Z}$ is a direct summand of*

$$H_{2pt-2}(E_p(K_{3,1}(f)); \mathbb{Z}).$$

Hence

$$H_{2pt-2}(K_{3,1}(\#_p f); \mathbb{Z}) = \mathbb{Z}/p\mathbb{Z} \oplus A$$

for some abelian group A .

- (5) *Furthermore, if $\mathbb{Z}/p^s\mathbb{Z}$ is a direct summand of $H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{Z})$, then*

$$H_{2tp^r-1}(E_{p^r}(\mathcal{K}_{3,1}(f)); \mathbb{Z}) = H_{2tp^r-1}(K_{3,1}(\#_{p^r} f); \mathbb{Z}) = \mathbb{Z}/p^{s+r}\mathbb{Z} \oplus A$$

for some abelian group A .

Assume that

$$H_j(K_{3,1}(f); \mathbb{Z}) = \mathbb{Z}/p^s\mathbb{Z} \oplus A$$

for some abelian group A . Label this $\mathbb{Z}/p^s\mathbb{Z}$ -summand (ambiguously) by

$$\langle f, j, \mathbb{Z}/p^s\mathbb{Z} \rangle.$$

Example 8.2. By [18], or section 3, if f is a non-trivial torus knot, then $\mathcal{K}_{3,1}(f)$ has the homotopy type of a circle. A direct application of Proposition 8.1 gives that

$$H_{2p-2}(Conf(\mathbb{R}^2, p) \times_{\Sigma_p} K_{3,1}(f)^p; \mathbb{Z}) = H_{2p-2}(K_{3,1}(\#_p f); \mathbb{Z}) = \langle \#_p f, 2p-2, \mathbb{Z}/p\mathbb{Z} \rangle \oplus A$$

for some abelian group A .

Next, consider the prime knot h given by a p/q -cabling of f .

Example 8.3. The examples here arise by an iterated cabling construction as follows. Since the symbol “ p ” is used in two distinct ways in the examples below, the knot h denoted $c_{p/q}(f)$ will also be denoted $c_{\alpha/\beta}(f)$, and thus

$$h = c_{p/q}(f) = c_{\alpha/\beta}(f).$$

There is a homotopy equivalence

$$\mathcal{K}_{3,1}(h) = \mathcal{K}_{3,1}(c_{\alpha/\beta}(f)) \rightarrow \mathcal{K}_{3,1}(f) \times S^1$$

by section 3, and [18].

Next, consider an m -fold iterated cabling h_m of f defined by

- $h_1 = c_{\alpha/\beta}(f)$,
- $h_{i+1} = c_{\alpha/\beta}(h_i)$, with
- $h_m = c_{\alpha/\beta}^m(f)$.

Then there are homotopy equivalences

$$\mathcal{K}_{3,1}(h_m) = \mathcal{K}_{3,1}(c_{\alpha/\beta}^m(f)) \rightarrow \mathcal{K}_{3,1}(f) \times (S^1)^m.$$

Assume that

$$H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{F}_p) = \langle f, 2t-1, \mathbb{Z}/p\mathbb{Z} \rangle \oplus A$$

for some abelian group A . Then the integer homology of $\mathcal{K}_{3,1}(h_m) = \mathcal{K}_{3,1}(f) \times (S^1)^m$ has a summand denoted (ambiguously) by

$$\langle f, 2t-1, \mathbb{Z}/p\mathbb{Z} \rangle \otimes H_*((S^1)^m; \mathbb{Z}).$$

Thus if $m \geq 1$, there are elements of order p in both odd as well as even degrees in the integer homology of $\mathcal{K}_{3,1}(h_m)$. If m is “large”, then there are many elements of order exactly p which are of both odd, and even degree. This fussiness concerning parity of degrees has consequences for higher torsion in homology.

The above remarks give examples of long knots with torsion of order exactly p concentrated in odd degrees for the integer homology of their path-components. One choice of f is a p/q -torus knot. The next proposition shows that p -torsion of order exactly p in the homology of $\mathcal{K}_{3,1}(g)$ gives rise higher p^s -torsion in the homology of other components related components as follows.

Recall that Example 8.3 provides instances of prime knots f such that

$$H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{Z}) = \langle f, 2t-1, \mathbb{Z}/p\mathbb{Z} \rangle \oplus A$$

for some abelian group A . Consider the long knot g given by the p^s -fold connected sum

$$g = f_1 \# f_2 \# \cdots \# f_{p^s} = \#_{p^s} f$$

for which $f_i = f$ as used in the next Proposition.

Proposition 8.4. *Let f denote a prime long knot such that*

$$H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{Z}) = \langle f, 2t-1, \mathbb{Z}/p\mathbb{Z} \rangle \oplus A$$

for some abelian group A . Let

$$g = f \# \cdots \# f = \#_{p^s} f.$$

Then

$$H_{2tp^s-1}(\mathcal{K}_{3,1}(g); \mathbb{Z}) = \langle \#_{p^s} f, 2tp^s-1, \mathbb{Z}/p^{s+1}\mathbb{Z} \rangle \oplus A$$

for some abelian group A .

Proof. Assume that the integer homology of $\mathcal{K}_{3,1}(f)$ has a non-trivial $\mathbb{Z}/p\mathbb{Z}$ summand in degree $2t - 1$ as guaranteed by example 8.3. Thus, in the mod- p reduction of the integer homology of $\mathcal{K}_{3,1}(f)$, there are classes x of degree $2t - 1$, the mod- p reduction of a class of order p , as well as a class y in degree $2t$ which corresponds to the contribution forced by x in the “Tor” term in the classical universal coefficient Theorem.

Since $H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{Z})$ has a $\mathbb{Z}/p\mathbb{Z}$ -summand, there are elements in the mod- p homology of $\mathcal{K}_{3,1}(f)$ with

- (1) x in $H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{Z}/p\mathbb{Z})$,
- (2) y in $H_{2t-1}(\mathcal{K}_{3,1}(f); \mathbb{Z}/p\mathbb{Z})$, and
- (3) the first Bockstein of y is x ,

$$\beta_1(y) = x.$$

A classical computation of the Bockstein spectral sequence gives that the $(s + 1)$ -st Bockstein is defined with

$$\beta_{s+1}(y^{p^s}) = x \cdot y^{-1+p^s} + I$$

in case $s \geq 1$ for which I denotes the indeterminacy of this operation. The Proposition follows as these classes survive in the Bockstein spectral sequence for $C(\mathbb{R}^2, X_{\mathcal{K}} \amalg \{*\})$. $\square$

Two concrete examples are listed next.

Example 8.5. (1) Let f denote a non-trivial torus knot with

$$H_1\mathcal{K}_{3,1}(f) = \mathbb{Z}.$$

Then the long knot

$$\#_p f = f \# \cdots \# f$$

satisfies the property that

$$\mathcal{K}_{3,1}(\#_p f) = E_p(\mathcal{K}_{3,1}(f))$$

with

$$H_{2p-2}(\mathcal{K}_{3,1}(\#_p f); \mathbb{Z}) = \langle \#_p f, 2p - 2, \mathbb{Z}/p^1\mathbb{Z} \rangle \oplus A$$

for some abelian group A .

- (2) Cable $\#_p f$ with a non-trivial α/β -torus knot, denoted $c_{\alpha/\beta}(\#_p f)$. There are homotopy equivalences

$$\mathcal{K}_{3,1}(c_{\alpha/\beta}(\#_p f)) \rightarrow \mathcal{K}_{3,1}(\#_p f) \times S^1$$

with the property that

$$H_{2p-1}(\mathcal{K}_{3,1}(c_{\alpha/\beta}(\#_p f)); \mathbb{Z}) = \langle c_{\alpha/\beta}(\#_p f), 2p - 1, \mathbb{Z}/p\mathbb{Z} \rangle \oplus A$$

for some abelian group A .

- (3) The p^s -fold connected sum of $c_{\alpha/\beta}(\#_p f)$,

$$\#_{p^s} c_{\alpha/\beta}(\#_p f)$$

has the property that

$$H_{2p^{s+1}-1}(\mathcal{K}_{3,1}(\#_{p^s} c_{\alpha/\beta}(\#_p f)); \mathbb{Z}) = \langle \#_{p^s} c_{\alpha/\beta}(\#_p f), 2p^{s+1} - 1, \mathbb{Z}/p^{s+1}\mathbb{Z} \rangle \oplus A$$

for some abelian group A .

The third part of Theorem 1.3 follows, thus concluding the proof.

Other ways in which higher torsion is created arises from properties of certain Lie elements in the Poisson algebra structure here. These Lie elements are analogous to the so-called Engel elements which are central to the solution of the restricted Burnside problem, and are given by

$$\{\cdots \{f, g\} f\} f \cdots f = ad^{p^s-1}(f)(g).$$

These same elements appear in work of J. C. Moore, and J. A. Neisendorfer as well as one of the current authors concerning higher torsion in the homotopy groups of mod- p^m Moore spaces. However, this connection will not be addressed here in order to preserve some sanity of the authors.

9. ON THE SUBSPACE GENERATED BY TORUS KNOTS

The purpose of this section is to describe the subspace $\mathcal{TK}_{3,1}$ of $\mathcal{K}_{3,1}$ generated by torus knots, as well as the operations of connected sums, cablings, and the action of the little two-cubes. An alternate description of the space $\mathcal{TK}_{3,1}$ is that it consists of precisely those long knots whose complements have JSJ-decompositions containing only Seifert fibred manifolds.

First define the space

$$\mathcal{T} = \coprod_{1 < p < q, (p,q)=1} \mathcal{K}_{3,1}(f(p, q))$$

where $f(p, q)$ denotes a p, q -torus knot. Thus $\mathcal{K}_{3,1}(f(p, q))$ has the homotopy type of S^1 . Consider the James construction

$$J(\mathcal{T} \amalg \{*\}) = \coprod_{0 \leq n} \mathcal{T}^n$$

with $\mathcal{T}^0 = \{*\}$, the base-point. Write

$$J_{\mathcal{K}} = \coprod_{1 \leq n} \mathcal{T}^n$$

with

$$J(\mathcal{T} \amalg \{*\}) = J_{\mathcal{K}} \amalg \{*\}.$$

Spaces Y_n are specified inductively in terms of $J_{\mathcal{K}}$ as follows.

- * $Y_0 = C_2(J_{\mathcal{K}} \amalg \{*\})$, and
- * $Y_{n+1} = \{([C_2(Y_n)] - Y_n) \times J_{\mathcal{K}}\} \amalg Y_n$.

Notice that Y_n is naturally a subspace of Y_{n+1} , and that all of these may be regarded as subspaces of $\mathcal{K}_{3,1}$ in the following way: There are induced maps

$$\mathcal{K}_{3,1} \times J(\mathcal{T} \amalg \{*\}) \rightarrow \mathcal{K}_{3,1}$$

induced by cabling.

Define

$$\mathcal{TK}_{3,1} = \cup_{n \geq 0} Y_n.$$

Notice that $\mathcal{TK}_{3,1}$ is the subspace of $\mathcal{K}_{3,1}$ which contains the path-components of p/q -torus knots, and which is closed under the operations of cabling, and sums. That is, there are induced maps $\mathcal{K}_{3,1} \times J(\mathcal{T} \amalg \{*\}) \rightarrow \mathcal{K}_{3,1}$ induced by cabling. There is an induced inclusion $\mathcal{TK}_{3,1} \rightarrow \mathcal{K}_{3,1}$.

The remainder of this section gives features of the homology of $\mathcal{TK}_{3,1}$. First notice that homology commutes with inductive co-limits, and so there are isomorphisms

$$\varinjlim H_*(Y_n) \rightarrow H_*(\varinjlim Y_n) \rightarrow H_*(\mathcal{TK}_{3,1}).$$

Recall the construction $\Gamma(X)$ as given in section 6.

To describe the homology of Y_n , restrict to field coefficients. $\mathbb{F}$, and recall the natural splitting of graded vectors spaces

$$\bar{H}_*(X; \mathbb{F}) \oplus \Gamma(X; \mathbb{F}) \rightarrow H_*(C_2(X; \mathbb{F}))$$

for a choice of graded vector space

$$\Gamma(X; \mathbb{F})$$

which is functor of $H_*(X; \mathbb{F})$ as listed in section 6. Thus $H_*(Y_{n+1})$ is given in terms of the construction $\Gamma(X; \mathbb{F})$ in case $X = Y_n$.

Proposition 9.1. *The natural map $H_*(Y_n) \rightarrow H_*(Y_{n+1})$ is a split monomorphism.*

Notice that the homology of $\mathcal{TK}_{3,1}$ exhibits a fractal-like behavior reflecting the geometry in Budney's theorem [5], and iterations of the constructions $\Gamma(X)$ as given in section 6. Namely, this homological behavior arises by first considering $Y_1 = C_2(J_K)$ together with the homology $H_*(C_2(X \amalg \{*\}); \mathbb{F})$ as given in section 6 as follows where

$$V = H_*(X; \mathbb{F}).$$

- (1) $S[\sigma^{-1}L[\sigma(V)]]$ if the characteristic of $\mathbb{F}$ is 0,
- (2) $S[\sigma^{-1}L^{(2)}[\sigma(V)]]$ if $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$, and
- (3) $S[\sigma^{-1}L^{(p)}[\sigma(V)] \oplus \sigma^{-2}W^p[\sigma(V)]]$ if $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ for odd primes p .

10. STEENROD-THOM REPRESENTABILITY OF HOMOLOGY CLASSES

Start with manifolds M which represent the mod-2 homology of the spaces $K_{3,1}(\mathcal{P})$ for prime knots $\mathcal{P}$. Consider the manifolds obtained by taking

- (1) finite products of M with finite products of circles,
- (2) the wreath product construction $S^1 \times_{\mathbb{Z}/2\mathbb{Z}} X^2$ for X one of the manifolds listed in 1 above, and
- (3) finite iterations of the constructions in 1, and 2 above by taking finite products of wreath product constructions, wreath product constructions of the resulting manifolds, taking finite products again, and continue.

Theorem 10.1. *Any class in the mod-2 homology of $\mathcal{K}_{3,1}$ is represented by the manifolds in 1-3 above.*

Proof. Suppose that the manifolds M_i represent the mod-2 homology of a space X . Then

- (1) products $M_1 \times M_2$, and
- (2) $S^1 \times_{\Sigma_2} (M_i)^2$

represent the mod-2 homology of $S^1 \times_{\Sigma_2} X^2$. Since the mod-2 homology of $Conf(\mathbb{R}^2, s) \times_{\Sigma_s} X^a$ is represented by iterates of these two constructions, the result follows. $\square$

11. CLOSED KNOTS AND HOMOLOGY

The purpose of this section is to use results of the earlier sections to give information about the space $\text{Emb}(S^1, S^3)$. Recall the homeomorphism

$$S^3 \times \text{Emb}_*(S^1, S^3) \rightarrow \text{Emb}(S^1, S^3)$$

for which $\text{Emb}_*(S^1, S^3)$ denotes the pointed embeddings. Thus $H_*(\text{Emb}(S^1, S^3)) = H_*(S^3) \otimes H_*(\text{Emb}_*(S^1, S^3))$ by Proposition 5.2.

Information giving the structure of the bundle $\mathcal{K}_{3,1} \rightarrow \text{Emb}_*(S^1, S^3) \rightarrow S^2$ was worked out earlier. That structure is used to provide information concerning $H_*(\text{Emb}_*(S^1, S^3))$ by a Mayer-Vietoris argument.

Let D_1 and D_2 be two discs in S^2 whose union is S^2 and whose intersection is S^1 . Let A_1 and A_2 be the preimages of D_1 and D_2 under the projection map $\text{Emb}_*(S^1, S^3) \rightarrow S^2$, then both A_1 and A_2 are homeomorphic to $\mathcal{K}_{3,1} \times D^2$. Consider the Mayer-Vietoris sequence for $\text{Emb}_*(S^1, S^3) = A_1 \cup A_2$ where $A_1 \cap A_2$ is homeomorphic to $S^1 \times \mathcal{K}_{3,1}$.

Identify $A_1 \equiv D^2 \times \mathcal{K}_{3,1}$ and $A_2 \equiv D^2 \times \mathcal{K}_{3,1}$ then the gluing map from $\partial A_1 = S^1 \times \mathcal{K}_{3,1} \rightarrow \partial A_2 = S^1 \times \mathcal{K}_{3,1}$ is the map $S^1 \times \mathcal{K}_{3,1} \ni (t, x) \mapsto (t, t^2 \cdot x) \in S^1 \times \mathcal{K}_{3,1}$. Thus the Mayer-Vietoris sequence has the form:

$$\cdots \rightarrow H_*(S^1 \times \mathcal{K}_{3,1}) \rightarrow H_*(D^2 \times \mathcal{K}_{3,1}) \oplus H_*(D^2 \times \mathcal{K}_{3,1}) \rightarrow H_*(\text{Emb}_*(S^1, S^3)) \rightarrow \cdots$$

where $H_*(S^1 \times \mathcal{K}_{3,1})$ is identified with $H_*\mathcal{K}_{3,1} \oplus H_{*-1}\mathcal{K}_{3,1}$, and $H_*(D^2 \times \mathcal{K}_{3,1})$ is identified with $H_*(\mathcal{K}_{3,1})$. Then the map

$H_n \mathcal{K}_{3,1} \oplus H_{n-1} \mathcal{K}_{3,1} \rightarrow H_n(\mathcal{K}_{3,1}) \oplus H_n(\mathcal{K}_{3,1})$ is given by the 2×2 matrix $\begin{pmatrix} I & 0 \\ I & 2\kappa_n \end{pmatrix}$ where $\mu : SO(2) \times \mathcal{K}_{3,1} \rightarrow \mathcal{K}_{3,1}$ is the $SO(2)$ -action on $\mathcal{K}_{3,1}$ and $\kappa_n : H_{n-1} \mathcal{K}_{3,1} \rightarrow H_n \mathcal{K}_{3,1}$ where $\kappa_n(x) = \mu_*(SO(2) \times x)$

Corollary 11.1. *There is a short exact sequence*

$$0 \rightarrow \text{coker}(2\kappa_n) \rightarrow H_n \text{Emb}_*(S^1, S^3) \rightarrow \text{ker}(2\kappa_{n-1}) \rightarrow 0$$

Example 11.2. *Some examples of κ_* .*

- If f is a torus knot $\mathcal{K}_{3,1}(f) \simeq S^1$ we identify $H_* \mathcal{K}_{3,1}(f)$ with $(\mathbb{Z}, \mathbb{Z}, 0, \dots)$. $SO(2) \times S^1 \rightarrow S^1$ is left multiplication, so the induced map on homology $\kappa_* : H_0 S^1 \rightarrow H_1 S^1$ is an isomorphism, in all other degrees κ_* is zero.
- If f is a connected sum of 2 distinct torus knots $\mathcal{K}_{3,1}(f) \simeq (S^1)^3$, $H_* \mathcal{K}_{3,1}(f) \simeq \Lambda^* \mathbb{Z}^3$ and κ_* can be identified with the wedge product $x \mapsto x \wedge (w_1 + w_2)$ where $\Lambda^1 \mathbb{Z}^3 \simeq \mathbb{Z}^3$ has basis w_1, w_2, w_3 .
- If f is a connected sum of 2 distinct hyperbolic knots $\mathcal{K}_{3,1}(f) \simeq (S^1)^5$, $H_* \mathcal{K}_{3,1}(f) \simeq \Lambda^* \mathbb{Z}^5$ and κ_* can be identified with the wedge product $x \mapsto x \wedge (w_1 + w_2)$ where $\Lambda^1 \mathbb{Z}^5 \simeq \mathbb{Z}^5$ has basis w_1, w_2, w_3, w_4, w_5 .

Corollary 11.3. *A knot $f : S^1 \rightarrow S^3$ is the unknot if and only if its component in $\text{Emb}(S^1, S^3)$ contains no 2-torsion in its homology.*

Proof. If f is the unknot, the component of f has the homotopy type of $V_{4,2} = S^3 \times S^2$ which has no torsion in its homology.

If f is non-trivial, then first consider its long knot component, $\mathcal{K}_{3,1}(f)$. This has a $\mathbb{Z}$ embedded in its fundamental group, embedded as the 2π rotation around the long axis [15]. We call the embedding $\mathbb{Z} \rightarrow \pi_1 \mathcal{K}_{3,1}(f)$ the Gramain map. In [6] it's shown that there is a map $\mathcal{K}_{3,1}(f) \rightarrow S^1$ which when composed with the Gramain map is not null-homotopic. The proof is analogous to the proof of Lemma 3.1, only a little more involved as the maps constructed in the proof of Lemma 3.1 do not suffice.

This allows us to deduce that $H_1 \mathcal{K}_{3,1}(f)$ contains a copy of the integers, generated by the Gramain element. Thus since the image of κ_1 is generated by twice the gramain element, $\text{coker}(2\kappa_1)$ must contain 2-torsion. $\square$

12. PROBLEMS

The purpose of this section is to list problems which arise naturally from the work above.

(1): The total homology of each path-component $\mathcal{K}_{3,1}(f)$ gives an invariant of the long knot f . The first homology group of each path-component of $\mathcal{K}_{3,1}$, or $\text{Emb}(S^1, S^3)$ distinguishes the unknot by 1.5. Neither the fundamental group nor the homology of $\mathcal{K}_{3,1}(f)$ is strong enough to distinguish all knots. For example, for all hyperbolic knots f , $\mathcal{K}_{3,1}(f) \simeq (S^1)^2$. Similarly, for all cables of torus knots g , $\mathcal{K}_{3,1}(g) \simeq (S^1)^2$.

Describe the equivalence relation on long knots given by $f \sim g$ if $\mathcal{K}_{3,1}(f) \simeq \mathcal{K}_{3,1}(g)$. This relation is some coarse ‘complexity’ invariant of the JSJ-tree of a knot complement and seems quite different than most traditional invariants of knots.

(2): A similar but cruder problem is to describe the knot types for which the total homology of the path-components are isomorphic.

(3): Which path components $\mathcal{K}_{3,1}(f)$ of prime knots f have torsion in their first homology group? Do any of these have p -torsion in their first homology group for p an odd prime? In some hyperbolic satellite cases, there is 2-torsion in the first homology group.

(4): Interpret the rational co-homology of $\mathcal{K}_{3,1}$ in terms of iterated integrals in the sense of Kohno-Kontsevich-Chen.

(5): Compare the Vassiliev invariants of braids as studied by T. Kohno [19], and the Lie algebra obtained from the descending central series for the fundamental groups of the spaces $\mathcal{C}_2(n) \times \mathcal{K}_{3,1}(f_1) \times \cdots \times \mathcal{K}_{3,1}(f_n)$ as well as the induced invariants for

$\mathcal{C}_2(n) \times_{\Sigma_f} (\mathcal{K}_{3,1}(f_1) \times \cdots \times \mathcal{K}_{3,1}(f_n))$. Use these constructions to analyze the behavior of $H_*(\mathcal{K}_{n,1})$. This problem approach is to consider the structure of each path component rather than trying to distinguish the components themselves.

(6): A natural connection between the space of long knots, and the mod-2 Steenrod algebra arises from the “group completion” of $\mathcal{K}_{3,1}$ given $\Omega B(\mathcal{K}_{3,1})$. Notice that the collapse map $X_{\mathcal{K}} \rightarrow \{*\}$ induces a map $p : C(\mathbb{R}^2, X_{\mathcal{K}} \amalg \{*\}) \rightarrow C(\mathbb{R}^2, S^0)$, and there is an induced map $p : C(\mathbb{R}^2, X_{\mathcal{K}} \amalg \{*\}) \rightarrow \mathbb{Z} \times BO$ induced by the regular representation bundle. Thus there are maps

$$\Omega B(\mathcal{K}_{3,1}) \rightarrow \Omega_0^2 S^2 \rightarrow BO$$

with composite denoted $\phi : \Omega B(\mathcal{K}_{3,1}) \rightarrow \mathbb{Z} \times BO$. The Thom spectrum of ϕ is a wedge of Eilenberg-Mac Lane spectra $H\mathbb{Z}/2$, and thus the mod-2 co-homology of the Thom spectrum $M\Omega B(\mathcal{K}_{3,1})$ is free over the mod-2 Steenrod algebra. Interpret the Steenrod operations in terms of knots.

(7): The unknot component $\mathcal{K}_{3,1}(f)$ is contractible. A natural question to ask is, does some ‘physical’ potential function exist on $\mathcal{K}_{3,1}$ having only one critical point in $\mathcal{K}_{3,1}(f)$? For an arbitrary knot f a finite-dimensional model-spaces for $\mathcal{K}_{3,1}(f)$ is given recursively by the work [6]. Can one find ‘natural’ potential functions on $\mathcal{K}_{3,1}$ lacking critical points outside of these model spaces? Similarly, there are the analogous questions for $\text{Emb}(S^1, S^3)$. These questions were raised by Freedman, He and Wang [13]. A recent preprint of Abrams, Cantarella and Fu [1] shows that certain natural potential functions on $\text{Emb}(S^1, \mathbb{R}^n)$ are minimized by geometric circles, thus giving some hope. If on the other hand, no such natural potential functions exist, one could hope for natural ‘Morse’ potential functions. In that case, the results of this paper and [6] could be used to predict the number and type of critical points.

REFERENCES

- [1] A. Abrams, J. Cantarella, J.H. Fu, *Circles minimize most knot energies*, submitted.
- [2] D. Altshuler, and L. Freidel, *On universal Vassiliev invariants*, Comm. Math. Phys., **170**(1995), 41-62.
- [3] D. Bar-Natan, *On the Vassiliev knot invariants*, Topology, **34**(1995), 423-472.
- [4] R. Bott, and C. Taubes, *On the self-linking of knots*, J. Math. Phys., **35**(1994), 5247-5287.
- [5] R. Budney, *Little cubes and long knots*, Topology, (to appear) (-), -.
- [6] R. Budney, *The topology of spaces of knots in dimension 3*, in preparation.
- [7] A. Cattaneo, P. Cotta-Ramusino, and R. Longoni, *Configuration spaces and Vassiliev invariants in any dimension*, Algebraic, and Geometric Topology, **2**(2002), 949-1000.
- [8] V. Chari, and A. Pressley, *A Guide to Quantum Groups*, Cambridge University Press, 1994.
- [9] F. R. Cohen, T. J. Lada and J. P. May, *The homology of iterated loop spaces*, Lecture Notes in Math., vol. **533**. Springer-Verlag (1976).
- [10] D. Eisenbud, W. Neumann, *Three-dimensional link theory and invariants of plane curve singularities*, Ann. Math. Stud. **110**.
- [11] M. Falk, R. Randell, *The lower central series of a fiber-type arrangement*, Invent. Math., **82** (1985), 77-88.
- [12] E. Fadell and L. Neuwirth, *Configuration spaces*, Math. Scand., **10** (1962), 111-118.
- [13] M.H. Freedman, Z.-X. He, Z. Wang, *Mobius energy of knots and unknots*, Ann. of Math., **139** (1994) 1-50.
- [14] T. Goodwillie, and M. Weiss, *Embeddings from the point of view of immersion theory II*, Geometric Topology, **3** (1999), 103-118.
- [15] A. Gramain, *Sur le groupe fondamental de l'espace des noeuds*, Ann. Inst. Fourier, Grenoble. **27,3** (1977), 29-44.
- [16] A. Haefliger, M. Hirsch, *On the existence and classification of differentiable embeddings*, Topology **2** (1963), 129-135.

- [17] A. Hatcher, *Homeomorphisms of sufficiently-large P^2 -irreducible 3-manifolds*, Topology, **15** (1976)
- [18] A. Hatcher, *Topological Moduli Spaces of Knots*, math.GT/9909095.
- [19] T. Kohno, *Vassiliev invariants and de Rham complex on the space of knots*, Contemp. Math. **179**(1994), 123-138.
- [20] M. Kontsevich, *Feynman diagrams and low dimensional topology*, Progress in Mathematics, vol. 2, 1990, 120.
- [21] P. Lambrechts, I. Volic, *The rational homotopy type of spaces of knots in codimension > 2* , preprint.
- [22] C. Lescop, *On configuration space integrals for links*, Invariants of knots and 3-manifolds (Kyoto, 2001), 183–199 (electronic), Geom. Topol. Monogr., **4**, Geom. Topol. Publ., Coventry, 2002.
- [23] G. I. Lehrer, *The cohomology of a regular semisimple variety*, J. Alg. 199(1998), 666-689.
- [24] G. I. Lehrer, G. B. Segal, *Homology stability for classical semisimple varieties*, Math. Zeit., (), –.
- [25] J. P. May, *The Geometry of Iterated Loop Spaces*, Lect. Notes in Math., **268**, Springer-Verlag, Berlin, 1972.
- [26] R. Palais, *Local triviality of the restriction map for embeddings*, Comment. Math. Helv. **34** (1960) 305–312.
- [27] M. Polyak, O. Viro, *On the Casson knot invariant*, Knots in Hellas '98, vol. 3(Delphi). J. Knot Theory Ramifications, **10** (2001), no. 5, 711–738.
- [28] H. Schubert, *Die eindeutige Zerlegbarkeit eines Knoten in Primknoten*, Sitzungsber. Akad. Wiss. Heidelberg, math.-nat. Kl., **3:57-167**. (1949)
- [29] D. Sinha, *The topology of spaces of knots*, math.AT/0202287].
- [30] D. Tamaki, *A dual Rothenberg-Steenrod spectral sequence*, Topology, **33**(4)(1994), 631-662.
- [31] H. F. Trotter *Non-invertible knots exist*. Topology **2** (1963) 275–280.
- [32] V. Tourtchine *On the other side of the bialgebra of chord diagrams*, arXiv QA/0411436 v2 9 Feb 2005.
- [33] V. Vassiliev, *Complements of discriminants of smooth maps: topology and applications*, Translations of Mathematical Monographs, **98**, American Mathematical Society, Providence, RI, 1992.
- [34] I. Volic, *Finite-type knot invariants and calculus of functors*. Brown University. (2003)
- [35] W. Wu, *On the isotopy of C^r -manifolds of dimension n in euclidean $(2n + 1)$ -space*. Sci. Record (N.S.) **2** (1958), 271–275.
- [36] H. Whitney, *Topological properties of differentiable manifolds*, Bull. Amer. Math. Soc., **43** (1937), 785–805.
- [37] H. Whitney, *The self-intersections of a smooth n -manifold in $2n$ -space*, Ann. of Math. (2) **45**, 220–246 (1944).
- [38] H. Whitney, *Differentiable Manifolds*, Ann. of Math. (2) **37** (1936), 645–680.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403-1221 U.S.A.
E-mail address: rybu@math.uoregon.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627 U.S.A.
E-mail address: cohf@math.rochester.edu