

# $\mathbb{Z}_n$ -manifolds in 4-dimensional graph-manifolds

A.Mozgova

Laboratoire d'analyse non linéaire et géométrie, Université d'Avignon, 33, rue Louis Pasteur, 84000 Avignon, France.

mozgova@univ-avignon.fr

## Abstract

A standard fact about two incompressible surfaces in an irreducible 3-manifold is that one can move one of them by isotopy so that their intersection becomes  $\pi_1$ -injective. By extending it on the maps of some 3-dimensional  $\mathbb{Z}_n$ -manifolds into 4-manifolds, we prove that any homotopy equivalence of 4-dimensional graph-manifolds with reduced graph-structures is homotopic to a diffeomorphism preserving the structures.

*Keywords:* graph-manifold,  $\pi_1$ -injective  $\mathbb{Z}_n$ -submanifold.

## 1 $\mathbb{Z}_n$ -submanifolds of 4-manifolds

A  $\mathbb{Z}_n$ -manifold of dimension  $k$  is an object  $Z$  obtained from a compact oriented  $k$ -manifold with boundary by identifying (with respect of the orientation)  $n$  isomorphic parts of boundary in such a way that if a point of the boundary component participate in the identification, then all the points of this component do. Every point of  $Z$  has a neighborhood isomorphic either to a  $k$ -cell, or to a  $k$ -semi-cell, or to a  $n$ -page book of  $k$ -cells. The identified parts form *the singular set* of  $Z$  denoted  $Z_s$ , and the closure of their complement is its *regular set* denoted  $Z'$ . Non-identified boundary components form the boundary  $\partial Z$ .


For example, by identifying boundaries of three oriented surfaces each one having one boundary component, one obtain a 2-dimensional  $\mathbb{Z}_3$ -manifold without boundary. Mapping cylinder of  $n$ -fold covering  $S^1 \rightarrow S^1$  gives an example of 2-dimensional  $\mathbb{Z}_n$ -manifold with boundary; it appears in 3-dimensional Seifert manifolds as the preimage of an arc in the base orbifold going from the projection of a singular fiber to the boundary.

A standard fact about two incompressible surfaces in an irreducible 3-manifold is that one can move one of them by isotopy so that their intersection becomes  $\pi_1$ -injective. Under natural homotopic assumption, this remains true for the map of 3-manifold and 3-dimensional submanifold of 4-manifold (Proposition 2.B.2 of [5] and Proposition 1 of [4]). The following lemma extends it on the maps of some 3-dimensional  $\mathbb{Z}_n$ -manifolds into 4-manifolds.

To fix the notations, let  $c : S^1 \rightarrow S^1$  be  $n$ -fold covering. Its mapping cylinder is a 2-dimensional  $\mathbb{Z}_n$ -manifolds  $Map^n(S^1, S^1) = (S^1 \times [0, 1]) / (S^1 \times \{1\}) = c(S^1 \times \{1\})$ .

Let  $Z = S^1 \times Map^n(S^1, S^1)$ , it is a 3-dimensional  $\mathbb{Z}_n$ -manifold with boundary  $\partial Z = S^1 \times \partial Map^n(S^1, S^1)$ , singular set  $Z_s = S^1 \times Map^n(S^1, S^1)_s$  and regular set  $Z' = S^1 \times (S^1 \times [0; 1])$ .

**Lemma 1.** *Let  $W$  be a compact smooth oriented 4-manifold with  $\pi_2(W) = \pi_3(W) = 0$  and  $M$  be a compact oriented  $\pi_1$ -injective 3-submanifold with  $\pi_2(M) = 0$ . Let  $f : (Z, \partial Z) \rightarrow (W, W \setminus M)$  be a  $\pi_1$ -injective map.*



*Then one can move  $f$  by homotopy (that is constant on  $\partial Z$ ) so that  $f(Z_s) \cap M = \emptyset$  and each connected component of  $F = f^{-1}(M)$  is a  $\pi_1$ -injective torus in  $M$ .*

$Z$ -dimensional  $Z_n$ -manifold whose embedding  $(F, F_s) \rightarrow (Z \setminus \partial Z, Z_s)$  induces an embedding of regular sets  $F' \rightarrow Z'$ . As  $M$  is closed and  $Z'$  is compact,  $F'$  has a finite number of connected components ([1], corollary 17.2(IV)).

In  $Z = S^1 \times \text{Map}^n(S^1, S^1)$  consider the subspace  $G = \{0\} \times \text{Map}^n(S^1, S^1)$  where  $0 \in S^1$ . We have  $(G, G_s) \subset (Z, Z_s)$  and  $G' \subset Z'$ . So  $F'$  (which is union of surfaces with boundaries) and  $G'$  (which is an annulus) are embedded in  $Z' = S^1 \times S^1 \times I$ . Denote the boundary component of  $G'$  lying  $\partial Z$  by  $\partial_0 G'$  and the other one by  $\partial_1 G'$ .




Figure 1: Regular parts, the covering  $\partial_1 Z' \rightarrow Z_s$ , and  $Z_s \cup G \cup F$

**Step 1: elimination of trivial circles of  $\partial F'$ .** Denote the boundary component of  $Z'$  giving  $\partial Z$  by  $\partial_0 Z'$  and the other one (participating in the identification) by  $\partial_1 Z'$ . Suppose that there is a circle of  $\partial F'$  that is trivial in  $\partial_1 Z'$ . Then its projection on  $Z_s$  is trivial, too. As its projection is embedded in  $Z_s \simeq T^2$  (because  $\partial F'$  is the preimage of  $F_s$  which is embedded), it bounds an embedded disk there. Now we can proceed as in Proposition 1 of [4]: take a map of 2-disk in  $M$  bounding the same loop. As  $\pi_2(W) = 0$ , the union of these two disks bounds a map of 3-disk  $\alpha : D^3 \rightarrow W$ , which can be separated from  $M$ . Denote by  $N$  a small book neighborhood of  $D^2 \subset Z_s$  in  $Z$ . The map  $f$  will not be changed on  $\overline{Z \setminus N}$ ; on  $D^2$  the homotopy of  $f$  will be the pushing across  $\alpha(D^3)$ , and  $N \setminus D$  will serve to rely the new map on  $D^2$  and the old map on  $Z \setminus N$ , doing it separately on each leaf of the book.




Figure 2: Changing of  $f$  on  $D^2 \subset Z_s$ .

This homotopy of  $f$  will eliminate the trivial circle from  $F'$ :




Figure 3: Surgery on a trivial loop of  $\partial F'$ .

**Step 2.** Reduction of  $F$  to a union of closed surfaces and annuli. Now all the circles  $\partial F' \subset \partial_1 Z' \simeq T^2$  are non-trivial and embedded, hence, parallel. They are either parallel to  $\partial_1 G'$ , or not, which corresponds respectively to  $\partial F' \cap \partial_1 G' = \emptyset$  or  $\partial F' \cap \partial_1 G' \neq \emptyset$ . Take one connected component of  $F'$ , denote it still by  $F'$ .

We will separate two cases:  $F' \cap G' = \emptyset$  or  $F' \cap G' \neq \emptyset$ .

Case 1:  $F' \cap G' = \emptyset$ . It implies  $F' \subset \overline{(Z' \setminus G')} \simeq S^1 \times I \times I$ . Note that  $\overline{(Z' \setminus G')} \subset Z'$  is  $\pi_1$ -injective but  $F' \subset Z'$  is not, so neither is  $F' \subset \overline{(Z' \setminus G')}$ .

In this case  $(F', \partial F') \rightarrow (S^1 \times I \times I, \partial(S^1 \times I \times I))$ ,  $\partial F'$  going to a curve parallel to the generator of  $\pi_1(S^1 \times I \times I) = H_1(S^1 \times I \times I)$ . We need to find an embedded disk in  $S^1 \times I \times I$  for the trivializing loop. As  $\partial F'$  bounds  $F' \subset S^1 \times I \times I$ ,  $\partial F' \sim 0$  in  $H^1(S^1 \times I \times I, \mathbb{Z}_2)$ . As all the curves of  $\partial F'$  are  $\neq 0$  and parallel in  $\partial(S^1 \times I \times I)$ , they are in even number: say  $m$  with one orientation,  $m$  with the opposite one.  $(F', \partial F') \rightarrow (S^1 \times I \times I, \partial(S^1 \times I \times I))$ , so each  $\partial F'$  is sent to a generator of  $\pi_1(S^1 \times I \times I)$ . Take an embedded disk  $D = \{0\} \times I \times I \subset (S^1 \times I \times I)$ . We have then  $(D, \partial D) \rightarrow (S^1 \times I \times I, \partial(S^1 \times I \times I))$  and  $D \cap F'$  is the union of circles and annuli. Take the innermost circle of  $D \cap F'$ ; if it is  $\sim 0$  in  $F'$ , we have a disk for the surgery, otherwise move  $D$  by isotopy to eliminate it, and so on. After treating all the circles either we have a disk for the surgery, or there are only arcs in  $D \cap F'$ . Take two components of  $\partial F$  corresponding to the arc whose ends are neighbouring in  $\partial D$ , denote them by  $a$  and  $a^{-1}$ . Then  $F'$  can be presented as  $F' = A \cup_{\gamma} F''$  where  $A$  is an annulus with a hole,  $\partial A = a \cup a^{-1} \cup \gamma$  and  $F''$  is the remaining part.




Figure 4:  $F'$  as union of an annulus with a hole and  $F''$ .

Denote by  $A'$  the annulus in  $\partial(S^1 \times I \times I)$  lying between  $a$  and  $a^{-1}$ . Note that  $\gamma \sim aa^{-1} \sim 0$  in  $\pi_1(S^1 \times I \times I)$  (but  $\gamma \neq 0$  in  $F'$  as  $F' \neq S^1 \times I$ ). The arcs  $A \cap D$  and  $A' \cap D$  bound a disk in  $D$ , whose interior does not intersect  $F'$ .




Figure 5:  $D \cap F'$  in  $D$  and  $D$  in  $S^1 \times I \times I$ .

Push  $A \cup A'$  along its normal bundle toward the inside of this disk, living  $\gamma$  unchanged, and do the surgery of the pushed part (which is a torus with a hole) on the interior disk. We obtain a disk  $\tilde{D}$  embedded in  $S^1 \times I \times I$  such that  $\partial \tilde{D} = \gamma$  and  $\tilde{D} \cap F' = \gamma$ . Hence we can move  $f$  by

of an annulus and  $F' \cap D'$ . Doing this for every pair of components of  $\partial F'$  corresponding to an arc with neighboring (in  $\partial D$ ) ends, we will reduce  $F'$  to a union of closed surfaces (which will be treated as in the closed manifold case) and annuli.

Case 2:  $F' \cap G' \neq \emptyset$  is a 1-dim manifold (embedded in both  $F'$  and  $G'$ ); recall that  $G'$  was chosen such that  $(F_s \cap G_s = \emptyset \Leftrightarrow \partial F' \cap \partial_1 G' = \emptyset)$ .

Case 2.1:  $F' \cap G' \neq \emptyset$  but  $F' \cap \partial_1 G' = \emptyset$ . Then  $\partial F'$  is a union of circles parallel to  $\partial_1 G'$ ; and  $F' \cap G'$  is a closed 1-manifold. Circles of  $F' \cap G'$  that are trivial in  $\pi_1(F')$ , can be eliminated by moving  $G'$  (without moving  $\partial_1 G'$ ). So, if all the circles of  $F' \cap G'$  are trivial in  $\pi_1(F')$ , we are back to the Case 1. If there is a circle of  $F' \cap G'$  that is non-trivial in  $\pi_1(F')$ , it is either parallel to a component of  $\partial F'$  or not.

- If there exists a circle not parallel to a component of  $\partial F'$ , then it is embedded in  $G'$  and not parallel to its boundary, so it bounds a disk in  $G'$ . We can do the surgery on this disk (choosing the innermost one).

- If all circles of  $F' \cap G'$  are parallel to a component of  $\partial F'$ , then cutting  $Z'$  along  $G'$ , we'll have some number of  $\pi_1$ -injective annuli embedded in  $S^1 \times I \times I$ , boundary going to boundary, hence isotopic to annuli in  $\partial(S^1 \times I \times I)$  by an isotopy that is trivial on the boundary. Use these isotopies in  $Z'$  to move  $G'$  in order to separate it from  $F'$ , and we are again in the settings of the Case 1.

Case 2.2:  $F' \cap G' \neq \emptyset$  and  $\partial F' \cap \partial_1 G' \neq \emptyset$ . If the components of  $\partial F'$  are homotopic to  $\partial_1 G'$ , move  $G'$  by isotopy to separate  $\partial_1 G'$  from  $\partial F'$ , and we are in the previous case. If the curves of  $\partial F'$  are not homotopic to  $\partial_1 G'$ , choose an embedded curve  $\alpha \subset \partial_1 Z'$ , parallel to the curves  $\partial F'$  (different from them) and make  $\alpha \times I \subset Z'$  be the new  $G'$ . Then, we are back to Case 2.1.

**Step 3: the image of the singular set  $Z_s$  can be separated from  $M$ .** Now every connected component of  $F'$  is a torus or an annulus. Let us show that all the annuli can be eliminated. To simplify the notations,  $F$  will stand for  $F'$  without tori-components.

Fix a decomposition of  $Map^n(S^1, S^1)$  by a wedge of intervals as follows:




Figure 6: Decomposition of  $Map^n(S^1, S^1)$  on sheets.

After multiplying by  $S^1$ , it gives a wedge of annuli  $\bigvee S_i$  (identified along one boundary component) embedded in  $Z$  and decomposing  $Z$  into sheets  $I \times I \times S^1$ , in which the corresponding parts of  $F$  are manifolds. One generator of  $\pi_1(Z_s)$  is given by the singular circle of  $Map^n(S^1, S^1)$ . The loop  $(\bigvee S_i) \cap Z_s$  corresponds to the second generator, and can be chosen not to be parallel to  $F_s$ . Then every connected component of  $F$  intersects a decomposing annulus  $S_i$  by arcs. Take an arc coming from a component  $F_1 \subset F'$ , and suppose it is the innermost one in  $S_i$  (i.e. its union with an arc from  $\partial S_i$  bounds a disk in  $S_i$  that does not contain other arcs from  $F \cap S_i$ ); denote it by  $\alpha \subset F_1 \cap S_1$ . Then the arcs  $F_1 \cap S_j$  are the innermost ones in  $S_j \forall j \neq i$ . The product of  $I$  with the union of the corresponding disks in  $S_1$  and  $S_2$  (see Fig. 7 below)




Figure 7: Disks in the sheets.

allows to change  $f$  by homotopy so that the new image of the singular set  $Z_s$  coincide with the image of  $I \times (F_1 \cap S_1)$ . Hence by pushing off along the normal bundle of  $M$ ,  $\alpha \times I$  can be

$S_1$  and  $S_2$  can be pushed away from  $M$ ). Taking the product of the union of disks in  $S_k$  and  $S_{k+1}$  by  $I$  and doing the same homotopy for the sheet lying between  $S_k$  and  $S_{k+1}$ , one will eliminate all the annuli of  $F$ .

□

## 2 Application to 4-dim graph-manifolds

### 2.1 Seifert manifolds

Following [6, 7], we say that an orientable 4-manifold  $S$  with boundary is a *Seifert manifold* if it has the structure of a fibered orbifold  $\pi : S \rightarrow B$  over a 2-orbifold with generic fiber  $T^2$ , and  $S$  and  $\partial S$  are non-singular as orbifolds. Note that  $B$  as orbifold has no boundary, but the underlying surface of  $B$  does.

#### Local picture.

Any point  $b \in B$  has a neighborhood of type  $D = D^2/G$  where  $G$  is a finite subgroup of  $\mathcal{O}(2)$  corresponding to the stabilizer of  $p$ . Then  $\pi^{-1}(D) = (T^2 \times D^2)/G$  where the action of  $G$  is free and is a lifting of an action of  $G$  on  $D^2$  (that is  $\pi|_{\pi^{-1}(D)} : (T^2 \times D^2)/G \rightarrow D^2/G$  is induced by the canonical projection  $T^2 \times D^2 \rightarrow D^2$ ).

Case 0:  $G = \{1\}$ .

In that case  $p$  is non-singular and  $\pi^{-1}(p) = T^2$  is a regular fiber. The preimage of an arc in  $B$  that joins  $p$  with  $\partial B$  is  $T^2 \times I$ .


Case 1:  $G = \mathbb{Z}_m = \langle g \mid g^m = 1 \rangle$  and the action is given by  $g(x, y, z) = (x - a/m, y - b/m, ze^{\frac{2\pi i}{m}})$ , where  $(x, y) \in T^2$ ,  $z \in D^2 = \{z \in \mathbb{C} \mid |z| \leq 1\}$ , and  $m, a, b$  being mutually relatively prime. Then  $p$  is a cone point of angle  $2\pi/m$  and  $\pi^{-1}(p) = T^2$ . The covering (regular fiber)  $\rightarrow$  (singular fiber) corresponds to the subgroup  $a\mathbb{Z} \oplus b\mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$ . As  $(a, b) = 1$ , we can choose a basis in the  $\pi_1$  of the regular fiber such that the covering corresponds to the subgroup  $\mathbb{Z} \oplus n\mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$ . Then the covering  $T^2 \times D \rightarrow (T^2 \times D^2)/G$  splits as

$$\begin{array}{ccc} S^1 & \times & (S^1 \times D^2) \\ \text{id} \downarrow & & \downarrow \text{n-fold covering} \\ S^1 & \tilde{\times} & (S^1 \times D^2) \end{array}$$

(the bundle  $S^1 \tilde{\times} (S^1 \times D^2)$  being trivial because  $(T^2 \times D^2)/G$  is orientable). That means that  $(T^2 \times D^2)/G = S^1 \times (3\text{-dimensional model})$ . So in this case, the preimage of an arc in  $B$  that joins  $p$  with  $\partial B$  is the  $\mathbb{Z}_n$ -manifold  $S^1 \times \text{Map}^n(S^1, S^1)$ .


Case 2:  $G = \mathbb{Z}_2 = \langle \tau \mid \tau^2 = 1 \rangle$  and the action is given by  $\tau \cdot (x, y, z) = (x + 1/2, -y, \bar{z})$ . Then  $\pi^{-1}(p) = K^2$ ,  $p$  lies on a reflector circle and  $\pi^{-1}(D)$  is a twisted  $D^2$ -bundle over  $K^2$ . The preimage of an arc in  $B$  that joins the point  $p$  with  $\partial B$  is in this case a manifold  $K^2 \tilde{\times} I$ .

Case 3:  $G = D_{2m} = \langle \tau, g \mid \tau^2 = g^m = 1, \tau g \tau^{-1} = g^{-1} \rangle$  and the action is given by  $\tau \cdot (x, y, z) = (x + 1/2, -y, \bar{z})$  and  $g \cdot (x, y, z) = (x, y - b/m, ze^{\frac{2\pi i}{m}})$ . Then  $p$  is a corner reflector of angle  $\pi/m$  and  $\pi^{-1}(p) = K^2$  which is  $m$ -fold covered by  $K^2$  over a regular point of the reflector circle.



The preimage of an arc in  $B$  that joins  $p$  with  $\partial B$  is the mapping cylinder of a covering  $T^2 \rightarrow K^2$  which is a 2-fold covering over the loop reversing the orientation of  $K^2$  and a  $m$ -fold covering over the other loop. This is not a  $\mathbb{Z}_k$ -manifold, but one can join  $p$  with  $\partial B$  by 2 consecutive arcs  $\alpha$  and  $\beta$ , the first one lying on the reflector circle and not containing another corner reflector, the second one joining the other end of  $\alpha$  with  $\partial B$ . Then  $\pi^{-1}(\alpha) = S^1 \tilde{\times} \text{Map}^n(S^1, S^1)$  and  $\pi^{-1}(\beta) = I \tilde{\times} K^2$ .

As in dim 3, on the fundamental group level the singular fibers create roots of the regular fiber.



$S = M \cup M_S$ , where  $M$  is a  $T^2$ -bundle over a surface and  $M_S$  is its Seifert-part. Note that  $0 \rightarrow \pi_1(M) \rightarrow \pi_1(S)$ .

**Theorem** ([9], [8], [6], [7].) *Let  $S, S'$  be 4-dimensional closed orientable Seifert bundles with hyperbolic bases. Then  $\pi_1(S) = \pi_1(S')$  if and only if there is a fiber-preserving diffeomorphism between  $S$  and  $S'$ .*

By taking the doubles, the previous theorem implies that if two Seifert manifolds with boundaries  $(S, \partial S)$  and  $(S', \partial S')$  over hyperbolic 2-orbifolds are homotopy equivalent rel boundary then  $S$  and  $S'$  are diffeomorphic.

## 2.2 4-dimensional graph-manifolds

A block is a Seifert bundle (with boundary) over a hyperbolic 2-orbifold. A graph-manifold structure on a compact closed oriented 4-manifold is a decomposition as a union of blocks, glued by diffeomorphisms of the boundary. Note that the boundary of a block has the structure of a  $T^2$ -bundle over a circle. A graph-manifold structure is *reduced* if none of the glueing maps are isotopic to fiber-preserving maps of  $T^2$ -bundles. Any graph-structure give rise to a reduced one by forming blocks glued by bundle maps into larger blocks. Like in the non-singular case, 4-dimensional graph-manifolds are aspherical, their Euler characteristic is 0 (because the blocks are finitely covered by  $T^2$ -bundles over hyperbolic surfaces, hence have  $\chi = 0$ , and the glueings are made along 3-manifolds), and can be smoothed.

**Theorem 1.** *Any homotopy equivalence of closed oriented 4-manifolds with reduced graph-structures is homotopic to a diffeomorphism preserving the structures.*

We will say that a  $\pi_1$ -injective map between the blocks  $f : S \rightarrow S'$  is *fiber covering* if in the fundamental groups it sends the (normal) fiber subgroup of  $\pi_1(S)$  into the (normal) fiber subgroup of  $\pi_1(S')$ . To extend the proof of the non-singular case ([4]), one has to show that any  $\pi_1$ -injective map  $f : W = \cup W_i \rightarrow W' = \cup W'_k$  of graph-manifolds with reduced graph-structures is homotopic to  $\bigcup f_i$ , where each  $f_i : (W_i, \partial W_i) \rightarrow (W'_j, \partial W'_j)$  is a fiber covering map. The missing step is the following

**Proposition 1.** *Let  $S$  be a block and  $W$  be a 4-dim graph-manifold with reduced graph-structure. Then any  $\pi_1$ -injective map  $f : S \rightarrow W$  is homotopic to a map into one block of  $W$ .*

*Proof.* In the base of  $\pi : S \rightarrow B$ , take the wedge of circle on which  $B$  retracts. Join its core by arcs with every cone point and every reflector circle. As  $S$  retracts on

$$S' = \pi^{-1}(\text{wedge of circles} + \text{wedge of arcs} + \text{reflector circles}),$$

we have to show that  $f|_{S'}$  is homotopic to a map into one bloc of  $W$ .

**Step 1:**  $f|_{\pi^{-1}(\text{wedge of circles})}$  is homotopic to a map in one block of  $W$ . To make the reasonings as in the singular case, one only needs to show that up to homotopy rel boundary, any  $\pi_1$ -injective map  $g : (T^2 \times I, \partial(T^2 \times I)) \rightarrow (S, \partial S)$  is either fiber-covering or is a map into  $\partial S$ . Note that  $S$  is finitely covered by  $\tilde{S}$  which is a  $T^2$ -bundle over a hyperbolic surface (with boundary). As  $g$  sends boundary to boundary,  $Im g_* \subset \pi_1(\tilde{S}) \subset \pi_1(S)$ , hence  $g$  lifts to a map  $\tilde{g} : (T^2 \times I, \partial) \rightarrow (\tilde{S}, \partial\tilde{S})$ , to which Lemma 2 ([4]) applies.

**Step 2:**  $f|_{\pi^{-1}(\text{wedge of arcs} + \text{reflector circles})}$  is homotopic to a map in one block of  $W$ . Denote by  $M$  the block of  $W$  in which  $f(\pi^{-1}(\text{wedge of circles}))$  lies and denote  $p$  the point of  $B$  which is the common core of wedge of circles and wedge of arcs. Then  $f|_{\pi^{-1}(p)}$  is a covering of the regular fiber of  $M$ .

**Step 2.1:** arcs to the cone points. The preimage of such an arc is  $Z = S^1 \times Map^n(S^1, S^1)$ , and  $f|_Z$  sends  $\partial Z$  into  $M$ , hence  $f(\partial Z)$  does not intersect the decomposing submanifolds of  $W$ .

and every component of  $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$  is a torus  $\pi_1$ -injectively embedded in  $Z = S^1 \times S^1 \times I$ . Hence  $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$  is  $\sqcup T^2$  that are parallel to  $\partial Z$  and hence are sent by  $f$  on the covering of regular fiber of  $M$ . Take the first  $T_1^2$  from  $\sqcup T^2$  counting from  $\partial Z$ , say it comes from  $M_\varphi \subset M \cap M'$ . If it exists, denote the next torus from  $\sqcup T^2$  by  $T_2^2$ . The restriction of  $f$  on  $T^2 \times I \subset Z$  lying between  $T_1^2$  and  $T_2^2$  is a  $\pi_1$ -injective map  $f_1 : (T^2 \times I, \partial) \rightarrow (M', \partial M')$ . As  $f(T_i^2)$  is a covering of the regular fiber of  $M$  and the graph-structure is reduced,  $f(T_i^2)$  is not a covering of the regular fiber of the neighbouring block  $M'$ , hence  $f(T_2^2) \subset M_\varphi$  and  $f_1$  is homotopic rel boundary to a map into  $M_\varphi$ . Continuing like this, we change  $f|_Z$  so that  $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$  contains at most one manifold component. If there are none, we are done. If there is one such component, we have a  $\pi_1$ -injective map  $f : (Z, \partial Z, Z_s) \rightarrow (M \cup M', M, M')$ .




Figure 8:  $f(S^1 \times \text{Map}^n(S^1, S^1))$  in the blocs  $M$  and  $M'$

As the torus  $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$  is parallel to  $\partial Z$ , we have a  $\pi_1$ -injective map  $f' : (Z, \partial Z) \rightarrow (M', \partial M')$ , such that  $f'(\partial Z)$  is not fiber-covering.

Case 1:  $f'$  lifts to a map into  $\tilde{M}'$ . If the torus  $\tilde{f}'(\partial Z)$  corresponds to a  $(\mathbb{Z} \oplus \mathbb{Z})$ -subgroup  $\langle \alpha^n, \beta \rangle$  in  $\pi_1(W)$ , then  $\tilde{f}'(Z_s)$  corresponds to the  $(\mathbb{Z} \oplus \mathbb{Z})$ -subgroup  $\langle \alpha, \beta \rangle$ . Denote  $p'$  the projection of  $\tilde{M}'$ . If  $p'_*(\alpha^n) = 0$  then  $p'_*(\alpha) = 0$ , too because  $\pi_1(\tilde{B}')$  is free. Hence  $\alpha$  is homotopic to a loop in the fiber of  $\tilde{M}'$ . If  $p'_*(\alpha^n) \neq 0$  then  $p'_*(\alpha) \neq 0$ , and we can apply

**Fact 1.** *Let  $B$  be a surface with boundary,  $S$  a component of  $\partial B$ . Then the image  $\pi_1(S) \rightarrow \pi_1(B)$  is root closed if and only if  $B$  is not a Möbius band and roots are squared.*

*Proof.* First note that by [3] roots are unique in free groups. If  $\partial B$  has more than one component, then  $S$  corresponds to a primitive element of  $\pi_1(B)$ , and the statement comes from [3] ( $a^k = s^r$  implies that  $a, s$  are powers of a common element, and as  $a$  is primitive,  $s$  is a power of  $a$ ).

Now suppose  $\partial B$  has one component. Denote  $M = \text{Map}^k(S^1, S^1)$ . Suppose there exists  $a \in \pi_1(B)$  such that  $a \notin \pi_1(\partial B)$  and  $a^k \in \pi_1(\partial B)$ . It gives a  $\pi_1$ -injective map  $(M, \partial M) \rightarrow (B, \partial B)$  such that  $\pi_1(M) \rightarrow a$  and  $\pi_1(\partial M) \rightarrow a^k$ . Attach disks to  $M$  and  $B$  to get a  $\pi_1$ -injective map  $M \cup_{\partial M} D^2 \rightarrow B \cup_{\partial B} D^2$ . As  $a^k$  is a finite power of a generator of  $\pi_1(B)$ ,  $\pi_1(M \cup_{\partial M} D^2) \rightarrow \pi_1(B \cup_{\partial B} D^2)$  is of finite index. As  $\pi_1(M \cup_{\partial M} D^2) = \mathbb{Z}_k$  and  $(B \cup_{\partial B} D^2)$  is closed,  $(B \cup_{\partial B} D^2) \sim \mathbb{R}P^2$ , because  $\mathbb{R}P^2$  is the only closed surface with finite non-trivial  $\pi_1$ . Hence  $k = 2$  and  $B$  is the Möbius band.  $\square$

Hence if  $p'(\alpha^n)$  is homotopic to a loop in  $\partial \tilde{B}'$ , then so is  $p'(\alpha)$ . Hence  $\alpha$  is homotopic to a loop in  $\partial \tilde{M}'$ , and  $f'$  is homotopic to a map into  $\partial M'$ .

Case 2:  $f'$  does not lift to a map into  $\tilde{M}'$ . Then  $\alpha^n$  have roots on the singular fibers of  $M'$ , hence  $\alpha^n$  is homotopic to a loop in the regular fiber of  $M'$ . Denote the Seifert projection of  $M'$  by  $\pi'$ . As the graph-structure is reduced and  $\langle \alpha^n, \beta \rangle$  corresponds to the regular fiber of  $M$ ,  $\pi'_*(\langle \alpha^n, \beta \rangle) \neq 0$ , hence  $\pi'_*(\beta) \neq 0$ . Which means that  $\alpha$  can not commute with  $\beta$  because the roots of the regular fiber do not commute with the elements of the base.

**Step 2.2: the image of a reflector circle without corner reflectors lies in one block.** The preimage of the reflector circle is twisted  $K^2$ -bundle over  $S^1$ . Applying the previous step to the preimage of the arc between  $\partial B$  and the reflector circle, we see that  $f$  restricted to the fiber of this bundle is homotopic to a map into the block  $M$ . Take in  $B$  a loop  $\gamma$  near the reflector

covers the regular fiber of the block  $M$ , hence the whole  $f(M_{\pm id})$  can be shrunk by homotopy into  $M$ , in particular  $f(\gamma)$  does. Hence the image of the whole 1-skeleton of  $\pi^{-1}$ (reflector circle) can be shrunk by homotopy into  $M$ , and the asphericity implies the statement.

**Step 2.3 : case of reflector circles having corner reflectors.** The preimage of an arc on a reflector circle between a corner reflector and a neighboring point is  $\tilde{Z} = S^1 \tilde{\times} \text{Map}^n(S^1, S^1)$ . Denote as before  $\partial\tilde{Z} = S^1 \tilde{\times} \partial(\text{Map}^n(S^1, S^1))$  and  $\tilde{Z}_s = S^1 \tilde{\times} (\text{Map}^n(S^1, S^1))_s$ . The preimage of an arc in  $B$  between the reflector circle and the boundary is  $I \tilde{\times} K^2$ , and by previous its image by  $f$  can be shrunk into  $M$ .




Figure 9: Access to a corner reflector by two arcs.

Let us show that  $f(\tilde{Z})$  such that  $f(\partial\tilde{Z}) \subset M$  can be shrunk into  $M$ . The double covering  $p : Z \rightarrow \tilde{Z}$  induces a  $\pi_1$ -injective map  $fp : Z \rightarrow W$  which by previous can be shrunk into  $M$  (by a homotopy which is constant on  $\partial Z$ ). Let us show that the whole mapping cylinder of  $p$ ,  $\text{Map}_p(Z, \tilde{Z})$  can be shrunk into  $M$ . For this, note that the mapping cylinders of the 2-fold coverings  $p|_{\partial Z}$  and  $p|_{Z_s}$  are  $(I \tilde{\times} K^2)$ 's with boundaries lying in  $M$ , hence both can be shrunk into  $M$ . It remains to remark that the union of these mapping cylinders with  $Z$  contains a 1-skeleton of  $\text{Map}_p(Z, \tilde{Z})$ .

□

## References

- [1] R. Abraham, J. Robbin, *Transversal mappings and flows*, New York-Amsterdam 1967.-161 p.
- [2] J. A. Hillman, *Four-manifolds, geometries and knots*, Geometry & Topology Monographs, 5. Geometry & Topology Publications, Coventry, 2002.-379 p.
- [3] R. C. Lyndon, M. P. Schützenberger, *The equation  $a^M = b^N c^P$  in a free group*, Michigan Math. J. **9** (1962), pp. 289–298.
- [4] A. Mozgova, *Non-singular graph-manifolds of dimension 4*, available on <http://arXiv.org/abs/math.GT/0411335>
- [5] J. Stallings, *Group theory and three-dimensional manifolds*, A James K. Whittemore Lecture in Mathematics given at Yale University, 1969. Yale Mathematical Monographs, 4. Yale University Press, New Haven, Conn.-London, 1971.-65 p.
- [6] M. Ue, *Geometric 4-manifolds in the sense of Thurston and Seifert 4-manifolds. I*, J. Math. Soc. Japan **42** (1990), no. 3, pp. 511–540.
- [7] M. Ue, *Geometric 4 manifolds in the sense of Thurston and Seifert 4 manifolds. II*, J. Math. Soc. Japan **43** (1991), no. 1, pp. 149–183.
- [8] E. Vogt, *Vierdimensionale Seifertsche Faserräume*, Dissertation zur Erlangung des Grades eines Doktors der Naturwissenschaften der Abteilung für Mathematik an der Ruhr-Universität Bochum, Ruhr-Universität, Bochum, 1970. 151 p.
- [9] H. Zieschang, *On toric fiberings over surfaces*, Math. Notes **5** (1969), pp. 341–345.