

# Stasis Points and Approximating Two-Cycles

Stewart D. Johnson  
Department of Mathematics and Statistics  
Williams College

November 9, 2018

## Abstract

Generically, every fixed point for the differential inclusion  $x' \in \text{Convex Hull}\{f_1(x), f_2(x)\}$  can be approximated by an arbitrarily small two-cycle for the inclusion  $x' \in \{f_1(x), f_2(x)\}$ , where  $f_1, f_2$  are a  $C^1$  flows on  $\mathbb{R}^n$ .

# 1 Two-Flows, Stasis Points and Two-Cycles

Consider a controlled system that switches freely between two autonomous flows  $x' = f_1(x)$  and  $x' = f_2(x)$  where  $x \in \mathbb{R}^n$  and  $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is  $C^1$ . This is the differential inclusion

$$x' \in \{f_1(x), f_2(x)\}.$$

Trajectories are piecewise  $C^1$  functions  $x(t)$  satisfying this inclusion.

A **stasis point** is a point  $x$  at where the flows are anti-parallel:

$$f_1(x) \cdot f_2(x) = -|f_1(x)| \cdot |f_2(x)|.$$

The set of stasis points is the set of possible fixed points for the relaxed differential inclusion

$$x' \in \text{Convex Hull}(\{f_1(x), f_2(x)\}).$$

A **two-cycle**  $x(t)$  is a periodic trajectory  $x(t) = x(\delta + t)$  with  $0 < \kappa < \delta$  and

$$\begin{aligned} x' &= f_1(x) & \text{for } 0 \leq t < \kappa \\ \text{and } x' &= f_2(x) & \text{for } \kappa \leq t < \delta \end{aligned}$$

It will be shown that neighborhoods of stasis points generically contain two-cycles.

# 2 Loops in the Action

An  $C^2$  **action** of  $\mathbb{R}$  on  $\mathbb{R}^n$  is a map  $\Phi : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$  which is  $C^2$  in all variables. A **loop** is a point and time  $x_0, t_0$  with  $\Phi(x_0, t_0) = \Phi(x_0, t_0 + \delta)$  for some  $\delta > 0$ .

A point  $x_0$  is **stationary** at time  $t_0$  if  $\frac{\partial}{\partial t}\Phi(x_0, t_0) = \vec{0}$ . Note that a point that remains stationary for some open time interval,  $\frac{\partial}{\partial t}\Phi(x_0, t) \equiv \vec{0}$  for  $\alpha < t < \beta$ , is a (degenerate) loop of all sufficiently small periods.

The following theorem states that actions with stationary points generically have small loops nearby.

**Theorem 1** *Let  $\Phi : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$  be a  $C^2$  action. Suppose  $\exists x_0, t_0$  with  $\frac{\partial}{\partial t}\Phi(x_0, t_0) = \vec{0}$ . If*

$$\frac{\partial^2}{\partial x \partial t}\Phi(x_0, t_0)$$

is non-singular then  $\forall \epsilon, \exists z_0, \tau_0, \delta > 0$  with  $\Phi(z_0, \tau_0) = \Phi(z_0, \tau_0 + \delta)$  and

$$|\Phi(z_0, t) - \Phi(x_0, t_0)| < \epsilon$$

for  $\tau_0 \leq t < \tau_0 + \delta$ .

**Proof:**

Without loss of generality we take  $x_0 = \vec{0}$  and  $t_0 = 0$ .

Let

$$G(x, \delta) = \begin{cases} \frac{1}{2\delta} (\Phi(x, \delta) - \Phi(x, -\delta)) & \text{if } \delta \neq 0 \\ \frac{\partial}{\partial t} \Phi(x, 0) & \text{if } \delta = 0 \end{cases}$$

So  $G$  is  $C^1$  at  $x = \vec{0}, \delta = 0$ , and  $\frac{\partial G}{\partial x}(\vec{0}, 0) = \frac{\partial^2 \Phi}{\partial x \partial t}(\vec{0}, 0)$  is non-singular by assumption. Therefore the Implicit Function Theorem applies, and since  $G(\vec{0}, 0) = \vec{0}$  there exist solutions  $x$  to  $G(x, \delta) = \vec{0}$  for  $\delta$  sufficiently small.

For  $\delta \neq 0$ , a point  $x$  with  $G(x, \delta) = \vec{0}$  has  $\Phi(x_0, \delta) = \Phi(x_0, -\delta)$ , which is a loop. ■

### 3 Two-Cycle Approximation to Stasis

It is known that relaxed solutions to differential inclusions can be arbitrarily approximated by non-relaxed solutions [1, 2]. Here it is shown that in the case of autonomous two-flows and relaxed fixed points, the approximating trajectory can be taken as a two-cycle. For flows  $f_1, f_2$ , the idea is to take  $\mathbb{R}^n$  modulo the trajectories of  $f_1$ , and consider the remaining (non-autonomous) action of  $f_2$  on  $\mathbb{R}^{n-1}$ . Loops in this resulting action will pull back to two-cycles in the original system.

**Theorem 2** Consider two  $C^1$  flows  $f_1, f_2 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , and a stasis point  $x_0$  with  $k_1 f_1(x_0) + k_2 f_2(x_0) = \vec{0}$ ,  $k_i > 0$ . If  $\frac{d}{dx} (k_1 f_1 + k_2 f_2)$  is non-singular at  $x_0$ , then there exist two-cycles near  $x_0$ .

**Proof:** Without loss of generality we take  $x_0 = \vec{0}$ . Since  $f_1$  is  $C^1$  and non-zero at near  $\vec{0}$ , there is a local diffeomorphism  $\rho : \mathbb{R}^n \rightarrow \mathbb{R}^n$  with  $\frac{d\rho}{dx} \cdot f_1 \equiv (1, 0, \dots, 0)$  near  $\vec{0}$ . Then  $g_2 = \frac{d\rho}{dx} \cdot f_2 \circ \rho^{-1}$  is a  $C^1$  flow with  $g_2(\vec{0}) = (-\frac{k_1}{k_2}, 0, \dots, 0)$ .

We construct the  $\mathbb{R}^{n-1}$  action  $\Phi$  as follows: given  $y = (a_1, \dots, a_{n-1}) \in \mathbb{R}^{n-1}$  let  $x(t) = (x_1(t), \dots, x_n(t))$  be the trajectory under the flow  $g_2$  with  $x(0) =$

$(0, a_1, \dots, a_{n-1}) \in \mathbb{R}^n$ . Taking  $\Phi(y, t) = (x_2(t), \dots, x_n(t))$  defines a  $C^2$  action on  $\mathbb{R}^{n-1}$  with  $\frac{\partial}{\partial t}\Phi(\vec{0}, 0) = \vec{0}$ . It is straightforward to check that non-singularity of  $\frac{d}{dx}(\alpha f_1 + \beta f_2)$  at  $\vec{0} \in \mathbb{R}^n$  implies that of  $\frac{\partial^2}{\partial x \partial t}\Phi$  at  $(\vec{0}, 0) \in \mathbb{R}^{n-1} \times \mathbb{R}$ .

By Theorem 1, for all  $\epsilon > 0$  there exist  $z = (z_1, \dots, z_{n-1})$  near  $\vec{0} \in \mathbb{R}^{n-1}$  with  $\Phi(z, \alpha) = \Phi(z, \beta)$  for some  $|\alpha|, |\beta| < \epsilon$ ,  $\alpha \neq \beta$ . This corresponds to a trajectory  $x(t)$  in under  $g_2$  on  $\mathbb{R}^n$  with  $x(0) = (0, z_1, \dots, z_{n-1})$ . Now  $x(\alpha)$  and  $x(\beta)$  are equal except for their first coordinate, which cannot be zero for sufficiently small  $\epsilon$  because  $g_2$  is non-zero near  $\vec{0}$ . Thus  $x(\alpha) - x(\beta)$  is parallel to the constant flow  $g_1$  and connecting these points with a segment of the  $g_1$  flow completes the two-cycle. This maps to a two-cycle for  $f_1$  and  $f_2$  under  $\rho^{-1}$ . ■

## 4 Three Questions

For  $\mathbb{R}^2$ , every two cycle contains a stasis point [3]. Is a similar statement true for  $\mathbb{R}^n$ ?

Given a pair of flows  $f_1, f_2$ , the stasis points generically form one dimensional curves. For  $n = 2$ , an approximating two cycle will cut across this curve twice, and so the set of approximating two-cycles is parameterized by pairs of nearby points on the stasis curve [3]. Is there a similar organization of two-cycles in  $\mathbb{R}^n$ ?

Finally, a natural question is to extend this to  $N$ -flows. That is, for a differential inclusion  $x' \in \{f_1(x), f_2(x), f_3(x)\}$ , can a relaxed fixed point always be approximated by a three-cycle?

## References

- [1] A. F. Filippov, Differential equations with discontinuous righthand sides, Kluwer Academic, Dordrecht, The Netherlands, 1988.
- [2] B. Ingalls, E.D. Sontag, Y. Wang, *An infinite time relaxation theorem for differential inclusions*, Proc. of AMS, Vol 131, **2**, Pages 487-499, 2002.
- [3] S.D. Johnson, *Stasis and two-cycles*, SIAM J of Op Control, to appear.