AVERAGES OVER CLASSICAL COMPACT LIE GROUPS AND WEYL CHARACTERS

PAUL-OLIVIER DEHAYE

ABSTRACT. We compute $\mathbb{E}_G(\prod_i \operatorname{tr}(g^{\lambda_i}))$, where $G = \operatorname{Sp}(2n)$ or SO(m) (m = 2n, 2n + 1) with Haar measure. This was first obtained by Diaconis and Shahshahani [9], but our proof is more selfcontained and gives a combinatorial description for the answer. We also consider how averages of general symmetric functions $\mathbb{E}_G \Phi_n$ are affected when we introduce a Weyl character χ_{λ}^G into the integrand. We show that the value of $\mathbb{E}_G \chi_{\lambda}^G \Phi_n / \mathbb{E}_G \Phi_n$ approaches a constant for large n. More surprisingly, the ratio we obtain only changes with Φ_n and λ and is independent of the Cartan type of G. Even in the unitary case, Bump and Diaconis [4] have obtained the same ratio. Finally, those ratios can be combined with asymptotics for $\mathbb{E}_G \Phi_n$ due to Johansson [11] and provide asymptotics for $\mathbb{E}_G \chi_{\lambda}^G \Phi_n$.

1. INTRODUCTION

Historically, the study of integrals of class functions over compact classical Lie groups with respect to Haar measure has been important for many areas of mathematics and physics. We will not even attempt to describe the relevance of this problem to physics, but refer the reader to the introduction of Mehta's book [15]. On the mathematics side, we would like to mention at least the following works:

• The Heine-Szegö identity and its relations to the strong Szegö limit theorem. This identity expresses averages over unitary groups as determinants of Toeplitz matrices (see Bump and Diaconis [4]), while the strong Szegö limit theorem gives asymptotics for such determinants (see the book by Böttcher and Silbermann [1]).

Date: November 13, 2018.

²⁰⁰⁰ Mathematics Subject Classification. Primary: 15A52; Secondary: 05E5, 20G05.

Key words and phrases. random matrices, classical invariant theory, Schur-Weyl duality, symmetric functions.

This research was supported in part by the NSF grant FRG DMS-0354662.

- The study of averages of characteristic polynomials over compact classical Lie groups. Keating and Snaith conjectured that their calculations of those averages would serve as good predictors for moments of the Riemann ζ function [13, unitary case] and other data extracted from *L*-functions [12, other classical groups]. Our personal interest in Random Matrix theory sparks from this connection with Number Theory.
- Diaconis and Shahshahani's work [9] on averages of products of traces, and further refinements by Johansson [11]. Those papers have a very probabilistic flavor, and rely on separate work for their most important result. Indeed, the answer to their computations turns out to be expressible as values of characters of the Brauer algebra. Those were evaluated by Ram [19, 20], and are given by a rather complicated-looking function g in [9, Theorem 4].

The first goal of this paper will be to offer with Theorem 1 a selfcontained proof of the results of Diaconis and Shahshahani, and even a combinatorial interpretation for the mysterious g function that they obtained. If the reader only wants to understand the proof of this theorem, it might be helpful to observe that Propositions 1 and 2 include a γ that will only be useful for Theorem 3. The reader could thus safely assume that $\gamma = (0, 0, \dots)$ and still see a full proof of the following statement.

Theorem 1. Let λ be a partition, $\lambda \vdash k$ and $n \geq k$. Let $\epsilon = 1$ when $G = \operatorname{Sp}(2n)$ and $\epsilon = 0$ when $G = \operatorname{SO}(2n)$ or $\operatorname{SO}(2n+1)$. If

$$\mathbf{p}_{\lambda}(g) := \prod_{i \in \mathbb{N}} \operatorname{tr}(g^{\lambda_i})$$

then

$$\mathbb{E}_G \mathbf{p}_{\lambda} = \operatorname{sgn}(\lambda)^{\epsilon} g(\lambda),$$

where $g(\lambda)$ is defined to be the number of matchings of k points preserved under the action of a given element of S_k of cycle type λ .

We remind the reader that a *matching* of a set S is a perfect partition of S into pairs.

If we are willing to restrict the integrand to have $\lambda_i = 1$ for all i, Rains [18, Theorem 3.4] has proved this result in the full range for n. We present only the symplectic case of his result. In our notation, he proved that $\mathbb{E}_{\operatorname{Sp}(2n)}\mathbf{p}_{\lambda}(g)$ with $\lambda = (1, 1, \dots, 1) \vdash k$ is equal to the number of fixed-point-free involutions of length k with no decreasing subsequence of length greater than 2n.

 $\mathbf{2}$

In the stable range¹, he is effectively counting the number of fixedpoint-free involutions of length k, i.e. the number of matchings on kpoints preserved by the identity permutation on those k points.

The problem of Theorem 1 was also solved in full generality by Pastur and Vasilchuk [16], although their method of proof is arguably more complicated. We will sketch it in the orthogonal case. Let $F : SO(m) \to \mathbb{R}$ be a continuously differentiable function and X be any $n \times n$ real antisymmetric matrix. By left-invariance of Haar measure, $\mathbb{E}_{g \in SO(m)}F(e^{tX}g)$ is independent of the real parameter t and so $\mathbb{E}_{g \in SO(m)}(F'(g)Xg) = 0$, where F' is the derivative of F. This expression can then be expanded and used to reduce the main expression to simpler ones.

We would like to point out that our proof of Theorem 1 involves the hyperoctahedral group \mathcal{B}_k . Both Stolz [22] and Rains [17] have already used the same group for this computation.

We now turn to a more complicated problem.

Let G be U(n), SO(2n), SO(2n+1) or Sp(2n) and let $\Phi_{n,f}$ be a class function on G, essentially defined by $\Phi_{n,f}(g) = \prod_i e^{f(t_i)}$, where $\{t_i\}$ is a subset of eigenvalues of g. There are extra technical conditions on $\Phi_{n,f}$, but these will be introduced just before the statement of Theorem 3, Section 3.

The strong Szegö limit theorem gives the asymptotics and the rate of convergence of $\lim_{n\to\infty} (\mathbb{E}_{\mathrm{U}(n)} \Phi_{n,f})$. Johansson [11] was the first to generalize this theorem to the other classical groups.

The second goal of this paper will be to study how those averages and asymptotics are affected when we introduce irreducible characters of G into the integrand. The characters χ_{λ}^{G} were constructed by Weyl for the compact classical Lie groups using his Character Formula. By the Peter-Weyl theorem, these characters form a basis of the Hilbert space of class functions on G and are thus very natural to consider.

Theorem 3 will show that the ratio

$$rac{\mathbb{E}_{G} oldsymbol{\chi}_{\lambda}^{G} oldsymbol{\Phi}_{n,f}}{\mathbb{E}_{G} oldsymbol{\Phi}_{n,f}}$$

approaches a limit when n >> 0. This extends the corresponding results for the unitary groups due to Bump and Diaconis [4] to other classical groups. Remarkably, our ratio is independent of the Cartan type of the group G and equal to the ratio they obtained for the unitary groups. It only varies with f and λ and can also be seen as the value

¹See page 7.

achieved by the Schur polynomial s_{λ} after setting the values of power polynomials to some Fourier coefficients of f.

A different point of view is offered in Bump, Diaconis and Keller [5]: we can modify the Haar measure dg into $\chi^G_{\lambda} \chi^G_{\lambda} dg$. We know that $\chi^G_{\lambda} \chi^G_{\lambda}$ is always positive and of mass 1 by orthogonality of Weyl characters hence $\chi^G_{\lambda} \overline{\chi^G_{\lambda}} dg$ is a measure. With this point of view, Theorem 3 would thus partially explain how the average of $\Phi_{n,f}$ with respect to Haar measure dg is modified when *twisting* the Haar measure by a character (see the last two remarks on page 16).

Thirdly, we would like to mention the recent preprint of Bump and Gamburd [6]. They showed how many of the integrals useful for Number Theory can be computed in a unified way. An example of such an integral would be

$$\int_{\mathrm{U}(n)}\prod_i\Lambda_g(e^{\alpha_i})\,\mathrm{d}g,$$

where $\Lambda_g(\cdot)$ is the characteristic polynomial of g, and the α_i 's are points on the unit circle. The importance of integrals of this type originates from the work of Keating and Snaith [12, 13], where the integrals have been shown to predict the moments of $\zeta(\cdot)$ and of L-functions.

The method of Bump and Gamburd is based on symmetric function theory and classical results (Weyl Character Formula, Littlewood Branching Rules of Theorem 2, page 8, and Cauchy Identity). The reader is referred to their introduction for a much more comprehensive survey of all the results their method is known to produce, and how (if) they were proved before.

This type of work is useful because it consolidates a wide array of methods into one more systematic technique.

In that same vein, we hope that this paper can complement theirs to get closer to a more universal method. Indeed, we have shown how to introduce elements of the basis of symmetric functions into the integrand, an interesting step for that goal. Further steps are taken in the author's Ph.D. thesis and associated paper [7].

Section 2 will first go over notation, then introduce the reader to the representation theory of the compact classical Lie groups (Weyl characters and Branching Rules). Section 3 will contain all of the proofs. It will also present the statement of Theorem 3, and then shortly discuss its significance in relation to the rest of the literature.

The author is pleased to thank Daniel Bump and Persi Diaconis for numerous stimulating discussions. Alex Gamburd clarified some of the technical details of Section 2.2 and suggested some of the references. Finally, I would like to thank the people in my entourage for their unfaltering support.

2. Representation theory of the classical groups

We now introduce Weyl characters and the branching rules between different classical compact Lie groups. We follow the expositions of [6] and [14], but our notation is closer to [6].

2.1. Notation.

Partitions. A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ is a finite decreasing sequence of non-negative integers. We define the weight $|\lambda|$ of λ to be the sum $\sum \lambda_i$. If this weight is k, we also use the notation $\lambda \vdash k$. The length $l(\lambda)$ of λ is the maximal i such that $\lambda_i \neq 0$. The conjugate of λ is denoted λ' . We say that a partition is even if all of its parts λ_i are even. We define the union $\lambda \cup \mu$ to be the partition of $|\lambda| + |\mu|$ whose parts are the union of the parts of λ and μ . There is a partial ordering on partitions: $\lambda \subseteq \mu$ iff $\lambda_i \leq \mu_i$ for all i. Finally, we define the $\lambda(i)$'s so that $(i^{\lambda(i)}) = (\lambda_1, \lambda_2, \dots, \lambda_n)$, i.e. $\lambda(i)$ counts the number of λ_j 's equal to i.

Symmetric group. The symmetric group on k points will be S_k . If $\lambda \vdash k$, elements of type λ are the elements whose cycle types correspond to the partition λ . We use \mathcal{C}_{λ} for the conjugacy class of those elements. We denote a centralizer in the group G by $C_G(\cdot)$, and by z_{λ} the order of the centralizer of an element of \mathcal{C}_{λ} . As usual, the irreducible characters χ_{λ} of S_k are indexed by partitions $\lambda \vdash k$. We sometimes abuse notation and take $\chi_{\lambda}(\mu)$ to mean the value of χ_{λ} on \mathcal{C}_{μ} . If χ_{λ} and χ_{μ} are characters of $S_{|\lambda|}$ and $S_{|\mu|}$, their product $\chi_{\lambda} \odot \chi_{\mu}$ in the character ring of symmetric groups will be the character $\operatorname{Ind}_{S_{|\lambda|} \times S_{|\mu|}}^{S_{|\lambda|} \times |\mu|} (\chi_{\lambda} \times \chi_{\mu})$ (see Sagan's book [21] for all aspects of the representation theory of symmetric groups, and page 164 for the product of characters $\chi_{\lambda} \odot \chi_{\mu}$).

Classical groups. Let J be the $2n \times 2n$ matrix given by

$$J = \begin{pmatrix} 0 & -\operatorname{Id}_n \\ \operatorname{Id}_n & 0 \end{pmatrix}.$$

We would like to introduce a few classical groups:

$$U(n) = \{g \in M_n(\mathbb{C}) \mid gg^* = I\}, \\O(n) = \{g \in U(n) \mid gg^t = I\}, \\SO(n) = \{g \in O(n) \mid \det(g) = 1\}, \\Sp(2n) = \{g \in U(2n) \mid gJg^t = J\}.$$

If G is one of those groups, it is compact for the topology induced by $M_n(\mathbb{C})$ or $M_{2n}(\mathbb{C})$. We can thus consider its Haar measure dg and normalize it so the total volume of G is 1. We write $\mathbb{E}_G f$ for $\int_G f(g) dg$.

Symmetric polynomials and power characters. Let $\mathbb{C}[x_1, \dots, x_m]^{\mathbb{S}_m}$ be the ring of symmetric polynomials in m variables. We define the power polynomials $p_i(x_1, \dots, x_m) = x_1^i + \dots + x_m^i$ and $p_\lambda(x_1, \dots, x_m) =$ $\prod_i p_{\lambda_i}(x_1, \dots, x_m)$. By abuse of notation, we also denote by p_λ the generalized character of $\mathbb{S}_{|\lambda|}$ that is the indicator function with value z_λ on the conjugacy class of type λ (see Sagan [21, page 162]). The difference in the arguments of p_λ should prevent any ambiguity. Note that the polynomial p_λ is the image of the character p_λ under the characteristic map (see Bump's book [2, Theorem 39.1]). Finally, we define the characters \mathbf{p}_λ of G = U(m), O(m), SO(m) or Sp(m = 2n)by $\mathbf{p}_\lambda(g) := p_\lambda(t_1, t_2, \dots, t_m)$ where the t_i 's are all the eigenvalues of g. There is an obvious interpretation of those generalized characters in terms of the trace. For instance, we have $\mathbf{p}_{(3,1,1)}(g) = \operatorname{tr}(g^3) \cdot (\operatorname{tr} g)^2$.

2.2. Weyl characters. Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition. Let *i* and *j* be indices running between 1 and *n*. Guided by the Weyl Character Formula, we define the following symmetric functions of $\{x_1, \dots, x_n\}$, actually polynomials in $\mathbb{Z}[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$:

$$s_{\lambda}(x_{1}, \cdots, x_{n}) = \frac{\left|x_{i}^{\lambda_{j}+n-j}\right|}{\left|x_{i}^{n-j}\right|},$$

$$\chi_{\lambda}^{\mathrm{SO}(2n+1)}(x_{1}, \cdots, x_{n}) = \frac{\left|x_{i}^{\lambda_{j}+n-j+1/2} - x_{i}^{-(\lambda_{j}+n-j+1/2)}\right|}{\left|x_{i}^{n-j+1/2} - x_{i}^{-(n-j+1/2)}\right|},$$

$$\chi_{\lambda}^{\mathrm{Sp}(2n)}(x_{1}, \cdots, x_{n}) = \frac{\left|x_{i}^{\lambda_{j}+n-j+1} - x_{i}^{-(\lambda_{j}+n-j+1)}\right|}{\left|x_{i}^{n-j+1} - x_{i}^{-(n-j+1)}\right|}.$$

The $s_{\lambda}(\cdot)$ are the regular Schur polynomials that appear in the representation theory of the symmetric group. Take $g \in U(n)$ (resp. SO(2n+1) or Sp(2n)). Label the eigenvalues of g by $\{t_1, \dots, t_n\}$ (resp. $\{t_1, t_1^{-1}, \dots, t_n, t_n^{-1}, 1\}$ or $\{t_1, t_1^{-1}, \dots, t_n, t_n^{-1}\}$). This allows us to define the functions $\mathbf{s}_{\lambda}(g), \boldsymbol{\chi}_{\lambda}^{Sp(2n)}(g)$ or $\boldsymbol{\chi}_{\lambda}^{SO(2n+1)}(g)$ through the values of the respective function on the subset $\{t_1, \dots, t_n\}$.

When $G = \mathrm{SO}(2n+1)$ (resp. $\mathrm{Sp}(2n)$), Weyl showed that the character $\boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n+1)}$ (resp. $\boldsymbol{\chi}_{\lambda}^{\mathrm{Sp}(2n)}$) is irreducible when $l(\lambda) \leq n$. This is called the *stable range* for λ^2 .

Due to the involution in the Dynkin diagram of type D_n , the case of $\boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)}$ is actually special. We will again define $\boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)}(g)$ as the value of a function $\boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)}$ on an appropriate subset of n eigenvalues of g. The difference in this case is that we only have $\lambda_1 \geq \lambda_2 \geq \cdots \geq |\lambda_n|$ for the index set. If λ is a regular partition, we define $\lambda_+ := \lambda = (\lambda_1, \lambda_2, \cdots, \lambda_n)$ and $\lambda_- := (\lambda_1, \lambda_2, \cdots, -\lambda_n)$. The characters $\boldsymbol{\chi}_{\lambda^+}^{\mathrm{SO}(2n)}$ and $\boldsymbol{\chi}_{\lambda^-}^{\mathrm{SO}(2n)}$ are exchanged by the involution on the Dynkin diagram, i.e by conjugation by an element of O(2n) of negative determinant³.

The Weyl character formula defines the functions

$$\frac{\chi_{\lambda}^{\text{SO}(2n)}(x_{1},\cdots,x_{n})}{\left|x_{i}^{\lambda_{j}+n-j}+x_{i}^{-(\lambda_{j}+n-j)}\right|+\left|x_{i}^{\lambda_{j}+n-j}-x_{i}^{-(\lambda_{j}+n-j)}\right|}{\left|x_{i}^{n-j}+x_{i}^{-(n-j)}\right|}.$$

If we set $\chi_{\lambda}^{O(2n)} := \chi_{\lambda^+}^{SO(2n)} + \chi_{\lambda^-}^{SO(2n)}$ when $\lambda_n \neq 0$ and $\chi_{\lambda}^{O(2n)} := \chi_{\lambda^+}^{SO(2n)}$ otherwise, then

$$\chi_{\lambda}^{\mathcal{O}(2n)}(x_1,\cdots,x_n) = \frac{\left|x_i^{\lambda_j+n-j} + x_i^{-(\lambda_j+n-j)}\right|}{\left|x_i^{n-j} + x_i^{-(n-j)}\right|}.$$

The character $\boldsymbol{\chi}_{\lambda}^{\mathrm{O}(2n)}(g)$ is defined similarly by evaluating $\chi_{\lambda}^{\mathrm{O}(2n)}$ on eigenvalues.

It is still a consequence of Weyl's work that $\boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)}$ is an irreducible character of $\mathrm{SO}(2n)$ when $l(\lambda) \leq n$. However, $\boldsymbol{\chi}_{\lambda}^{\mathrm{O}(2n)}$ will merely be the character of the representation of $\mathrm{SO}(2n)$ which is obtained by

 $^{^{2}}$ The book of Goodman and Wallach [10, Chapter 10] is the standard reference for this. See also the paper of Koike and Terada [14].

³It might be helpful for the reader to observe that in the odd orthogonal case, $O(2n+1) \cong SO(2n+1) \times \mathbb{Z}/2$ so the involution acts trivially.

restricting an irreducible representation of O(2n) to SO(2n), not the character of a representation of O(2n).

For the sake of uniformity in the orthogonal case, we will sometimes want to use $\boldsymbol{\chi}_{\lambda}^{O(2n+1)} := \boldsymbol{\chi}_{\lambda}^{SO(2n+1)}$.

We also use the notational shortcut χ_{λ}^{G} where G is one of the Lie groups defined above. It might be good at this point to remind the reader that χ_{λ} denotes a character of a symmetric group.

The irreducibility of the various characters considered guarantees certain orthogonality properties, which we will only describe as needed in the proofs.

2.3. Branching rules. Let G = SO(m) or Sp(m). Since $G \subset U(m)$, the restriction of \mathbf{s}_{λ} to G is a class function for G and can be expressed as a sum of $\boldsymbol{\chi}_{\mu}^{G}$'s. The branching rules describe more precisely how to do that (see the paper of Koike and Terada [14, page 492] for a modern and complete proof).

Theorem 2 (Littlewood). Let λ be a partition of length less than or equal to n. Then

$$\begin{aligned} s_{\lambda} \Big|_{\operatorname{Sp}(2n)}^{\operatorname{U}(2n)} &= \sum_{\mu \subseteq \lambda} \left(\sum_{\nu \text{ even}} c_{\nu'\mu}^{\lambda} \right) \chi_{\mu}^{\operatorname{Sp}(2n)}, \\ s_{\lambda} \Big|_{\operatorname{SO}(2n+1)}^{\operatorname{U}(2n+1)} &= \sum_{\mu \subseteq \lambda} \left(\sum_{\nu \text{ even}} c_{\nu\mu}^{\lambda} \right) \chi_{\mu}^{\operatorname{O}(2n+1)} \\ s_{\lambda} \Big|_{\operatorname{SO}(2n)}^{\operatorname{U}(2n)} &= \sum_{\mu \subseteq \lambda} \left(\sum_{\nu \text{ even}} c_{\nu\mu}^{\lambda} \right) \chi_{\mu}^{\operatorname{O}(2n)}, \end{aligned}$$

where $\mathbf{s}_{\lambda} \downarrow_{G}^{\mathrm{U}(n)}$ indicates the restriction to G of the character \mathbf{s}_{λ} of $\mathrm{U}(n)$ and $c_{\nu\mu}^{\lambda}$ are the Littlewood-Richardson coefficients.

Remark. This is where the eigenvalue 1 "disappears" in the SO(2n+1) case. Let $g \in SO(2n+1) \subset U(2n+1)$, with eigenvalues $\{1, t_1, \dots, t_n, t_1^{-1}, \dots, t_n^{1}\}$. The left-hand side is

$$\mathbf{s}_{\lambda}(g) = s_{\lambda}(1, t_1, \cdots, t_n, t_1^{-1}, \cdots, t_n^{-1}),$$

while the right-hand side only involves terms of the form

$$\boldsymbol{\chi}^{\mathcal{O}(2n+1)}_{\mu}(g) = \chi^{\mathcal{O}(2n+1)}_{\mu}(t_1,\cdots,t_n).$$

3. Proofs

We will now present the main derivation. This is vaguely similar to a few steps of the proof of [8, Theorem 2.1] in the unitary case.

Proposition 1. Let $\lambda \vdash k$ and $n \geq k$. Then

$$\mathbb{E}_{\mathrm{Sp}(2n)} \boldsymbol{\chi}_{\gamma}^{\mathrm{Sp}(2n)} \mathbf{p}_{\lambda} = \sum_{\substack{\beta' \text{ even} \\ \gamma \cup \beta \vdash k}} \langle \chi_{\gamma} \odot \chi_{\beta}, p_{\lambda} \rangle_{\mathcal{S}_{k}} \,.$$

Similarly (but with β instead of β'), we have

$$\mathbb{E}_{\mathrm{SO}(2n+1)}\boldsymbol{\chi}_{\gamma}^{\mathrm{SO}(2n+1)}\mathbf{p}_{\lambda} = \sum_{\substack{\beta \text{ even}\\\gamma\cup\beta\vdash k}} \langle \chi_{\gamma}\odot\chi_{\beta}, p_{\lambda}\rangle_{\mathcal{S}_{k}} = \mathbb{E}_{\mathrm{SO}(2n)}\boldsymbol{\chi}_{\gamma}^{\mathrm{SO}(2n)}\mathbf{p}_{\lambda}$$

Note: when $|\gamma| > |\lambda| = k$ or when $k - |\gamma|$ is odd, those sums are indeed trivial and give a value of 0.

Proof. The general method of proof is to use the branching rules from Section 2.3 to eventually transfer the problem to a symmetric group.

For definiteness, we will only prove this for Sp(2n) and discuss at the end the minor changes needed in the orthogonal cases. Let $g \in \text{Sp}(2n)$ have eigenvalues $\{t_1, t_1^{-1}, \dots, t_n, t_n^{-1}\}$. Then

$$\begin{aligned} \mathbf{p}_{\lambda}(g) &= \sum_{\mu \vdash k} \chi_{\mu}(\lambda) \mathbf{s}_{\mu}(g) \\ &= \sum_{\mu \vdash k} \chi_{\mu}(\lambda) \sum_{\nu \subseteq \mu} \left(\sum_{\beta' \text{ even}} c_{\nu\beta}^{\mu} \right) \boldsymbol{\chi}_{\nu}^{\mathrm{Sp}(2n)}(g), \end{aligned}$$

where the first line follows from the usual decomposition of power polynomials into Schur polynomials given by the character table of a symmetric group (see Sagan [21, Equation (4.23)]). The second line follows by applying the branching rule for each $\mu \vdash k$. The branching rule is only valid when $l(\mu) \leq n$. This explains our final restriction of $n \geq k$. We know that $\mathbb{E}_{\mathrm{Sp}(2n)} \chi_{\gamma}^{\mathrm{Sp}(2n)} \chi_{\nu}^{\mathrm{Sp}(2n)} = 1$ when $\gamma = \nu$ and 0 otherwise

We know that $\mathbb{E}_{\text{Sp}(2n)} \boldsymbol{\chi}_{\gamma}^{\text{Sp}(2n)} \boldsymbol{\chi}_{\nu}^{\nu} = 1$ when $\gamma = \nu$ and 0 otherwise (this is a consequence of the theory of the Weyl Character formula). Hence

$$\mathbb{E}_{\mathrm{Sp}(2n)} \boldsymbol{\chi}_{\gamma}^{\mathrm{Sp}(2n)} \mathbf{p}_{\lambda} = \sum_{\mu \vdash k} \left(\chi_{\mu}(\lambda) \sum_{\beta' \text{ even}} c_{\gamma\beta}^{\mu} \right),$$

where the condition that $\nu = \gamma \subseteq \mu$ is still present implicitly in the Littlewood-Richardson coefficient ($c^{\mu}_{\gamma\beta} = 0$ if $\gamma \not\subseteq \mu$). For the same reason, we see that this sum is trivial when $|\gamma| > |\mu| = k$.

The final statement follows from observing that $\sum_{\mu \vdash k} c^{\mu}_{\gamma\beta} \chi_{\mu} = \chi_{\gamma} \odot \chi_{\beta}$ and $\chi(\lambda) = \langle \chi, p_{\lambda} \rangle_{S_k}$.

For the orthogonal groups, the only difference is that two characters will pop up when $\lambda_n \neq 0$. Let m = 2n or 2n + 1. The branching rules will involve $\boldsymbol{\chi}_{\lambda}^{O(m)}$ while the twist that we introduce comes from a character of type $\boldsymbol{\chi}_{\lambda}^{SO(m)}$. Fortunately, all we need for the same proof to work is $\mathbb{E}_{G} \boldsymbol{\chi}_{\lambda}^{O(m)} \boldsymbol{\chi}_{\lambda}^{SO(m)} = 1$:

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\lambda}^{\mathrm{O}(2n)} \boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)} = \mathbb{E}_{G} \boldsymbol{\chi}_{\lambda^{+}}^{\mathrm{SO}(2n)} \boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)} + \mathbb{E}_{G} \boldsymbol{\chi}_{\lambda^{-}}^{\mathrm{SO}(2n)} \boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n)}$$

$$= 1 + 0 \text{ by orthonormality for SO}(2n).$$

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\lambda}^{\mathrm{O}(2n+1)} \boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n+1)} = \mathbb{E}_{G} \boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n+1)} \boldsymbol{\chi}_{\lambda}^{\mathrm{SO}(2n+1)}$$

$$= 1 \text{ by orthonormality for SO}(2n+1).$$

We would like to remind the reader at this point of a few facts from the representation theory of the symmetric group.

Lemma 1. Let sgn be the sign character in S_k .

(1) If $\beta \vdash k$, then $\chi_{\beta'} = \operatorname{sgn} \otimes \chi_{\beta}$, (2) If $\beta \vdash k$, then

$$p_{\beta} \otimes \operatorname{sgn} = \operatorname{sgn}(\beta) p_{\beta}$$

(3) Restrict k to be even. Then

$$\sum_{\substack{\beta \text{ even}\\\beta\vdash k}} \chi_{\beta} = \operatorname{Ind}_{\mathcal{B}_{k}}^{\mathcal{S}_{k}} 1,$$

where \mathfrak{B}_k is the centralizer of the chosen permutation (1,2) $(3,4)\cdots(k-1,k)$ in \mathfrak{S}_k .

(4) Restrict k to be even. Then

$$\operatorname{sgn} \otimes \operatorname{Ind}_{\mathcal{B}_k}^{\mathcal{S}_k} 1 = \operatorname{Ind}_{\mathcal{B}_k}^{\mathcal{S}_k} (\operatorname{Res}_{\mathcal{B}_k}^{\mathcal{S}_k} \operatorname{sgn}).$$

Proof. (1) This is in Bump's book [2, Theorem 39.3].(2) This is immediate:

$$p_{\lambda} \otimes \operatorname{sgn} = \sum_{\mu \vdash k} \chi_{\mu}(\lambda) \chi_{\mu} \otimes \operatorname{sgn} = \sum_{\mu \vdash k} \chi_{\mu}(\lambda) \chi_{\mu'}$$

and, by part 1,

$$=\sum_{\mu\vdash k}\chi_{\mu'}(\lambda)\chi_{\mu}=\sum_{\mu\vdash k}\chi_{\mu}(\lambda)\operatorname{sgn}(\lambda)\chi_{\mu}$$

(3) See [2, Theorem 45.4].

(4) This is a consequence of Frobenius Reciprocity.

This lemma leads immediately to a second version of Proposition 1.

Proposition 2. Let $\lambda \vdash k$ and $n \geq k$. Let $\epsilon = 1$ when G = Sp(2n)and $\epsilon = 0$ when G = SO(2n) or SO(2n+1). Then

$$\mathbb{E}_{G}\boldsymbol{\chi}_{\gamma}^{G}\mathbf{p}_{\lambda} = \left\langle \operatorname{Ind}_{\mathcal{S}_{|\gamma|} \times \mathcal{B}_{k-|\gamma|}}^{\mathcal{S}_{k}} \left(\chi_{\gamma} \otimes \operatorname{sgn}^{\epsilon} \right), p_{\lambda} \right\rangle_{\mathcal{S}_{k}},$$

where by a slight abuse of notation, we confuse sgn and $\operatorname{Res}_{\mathcal{B}_k}^{\mathcal{S}_k}$ sgn.

Proof. All the steps required are applications of Lemma 1 to the statement of Proposition 1.

$$\begin{split} \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda} &= \sum_{\substack{\beta \text{ even}\\\gamma \cup \beta \vdash k}} \left\langle \chi_{\gamma} \odot \left(\mathrm{sgn}^{\epsilon} \right) \otimes \chi_{\beta}, p_{\lambda} \right\rangle_{\mathcal{S}_{k}} \\ &= \left\langle \chi_{\gamma} \odot \left(\mathrm{sgn}^{\epsilon} \otimes \mathrm{Ind}_{\mathcal{B}_{k-|\gamma|}}^{\mathcal{S}_{k-|\gamma|}} 1 \right), p_{\lambda} \right\rangle_{\mathcal{S}_{k}} \end{split}$$

We now apply Lemma 1.4 to get the result stated.

3.1. Discussion of Theorem 1. As a special case to Proposition 2, we are now ready to compute integrals of traces directly, without involving the Brauer algebra as in Ram [20].

Proof of Theorem 1. We want here to compute $\mathbb{E}_{G}\mathbf{p}_{\lambda}$, so we are now in the simplest case of Proposition 2, when $|\gamma| = 0$. When k is odd, there is simply no matching on k points. On the other hand, it was a consequence of Proposition 1 that $\mathbb{E}_{G}\mathbf{p}_{\lambda} = 0$ as $k - |\gamma| = k$ is odd. We can thus restrict our attention to the k even case. We have thanks to Lemma 1 that

$$\mathbb{E}_{G}\mathbf{p}_{\lambda} = \left\langle \operatorname{Ind}_{\mathcal{B}_{k}}^{S_{k}} 1, p_{\lambda} \otimes \operatorname{sgn}^{\epsilon} \right\rangle_{S_{k}} \\
= \operatorname{sgn}(\lambda)^{\epsilon} \left\langle 1, \operatorname{Res}_{\mathcal{B}_{k}}^{S_{k}} p_{\lambda} \right\rangle_{\mathcal{B}_{k}} \\
= \frac{z_{\lambda} \operatorname{sgn}(\lambda)^{\epsilon}}{|\mathcal{B}_{k}|} \# \left\{ \sigma \in C_{S_{k}}((1,2)\cdots(k-1,k)) \mid \operatorname{type}(\sigma) = \lambda \right\},$$

since p_{λ} is an indicator function for the conjugacy class of permutations of type λ in S_k .

If $\sigma \in C_{S_k}((1,2)\cdots(k-1,k))$ then σ preserves the matching $\{\{1,2\}, \cdots, \{k-1,k\}\}$, i.e. it sends a pair to a pair. We use this to switch to the language of matchings.

$$\mathbb{E}_{G}\mathbf{p}_{\lambda} = \frac{\operatorname{sgn}(\lambda)^{\epsilon}}{|\mathcal{C}_{\lambda}|} \frac{|\mathcal{S}_{k}|}{|\mathcal{B}_{k}|} \# \{ \sigma \in C_{\mathcal{S}_{k}}((1,2)(3,4)\cdots(k-1,k)) \cap \mathcal{C}_{\lambda} \}$$

$$= \frac{\operatorname{sgn}(\lambda)^{\epsilon}}{|\mathcal{C}_{\lambda}|} \sum_{\operatorname{matching } M \text{ of } k \text{ points}} \# \{ \sigma \in \mathcal{C}_{\lambda} \mid \sigma(M) = M \}$$

$$= \frac{\operatorname{sgn}(\lambda)^{\epsilon}}{|\mathcal{C}_{\lambda}|} \# \{ (M,\sigma) \mid M \text{ a matching of } k \text{ points},$$

$$\sigma \in \mathcal{C}_{\lambda}, \sigma(M) = M \}$$

$$= \frac{\operatorname{sgn}(\lambda)^{\epsilon}}{|\mathcal{C}_{\lambda}|} \sum_{\sigma \in \mathcal{C}_{\lambda}} \# \{ \operatorname{matchings preserved by } \sigma \}.$$

The last steps make use of a double-counting argument. All the summands in the last line are equal, and there are $|\mathcal{C}_{\lambda}|$ of them so we have

$$\mathbb{E}_G \mathbf{p}_{\lambda} = \operatorname{sgn}(\lambda)^{\epsilon} g(\lambda),$$

where $g(\lambda)$ is the number of matchings preserved by a permutation of cycle type λ .

As mentioned earlier, this offers a combinatorial interpretation for a result first proved by Diaconis and Shahshahani [9]. Naturally, we have to check that our definition of g agrees with the definition they gave. This is a purely combinatorial problem.

Proposition 3. Let $\lambda \vdash k$. Then $g(\lambda) = \prod_j g_j(\lambda(j))$, where $g_j(\cdot)$ is given by

$$if j is odd \quad g_j(a) = \begin{cases} 0 & \text{if } a \text{ is odd} \\ j^{a/2}(a-1)(a-3)\cdots 1 & \text{if } a \text{ is even} \end{cases}$$
$$if j is even \quad g_j(a) = \sum_t \binom{a}{2t} j^t (2t-1)(2t-3)\cdots 1$$

Proof. From our combinatorial definition of g, it is immediate that $g(\lambda) = \prod_j g((j^{\lambda(j)}))$. All we have left to prove is $g((j^a)) = g_j(a)$.

if j is odd: : Take $\sigma \in C_{(j^a)}$. Since each cycle of σ is of odd length, any matching of points preserved by σ must match cycles as well. If a is odd there is no such matching. If a is even, any matching of points will also match cycles. There are $(a-1)(a-3)\cdots 1$ possible matchings of cycles. Once a matching of cycles is chosen, we still have to decide on how to match points in each individual pair of cycles. There are j choices for each of the a/2 pairs of cycles. if j is even: : This is more subtle, as matchings of points inside the same cycle are allowed. Say there are 2t cycles whose points are matched with points in another cycle (the *external* cycles) and thus a - 2t cycles whose points are matched with a point within the same cycle (the *internal* cycles). There are $\binom{a}{2t}$ ways of choosing which cycles will be external, and then $(2t-1)(2t-3)\cdots 1$ ways of matching external cycles. Once we have a pair of external cycles, there are j ways of matching points between the two cycles. On the other hand, there is a unique way of matching points within an internal cycle: a point has to be paired with the point most distant for the ordering given by the cycle.

3.2. Discussion of Theorem 3. Let $\mathbb{T} = \{t \in \mathbb{C} \mid |t| = 1\}$, and let $\sigma(t) = \sum_{i \in \mathbb{Z}} d_i t^i = \exp\left(\sum_{i \in \mathbb{Z}} c_i t^i\right) = e^{f(t)}$ be a function on \mathbb{T} . We will always assume $f(t^{-1}) = f(t)$ (i.e. $c_i = c_{-i}$).

We will also need two extra conditions:

Condition (A):

$$\sum |c_i| < \infty$$

Condition (B):

$$\sum |i||c_i|^2 < \infty$$

Those conditions were already relevant to the work of Bump and Diaconis [4], and the whole field of Toeplitz matrices⁴.

One can define a class function $\Phi_{n,f}(g)$ on G as

$$\Phi_{n,f}(g) = e^{nc_0} \exp\left(\sum_{i>0} c_i \mathbf{p}_{(i)}(g)\right).$$

A possibly more intuitive definition (but only valid when $G = \operatorname{Sp}(2n)$ or $G = \operatorname{SO}(2n)$) is $\Phi_{n,f}(g) = \prod_{k=1}^{n} \sigma(t_k)$, where the product is taken over half of the eigenvalues of g, one in each conjugate pair. The symmetry condition $f(t^{-1}) = f(t)$ guarantees that $\Phi_{n,f}$ is independent of the chosen subset of eigenvalues. When $G = \operatorname{SO}(2n+1)$, the product expression becomes slightly more complicated because of the eigenvalue 1.

⁴The book by Böttcher and Silbermann [1] gives a very clear introduction to the analytic theory of Toeplitz matrices. Theorem 5.2 in [1] uses those conditions. Sets of functions satisfying Conditions (A) and (B) are denoted $W(\mathbb{T})$ and $B_2^{1/2}(\mathbb{T})$ respectively.

Theorem 3. Assume that f satisfies Condition (A). For simplicity of notation, take $\boldsymbol{\chi}_{\gamma}^{G} = \boldsymbol{\chi}_{\gamma}^{\mathrm{SO}(2n+1)}$ (resp. $\boldsymbol{\chi}_{\gamma}^{\mathrm{Sp}(2n)}, \boldsymbol{\chi}_{\gamma}^{\mathrm{SO}(2n)}$) if $G = \mathrm{SO}(2n+1)$ (resp. $\mathrm{Sp}(2n), \mathrm{SO}(2n)$). Then

$$\lim_{n \to \infty} \frac{\mathbb{E}_G \boldsymbol{\chi}_{\gamma}^G \boldsymbol{\Phi}_{n,f}}{\mathbb{E}_G \boldsymbol{\Phi}_{n,f}} = R(\gamma, (c_i)),$$

with

$$R(\gamma, (c_i)) = \sum_{\lambda \vdash |\gamma|} \chi_{\gamma}(\lambda) \left(\prod_{i=1}^{\infty} \frac{c_i^{\lambda(i)}}{\lambda(i)!} \right)$$
$$= s_{\gamma} \Big|_{p_i:=ic_i},$$

where the last expression is a specialization for the Schur polynomial s_{γ} when the value of the power polynomials is set using the Fourier coefficients c_i .

We delay comments on this Theorem to page 16 and start with the proof.

Proof. As a first approximation to $\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \boldsymbol{\Phi}_{n,f}$, we will actually study $\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda}$ for $\lambda \vdash k \leq n$. It will be useful to split up λ into subpartitions. To avoid confusion with notation previously used for partition parts $(\lambda_{1}, \lambda_{2}, \cdots, \lambda_{n})$, we will use $\lambda_{a} \cup \lambda_{b} = \lambda$ in this proof only.

We start from the final equation in Proposition 2 and apply Frobenius Reciprocity to get

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda} = \left\langle \chi_{\gamma} \otimes \operatorname{Res}_{\mathcal{B}_{k-|\gamma|}}^{\mathcal{S}_{k-|\gamma|}} \operatorname{sgn}^{\epsilon}, \operatorname{Res}_{\mathcal{S}_{|\gamma|} \times \mathcal{B}_{k-|\gamma|}}^{\mathcal{S}_{k}} p_{\lambda} \right\rangle_{\mathcal{S}_{|\gamma|} \times \mathcal{B}_{k-|\gamma|}} \\ = \frac{z_{\lambda}}{|\mathcal{S}_{|\gamma|}||\mathcal{B}_{k-|\gamma|}|} \sum_{\substack{(\rho_{a},\rho_{b}) \in \mathcal{S}_{|\gamma|} \times \mathcal{B}_{k-|\gamma|} \\ \operatorname{type}(\rho_{a}) = \lambda_{a} \vdash |\gamma| \\ \operatorname{type}(\rho_{b}) = \lambda_{b} \vdash k-|\gamma|} \chi_{\gamma}(\rho_{a}) \operatorname{sgn}^{\epsilon}(\rho_{b}),$$

where $\epsilon = 1$ when $G = \operatorname{Sp}(2n)$ and 0 otherwise. We now sum over conjugacy classes (i.e. cycle types) instead. The correction factor for the ρ_a 's of type λ_a will be $\frac{|\mathcal{S}_{|\gamma|}|}{z_{\lambda_a}} = |\mathcal{C}_{\lambda_a}|$, so

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda} = \frac{z_{\lambda}}{|\mathcal{B}_{k-|\gamma|}|} \sum_{\substack{\lambda_{a} \vdash |\gamma| \\ \lambda_{a} \cup \lambda_{b} = \lambda}} \frac{\chi_{\gamma}(\lambda_{a}) \operatorname{sgn}(\lambda_{b})^{\epsilon}}{z_{\lambda_{a}}} |\mathcal{B}_{k-|\gamma|} \cap \mathcal{C}_{\lambda_{b}}|.$$

Observe from the proof of Theorem 1, with λ replaced by λ_b , that

$$\mathbb{E}_{G} \mathbf{p}_{\lambda_{b}} = rac{z_{\lambda_{b}} \operatorname{sgn}(\lambda_{b})^{\epsilon}}{|\mathcal{B}_{k-|\gamma|}|} |\mathcal{B}_{k-|\gamma|} \cap \mathfrak{C}_{\lambda_{b}}|.$$

The hypothesis $n \geq |\lambda_b|$ of Theorem 1 is automatically satisfied since we already assume $n \geq |\lambda|$ and $\lambda = \lambda_a \cup \lambda_b$.

We now have the much simpler

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda} = \sum_{\substack{\lambda_{a} \vdash |\gamma| \\ \lambda_{a} \cup \lambda_{b} = \lambda}} \frac{z_{\lambda}}{z_{\lambda_{a}} z_{\lambda_{b}}} \chi_{\gamma}(\lambda_{a}) \mathbb{E}_{G} \mathbf{p}_{\lambda_{b}}$$

or even

(1)
$$\mathbb{E}_{G}\boldsymbol{\chi}_{\gamma}^{G}\mathbf{p}_{\lambda} = \sum_{\substack{\lambda_{a} \vdash |\gamma| \\ \lambda_{a} \cup \lambda_{b} = \lambda}} \frac{\lambda!}{\lambda_{a}!\lambda_{b}!} \chi_{\gamma}(\lambda_{a}) \mathbb{E}_{G}\mathbf{p}_{\lambda_{b}}$$

where $\lambda! = \prod_{i \ge 1} (\lambda(i)!)$.

We can now deal with $\mathbb{E}_G \chi_{\gamma}^G \Phi_{n,f}$. As in *Toeplitz minors* [4], absolute convergence is guaranteed by Condition (A), the bound $|\operatorname{tr}(g^i)| \leq m$ when $g \in \mathrm{U}(m)$, SO(m) or Sp(m) and compactness of those groups:

$$\mathbb{E}_G oldsymbol{\chi}^G_\gamma oldsymbol{\Phi}_{n,f} ~\leq~ \int_G \max_{g \in G} (|oldsymbol{\chi}^G_\gamma|) \exp\left(\sum_{i \geq 0} |c_i| |\operatorname{tr}(g^i)|
ight).$$

We are thus allowed to permute sums and products in the full expansion of $\Phi_{n,f}$:

$$\begin{split} \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \boldsymbol{\Phi}_{n,f} &= e^{nc_{0}} \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \exp\left(\sum_{i>0} c_{i} \mathbf{p}_{(i)}\right) \\ &= e^{nc_{0}} \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \prod_{i=1}^{\infty} \sum_{j=0}^{\infty} \frac{(c_{i} \mathbf{p}_{(i)})^{j}}{j!} \\ &= e^{nc_{0}} \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \sum_{(\alpha_{i})} \prod_{i=1}^{\infty} \frac{(c_{i} \mathbf{p}_{(i)})^{\alpha_{i}}}{\alpha_{i}!} \\ &= e^{nc_{0}} \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \sum_{(\alpha_{i})} \prod_{i=1}^{\infty} \frac{c_{i}^{\alpha_{i}}}{\alpha_{i}!} \mathbf{p}_{(i^{\alpha_{i}})} \\ &= e^{nc_{0}} \sum_{\substack{(\alpha_{i})\\\lambda:=(i^{\alpha_{i}})}} \left(\prod_{i=1}^{\infty} \frac{c_{i}^{\alpha_{i}}}{\alpha_{i}!}\right) \mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda}, \end{split}$$

From this definition of λ , we observe that $\lambda(j) = \alpha_j$, which explains the notation: $\alpha_j \ll \lambda_j$ in general.

Once $n \geq |\lambda|$, we are allowed to substitute for every term $\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \mathbf{p}_{\lambda}$ the r.h.s. of Equation (1). For a given n, this only applies for the terms at the head of the series, but any term in the series will eventually be substituted, when $n \geq |\lambda|$. Combined with absolute convergence, this guarantees the asymptotics

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \boldsymbol{\Phi}_{n,f} \overset{n \to \infty}{\sim} e^{nc_{0}} \sum_{(\alpha_{i})} \left(\left(\prod_{i=1}^{\infty} \frac{c_{i}^{\alpha_{i}}}{\alpha_{i}!} \right) \sum_{\substack{\lambda_{a} \vdash |\gamma| \\ \lambda_{a} \cup \lambda_{b} = (i^{\alpha_{i}}) =: \lambda}} \frac{\lambda!}{\lambda_{a}! \lambda_{b}!} \chi_{\gamma}(\lambda_{a}) \mathbb{E}_{G} \mathbf{p}_{\lambda_{b}} \right)$$

We now switch the sums, and change the index of one sum from (α_i) with $(i^{\alpha_i}) = \lambda$ to (β_i) with $(i^{\beta_i}) = \lambda_b$. This implies $\lambda_a(j) + \beta_j = \lambda(j) = \alpha_j$. We get

$$\mathbb{E}_{G} \boldsymbol{\chi}_{\gamma}^{G} \boldsymbol{\Phi}_{n,f} \stackrel{n \to \infty}{\sim} e^{nc_{0}} \sum_{\lambda_{a} \vdash |\gamma|} \left(\left(\frac{\chi_{\gamma}(\lambda_{a})}{\lambda_{a}!} \prod_{i=1}^{\infty} c_{i}^{\lambda_{a}(i)} \right) \sum_{(\beta_{i})} \left(\prod_{i=1}^{\infty} \frac{c_{i}^{\beta_{i}}}{\beta_{i}!} \right) \mathbb{E}_{G} \mathbf{p}_{(i^{\beta_{i}})} \right) \\
= R(\gamma, (c_{i})) \mathbb{E}_{G} \boldsymbol{\Phi}_{n,f},$$

and finally

$$\lim_{n \to \infty} \frac{\mathbb{E}_G \boldsymbol{\chi}_{\gamma}^G \boldsymbol{\Phi}_{n,f}}{\mathbb{E}_G \boldsymbol{\Phi}_{n,f}} = R(\gamma, (c_i)) = \sum_{\lambda \vdash |\gamma|} \chi_{\gamma}(\lambda) \left(\prod_{i=1}^{\infty} \frac{c_i^{\lambda(i)}}{\lambda(i)!} \right)$$

The specialization expression now follows from the usual decomposition of power polynomials into Schur polynomials given by the character table of a symmetric group (see Sagan [21, Equation (4.23)]).

Remarks.

- As mentioned earlier, this ratio $R(\gamma, (c_i))$ already appears in Theorem 6 of Bump and Diaconis [4], when G = U(n). It is striking that this ratio is independent of the Cartan type of G.
- The authors went a bit further in [4] and modified the integrand using two characters (one of them appeared conjugated). There is no real need to do this here, as the characters χ_{λ}^{G} are real in the non-unitary cases, and we would just end up with a product of two characters. Koike and Terada [14, Corollary 2.5.3] have shown that the multiplication rules are also essentially⁵

⁵This is only valid for $n \ge l(\mu) + l(\nu)$, and the case G = SO(2n) is slightly different.

independent of the Cartan type of G, i.e. that

$$oldsymbol{\chi}^G_\mu\cdotoldsymbol{\chi}^G_
u=\sum_\lambda c^\lambda_{\mu
u}oldsymbol{\chi}^G_\lambda.$$

This can be combined with Theorem 3 to show that there will also be an asymptotic ratio for $\frac{\mathbb{E}_{G} \chi_{\mu}^{G} \chi_{\nu}^{G} \Phi_{n,f}}{\mathbb{E}_{G} \Phi_{n,f}}$, independent of the Cartan type of G.

• Johansson [11, Theorem 3.8.i with $\eta = i$] was the first to generalize the strong Szegö limit theorem to all the classical groups. He found asymptotics for $\mathbb{E}_G \Phi_{n,f}$ as $n \to \infty$. Bump and Diaconis [3] later found a new proof of Johansson's result that actually inspired our own work and an extension of this result. We state here a weaker version of Johansson's result in a style closer to our own. Note that this is the first time we need Condition (B).

Theorem 4 (Johansson [11], Bump and Diaconis [3]). Let $f(t) = \sum_{i} c_i t^i$ satisfy Conditions (A) and (B) in addition to the usual symmetry condition $f(t) = f(t^{-1})$. Then

$$\mathbb{E}_{SO(2n+1)} \Phi_{n,f} = \exp\left(\sum_{i=1}^{\infty} \frac{ic_i^2}{2} - \sum_{i=1}^{\infty} c_{2i-1} + o(1)\right)$$
$$\mathbb{E}_{Sp(2n)} \Phi_{n,f} = \exp\left(\sum_{i=1}^{\infty} \frac{ic_i^2}{2} - \sum_{i=1}^{\infty} c_{2i} + o(1)\right)$$
$$\mathbb{E}_{SO(2n)} \Phi_{n,f} = \exp\left(\sum_{i=1}^{\infty} \frac{ic_i^2}{2} + \sum_{i=1}^{\infty} c_{2i} + o(1)\right)$$

We can thus combine Theorems 3 and 4 to get the asymptotics for $\mathbb{E}_{G} \chi_{\gamma}^{G} \Phi_{n,f}$, i.e. for the Haar measure twisted by a character of type χ_{λ}^{G} .

References

- Albrecht Böttcher and Bernd Silbermann. Introduction to large truncated Toeplitz matrices. Universitext. Springer-Verlag, New York, 1999.
- [2] Daniel Bump. *Lie groups*, volume 225 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2004.
- [3] Daniel Bump and Persi Diaconis. A Szegö limit theorem on the classical groups. Private communication, 4 pages.
- [4] Daniel Bump and Persi Diaconis. Toeplitz minors. J. Combin. Theory Ser. A, 97(2):252-271, 2002. Erratum for the proof of Theorem 4 available at http://sporadic.stanford.edu/bump/correction.ps.

- [5] Daniel Bump, Persi Diaconis, and Joseph B. Keller. Unitary correlations and the Fejér kernel. Math. Phys. Anal. Geom., 5(2):101–123, 2002.
- [6] Daniel Bump and Alex Gamburd. On the averages of characteristic polynomials from classical groups, math-ph/0502043. To appear in Communications in Mathematical Physics.
- [7] Paul-Olivier Dehaye. On an identity of ((Bump and Diaconis) and (Tracy and Widom)), math.CO/0601348. Submitted for publication.
- [8] Persi Diaconis and Steven N. Evans. Linear functionals of eigenvalues of random matrices. *Trans. Amer. Math. Soc.*, 353(7):2615–2633 (electronic), 2001.
- [9] Persi Diaconis and Mehrdad Shahshahani. On the eigenvalues of random matrices. J. Appl. Probab., 31A:49–62, 1994. Studies in applied probability.
- [10] Roe Goodman and Nolan R. Wallach. Representations and invariants of the classical groups, volume 68 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 1998.
- [11] Kurt Johansson. On random matrices from the compact classical groups. Ann. of Math. (2), 145(3):519–545, 1997.
- [12] Jon P. Keating and Nina C. Snaith. Random matrix theory and L-functions at s = 1/2. Comm. Math. Phys., 214(1):91–110, 2000.
- [13] Jon P. Keating and Nina C. Snaith. Random matrix theory and $\zeta(1/2 + it)$. Comm. Math. Phys., 214(1):57–89, 2000.
- [14] Kazuhiko Koike and Itaru Terada. Young-diagrammatic methods for the representation theory of the classical groups of type B_n , C_n , D_n . J. Algebra, 107(2):466–511, 1987.
- [15] Madan Lal Mehta. Random matrices. Academic Press Inc., Boston, MA, second edition, 1991.
- [16] Leonid Pastur and Vladimir Vasilchuk. On the moments of traces of matrices of classical groups. Comm. Math. Phys., 252(1-3):149–166, 2004.
- [17] Eric M. Rains. Topics in probability on compact Lie groups. PhD thesis, Harvard University, 1995.
- [18] Eric M. Rains. Increasing subsequences and the classical groups. *Electron. J. Combin.*, 5:Research Paper 12, 9 pp. (electronic), 1998.
- [19] Arun Ram. Characters of Brauer's centralizer algebras. Pacific J. Math., 169(1):173–200, 1995.
- [20] Arun Ram. A "second orthogonality relation" for characters of Brauer algebras. European J. Combin., 18(6):685–706, 1997.
- [21] Bruce E. Sagan. The symmetric group. The Wadsworth & Brooks/Cole Mathematics Series. 1991. Representations, combinatorial algorithms, and symmetric functions.
- [22] Michael Stolz. On the Diaconis-Shahshahani method in random matrix theory. J. Algebraic Combin., 22(4):471–491, 2005.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, CA *E-mail address*: pdehaye@math.stanford.edu