

DYNAMICS OF THE MAPPING CLASS GROUP ACTION ON THE VARIETY OF $\mathrm{PSL}_2 \mathbb{C}$ CHARACTERS

JUAN SOUTO AND PETER STORM

ABSTRACT. We study the action of the mapping class group $\mathrm{Mod}(S)$ on the boundary $\partial \mathcal{Q}$ of quasifuchsian space $\mathcal{Q}$. Among other results, $\mathrm{Mod}(S)$ is shown to be topologically transitive on the subset $\mathcal{C} \subset \partial \mathcal{Q}$ of manifolds without a conformally compact end. We also prove that any open subset of the character variety $\mathcal{X}(\pi_1(S), \mathrm{PSL}_2 \mathbb{C})$ intersecting $\partial \mathcal{Q}$ does not admit a nonconstant $\mathrm{Mod}(S)$ -invariant meromorphic function. This is related to a question of Goldman.

1. INTRODUCTION

Let S be a closed oriented surface of genus $g \geq 2$ and let Γ be its fundamental group. The mapping class group

$$\mathrm{Mod}(S) = \mathrm{Diff}(S) / \mathrm{Diff}_0(S) = \mathrm{Out}(\Gamma)$$

of S acts on the character variety

$$\mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C}) = \mathrm{Hom}(\Gamma, \mathrm{PSL}_2 \mathbb{C}) // \mathrm{PSL}_2 \mathbb{C}$$

by precomposition. Quasifuchsian space $\mathcal{Q}$ is the open cell in the character variety $\mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$ formed by the conjugacy classes of faithful representations with convex cocompact image. It is invariant under the mapping class group, and the action $\mathrm{Mod}(S) \curvearrowright \mathcal{Q}$ is properly discontinuous. On the other hand, our first result shows that the action $\mathrm{Mod}(S) \curvearrowright \partial \mathcal{Q}$ has very complicated dynamics.

Theorem 1. *There is a $\mathrm{Mod}(S)$ -invariant closed nowhere dense topologically perfect set $\overline{\mathcal{C}} \subset \partial \mathcal{Q}$ such that:*

1. *The points $\rho \in \partial \mathcal{Q}$ with $\overline{\mathcal{C}} \subset \overline{\mathrm{Mod}(S) \cdot \rho}$ form a dense G_δ -set.*
2. *The action of $\mathrm{Mod}(S)$ on $\overline{\mathcal{C}}$ is topologically transitive.*

In particular, Theorem 1 implies that any continuous $\mathrm{Mod}(S)$ -invariant function on $\partial \mathcal{Q}$ is constant. Recall that the action of a group on a locally compact Hausdorff separable topological space is topologically

The first author is supported by a postdoctoral fellowship of the Deutsche Forschungsgemeinschaft. The second author is supported by a National Science Foundation Postdoctoral Fellowship.

transitive if the translates of any two open sets intersect, or equivalently, if there is a dense orbit.

The group $\mathrm{PSL}_2 \mathbb{C}$ is the group of orientation preserving isometries of hyperbolic 3-space $\mathbb{H}^3$ and it is a well known fact that ρ is faithful and that the action $\rho(\Gamma) \curvearrowright \mathbb{H}^3$ is free and properly discontinuous for each $\rho \in \overline{\mathcal{Q}}$. In particular $M_\rho = \mathbb{H}^3 / \rho(\Gamma)$ is an orientable hyperbolic manifold homotopy equivalent to S . From this point of view, the set $\mathcal{C}$ provided by Theorem 1 is the set of all $\rho \in \overline{\mathcal{Q}}$ such that the boundary of the convex core of the associated hyperbolic manifold M_ρ does not have a compact component. (Recall that the convex core of a hyperbolic manifold is the smallest closed convex subset whose inclusion is a homotopy equivalence.)

A special role is played by geometrically finite representations $\rho \in \mathcal{C}$ where the boundary of the convex core of M_ρ is a collection of thrice-punctured spheres. We call these representations full maximal cusps. The proof of Theorem 1 involves studying the dynamics of the $\mathrm{Mod}(S)$ -action near these points. For example, the techniques of Theorem 1 are used to prove that for any open neighborhood $U \subset \partial \mathcal{Q}$ of a full maximal cusp, the orbit $\mathrm{Mod}(S).U$ is dense in $\partial \mathcal{Q}$ (see Corollary 9).

It is well known that two representations $\rho, \rho' \in \mathcal{Q}$ are close if the associated hyperbolic manifolds M_ρ and $M_{\rho'}$ are bi-Lipschitz with a small bi-Lipschitz constant. This is why geometric invariants of the manifold M_ρ , for example the volume of the convex core, the injectivity radius, the bottom of the Laplacian, or the Hausdorff dimension of the limit set, are continuous functions on quasifuchsian space. In each case it was previously known via different methods that these quantities are no longer continuous on $\overline{\mathcal{Q}}$. We derive from Theorem 1 a unified proof of this fact:

Theorem 2. *The volume of the convex-core, the injectivity radius, the bottom of the Laplacian and the Hausdorff dimension of the limit set do not vary continuously on $\overline{\mathcal{Q}}$.*

We also apply Theorem 1 to study $\mathrm{Mod}(S)$ -invariant meromorphic functions defined on subsets of the character variety $\mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$. In [Gol2], Goldman proved that every $\mathrm{Mod}(S)$ -invariant meromorphic functions defined on the whole of $\mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$ must be constant. This result motivates the question of which connected open subsets U of $\mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$ admit nonconstant $\mathrm{Mod}(S)$ -invariant meromorphic functions. (A weaker form of this question can be found in [Gol2, Sec.1.4].) Goldman deduced the nonexistence of invariant nonconstant meromorphic functions from the ergodicity of the $\mathrm{Mod}(S)$ -action on the (real) subvariety $\mathcal{X}(\Gamma, \mathrm{SU}_2)$ [Gol1]. In particular his result applies to every

connected open set $U \subset \mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$ containing unitary representations. We obtain the following analogue of Goldman's result:

Theorem 3. *Let $U \subset \mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$ be a $\mathrm{Mod}(S)$ -invariant connected open set. If U contains both (faithful) convex cocompact representations and indiscrete representations then any $\mathrm{Mod}(S)$ -invariant meromorphic function on U is a constant function.*

After some preliminaries in Section 2 we present in Section 3 a concrete construction of hyperbolic 3-manifolds which will be one of the key ingredients in the proof of Theorem 1. Full maximal cusps are the other key ingredient. In particular, in Section 4 we deduce that full maximal cusps are dense in $\mathcal{C}$. The proof uses techniques developed by McMullen [McM1], Canary, Culler, Hersonsky, and Shalen [CCHS, CH]. Theorem 2 is proved in Section 5 and Theorem 3 is proved in the last section.

The first author would like to thank the members of the Mathematics Department of the University of Chicago for their hospitality. The second author would like to thank Richard Canary and Misha Kapovich for helpful conversations on these topics.

2. PRELIMINARIES

We refer to [HP] for basic facts about the character variety and to [And] for a survey about the deformation theory of discrete subgroups of $\mathrm{PSL}_2 \mathbb{C}$. If H is a finitely generated non-virtually abelian torsion-free group then $\mathrm{Hom}(H, \mathrm{PSL}_2 \mathbb{C})$ is a complex algebraic variety on which $\mathrm{PSL}_2 \mathbb{C}$ acts by conjugacy. The character variety

$$\mathcal{X}(H, \mathrm{PSL}_2 \mathbb{C}) = \mathrm{Hom}(H, \mathrm{PSL}_2 \mathbb{C}) // \mathrm{PSL}_2 \mathbb{C}$$

is the quotient of $\mathrm{Hom}(H, \mathrm{PSL}_2 \mathbb{C})$ under this action in the sense of invariant theory. We remind the reader that $\mathcal{X}(H, \mathrm{PSL}_2 \mathbb{C})$ does not coincide with the set theoretic quotient $\mathrm{Hom}(H, \mathrm{PSL}_2 \mathbb{C}) / \mathrm{PSL}_2 \mathbb{C}$. However, the set of conjugacy classes of discrete faithful representations is contained in a smooth open manifold in $\mathcal{X}(H, \mathrm{PSL}_2 \mathbb{C})$ [HP, Sec.4].

Notation. *The Greek letters ρ and σ (possibly with decoration) will be used to indicate conjugacy classes of representations. Thus ρ and σ will be elements of the appropriate character variety. The notation $\rho(H)$ will indicate the image of H under any fixed homomorphism of the conjugacy class ρ .*

We identify the group $\mathrm{PSL}_2 \mathbb{C}$ with the group of orientation preserving isometries of hyperbolic 3-space $\mathbb{H}^3$. We will denote the convex core of a hyperbolic manifold M by $CC(M)$. The convergence of sequences

in $\mathcal{X}(H, \mathrm{PSL}_2 \mathbb{C})$ is said to be *algebraic convergence*. Let $\rho_i \rightarrow \rho$ be an algebraically convergent sequence. If the representations ρ_i are discrete and faithful for all i , then it is well known that ρ is discrete and faithful as well. Moreover, up to a choice of a subsequence, the groups $\rho_i(H)$ converge in the Chabauty topology to a discrete subgroup H_G of $\mathrm{PSL}_2 \mathbb{C}$ which contains the image of ρ ; H_G is the *geometric limit* of the sequence $\{\rho_i(H)\}$.

In this paper we will mainly consider discrete and faithful representations of the fundamental group Γ of a closed surface S . A faithful and discrete representation $\rho \in \mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$ induces a privileged homotopy class of homotopy equivalences $S \rightarrow M_\rho$. A theorem of Bonahon [Bon] ensures that M_ρ is homeomorphic to a trivial interval bundle over S . Moreover, the homotopy equivalence $S \rightarrow M_\rho$ determines a unique isotopy class of orientation preserving homeomorphisms $S \times (-1, 1) \rightarrow M_\rho$. In other words, M_ρ has a positive end (the top end) and a negative end (the bottom end). Observe that orientation reversing elements in $\mathrm{Mod}(S)$ extend in a canonical way to orientation preserving homeomorphisms of $S \times (-1, 1)$ which interchange the top and bottom ends.

The top (resp. bottom) end of M_ρ is *conformally compact* if there is a single compact component of $\partial CC(M_\rho)$ facing the top (resp. bottom) end of M_ρ , or equivalently if there are compact embedded convex surfaces exiting the top (resp. bottom) end of M_ρ .

Quasifuchsian space $\mathcal{Q} \subset \mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$ is the open set of conjugacy classes of faithful convex cocompact representations. The closure $\overline{\mathcal{Q}}$ of quasifuchsian space $\mathcal{Q}$ consists of faithful representations with discrete image and hence it is contained in an open submanifold of $\mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$. Let $\partial \mathcal{Q}$ denote the boundary of quasifuchsian space $\overline{\mathcal{Q}} - \mathcal{Q} \subset \mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$. Sullivan [Sul] proved that $\partial \mathcal{Q}$ is in fact the closure of $\mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C}) - \overline{\mathcal{Q}}$, i.e. $\partial \mathcal{Q}$ is the frontier of $\overline{\mathcal{Q}}$. We observe that $\rho \in \overline{\mathcal{Q}}$ lies within quasifuchsian space if and only if it has two conformally compact ends. The following $\mathrm{Mod}(S)$ -invariant subset of $\partial \mathcal{Q}$ plays a central role in the present paper:

$$\mathcal{C} = \{\rho \in \partial \mathcal{Q} \mid M_\rho \text{ has no conformally compact end}\}$$

3. THE MAIN CONSTRUCTION

In this section we present a construction which is the core of the proof of Theorem 1. The main building pieces in our construction are so called maximal cusps. Fix a compact hyperbolic surface S with fundamental group Γ .

We will say that a discrete finitely generated subgroup of $\mathrm{PSL}_2 \mathbb{C}$ is a *full maximal cusp* if it is geometrically finite and every component of the boundary of the convex core of the associated hyperbolic manifold is a thrice punctured sphere. (These groups are often referred to simply as a “maximal cusps”. The word “full” has been added for emphasis.) Observe that $\rho \in \mathcal{C}$ if $\rho(\Gamma)$ is a full maximal cusp. We will say a representation $\rho \in \partial \mathcal{Q}$ is a *one sided maximal cusp* if it is geometrically finite, it has one conformally compact end, and each component of $\partial CC(M_\rho)$ facing the end of M_ρ which is not conformally compact is a thrice punctured sphere.

Maximal cusps are very convenient when making concrete constructions because any pair of totally geodesic thrice punctured spheres in any pair of hyperbolic 3-manifolds are isometric. In particular, if M and M' are hyperbolic manifolds whose convex-cores boundaries $\partial CC(M), \partial CC(M')$ contain thrice punctured spheres X and X' and $\phi : X \rightarrow X'$ is a homeomorphism, then ϕ is isotopic to an isometry which we denote again by ϕ . Hence there is a hyperbolic manifold N which is covered by M and M' , and whose convex core is isometric to $CC(M) \cup_\phi CC(M')$.

The second main ingredient in our constructions is the following lemma, which is a consequence of Thurston’s Dehn-filling Theorem [Thu1, Ch.4][BO, Com].

Lemma 4. *Let $H < \mathrm{PSL}_2 \mathbb{C}$ be a geometrically finite group such that $\mathbb{H}^3/H$ is homeomorphic to $S \times (0, 1) - \mathcal{P}$ where $\mathcal{P}$ is an unlinked collection of disjoint simple closed curves. Then the group H is the geometric limit of geometrically finite groups H_n isomorphic to $\pi_1(S)$. Moreover, if H is a full (resp. one sided) maximal cusp, then the H_n can be chosen to be full (resp. one sided) maximal cusps. $\square$*

Recall that a collection of disjoint simple closed curves in $S \times [0, 1]$ is unlinked if every curve is contained in an embedded boundary parallel surface which is disjoint from all the other curves. The groups H_n in the statement of Lemma 4 are obtained by performing a hyperbolic $(1, n)$ -Dehn surgery on each rank two cusp of H .

Lemma 5. *Let $H, H_n < \mathrm{PSL}_2 \mathbb{C}$ be as in the statement of Lemma 4, $\rho_n : \Gamma \rightarrow H_n$ isomorphisms and $\rho \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$ be discrete and faithful such that $\rho(\Gamma) < H$ is the fundamental group of an embedded level surface*

$$S \times \{t\} \subset S \times (0, 1) - \mathcal{P} \cong \mathbb{H}^3/H.$$

Then there are automorphisms α_n of Γ such that $\rho_n \circ \alpha_n$ converges algebraically to ρ .

Proof. Since $H_n \longrightarrow H$ geometrically, for any a compact submanifold $K \subset \mathbb{H}^3/H$ there is a sequence of smooth embeddings $\phi_n : K \longrightarrow \mathbb{H}^3/H_n$ which converge in the $\mathcal{C}^\infty$ -topology to isometric embeddings [McM2, Sec.2.2]. Since $\mathbb{H}^3/H_n$ is obtained by $(1, n)$ -Dehn surgery on $\mathbb{H}^3/H$, it is clear that by choosing a sufficiently large compact submanifold K the restricted homomorphism

$$(\phi_{n*})|_{\rho(\Gamma)} : \rho(\Gamma) \longrightarrow H_n < \mathrm{PSL}_2 \mathbb{C}$$

will be well-defined and injective, and thus an isomorphism. Since the maps ϕ_n are converging to isometries, the sequence of homomorphisms $(\phi_{n*})|_{\rho(\Gamma)}$ is converging to the identity map. Therefore

$$\phi_{n*} \circ \rho : \Gamma \longrightarrow H_n$$

is a sequence of discrete faithful representations converging to ρ . Define α_n to be $\rho_n^{-1} \circ \phi_{n*} \circ \rho$. This proves the lemma. $\square$

The following is the main result of the present section:

Proposition 6. *Let $\rho, \bar{\rho} \in \partial \mathcal{Q} - \mathcal{C}$ be one sided maximal cusps and $\rho^{\mathcal{C}}, \bar{\rho}^{\mathcal{C}} \in \mathcal{C}$ full maximal cusps. There exist sequences of representations $\{\rho_i^{\mathcal{Q}}\}_i$ in quasifuchsian space $\mathcal{Q}$, one sided maximal cusps $\{\rho_i^{\mathcal{Q}-\mathcal{C}}\}_i$ in $\mathcal{Q} - \mathcal{C}$, full maximal cusps $\{\rho_i^{\mathcal{C}}\}_i$ in $\mathcal{C}$, and sequences of mapping classes $\{\alpha_i\}$, $\{\beta_i\}$ and $\{\gamma_i\} \subset \mathrm{Mod}(S)$ with:*

1. $\rho = \lim_i \rho_i^{\mathcal{Q}}$ and $\bar{\rho} = \lim_i \alpha_i \cdot \rho_i^{\mathcal{Q}}$.
2. $\rho = \lim_i \rho_i^{\mathcal{Q}-\mathcal{C}}$ and $\rho^{\mathcal{C}} = \lim_i \beta_i \cdot \rho_i^{\mathcal{Q}-\mathcal{C}}$.
3. $\rho^{\mathcal{C}} = \lim_i \rho_i^{\mathcal{C}}$ and $\bar{\rho}^{\mathcal{C}} = \lim_i \gamma_i \cdot \rho_i^{\mathcal{C}}$.


Proof. The proofs of all three claims of Proposition 6 are very similar, so we will show only the first one. Moreover, in order to simplify notation we will denote the sequence in $\mathcal{Q}$ which we are going to construct by $\{\rho_i\}$ instead of $\{\rho_i^{\mathcal{Q}}\}$. Moreover, up to composition with an orientation reversing element in $\mathrm{Mod}(S)$, we may assume that the top end of M_ρ and the bottom end of $M_{\bar{\rho}}$ are conformally compact.

By Lemma 5 it suffices to construct a hyperbolic manifold N homeomorphic to the complement in $S \times (0, 1)$ of an unlinked collection of simple curves, which is (locally isometrically) covered by both M_ρ and $M_{\bar{\rho}}$.

The convex cores $CC(M_\rho)$ and $CC(M_{\bar{\rho}})$ are by definition homeomorphic to

$$(S \times [-1, 1]) - (\mathcal{P}_T \times \{-1\}) \text{ and } (S \times [-1, 1]) - (\mathcal{P}_B \times \{1\}),$$

where $\mathcal{P}_T$ and $\mathcal{P}_B$ are pants decompositions of S . We choose sufficiently complicated homeomorphisms $\phi_T : S \rightarrow S$ and $\phi_B : S \rightarrow S$ such that the two collections $\phi_T(\mathcal{P}_T)$ and $\phi_B(\mathcal{P}_B)$ bind the surface S . Thurston's

FIGURE 1. constructing the manifold N

hyperbolization theorem [Ota] implies that there is a geometrically finite hyperbolic 3-manifold $S \rightarrow M$ whose convex core is homeomorphic to

$$(S \times [-1, 1]) - (\phi_B(\mathcal{P}_B) \times \{-1\} \cup \phi_T(\mathcal{P}_T) \times \{1\}).$$

The middle manifold M is a full maximal cusp. M_ρ and $M_{\bar{\rho}}$ are one sided maximal cusps. Therefore the convex cores of M_ρ , $M_{\bar{\rho}}$ and M can be glued together to obtain the convex core $CC(N)$ of a hyperbolic 3-manifold N covered by M_ρ and $M_{\bar{\rho}}$ and homeomorphic to

$$(S \times [-3, 3]) - (\phi_B(\mathcal{P}_B) \times \{-1\} \cup \phi_T(\mathcal{P}_T) \times \{1\}).$$

Two applications of Lemma 5 concludes the proof of Proposition 6. $\square$

4. TOPOLOGICAL DYNAMIC OF $\text{Mod}(S) \curvearrowright \partial\mathcal{Q}$

In this section we derive Theorem 1 from the results in the last section and the ubiquity of maximal cusps in $\partial\mathcal{Q}$. We will use techniques developed by McMullen [McM1], Canary-Culler-Hersonsky-Shalen [CCHS], and Canary-Hersonsky [CH]. We begin by using recent work [BBES, CCHS] to generalize a theorem of McMullen's.

Theorem 7. (McMullen [McM1]) *One sided maximal cusps form a dense subset of $\partial\mathcal{Q}$.*

Proof. A standard Baire category argument shows that the set of representations $\rho \in \partial\mathcal{Q}$ without parabolic elements forms a dense subset of $\partial\mathcal{Q}$ (see [McM1, Cor1.5] or [CCHS, Lem15.2]). If the limit set of such a representation ρ is the entire sphere at infinity, then ρ is an algebraic limit of full maximal cusps [CCHS, Thm6.1]. Proposition 6(2) shows that a full maximal cusp is always an algebraic limit of one sided maximal cusps in $\partial\mathcal{Q} - \mathcal{C}$. Therefore it only remains to consider the case where M_ρ has a conformally compact end.

Let us assume without a loss of generality that the top end of M_ρ is conformally compact. By [BBES], the representation $\rho \in \partial\mathcal{Q}$ is a strong limit of quasiconformal representations ρ_i . By [KT], the conformal structure at infinity of the top end of M_ρ must be the limit in Teichmüller space of the conformal structures at infinity of the top ends of the M_{ρ_i} . Therefore altering the manifolds M_{ρ_i} by a K_i -bi-Lipschitz deformation, where $K_i \rightarrow 1$, produces a sequence of quasifuchsian manifolds in a Bers slice converging strongly to M_ρ . We may now apply [McM1] to conclude that ρ is an algebraic limit of one sided maximal cusps. $\square$

From Proposition 6 and Theorem 7 we derive the following ostensibly stronger result:

Proposition 8. *In every open neighborhood of $\partial\mathcal{Q}$ we find a $\bar{\rho}$ with the following property: For every full maximal cusp $\rho^{\mathcal{C}}$ there is a sequence $\{\alpha_i\}$ in $\text{Mod}(S)$ with*

$$\rho^{\mathcal{C}} = \lim_i \alpha_i \cdot \bar{\rho}$$

Proof. Observe that the set of all full maximal cusps is countable. In particular there is a sequence of full maximal cusps $\{\rho_j\} \subset \mathcal{C}$ such that every full maximal cusp appears infinitely often in the sequence.

Moreover, we can fix open neighborhoods V_j of ρ_j in $\partial\mathcal{Q}$ such that when a subsequence ρ_{j_i} converges then for every choice of $\rho'_{j_i} \in V_{j_i}$ the sequence ρ'_{j_i} also converges to the same limit.

Given an open neighborhood U_1 in $\partial\mathcal{Q}$ we obtain from Proposition 6(2) and Theorem 7 a one sided maximal cusp $\sigma_1 \in U_1 \subset \partial\mathcal{Q}$ and $\alpha_1 \in \text{Mod}(S)$ with $\alpha_1 \cdot \sigma_1 \in V_1$. Let U_2 be an open and relatively compact neighborhood of σ_1 in $U \cap \alpha_1^{-1}(V_1)$. Applying again the same argument we find again a one sided maximal cusp $\sigma_2 \in U_2$ and $\alpha_2 \in \text{Mod}(S)$ with $\alpha_2 \cdot \sigma_2 \in V_2$. Let again U_3 be an open and relatively compact neighborhood of σ_2 in $U_2 \cap \alpha_2^{-1}(V_2)$. Inductively we find sequences of one sided maximal cusps $\{\sigma_i\}$, $\{\alpha_i\}$ in $\text{Mod}(S)$ and open sets $\{U_i\}$ with U_{i+1} relatively compact in U_i , $\sigma_i \in U_i$, and $\alpha_i(U_i) \subset V_i$. By our choice of the neighborhoods V_i we derive that every $\bar{\rho}$ in $\cap_i U_i$ has the desired property. $\square$

By reversing the logic of Proposition 8 we obtain

Corollary 9. *If $W \subset \partial\mathcal{Q}$ is an open set containing a full maximal cusp then $\text{Mod}(S) \cdot W$ is a dense open subset of $\partial\mathcal{Q}$. Moreover, the set*

$$\{\rho \in \partial\mathcal{Q} \mid \overline{\text{Mod}(S) \cdot \rho} \text{ contains every full maximal cusp}\}$$

is a dense G_δ -set.

Proof. The first sentence of the corollary follows immediately from Proposition 8. To prove the second, let us reuse the notation $\{\rho_j\}$ and $\{V_j\}$ from the proof of Proposition 8. Then

$$\bigcap_j \text{Mod}(S) \cdot V_j = \{\rho \in \partial\mathcal{Q} \mid \overline{\text{Mod}(S) \cdot \rho} \text{ contains every full maximal cusp}\}.$$

From this the corollary follows. $\square$

Recall that $\mathcal{C} \subset \partial\mathcal{Q}$ is the set of manifolds without a conformally compact end. Below we show:

Lemma 10. *Full maximal cusps are dense in $\mathcal{C}$.*

Before showing Lemma 10 we will prove Theorem 1:

Theorem 1. *The closure $\overline{\mathcal{C}}$ of $\mathcal{C}$ is a $\text{Mod}(S)$ -invariant nowhere dense topologically perfect set such that:*

1. *The points $\rho \in \partial\mathcal{Q}$ with $\overline{\mathcal{C}} \subset \overline{\text{Mod}(S) \cdot \rho}$ form a dense G_δ -set.*
2. *The action of $\text{Mod}(S)$ on $\overline{\mathcal{C}}$ is topologically transitive.*

Proof. The set $\mathcal{C}$ is $\text{Mod}(S)$ -invariant by definition, implying the same for its closure $\overline{\mathcal{C}}$. The set $\mathcal{C}$ is topologically perfect since full maximal cusps are dense in $\mathcal{C}$ and Proposition 6(3) implies full maximal cusps are not isolated. Claim (1) follows immediately from Lemma 10 and Corollary 9.

We now prove claim (2). Let U^1 and U^2 be open sets in $\mathcal{C}$. By Lemma 10 there are full maximal cusps $\rho^1 \in U^1$ and $\rho^2 \in U^2$ and by Proposition 6 there exist sequences $\{\rho_i\}$ in $\mathcal{C}$ and $\{\alpha_i\}$ in $\text{Mod}(S)$ such that

$$\rho^1 = \lim_i \rho_i \quad \text{and} \quad \rho^2 = \lim_i \alpha_i \cdot \rho_i$$

This shows that for sufficiently large i , $(\alpha_i \cdot U^2) \cap U^1$ is not empty and hence that the action of $\text{Mod}(S)$ on $\mathcal{C}$ is topologically transitive. This implies topological transitivity on $\overline{\mathcal{C}}$.

It remains to prove that $\overline{\mathcal{C}}$ is nowhere dense. Pick a representation $\sigma \in \partial\mathcal{Q} - \mathcal{C}$ with no parabolic elements and exactly one conformally compact end (i.e. a singly degenerate representation without parabolics). Suppose that $\sigma \in \overline{\mathcal{C}}$. Then by Lemma 10 there is a sequence of full maximal cusps $\tau_i \rightarrow \sigma$. Since σ has no parabolic elements, it follows from a theorem of Evans that the manifolds M_{τ_i} converge geometrically to M_σ [Ev]. Since the top end of M_σ is conformally compact, there is a strictly convex surface Σ embedded in the top end of M_σ . Use the almost isometric embeddings provided by geometric convergence to push the surface Σ into M_{τ_i} . For sufficiently large i this yields a strictly convex embedded surface in M_{τ_i} , showing that the manifolds M_{τ_i} eventually have a conformally compact end. Since they are all full maximal cusps, this is a contradiction. Therefore σ is not in $\overline{\mathcal{C}}$.

Let U be a neighborhood of σ in the complement of $\overline{\mathcal{C}}$. By Proposition 8 there is a representation in U whose orbit limits onto all of $\mathcal{C}$. As $\text{Mod}(S) \cdot U \cap \overline{\mathcal{C}} = \emptyset$, this proves that $\overline{\mathcal{C}}$ is nowhere dense. $\square$

It remains only to prove Lemma 10:

Proof of Lemma 10. We divide the proof of Lemma 10 in four cases.

Case 1: Assume ρ is geometrically finite.

By Lemma 5, it suffices to show that there is a hyperbolic manifold covered by M_ρ which is the geometric limit of a sequence of full maximal cusps in $\mathcal{C}$.

The convex core $CC(M_\rho)$ of M_ρ is homeomorphic to

$$(S \times [-1, 1]) - (\mathcal{P}_- \times \{-1\} \cup \mathcal{P}_+ \times \{1\}),$$

where $\mathcal{P}_\pm$ are nonempty collections of disjoint, nonparallel simple closed curves. We consider the embedding

$$\begin{aligned} (S \times [-1, 1]) - (\mathcal{P}_- \times \{-1\} \cup \mathcal{P}_+ \times \{1\}) &\longrightarrow \\ (S \times [-1, 2]) - (\mathcal{P}_- \times \{-1\} \cup \mathcal{P}_+ \times \{1\}) &=: N \end{aligned}$$

and the induced morphism

$$\Pi : \mathcal{X}(\pi_1(N), \mathrm{PSL}_2 \mathbb{C}) \rightarrow \mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$$

By [BBES, Lem.5.1] there is an open set in $\Pi^{-1}(\rho)$ which consists of discrete and faithful representations. Moreover, the marked conformal structure at infinity corresponding to the bottom end of any manifold in the fiber $\Pi^{-1}(\rho)$ is equivalent to the marked conformal structure at infinity corresponding to the bottom end of M_ρ .

On the other hand, by deforming the added generator in any $\mathbb{Z} \times \mathbb{Z}$ subgroup of $\pi_1(N)$, we see that every connected component of $\Pi^{-1}(\rho)$ contains indiscrete representations. For a generic $\sigma \in \Pi^{-1}(\rho)$ the only parabolic elements in $\sigma(\pi_1(N))$ correspond to curves in N freely homotopic into

$$(\mathcal{P}_- \times \{-1\} \cup \mathcal{P}_+ \times \{1\}) \subset N,$$

i.e. a generic point is minimally parabolic. (This is again a Baire category argument; compare with [McM1, Cor1.5] or [CCHS, Lem15.2].) We deduce that there is a discrete and faithful $\sigma \in \Pi^{-1}(\rho)$ such that the top end of M_σ has empty conformal boundary.

Now would now like to apply [CH, Thm.9.1] to conclude that σ is the algebraic limit of a sequence σ_i in $\mathcal{X}(\pi_1(N), \mathrm{PSL}_2 \mathbb{C})$ such σ_i is discrete, faithful, and the top end of M_{σ_i} is maximally cusped. The only technicality is verifying condition 2 of [CH, Thm.9.1] (finding short pants decompositions). This can be accomplished by using the proof of [CCHS, Prop.4.1], where the application of Sullivan rigidity goes through unchanged because the conformal structure at infinity corresponding to the bottom end of any manifold in $\Pi^{-1}(\rho)$ is equivalent to the conformal structure at infinity of the bottom end of M_ρ . Therefore [CH, Thm.9.1] can be used.

Since M_σ is minimally parabolic, we may apply a theorem of Evans [Ev] to conclude that the manifolds M_{σ_i} also converge geometrically to M_σ . Lemma 4 implies now that for all i the manifold M_{σ_i} is the geometric limit of a sequence M_i^j of geometrically finite hyperbolic manifolds homeomorphic to $S \times (0, 1)$ such that the top end of each M_i^j is maximally cusped. Taking a diagonal sequence we deduce that

M_σ is the geometric limit of a sequence of geometrically finite manifolds whose top ends are maximally cusped. Since M_σ is covered by M_ρ , we may apply Lemma 5 to conclude that

M_ρ is an algebraic limit of geometrically finite manifolds in $\mathcal{C}$ whose top ends are maximally cusped.

Therefore, we may without a loss of generality assume the the top end of M_ρ is maximally cusped. Now apply the above argument again to M_ρ , swapping the roles of the bottom end and the top end. This yields a sequence of full maximal cusps converging algebraically to M_ρ . This completes the proof of *Step 1*.

Case 2: M_ρ has no cusps.

In this case M_ρ must have an empty domain of discontinuity. We may then directly apply [CH, Thm.10.1] to conclude that M_ρ is the algebraic limit of a sequence of full maximal cusps.

Case 3: M_ρ has at least one rank one cusp in each end.

By *Step 1*, it suffices to approximate M_ρ by geometrically finite manifolds in $\mathcal{C}$. Let P^T (resp. P^B) be a nonempty collection of disjoint essential annuli on S such that there is a relative homeomorphism

$$(S \times (0, 1), (P^T \times [.8, 1)) \cup (P^B \times (0, .2])) \longrightarrow (M_\rho, \text{cusps of } M_\rho).$$

By [BBES, Cor.3.2], M_ρ is the algebraic and geometric limit of a sequence of geometrically finite manifolds M_{ρ_i} such that for each i there is a relative *homotopy equivalence*

$$(S \times (0, 1), (P^T \times [.8, 1)) \cup (P^B \times (0, .2])) \longrightarrow (M_{\rho_i}, \text{cusps of } M_{\rho_i}).$$

To see that the manifolds M_{ρ_i} are in $\mathcal{C} \subset \mathcal{Q}$, it suffices to show that the rank one cusps cannot flip ends in the limit. This follows from the fact that M_ρ is both the algebraic and the geometric limit of the M_{ρ_i} .

Case 4: M_ρ has all of its rank one cusps in exactly one end.

Assume without a loss of generality that M_ρ has rank one cusps in its bottom end. Then the top end of M_ρ is necessarily geometrically infinite. The proof begins as in Case 3. Let P^B be a nonempty collection of disjoint essential annuli on S such that there is a relative homeomorphism

$$(S \times (0, 1), P^B \times (0, .2]) \longrightarrow (M_\rho, \text{cusps of } M_\rho).$$

As above, applying [BBES, Cor.3.2] yields that M_ρ is the algebraic and geometric limit of a sequence of geometrically finite manifolds M_{ρ_i} such that for each i there is a relative homotopy equivalence

$$(S \times (0, 1), P^B \times (0, .2]) \longrightarrow (M_{\rho_i}, \text{cusps of } M_{\rho_i}).$$

Again using the fact that M_{ρ_i} converges both algebraically and geometrically to M_ρ , after possibly passing to a subsequence we may assume these relative homotopy equivalences are homotopic to relative homeomorphisms.

Let $\mathcal{E}$ denote the geometrically finite ends of $M_\rho - \{\text{cusps}\}$. As in the proof of Theorem 7, we can deform each M_{ρ_i} by a K_i -bi-Lipschitz diffeomorphism, where $K_i \rightarrow 1$, to obtain constant conformal structures at infinity for the $\mathcal{E}$ -ends of the deformed $M_{\rho'_i} - \{\text{cusps}\}$. (Which by geometric convergence means that these conformal structures are equal to the conformal structure at infinity for the $\mathcal{E}$ -ends of $M_\rho - \{\text{cusps}\}$.) This allows us to apply the proof of [CCHS, Prop.4.1] to obtain “increasingly short pinchable pants decompositions.” We may then apply [CH, Thm.9.1] to find an M_ρ convergent sequence of geometrically finite manifolds in $\mathcal{C}$ whose top ends are maximally cusped. This reduces the proof to Step 1. $\square$

5. GEOMETRIC INVARIANTS NEVER VARY CONTINUOUSLY

In this section we quickly prove Theorem 2.

Theorem 2. *The volume of the convex-core, the injectivity radius, the bottom of the Laplacian and the Hausdorff dimension of the limit set do not vary continuously on $\overline{\mathcal{Q}}$.*

Each part of Theorem 2 was known previously. We present merely a unified proof.

Proof. All these invariants are $\text{Mod}(S)$ -invariant functions $\partial\mathcal{Q} \rightarrow \mathbb{R} \cup \{\infty\}$. In particular we derive from Theorem 1 (3) and (4) that they either are constant or fail to be continuous. In particular, it suffices to find $\rho, \rho' \in \partial\mathcal{Q}$ for which they do not take the same value. Let ρ be a full maximal cusp and ρ' the cyclic cover of the mapping torus of a pseudo-Anosov $\phi \in \text{Mod}(S)$; ρ' is in the boundary of quasifuchsian space by the work of Thurston [Thu2]. We know that

$$\text{vol}(CC(M_\rho)) < \infty \text{ and } \text{vol}(CC(M_{\rho'})) = \infty$$

because ρ is geometrically finite and ρ' is not. Geometric finiteness of ρ implies that Λ_ρ has Hausdorff dimension less than 2 while $\Lambda_{\rho'} = \mathbb{C}P^1$ has dimension 2. The relation between the dimension of the limit set and the bottom of the Laplacian shows that these two quantities also differ [CMT]. The manifold $M_{\rho'}$ covers a compact manifold and hence has positive injectivity radius while M_ρ has cusps and hence its injectivity radius vanishes. $\square$

From this we see that it is in fact impossible to define a nontrivial purely geometric invariant which varies continuously.

6. $\text{MAP}(S)$ -INVARIANT MEROMORPHIC FUNCTIONS

In this section we prove Theorem 3. As a warm up we first consider meromorphic functions on open subsets of $\mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$.

Theorem 11. *Let $U \subset \mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$ be a $\text{Mod}(S)$ -invariant connected open set. If U contains both (faithful) convex cocompact representations and indiscrete representations then any $\text{Mod}(S)$ -invariant meromorphic function on U is a constant function.*

Note that indiscrete representations are dense in the complement of quasifuchsian space.

Proof. Without a loss of generality we may assume that U is a manifold. The set U intersects the interior and the exterior of $\overline{\mathcal{Q}}$ and hence $\partial\mathcal{Q} \cap U \neq \emptyset$. We claim that every continuous $\text{Mod}(S)$ -invariant function $f : U \rightarrow \mathbb{C}$ is constant on $\partial\mathcal{Q} \cap U$. Given full maximal cusps $\rho, \bar{\rho} \in \partial\mathcal{Q}$ we obtain from Proposition 6 a sequence $\{\rho_i\} \subset \mathcal{Q}$ and a sequence of mapping classes $\{\alpha_i\} \subset \text{Mod}(S)$ with

$$\lim_i \rho_i = \rho \text{ and } \lim_i \alpha_i \cdot \rho_i = \bar{\rho}.$$

We have $\rho_i \in U$ for sufficiently large i . The continuity and the $\text{Mod}(S)$ -invariance of f imply that $f(\rho) = f(\bar{\rho})$. Since maximal cusps are dense in $\partial\mathcal{Q} \cap U$, the claim follows.

Assume now that $f : U \rightarrow \mathbb{C}$ is meromorphic and $\text{Mod}(S)$ -invariant. The divisor D of poles of f is either empty or has complex co-dimension 1. The open set $U - D$ is open, connected, $\text{Mod}(S)$ -invariant and intersects again $\partial\mathcal{Q}$. In particular we may assume without loss of generality that f is holomorphic. As we have proved above f is constant on the separating set $\partial\mathcal{Q} \cap U$. This implies that f is everywhere constant. $\square$

We discuss now the relation between the character varieties $\mathcal{X}(\Gamma, \text{SL}_2 \mathbb{C})$ and $\mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$ and prove Theorem 3. The homomorphism $\text{SL}_2 \mathbb{C} \rightarrow \text{PSL}_2 \mathbb{C}$ induces a map

$$p : \mathcal{X}(\Gamma, \text{SL}_2 \mathbb{C}) \rightarrow \mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$$

which maps onto the connected component of $\mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$ containing $\overline{\mathcal{Q}}$ [HP, Rmk.4.3][Cu]. Recall that the closure of quasifuchsian space $\overline{\mathcal{Q}} \subset \mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$ is contained in a smooth open $\text{Mod}(S)$ -invariant manifold $\mathcal{O} \subset \mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C})$ [HP, Sec.4][Gol2]. After possibly shrinking $\mathcal{O}$ slightly we may assume that

$$p|_{p^{-1}(\mathcal{O})} : p^{-1}(\mathcal{O}) \rightarrow \mathcal{O}$$

is a Galois cover with covering transformation group $H_1(S, \mathbb{Z}/2\mathbb{Z})$ [HP]. See Goldman [Gol2] for properties of $\mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$.

Theorem 3. *Let $U \subset \mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$ be a $\mathrm{Mod}(S)$ -invariant connected open set. If U contains both (faithful) convex cocompact representations and indiscrete representations then any $\mathrm{Mod}(S)$ -invariant meromorphic function on U is a constant function.*

Again note that indiscrete representations are dense in the complement of the set of convex cocompact representations.

Proof. As above we may assume without loss of generality that U is a manifold contained in $\mathcal{O}$. Moreover, it suffices to show that a holomorphic $\mathrm{Mod}(S)$ -invariant function f on U is constant.

The conditions on U imply that there is some $\sigma \in \partial \mathcal{Q} \cap p(U)$ since the image under p of a faithful convex cocompact representation lies in $\mathcal{Q}$ and the image of an indiscrete representation is again indiscrete. Choose a neighborhood V of σ and $V' \subset U$ open such that $p|_{V'} : V' \rightarrow V$ is a homeomorphism. Since the actions of $\mathrm{Mod}(S)$ on $\mathcal{X}(\Gamma, \mathrm{SL}_2 \mathbb{C})$ and $\mathcal{X}(\Gamma, \mathrm{PSL}_2 \mathbb{C})$ are compatible we obtain that the restriction of p to the open $\mathrm{Mod}(S)$ -invariant set

$$U' = \cup_{\alpha \in \mathrm{Mod}(S)} \alpha V'$$

is a covering. It suffices to show that the restriction of f to U' is constant. Given $n \in \mathbb{N}$ we consider the holomorphic function on $p(U')$ given by

$$S_n(z) = \sum_{\{z' \in U' \mid p(z')=z\}} f^n(z').$$

We claim that S_n is constant for all n . The function S_n is holomorphic and $\mathrm{Mod}(S)$ -invariant, but we cannot directly apply Theorem 11 since $p(U')$ may not be connected. However, every connected component of $p(U')$ intersects $\partial \mathcal{Q}$ and therefore the same proof as in Theorem 11 applies to show that S_n is locally constant.

The following lemma shows that S_n being locally constant for all n implies that f itself is constant on U' .

Lemma 12. *Let $\Omega \subset \mathbb{C}^r$ be a connected open set and $f_1, \dots, f_k : \Omega \rightarrow \mathbb{C}$ holomorphic such that for all n the function $S_n(z) = \sum f_i(z)^n$ is constant, then all the functions f_i are constant.*

Proof. Before going further observe that it suffices to consider the case $r = 1$, because a nonconstant holomorphic function will have a non-constant restriction to some holomorphic disk. Seeking a contradiction assume that the lemma fails to be true. One can assume without loss of generality that none of the functions f_i are constant and that no

two of the functions f_i and f_j are proportional. We may further assume, up to reducing Ω , that for all i and z we have $f_i(z) \neq 0$ and $df_i(z) \neq 0$. Moreover, the assumption that no two of the functions are proportional implies that, up to relabeling, there is some z_0 satisfying $|f_1(z_0)| > |f_2(z_0)| \geq \cdots \geq |f_k(z_0)|$. Multiplying by a suitable scalar we may assume that $f_1(z_0) = 1$. Computing the derivative of S_n at the point 0 we obtain the following identity for all n :

$$0 = dS_n(z_0) = n(df_1(z_0) + f_2(z_0)^{n-1}df_2(z_0) + \cdots + f_k(z_0)^{n-1}df_k(z_0))$$

Dividing by n and taking a limit $n \rightarrow \infty$ we derive that $df_1(z_0) = 0$. This is a contradiction. $\square$

As mentioned above this concludes the proof of Theorem 3 $\square$

REFERENCES

- [And] J. Anderson, A brief survey of the deformation theory of Kleinian groups, in *Geometry and Topology Monographs*, Volume 1: The Epstein Birthday Schrift (1998), 23 - 49.
- [Bon] F. Bonahon, Bouts des variétés hyperboliques de dimension 3, *Ann. of Math.* 124 (1986), 71-158.
- [BO] F. Bonahon, J.P. Otal, Variétés hyperboliques à géodésiques arbitrairement courtes, *Bull. London Math. Soc.* 20 (1988), 255-261.
- [BBES] J. Brock, K. Bromberg, R. Evans and J. Souto, Tameness on the boundary and Ahlfors' measure conjecture, *Pub. Math. IHES* 2003.
- [CCHS] R. Canary, M. Culler, S. Hersonsky and P. Shalen, Approximation by maximal cusps in boundaries of deformation spaces of Kleinian groups, *J. of Diff. Geom.*, vol.64 (2003), 57-109.
- [CH] R. Canary and S. Hersonsky, Ubiquity of geometric finiteness in boundaries of deformation spaces of hyperbolic 3-manifolds, to appear in the *American Journal of Mathematics*.
- [CMT] R. Canary, Y. Minsky and E. Taylor, Spectral theory, Hausdorff dimension and the topology of hyperbolic 3-manifolds, *J. Geom. Analysis*.
- [Com] T. Comar, Hyperbolic Dehn surgery and convergence of Kleinian groups, PhD thesis, University of Michigan (1996).
- [Cu] M. Culler, Lifting representations to covering groups, *Adv. in Math.* 59(1986), 64-70.
- [Ev] R. Evans, Weakly Type-Preserving Sequences and Strong Convergence, *Geometriae Dedicata*, vol.108, no.1 (2004), 71-92.
- [Gol1] W. Goldman, Ergodic theory on moduli spaces, *Ann of Math.* 146 (1997), 1-33.
- [Gol2] W. Goldman, The complex-symplectic geometry of $SL(2, \mathbb{C})$ -characters over surfaces, preprint 2003.
- [HP] H. Heusener and J. Porti, The variety of characters in $PSL_2 \mathbb{C}$, preprint 2003.
- [KT] S. Kerckhoff and W. Thurston, Noncontinuity of the action of the modular group at Bers' boundary of Teichmüller space, *Invent. Math.* 100 (1990), no. 1, 25-47.

- [McM1] C. McMullen, Cusps are dense, *Ann. of Math.* (2) 133 (1991), no. 1, 217-247.
- [McM2] C. McMullen, *Renormalization and 3-Manifolds Which Fiber Over the Circle*, *Annals of Math Studies* 142, Princeton University Press, 1996.
- [Ota] J.-P. Otal, Thurston's hyperbolization of Haken manifolds, *Surveys in differential geometry*, Vol. III (1998), 77-194.
- [Sul] D. Sullivan, Quasiconformal homeomorphisms and dynamics II: Structural stability implies hyperbolicity of Kleinian groups, *Acta Math.* 155 (1985), 243-260.
- [Thu1] W. Thurston, The topology and geometry of 3-manifolds. Available from the MSRI website www.msri.org, 1976-1979. Princeton Univ. lecture notes.
- [Thu2] W. Thurston, Hyperbolic Structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle. preprint.