GENERALIZED CLUSTER COMPLEXES AND COXETER COMBINATORICS

SERGEY FOMIN AND NATHAN READING

ABSTRACT. We introduce and study a family of simplicial complexes associated to an arbitrary finite root system and a nonnegative integer parameter m. For m=1, our construction specializes to the (simplicial) generalized associahedra or, equivalently, to the cluster complexes for the cluster algebras of finite type.

Our computation of the face numbers and h-vectors of these complexes produces the enumerative invariants defined in other contexts by C. A. Athanasiadis, suggesting links to a host of well studied problems in algebraic combinatorics of finite Coxeter groups, root systems, and hyperplane arrangements.

Recurrences satisfied by the face numbers of our complexes lead to combinatorial algorithms for determining Coxeter-theoretic invariants. That is, starting with a Coxeter diagram of a finite Coxeter group, one can compute the Coxeter number, the exponents, and other classical invariants by a recursive procedure that only uses most basic graph-theoretic concepts applied to the input diagram.

Contents

1.	Introduction	6
2.	Cluster complexes of finite type	Ģ
3.	Generalized cluster complexes	4
4.	Examples	6
5.	The classical types	8
	Proof of Theorem 3.4	14
7.	Proof of Theorem 3.7	15
8.	Face numbers	17
9.	Proof of Theorem 8.5	21
10.	h-vectors	24
11.	Euler characteristic	25
12.	Reciprocal face numbers	27
	Combinatorics of Coxeter-theoretic invariants	28
14.	Fake Coxeter invariants	36
Ack	knowledgments	38
	ferences	39

Date: May 4, 2005. Revised July 25, 2005.

²⁰⁰⁰ Mathematics Subject Classification. Primary 20F55, Secondary 05A15, 52B05, 55U10.

Key words and phrases. Root system, associahedron, cluster complex, Coxeter group, Coxeter number, polygon dissections, Catalan numbers, Fuss numbers, Kirkman-Cayley numbers.

This work was partially supported by NSF grants DMS-0245385 (S.F.) and DMS-0202430 (N.R.).

1. Introduction

In the first part of this paper (Sections 2–12), we introduce and study a simplicial complex $\Delta^m(\Phi)$ associated to a finite root system Φ and a nonnegative integer parameter m. For m=1, our construction specializes to the (simplicial) generalized associahedra $\Delta(\Phi)$ introduced in [13] and identified in [14] as the underlying complexes for the cluster algebras of finite type. We enumerate the faces of the complexes $\Delta^m(\Phi)$ and determine their Euler characteristics. For the classical types in the Cartan-Killing classification, we provide explicit combinatorial descriptions of these complexes in terms of dissections of a convex polygon into smaller polygons. In types A_n and B_n , we rediscover the constructions given by E. Tzanaki [29]. Enumerative combinatorial invariants of the complexes $\Delta^m(\Phi)$ provide natural generalizations of the Fuss-Catalan, Kirkman-Cayley, and Przytycki-Sikora numbers to arbitrary types, suggesting connections with a host of well studied problems in the algebraic combinatorics of finite Coxeter groups, root systems, and hyperplane arrangements.

The second part of the paper (Sections 13–14) is devoted to combinatorial algorithms for determining Coxeter-theoretic invariants. Starting with a Coxeter diagram of a finite Coxeter group (or with the corresponding Dynkin diagram or Cartan matrix), we compute the Coxeter number, the exponents, and other related invariants by a procedure (in fact, by several alternative procedures) which is entirely combinatorial in nature. That is, these procedures only use the most elementary graph-theoretic concepts applied to the input diagram, and do not involve, in any way, the Coxeter group itself, the associated root system, the root lattice, or any other group-theoretic, lattice-theoretic, or Lie-theoretic notions and constructions. The crucial ingredients in all of these calculations are some identities for the face numbers of the generalized cluster complexes established in the first part of the paper. Formally extending these procedures, one can calculate "fake" Coxeter invariants (such as a "fake Coxeter number") of various infinite Coxeter groups.

The only prerequisites for this paper are the fundamentals of root systems and finite Coxeter groups; see, e.g., [12, Lectures 1–2], or the standard references [6, 17].

The paper is organized as follows. Section 2 reviews the main notions related to the generalized associahedra $\Delta(\Phi)$. Readers familiar with the original source [13, Section 3] or with the survey in [12, Section 4.3] may proceed directly to Section 3. There, we define the generalized cluster complexes $\Delta^m(\Phi)$ and describe their basic structural properties. The proofs of two of these results (Theorems 3.4 and 3.7) are given in Sections 6 and 7, respectively, and can be skipped at first reading. Section 4 presents some examples of the complexes $\Delta^m(\Phi)$. Combinatorial models for $\Delta^m(\Phi)$ when Φ is of classical type (i.e., A_n , B_n , C_n , or D_n) are given in Section 5. Recursions and explicit multiplicative formulas for the face numbers of $\Delta^m(\Phi)$ are given in Section 8; these formulas are proved in Section 9. In Section 10, we compute the h-vector of $\Delta^m(\Phi)$, recovering the m-Narayana numbers introduced by C. A. Athanasiadis [3]. The reduced Euler characteristic of $\Delta^m(\Phi)$ is determined in Section 11. In Section 12, we enumerate the m-analogues of "positive clusters." Section 13 presents several alternative algorithms for computing classical invariants of a root system Φ (or the associated Coxeter group W). In Section 14, these algorithms are applied to calculate "fake invariants" of some infinite Coxeter groups.

2. Cluster complexes of finite type

This section reviews the basic facts from [13, Section 3] concerning (simplicial) generalized associahedra; alternatively, see [12, Section 4.3]. The only difference is that we do *not* assume that the underlying root system Φ is crystallographic. This does not in fact create any additional complications: as noted in [12], the constructions in [13] extend verbatim to the non-crystallographic case.

Let Φ be a finite root system of rank n. We denote by $\Phi_{>0}$ the set of positive roots in Φ . The set of simple roots in Φ is denoted by $\Pi = \{\alpha_i : i \in I\}$, where I is an n-element indexing set. (The standard choice is $I = \{1, \ldots, n\}$.) Accordingly, $-\Pi = \{-\alpha_i : i \in I\}$ is the set of negative simple roots. The set $S = \{s_i : i \in I\}$ of simple reflections corresponding to the simple roots α_i generates a finite reflection group W. The pair (W, S) is a Coxeter system; the Coxeter group W naturally acts on the set of roots Φ .

Let us temporarily assume that the root system Φ is *irreducible*. Let $I = I_+ \cup I_-$ be a decomposition of I such that the sets I_+ and I_- are disjoint, and each of them labels a totally disconnected set of vertices in the Coxeter diagram of Φ .

The ground set for the cluster complex $\Delta(\Phi)$ is the set

$$\Phi_{>-1} = \Phi_{>0} \cup (-\Pi)$$

of almost positive roots. We define the involutions $\tau_{\pm}:\Phi_{\geq -1}\to\Phi_{\geq -1}$ by

$$\tau_{\varepsilon}(\alpha) = \begin{cases} \alpha & \text{if } \alpha = -\alpha_i, \text{ for } i \in I_{-\varepsilon}; \\ \left(\prod_{i \in I_{\varepsilon}} s_i\right)(\alpha) & \text{otherwise,} \end{cases}$$

for $\varepsilon \in \{+, -\}$. The product $R = \tau_- \tau_+$ can be viewed as a deformation of the Coxeter element in W. We denote by $\langle R \rangle$ the cyclic group generated by R.

Let h denote the Coxeter number of W and let w_o be the longest element of W.

Lemma 2.1 ([13, Theorem 2.6, Proposition 2.5]). The order of R is (h + 2)/2 if $w_{\circ} = -1$, and is h + 2 otherwise. Every $\langle R \rangle$ -orbit in $\Phi_{\geq -1}$ has a nonempty intersection with $(-\Pi)$. These intersections are precisely the $\langle -w_{\circ} \rangle$ -orbits in $(-\Pi)$.

The following theorem is a reformulation of results in [13, Section 3.1].

Theorem 2.2. There is a unique symmetric binary relation on $\Phi_{\geq -1}$ (called "compatibility") such that:

- α and β are compatible if and only if $R(\alpha)$ and $R(\beta)$ are compatible;
- a negative simple root $-\alpha_i$ is compatible with a positive root β if and only if the simple root expansion of β does not involve α_i .

Following [13, p. 983], we define the cluster complex $\Delta(\Phi)$ (of type Φ) as the clique complex for the compatibility relation. That is, a subset of roots in $\Phi_{\geq -1}$ forms a simplex in $\Delta(\Phi)$ if and only if every pair of roots in this subset is compatible.

If Φ is *reducible*, with irreducible components Φ_1, \ldots, Φ_l , then

$$\Phi_{\geq -1} = \bigcup_{j} (\Phi_j)_{\geq -1}$$

(disjoint union). We declare two roots in $\Phi_{\geq -1}$ compatible if and only if they either belong to different components, or belong to the same component and are compatible within it. Thus, the simplicial complex $\Delta(\Phi)$ is the *join* of the complexes $\Delta(\Phi_i)$.

The simplicial complex $\Delta(\Phi)$ is homeomorphic to a sphere [13]. Moreover, it can be explicitly realized as a boundary of a convex polytope [9], a polar dual to the (simple) generalized associahedron of type Φ . This is why $\Delta(\Phi)$ is sometimes referred to as the *simplicial* generalized associahedron (of type Φ).

By [14, Theorem 1.13], the definition of $\Delta(\Phi)$ given above is equivalent to the algebraic definition [14] of a cluster complex for a cluster algebra of finite type. Although inspired by cluster algebra theory, this paper does not rely on any of its results; cf. Remark 3.13.

3. Generalized cluster complexes

Let m be a nonnegative integer. In this section, we define and begin to study the main object of this paper, the generalized cluster complex $\Delta^m(\Phi)$. The ground set of $\Delta^m(\Phi)$ is the set $\Phi_{\geq -1}^m$ of colored almost positive roots. It consists of m copies of the set $\Phi_{>0}$ of positive roots in Φ together with one copy of the negative simple roots. It will be convenient to use the following notation for the elements of $\Phi_{\geq -1}^m$. For each $\alpha \in \Phi_{>0}$, let $\alpha^1, \ldots, \alpha^m$ denote the m "colored" copies of α occurring in $\Phi_{\geq -1}^m$. Each negative simple root α occurs in $\Phi_{\geq -1}^m$ as α^1 . Thus,

$$\Phi_{\geq -1}^m = \{ \alpha^k : \alpha \in \Phi_{>0}, k \in \{1, \dots, m\} \} \cup \{ (-\alpha_i)^1 : i \in I \}.$$

The simplicial complex $\Delta^m(\Phi)$ is defined using the binary compatibility relation on $\Phi^m_{\geq -1}$. This relation can be defined in (at least) two different ways, which we will now describe. As in the non-colored (m=1) case, we assume that Φ is irreducible, since the reducible case can be obtained by taking joins.

For a root $\beta \in \Phi_{>-1}$, let $d(\beta)$ denote the smallest d such that

$$\underbrace{R(R(R(\cdots R(\beta)\cdots)))}_{d \text{ times}}$$

is a negative root. In particular, $d(\beta) = 0$ if β is negative simple.

Definition 3.1. Two colored roots $\alpha^k, \beta^l \in \Phi^m_{\geq -1}$ are called *compatible* if and only if one of the following conditions is satisfied:

- k > l, $d(\alpha) \le d(\beta)$, and the roots $R(\alpha)$ and β are compatible (in the original "non-colored" sense of Theorem 2.2);
- $k < l, d(\alpha) \ge d(\beta)$, and the roots α and $R(\beta)$ are compatible;
- k > l, $d(\alpha) > d(\beta)$, and the roots α and β are compatible;
- $k < l, d(\alpha) < d(\beta)$, and the roots α and β are compatible;
- k = l, and the roots α and β are compatible.

Lemma 3.2. The compatibility relation on $\Phi^m_{\geq -1}$ is symmetric.

Proof. Immediate from Definition 3.1 and the symmetry of the compatibility relation on $\Phi_{\geq -1}$.

We next define R_m , the m-analogue of R, and prove an m-analogue of Theorem 2.2.

Definition 3.3. For $\alpha^k \in \Phi^m_{>-1}$, we set

$$R_m(\alpha^k) = \begin{cases} \alpha^{k+1} & \text{if } \alpha \in \Phi_{>0} \text{ and } k < m; \\ (R(\alpha))^1 & \text{otherwise.} \end{cases}$$

Theorem 3.4. The compatibility relation on $\Phi^m_{>-1}$ has the following properties:

- (i) α^k is compatible with β^l if and only if $R_m(\alpha^k)$ is compatible with $R_m(\beta^l)$;
- (ii) $(-\alpha_i)^1$ is compatible with β^l if and only if the simple root expansion of β does not involve α_i .

Furthermore, conditions (i)–(ii) uniquely determine this relation.

The proof of Theorem 3.4 is given in Section 6. It does not rely on any statements proved in between.

Corollary 3.5. Simultaneously replacing $R = \tau_- \tau_+$ by $R = \tau_+ \tau_-$ and changing the colors by $i \mapsto m - i + 1$ does not change the compatibility relation on $\Phi^m_{>-1}$.

Proof. Immediate from Theorem 3.4, since all we are really doing is replacing R_m by $(R_m)^{-1}$.

Corollary 3.6. If $m' \leq m$, then $\Delta^{m'}(\Phi)$ is a (vertex-induced) subcomplex of $\Delta^{m}(\Phi)$.

Proof. Follows directly from Theorem 3.4.

For a reducible root system $\Phi = \Phi_1 \times \cdots \times \Phi_l$, compatibility is defined by analogy with the non-colored case. Two colored roots $\alpha^k \in (\Phi_i)_{\geq -1}^m$ and $\beta^l \in (\Phi_j)_{\geq -1}^m$ are compatible if either $i \neq j$ or else i = j and α^k and β^l are compatible as elements of $(\Phi_i)_{\geq -1}^m$.

For a non-empty subset $J \subset I$, let Φ_J denote the parabolic root subsystem of Φ spanned by the simple roots α_j for $j \in J$. For $i \in I$, we denote $\langle i \rangle \stackrel{\text{def}}{=} I - \{i\}$. Thus $\Phi_{\langle i \rangle}$ consists of all roots in Φ whose simple root expansion does not involve α_i .

The compatibility relation is preserved under restriction to a parabolic subsystem:

Theorem 3.7. If $\alpha^k, \beta^l \in \Phi^m_{\geq -1}$ and $\alpha, \beta \in \Phi_J$, then α and β are compatible in $\Phi^m_{\geq -1}$ if and only if they are compatible in $(\Phi_J)^m_{\geq -1}$.

The proof of Theorem 3.7 is given in Section 7 using the classification of finite Coxeter groups together with the combinatorial models for generalized cluster complexes of types ABD presented in Section 5.

Definition 3.8. The generalized cluster complex $\Delta^m(\Phi)$ is the clique complex for the compatibility relation on $\Phi^m_{\geq -1}$. That is, the ground set for $\Delta^m(\Phi)$ is the set of colored almost positive roots, and a subset of such roots forms a simplex if and only if any two of them are compatible.

If Φ is irreducible, Corollary 3.5 implies that the complex $\Delta^m(\Phi)$ is invariant, up to an isomorphism, under interchanging τ_+ and τ_- . If Φ is reducible, $\Phi = \Phi_1 \times \cdots \times \Phi_l$, then $\Delta^m(\Phi)$ is, by design, the join of the complexes $\Delta^m(\Phi_i)$.

As in the non-colored case, all maximal simplices of $\Delta^m(\Phi)$ have cardinality n:

Theorem 3.9. The simplicial complex $\Delta^m(\Phi)$ is pure of dimension n-1.

Proof. Let F be a maximal face of $\Delta^m(\Phi)$ and let α^k be a colored root in F. Choose i so that $R_m^i(\alpha^k)$ is a negative simple root $(\alpha')^1$. Then $F' \stackrel{\text{def}}{=} R_m^i(F)$ is a maximal face of $\Delta^m(\Phi)$ containing $(\alpha')^1$. By Theorem 3.4, all other roots in F' are of the form β^l for $\beta \in (\Phi_{\langle i \rangle})_{\geq -1}$, and these roots form a maximal face F'' in $\Delta^m(\Phi_{\langle i \rangle})$. (Here we use Theorem 3.7.) By induction, this simplex is of dimension n-2, so F is of dimension n-1. The base of the induction (n=1) is trivial; cf. Example 4.3 below.

Proposition 3.10. Each codimension 1 face of $\Delta^m(\Phi)$ is contained in exactly m+1 faces of maximal dimension.

Proof. Let F be a codimension 1 face of $\Delta^m(\Phi)$ and let α^k , i, α' , F' and F'' be defined as in the proof of Theorem 3.9. Then F'' is a codimension 1 face of $\Delta^m(\Phi_{\langle i \rangle})$, which by induction is contained in exactly m+1 faces of maximal dimension. Adjoining $(\alpha')^1$ to each of these faces and applying R_m^{-i} , one obtains all maximal faces containing F. \square

Proposition 3.11. The link of any face in $\Delta^m(\Phi)$ is isomorphic to a join of generalized cluster complexes of the form $\Delta^m(\Phi_J)$, for some irreducible parabolic root subsystems $\Phi_j \subset \Phi$.

Proof. It is enough to verify the statement for links of vertices. Applying R_m to a vertex until it becomes a negative simple root, the statement follows.

Remark 3.12. Assume that Φ is a crystallographic root system, and let Φ^{\vee} be the root system dual to Φ . As shown in [13, Proposition 3.3], two roots $\alpha, \beta \in \Phi_{\geq -1}$ are compatible if and only if the corresponding coroots α^{\vee} and β^{\vee} are compatible in $\Phi^{\vee}_{\geq -1}$. Thus the cluster complex $\Delta(\Phi)$ is canonically isomorphic to $\Delta(\Phi^{\vee})$. It follows directly from Definition 3.1 that the same is true about $\Delta^m(\Phi)$ and $\Delta^m(\Phi^{\vee})$. In particular, the generalized cluster complexes of types B_n and C_n are canonically isomorphic. In view of this, we do not consider type C_n in what follows.

Remark 3.13. It would be very interesting to find a generalization of the notion of a cluster complex of a cluster algebra of finite type (as defined in [14]) that would yield the complexes $\Delta^m(\Phi)$. This might also lead to an extension of the concept to infinite types.

4. Examples

Here we illustrate the definition of a generalized cluster complex (see Definition 3.8) by considering the special cases where $m \le 1$ or $n \le 2$.

Example 4.1 (m = 0). The ground set of $\Delta^0(\Phi)$ consists of the n negative simple roots in Φ . Any two such roots are compatible. Hence $\Delta^0(\Phi)$ is an (n-1)-dimensional simplex.

Example 4.2 (m=1). By design, $\Delta^1(\Phi) = \Delta(\Phi)$ is the cluster complex of (finite) type Φ . In particular, in type A_n we obtain Stasheff's associahedron, while in type B_n/C_n we get the Bott-Taubes' cyclohedron. See [12, 13] or Section 5 below for further details.

Example 4.3 (n = 1). For a root system Φ of type A_1 , the simplicial complex $\Delta^m(\Phi)$ consists of m+1 disconnected points $(-\alpha_1)^1, \alpha_1^1, \ldots, \alpha_1^m$.

Example 4.4 (n = 2). Let Φ be a root system of type $I_2(a)$. The (1-dimensional) simplicial complex $\Delta^m(\Phi)$ is an (m + 1)-regular graph on am + 2 vertices. Figure 1 shows this graph for a root system of type $I_2(4) = B_2$ and m = 3. The figure is color-coded to highlight the subgraphs corresponding to m = 2 and m = 1 and thus to illustrate how $\Delta^m(B_2)$ changes with increasing m (cf. Corollary 3.6). The simple roots are α_1 and α_2 , with $I_+ = \{1\}$ and $I_- = \{2\}$. The other two roots are denoted by $\beta_1 = 2\alpha_1 + \alpha_2$ and $\beta_2 = \alpha_1 + \alpha_2$. The black edges are the edges of $\Delta^1(B_2)$, the black and dark-gray edges form $\Delta^2(B_2)$, and the edges of all colors form $\Delta^3(B_2)$.




FIGURE 1. Root system of type B_2 and the complex $\Delta^3(B_2)$

Figure 2 shows the same graph, drawn so as to make its symmetry apparent. The map R_m acts by a $\frac{2\pi}{7}$ counterclockwise rotation of this picture.




FIGURE 2. Representation of $\Delta^3(B_2)$ showing 7-fold symmetry

In general, the graph $\Delta^m(I_2(a))$ can be constructed in the plane as follows. Take the vertex set to be the integers modulo (am+2), identified with the vertices of a regular (am+2)-gon so that $0,1,2,\ldots$ lists them in clockwise order. For a odd, the edge set has (am+2)-fold rotational symmetry and connects each vertex v to the m+1 vertices of the form $v+\frac{a-1}{2}m+j$, for $j=0,\ldots,m$. Figure 3 shows this graph for a=5 and m=4. For a even, fix an odd integer i. The edge set of $\Delta^m(I_2(a))$ has (am/2+1)-fold rotational symmetry, and connects 0 to the vertices $i, i+2, \ldots, i+2m$. Figure 2 is $\Delta^3(B_2)$ drawn in this style with i=3.




FIGURE 3. $\Delta^4(I_2(5))$

Remark 4.5. In the special case n=m=2, the complex $\Delta^2(I_2(a))$ can be obtained from the boundary of a regular (2a+2)-gon by adding an edge connecting every pair of antipodal vertices, as illustrated in Figure 4. Thus, $\Delta^2(I_2(a))$ is the 1-skeleton of a polygonal subdivision of \mathbb{RP}^2 . It would be interesting to determine which complexes $\Delta^m(\Phi)$ (perhaps all of them?) can be realized as skeleta of polyhedral (n+m-2)-dimensional manifolds.

5. The classical types

In this section, we describe combinatorial models realizing the generalized cluster complexes $\Delta^m(\Phi)$ for the classical types A_n , B_n/C_n , and D_n . Each of these models is constructed in terms of collections of non-intersecting diagonals in a certain convex polygon. In every case, the isomorphism between the suggested model and the root-theoretic description of $\Delta^m(\Phi)$ can be verified by straightforward induction on m. The base of the induction is the non-colored (m=1) case, described in [13, Section 3.5].




FIGURE 4. $\Delta^2(B_2)$ in the real projective plane

For the types A_n and B_n , we rediscover the constructions given earlier by E. Tzanaki [29]. The type- A_n construction was inspired by the paper [22] by J. H. Przytycki and A. S. Sikora.

5.1. **Type** A_n . It is easy to see that a convex polygon \mathbf{P} with N vertices can be dissected into (convex) (m+2)-gons by pairwise non-crossing diagonals if and only if $N \equiv 2 \mod m$. (Here and in what follows, "non-crossing" means that diagonals have no common points in the *interior* of \mathbf{P} .) Let \mathbf{P} be a convex polygon with (n+1)m+2 vertices. To simplify descriptions, we take \mathbf{P} to be regular. A diagonal of \mathbf{P} is called m-allowable if it cuts \mathbf{P} into two polygons each of which can be dissected into (m+2)-gons. That is, an m-allowable diagonal connects two vertices such that each of the two paths connecting them along the perimeter of \mathbf{P} passes through a number of vertices that is divisible by m.

Let us now define a simplicial complex $\Delta(m,n)$ on the set of all m-allowable diagonals in \mathbf{P} . Such diagonals form a simplex in $\Delta(m,n)$ if and only if they are pairwise non-crossing. The complex $\Delta(m,n)$ is easily seen to be pure, with maximal simplices formed by n-tuples of diagonals that cut \mathbf{P} into (m+2)-gons.

We now construct an isomorphism between $\Delta(m,n)$ and the generalized cluster complex $\Delta^m(\Phi)$ for a root system Φ of type A_n . Under the isomorphism, R_m will correspond to a clockwise rotation of \mathbf{P} taking P_2 to P_1 , etc. To avoid additional notation, we refer to this clockwise rotation as R_m while defining the isomorphism.

We use the standard labeling of the simple roots in Φ (see, e.g., [6]) and take I_+ to be the odd indices. Let $P_1, P_2, \ldots, P_{(n+1)m+2}$ be the vertices of \mathbf{P} , labeled counterclockwise. For $1 \leq i \leq \frac{n+1}{2}$, identify the negative simple root $-\alpha_{2i-1}$ with the diagonal of \mathbf{P} connecting $P_{(i-1)m+1}$ to $P_{(n+1-i)m+2}$. For $1 \leq i \leq \frac{n}{2}$, identify $-\alpha_{2i}$ with the diagonal connecting P_{im+1} to $P_{(n+1-i)m+2}$. Collectively, these n diagonals (each of them m-allowable) form what we call the m-snake (cf. [13, Figure 3]). The

positive roots of Φ are $\alpha_{ij} = \alpha_i + \cdots + \alpha_j$ for each i and j with $1 \leq i \leq j \leq n$. For each such i and j, there are exactly m diagonals which are m-allowable and intersect the diagonals $-\alpha_i, \ldots, -\alpha_j$ and no other diagonals in the m-snake. This collection of diagonals is of the form $R_m^0 D, R_m^1 D, \ldots, R_m^{m-1} D$ for some diagonal D. For $1 \leq k \leq m$, we identify α_{ij}^k with $R_m^{k-1} D$. Figure 5 shows the m-snake for n=4 and m=5, along with the diagonals identified with the colored roots $\alpha_{24}^1, \ldots, \alpha_{24}^5$.




FIGURE 5. The 5-snake for A_4

For m=1, the complex $\Delta(1,n)\cong\Delta(A_n)$ is, by definition (see, e.g., [12] and references therein), the dual complex for (the boundary of) the ordinary n-dimensional associahedron, also known as the Stasheff polytope. The first non-trivial example with m>1 is $\Delta(2,2)\cong\Delta^2(A_2)$, the complex of quadrangulations of an octagon. (Cf. Example 4.4 with m=2 and a=3, and also Remark 4.5 with a=3.) This 1-dimensional complex is a non-planar 3-regular graph with 8 vertices and 12 edges. Thus $\Delta(2,2)$ is homotopy equivalent to a wedge of 5 circles.

5.2. **Type** B_n/C_n . When Φ is of type B_n (or C_n —cf. Remark 3.12), the complex $\Delta^m(\Phi)$ can be realized as follows. Let **P** be a centrally symmetric regular polygon

with 2nm + 2 vertices. The map R_m will correspond to a $\frac{\pi}{nm+1}$ clockwise rotation about the center of **P**. The vertices of the complex are of two kinds:

- the *diameters* of **P**, i.e., diagonals connecting antipodal vertices (all such diagonals are m-allowable in the sense of Section 5.1);
- the pairs (D, D') of distinct m-allowable diagonals of \mathbf{P} such that D is related to D' by a half-turn about the center of \mathbf{P} .

Two vertices are called "non-crossing" if no diagonal representing one vertex crosses a diagonal representing the second vertex. The simplices in the complex are the sets of pairwise non-crossing vertices. It is easily verified that this is a pure complex of dimension n. Its maximal simplices correspond to centrally symmetric dissections of \mathbf{P} into (m+2)-gons.

Figure 6 illustrates this complex for n = m = 2. The edges of the graph correspond to centrally symmetric quadrangulations of a 10-gon.




FIGURE 6. Combinatorial realization of $\Delta^2(B_2)$

For m=1, this construction specializes to one of the common definitions of the n-dimensional cyclohedron, or $Bott-Taubes\ polytope$ (see, e.g., [12, 13] and references therein).

In order to describe an isomorphism between this complex and $\Delta^m(B_n)$, we relate the former to $\Delta(m,n)$. Label the simple roots α_1,\ldots,α_n so that α_n is the only simple root in its W-orbit. The negative simple roots correspond to the orbits, under central symmetry, of the diagonals on the m-snake of type A_{2n-1} (see Section 5.1). If $\beta_1,\ldots,\beta_{2n-1}$ are the simple roots of A_{2n-1} , then $-\alpha_i$ is encoded by the pair of diagonals corresponding to $-\beta_i$ and $-\beta_{2n-i}$, for $1 \leq i \leq n-1$, while $-\alpha_n$ is encoded by the diameter corresponding to $-\beta_n$.

The positive roots of B_n can be divided into three categories:

- (I) $\alpha_i + \alpha_{i+1} + \dots + \alpha_j$ for $i \le j < n$
- (II) $\alpha_i + \alpha_{i+1} + \dots + \alpha_n$ for $i \leq n$, and
- (III) $\alpha_i + \alpha_{i+1} + \dots + \alpha_{j-1} + 2\alpha_j + 2\alpha_{j+1} + \dots + 2\alpha_n$ for $i < j \le n$.

A colored positive root α^k for α in category (I) is encoded by the pair consisting of the diagonal for $(\beta_i + \beta_{i+1} + \cdots + \beta_j)^k$ and the diagonal for $(\beta_{2n-j} + \beta_{n-j+1} + \cdots + \beta_{2n-i})^k$. For α in category (II), α^k is encoded by the diameter corresponding to $(\beta_i + \beta_{i+1} + \cdots + \beta_{2n-i})^k$. For α in category (III), α^k is encoded by the diagonal corresponding to $(\beta_i + \beta_{i+1} + \cdots + \beta_{2n-j})^k$ and the diagonal corresponding to $(\beta_j + \beta_{j+1} + \cdots + \beta_{2n-j})^k$.

5.3. **Type** D_n . Let **P** be a regular polygon with 2(n-1)m+2 vertices. The vertices in the combinatorial realization of $\Delta^m(D_n)$ fall into two groups. The vertices in the first group correspond one-to-one to pairs of distinct non-diameter m-allowable diagonals in **P** related by a half-turn. In the second group, each vertex is indexed by a diameter of **P**, together with one of two *flavors*, which we will call "dashed" and "gray," and picture accordingly. Thus, each diameter occurs twice, in each of the two flavors. We label the vertices of **P** counterclockwise:

$$P_1, P_2, \dots, P_{(n-1)m+1}, -P_1, -P_2, \dots, -P_{(n-1)m+1}.$$

We call $[P_1, -P_1]$ the primary diameter. By construction, the map R_m acts by rotating P_2 clockwise to P_1 and switching the flavor of certain diameters. Specifically, R_m preserves flavor when applied to a diameter of the form $[P_k, -P_k]$ (for $1 \le k \le (n-1)m+1$) unless k=1 or $k \equiv 2 \mod m$. If k=1 or $k \equiv 2 \mod m$, the flavor is switched.

By analogy with types A_n and B_n , the complex $\Delta^m(D_n)$ is realized as a clique complex for a certain "compatibility" relation defined as follows. Two vertices at least one of which is not a diameter are compatible or not according to precisely the same rules as in the type- B_n case. A more complicated condition determines whether two diameters are compatible. Two diameters with the same location and different flavors are compatible. Two diameters at different locations are compatible if and only if applying R_m repeatedly until either of them is in position $[P_1, -P_1]$ results in diameters of the same flavor. Explicitly, let D be a flavored diameter and let \overline{D} denote D with its flavor reversed. Then for $1 \leq k \leq (n-1)m$, the flavored diameter $R_m^k(D)$ is compatible with D (and incompatible with \overline{D}) if and only if $\left\lceil \frac{k}{m} \right\rceil$ is even.

Figure 7 shows the 12 maximal simplices of $\Delta^2(D_3)$ containing the gray diameter $[P_1, -P_1]^{gr}$. The points $\pm P_1$ are marked with a white dot. Some edges of \mathbf{P} are broken, indicating the locations at which diameters change flavors under the action of R_m . Specifically, P_i and P_{i-1} are connected by a broken edge if and only if R_m maps a flavored diameter $[P_i, -P_i]$ to the opposite-flavored diameter $[P_{i-1}, -P_{i-1}]$. To illustrate the rules of compatibility for pairs of diameters, Figure 8 shows the orbit of a maximal simplex under the action of R_m .

We next describe an isomorphism between the complex defined above and the original construction of $\Delta^m(D_n)$. Label the simple roots $\alpha_1, \ldots, \alpha_n$ so that α_{n-1} and α_n correspond to the ends of the fork in the Dynkin diagram of type D_n . The negative simple roots correspond to the orbits, under central symmetry, of the diagonals in the m-snake for A_{2n-3} , except that α_n and α_{n-1} both correspond to the same diameter (in two different flavors). Specifically, if $\beta_1, \ldots, \beta_{2n-3}$ are the simple roots of A_{2n-3} , then $-\alpha_i$ is encoded by the pair of diagonals corresponding to $-\beta_i$ and $-\beta_{2n-i-2}$, for $1 \le i \le n-2$. The dashed copy of the diameter corresponding to $-\beta_n$




FIGURE 7. Some maximal simplices in $\Delta^2(D_3)$




FIGURE 8. The orbit of a maximal simplex in $\Delta^2(D_3)$

encodes $-\alpha_{n-1}$, while $-\alpha_n$ is encoded by the gray diameter corresponding to $-\beta_n$. We label the vertices of **P** so that this is the primary diameter $[P_1, -P_1]$.

The positive roots of D_n can be divided into two categories:

(I)
$$\alpha_i + \alpha_{i+1} + \dots + \alpha_j$$
 for $i \le j < n$
(II) $\alpha_i + \alpha_{i+1} + \dots + \alpha_{n-2} + \alpha_j + \alpha_{j+1} + \dots + \alpha_n$ for $i < j \le n$.

A colored positive root α^k for α in category (I) is encoded by the pair consisting of the diagonal for $(\beta_i + \beta_{i+1} + \dots + \beta_j)^k$ and the diagonal for $(\beta_{2n-j-2} + \beta_{2n-j-1} + \dots + \beta_{2n-i-2})^k$ if j < n-1. If $\alpha = \alpha_i + \dots + \alpha_{n-1}$ then α^k is encoded by the gray diameter corresponding to $(\beta_i + \beta_{i+1} + \dots + \beta_{2n-i-2})^k$.

For α in category (II), α^k is encoded by the diagonal corresponding to $(\beta_i + \beta_{i+1} + \cdots + \beta_{2n-j-2})^k$ and the diagonal corresponding to $(\beta_j + \beta_{j+1} + \cdots + \beta_{2n-i-2})^k$ if j < n.

If $\alpha = \alpha_i + \cdots + \alpha_{n-2} + \alpha_n$ then α^k is encoded by the dashed diameter corresponding to $(\beta_i + \beta_{i+1} + \cdots + \beta_{2n-i-2})^k$.

Remark 5.1. In types A_n and B_n , the combinatorial models for the complex $\Delta^m(\Phi)$ presented in Sections 5.1 and 5.2 describe this complex as a subcomplex of the ordinary cluster complex $\Delta(\Psi)$ for a larger root system Ψ of type A_{nm+m-1} or B_{nm} , respectively. It is not clear whether such an embedding exists for the type D_n .

6. Proof of Theorem 3.4

The proof makes use of the following results, which can be extracted from [13].

Lemma 6.1.

- (1) Each R-orbit in $\Phi_{\geq -1}$ either has size h+2 and contains two negative simple roots, or has size (h+2)/2 and contains one negative simple root.
- (2) If h is even, then for every orbit of size h+2, the two negative simple roots are placed symmetrically in the orbit. That is, applying R to one of the negative simple roots (h+2)/2 times yields the other negative simple root.
- (3) If h is odd, then necessarily all R-orbits have size h + 2. The two negative simple roots in each orbit are placed so that applying R to one of the negative simple roots (h + 3)/2 times yields the other negative simple root. Then applying R an additional (h + 1)/2 times yields the original negative simple root.
- (4) In every orbit, positive simple roots are adjacent to their negatives. When h is odd, both positive simple roots in an orbit are located on the longer of the two paths between the two negative simple roots.

If m=1, then both characterizations of compatibility specialize to the non-colored version, so from now on we assume that m>1. Let $\alpha^k, \beta^l \in \Phi^m_{\geq -1}$. Without loss of generality, let $d(\alpha) \leq d(\beta)$. Proving condition (i) requires us to consider several cases. First, if neither α nor β is negative simple and neither k nor l equals m, then condition (i) is immediate.

Let us consider the cases where neither α nor β is negative simple but k=m or l=m or both. If k=l=m, then α^k and β^l are compatible if and only if α and β are compatible, and $R_m(\alpha^k) = R(\alpha)^1$ is compatible with $R_m(\beta^l) = R(\beta)^1$ if and only if $R(\beta)$ and $R(\alpha)$ are compatible, so (i) follows by the R-invariance of non-colored compatibility. If m=k>l, then $R_m(\alpha^k) = R(\alpha)^1$ and since $d(R(\alpha)) < d(\alpha) \le d(\beta)$, $R(\alpha)^1$ is compatible with $R_m(\beta^l) = \beta^{l+1}$ if and only if $R(\alpha)$ is compatible with β , and this is exactly the requirement for α^k to be compatible with β^l .

For k < l = m there are two cases: $d(\alpha) < d(\beta)$ and $d(\alpha) = d(\beta)$. If $d(\alpha) < d(\beta)$, then α^k is compatible with β^l if and only if α is compatible with β . Also, $d(\alpha) \le d(R(\beta))$, so $R_m(\alpha^k) = \alpha^{k+1}$ is compatible with $R_m(\beta^l) = R(\beta)^1$ if and only if $R(\alpha)$ and $R(\beta)$ are compatible, and (i) follows by R-invariance of non-colored compatibility. If $d(\alpha) = d(\beta)$, then α^k is compatible with β^l if and only if α is compatible with $R(\beta)$. Now $d(\alpha) > d(R(\beta))$, so $R_m(\alpha^k) = \alpha^{k+1}$ is compatible with $R_m(\beta^l) = R(\beta)^1$ if and only if α and $R(\beta)$ are compatible.

If both α and β are negative simple, then k = l = 1, $R_m(\alpha^1) = R(\alpha)^1$, and $R_m(\beta^1) = R(\beta)^1$. Now (i) is immediate by R-invariance.

If only one of the two roots is negative simple, then $d(\alpha) \leq d(\beta)$ implies that the negative simple root must be α , and in particular k = 1 and $d(\alpha) < d(\beta)$. The colored roots α^1 and β^l are compatible if and only if α and β are compatible. If l = m, then $R_m(\alpha^1) = R(\alpha)^1$ is compatible with $R_m(\beta^m) = R(\beta)^1$ if and only if $R(\alpha)$ and $R(\beta)$ are compatible, and (i) follows by R-invariance. We now assume l < m. If $d(R(\alpha)) \geq d(\beta)$, then $R_m(\alpha^1) = R(\alpha)^1$ is compatible with $R_m(\beta^l) = \beta^{l+1}$ if and only if $R(\alpha)$ and $R(\beta)$ are compatible, so (i) follows by R-invariance.

The condition $d(R(\alpha)) \geq d(\beta)$ will hold for most negative simple roots α and positive roots β . The only way for it to fail is if h is odd, $d(\beta) = (h+1)/2$ and $d(R(\alpha)) = (h-1)/2$. In this case, $R_m(\alpha^1) = R(\alpha)^1$ is compatible with $R_m(\beta^l) = \beta^{l+1}$ if and only if $R(\alpha)$ is compatible with β . Thus condition (i) in this case amounts to verifying that α and β are compatible if and only if $R(\alpha)$ and β are compatible. Because $d(\beta) = (h+1)/2$, the root β lies on the long path between two negative simple roots in its R-orbit and is adjacent to a negative simple root. Therefore β must be a (positive) simple root. If $\beta = -\alpha$, then α and β are not compatible, and $R(\alpha) = \beta$ so $R(\alpha)$ and β are not compatible. If $\beta \neq -\alpha$, then α and β are compatible and $R^{-1}(\beta)$ is a negative simple root distinct from α , so $R^{-1}(\beta)$ and α are compatible. Therefore $R(\alpha)$ and β are compatible.

If α is negative simple, then the third condition in Definition 3.1 always applies, and condition (ii) follows.

It remains to prove the last statement of Theorem 3.4. The following is a generalization of Lemma 2.1.

Lemma 6.2. The order of R_m is (mh+2)/2 if $w_o = -1$, and is mh+2 otherwise. Every $\langle R_m \rangle$ -orbit in $\Phi^m_{\geq -1}$ has a nonempty intersection with the set $(-\Pi)^1 \stackrel{\text{def}}{=} \{(-\alpha_i)^1\}_{i \in I}$. These intersections are precisely the $\langle -w_o \rangle$ -orbits in $(-\Pi)^1$.

Proof. Immediate from Lemma 2.1 and Definition 3.3.

It follows from Lemma 6.2 that conditions (i) and (ii) uniquely define the compatibility relation on $\Phi^m_{>-1}$.

7. Proof of Theorem 3.7

Type A_n . It is sufficient to treat the case $J = \langle i \rangle$. Let $\alpha^k \in \Phi^m_{\geq -1}$. By Theorem 3.4, $\alpha \in \Phi_{\langle i \rangle}$ if and only if the m-allowable diagonal in \mathbf{P} corresponding to α^k does not intersect the diagonal D corresponding to $(-\alpha_i)^1$. The diagonal D dissects \mathbf{P} into two polygons $\mathbf{P}_{\langle i \rangle}$ and $\mathbf{P}_{\geq i}$. The remaining negative simple roots (besides $-\alpha_i$) form two m-snakes, one in each of the two polygons.

The diagonal corresponding to a colored positive root α_{ij}^k is uniquely determined as the kth (in clockwise order) of the exactly m diagonals which intersect the diagonals for $-\alpha_i, -\alpha_{i+1}, \ldots, \alpha_j$ and no other diagonals in the m-snake. Thus the correspondence between $(\Phi_{\langle i \rangle})_{\geq -1}^m$ and the diagonals in the polygons $\mathbf{P}_{< i}$ and $\mathbf{P}_{> i}$ is identical to the correspondence between colored roots $\alpha^k \in \Phi_{\geq -1}^m$ with $\alpha \in \Phi_{\langle i \rangle}$ and diagonals in \mathbf{P} .

Type B_n . Theorem 3.7 for Φ of type B_n follows easily from the type- A_{2n-1} case by identifying parabolic root subsystems of B_n with parabolic root subsystems of A_{2n-1} which are fixed by the involution $\beta_i \mapsto \beta_{2n-i}$.

Type D_n . Again, it suffices to consider the case $J = \langle i \rangle$. First, assume that $i \notin \{n-1,n\}$. By Theorem 3.4, $\alpha^k \in \Phi^m_{\geq -1}$ has $\alpha \in \Phi_{\langle i \rangle}$ if and only if the corresponding diameter or diagonal pair in **P** does not intersect the diagonal pair $\{D,D'\}$ corresponding to $(-\alpha_i)^1$. These two diagonals dissect **P** into three polygons, one of which is centrally symmetric and two of which are related to each other by central symmetry. The proposition follows in this case by an argument analogous to the type A_n case.

By symmetry, we need only consider one additional case: i = n. The colored almost positive roots compatible with $-\alpha_n$ correspond to the dashed primary diameter, all gray non-primary diameters, and all pairs of diagonals which do not intersect the primary diameter. The parabolic root subsystem $\Phi_{\langle i \rangle}$ is of type A_{n-1} . We will realize $\Delta^m(\Phi_{\langle i \rangle})$ as the complex of non-crossing m-allowable diagonals in a convex polygon \mathbf{P}' with the vertices

$$P_1, P_2, \ldots, P_{(n-1)m+1}, -P_1, Q_1, Q_2, \ldots, Q_m$$

in counterclockwise order. The vertices P_i are vertices of ${\bf P}$ and the vertices Q_i are new.

For a diagonal pair $\{D, D'\}$ in \mathbf{P} which does not intersect the primary diameter, exactly one of the diagonals D and D' connects vertices in $\{P_1, P_2, \ldots, P_{(n-1)m+1}, -P_1\}$, and is therefore a diagonal in \mathbf{P}' . The dashed primary diameter of \mathbf{P} corresponds to the diagonal $[P_1, -P_1]$ of \mathbf{P}' . The correspondence between gray diameters in \mathbf{P} and the remaining diagonals of \mathbf{P}' is as follows. The gray diameter $[P_i, -P_i]$ in \mathbf{P} is mapped to the diagonal $[P_i, Q_{k(i)}]$ in \mathbf{P}' for $k(i) \equiv (i-1) \mod m$. Figure 9 illustrates this correspondence (for m=3 and Φ of type D_5) by showing a maximal simplex in $\Delta^m(\Phi_{\langle i\rangle})$ represented as a collection of gray diameters in \mathbf{P} and as the corresponding collection of diagonals in \mathbf{P}' .




Figure 9

It is immediate that if two diagonal pairs in \mathbf{P} do not intersect the primary diameter, then they intersect each other if and only if the corresponding diagonals in \mathbf{P}' intersect each other. It is also immediate that a diagonal pair in \mathbf{P} that does

not intersect the primary diameter intersects a gray diameter in \mathbf{P} if and only if the corresponding diagonals in \mathbf{P}' intersect each other. Intersections with the dashed primary diameter of \mathbf{P} also correspond to intersections with the diagonal $[P_1, -P_1]$ in \mathbf{P}' . One can check that for 1 < i < j < (n-1)m+1, a gray diameter $[P_i, -P_i]$ is compatible with a gray diameter $[P_j, -P_j]$ if and only if the fractional part of $\frac{i-2}{m}$ is greater than or equal to the fractional part of $\frac{j-2}{m}$. This is exactly the requirement that $[P_i, Q_{k(i)}]$ and $[P_i, Q_{k(j)}]$ do not intersect. Thus our correspondence also maps the compatibility relation among diameters and m-allowable diagonal pairs in \mathbf{P} compatible with the gray primary diameter to the compatibility relation among m-allowable diagonals of \mathbf{P}' .

It remains to verify that the process of mapping from $(\Phi_{\langle i \rangle})_{\geq -1}^m$ to diagonal pairs and flavored diameters in \mathbf{P} , then mapping to diagonals in \mathbf{P}' produces the same result as the process of mapping directly from $(\Phi_{\langle i \rangle})_{\geq -1}^m$ to diagonals in \mathbf{P}' by the type A_{n-1} construction. This is a straightforward check.

Exceptional types. For all exceptional types (including the non-crystallographic ones), Theorem 3.7 was checked with the help of a computer. In Definition 3.1, the compatibility of α^k and β^l depends on the comparison between k and l but not on the actual values of k and l. Thus it is sufficient to check the theorem in the case m = 2, so that in particular, for each type the theorem reduces to a finite computation. \square

Dihedral types $I_2(a)$. In this case, the statement is trivial (cf. Examples 4.3–4.4). \square

8. Face numbers

In this section, we enumerate the faces of various dimensions in the generalized cluster complexes $\Delta^m(\Phi)$. Let $f_k(\Phi, m)$ denote the kth face number $\Delta^m(\Phi)$ or, more precisely, the number of k-element simplices (i.e., (k-1)-dimensional faces) in $\Delta^m(\Phi)$. We denote by $N(\Phi, m) = f_n(\Phi, m)$ the number of maximal simplices in $\Delta^m(\Phi)$. We note that $f_0(\Phi) = 1$ for any Φ .

Example 8.1 (n = 1). In view of Example 4.3, we have

$$f_1(A_1,m) = m+1.$$

Example 8.2 (n = 2). By Example 4.4, the complex $\Delta^m(I_2(a))$ is an (m+1)-regular graph on am + 2 vertices. Hence

$$f_1(I_2(a), m) = am + 2,$$

 $f_2(I_2(a), m) = \frac{(am + 2)(m + 1)}{2}.$

The face numbers $f_k(\Phi, m)$ satisfy the following recurrences which generalize [13, Proposition 3.7].

Proposition 8.3. Let k be a positive integer. For a reducible root system $\Phi = \Phi_1 \times \Phi_2$,

(8.1)
$$f_k(\Phi_1 \times \Phi_2, m) = \sum_{k_1 + k_2 = k} f_{k_1}(\Phi_1, m) f_{k_2}(\Phi_2, m).$$

In particular,

$$N(\Phi_1 \times \Phi_2, m) = N(\Phi_1, m) N(\Phi_2, m).$$

If Φ is irreducible, then

(8.2)
$$f_k(\Phi, m) = \frac{mh + 2}{2k} \sum_{i \in I} f_{k-1}(\Phi_{\langle i \rangle}, m).$$

In particular,

(8.3)
$$N(\Phi, m) = \frac{mh + 2}{2n} \sum_{i \in I} N(\Phi_{\langle i \rangle}, m).$$

Proof. Formula (8.1) follows from the fact that $\Delta^m(\Phi_1 \times \Phi_2)$ is a join of $\Delta^m(\Phi_1)$ and $\Delta^m(\Phi_2)$. The proof of (8.2) is analogous to the proof of [13, Proposition 3.7]. Let us count in two different ways the number of pairs (α^j, S) where S is a k-element simplex in $\Delta^m(\Phi)$ and $\alpha^j \in S$. On the one hand, the number of such pairs is $k \cdot f_k(\Phi, m)$. On the other hand, combining Lemma 6.2, Theorem 3.4, and Theorem 3.7, we conclude that the colored roots α^j belonging to each $\langle R_m \rangle$ -orbit $\Omega \subset \Phi^m_{\geq -1}$ contribute

$$\frac{mh+2}{2} \sum_{-\alpha_i \in \Omega} f_{k_1}(\Phi_{\langle i \rangle}, m)$$

to the count (per negative simple root $(-\alpha)^1$), implying (8.2).

It follows from (8.2) by induction on m that $f_k(\Phi, m)$ is a polynomial in m of degree k. In this section, we calculate these polynomials for all irreducible root systems Φ . (The formulas for reducible cases follow at once.) It turns out that, with a single exception of $f_4(E_8, m)$, the face numbers $f_k(\Phi, m)$ factor over the rationals into linear factors.

We present the special case k = n first.

For the rest of this section, we assume that the root system Φ is irreducible. Let e_1, \ldots, e_n be the *exponents* of Φ .

Proposition 8.4. The number of maximal simplices in $\Delta^m(\Phi)$ is given by

(8.4)
$$N(\Phi, m) = \prod_{i} \frac{mh + e_i + 1}{e_i + 1}.$$

Proposition 8.4 is a special case of the general formula for the face numbers $f_k(\Phi, m)$ given in Theorem 8.5 below.

The product appearing in the right-hand side of (8.4) comes up in a variety of contexts related to the combinatorics of the root system Φ , the corresponding Coxeter group W, and associated Lie algebras and Lie groups. We refer the reader to [2] and references therein; see also the discussion in Section 10 below.

One would hope for a uniform formula for the face numbers in the spirit of (8.4). This does not quite happen. The factorization of $f_k(\Phi, m)$ into irreducible polynomials in m yields a subset of the factors appearing on the right-hand side of (8.4) together with, in types D_n $(n \ge 4)$, E_6 , E_7 , E_8 , F_4 , and H_4 , one additional "mysterious" factor, which we denote by $c_f(\Phi, k, m)$. We emphasize that $c_f(\Phi, k, m)$ is not a "fudge factor" needed to force our formulas to look like (8.4). Rather, it is forced upon us by the unique factorizations of the polynomials $f_k(\Phi, m)$ into irreducibles.

Theorem 8.5. The face numbers of the generalized cluster complex $\Delta^m(\Phi)$ for an irreducible root system Φ are given by

(8.5)
$$f_k(\Phi, m) = c_f(\Phi, k, m) \binom{n}{k} \prod_{\substack{\text{level}(e_i) \le k}} \frac{mh + e_i + 1}{e_i + 1}.$$

Here, the "levels" of exponents e_i are defined in Table 1, with each column showing the exponents of appropriate type, each of them placed in the row corresponding to its level. The factor $c_f(\Phi, k, m)$ is a polynomial in m given as follows:

- in types A_n , B_n , H_3 , and $I_2(m)$, we have $c_f(\Phi, k, m) = 1$;
- in type D_n , we have:

$$c_f(D_n, k, m) = \begin{cases} 1 & \text{if } k \in \{0, 1, n - 1, n\}; \\ \frac{n(n-1)m + k(k-1)m + kn}{kn} & \text{otherwise;} \end{cases}$$

• in types E_6 , E_7 , E_8 , F_4 , and H_4 , the factors $c_f(\Phi, k, m)$ are shown in Table 2. In particular, for the classical types A_n , B_n , and D_n , the face numbers are given by

(8.6)
$$f_k(A_n, m) = \frac{1}{k+1} \binom{n}{k} \binom{(n+1)m+k+1}{k},$$

(8.7)
$$f_k(B_n, m) = \binom{n}{k} \binom{nm+k}{k},$$

(8.8)
$$f_k(D_n, m) = \binom{n}{k} \binom{(n-1)m+k}{k} + \binom{n-2}{k-2} \binom{(n-1)m+k-1}{k}.$$

level	A_n	B_n	D_n	E_6	E_7	E_8	F_4	H_3	H_4	$I_2(a)$
1	1	1	1	1	1	1	1	1	1	1
2	2	3						5		a-1
3	3	5	3	4	5	7	5,7	9	11, 19	
4	4	7	5	5	7		11		29	
5	5	9	7	7,8	9	11, 13				
6	6	11	9	11	11, 13	17				
7	7	13	11		17	19, 23				
8	8	15	13			29				
:	:	:	÷							
n-1	n-1	2n-3	2n-5, n-1							
n	n	2n-1	2n-3							

Table 1. Levels of exponents

The type A_n case of Theorem 8.5 is due to J. H. Przytycki and A. S. Sikora [22]. The type B_n case is due to E. Tzanaki [29]. The formula (8.8) appeared in [30] in connection with a simplicial complex non-isomorphic to $\Delta^m(D_n)$.

For comparison, Table 2 also includes the factors $c_f(\Phi, k, m)$ for the type D_8 .

k	D_8	E_6	E_7	E_8	F_4	H_4
0	1	1	1	1	1	1
1	1	1	1	1	1	1
2	$\frac{29m}{8} + 1$	$\frac{14m}{5} + 1$	$\frac{7m}{2} + 1$	$\frac{35m}{8} + 1$	$\frac{13m}{6} + 1$	$\frac{31m}{12} + 1$
3	$\frac{31m}{12} + 1$	$\frac{9m}{4} + 1$	$\frac{27m}{10} + 1$	$\frac{45m}{14} + 1$	1	1
4	$\frac{17m}{8} + 1$	$\frac{5m}{3} + 1$	$\frac{21m}{10} + 1$	$\frac{46m^2}{7} + \frac{179m}{35} + 1$	1	1
5	$\frac{19m}{10} + 1$	1	$\frac{23m}{14} + 1$	2m + 1		
6	$\frac{43m}{24} + 1$	1	1	$\frac{13m}{8} + 1$		
7	1		1	1		
8	1			1		

Table 2. The factors $c_f(\Phi, k, m)$

The constant term of the factor $c_f(\Phi, k, m)$ as a polynomial in m is always 1. The degree of $c_f(\Phi, k, m)$ is equal to k minus the number of exponents at levels $\leq k$. This degree is ≤ 1 (i.e., $c_f(\Phi, k, m)$ equals 1 or is linear) in every case except for $c_f(E_8, 4, m)$, which is quadratic in m.

We call the face numbers $f_k(A_n, m)$ for the generalized cluster complex of type A_n the Przytycki-Sikora numbers. Thus, $f_k(A_n, m)$ is the number of k-tuples of pairwise-noncrossing diagonals that dissect a given convex ((n+1)m+2)-gon \mathbf{P} into smaller ((l+1)m+2)-gons $(0 \le l \le n)$.

The Przytycki-Sikora numbers (8.6) generalize both the Fuss numbers

(8.9)
$$N(A_n, m) = \frac{1}{n+1} \binom{(n+1)(m+1)}{n} = \frac{1}{nm+m+1} \binom{(n+1)(m+1)}{n+1},$$

which count the maximal simplices in $\Delta^m(A_n)$, and the Kirkman-Cayley numbers

(8.10)
$$f_k(A_n, 1) = \frac{1}{k+1} \binom{n}{k} \binom{n+k+2}{k},$$

the face numbers of an ordinary associahedron.

Both the Fuss numbers and the Kirkman-Cayley numbers (each of which in turn generalize the Catalan numbers) come up in many algebraic and combinatorial contexts; see, for example, [22, 24, 26]. For example, the Kirkman-Cayley numbers count plane trees with n + k + 3 vertices, n + 2 of which are leaves, such that no vertex has exactly one successor [28, Exercise 6.33.c]. The Fuss numbers:

- are dimensions of the Fuss-Catalan algebras of V. Jones and D. Bisch [5];
- count noncrossing partitions into blocks of size m+1 (see, e.g., [22]);
- count paths on the coordinate plane which connect the points (0,0) and ((m+1)n,0), have steps of the form (1,1) and (1,-m), and never dive below the x-axis (see, e.g., [15]);
- count plane (m+1)-ary trees with (n+1)(m+1)+1 vertices (see, e.g., [28, Proposition 6.2.1]);

• count plane rooted trees with (m-1)[(n+1)(m+1)+1] black vertices and (n+1)(m+1)+1 white vertices such that every black vertex is a leaf and every white vertex has m-1 black successors (see, e.g., [18]).

The Przytycki-Sikora numbers have interpretations in most of these contexts. It would be interesting to find analogues of these interpretations for other classical types (or even arbitrary finite types).

We conclude this section by a curious observation. The following result appeared in [29, Corollaries 3.2 and 3.4], with a different proof.

Corollary 8.6. In the combinatorial model of the complex $\Delta^m(B_n)$ presented in Section 5.2, the number of k-element simplices that include a diameter is equal to $\binom{n-1}{k-1}\binom{nm+k}{k}$.

Proof. The number of diameters in **P** is nm + 1. By formula (8.6), the number of k-simplices in $\Delta^m(B_n)$ which include a diameter is equal to

$$(nm+1)f_{k-1}(A_{n-1},m) = \frac{nm+1}{k} \binom{n-1}{k-1} \binom{nm+k}{k-1} = \binom{n-1}{k-1} \binom{nm+k}{k},$$
 as claimed.

Since

$$\binom{n-1}{k-1}\binom{nm+k}{k} = \frac{k}{n}\binom{n}{k}\binom{nm+k}{k} = \frac{k}{n}f_k(B_n, m),$$

Corollary 8.6 can be restated as saying that the proportion of k-element simplices in $\Delta^m(B_n)$ that include a diameter is equal to $\frac{k}{n}$. The latter statement has the following probabilistic reformulation.

Corollary 8.7. Fix k. Choose a k-element simplex in $\Delta^m(B_n)$ uniformly at random. It consists of k pairs of centrally symmetric (possibly identical) diagonals. Pick one of these k uniformly at random. Then the probability that it is a diameter is equal to $\frac{1}{n}$.

It would be interesting to find a direct probabilistic proof of Corollary 8.7, leading to a simpler proof of (8.7).

9. Proof of Theorem 8.5

Exceptional types E_6 , E_7 , F_4 , G_2 , H_3 , and H_4 . For each of these types, Theorem 8.5 has been verified on a computer by calculating the face numbers $f_k(\Phi, m)$ using the recurrences in Proposition 8.3.

Type
$$I_2(a)$$
. Immediate from Example 8.2.

Type A_n . In light of the isomorphism between $\Delta(m, n)$ and $\Delta^m(A_n)$ (see Section 5.1), formula (8.6) is equivalent to [22, Corollary 2].

Type B_n . In view of the combinatorial model for $\Delta^m(B_n)$ presented in Section 5.2, formula (8.7) is equivalent to [29, Theorem 1.2].

The proofs for the types A_n and B_n given in [22] and [29] are combinatorial (bijective). Rather than deducing (8.6) and (8.7) directly from the recurrences in Proposition 8.3, those proofs proceed by establishing bijections with combinatorial objects

that are easily enumerated by the right-hand-sides of (8.6) and (8.7). Our proof for the type D_n presented below is combinatorial as well.

Type D_n . This is the hardest case. The proof given here relies in large part on ideas developed by E. Tzanaki in an earlier analysis [30] of an alternative type-D analogue of $\Delta(m, n)$.

We will work with the combinatorial model of $\Delta^m(D_n)$ described in Section 5.3. We first define, for each diameter or diagonal pair, a starting vertex. For a diameter, the starting vertex is always the positive vertex (i.e., P_l as opposed to $-P_l$). A diagonal pair has four vertices, and the starting vertex is the unique vertex which is positive and from which one can travel along a diagonal in the pair while keeping the center of \mathbf{P} on the left. We will need the following lemma about the maximal faces of $\Delta^m(D_n)$ which consist entirely of diameters.

Lemma 9.1. For n > 2, given n diameters of **P**, the following are equivalent.

- (i) There exists a choice of flavor for each diameter such that the flavored diameters are pairwise compatible.
- (ii) If $1 \le a_1 \le a_2 \le \cdots \le a_n \le (n-1)m+1$ are the indices of the starting vertices of the diameters, then for each $1 \le j \le n$, we have $a_{j+1} a_j \le m$ (with the convention $a_{n+1} = a_1 + (n-1)m+1$).

If these conditions hold, then there are exactly two ways to assign flavors to the diameters so that they are pairwise compatible. These two ways of assigning flavors are related by switching the flavor of each of the n diameters.

Proof. First consider the case $a_1=1$, so that one of these diameters is primary. All of the diameters must be of the same flavor if they are all to be compatible with a diameter in primary position. Take any two of the diameters not in primary position and let i and k be the indices of their starting vertices, with i < k. Then the two (same-flavored) diameters are compatible if and only if, when $[P_i, -P_i]$ is moved to the primary position by R_m , and R_m is applied to $[P_j, -P_j]$ the same number of times, then both of them experience the same number of color changes (mod 2). As noted in Section 7, this occurs if and only if the fractional part of $\frac{i-2}{m}$ is greater than or equal to the fractional part of $\frac{j-2}{m}$. Thus the n diameters are pairwise compatible (when flavored identically) if and only if

frac. part
$$\left(\frac{a_2-2}{m}\right) \ge$$
 frac. part $\left(\frac{a_3-2}{m}\right) \ge \cdots \ge$ frac. part $\left(\frac{a_n-2}{m}\right)$.

Since $a_1 = 1$, this is now easily checked to be equivalent to (ii). Furthermore, there are two choices of flavorings, either all gray or all dashed.

If $a_1 > 1$, then rotate the set of diameters clockwise until $a_1 = 1$ and apply the argument above. If the rotated diagonals are pairwise compatible when flavored identically, apply R_m^{-1} repeatedly to undo the rotation. This will define a flavoring of the n diagonals such that they are pairwise compatible.

We are now prepared to complete the proof of (8.8). Given a face F in $\Delta^m(D_n)$ containing at least one diameter, call F a gray face if for the minimal i such that F contains a diameter in position $[P_i, -P_i]$, there is a gray diameter in that position.

Similarly define a *dashed face*. A face with two coinciding diameters is both gray and dashed.

Following [30], we rewrite (8.8) as the sum of the expressions

(9.1)
$${n-1 \choose k} {(n-1)m+k \choose k} + {n-2 \choose k-2} {(n-1)m+k \choose k}, \text{ and }$$

$$(9.2) 2\binom{n-2}{k-2}\binom{(n-1)m+k}{k} - \binom{n-2}{k-2}\binom{(n-1)m+k-1}{k-1},$$

and identify the terms of these expressions. The first term of (9.1) is $f_k(B_{n-1}, m)$, which also counts the k-element faces of $\Delta^m(D_n)$ with at most one gray diameter and no dashed diameters. By [29, Corollary 3.4], the second term is the number of faces of $\Delta^m(B_{n-1})$ with k elements, exactly one of which is a diameter. This also counts the k-element faces of $\Delta^m(D_n)$ with exactly one dashed diameter and no gray diameters. Thus (9.1) counts faces of $\Delta^m(D_n)$ with at most one diameter. Therefore (9.2) counts faces of $\Delta^m(D_n)$ with two or more diameters. The second term of (9.2) is the number of faces of $\Delta^m(B_{n-1})$ with k-1 elements, exactly one of which is a diameter, which also counts the k-element faces of $\Delta^m(D_n)$ with a pair of coinciding diameters. Thus the first term of (9.2) is the number of gray k-element faces with two or more diameters plus the number of dashed k-element faces with two or more diameters. (This double-counts the faces with a pair of coinciding diameters.)

To complete the proof, it remains to show that the number of gray (respectively dashed) k-element faces containing two or more diameters is

$$\binom{n-2}{k-2}\binom{(n-1)m+k}{k}.$$

This is done by giving a bijection between gray faces with two or more diameters and pairs

$$((\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_{n-2}), (a_1, a_2, \ldots, a_k)),$$

where each ε_i is either 0 or 1, with $\varepsilon_i = 1$ exactly k-2 times, and the a_j satisfy $1 \le a_1 \le a_2 \le \cdots \le a_k \le (n-1)m+1$. Given a gray face F with k vertices, including two or more diameters, the sequence of a_j 's is the set of indices of the starting vertices of the diameters or diagonal pairs in F, written in weakly increasing order.

For k=2, nothing more is needed to define the map, so assume k>2. Now, take the smallest $j \in [k]$ such that $a_{j+1}-a_j>m$ (if such a j exists), where a_{k+1} is understood to mean $a_1+(n-1)m+1$. If the diagonal $[P_{a_j}, P_{a_j+m+1}]$ is in F then set $\varepsilon_1=1$. Otherwise set $\varepsilon_1=0$. The remaining k-1 of k vertices of F are a face in $\Delta^m(D_{n-1})$, realized in the polygon \mathbf{P}' obtained from \mathbf{P} by deleting the vertices

$$\pm P_{a_j+1}, \pm P_{a_j+2}, \dots, \pm P_{a_j+m}.$$

By induction, determine the remaining ε_i 's.

If there is no $j \in [k]$ with $a_{j+1} - a_j > m$, then we must have k = n, so set all the ε_i 's to be 1. Furthermore, in this case, all of the elements of F must be diagonals.

To see that this is a bijection, we define an inverse. Suppose we are given a pair $((\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{n-2}), (a_1, a_2, \dots, a_k))$ and we wish to construct a face F. If k = 2, then F consists of the gray diameter $[P_{a_1}, -P_{a_1}]$ and the diameter $[P_{a_2}, -P_{a_2}]$, the latter flavored in the unique way which makes the two compatible. Otherwise, take

the smallest $j \in [k]$ such that $a_{j+1} - a_j > m$ (if such a j exists). If $\varepsilon_1 = 1$ then use $((\varepsilon_2, \ldots, \varepsilon_{n-2}), (a_1, a_2, \ldots, \hat{a}_j, \ldots, a_k))$ to inductively determine a (k-1)-element face F' of $\Delta_m(D_{n-1})$ realized in the polygon \mathbf{P}' described above. Then F is the face of $\Delta_m(D_n)$ obtained by adjoining the diagonal $[P_{a_j}, P_{a_j+m+1}]$ and its symmetric diagonal to F'. If $\varepsilon_1 = 0$, then use $((\varepsilon_2, \ldots, \varepsilon_{n-2}), (a_1, a_2, \ldots, a_k))$ to inductively determine a k-element face F' of $\Delta_m(D_{n-1})$ realized in \mathbf{P}' . Then F is the face of $\Delta_m(D_n)$ which agrees with F'.

If no such j exists, then n must equal k. Consider the collection of n diagonals whose starting vertices are given by the a_i 's. By Lemma 9.1, there is a unique way of assigning flavors to each of the diameters in the collection so as to obtain a gray face of $\Delta_m(D_n)$.

10. h-vectors

We define the f-polynomial $F(\Phi, m, x)$ in the formal variable x by

$$F(\Phi, m, x) = \sum_{k=0}^{n} f_{n-k}(\Phi, m) x^{k}.$$

We then define the h-polynomial $H(\Phi, m, x)$ by

$$H(\Phi, m, x) = F(\Phi, m, x - 1).$$

The corresponding h-vector $(h_0, \ldots, h_n) = (h_0(\Phi, m), \ldots, h_n(\Phi, m))$ is then defined by

$$H(\Phi, m, x) = \sum_{k=0}^{n} h_{n-k}(\Phi, m) x^{k}.$$

For m=1, we recover the h-vector of the associahedron of type Φ , whose coefficients are the (generalized) Narayana numbers of type Φ . These Narayana numbers have been shown [13, 19, 23, 25] to count various combinatorial objects associated with the root system Φ , the Coxeter group W, the corresponding semisimple Lie algebra, etc. See [12, Section 5.2] or [3] for a survey of these results.

For a general m, the numbers $h_k(\Phi, m)$ turn out to coincide with the m-analogues of the Narayana numbers introduced and studied by C. A. Athanasiadis [3]. The latter numbers, herein denoted $H_k(\Phi, m)$, count various objects associated with a root system Φ and a positive integer m according to certain statistics whose values are recorded by k. We next sketch the definition of the numbers $H_k(\Phi, m)$, referring the reader to [3] for further details and references.

The extended Catalan arrangement $\operatorname{Cat}_{\Phi}^m$ (see [1, 20]) is the collection of affine hyperplanes $H_{\alpha,j}$ defined by the equations $\langle \alpha, x \rangle = j$ for $\alpha \in \Phi$ and $j = 0, 1, \ldots, m$. The discrete torus T^m is the coroot lattice \check{Q} of Φ modulo its dilation $(mh+1)\check{Q}$. The group W naturally acts on T^m ; the rank of a W-orbit is the rank (as a reflection group) of the stabilizer of any point in the orbit. The root poset of Φ is the partial order on $\Phi_{>0}$ such that $\beta \leq \gamma$ if and only if $\gamma - \beta$ is a nonnegative linear combination of simple roots.

The number $H_k(\Phi, m)$ counts:

• the regions R of $\operatorname{Cat}_{\Phi}^m$ in the fundamental chamber such that n-k walls of R of the form $H_{\alpha,m}$ separate R from the fundamental alcove;

- the W-orbits of rank n-k in T^m ;
- collections of nested order filters in the root poset of Φ satisfying certain technical conditions (see [3, (1.2)–(1.3)]) and having n-k indecomposable elements of rank m.

Conjecture 10.1. For any finite crystallographic root system Φ , we have $h_k(\Phi, m) = H_k(\Phi, m)$.

Conjecture 10.1 can be verified for the classical series ABCD by comparing the formulas (10.1)–(10.3) below with their counterparts in [3]. This observation has already been made in [30]. The conjecture has not been completely verified because the numbers $H_k(\Phi, m)$ have not been computed for the exceptional types.

The h-numbers $h_k(\Phi, m)$ are given by multiplicative formulas very similar to the corresponding formulas for the face numbers.

Theorem 10.2. For an irreducible root system Φ and an integer $0 \le k \le n$, we have

$$h_k(\Phi, m) = c_h(\Phi, k, m) \binom{n}{k} \prod_{\substack{\text{level}(e_i) \le k}} \frac{mh - e_i + 1}{e_i + 1},$$

where the factor $c_h(\Phi, k, m)$ is a polynomial in m given as follows:

- in types A_n , B_n , H_3 , and $I_2(m)$, we have $c_h(\Phi, k, m) = 1$;
- in type D_n , we have

$$c_h(D_n, k, m) = \begin{cases} 1 & \text{if } k \in \{0, 1, n-1, n\}; \\ \frac{(n^2 - n + k^2 - k)(mn - m + 1)}{kn(n - 1)} - 1 & \text{otherwise}; \end{cases}$$

• in types E_6 , E_7 , E_8 , F_4 , and H_4 , the factors $c_h(\Phi, k, m)$ are shown in Table 3. In particular, for the types A_n , B_n , and D_n , we have:

(10.1)
$$h_k(A_n, m) = \frac{1}{k+1} \binom{n}{k} \binom{(n+1)m}{k}$$

(10.2)
$$h_k(B_n, m) = \binom{n}{k} \binom{nm}{k}$$

(10.3)
$$h_k(D_n, m) = \binom{n}{k} \binom{(n-1)m}{k} + \binom{n-2}{k-2} \binom{(n-1)m+1}{k}.$$

Proof. As in the case of Theorem 8.5, the type $I_2(a)$ is an easy calculation, and the types E_6 , E_7 , F_4 , G_2 , H_3 , and H_4 can be verified by computer. The formulas in the classical types ABD can be derived from their counterparts in Theorem 8.5 using Chu-Vandermonde summation. Details are omitted.

11. Euler Characteristic

Recall that the reduced Euler characteristic of a simplicial complex Δ that has f_k faces of dimension k-1, for $k=0,1,\ldots,n$ (including one empty face for $f_0=1$), is defined by

$$\tilde{\chi}(\Delta) = \sum_{k \ge 0} (-1)^{k-1} f_k.$$

k	D_8	E_6	E_7	E_8	F_4	H_4
0	1	1	1	1	1	1
1	1	1	1	1	1	1
2	$\frac{203m-27}{56}$	$\frac{42m-8}{15}$	$\frac{63m-11}{18}$	$\frac{105m-17}{24}$	$\frac{78m-23}{36}$	$\frac{465m-149}{180}$
3	$\frac{217m - 53}{84}$	$\frac{18m-5}{8}$	$\frac{27m-7}{10}$	$\frac{45m-11}{14}$	1	1
4	$\frac{119m - 39}{56}$	$\frac{30m-13}{18}$	$\frac{63m-23}{30}$	$\frac{10350m^2 - 6675m + 1084}{1575}$	1	1
5	$\frac{133m-51}{70}$	1	$\frac{207m-103}{126}$	$\frac{30m-13}{15}$		
6	$\frac{301m-125}{168}$	1	1	$\frac{195m - 107}{120}$		
7	1		1	1		
8	1			1		

Table 3. The factors $c_h(\Phi, k, m)$

Thus $\tilde{\chi}(\Delta) = \chi(\Delta) - 1$, where $\chi(\Delta)$ is the usual Euler characteristic. It is well-known that the Euler characteristic of an (n-1)-dimensional complex is $(-1)^{n-1}h_n$. Using the k = n case of Theorem 10.2 and the fact that $e_i = h - e_{n-i}$, we obtain:

Proposition 11.1. The reduced Euler characteristic of $\Delta^m(\Phi)$ is given by

(11.1)
$$\tilde{\chi}(\Delta^m(\Phi)) = (-1)^{n-1} N(\Phi, m-1).$$

Thus, it is equal, up to a sign, to the number of maximal simplices in the complex $\Delta^{m-1}(\Phi)$.

Formula (11.1) can be restated as

(11.2)
$$\sum_{k=0}^{n} (-1)^{n-k} f_k(\Phi, m) = N(\Phi, m-1).$$

In the special case m=1, Proposition 11.1 is a corollary of the following statement proved in [13]: the simplicial complex $\Delta^1(\Phi)$ is homeomorphic to an (n-1)-dimensional sphere. Note that $N(\Phi,0)=1$ (see Example 4.1).

Remark 11.2. Our proof of Proposition 11.1 relies on a type-by-type calculation. Is there a conceptual and/or geometric proof? For example, there may be a way to remove $N(\Phi, m-1)$ open facets from $\Delta^m(\Phi)$ and obtain a contractible complex.

The following conjecture was suggested by our discussions with Hugh Thomas in December 2004.

Conjecture 11.3. The complex $\Delta^m(\Phi)$ is shellable, hence Cohen-Macaulay, and homotopy equivalent to a wedge of spheres.

Conjecture 11.3 has been proved by E. Tzanaki [29, Theorem 1.4] in types A_n and B_n .

12. RECIPROCAL FACE NUMBERS

Inspired by the construction of "positive clusters" in [13], we define the reciprocal face numbers $f_k^+(\Phi, m)$ as follows. Since $f_k(\Phi, m)$ is a polynomial in m, one can extend the definition of $f_k(\Phi, m)$ to the non-positive integer values of m. Let us define

(12.1)
$$f_k^+(\Phi, m) = (-1)^k f_k(\Phi, -m-1).$$

For k = n, we set

(12.2)
$$N^{+}(\Phi, m) = f_{n}^{+}(\Phi, m) = (-1)^{n} N(\Phi, -m - 1).$$

Our choice of notation N^+ is explained by Proposition 12.4 below.

The definition (12.2) combined with Propositions 8.3 and 8.4 yields the following formulas.

Corollary 12.1. For a reducible root system $\Phi = \Phi_1 \times \Phi_2$,

(12.3)
$$N^{+}(\Phi_{1} \times \Phi_{2}, m) = N^{+}(\Phi_{1}, m) N^{+}(\Phi_{2}, m).$$

If Φ is irreducible, then

(12.4)
$$N^{+}(\Phi, m) = \frac{(m+1)h-2}{2n} \sum_{i \in I} N^{+}(\Phi_{\langle i \rangle}, m).$$

Corollary 12.2. For an irreducible root system Φ ,

(12.5)
$$N^{+}(\Phi, m) = \prod_{i} \frac{mh + e_{i} - 1}{e_{i} + 1}$$

We note that the recurrence (12.4) is new even for m=1, with the numbers $N^+(\Phi)=N^+(\Phi,1)$ counting positive clusters in $\Delta(\Phi)$ (cf. [13, Proposition 3.9]).

Analogues of (12.4) and (12.5) for the general numbers $f_k^+(\Phi, m)$ can be obtained directly from (12.1) and similar formulas for the face numbers $f_k(\Phi, m)$.

Proposition 12.3. We have

(12.6)
$$N(\Phi, m) = \sum_{J \subseteq I} N^{+}(\Phi_{J}, m),$$

where, by convention, $N^+(\Phi_J, m) = 1$ for the empty subset $J = \emptyset$.

Proof. As shown in [2, Corollary 1.3] (cf. discussion in Section 10), the number $N(\Phi, m)$ (resp., $N^+(\Phi, m)$) counts all (resp., bounded) regions of the extended Catalan arrangement $\operatorname{Cat}_{\Phi}^m$ contained in the fundamental chamber of the corresponding Coxeter arrangement. Then (12.6) follows, for example, from [4, Lemma 5.3].

The following generalization of [13, Proposition 3.9] provides a direct combinatorial interpretation of the numbers $N^+(\Phi, m)$ in terms of generalized cluster complexes.

Proposition 12.4. The number of maximal simplices in $\Delta^m(\Phi)$ which only involve positive colored roots is equal to $N^+(\Phi, m)$.

Proof. Let $\tilde{N}^+(\Phi, m)$ denote the number of maximal simplices in $\Delta^m(\Phi)$ consisting of positive colored roots. Theorem 3.7 implies that $N(\Phi, m) = \sum_{J \subseteq I} \tilde{N}^+(\Phi_J, m)$, and the claim follows by Proposition 12.3.

Remark 12.5. It is natural to suggest that the relationship (12.2) between the quantities $N(\Phi, m)$ and $N^+(\Phi, m)$ is an instance of Ehrhart reciprocity [27, Theorem 4.6.26. This suggestion has been confirmed by C. A. Athanasiadis and E. Tzanaki in [4, Section 7].

13. Combinatorics of Coxeter-Theoretic invariants

A Coxeter diagram is an undirected graph G without loops and multiple edges, in which every edge is labeled by an integer ≥ 3 . (In general Coxeter group theory, labels equal to ∞ are also allowed, but we restrict our attention to integer labels.) By convention, each missing edge (i.e., each pair of vertices in G not connected by an edge) is thought of as an edge labeled by 2. The number of vertices in a Coxeter diagram is called its rank.

Let G be a Coxeter diagram of rank n for an irreducible finite Coxeter group W. There is a host of Coxeter-theoretic invariants associated with G and the corresponding root system Φ :

- the exponents e_1, \ldots, e_n ;
- the Coxeter number $h = \frac{2}{n} \sum_{i=1}^{n} e_i$; the number of roots $|\Phi| = nh$;
- the order of the group $|W| = \prod (e_i + 1)$;
- the generalized Catalan number $N(\Phi) = \prod \frac{h + e_i + 1}{e_i + 1}$;
- the face numbers $f_k(G, m)$ of the simplicial complex $\Delta^m(\Phi, m)$;

and many others. Our goal in this section is to develop purely combinatorial procedures for computing all these invariants directly from G, i.e., procedures that do not explicitly involve the root system Φ , the Coxeter group W, or any other algebraic or Lie-theoretic concepts.

Remark 13.1. Since all quantities listed above can be expressed in terms of the exponents e_1, \ldots, e_n , computing the exponents would accomplish the task. However, determining them directly from the Coxeter diagram G is not at all straightforward.

In turn, the computation of exponents can be reduced to the calculation of a Coxeter number, as follows. Use Proposition 8.3 to recursively compute the generalized Fuss number

$$N(G, m) = N(\Phi, m) = \prod \frac{mh + e_i + 1}{e_i + 1}.$$

Then, viewing N(G,m) as a polynomial in m, find its roots $-\frac{e_i+1}{h}$ and determine the exponents e_i .

Remark 13.2. As a rule of thumb, any use of a "Coxeter number" or "exponents" in the reducible case should be viewed with skepticism. In particular, the formulas above in this section expressing other invariants in terms of the Coxeter number and/or the exponents make no sense in the reducible case. In what follows, we do not consider Coxeter numbers or exponents for any disconnected diagrams.

We will describe several combinatorial algorithms for computing invariants of Coxeter diagrams associated with finite Coxeter groups. Each of these algorithms can then be used to define "fake" Coxeter-theoretic invariants of more general Coxeter diagrams. Such generalizations are discussed in Section 14.

All our algorithms are recursive: the computation of invariants of G relies upon prior calculation of similar invariants for induced subgraphs of G. The base of recursion is provided by the special cases $n \leq 2$, together with setting $f_0(G, m) = 1$ for any G.

For n = 0, $G = \emptyset$, we have

$$N(G, m) = N^{+}(G, m) = 1.$$

For n = 1, $G = (\bullet)$ (see Example 8.1), we postulate:

(13.1)
$$h = 2,$$

$$e_1 = 1,$$

$$N(G, m) = f_1(G, m) = m + 1,$$

$$N^+(G, m) = m.$$

For n=2, $G=(\bullet \stackrel{a}{----} \bullet)$ (see Example 8.2), we postulate:

(13.3)
$$h = a,$$

$$\{e_1, e_2\} = \{1, a - 1\},$$

$$f_1(G, m) = am + 2,$$

(13.4)
$$N(G,m) = f_2(G,m) = \frac{(am+2)(m+1)}{2},$$

(13.5)
$$N^{+}(G,m) = \frac{(am+a-2)m}{2}.$$

We note that formulas (13.3)–(13.5) hold just as well in the reducible case a=2, where G is of type $A_1 \times A_1$, and we have $f_1(A_1 \times A_1) = 2m + 2$ and $N(A_1 \times A_1) = (m+1)^2$.

We next present four different combinatorial procedures, each of which determines the invariants of a connected Coxeter diagram G of rank $n \geq 3$ assuming that such invariants for all proper induced connected subgraphs of G are already known.

13.1. The Euler characteristic method. We start with the basic recurrence (8.2):

(13.6)
$$f_k(G,m) = \frac{mh+2}{2k} \sum_{G' \leq G} f_{k-1}(G',m),$$

where $k \in \{1, ..., n\}$ and the summation is over all induced subgraphs $G' \subset G$ obtained by removing one vertex from G. Whenever G' is disconnected, with connected components $G_1, ..., G_s$, we use the formula

(13.7)
$$f_{k-1}(G',m) = \sum_{k_1 + \dots + k_s = k-1} \prod_{i=1}^s f_{k_i}(G_i,m)$$

(see (8.1)) to calculate $f_{k-1}(G', m)$.

We note that the number h = h(G) is undetermined as of yet. To find the value of h, we use (11.2):

(13.8)
$$\sum_{k=0}^{n} (-1)^{n-k} f_k(G, m) = f_n(G, m-1).$$

Substituting (13.6) into (13.8) gives a linear equation for h: (13.9)

$$(-1)^n + \sum_{G' \leq G} \left(-\frac{(m-1)h+2}{2n} f_{n-1}(G', m-1) + (mh+2) \sum_{k=1}^n \frac{(-1)^{n-k}}{2k} f_{k-1}(G', m) \right) = 0.$$

A priori, solving this equation for h results in a rational function in m. However, if G is a Coxeter diagram for a finite Coxeter group (and in some other cases as well), the answer miraculously turns out to be a constant—in fact, a positive integer.

Example 13.3 (n = 3). Let G be a connected Coxeter diagram of rank 3 with edge labels $a_1, a_2, a_3 \ge 2$, as shown in Figure 10.




Figure 10

We denote $a = a_1 + a_2 + a_3$. The values of a corresponding to finite irreducible Coxeter groups are shown in Table 4.

type of G	A_3	B_3	H_3
a	8	9	10

Table 4

The "maximal parabolic" subgraphs $G' \leq G$ are of types $I_2(a_1)$, $I_2(a_2)$, and $I_2(a_3)$, respectively. Their contributions to the outer sum in (13.9) are of the form

$$-\frac{(m-1)h+2}{6}f_2(I_2(a_i), m-1) + (mh+2)\sum_{k=1}^3 \frac{(-1)^{3-k}}{2k} f_{k-1}(I_2(a_i), m)$$

$$= -\frac{(m-1)h+2}{6} \frac{(a_i(m-1)+2)m}{2}$$

$$+ (mh+2)\left(\frac{1}{2} - \frac{1}{4}(a_im+2) + \frac{1}{6} \frac{(a_im+2)(m+1)}{2}\right)$$

$$= \frac{-a_imh - 2a_im + 4mh + 4}{12}.$$

Thus (13.9) becomes

$$-1 - \frac{(a_1 + a_2 + a_3)m(h+2)}{12} + mh + 1 = 0$$

yielding

$$(13.10) h = \frac{2a}{12 - a}.$$

We can now use this value of h to compute the face numbers by means of (13.6):

(13.11)
$$f_1(G,m) = \frac{3(am-a+12)}{12-a},$$

(13.12)
$$f_2(G,m) = \frac{(am-a+12)(am+6)}{2(12-a)},$$

(13.13)
$$N(G,m) = f_3(G,m) = \frac{(m+1)(am+6)(am-a+12)}{6(12-a)}$$

The roots of N(G, m) are $\{\mu_1, \mu_2, \mu_3\} = \{-1, -\frac{6}{a}, -\frac{12-a}{a}\}$. Since $\mu_i = -\frac{e_i+1}{h}$, we recover the exponents by $e_i = -h\mu_i - 1$:

(13.14)
$$\{e_1, e_2, e_3\} = \left\{1, \frac{a}{12 - a}, \frac{3a - 12}{12 - a}\right\}.$$

We also use (12.2) to get

(13.15)
$$N^{+}(G,m) = \frac{m(am+a-6)(am+2a-12)}{6(12-a)}.$$

13.2. Symmetry-based method. This method does not require a recursive computation of all polynomials $f_k(G, m)$, but only N(G, m). It exploits the fact that the set of exponents is invariant with respect to the reflection

$$e_i \mapsto h - e_i$$
.

Equivalently, the set $\{\mu_i\}$ of (simple) roots of the polynomial N(G, m) is invariant with respect to the reflection

$$\mu_i \mapsto -\frac{h+2}{h} - \mu_i.$$

Recall once again the recurrence

(13.17)
$$N(G,m) = \frac{mh+2}{2n} \sum_{G' \leq G} N(G',m).$$

It follows by induction on n that N(G,-1)=0 for any non-empty G. Hence

(13.18)
$$Q(G,m) = \frac{1}{m+1} \sum_{G' \leq G} N(G',m)$$

is a polynomial in m (of degree n-2). Furthermore, its set of roots is still invariant under the reflection (13.16), since we removed two symmetric roots -1 and $-\frac{2}{h}$ from N(G, m).

Note that the computation of Q(G, m) by the formula (13.18) does not require the value of h. So we find the polynomial Q(G, m), and then determine the average of

its roots. This average is equal to $-\frac{h+2}{2h}$, so the negative sum of the n-2 roots of Q(G,m) is equal to

(13.19)
$$\frac{\text{coefficient of } m^{n-3} \text{ in } Q(G,m)}{\text{coefficient of } m^{n-2} \text{ in } Q(G,m)} = \frac{(n-2)(h+2)}{2h}.$$

We then find h from this linear equation, and determine the exponents.

Example 13.4 (n = 3). Let G be a connected Coxeter diagram of rank 3 with edge labels a_1, a_2, a_3 (cf. Example 13.3). As before, we use the notation $a = a_1 + a_2 + a_3$. Combining (13.18) with (13.4), we obtain:

$$Q(G,m) = \frac{1}{m+1} \sum_{i=1}^{3} \frac{(a_i m + 2)(m+1)}{2} = \frac{am+6}{2}.$$

Hence (13.19) becomes

$$\frac{6}{a} = \frac{h+2}{2h},$$

implying (13.10). We conclude that the roots of N(G, m) are $-\frac{6}{a}$ (the only root of Q(G, m)), -1 (always a root), and $-\frac{12-a}{a}$ (obtained from -1 using (13.16)). The exponents (13.14) are then found as in Example 13.3.

13.3. Reciprocity-based method. This method uses the inclusion-exclusion formula for the reciprocal face numbers $N^+(G, m)$ as the additional equation needed to recursively compute the numbers N(G, m) if the Coxeter number h is unknown. For the simplest version of this method, one only needs to recursively compute the numbers N(G) = N(G, 1) and $N^+(G) = N^+(G, 1) = (-1)^n N(G, -2)$. That is, this version only involves the numbers $f_k(G, m)$ for k = n and $m \in \{1, -2\}$.

The idea is very simple. Proposition 12.3 and formula (13.6) give:

(13.20)
$$N(G) = \frac{h+2}{2n} \sum_{G' \in G} N(G'),$$

(13.21)
$$N^{+}(G) = \frac{h-1}{n} \sum_{G' \leq G} N^{+}(G'),$$

(13.22)
$$N(G) = \sum_{H \subset G} N^{+}(H),$$

where H ranges over all induced subgraphs of G (including G and the empty subgraph). Since the values of N(H) and $N^+(H)$ for all proper subgraphs of G are presumed known, it remains to find the three unknowns N(G), $N^+(G)$, and h from the three linear equations (13.20)–(13.22).

Example 13.5 (n = 3). Let G be a connected Coxeter diagram of rank 3 with edge labels a_1, a_2, a_3 (cf. Examples 13.3 and 13.4).

The values of N(H) and $N^+(H)$ for graphs H of rank 1 or 2 are given by (13.1) and (13.4) with m = 1 and m = -2:

$$H=(ullet)$$
 $N(H)=2$ $N^+(H)=1$
$$H=(ullet -\frac{b}{} ullet)$$
 $N(H)=b+2$ $N^+(H)=b-1$

We continue to use the notation $a = a_1 + a_2 + a_3$. Equations (13.20)–(13.22) become:

$$N(G) = \frac{h+2}{6} (a+6),$$

$$N^{+}(G) = \frac{h-1}{3} (a-3),$$

$$N(G) = N^{+}(G) + (a-3) + 3 + 1.$$

Solving these equations for h, N(G), and $N^+(G)$ yields $h = \frac{2a}{12-a}$, $N(G) = \frac{4(a+6)}{12-a}$, and $N^+(G) = \frac{(a-3)(a-4)}{12-a}$, in agreement with (13.10), (13.13), and (13.15) (with m = 1). The exponents are found as before.

The more general version of the reciprocity-based method retains the parameter m. Let G be a connected Coxeter diagram of rank $n \geq 3$. As before, we start with the relevant versions of the basic recurrence (13.6), namely (13.17) and

(13.23)
$$N^{+}(G,m) = \frac{h(m+1)-2}{2n} \sum_{G' \leqslant G} N^{+}(G',m).$$

Substituting (13.22) into both sides of (13.17), we obtain:

$$\sum_{H \subset G} N^{+}(H, m) = \frac{mh + 2}{2n} \sum_{H \subset G} (n - |H|) N^{+}(H, m),$$

where |H| denotes the rank of H. Solving for $N^+(G,m)$, we get

(13.24)
$$N^{+}(G,m) = \sum_{H \subseteq G} \frac{mh}{2n} (n - |H|) N^{+}(H,m).$$

Equating the right-hand sides of (13.24) and (13.23) and solving for h yields

(13.25)
$$h = 2 \frac{(n-2) \sum_{G' \leqslant G} N^+(G', m) + \sum_{H \ll G} |H| N^+(H, m)}{m \sum_{H \ll G} (n - |H|) N^+(H, m) - \sum_{G' \leqslant G} N^+(G', m)},$$

where $H \ll G$ means that $H \subset G$ and |H| < |G| - 1.

13.4. The invariants M(G). We next present a simplified version of the reciprocity-based method that exploits the fact that the Coxeter diagrams of finite Coxeter groups are *sparse*, i.e., they have far fewer edges than $\binom{n}{2}$. The method is based on recursive computation of the following invariant.

Definition 13.6. For a non-empty Coxeter graph G of rank n, define

(13.26)
$$M(G) = n \lim_{m \to 0} \frac{N^+(G, m)}{m}.$$

If G is empty, set M(G) = 0.

Remark 13.7. Note that (13.2), (13.5), and (13.23) imply $N^+(G,0) = 0$, so M(G) is well defined. For the same reasons, M(G) = 0 if G is disconnected.

Corollary 13.8. If G is the Coxeter diagram of an irreducible finite Coxeter group with exponents e_1, \ldots, e_n and Coxeter number h, then

(13.27)
$$M(G) = \frac{nh}{2} \prod_{e_i \neq 1} \frac{e_i - 1}{e_i + 1}.$$

Proof. Follows from (12.5) and (13.26).

type of G	A_n	B_n	D_n	E_6	E_7	E_8	F_4	H_3	H_4	$I_2(a)$
M(G)	1	n	n-2	7	16	44	10	8	42	a-2

Table 5. Values of M(G) for finite Coxeter groups

The values of M(G) for all finite irreducible Coxeter groups are listed in Table 5. All of these values are integers. An explanation of this integrality is provided by the following interpretation of the invariants M(G), due to F. Chapoton.

Proposition 13.9 ([8]). Let W be a finite irreducible Coxeter group with Coxeter diagram G. Then the number of reflections in W which do not belong to any proper parabolic subgroup W_J is equal to M(G). Accordingly, the number of all reflections in W is equal to

(13.28)
$$\frac{nh}{2} = \sum_{H \subset G} M(H) \,.$$

Proposition 13.9 can be restated as saying that in a finite root system Φ , the number of positive roots whose simple root expansion has full support (i.e., involves every simple root) is equal to M(G), where G is the corresponding Coxeter diagram.

Proof. The inclusion-exclusion relation (13.28) uniquely defines the numbers M(G), and is satisfied by the number of reflections not lying in a proper parabolic subgroup. Consequently, it suffices to prove (13.28) in order to prove the Proposition.

Substituting (12.6) into both sides of (8.3), we obtain:

$$\sum_{H \in G} N^{+}(H, m) = \frac{mh + 2}{2n} \sum_{H \in G} (n - |H|) N^{+}(H, m),$$

implying

$$\frac{mh}{2} \sum_{H \subset G} N^{+}(H, m) = \frac{mh + 2}{2n} \sum_{H \subset G} |H| N^{+}(H, m).$$

Dividing by m and substituting m=0, we get (the only surviving term on the left-hand side is the one for $H=\emptyset$):

$$\frac{h}{2} = \frac{1}{n} \sum_{H \subset G} M(H) \,,$$

and (13.28) follows.

It would be interesting to find a direct proof of Proposition 13.9 based on Definition 13.6 and one of the combinatorial interpretations of $N^+(G, m)$, rather than on calculations involving product formulas and/or recurrence relations.

It follows from (12.4) that the numbers M(G) satisfy the recurrence

(13.29)
$$M(G) = \frac{h-2}{2(n-1)} \sum_{G' \leq G} M(G').$$

There is also a more complicated recurrence for these numbers that does not require knowing the Coxeter number h.

Lemma 13.10. Let G be the Coxeter diagram of a finite irreducible Coxeter group of rank $n \geq 3$. Set

(13.30)
$$\Sigma_1 = \sum_{G' \leqslant G} M(G'), \qquad \Sigma_2 = \sum_{\substack{H \subset G \\ 2 \leqslant |H| \leqslant n-1}} M(H).$$

Then

(13.31)
$$M(G) = \frac{\sum_{1} \sum_{2}}{n(n-1) - \sum_{1}}.$$

Proof. Identities (13.28) and (13.29) can be written as

(13.32)
$$h = \frac{2}{n} (M(G) + \Sigma_2 + n),$$

(13.33)
$$M(G) = \frac{h-2}{2(n-1)} \Sigma_1.$$

Substituting (13.32) into (13.33) yields (13.31).

Formulas (13.30)–(13.31) can be used to recursively compute the values M(G). In view of Remark 13.7, the sums in (13.30) can be restricted to *connected* subgraphs G' and H, simplifying the calculations.

The Coxeter numbers are then recovered using (13.28) or, equivalently, (13.32).

Example 13.11. For G of type $I_2(a)$, one has M(G) = a-2. For a 3-vertex diagram with edge labels a_1, a_2, a_3 (cf. Examples 13.3, 13.4 and 13.5), with $a = a_1 + a_2 + a_3$, we get $\Sigma_1 = \Sigma_2 = a - 6$. Substituting this into (13.31)–(13.32), we obtain

(13.34)
$$M(G) = \frac{(a-6)^2}{12-a}, \qquad h(G) = \frac{2a}{12-a},$$

matching the results of our earlier calculations.

Example 13.12. Now let

$$G = (\bullet \xrightarrow{b_1} \bullet \xrightarrow{b_2} \bullet \xrightarrow{b_3} \bullet)$$

be a linear Coxeter diagram of rank 4. (Here b_1, b_2, b_3 are integers ≥ 3 .) Us-

ing (13.34), we compute:

$$\Sigma_1 = \frac{(b_1 + b_2 - 4)^2}{10 - b_1 - b_2} + \frac{(b_2 + b_3 - 4)^2}{10 - b_2 - b_3},$$

$$\Sigma_2 = b_1 + b_2 + b_3 - 6 + \Sigma_1.$$

The values of M(G) and h(G) are then obtained using (13.31)–(13.32). The resulting formula for the Coxeter number h(G) is a cumbersome rational expression in b_1, b_2, b_3 . A diligent reader might want to verify that, in the special cases where G is of types A_4 , B_4 , F_4 , or H_4 , the outcomes coincide with the known values of Coxeter numbers.

14. Fake Coxeter invariants

Each of the methods described in Section 13 can, in principle, be applied to a very general Coxeter diagram G. The resulting "fake Coxeter invariants" may or may not be meaningful. Be that as it may, it is tempting to calculate these invariants in the simplest non-finite cases; to catalogue the infinite types where such invariants are "nice;" and possibly find intrinsic interpretations for them. One also wonders what the limits of applicability are for each of the methods of Section 13, and how their outputs compare if more than one of them works.

Examples 13.3, 13.4, 13.5, and 13.11 illustrate the methods in the n=3 case, and provide the first example in which all of them break down: when a=12, the denominator in the formula for h vanishes. (For a>12, we get $h=\frac{2a}{12-a}<0$, a negative "fake Coxeter number.") Here is another such example.

Example 14.1. Let $G = K_4$ be the complete graph on 4 vertices, with all 6 edges labeled by 3. Then each 3-vertex subgraph G' < G is of affine type \tilde{A}_2 , so (13.34) gives M(G') = 3. Then (13.30) yields $\Sigma_1 = 4 \cdot 3 = 12$, and the denominator in (13.31) vanishes. So the method based on Lemma 13.10 does not work for $G = K_4$. In fact, none of the methods described above will work in this case. Consequently, the methods will fail for any Coxeter diagram that has K_4 as an induced subgraph. Another similar example is the 4-cycle with edge labels 3, 4, 3, 4 (in this order).

Actually, each of our three main methods (see Sections 13.1–13.3) will typically fail in less drastic ways. In the Euler characteristic method and the (general) reciprocity-based method, solving for h can yield a non-constant rational function of m; see (13.9) and (13.25), respectively. In the symmetry-based method, the polynomial Q(G, m) can fail to be symmetric with respect to the involution (13.16). In these cases, the methods yield an answer which should be viewed with some skepticism. In what follows, references to the failure of one of the three methods mean that h is non-constant or that symmetry fails. (See Remark 14.4 for a discussion of the special versions of the reciprocity-based method, described in Sections 13.3–13.4.)

We now summarize the results of applying our methods to various Coxeter diagrams of infinite type, including the affine diagrams. The only infinite family for which the methods succeed wonderfully is the affine series of type \tilde{A} .

Proposition 14.2. For a Coxeter diagram of type \tilde{A}_{n-1} with $n \geq 3$, the fake invariants arising from all methods described in Section 13 agree, and coincide with

the corresponding invariants of type B_n . That is, h = 2n and the exponents are $1, 3, \ldots 2n - 1$.

Proof. Let the Coxeter diagram G be a "simply-laced n-cycle," i.e., an unoriented graph with the vertices $1, 2, \ldots, n$ such that the labels of the edges $\{1, 2\}, \ldots, \{n-1, n\}, \{n, 1\}$ are equal to 3, and the labels of all other edges are 2. To prove Proposition 14.2, one needs to verify the following statements for this particular diagram G and the invariants given by the type- B_n formulas: the recurrence (13.6); the formula (13.8) for the "fake Euler characteristic;" and the inclusion-exclusion formula (13.22). This is a straightforward verification, omitted here. (Note that each connected induced subgraph of G is of type A.)

Remark 14.3. The recurrences that determine the fake invariants of type \tilde{A}_{n-1} are different from the type- B_n recurrences, so the coincidence of the final answers is somewhat mysterious. On the other hand, the fact that the n-cycle is combinatorially related to the type- B_n cluster complex (the n-dimensional cyclohedron) is well known: there is a different generalization of associahedra [7, 10, 21] which produces the cyclohedron in case of an n-cycle. It would be interesting to clarify the relationship between the two constructions, and provide a conceptual explanation for Proposition 14.2.

Our computations suggest that the complete list of connected Coxeter diagrams for which all methods described in Sections 13.1–13.3 produce *positive integer* (fake) exponents is as follows:

- Coxeter diagrams of finite irreducible Coxeter groups;
- Coxeter diagrams of affine type \tilde{A} ;
- diagrams of rank 3 with the sum of edge labels equal to 8, 9, 10, or 11.

Remark 14.4. The discussion below does not refer to the methods based on (13.20)–(13.22) and on Lemma 13.10, respectively. This is because these methods produce results which are, by construction, the specializations (at m = 1 and m = 0, respectively) of the results delivered by the general reciprocity method of Section 13.3 based on the recurrences (13.23) and (13.25). So whenever the general reciprocity method produces a fake Coxeter number h that is a non-constant rational function in m, the specializations of this rational function at m = 0 and m = 1 should be viewed with suspicion. Still, it is tempting to mention some instances of Coxeter diagrams of affine type where the general reciprocity method fails whereas the M(G)-based method of Section 13.4 gives a positive integer value of h. They are:

- type \tilde{B}_5 , with h = 22 and M(G) = 26 (in type \tilde{B}_4 , h is a fraction);
- type \tilde{C}_n , with h = 3n + 4 and M(G) = 3n + 2 (but irrational exponents);
- type \tilde{E}_8 , with h = 98 and M(G) = 306.

The fake Coxeter number, if defined, is always a rational number (by design), and sometimes even an integer. On the other hand, in most cases, some of the fake exponents are irrational, as the generalized Fuss-Catalan invariant N(G, m) would not factor into linear polynomials in m (over \mathbb{Q}). Still, N(G, m) would sometimes be an integer-valued polynomial, potentially allowing for a meaningful interpretation.

For the affine types other than A, calculations give the following results.

The types \tilde{B}_2/\tilde{C}_2 and \tilde{G}_2 fall into the n=3 case mentioned above, with a=10and a=11, respectively. In type \tilde{B}_2/\tilde{C}_2 , we get h=10, with exponents 1,5,9—the same invariants as in type H_3 . In type \tilde{G}_2 , we get h=22, with exponents 1, 11, 21.

In type \tilde{B}_3 , all methods yield $h = \frac{76}{5}$, with exponents $1, \frac{33}{5}, \frac{43}{5}, \frac{71}{5}$. In type \tilde{C}_3 , all methods yield h = 13, with exponents $1, \frac{13-\sqrt{17}}{2}, \frac{13+\sqrt{17}}{2}, 12$. Cf. Example 13.12 with $b_1 = 4$, $b_2 = 3$, $b_3 = 4$.

The type D_4 is the only ambiguous case that we have found. The reciprocity method gives h = 14 with the exponents 1, 6, 6, 9, 13. This sequence of exponents is not symmetric, and accordingly, the symmetry-based method fails. However, the answer produced by blindly applying the formulas from the symmetry-based method still agrees with the result of the reciprocity method. The Euler characteristic method fails.

For all other affine types, computations up to rank 12 suggest that all methods fail. Beyond rank 3, our tests succeeded in a few isolated examples of non-affine infinite type, although they did not ordinarily produce integer values of h. Here are a few typical examples.

For a 4-cycle with one edge labeled 4 and the others labeled 3, we get h = 43/2with exponents $1, \frac{43-\sqrt{145}}{4}, \frac{43+\sqrt{145}}{4}, \frac{41}{2}$. For a 5-cycle with similar labels, all tests fail. For a 4-cycle with one edge labeled 5 and the others labeled 3, all methods yield h = -22 with exponents $1, -11 + 2\sqrt{3}, -11 - 2\sqrt{3}, -23$. We also found two more examples with positive integer fake Coxeter numbers:

$$\bullet \xrightarrow{3} \bullet \xrightarrow{4} \bullet \xrightarrow{4} \bullet \qquad h = 98$$

$$\bullet \xrightarrow{4} \bullet \xrightarrow{3} \bullet \xrightarrow{5} \bullet \qquad h = 104$$

We note that in all cases we have tried, if all methods work, then they all agree.

ACKNOWLEDGMENTS

We are grateful to Christos Athanasiadis for valuable comments on the preliminary versions of this paper, for explaining some of his work, and for making available his paper [4] (joint with E. Tzanaki) which we cite in Section 12.

Christian Krattenthaler, David Speyer and Hugh Thomas contributed several suggestions which improved the quality of the final version. We also thank Egon Schulte, Richard Stanley, John Stembridge, and Andrei Zelevinsky for helpful conversations.

The first version of this paper was circulated at the workshop "Braid groups, clusters and free probability" held at the American Institute of Mathematics on January 10–14, 2005. We thank the organizers (Jon McCammond, Alexandru Nica, and Victor Reiner) as well as the rest of participants for stimulating discussions.

This paper was completed in the spring of 2005, when S.F. was visiting the Mittag-Leffler Institute in Djursholm, Sweden.

Our computations of (fake) Coxeter invariants were done with Maple using John Stembridge's package coxeter. The Maple source code is currently available at http://www.math.lsa.umich.edu/~nreading/papers/gcccc/.

REFERENCES

- [1] C. A. Athanasiadis, Deformations of Coxeter hyperplane arrangements and their characteristic polynomials, *in:* Arrangements Tokyo 1998 (ed. M. Falk and H. Terao, Kinokuniya, Tokyo), *Adv. Stud. Pure Math.* **27** (2000), 1–26.
- [2] C. A. Athanasiadis, Generalized Catalan numbers, Weyl groups and arrangements of hyperplanes, Bull. London Math. Soc. **36** (2004), 294–302.
- [3] C. A. Athanasiadis, On a refinement of the generalized Catalan numbers for Weyl groups, Tran. Amer. Math. Soc. **357** (2005), 179-196.
- [4] C. A. Athanasiadis and E. Tzanaki, On the enumeration of positive cells in generalized cluster complexes and Catalan hyperplane arrangements, preprint, March 24, 2005.
- [5] D. Bisch and V. Jones, Algebras associated to intermediate subfactors, *Invent. Math.* 128 (1997), 89–157.
- [6] N. Bourbaki, Lie groups and Lie algebras, Chapters 4-6, Springer-Verlag, Berlin, 2002.
- [7] M. Carr and S. Devadoss, Coxeter complexes and graph-associahedra, preprint math.QA/ 0407229, July 2004.
- [8] F. Chapoton, Sur le nombre de reflexions pleines dans les groupes de Coxeter finis, preprint math.RT/0405371, May 2004.
- [9] F. Chapoton, S. Fomin and A. Zelevinsky, Polytopal realizations of generalized associahedra, Canad. Math. Bull. 45 (2002), 537–566.
- [10] M. Davis, T. Januszkiewicz, and R. Scott, Fundamental groups of blow-ups, Adv. Math. 177 (2003), 115–179.
- [11] D. Ž. Djoković, On conjugacy classes of elements of finite order in compact or complex semisimple Lie groups, *Proc. Amer. Math. Soc.* **80** (1980), 181–184.
- [12] S. Fomin and N. Reading, Root systems and generalized associahedra, lecture notes for the IAS/Park City Graduate Summer School in Geometric Combinatorics, 2004, available at http://www.math.lsa.umich.edu/~fomin/papers.html.
- [13] S. Fomin and A. Zelevinsky, Y-systems and generalized associahedra, Ann. of Math. 158 (2003), 977–1018.
- [14] S. Fomin and A. Zelevinsky, Cluster algebras II: Finite type classification, *Invent. Math.* **154** (2003), 63–121.
- [15] R. L. Graham, D. E. Knuth, and O. Patashnik, Concrete mathematics. A foundation for computer science, 2nd edition. Addison-Wesley Publishing Company, Reading, MA, 1994.
- [16] M. D. Haiman, Conjectures on the quotient ring by diagonal invariants, *J. Algebraic Combin.* **3** (1994) 17–76.
- [17] J. Humphreys, Reflection Groups and Coxeter Groups, Cambridge Univ. Press, 1990.
- [18] I. Pak, Reduced decompositions of permutations in terms of star transpositions, generalized Catalan numbers and k-ary trees, Discrete Math. **204** (1999), 329–335.
- [19] D. I. Panyushev, ad-nilpotent ideals of a Borel subalgebra: generators and duality, *J. Algebra* **274** (2004), 822–846.
- [20] A. Postnikov and R. P. Stanley, Deformations of Coxeter hyperplane arrangements, J. Combin. Theory Ser. A 91 (2000) 544–597
- [21] A. Postnikov, Permutohedra, associahedra, and beyond (extended abstract), preprint available at http://www-math.mit.edu/~apost/papers.html, November 2004.
- [22] J. H. Przytycki and A. S. Sikora, Polygon dissections and Euler, Fuss, Kirkman, and Cayley numbers, J. Combin. Theory Ser. A 92 (2000), 68–76.
- [23] V. Reiner and V. Welker, On the Charney-Davis and Neggers-Stanley conjectures, J. Combin. Theory Ser. A 109 (2005), 247–280.
- [24] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, published electronically at http://www.research.att.com/~njas/sequences/, sequence A001764.
- [25] E. Sommers, B-stable ideals in the nilradical of a Borel subalgebra, Canad. Math. Bull., to appear.
- [26] R. P. Stanley, Polygon dissections and standard Young tableaux, J. Combin. Theory Ser. A 76 (1996), 175–177.

- [27] R. P. Stanley, Enumerative combinatorics, vol. 1, Cambridge University Press, Cambridge, 1997.
- [28] R. P. Stanley, *Enumerative combinatorics*, vol. 2, Cambridge University Press, Cambridge, 1999.
- [29] E. Tzanaki, Polygon dissections and some generalizations of cluster complexes, preprint math.CO/0501100, version 2, March 2005.
- [30] E. Tzanaki, Polygon dissections and some generalizations of cluster complexes for the classical reflection groups, preprint math.CO/0501100, version 1, January 2005.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA *E-mail address*: fomin@umich.edu, nreading@umich.edu