

# RESOLUTION OF SINGULARITIES FOR A FAMILY OF THIRD-ORDER DIFFERENTIAL SYSTEMS WITH SMALL MEROMORPHIC SOLUTION SPACES

YUSUKE SASANO

## 0. ABSTRACT

Regarding the resolution of singularities for the differential equations of Painlevé type, there are important differences between the second-order Painlevé equations and those of higher order. Unlike the second-order case, in higher order cases there may exist some meromorphic solution spaces with codimension 2. In this paper, we will give an explicit global resolution of singularities for a 3-parameter family of third-order differential systems with meromorphic solution spaces of codimension 2.

## 1. INTRODUCTION

In 1979, K. Okamoto constructed the spaces of initial conditions of Painlevé equations, which can be considered as the parametrized spaces of all solutions, including the meromorphic solutions [1]. They are constructed by means of successive blowing-up procedures at singular points. For Painlevé equations, the dimension of the space of meromorphic solutions through any singular point is always codimension one. But, in the case of higher order Painlevé equations, the space of meromorphic solutions through a singular point may be of codimension greater than or equal to 2 [5]. In this paper, we will give an explicit resolution of singularities for a 3-parameter family of third-order differential systems with meromorphic solution spaces of codimension 2. For second-order Painlevé equations, we can obtain the entire space of initial conditions by adding subvarieties of codimension 1 (equivalently, of dimension 1) to the space of initial conditions of holomorphic solutions [3]. But in the case of third-order differential equations, we need to add codimension 2 subvarieties to the space in addition to codimension 1 subvarieties. In order to resolve singularities, we need to both blow up and blow down. Moreover, to obtain a smooth variety by blowing-down, we need to resolve for a pair of singularities.

## 2. STATEMENT OF MAIN RESULTS

Let us consider a 3-parameter family of third-order differential system:

$$\begin{cases} \frac{dx}{dt} = x(t - x - 2z) + \alpha_1 \\ \frac{dy}{dt} = y(-t + y + 2z) + \alpha_2 \\ \frac{dz}{dt} = z(t - 2y - z) + \alpha_3. \end{cases} \quad (1)$$

Here,  $x, y$  and  $z$  denote unknown complex variables and  $\alpha_1, \alpha_2$  and  $\alpha_3$  are constant parameters. The last two equations of this system (1) are equivalent to the fourth Painlevé equations with unknown variables  $y$  and  $z$ . For each solution  $z(t)$  of the Painlevé IV equations, the first equations of this system (1) give Riccati equations with unknown variable  $x$ .

From the viewpoint of symmetry, it is interesting to point out that our differential system (1) is invariant under the following transformations:

$$s_1 : (x, y, z, t; \alpha_1, \alpha_2, \alpha_3) \rightarrow (x, y - \frac{\alpha_3}{z}, z, t; \alpha_1, \alpha_2 + \alpha_3, -\alpha_3),$$

$$s_2 : (x, y, z, t; \alpha_1, \alpha_2, \alpha_3) \rightarrow (x - \frac{\alpha_2}{y}, y, z + \frac{\alpha_2}{y}, t; \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2).$$

In order to characterize this differential system (1), we can prove the following two propositions by direct calculation.

**Notation.**

- $W = \langle s_1, s_2 \rangle$ ,
- $\mathcal{V} := \{(dx/dt = f_1(x, y, z, t; a_i), dy/dt = f_2(x, y, z, t; b_i), dz/dt = f_3(x, y, z, t; c_i))\}$ , where  $f_i(X, Y, Z, t; e_i) = e_1X^2 + e_2Y^2 + e_3Z^2 + e_4XY + e_5XZ + e_6YZ + e_7tX + e_8tY + e_9tZ + e_{10}$  with  $e_i \in \mathbb{C}$  ( $1 \leq i \leq 10$ ),
- $\mathcal{H} \cong \mathbb{P}^2(\subset \mathbb{P}^3)$ : boundary divisor of  $\mathbb{P}^3$ .

**Proposition 2.1.** *The differential systems in  $\mathcal{V}$  with symmetry under  $W$  are written as follows:*

$$\begin{cases} \frac{dx}{dt} = a_1x^2 + (a_1 - c_3)z^2 + a_5xz + a_6yz + a_7tx + (a_7 + b_8)tz \\ \hspace{15em} + \alpha_2(b_2 - a_6) + \alpha_3a_6 + a_{10} \\ \frac{dy}{dt} = b_2y^2 + (a_5 - 2a_1)xy + (2a_1 - a_5 - 2c_3)yz + b_8ty + \alpha_2(a_5 - 2a_1) + \alpha_2c_3 \\ \frac{dz}{dt} = c_3z^2 + (2a_1 - a_5)xz - 2b_2yz - b_8tz + (\alpha_2 - \alpha_3)b_2. \end{cases} \quad (2)$$

**Proposition 2.2.** *The last two equations of this system (2) are equivalent to the fourth Painlevé systems with unknown variables  $y$  and  $z$  if and only if*

$$a_5 = 2a_1.$$

**Remark 2.1.** After we recall the notion of local index in § 3, we will see that the condition  $a_5 = 2a_1$  is equivalent to the following condition (A):

(A) The differential system (2) has local index  $(-a_1, -a_1, -a_1)$  at the accessible singular point  $P_3 = \{[z_0 : z_1 : z_2 : z_3] = [0 : 1 : 0 : 0]\} \in \mathcal{H} \subset \mathbb{P}^3$ . Here  $[z_0 : z_1 : z_2 : z_3]$  is the homogeneous coordinate of  $\mathbb{P}^3$ .

**Remark 2.2.** In 1998, Noumi and Yamada [2] proposed the system of type  $A_4^{(1)}$ , which can be considered as a 4-parameter family of fourth-order coupled Painlevé IV system in dimension 4, and which is given as follows:

$$\begin{cases} \frac{dx}{dt} = x^2 + 2xy + 2xw - tx + \beta_1 \\ \frac{dy}{dt} = -y^2 - 2xy - 2yw + ty + \beta_2 \\ \frac{dz}{dt} = z^2 + 2zw + 2xy - tz + \beta_1 + \beta_3 \\ \frac{dw}{dt} = -w^2 - 2zw + tw + \beta_4. \end{cases}$$

This system reduces to the system (1) by setting  $x = 0$  and  $\beta_1 = 0$ .

In order to consider the phase space for the system (1), let us take the compactification  $\mathbb{P}^3$  of  $\mathbb{C}^3$  with the natural embedding  $(x, y, z) = (z_1/z_0, z_2/z_0, z_3/z_0)$ . Moreover we denote the hypersurface of  $\mathbb{P}^3$  by  $\mathcal{H} = \{z_0 = 0\} \simeq \mathbb{P}^2 \subset \mathbb{P}^3$ . Fixing parameter  $\alpha_i$  and setting  $B = \text{Spec}\mathbb{C}[t] \cong \mathbb{C}$ , consider the product  $\mathbb{P}^3 \times B \cong \mathbb{P}^3 \times \mathbb{C}$  and extend the regular vector field on  $\mathbb{C}^3 \times B$  to a rational vector field  $\tilde{v}$  on  $\mathbb{P}^3 \times B$ . By direct calculation, this rational vector field  $\tilde{v}$  has seven accessible singular points on the boundary divisor  $\mathcal{H} \times \{t\} \subset \mathbb{P}^3 \times \{t\}$  for each  $t \in B$  (See Figure 1).




FIGURE 1

**Theorem 2.1.** *After a series of explicit blowing-ups and blowing-downs of  $\mathbb{P}^3 \times B$ , we obtain a smooth projective family of 3-fold  $\pi : \mathcal{X} \rightarrow B$  and a birational morphism  $\varphi : \mathcal{X} \rightarrow \mathbb{P}^3 \times B$  which make the following diagram commutative:*

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\varphi} & \mathbb{P}^3 \times B \\ \pi \downarrow & & \downarrow \\ B & = & B, \end{array}$$

and satisfy the following conditions:

- (1) *The phase space  $\mathcal{X}$  over  $B = \mathbb{C}$  for the vector field  $v$  in (1) is obtained by gluing eight copies of  $\mathbb{C}^3 \times \mathbb{C}$ :*

$$\begin{aligned} U_0 \times \mathbb{C} &= \mathbb{C}^3 \times \mathbb{C} \ni (x, y, z, t), \\ U_j \times \mathbb{C} &= \mathbb{C}^3 \times \mathbb{C} \ni (x_j, y_j, z_j, t) \quad (j = 1, 2, \dots, 7), \end{aligned}$$

via the following rational transformations:

- 1)  $x_1 = \frac{1}{x+z}$ ,  $y_1 = (yz + \alpha_2)z$ ,  $z_1 = \frac{1}{z}$ ,
- 2)  $x_2 = x + z$ ,  $y_2 = (yz + \alpha_2)z$ ,  $z_2 = \frac{1}{z}$ ,
- 3)  $x_3 = \frac{1}{x}$ ,  $y_3 = y$ ,  $z_3 = z$ ,
- 4)  $x_4 = (xz - \alpha_1)z$ ,  $y_4 = ((y + z - t)z + 1 - \alpha_2 - \alpha_3)z$ ,  $z_4 = \frac{1}{z}$ ,

- 5)  $x_5 = \frac{1}{x}$ ,  $y_5 = \frac{1}{y}$ ,  $z_5 = (yz - \alpha_3)y$ ,  
6)  $x_6 = x$ ,  $y_6 = \frac{1}{y}$ ,  $z_6 = (yz - \alpha_3)y$ ,  
7)  $x_7 = \frac{1}{z(xz - \alpha_1)}$ ,  $y_7 = z((y + z - t)z + 1 - \alpha_2 - \alpha_3)$ ,  $z_7 = \frac{1}{z}$ .
- (2) Each fiber  $(\mathcal{X} \setminus \mathcal{D})_t$  of the morphism  $\pi : \mathcal{X} \setminus \mathcal{D} \rightarrow B$  has a stratification

$$(\mathcal{X} \setminus \mathcal{D})_t = \mathbb{C}^3 \sqcup \mathbb{C}^2 \sqcup \mathbb{C}^2 \sqcup \mathbb{C}^2 \sqcup \mathbb{C}^2 \sqcup \mathbb{C}^1 \sqcup \mathbb{C}^1 \sqcup \mathbb{C}^1,$$

where  $\mathbb{C}^3$  is the original affine open subset of  $\mathbb{P}^3 \setminus \mathcal{H}$ . The three-dimensional space is a parameter space of the holomorphic solutions, and each two-dimensional or one-dimensional space is a parameter space of a two-parameter or a one-parameter family of meromorphic solutions, respectively.




FIGURE 2

**Remark 2.3.** The surface defined by  $x = 0$  is the minimal compactification of the space of initial conditions of Painlevé IV equation (see Figure 2). Actually, the ordinary differential system (1) can be reduced to Painlevé IV equation if one sets  $x \equiv 0$  and  $\alpha_1 = 0$ . However, the total differential system (1) is not a product of the fourth Painlevé equation and a Riccati equation.

**Remark 2.4.** It is still an open question whether we can resolve the accessible singularities for any differential system in (2) with the condition  $a_5 = 2a_1$ .

### 3. ACCESSIBLE SINGULARITY

Let us review the notion of accessible singularity [4]. Let  $B$  be a connected open domain in  $\mathbb{C}$  and  $\pi : \mathcal{W} \rightarrow B$  a smooth proper holomorphic map. We assume that  $\mathcal{H} \subset \mathcal{W}$  is a normal crossing divisor which is flat over  $B$ . Let us consider a rational vector field  $\tilde{v}$  on  $\mathcal{W}$  satisfying the condition

$$\tilde{v} \in H^0(\mathcal{W}, \Theta_{\mathcal{W}}(-\log \mathcal{H})(\mathcal{H})).$$

Fixing  $t_0 \in B$  and  $P \in \mathcal{W}_{t_0}$ , we can take a local coordinate system  $(x_1, x_2, \dots, x_n)$  of  $\mathcal{W}_{t_0}$  centered at  $P$  such that  $\mathcal{H}_{smooth}$  can be defined by the local equation  $x_1 = 0$ . Since  $\tilde{v} \in H^0(\mathcal{W}, \Theta_{\mathcal{W}}(-\log \mathcal{H})(\mathcal{H}))$ , we can write down the vector field  $\tilde{v}$  near  $P = (0, 0, \dots, 0, t_0)$  as follows:

$$\tilde{v} = \frac{\partial}{\partial t} + a_1 \frac{\partial}{\partial x_1} + \frac{a_2}{x_1} \frac{\partial}{\partial x_2} + \dots + \frac{a_n}{x_1} \frac{\partial}{\partial x_n}$$

This vector field defines the following system of differential equations

$$\begin{cases} \frac{dx_1}{dt} = a_1(x_1, x_2, \dots, x_n, t) \\ \frac{dx_2}{dt} = \frac{a_2(x_1, x_2, \dots, x_n, t)}{x_1} \\ \cdot \\ \cdot \\ \cdot \\ \frac{dx_n}{dt} = \frac{a_n(x_1, x_2, \dots, x_n, t)}{x_1} \end{cases} \quad (3)$$

Here  $a_i(x_1, \dots, x_n, t)$  ( $i = 1, 2, \dots, n$ ) are holomorphic functions near  $P = (0, 0, \dots, 0, t_0)$ .

**Definition 3.1.** *With the above notation, assume that the rational vector field  $\tilde{v}$  on  $\mathcal{W}$  satisfies the condition*

$$\tilde{v} \in H^0(\mathcal{W}, \Theta_{\mathcal{W}}(-\log \mathcal{H})(\mathcal{H})).$$

*We say that  $\tilde{v}$  has an accessible singular point at  $P = (0, 0, \dots, 0, t_0)$  if*

$$x_1 = 0 \text{ and } a_i(0, 0, \dots, 0, t_0) = 0 \text{ for every } i, 2 \leq i \leq n.$$

If  $P \in \mathcal{H}_{smooth}$  is not an accessible singular point, all solutions of the ordinary differential equation passing through  $P$  are vertical solutions, that is, the solutions are contained in the fiber  $\mathcal{W}_{t_0}$  over  $t = t_0$ . If  $P \in \mathcal{H}_{smooth}$  is an accessible singular point, there may be a solution of (3) which passes through  $P$  and goes into the interior  $\mathcal{W} - \mathcal{H}$  of  $\mathcal{W}$ .

Let us recall the notion of local index [4]. When we construct the phase spaces of the higher order Painlevé equations, an object that we call the local index is the key for determining when we need to make a blowing-up of an accessible singularity or a blowing-down to a minimal phase space. In the case of equations of higher order with favorable properties, for example the systems of type  $A_4^{(1)}$  [2], the local index at the accessible singular point corresponds to the set of orders that appears in the free parameters of formal solutions passing through that point [5].

**Definition 3.2.** *Let  $v$  be an algebraic vector field which is given by (3) and  $(X, Y, Z)$  be a boundary coordinate system in a neighborhood of an accessible singular point  $P = (0, 0, 0, t)$ . Assume that the system is written as*

$$\begin{cases} \frac{dX}{dt} = a + f_1(X, Y, Z, t) \\ \frac{dY}{dt} = \frac{bY + f_2(X, Y, Z, t)}{X} \\ \frac{dZ}{dt} = \frac{cZ + f_3(X, Y, Z, t)}{X} \end{cases}$$

near the accessible singular point  $P$ , where  $a, b$  and  $c$  are nonzero constants. We say that the vector field  $v$  has local index  $(a, b, c)$  at  $P$  if  $f_1(X, Y, Z, t)$  is a polynomial which vanishes at  $P = (0, 0, 0, t)$  and  $f_i(X, Y, Z, t)$  ( $i = 2, 3$ ) are polynomials of order 2 in  $X, Y, Z$ . Here  $f_i \in \mathbb{C}[X, Y, Z, t]$  ( $i = 1, 2, 3$ ).

**Remark 3.1.** We are interested in the case with local index  $(1, \frac{b}{a}, \frac{c}{a}) \in \mathbb{Z}^3$ . If each component of  $(1, \frac{b}{a}, \frac{c}{a})$  has the same sign, we may resolve the accessible singularity by blowing-up finitely many times. But when different signs appear, we may need to both blow up and blow down.

In our case, there exist 7 accessible singular points of this system on the boundary divisor of  $\mathbb{P}^3$ . They are listed as follows.

Singular point	$P_1$	$P_2$	$P_3$	$P_4$
$[z_0 : z_1 : z_2 : z_3]$	$[0 : 0 : 0 : 1]$	$[0 : -1 : 0 : 1]$	$[0 : 1 : 0 : 0]$	$[0 : 0 : -1 : 1]$
Type (dim. of sol.)	$\circ$ (dim. 1)	$\bullet$ (dim. 2)	$\bullet$ (dim. 2)	$\bullet$ (dim. 2)
Type of local index	$(-1, +3, +1)$	$(+1, +3, +1)$	$(+1, +1, +1)$	$(-3, -3, -1)$

Singular point	$P_5$	$P_6$	$P_7$
$[z_0 : z_1 : z_2 : z_3]$	$[0 : -1 : 1 : 0]$	$[0 : 0 : 1 : 0]$	$[0 : -3 : -1 : 1]$
Type (dim. of sol.)	$\circ$ (dim. 1)	$\bullet$ (dim. 2)	$\star$ (dim. 1)
Type of local index	$(+1, -1, -3)$	$(-1, -1, -3)$	$(+3, -3, -1)$

Here, we remark that there are 3 types of accessible singularities. We denote the mark by  $\bullet$  an accessible singular point into which two dimensional meromorphic solutions flow, and the marks by  $\circ$  and  $\star$  accessible singular points into which one dimensional meromorphic solutions flow.

#### 4. RESOLUTION OF ACCESSIBLE SINGULARITIES

Let  $P$  be an accessible singular point in the boundary divisor  $\mathcal{H}$ . Rewriting the systems in a local coordinate system, the right hand side of each differential equation has poles along  $\mathcal{H}$ . If we resolve the accessible singular point  $P$  and the right hand side of each differential equation becomes holomorphic in new coordinate system, then we can use Cauchy's existence and uniqueness theorem of solutions.

##### Notation

$(u, v, w) = (\frac{x}{z}, \frac{y}{z}, \frac{1}{z})$ ,  $(p, q, r) = (\frac{1}{x}, \frac{y}{x}, \frac{z}{x})$ ,  $(l, m, n) = (\frac{x}{y}, \frac{1}{y}, \frac{z}{y})$ ,  $(\tilde{u}, v, w) = (u + 1, v, w)$ .

##### 4.1. Resolution of accessible singular points $P_1, P_2$ and $P_3$

Let us start by summarizing the steps which are needed to resolve the accessible singular points of  $\tilde{v}$  at  $P_1 : (u, v, w) = (0, 0, 0)$ ,  $P_2 : (\tilde{u}, v, w) = (0, 0, 0)$  and  $P_3 : (p, q, r) = (0, 0, 0)$ .

- (1) **Step 1:** We will blow up at two points  $P_2, P_3$  (see Figure 3).
- (2) **Step 2:** We will blow up along the curve  $C_1$  (see Figure 3).
- (3) **Step 3:** We will blow down the surface  $F$  (see Figure 3).
- (4) **Step 4:** We will blow up along the curve  $C_4$  (see Figure 3).
- (5) **Step 5:** We will blow up along the curve  $C_5$  (see Figure 3).

In a neighborhood of  $P_1$ , the system is written as




FIGURE 3

$$\begin{cases} \frac{du}{dt} = \frac{-u - u^2 + 2uv}{w} + \alpha_1 w - \alpha_3 u w \\ \frac{dv}{dt} = -2tv + \frac{3v + 3v^2}{w} + \alpha_2 w - \alpha_3 v w \\ \frac{dw}{dt} = 1 + 2v - tw - \alpha_3 w^2. \end{cases}$$

In a neighborhood of  $P_2$ , the system is written as

$$\begin{cases} \frac{d\tilde{u}}{dt} = \frac{\tilde{u} - \tilde{u}^2 - v(2 - 2\tilde{u})}{w} + \alpha_3 w + \alpha_1 w - \alpha_3 \tilde{u} w \\ \frac{dv}{dt} = -2tv + \frac{3v + 3v^2}{w} + \alpha_2 w - \alpha_3 v w \\ \frac{dw}{dt} = -\alpha_3 w^2 - tw + 2v + 1. \end{cases}$$

In a neighborhood of  $P_3$ , the system is written as

$$\begin{cases} \frac{dp}{dt} = -\alpha_1 p^2 - tp + 2r + 1 \\ \frac{dq}{dt} = \frac{q + q(-2tp + q + 4r)}{p} + \alpha_2 p - \alpha_1 p q \\ \frac{dr}{dt} = \frac{r + r(r - 2q)}{p} - \alpha_1 p r + \alpha_3 p. \end{cases}$$

**4.1.1.** We blow up at two points  $P_2, P_3$ .

We blow up at the point  $P_2$

$$U_1 = \frac{\tilde{u}}{w}, \quad V_1 = \frac{v}{w}, \quad W_1 = w.$$

We blow up at the point  $P_3$

$$x_3 = p, \quad y_3 = \frac{q}{p}, \quad z_3 = \frac{r}{p}.$$

We have resolved the accessible singular point  $P_3$ .

**4.1.2.** We blow up along the curve  $C_1 = \{(u, v, w) | v = w = 0\} \subset \mathbb{P}^1$

$$u_1 = u, \quad v_1 = \frac{v}{w}, \quad w_1 = w.$$

In a neighborhood of  $\{(u_1, v_1, w_1) = (0, 0, 0)\}$ , the system is written as

$$\begin{cases} \frac{du_1}{dt} = 2u_1v_1 + \frac{-u_1 - (u_1)^2}{w_1} + \alpha_1w_1 - \alpha_3u_1w_1 \\ \frac{dv_1}{dt} = -tv_1 + (v_1)^2 + \frac{2v_1}{w_1} + \alpha_2 \\ \frac{dw_1}{dt} = 1 - tw_1 + 2v_1w_1 - \alpha_3(w_1)^2. \end{cases}$$

**4.1.3.** We blow down the surface  $F = \{(u_1, v_1, w_1) | w_1 = 0\} \cong \mathbb{P}^1 \times \mathbb{P}^1$

$$u_2 = \frac{w_1}{u_1 + 1}, \quad v_2 = v_1, \quad w_2 = w_1.$$

In a neighborhood of  $\{(u_2, v_2, w_2) = (0, 0, 0)\}$ , the system is written as

$$\begin{cases} \frac{du_2}{dt} = 1 - tu_2 + \frac{2(u_2)^2v_2}{w_2} - \alpha_1(u_2)^2 - \alpha_3(u_2)^2 \\ \frac{dv_2}{dt} = -tv_2 + (v_2)^2 + \frac{2v_2}{w_2} + \alpha_2 \\ \frac{dw_2}{dt} = 1 - tw_2 + 2v_2w_2 - \alpha_3(w_2)^2. \end{cases} \quad (4)$$

The resolution process from Step 2 to Step 3 is well-known as flop. In order to resolve the accessible singular point  $P_1$  and obtain a holomorphic coordinate system, we need to blow down the surface  $F \cong \mathbb{P}^1 \times \mathbb{P}^1$  along the  $\mathbb{P}^1$ -fiber. After we blow down the surface  $F$ , the system (4) has the local index  $(0, +2, +1)$  at the point  $\{(u_2, v_2, w_2) = (0, 0, 0)\}$ .

**4.1.4.** We blow up along the curve  $C_4 = \{(u_2, v_2, w_2) | v_2 = w_2 = 0\} \subset \mathbb{P}^1$

$$u_3 = u_2, \quad v_3 = \frac{v_2}{w_2}, \quad w_3 = w_2.$$

**4.1.5.** We blow up along the curve  $C_5 = \{(u_3, v_3, w_3) | v_3 + \alpha_2 = w_3 = 0\} \subset \mathbb{P}^1$

$$x_1 = u_3, \quad y_1 = \frac{v_3 + \alpha_2}{w_3}, \quad z_1 = w_3.$$

We have resolved the accessible singular point  $P_1$ .

**4.2. Resolution of accessible singular points  $P_5 : (l, m, n) = (0, -1, 0)$  and  $P_6 : (l, m, n) = (0, 0, 0)$**

We can resolve accessible singular points  $P_5, P_6$  by the same way of 4.1. By the birational transformation of 4.1 and 4.2, we have resolved five accessible singular points  $P_1, P_2, P_3, P_5$  and  $P_6$ . There are only two accessible singular points  $P_4, P_7$  in  $\mathbb{F}_1$ -surface. Here  $\mathbb{F}_1$ -surface is obtained by blowing up one time at a point in  $\mathbb{P}^2$ . These two accessible singular points are on the same fiber in  $\mathbb{F}_1$ -surface (See figure 4).




FIGURE 4

### 4.3. Resolution of accessible singular points $P_4$ and $P_7$

Let us start by summarizing the steps which are needed to resolve the accessible singular points of  $\tilde{v}$  at  $P_4$  and  $P_7$ :

$$P_4 = \{(u, v, w) = (0, -1, 0)\}, \quad P_7 = \{(u, v, w) = (-3, -1, 0)\}.$$

- (1) **Step 6:** We will blow up along the curve  $C_6$  and blow down the surface  $S_1$  (see Figure 4).
- (2) **Step 7:** We will blow up along the curve  $C_7$  and blow down the surface  $S_2$  (see Figure 4).
- (3) **Step 8:** We will blow up along the curve  $C_8$  and blow down the surface  $S_3$  (see Figure 4).
- (4) **Step 9:** We will blow up along the curve  $C_9$  (see Figure 4).
- (5) **Step 10:** We will blow up along the curve  $C_{10}$  (see Figure 4).
- (6) **Step 11:** We will blow up along the curve  $C_{11}$  (see Figure 4).

In a neighborhood of  $P_7$ , the system is written as

$$\begin{cases} \frac{dl_1}{dt} = \frac{3l_1 - (l_1)^2 - 6m_1 + 2l_1m_1}{n_1} + n_1\alpha_1 + 3n_1\alpha_3 - l_1n_1\alpha_3 \\ \frac{dm_1}{dt} = \frac{-3m_1 + 3(m_1)^2}{n_1} + 2t - 2tm_1 + n_1\alpha_2 + n_1\alpha_3 - m_1n_1\alpha_3 \\ \frac{dn_1}{dt} = -1 + 2m_1 - tn_1 - \alpha_3(n_1)^2. \end{cases}$$

In a neighborhood of  $P_4$ , the system is written as

$$\begin{cases} \frac{dL_1}{dt} = \frac{-3L_1 - (L_1)^2 + 2L_1M_1}{N_1} + \alpha_1N_1 - \alpha_3L_1N_1 \\ \frac{dM_1}{dt} = 2t - 2tM_1 + \frac{-3M_1 + 3(M_1)^2}{N_1} + \alpha_2N_1 + \alpha_3N_1 - \alpha_3M_1N_1 \\ \frac{dN_1}{dt} = -1 + 2M_1 - tN_1 - \alpha_3(N_1)^2. \end{cases}$$

**4.3.1.** We blow up along the curve  $C_6 = \{(L_1, M_1, N_1) | L_1 = N_1 = 0\} \subset \mathbb{P}^1$  and blow down the surface  $S_1 = \{(l_1, m_1, n_1) | n_1 = 0\}$

$$\begin{aligned} L_2 &= \frac{L_1}{N_1}, & M_2 &= M_1, & N_2 &= N_1, \\ l_2 &= \frac{n_1}{l_1 - 3}, & m_2 &= m_1, & n_2 &= n_1. \end{aligned}$$

In a neighborhood of  $\{(l_2, m_2, n_2) = (0, 0, 0)\}$ , the system is written as

$$\begin{cases} \frac{dl_2}{dt} = 1 - tl_2 + \frac{2l_2}{n_2} - \alpha_1(l_2)^2 \\ \frac{dm_2}{dt} = 2t - 2tm_2 - \frac{3m_2 - 3(m_2)^2}{n_2} + \alpha_2n_2 + \alpha_3n_2 - \alpha_3m_2n_2 \\ \frac{dn_2}{dt} = -1 + 2m_2 - tn_2 - \alpha_3(n_2)^2. \end{cases}$$

**4.3.2.** We blow up along the curve  $C_7 = \{(L_2, M_2, N_2) | L_2 = N_2 = 0\} \subset \mathbb{P}^1$  and blow down the surface  $S_2 = \{(l_2, m_2, n_2) | n_2 = 0\}$

$$\begin{aligned} L_3 &= \frac{L_2}{N_2}, & M_3 &= M_2, & N_3 &= N_2, \\ l_3 &= l_2n_2, & m_3 &= m_2, & n_3 &= n_2. \end{aligned}$$

In a neighborhood of  $\{(l_3, m_3, n_3) = (0, 0, 0)\}$ , the system is written as

$$\begin{cases} \frac{dl_3}{dt} = -2tl_3 + \frac{l_3 + 2l_3m_3 - \alpha_2(l_3)^2}{n_3} + n_3 - \alpha_3l_3n_3 \\ \frac{dm_3}{dt} = 2t - 2tm_3 - \frac{3m_3 - 3(m_3)^2}{n_3} + \alpha_2n_3 + \alpha_3n_3 - \alpha_3m_3n_3 \\ \frac{dn_3}{dt} = -1 + 2m_3 - tn_3 - \alpha_3(n_3)^2. \end{cases}$$

**4.3.3.** We blow up along the curve  $C_8 = \{(L_3, M_3, N_3) | L_3 = N_3 = 0\} \subset \mathbb{P}^1$  and blow down the surface  $S_3 = \{(l_3, m_3, n_3) | n_3 = 0\}$

$$\begin{aligned} L_4 &= \frac{L_3 - \alpha_1}{N_3}, & M_4 &= M_3, & N_4 &= N_3, \\ l_4 &= \frac{l_3n_3}{1 - \alpha_1l_3}, & m_4 &= m_3, & n_4 &= n_3. \end{aligned}$$

**4.3.4.** We blow up along the curve  $C_9 = \{(L_4, M_4, N_4) | M_4 = N_4 = 0\} \cup \{(l_4, m_4, n_4) | m_4 = n_4 = 0\} \cong \mathbb{P}^1$

$$L_5 = L_4, \quad M_5 = \frac{M_4}{N_4}, \quad N_5 = N_4,$$

$$l_5 = l_4, \quad m_5 = \frac{m_4}{n_4}, \quad n_5 = n_4.$$

**4.3.5.** We blow up along the curve  $C_{10} = \{(L_5, M_5, N_5) | M_5 - t = N_5 = 0\} \cup \{(l_5, m_5, n_5) | m_5 - t = n_5 = 0\} \cong \mathbb{P}^1$

$$L_6 = L_5, \quad M_6 = \frac{M_5 - t}{N_5}, \quad N_6 = N_5,$$

$$l_6 = l_5, \quad m_6 = \frac{m_5 - t}{n_5}, \quad n_6 = n_5.$$

**4.3.6.** We blow up along the curve  $C_{11} = \{(L_6, M_6, N_6) | M_6 + 1 - \alpha_2 - \alpha_3 = N_6 = 0\} \cup \{(l_6, m_6, n_6) | m_6 + 1 - \alpha_2 - \alpha_3 = n_6 = 0\} \cong \mathbb{P}^1$

$$x_4 = L_6, \quad y_4 = \frac{M_6 + 1 - \alpha_2 - \alpha_3}{N_6}, \quad z_4 = N_6,$$

$$x_7 = l_6, \quad y_7 = \frac{m_6 + 1 - \alpha_2 - \alpha_3}{n_6}, \quad z_7 = n_6.$$

We have resolved the accessible singular points  $P_4$  and  $P_7$ .

*Acknowledgement.* The author thanks Professors M.-H. Saito, K. Takano and Y. Yamada for giving helpful suggestions and encouragement, and Professor W. Rossman for checking English.

## REFERENCES

- [1] K. Okamoto, *Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé, Espaces des conditions initiales*, Japan. J. Math., **5**, 1979, 1–79.
- [2] M. Noumi and Y. Yamada, *Higher order Painlevé equations of type  $A_l^{(1)}$* , Funkcial. Ekvac., **41**, 1998, 483–503.
- [3] T. Shioda and K. Takano, *On some Hamiltonian structures of Painlevé systems I*, Funkcial. Ekvac., **40**, 1997, 271–291.
- [4] Y. Sasano, *Coupled Painlevé II systems in dimension 4 and the systems of type  $A_4^{(1)}$* , preprint.
- [5] N. Tahara, *An augmentation of the phase space of the system of type  $A_4^{(1)}$* , Kyushu J. Math. **58**, 2004, 393–425.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KOBE UNIVERSITY, KOBE, ROKKO, 657-8501, JAPAN

*E-mail address:* sasano@math.kobe-u.ac.jp