

ON THE SLICING GENUS OF LEGENDRIAN KNOTS

HAO WU

ABSTRACT. We apply Heegaard-Floer homology theory to establish generalized slicing Bennequin inequalities closely related to a recent result of T. Mrowka and Y. Rollin proved using Seiberg-Witten monopoles.

1. INTRODUCTION

Let ξ be an oriented 2-plane distribution on an oriented 3-manifold M . (Unless otherwise specified, all 3-manifolds in this paper are closed, connected and oriented.) ξ is said to be a contact structure on M if there is a 1-form α on M so that $\xi = \ker \alpha$, $d\alpha|_{\xi} > 0$ and $\alpha \wedge d\alpha > 0$. Such a 1-form is called a contact form for ξ . And (M, ξ) is called a contact 3-manifold. A knot K in a contact 3-manifold (M, ξ) is called a Legendrian knot if it's tangent to ξ . (Unless otherwise specified, all the knots in this paper are oriented.) (M, ξ) is said to be overtwisted if there is an embedded disk D in M s.t. ∂D is Legendrian, but D is transverse to ξ along ∂D . If (M, ξ) is not overtwisted, then it's called tight. For example, the standard contact structure ξ_{st} on S^3 given by the complex tangencies of the unit 3-sphere in \mathbb{C}^2 is tight. Overtwisted contact structures are kind of "soft", and are completely classified up to isotopy by the homotopy type of the underlying 2-plane distribution. (See [2].) Tight contact structures display more rigidity, and possess more interesting properties.

There are two "classical" invariants, the Thurston-Bennequin number $tb(K)$ and the rotation number $r(K)$, for a Legendrian knot K in (S^3, ξ_{st}) . These are generalized to null-homologous Legendrian knots in any contact 3-manifold (c.f. [4]). Let K be a null-homologous Legendrian knot in a contact 3-manifold (M, ξ) , and $\Sigma \subset M$ a Seifert surface of K . Let K' be a knot obtained by pushing K slightly in the direction of a vector field that is transverse to ξ along K . Then the Thurston-Bennequin number $tb(K, \Sigma)$ is defined to be the intersection number $\#(K' \cap \Sigma)$. Let u be the positive unit tangent vector field of K . Then the rotation number $r(K, \Sigma)$ is defined to be the pairing $\langle c_1(\xi, u), [\Sigma] \rangle$, where $[\Sigma] \in H_2(M, K)$ is the relative homology class represented by Σ . If we reverse the orientation of K , then $tb(K, \Sigma)$ is unchanged, and $r(K, \Sigma)$ changes sign. Note that $tb(K, \Sigma)$ and $r(K, \Sigma)$ depend on Σ only through the relative homology class $[\Sigma]$. If M is a homology sphere, then $H_2(M, K) = \mathbb{Z}$, and tb, r are independent of Σ . In this case, we suppress Σ from the notation.

In [1], D. Bennequin proved the following Bennequin inequality:

For any Legendrian knot K in (S^3, ξ_{st}) ,

$$(1) \quad tb(K) + |r(K)| \leq 2g(K) - 1,$$

where $g(K)$ is the genus of K .

In [4], Y. Eliashberg generalized (1) to any tight contact 3-manifold, and get:

For any null-homologous Legendrian knot K in a tight contact 3-manifold, and any Seifert surface Σ of K ,

$$(2) \quad tb(K, \Sigma) + |r(K, \Sigma)| \leq -\chi(\Sigma).$$

In [16], L. Rudolph strengthened (1) to the slicing Bennequin inequality:

For any Legendrian knot K in (S^3, ξ_{st}) ,

$$(3) \quad tb(K) + |r(K)| \leq 2g_s(K) - 1,$$

where $g_s(K)$ is the slicing genus of K .

Let W be an oriented 4-manifold with connected boundary $\partial W = M$, and ξ a contact structure on M . Assume that K is a Legendrian knot in (M, ξ) , and Σ is an embedded surface in W bounded by K . One can extend the definition of tb and r to this situation (c.f. [9]). Let v be a vector field on M transverse to ξ . Extend v to a vector field on W , and denote by $\{\varphi_t\}$ the flow of this extended vector field. For a small $\varepsilon > 0$, let $K' = \varphi_\varepsilon(K)$ and $\Sigma' = \varphi_\varepsilon(\Sigma)$. Then the intersection number $\#(\Sigma \cap \Sigma')$ is well defined, and we define the Thurston-Bennequin number to be $tb(K, \Sigma) = \#(\Sigma \cap \Sigma')$. Note that $tb(K, \Sigma)$ depends on Σ only through the relative homology class $[\Sigma] \in H_2(W, K)$, and, when $\Sigma \subset M$, this definition coincide with the previous definition of tb . Assume $\mathfrak{s} \in Spin^C(W)$, and there is an isomorphism $h : \mathfrak{s}|_M \rightarrow \mathfrak{t}_\xi$, where \mathfrak{t}_ξ is the canonical $Spin^C$ -structure on M associated to ξ . Choose a complex structure on ξ . Then $\det(\mathfrak{t}_\xi)$ is canonically isomorphic to ξ , and h induces an isomorphism $\det(h) : \det(\mathfrak{s})|_M \rightarrow \xi$. Let u be the positive unit tangent vector field of K . We define the rotation number to be $r(K, \Sigma, \mathfrak{s}, h) = \langle c_1(\det(\mathfrak{s}), \det(h)^{-1}(u)), [\Sigma] \rangle$. Note that $r(K, \Sigma, \mathfrak{s}, h)$ depends on Σ only through the relative homology class $[\Sigma] \in H_2(W, K)$, depends on the pair (\mathfrak{s}, h) only through the isomorphism type of it in $Spin^C(W, \xi)$, and, again, when $\Sigma \subset M$, r is independent of (\mathfrak{s}, h) and coincide with the previous definition of the rotation number. As before, under the reversal of the orientation of K , tb is unchanged, and r changes sign. In the special case that there is a symplectic form ω on W such that (W, ω) is a weak symplectic filling of (M, ξ) , i.e., $\omega|_\xi > 0$, this symplectic form ω determines a canonical $Spin^C$ -structure \mathfrak{s}_ω on W and a canonical isomorphism $h_\omega : \mathfrak{s}_\omega|_M \rightarrow \mathfrak{t}_\xi$. We write $r(K, \Sigma, \omega) = r(K, \Sigma, \mathfrak{s}_\omega, h_\omega)$.

In [9], T. Mrowka and Y. Rollin prove the following generalized slicing Bennequin inequality using Seiberg-Witten monopole invariants.

Theorem 1.1 ([9], Theorem A). *Let W be an oriented 4-manifold with connected boundary $\partial W = M$, and ξ a contact structure on M . Let K be a Legendrian knot in (M, ξ) , and Σ an embedded surface in W bounded by K . Assume there is an element $(\mathfrak{s}, h) \in Spin^C(W, \xi)$, such that $SW(\mathfrak{s}, h) \neq 0$. Then*

$$(4) \quad tb(K, \Sigma) + |r(K, \Sigma, \mathfrak{s}, h)| \leq -\chi(\Sigma).$$

Specially, when (W, ω) is a weak symplectic filling of (M, ξ) (c.f. [8], Theorem 1.1), we have

$$(5) \quad tb(K, \Sigma) + |r(K, \Sigma, \omega)| \leq -\chi(\Sigma).$$

There are two approaches in the study of 3-dimensional gauge theory: the Seiberg-Witten-Floer approach by counting solutions to the Seiberg-Witten equation; and the Heegaard-Floer approach by counting holomorphic curves. Though the techniques used in these two approaches are quite different, it is conjectured that these give equivalent theories as their 4-dimensional counterparts do. In this paper, we use Heegaard-Floer homology to prove the following generalizations of the slicing Bennequin inequality, which further demonstrates the similarity between the two theories.

Theorem 1.2. *Let W be an oriented 4-manifold with connected boundary $\partial W = M$, ξ a contact structure on M , and K a Legendrian knot in (M, ξ) .*

(a) *If there is a $\text{Spin}^\mathbb{C}$ -structure \mathfrak{s} on W with $F_{W \setminus B, \mathfrak{s}|_{W \setminus B}}^+(c^+(\xi)) \neq 0$, where B is an embedded 4-ball in the interior of W , then there is an isomorphism $h : \mathfrak{s}|_M \rightarrow \mathfrak{t}_\xi$ such that, for any embedded surface Σ in W bounded by K ,*

$$(6) \quad tb(K, \Sigma) + |r(K, \Sigma, \mathfrak{s}, h)| \leq -\chi(\Sigma).$$

(b) *If (W, ω) is a weak symplectic filling of (M, ξ) , then, for any embedded surface Σ in W bounded by K ,*

$$(7) \quad tb(K, \Sigma) + |r(K, \Sigma, \omega)| \leq -\chi(\Sigma).$$

2. HEEGAARD-FLOER HOMOLOGY

In this section, we review aspects of the Heegaard-Floer theory necessary for the proof of Theorem 1.2.

2.1. Heegaard-Floer homology. In [14], P. Ozsváth and Z. Szabó defined the Heegaard-Floer homology groups of 3-manifolds. Given a connected oriented closed 3-manifold M and a $\text{Spin}^\mathbb{C}$ -structure \mathfrak{t} on M , there are four Heegaard-Floer homology groups associated to M : $HF^\infty(M, \mathfrak{t})$, $HF^-(M, \mathfrak{t})$, $HF^+(M, \mathfrak{t})$ and $\widehat{HF}(M, \mathfrak{t})$. The first three are $\mathbb{Z}[U]$ -modules, and the last one is a \mathbb{Z} -module. In this paper, we will mostly use $HF^+(M, \mathfrak{t})$. Moreover, given a $\mathbb{Z}[H^1(M)]$ -module \mathfrak{M} , there is the notion of \mathfrak{M} -twisted Heegaard-Floer homology $\underline{HF}^+(M, \mathfrak{t}; \mathfrak{M})$, which is a $\mathbb{Z}[U] \otimes \mathbb{Z}[H^1(M)]$ -module (c.f. [15]).

If \mathfrak{M}_1 and \mathfrak{M}_2 are two $\mathbb{Z}[H^1(M)]$ -modules, and $\theta : \mathfrak{M}_1 \rightarrow \mathfrak{M}_2$ is a homomorphism, then θ naturally induces a homomorphism

$$\Theta : \underline{HF}^+(M, \mathfrak{t}; \mathfrak{M}_1) \rightarrow \underline{HF}^+(M, \mathfrak{t}; \mathfrak{M}_2).$$

If we consider \mathbb{Z} as a $\mathbb{Z}[H^1(M)]$ -module, then $\underline{HF}^+(M, \mathfrak{t}; \mathbb{Z})$ is the (untwisted) Heegaard-Floer homology $HF^+(M, \mathfrak{t})$ defined with the appropriate coherent orientation system, and the $2^{b_1(M)}$ choices of $\mathbb{Z}[H^1(M)]$ -module structures on \mathbb{Z} correspond to the $2^{b_1(M)}$ coherent orientation systems on the moduli spaces (c.f. [14, 15]).

In [13], P. Ozsváth and Z. Szabó introduced the Heegaard-Floer homology twisted by a 2-form. More precisely, consider the polynomial ring

$$\mathbb{Z}[\mathbb{R}] = \left\{ \sum_{i=1}^k c_i T^{s_i} \mid k \in \mathbb{Z}_{\geq 0}, c_i \in \mathbb{Z}, s_i \in \mathbb{R} \right\}.$$

Let $[\omega] \in H^2(M; \mathbb{R})$. The action $e^{[\nu]} \cdot T^s = T^{s + \int_M \nu \wedge \omega}$, where $[\nu] \in H^1(M)$, gives $\mathbb{Z}[\mathbb{R}]$ a $\mathbb{Z}[H^1(M)]$ -module structure. Denote the module by $\mathbb{Z}[\mathbb{R}]_{[\omega]}$. Then the Heegaard-Floer homology of M twisted by $[\omega]$ is defined to be

$$\underline{HF}^+(M, \mathfrak{t}; [\omega]) = \underline{HF}^+(M, \mathfrak{t}; \mathbb{Z}[\mathbb{R}]_{[\omega]}).$$

In [15], P. Ozsváth and Z. Szabó deduced the following adjunction inequality:

Theorem 2.1 ([15], Theorem 7.1). *Let Σ be a close oriented surface embedded in a 3-manifold M with $g(\Sigma) \leq 1$, \mathfrak{t} a $Spin^{\mathbb{C}}$ -structure on M , and \mathfrak{M} a $\mathbb{Z}[H^1(M)]$ -module. If $\underline{HF}^+(M, \mathfrak{t}; \mathfrak{M}) \neq 0$, then*

$$(8) \quad |\langle c_1(\mathfrak{t}), [\Sigma] \rangle| \leq -\chi(\Sigma).$$

Note that, although the adjunction inequality is only prove for untwisted Heegaard-Floer homology in [15], the proof there readily adapts to the twisted case.

2.2. Homomorphisms induced by cobodisms. Let W be a cobodism from a 3-manifold M_1 to another 3-manifold M_2 , and \mathfrak{s} a $Spin^{\mathbb{C}}$ -structure on W . Then W and \mathfrak{s} induce a homomorphism

$$F_{W, \mathfrak{s}}^+ : HF^+(M_1, \mathfrak{s}|_{M_1}) \rightarrow HF^+(M_2, \mathfrak{s}|_{M_2}).$$

Let \mathfrak{M} be a $\mathbb{Z}[H^1(M_1)]$ -module, and $\delta : H^1(\partial W) \rightarrow H^2(W, \partial W)$ the connecting map in the long exact sequence of the pair $(W, \partial W)$. Define

$$\mathfrak{M}(W) = \mathfrak{M} \otimes_{\mathbb{Z}[H^1(M_1)]} \mathbb{Z}[\delta H^1(\partial W)],$$

where the action of $\mathbb{Z}[H^1(M_i)]$ on $\mathbb{Z}[\delta H^1(\partial W)]$ is induced by $e^{[\nu]} \mapsto e^{\delta([\nu])}$. Then W and \mathfrak{s} also induce a homomorphism

$$F_{W, \mathfrak{s}}^+ : \underline{HF}^+(M_1, \mathfrak{s}|_{M_1}; \mathfrak{M}) \rightarrow \underline{HF}^+(M_2, \mathfrak{s}|_{M_2}; \mathfrak{M}(W)).$$

The definition of this homomorphism depends on some auxiliary choices. So it's only define up to right action by units of $\mathbb{Z}[H^1(M_1)]$ and left action by units of $\mathbb{Z}[H^1(M_2)]$. Alternatively, we consider it as an equivalence class of homomorphisms from $\underline{HF}^+(M_1, \mathfrak{s}|_{M_1}; \mathfrak{M})$ to $\underline{HF}^+(M_2, \mathfrak{s}|_{M_2}; \mathfrak{M}(W))$, and denote this equivalence class by $[\underline{F}_{W, \mathfrak{s}}^+]$.

Specially, for an $[\omega] \in H^2(W; \mathbb{R})$, let $\mathfrak{M} = \mathbb{Z}[\mathbb{R}]_{[\omega|_{M_1}]}$. There is a natural homomorphism $\theta : \mathfrak{M}(W) \rightarrow \mathbb{Z}[\mathbb{R}]_{[\omega|_{M_2}]}$ induced by $T^s \otimes e^{\delta([\nu])} \mapsto T^{s + \int_{M_2} \nu \wedge \omega}$, where $[\nu] \in H^1(M_2)$. This map induces a homomorphism Θ between the Heegaard-Floer homologies of M_2 twisted by these two $\mathbb{Z}[H^1(M_2)]$ -modules. Composing it with $\underline{F}_{W, \mathfrak{s}}^+$, we get a homomorphism

$$\underline{F}_{W, \mathfrak{s}; [\omega]}^+ = \Theta \circ \underline{F}_{W, \mathfrak{s}}^+ : \underline{HF}^+(M_1, \mathfrak{s}|_{M_1}; [\omega|_{M_1}]) \rightarrow \underline{HF}^+(M_2, \mathfrak{s}|_{M_2}; [\omega|_{M_2}]).$$

Again, $\underline{F}_{W, \mathfrak{s}; [\omega]}^+$ is only defined up to multiplication by $\pm T^s$, and is consider as a equivalent class $[\underline{F}_{W, \mathfrak{s}; [\omega]}^+]$ (c.f. [13]).

The homomorphisms defined here satisfy the following composition laws:

Theorem 2.2 ([10], Theorems 3.4, 3.9). *Let W_1 be a cobodism from a 3-manifold M_1 to another 3-manifold M_2 , and W_2 a cobodism from M_2 to a third 3-manifold M_3 . Then $W = W_1 \cup_{M_2} W_2$ is a cobodism from M_1 to M_3 . We have:*

(a) *For any $\text{Spin}^{\mathbb{C}}$ -structures $\mathfrak{s}_i \in \text{Spin}^{\mathbb{C}}(W_i)$, $i = 1, 2$,*

$$(9) \quad F_{W_2, \mathfrak{s}_2}^+ \circ F_{W_1, \mathfrak{s}_1}^+ = \sum_{\{\mathfrak{s} \in \text{Spin}^{\mathbb{C}}(W) \mid \mathfrak{s}|_{W_i} \cong \mathfrak{s}_i\}} \pm F_{W, \mathfrak{s}}^+.$$

(b) *Let $\mathfrak{s} \in \text{Spin}^{\mathbb{C}}(W)$, and $\mathfrak{s}_i = \mathfrak{s}|_{W_i}$. For any $\mathbb{Z}[H^1(M_1)]$ -module \mathfrak{M} , there are representatives $\underline{F}_{W_1, \mathfrak{s}_1}^+ \in [\underline{F}_{W_1, \mathfrak{s}_1}^+]$ and $\underline{F}_{W_2, \mathfrak{s}_2}^+ \in [\underline{F}_{W_2, \mathfrak{s}_2}^+]$ such that*

$$(10) \quad [\underline{F}_{W, \mathfrak{s}}^+] = [\Pi \circ \underline{F}_{W_1, \mathfrak{s}_1}^+ \circ \underline{F}_{W_2, \mathfrak{s}_2}^+],$$

where Π is induced by the natural homomorphism from $\mathfrak{M}(W_1)(W_2)$ to $\mathfrak{M}(W)$.

Combine Theorem 2.2 and the blow-up formula ([10], Theorem 3.7), we have the following theorem.

Theorem 2.3 ([10], Theorems 3.4, 3.7). *Let W be a cobodism from a 3-manifold M_1 to another 3-manifold M_2 , and \mathfrak{s} a $\text{Spin}^{\mathbb{C}}$ -structure on W . Blow up an interior point of W . We get a new cobodism \widehat{W} from M_1 to M_2 . Let $\widehat{\mathfrak{s}}$ be the lift of \mathfrak{s} to \widehat{W} with $\langle c_1(\widehat{\mathfrak{s}}), [E] \rangle = -1$, where E is the exceptional sphere. Then $F_{W, \mathfrak{s}}^+ = F_{\widehat{W}, \widehat{\mathfrak{s}}}^+$.*

2.3. The contact invariant. In [12], P. Ozsváth and Z. Szabó defined the Ozsváth-Szabó invariants for contact 3-manifolds. For each contact 3-manifold (M, ξ) , it is an element $c(\xi)$ of the quotient $\widehat{HF}(-M, \mathfrak{t}_{\xi})/\{\pm 1\}$, where \mathfrak{t}_{ξ} is the $\text{Spin}^{\mathbb{C}}$ -structure associated to ξ . Let $\iota : \widehat{HF}(-M) \rightarrow HF^+(-M)$ be the natural map (c.f. [14]). We set $c^+(\xi) = \iota(c(\xi)) \in HF^+(-M, \mathfrak{t}_{\xi})/\{\pm 1\}$. This version of the Ozsváth-Szabó contact invariants is easier to use for our purpose. Given a $\mathbb{Z}[H^1(M)]$ -module \mathfrak{M} , one can similarly define the twisted Ozsváth-Szabó contact invariant $c^+(\xi; \mathfrak{M}) \in HF^+(-M, \mathfrak{t}_{\xi}; \mathfrak{M})/\mathbb{Z}[H^1(M)]^{\times}$, where $\mathbb{Z}[H^1(M)]^{\times}$ is the set of units of $\mathbb{Z}[H^1(M)]$. Specially, if $[\omega] \in H^2(M; \mathbb{R})$, then we have the $[\omega]$ -twisted invariant $c^+(\xi; [\omega]) \in HF^+(-M, \mathfrak{t}_{\xi}; [\omega])/\{\pm T^s \mid s \in \mathbb{R}\}$ (c.f. [13]). These contact invariants vanish when ξ is overtwisted. Following properties of the Ozsváth-Szabó contact invariants are needed for the proof of Theorem 1.2.

Proposition 2.4 ([6], Proposition 3.3). *Suppose that (M', ξ') is obtained from (M, ξ) by Legendrian surgery on a Legendrian link. Then we have $F_{W, \mathfrak{s}_0}^+(c^+(\xi')) = c^+(\xi)$, where W is the cobordism induced by the surgery and \mathfrak{s}_0 is the canonical $\text{Spin}^{\mathbb{C}}$ -structure associated to the symplectic structure on W . Moreover, $F_{W, \mathfrak{s}}^+(c^+(\xi')) = 0$ for any $\text{Spin}^{\mathbb{C}}$ -structure \mathfrak{s} on W with $\mathfrak{s} \not\cong \mathfrak{s}_0$.*

Theorem 2.5 ([13], Theorem 4.2). *Let (M, ξ) be a contact 3-manifold with a weak symplectic filling (W, ω) . Let B be an embedded 4-ball in the interior of W . Consider $W \setminus B$ as a cobodism from $-M$ to $-\partial B$. Then $\underline{F}_{W \setminus B, \mathfrak{s}_{\omega}|_{W \setminus B}; [\omega|_{W \setminus B}]}^+(c^+(\xi; [\omega|_M])) \neq 0$, where \mathfrak{s}_{ω} is the $\text{Spin}^{\mathbb{C}}$ -structure on W associated to ω .*

3. GENERALIZED SLICING BENNEQUIN INEQUALITIES

In this section, we adapt T. Mrowka and Y. Rollin's idea into the Heegaard-Floer setting, and prove Theorem 1.2.


Lemma 3.1 ([9]). *Let W be an oriented 4-manifold with connected boundary $\partial W = M$, ξ a contact structure on M , and \mathfrak{s} a $\text{Spin}^{\mathbb{C}}$ -structure on W with an isomorphism $h : \mathfrak{s}|_M \rightarrow \mathfrak{t}_\xi$. Assume K is a Legendrian knot in (M, ξ) , and $\Sigma \subset W$ is an embedded surface bounded by K . Then there are a Legendrian knot K' in (M, ξ) and an embedded surface $\Sigma' \subset W$ bounded by K' , such that $tb(K', \Sigma') \geq 1$, $\chi(\Sigma') \leq -1$, and $tb(K', \Sigma') + |r(K', \Sigma', \mathfrak{s}, h)| + \chi(\Sigma') = tb(K, \Sigma) + |r(K, \Sigma, \mathfrak{s}, h)| + \chi(\Sigma)$.*

Proof. Let p be a point on K . There is a neighborhood U of p so that $(U, \xi|_U) \cong (\mathbb{R}^3, \xi_0)$, where ξ_0 is the standard contact structure on \mathbb{R}^3 defined by $dz - ydx$. By the following Legendrian Reidemeister move, we create a pair of cusps on the front projection of $K \cap U$ (c.f. [5]).



FIGURE 1. Creating cusps

Near a cusp, connect sum K with a Legendrian righthand trefoil knot T_r in U with $tb(T_r) = 1$. We get a new Legendrian knot K_1 and an embedded surface Σ_1 in W bounded by K_1 , s.t., $tb(K_1, \Sigma_1) = tb(K, \Sigma) + 1$, $\chi(\Sigma_1) = \chi(\Sigma) - 1$, and $|r(K_1, \Sigma_1, \mathfrak{s}, h)| = |r(K, \Sigma, \mathfrak{s}, h)|$. Repeat this process, we will find a K' and a Σ' with the properties specified in the lemma. \square

FIGURE 2. Connect summing with T_r

Proof of Theorem 1.2. By Lemma 3.1, we only need prove the theorem for K and Σ with $tb(K, \Sigma) \geq 1$ and $\chi(\Sigma) \leq -1$. We assume these are true throughout the proof.

We prove part (a) first. Performing Legendrian surgery along K gives a symplectic cobordism (V, ω') from (M, ξ) to another contact 3-manifold (M', ξ') (c.f. [17, 18]). By Proposition 2.4, $F_{V, \mathfrak{s}_{\omega'}}^+(c^+(\xi')) = c^+(\xi)$. Let $\widetilde{W} = W \cup_M V$. Then, by Theorem 2.2,

we have

$$\sum_{\{\tilde{\mathfrak{s}} \in \text{Spin}^{\mathbb{C}}(\widetilde{W}) \mid \tilde{\mathfrak{s}}|_W \cong \mathfrak{s}, \tilde{\mathfrak{s}}|_V \cong \mathfrak{s}_{\omega'}\}} \pm F_{W, \tilde{\mathfrak{s}}}^+(c^+(\xi')) = F_{W, \mathfrak{s}}^+ \circ F_{V, \mathfrak{s}_{\omega'}}^+(c^+(\xi')) = F_{W, \mathfrak{s}}^+(c^+(\xi)) \neq 0.$$

Thus, there is an $\tilde{\mathfrak{s}} \in \text{Spin}^{\mathbb{C}}(\widetilde{W})$ with $\tilde{\mathfrak{s}}|_W \cong \mathfrak{s}$, and $\tilde{\mathfrak{s}}|_V \cong \mathfrak{s}_{\omega'}$, such that

$$F_{\widetilde{W} \setminus B, \tilde{\mathfrak{s}}|_{\widetilde{W} \setminus B}}^+(c^+(\xi')) \neq 0.$$

Let $h_1 : \tilde{\mathfrak{s}}|_W \rightarrow \mathfrak{s}$, $h_2 : \tilde{\mathfrak{s}}|_V \rightarrow \mathfrak{s}_{\omega'}$ be the above isomorphisms, and $h_3 : \mathfrak{s}_{\omega'}|_M \rightarrow \mathfrak{t}_{\xi}$ the natural isomorphism. And define $h : \mathfrak{s}|_M \rightarrow \mathfrak{t}_{\xi}$ by $h = h_3 \circ h_2 \circ h_1^{-1}$.

Capping off Σ by the core of the 2-handle, we get an embedded closed surface $\tilde{\Sigma}$ satisfying $\chi(\tilde{\Sigma}) = \chi(\Sigma) + 1 \leq 0$, $[\tilde{\Sigma}] \cdot [\tilde{\Sigma}] = tb(K, \Sigma) - 1 \geq 0$, and $\langle c_1(\tilde{\mathfrak{s}}), [\tilde{\Sigma}] \rangle = r(K, \Sigma, \mathfrak{s}, h)$. Next, blow up $tb(K, \Sigma) - 1$ points on the core of the 2-handle, we get a new 4-manifold \widehat{W} with a natural projection $\pi : \widehat{W} \rightarrow W$. Let $\widehat{\mathfrak{s}}$ be the lift of $\tilde{\mathfrak{s}}$ to \widehat{W} whose evaluation on each exceptional sphere is -1 , and $\widehat{\Sigma}$ be the lift of $\tilde{\Sigma}$ to \widehat{W} obtained by removing the exceptional spheres from $\pi^{-1}(\tilde{\Sigma})$. Then $\chi(\widehat{\Sigma}) = \chi(\tilde{\Sigma}) = \chi(\Sigma) + 1$, $[\widehat{\Sigma}] \cdot [\widehat{\Sigma}] = 0$, and $\langle c_1(\widehat{\mathfrak{s}}), [\widehat{\Sigma}] \rangle = r(K, \Sigma, \mathfrak{s}, h) + tb(K, \Sigma) - 1$. Also, by Theorem 2.3,

$$F_{\widehat{W} \setminus \widehat{B}, \widehat{\mathfrak{s}}|_{\widehat{W} \setminus \widehat{B}}}^+(c^+(\xi')) = F_{\widetilde{W} \setminus B, \tilde{\mathfrak{s}}|_{\widetilde{W} \setminus B}}^+(c^+(\xi')) \neq 0,$$

where $\widehat{B} \subset \widehat{W}$ is the pre-image of $B \subset W \subset \widetilde{W}$ under π . Since $[\widehat{\Sigma}] \cdot [\widehat{\Sigma}] = 0$, there is a neighborhood U of $\widehat{\Sigma}$ in \widehat{W} diffeomorphic to $\widehat{\Sigma} \times D^2$. Since the location of \widehat{B} does not affect the map $F_{\widehat{W} \setminus \widehat{B}, \widehat{\mathfrak{s}}|_{\widehat{W} \setminus \widehat{B}}}^+$, we assume that \widehat{B} is in the interior of U . Let $W_1 = \widehat{W} \setminus U$, and $W_2 = U \setminus \widehat{B}$. Then, by Theorem 2.2, there are maps

$$\underline{F}_{W_1, \widehat{\mathfrak{s}}|_{W_1}}^+ : HF^+(-M', \mathfrak{t}_{\xi'}) \rightarrow \underline{HF}^+(-\partial U, \widehat{\mathfrak{s}}|_{\partial U}; \mathbb{Z}(W_1)),$$

$$\underline{F}_{W_2, \widehat{\mathfrak{s}}|_{W_2}}^+ : \underline{HF}^+(-\partial U, \widehat{\mathfrak{s}}|_{\partial U}; \mathbb{Z}(W_1)) \rightarrow \underline{HF}^+(-\partial B, \widehat{\mathfrak{s}}|_{\partial B}; \mathbb{Z}(W_1)(W_2)),$$

such that $F_{\widehat{W} \setminus \pi^{-1}(B), \widehat{\mathfrak{s}}|_{\widehat{W} \setminus \pi^{-1}(B)}}^+ = \Theta \circ \underline{F}_{W_2, \widehat{\mathfrak{s}}|_{W_2}}^+ \circ \underline{F}_{W_1, \widehat{\mathfrak{s}}|_{W_1}}^+$, where

$$\Theta : \underline{HF}^+(-\partial B, \widehat{\mathfrak{s}}|_{\partial B}; \mathbb{Z}(W_1)(W_2)) \rightarrow HF^+(-\partial B, \widehat{\mathfrak{s}}|_{\partial B})$$

is induced by the natural projection $\theta : \mathbb{Z}(W_1)(W_2) \rightarrow \mathbb{Z}$. Specially, this implies $\underline{HF}^+(-\partial U, \widehat{\mathfrak{s}}|_{\partial U}; \mathbb{Z}(W_1)) \neq 0$. Note that $\partial U \cong \widehat{\Sigma} \times S^1$. Hence, by Theorem 2.1, we have

$$\langle c_1(\widehat{\mathfrak{s}}), [\widehat{\Sigma}] \rangle \leq -\chi(\widehat{\Sigma}),$$

that is

$$tb(K, \Sigma) + r(K, \Sigma, \mathfrak{s}, h) \leq -\chi(\Sigma).$$

Reverse the orientations of K and Σ , and repeat the whole argument. We get

$$tb(K, \Sigma) - r(K, \Sigma, \mathfrak{s}, h) \leq -\chi(\Sigma).$$

Thus,

$$tb(K, \Sigma) + |r(K, \Sigma, \mathfrak{s}, h)| \leq -\chi(\Sigma).$$

Now we use twisted Heegaard-Floer homology to prove part (b). Again, perform Legendrian surgery along K . This gives a new contact 3-manifold (M', ξ') with a weak symplectic filling $(\widehat{W}, \widehat{\omega})$ (c.f. [17, 18]). Define $\widehat{\Sigma}$ and \widehat{W} as above, i.e., by capping off Σ with the core of the 2-handle, and then blowing up $tb(K, \Sigma) - 1$ points on the core of the two handle. Let $\widehat{\omega}$ be the blown-up symplectic form on \widehat{W} . Then $(\widehat{W}, \widehat{\omega})$ is also a weak symplectic filling of (M', ξ') . Denote by $\widehat{\mathfrak{s}}$ the canonical $Spin^{\mathbb{C}}$ -structure associated to $\widehat{\omega}$. We have $\chi(\widehat{\Sigma}) = \chi(\Sigma) + 1$, $[\widehat{\Sigma}] \cdot [\widehat{\Sigma}] = 0$, and $\langle c_1(\widehat{\mathfrak{s}}), [\widehat{\Sigma}] \rangle = r(K, \Sigma, \omega) + tb(K, \Sigma) - 1$. Also, by Theorem 2.5, we have $\frac{F^+}{\widehat{W} \setminus \widehat{B}, \widehat{\mathfrak{s}}|_{\widehat{W} \setminus \widehat{B}}; [\widehat{\omega}|_{\widehat{W} \setminus \widehat{B}}]}(c^+(\xi'; [\widehat{\omega}|_{M'}])) \neq 0$. Again, assume that \widehat{B} is in the interior of U . Then, similar to part (a), $\frac{F^+}{\widehat{W} \setminus \widehat{B}, \widehat{\mathfrak{s}}|_{\widehat{W} \setminus \widehat{B}}; [\widehat{\omega}|_{\widehat{W} \setminus \widehat{B}}]}$ factors through $\underline{HF}^+(-\partial U, \widehat{\mathfrak{s}}|_{\partial U}; \mathbb{Z}[\mathbb{R}]_{[\widehat{\omega}|_{M'}]}(W_1))$, where $W_1 = \widehat{W} \setminus U$. So

$$\underline{HF}^+(-\partial U, \widehat{\mathfrak{s}}|_{\partial U}; \mathbb{Z}[\mathbb{R}]_{[\widehat{\omega}|_{M'}]}(W_1)) \neq 0,$$

and we apply Theorem 2.1 as above to prove part (b). \square

REFERENCES

- [1] D. Bennequin, *Entrelacements et équations de Pfaff*, Astérisque, **107-108** (1983), 87–161.
- [2] Y. Eliashberg, *Classification of overtwisted contact structures on 3-manifolds*, Invent. Math. **98** (1989), no. 3, 623–637.
- [3] *Filling by holomorphic discs and its applications*, Geometry of low-dimensional manifolds, 2 (Durham, 1989), 45–67, London Math. Soc. Lecture Note Ser., **151**, Cambridge Univ. Press, Cambridge, 1990.
- [4] Y. Eliashberg, *Legendrian and transversal knots in tight contact 3-manifolds*, Topological methods in modern mathematics (Stony Brook, NY, 1991), 171–193, Publish or Perish, Houston, TX, 1993.
- [5] D. Fuchs, S. Tabachnikov, *Invariants of Legendrian and transverse knots in the standard contact space*, Topology **36** (1997), no. 5, 1025–1053.
- [6] P. Ghiggini, *Ozsváth-Szabó invariants and fillability of contact structures*, arXiv:math.GT/0403367.
- [7] R. Gompf, A. Stipsicz, *4-manifolds and Kirby calculus*, American Mathematical Society, Providence, RI, 1999. xvi+558 pp. ISBN: 0-8218-0994-6.
- [8] P. Kronheimer, T. Mrowka, *Monopoles and contact structures*, Invent. Math. **130** (1997), no. 2, 209–255.
- [9] T. Mrowka, Y. Rollin, *Legendrian knots and monopoles*, arXiv:math.DG/0410559.
- [10] P. Ozsváth, Z. Szabó, *Holomorphic triangles and invariants for smooth four-manifolds*, arXiv:math.SG/0110169.
- [11] P. Ozsváth, Z. Szabó, *Holomorphic triangle invariants and the topology of symplectic four-manifolds*, arXiv:math.SG/0201049.
- [12] P. Ozsváth, Z. Szabó, *Heegaard Floer homology and contact structures*, arXiv:math.SG/0309326.
- [13] P. Ozsváth, Z. Szabó, *Holomorphic disks and genus bounds*, Geom. Topol. **8** (2004), 311–334 (electronic).
- [14] P. Ozsváth, Z. Szabó, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) **159** (2004), no. 3, 1027–1158.
- [15] P. Ozsváth, Z. Szabó, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. of Math. (2) **159** (2004), no. 3, 1159–1245.
- [16] L. Rudolph, *The slice genus and the Thurston-Bennequin invariant of a knot*, Proc. Amer. Math. Soc. **125** (1997), no. 10, 3049–3050.
- [17] A. Weinstein, *Contact surgery and symplectic handlebodies*, Hokkaido Math. J. **20** (1991), no. 2, 241–251.
- [18] H. Wu, *Legendrian Surgeries on Stabilized Legendrian Links*, arXiv:math.GT/0501074.

DEPARTMENT OF MATHEMATICS AND STATISTICS, LEDERLE GRADUATE RESEARCH TOWER, 710
NORTH PLEASANT STREET, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA 01003-9305, USA
E-mail address: `wu@math.umass.edu`