On the Hodge Conjecture for products of certain surfaces

José J. Ramón Mari Humboldt Uni. zu Berlin Institut für Mathematik Unter den Linden, 6 D - 10099 Berlin Germany E-mail: akalmahakal@hotmail.com, jjramon@mathematik.hu-berlin.de

September 11, 2018

Abstract

In this paper we prove the Hodge conjecture for arbitrary products of surfaces, $S_1 \times \cdots \times S_n$ such that $q(S_i) = 2$, $p_g(S_i) = 1$. We also prove the Hodge conjecture for arbitrary self-products of a K3 surface X such that the field $E = End_{hg}T(X)$ is CM.

0 Notation and preliminaries

Unless otherwise stated, we use the terms curve and surface to denote smooth projective curves and surfaces, resp. The term $p_g(S) = h^{2,0}(S)$ is called the geometric genus of S, and $q(S) = h^{1,0}(S) = \dim Alb(S)$ is known as the irregularity of S. For any complex projective manifold X, $H^k(X)$ will denote the group $H^k(X, \mathbb{Q})$ regarded as a (rational) Hodge structure of (pure) weight k. All Hodge structures appearing in this paper are rational and pure [2]; as usual, a Hodge cycle (of codimension p) or Hodge class of a Hodge structure V is an element $v \in V_{\mathbb{C}}^{p,p} \cap V$. We denote the subspace of Hodge cycles of V by $\mathcal{H}(V)$, and also $\mathcal{H}^p(X) = \mathcal{H}(H^{2p}(X))$ for X a smooth projective variety; consequently, $\mathcal{H}(X) = \bigoplus_{i=0}^{\dim(X)} \mathcal{H}^i(X)$ will denote the Hodge ring, or ring of Hodge classes of X.

We define the (rational) transcendental lattice T(S) of a surface S by the following orthogonal decomposition

(1)
$$H^2(S) = T(S) \oplus NS(S)_{\mathbb{Q}}$$

with respect to the cup-product. The cup-product induces, after a change of sign, a polarisation of the Hodge structure T(S) [2].

For V and W two (pure) Hodge structures of the same weight, we denote $Hom_{hg}(V, W)$ to be the space of linear maps from V to W respecting the Hodge structures. For an introduction see [2], [5].

For a Hodge structure V as above we define the **Hodge group** of V, Hg(V) to be the minimal \mathbb{Q} -defined algebraic subgroup of GL(V) such that $h(U(1)) \subset Hg(V)_{\mathbb{R}}$; here h is the representation corresponding to the Hodge bigraduation as in [2]. The following is basic in this paper:

Proposition 0.1 [2] [3] Let V be a polarisable Hodge structure. Then Hg(V) is reductive. As a result, the category of polarisable Hodge structures is semisimple abelian.

For an comprehensive survey on the Hodge conjecture for abelian varieties, as well as a detailed introduction on the Hodge group Hg(A), we refer the reader to [9] Appendix B.

Acknowledgments: I thank Prof. Bert van Geemen for a very useful and generous tutorial on the subject. I wish to express my gratitude to Prof. B.J. Totaro for valuable suggestions on the writing of this paper. I am grateful to the EU Research Training Network 'Arithmetic Algebraic Geometry' for their financial support during my PhD. The warm hospitality of the Isaac Newton Institute is gratefully acknowledged. I finally wish to thank Prof. G. E. Welters for introducing me to this subject.

1 Introduction

The purpose of this article is to prove the Hodge conjecture in two different situations of product of surfaces. The first one is the product $S_1 \times \cdots \times S_n$, where $q(S_i) = 2$ and $p_g(S_i) = 1$. (It turns out that these surfaces are birationally equivalent to abelian or elliptic isotrivial surfaces). This result generalises the Main Theorem in the author's PhD thesis [18].

The other case we consider is the following: take a K3 surface X; then the transcendental lattice T(X) is irreducible, and its endomorphism algebra is a number field $E = End_{hg}T(X)$, which can be either CM or totally real ([25] 1.5). We will prove that the Hodge conjecture for arbitrary powers of X follows from the Hodge conjecture for $X \times X$. In the case when E is CM, we use results of Mukai [14], together with an elementary lemma, to prove the Hodge conjecture for $X \times X$, and establish the result for X^n for all n by using invariant theory (see for instance [19] for similar arguments).

2 Surfaces S with $p_g = 1, q = 2$

This section is devoted to understanding the geometry of surfaces with $p_g = 1, q = 2$.

Proposition 2.1 Let S be a minimal surface with $p_g = 1, q = 2$. If S is not abelian, then S is of the form

$$S = (C' \times E')/G$$

where C' is a curve, E' is an elliptic curve and G acts faithfully on both components.

Proof: One has $\chi(\mathcal{O}_S) = 0 = 1 - q + p_g$. From Enriques' classification we see that S is non-ruled, and $K^2 \ge 0$. Also $e(S) \ge 0$ (see [1] Th. X.4), and by Noether's formula we get 0 = e(S), *i.e.* $b_2(S) = 6$, and so therefore $K^2 = 0$, which yields S elliptic. Finally, by [23] or [1] Exs. VI.22(4), VIII.22., we see that $S = (C' \times E')/G$ is a finite étale quotient such that g(E') = 1, and the proof is thus complete.

All the statements concerning motives are, unless otherwise stated, considered in the category of Chow motives modulo homological equivalence. We refer the reader to [21] for the basic notations and language.

Proposition 2.2 (Murre) [21] Let X be a surface. Then there exists a decomposition $h(X) = \bigoplus_{i=0}^{4} h^{i}(X)$; i.e. a Chow-Künneth decomposition exists in the case of surfaces (in fact, modulo rational equivalence).

Remark 2.3 From the above and the standard conjectures for abelian varieties [8], it follows that the Hodge classes on $X \times X$ inducing the projectors $H^{\bullet}(X) \twoheadrightarrow H^{i}(X) \subset H^{\bullet}(X)$ on a variety X which is a product of surfaces and abelian varieties are all algebraic, and thus X admits a decomposition $h(X) \simeq \bigoplus_{i=0}^{2\dim(X)} h^{i}(X)$ modulo homological equivalence. This result will be used throughout.

Proposition 2.4 Let S be a minimal, not abelian surface such that $p_g = 1, q = 2$. Notations being as in Proposition 2.1, the following cases hold:

- (a) either g(E'/G) = 1 and the Albanese map a induces an isomorphism $h(S) \simeq h(Alb S)$, or
- (b) g(E'/G) = 1 and the Albanese map a sends S onto a curve B. It turns out that B = C'/G and the Albanese fibration

$$S = (C' \times E')/G \to a(S) = B = C'/G$$

is the canonical projection.

Proof: The following argument holds in both cases [4] [23]: $H^1((C' \times E')/G) = H^1(C'/G) \oplus H^1(E'/G)$. Since $q(S) = \frac{1}{2}b_1(S)$, we have

(2)
$$q(S) = g(C'/G) + g(E'/G)$$

and so the following cases are possible.

(a) g(E'/G) = 1, g(C'/G) = 1. In this case G acts on E' by translations, and $A = C'/G \times E'/G$ is an abelian surface; the natural map

$$\phi: S = (C' \times E')/G \to C'/G \times E'/G = A$$

yields an isomorphism on H^1 by the above (and so on H^3); therefore Alb $S \sim A$, whence $h^1(S) \simeq h^1(A)$. On H^2 , the following holds:

$$H^{2}(S) = H^{2}(C' \times E')^{G} = H^{2}(C'/G \times E'/G),$$

for G acts freely on $C' \times E'$ and trivially on $H^{\bullet}(E')$; this proves that $\phi|H^2$ is an isomorphism, thus establishing the result.

(b) Let B = C'/G. In this case we have g(E'/G) = 0 and g(B) = 2. The natural map

$$p: S = (C' \times E')/G \to C'/G = B$$

satisfies q(S) = g(B) by Formula (2), and therefore coincides with the Albanese fibration [1]; see also [24] Ch. 9.

2.1 The case g(E'/G) = 0

Let S satisfy case (b) of Proposition 2.4, and let $H \subset G$ be the subgroup of translations on E'. Since $H - \{1\}$ coincides with the set of fixed-point-free transformations of E' in G, we have a split exact sequence (we now fix a section σ)

$$1 \to H \to G \leftrightarrows \mathbb{Z}_n \to 1,$$

where $\mathbb{Z}_n \hookrightarrow Aut_P(E')$ for P fixed point of a generator ϕ of $\sigma(\mathbb{Z}_n)$. Clearly $n \in \{2, 3, 4, 6\}$.

The following proposition is a reduction to the case $G = \mathbb{Z}_n, H = \{1\}$.

Proposition 2.5 Let C = C'/H, E = E'/H. If the natural action μ of $\mathbb{Z}_n = G/H$ on $C \times E$ is étale, then the natural map

$$\beta: S = (C' \times E')/G \to S' = (C \times E)/\mathbb{Z}_n$$

yields an isomorphism of motives $h(S) \simeq h(S')$.

Proof: The proof is similar to that of Proposition 2.4(b).

2.2 μ is free

We suppose g(E'/G) = 0, notations being as above. We are going to prove that this case meets the hypotheses of Proposition 2.5.

Remark 2.6 Consider the Hodge structure

$$V = \left[H^1(C') \otimes H^1(E')\right]^G = \left[H^1(C) \otimes H^1(E)\right]^{\mathbb{Z}_n}.$$

Then

$$H^2(S) = V \oplus \mathbb{Q}(-1)^{\oplus 2}$$

and V has Hodge numbers dim $V^{2,0} = 1 = \dim V^{0,2}$, dim $V^{1,1} = 2$.

Consider the action of G/H on JC; let ϕ be a generator of G/H such that $\phi^*|H^{1,0}(E) = \omega$ where $\omega = e^{2\pi i/n}$; let $Q_n(x)$ denote the cyclotomic polynomial of order n.

Theorem 2.7 Let $P := \ker Q_n(\phi_*)^0 \subset JC$. Then $\dim P = 1$ for n = 2 and $\dim P = 2$ for n = 3, 4, 6. The quotient map $C \to C/\mathbb{Z}_n = B$ is étale in all cases.

Proof of Theorem 2.7: Consider V as above. It is clear that

(3)
$$V = \left(H^1(P) \otimes H^1(E)\right)^{\mathbb{Z}_n}$$

In the case n = 2, ϕ acts on both vector spaces as -Id, so $V = H^1(P) \otimes H^1(E)$, whence dim P = 1 by inspection. For n = 3, 4, 6, let χ be the character of \mathbb{Z}_n such that $H^{1,0}(E) = \chi$; then $H^{1,0}(P) = a\chi \oplus b\overline{\chi}$. Inspecting Hodge numbers as above and using Remark 2.6 we find a = b = 1, which in turn yields dim P = 2.

From the above we conclude that the action μ of \mathbb{Z}_n on C has no fixed points. This follows from [20] Lemma 1.5; alternatively one can derive this result from several Riemann-Hurwitz type inequalities.

Corollary 2.8 The motive of a surface $S = (C' \times E')/G$ with $G \neq H$ is isomorphic to that of a surface $(C \times E)/\mathbb{Z}_n$ with $H = \{1\}$. In other words, the conclusion of Proposition 2.5 holds true always.

2.3 $h^2(S) \simeq h^2(A)$

We now consider S as above, *i.e.* with cyclic $G = \mathbb{Z}_n$, such that B = C/G is a genus 2 étale quotient, and find an abelian surface A such that an isomorphism of Hodge structures $H^2(S) \cong H^2(A)$ holds. The first step is to decompose P:

Lemma 2.9 The abelian surface P above splits as $P \sim E_1 \times E_1$.

Proof: Indeed, suppose that P is simple. Then $Hg(P \times E) = Hg(P) \times Hg(E)$ (due to F. Hazama; see e.g. [9]B.7.6.2; see also [11]), whence the Hodge structure $W = H^1(P) \otimes H^1(E)$ is irreducible (with dim $W^{2,0} = 2$). Hence W cannot contain V, which contradicts our hypothesis. Therefore P must split; using the \mathbb{Z}_n -decomposition of $H^1(P)$ from the Proof of Theorem 2.7 and an elementary argument we obtain $P \sim E_1 \times E_1$ for E_1 an elliptic curve, thereby completing the proof.

Let us get back to our $H^2(S)$. We had by Formula (3) and Lemma 2.9

(4)
$$H^2(S) = \mathbb{Q}(-1)^2 \oplus [H^1(P) \otimes H^1(E)]^{\mathbb{Z}_n} \subset \mathbb{Q}(-1)^2 \oplus [H^1(E_1) \otimes H^1(E)]^2.$$

Again, since the transcendental part of $H^1(E_1) \otimes H^1(E)$ has one-dimensional (2,0)-part (and is thus irreducible [5] [25]), by Formula (4) $H^2(S)$ and $H^2(E_1 \times E)$ differ only by powers of the Tate Hodge structure, which implies $H^2(S) \cong H^2(E_1 \times E)$ by counting dimensions. We have thus proven the following Proposition.

Proposition 2.10 Under the hypotheses of Proposition 2.4(b), the abelian surface $A = E_1 \times E$ is such that $H^2(S) \cong H^2(A)$ (as Hodge structures).

We now proceed to construct an algebraic cycle inducing the described isomorphism. The scheme is the following. Choose a ϕ -equivariant projection $u: JC \to P$, and consider the correspondence $\beta = (u_* \circ (alb_C)_*, id_E) \circ \pi^*$ from S to $P \times E$, where $\pi: C \times E \to (C \times E)/\mathbb{Z}_n = S$ is the natural projection. This correspondence from S to $P \times E$ realises the inclusion in Formula (4). The final step in this construction will be to cook up a correspondence from $P \times E \sim E_1 \times E_1 \times E$ sending the image of V onto $H^1(E_1) \otimes H^1(E)$ in $E_1 \times E$, which after composing can be easily extended to the sought-after isomorphism.

Lemma 2.11 Let E_1, E_2 be two elliptic curves. For every Hodge substructure V of $H^1(E_1 \times E_1) \otimes H^1(E_2)$ isomorphic to $H^1(E_1) \otimes H^1(E_2)$ there exists an algebraic correspondence α from $E_1 \times E_1 \times E_2$ to $E_1 \times E_2$ such that $\alpha_* V = H^1(E_1) \otimes H^1(E_2)$.

Proof of Lemma 2.11: It suffices to prove that every Hodge correspondence between $H^1(E_1) \otimes H^1(E_2)$ and $H^1(E_1 \times E_1) \otimes H^1(E_2)$ is algebraic. This follows from the Hodge conjecture for products of elliptic curves, due to Imai [9] [11] (see also Proposition 2.18 below.)

We are now ready to prove the following result:

Theorem 2.12 With the assumptions of this Section, the motives $h^2(S)$ and $h^2(E_1 \times E)$ are isomorphic (modulo homological equivalence).

Proof: To conclude the proof, consider the correspondence β above, which takes V to its image inside $H^1(P) \otimes H^1(E)$ of $P \times E \sim E_1 \times E_1 \times E$. Choose a projection

$$\alpha: H^1(E_1 \times E_1) \otimes H^1(E) \twoheadrightarrow H^1(E_1) \otimes H^1(E)$$

such that $\alpha | V$ is a (Hodge) isomorphism. α is algebraic by Lemma 2.11, and so the composition $\alpha \circ \beta$, also algebraic, yields the desired isomorphism.

Remark 2.13 An explicit isomorphism could be obtained by fiddling with ϕ^* as an element of $M_2(End(E_1) \otimes End(E))$, without the use of Lemma 2.11. We leave this to the reader.

2.4 The Hodge Conjecture for $S_1 \times \cdots \times S_m$, $p_g(S_i) = 1, q(S_i) = 2$

We are going to prove the following theorem:

Theorem 2.14 Let S_i be surfaces such that $p_q(S_i) = 1, q(S_i) = 2$ (S_i need not be minimal). Then the Hodge conjecture holds for $S_1 \times \cdots \times S_m$.

Remark 2.15 Let S be a surface such that $p_g = 1, q = 2$. In the former sections we have actually proven that the motive of such a surface (minimal or not) is generated (in the Tannakian sense, see [3]) by motives of abelian surfaces and elliptic curves.

The following lemma follows easily from [3](see also [9] Appendix B) and some linear algebra.

Lemma 2.16 Let A be an abelian variety of dimension ≥ 2 . Then $Hg(H^2(A)) = Hg(A)/\mu_2$. In particular, for A an abelian surface of simple CM type (F, Φ) one has $U_F(1) \simeq Hg(T(A)) = Hg(A)/\mu_2 = U_F(1)/\mu_2$.

There are two cases of a simple quartic CM field F [22] according to the Galois group of its normal closure N over \mathbb{Q} . We now solve the case $F \neq N$. In the case where F = N, there is essentially one CM-type, hence one isogeny type of abelian surfaces [18]. In the case when F is Galois and $Gal(F|\mathbb{Q}) = V_4$ the corresponding abelian surface is non-simple [22]. Therefore one is left with the cases F = N, $Gal(F|\mathbb{Q}) = C_4$ and $Gal(N|\mathbb{Q}) = D_{2,4}$.

Lemma 2.17 1. Let F be a simple CM field of degree 4, Galois over \mathbb{Q} (i.e. $Gal(F|\mathbb{Q}) = C_4$). Then

$$End_{hq}(T(A)) = F.$$

2. Let F be a CM field of degree 4 over \mathbb{Q} such that its normal closure has Galois group $D_{2,4}$ and A belong to the CM type (F, Φ) , where $\Phi = \{\sigma_1, \sigma_2\}$. Then $End_{hg}T(A) = E$, where $E \subset N$ is quartic CM and non-isomorphic to F. As a result one has the following isomorphism of algebraic groups over \mathbb{Q} :

$$U_F(1) \simeq U_E(1),$$

with E and F non-isomorphic number fields. In fact, if we write $F = F_0[\theta]$ where $\theta^2 = -\alpha$ for $\alpha \in F_0$ totally positive, then $E \cap \mathbb{R} = \mathbb{Q}\left(\sqrt{N_{F_0|\mathbb{Q}}(\alpha)}\right) \neq F_0$ (this follows from the condition on $Gal(N|\mathbb{Q})$ and also from the uniqueness of E up to automorphisms of $N|\mathbb{Q}$). Thus E and F mutually determine each other.

Proof of Lemma 2.17: One need only observe that the subfield E of \mathbb{C} spanned by the action of F^{\times} on T(A) (which can be read on $H^{2,0}(A)$) is quartic CM and not isomorphic to F. Indeed, the homomorphism of abstract groups

$$\rho: F^{\times} \to GL(H^{2,0}(A))$$

is described by $x \mapsto \rho(x) = \sigma_1(x)\sigma_2(x)$ where $\sigma_i|F_0$ are different. A little Galois theory shows that if θ is described as above and $\theta_1 \neq \pm \theta$ is an algebraic conjugate then $\theta_1^2 = \tau(\alpha)$ and $E = \mathbb{Q}[\theta + \theta_1]$. One can see that the element $(\theta + \theta_1)^2$ is a primitive element of the real quadratic extension $\mathbb{Q}(\sqrt{N_{F_0|\mathbb{Q}}(\alpha)}) \neq F_0$ since $Gal(N|\mathbb{Q}) = D_{2,4}$. The Lemma is thus established.

We state the following proposition and prove only the cases not included in Moonen and Zarhin [11]:

Proposition 2.18 Let A_i be abelian varieties of dimension 1 or 2. Then the Hodge conjecture holds for $A_1 \times \cdots \times A_r$ for r an arbitrary natural number.

Proof of Proposition 2.18: By Goursat's Lemma [9] [19] and the results of Hazama [6] and Moonen-Zarhin [11] one needs only prove the following statement. For A_i such that dim $A_i \leq 2$ and $\text{Hom}(A_1, A_2) = 0$, one has $Hg(A_1 \times A_2) = Hg(A_1) \times Hg(A_2)$. By Hazama [9], Moonen and Zarhin [11] the only case left is the following.

Let A_i be simple abelian varieties of CM type. Then $Hg(A_i) = U_{F_i}(1)$ and $Hg(A_1 \times A_2) \subset Hg(A_1) \times Hg(A_2)$ surjects onto both components, so either $Hg(A_1 \times A_2)$ is simple (and the projections are isogenies) or the former inclusion is an equality. Suppose that the projections are isogenies; in this case, $T(A_1) \otimes T(A_2)$ has a Hodge class (in fact, four such classes), and thus there is a Hodge isomorphism $T(A_1) \cong T(A_2)$. This implies that $E_i = End_{hg}T(A_i)$ are isomorphic number fields; in the case where the Galois group of $N_1 = F_1^{gal}$ over \mathbb{Q} is $D_{2,4}$, we have $E_1 \simeq E_2$ and it follows from Lemma 2.17 that $F_1 \simeq F_2$ as well. The Proposition follows in this case from [11] Proposition 4.2. For the remaining cases, there is only one CM type for F up to automorphisms of $F|\mathbb{Q}$ and the proof is similar.

Now Theorem 2.14 follows easily from Proposition 2.18 and Remark 2.15.

3 The case of powers of a K3 surface

Let X be a K3 surface, and let $\mathcal{H}^{\bullet}(X) \subset H^{\bullet}(X)$ be the ring of Hodge classes of X. Then $H^{\bullet}(X) = T(X) \oplus \mathcal{H}^{\bullet}(X)$. T(X) is an irreducible Hodge structure [5] [25], and if $E = End_{hq}T(X)$ we have an inclusion

$$E \hookrightarrow End_{\mathbb{C}}(H^{2,0}(X)) = \mathbb{C}$$

which renders E a number field. It can be shown that E is either totally real or CM [25].

The following proposition holds:

Proposition 3.1 The Hodge conjecture for X^n , for arbitrary n, holds if it holds for $X \times X$.

Proof: The ring of Hodge classes $\mathcal{H}^{\bullet}(X^n)$ is, by the above, generated by the Hodge classes in the tensor powers of T(X) up to order n and by pullbacks of algebraic classes on X via the canonical projections. Thus our result amounts to show that the ring of tensor invariants of the Hg(X)-module T(X) is generated by those of degree 2 as an algebra; it is known (see [25]) that $Hg(X)_{\mathbb{C}}$ is isomorphic to a product of special orthogonal or general linear groups, which shows (see [19]) that the ring of tensor invariants of Hg(X) is generated by the degree-2 invariants, thereby establishing the result.

We now prove the Hodge conjecture for self-products of a K3 surface X in the case where E is a CM field. We need the following elementary lemma.

Lemma 3.2 Let *E* be a *CM* number field. Then *E* is spanned as a vector space over \mathbb{Q} by elements $\alpha_i \in E$ such that $\alpha_i \overline{\alpha}_i = 1$.

Proof of Lemma 3.2: Let $\chi_0 : E^{\times} \to E^{\times}$ be given by $\chi_0(\alpha) = \alpha/\overline{\alpha}$. Suppose that the images of χ_0 do not span E over \mathbb{Q} ; then there exists $\theta \in E$ such that

$$\operatorname{Tr}_{E|\mathbb{O}}(\theta\chi_0(\alpha)) = 0 \text{ for all } \alpha \in E^{\times}.$$

Now let $\chi_{\sigma} = \sigma \circ \chi_0$ for $\sigma : E \hookrightarrow \mathbb{C}$ an embedding; by Artin's linear independence of characters, there are embeddings $\sigma \neq \tau$ such that $\chi_{\sigma} = \chi_{\tau}$, which amounts to saying that $\sigma(\alpha)/\tau(\alpha)$ is always real. It is not difficult to see that, since E is nonreal CM, this cannot hold if σ and τ are different; indeed, evaluating at α and $1 + \alpha$ for $\alpha \in E$ neither real nor purely imaginary, we see that $1 + \sigma(\alpha)$ does not belong to $\mathbb{R}(1 + \tau(\alpha))$, which leads to a contradiction, thereby establishing the Lemma.

We are now ready to prove our Theorem. See Morrison [12] for an earlier result in this direction.

Theorem 3.3 Let X be a K3 surface such that $E = End_{hg}T(X)$ is a CM field. Then the Hodge conjecture holds for arbitrary powers of X.

Proof of Theorem 3.3: By the above Lemma 3.2, it suffices to prove algebraicity for $\alpha \in E$ such that $\alpha \cdot \overline{\alpha} = 1$, *i.e.* for the Hodge isometries of the polarised Hodge structure (T(X), Q) [25]. This is a result established by Mukai, by refining former results on his theory of moduli:

Theorem 3.4 [14] Let X_1 and X_2 be K3 surfaces, and let $\psi : T(X_1) \to T(X_2)$ be a Hodge isometry. Then ψ is induced by an algebraic cycle.

References

- A. Beauville, *Complex Algebraic Surfaces*, 2nd English edition, Cambridge Univ. Press, 1996.
- [2] P. Deligne, La Conjecture de Weil pour les surfaces K3, Inv. Math 15 (1972), 206-226.
- [3] P. Deligne, J.S. Milne, A. Ogus, K.Y. Shih, *Hodge cycles, motives and Shimura varieties*, LNM 900. Springer-Verlag, Berlin-Heidelberg-New York, 1982.
- [4] E. Freitag, Über die Struktur der Funktionenkörper zu hyperabelschen Gruppen.I. J.Reine und Angew. Math 247(1971), 97-117.
- [5] B.L. van Geemen, Kuga-Satake varieties and the Hodge conjecture. The Geometry and Arithmetic of Algebraic Cycles. J.D. Lewis, B.Brent Gordon, S.Müller-Stach, N.Yui (eds.), NATO Science Series 548, World Scientific 2001.
- [6] F. Hazama, Algebraic cycles on nonsimple abelian varieties, Duke Math. J. Vol. 58 (1989) No. 1, pp.31-37.
- [7] U. Jannsen, Motives, numerical equivalence, and semisimplicity. Inv. Math. 107 (1992) 447-452.
- [8] S. Kleiman, Algebraic cycles and the Weil conjectures. In: Dix Exposés sur la Cohomologie des Schémas, North Holland, Amsterdam 1968, pp. 359-386.

- [9] J. D. Lewis, A survey of the Hodge conjecture, 2nd Edition, AMS Providence, RI 1998 (with an Appendix by B. Brent Gordon on the Hodge conjecture on abelian varieties.)
- [10] Yu I. Manin, Correspondences, motifs and monoidal transformations, Math. USSR Sbornik 6 (1968), 439-470.
- [11] B.J.J. Moonen, Yu G. Zarhin, Hodge cycles on abelian varieties of low dimension. Math.Ann. 315(1999), No. 4, 711-733.
- [12] D. R. Morrison, Algebraic cycles on products of surfaces. In: Proceedings, Algebraic Geometry Symposium, Tôhoku Univ. 1984, pp. 194-210.
- [13] S.Mukai, On the moduli space of bundles of K3 surfaces, I. Vector Bundles on Algebraic Varieties, Bombay Colloquium 1984, Oxford 1987.
- [14] S. Mukai, Vector Bundles on a K3 Surface. Proceedings of the ICM, Beijing 2002, Vol. II (invited lectures), pp. 485-502. Higher Education Press – World Scientific, Beijing 2002.
- [15] D. Mumford, Abelian Varieties, Oxford University Press, Oxford 1970.
- [16] D. Mumford, On Shimura's paper "Discontinuous groups and abelian varieties". Math. Ann. 181 (1969), 345-351.
- [17] J.P. Murre, On the motive of an algebraic surface, J. Reine und Angew. Math. 409(1990), 190-204.
- [18] J.J. Ramón Mari, On the Hodge conjecture for products of certain surfaces. PhD Thesis, U. Durham, UK 2003.
- [19] K. Ribet, Hodge classes on certain Abelian varieties, Amer. J. Math 115 (1983), 523-538.
- [20] C. Schoen, Hodge classes on self-products of a variety with an automorphism, Comp. Math. 65(1988) 3-32.
- [21] A.J. Scholl, *Classical Motives*, Proc. Symp. Pure Math. 55 (Seattle, WA 1991), Part 1, pp. 163-187. U. Jannsen, S. Kleiman, J. P. Serre (eds.) AMS, Providence, RI 1995.
- [22] G. Shimura, Y. Taniyama, Complex Multiplication of abelian varieties, Publ. Math. Soc. Japan, vol. 6. Math. Soc. Japan, 1961.
- [23] F. Serrano, The Picard group of a quasi-bundle, Manuscripta Mathematica 73 (1991), 63-82.
- [24] K. Ueno, Classification Theory of Algebraic Varieties and Compact Complex Spaces, LNM 439, Springer-Verlag, Berlin-Heidelberg, 1975.
- [25] Yu. G. Zarhin, Hodge groups of K3 surfaces, J. Reine und Angew. Math. 341 (1983), 193-220.