Probing mapping class groups using arcs

R. C. Penner

Departments of Mathematics and Physics/Astronomy University of Southern California Los Angeles, CA 90089

Abstract The action of the mapping class group of a surface on the collection of homotopy classes of disjointly embedded curves or arcs in the surface is discussed here as a tool for understanding Riemann's moduli space and its topological and geometric invariants. Furthermore, appropriate completions, elaborations, or quotients of the set of all such homotopy classes of curves or arcs give for instance Thurston's boundary for Teichmüller space or a combinatorial description of moduli space in terms of fatgraphs. Related open problems and questions are discussed.

Introduction

One basic theme in this paper on open problems is that the action of the mapping class group on spaces of measured foliations, and in particular on weighted families of curves and arcs, is calculable and captures the dynamics of homeomorphisms of the surface both on the surface itself and on its Teichmüller space. Another basic theme is that suitable spaces of arcs can be exploited to give group-theoretic and other data about the mapping class groups as well as their subgroups and completions. The author was specifically given the task by the editor of presenting open problems on his earliest and latest works in [23] and [46-49], which respectively develop these two themes and are also surveyed here.

1. Definitions

Let $F = F_{g,r}^s$ denote a smooth oriented surface of genus $g \ge 0$ with $r \ge 0$ boundary components and $s \ge 0$ punctures, where 2g-2+r+s > 0. The mapping class group MC(F)of F is the collection of isotopy classes of all orientation-preserving homeomorphisms of F, where the isotopies and homeomorphisms necessarily setwise fix the boundary ∂F and setwise fix the collection of punctures. Let PMC(F) < MC(F) denote the pure mapping class group whose homeomorphisms and isotopies pointwise fix each puncture and each boundary component.

But one aspect of Bill Thurston's seminal contributions to mathematics, [37] among other things provides a natural spherical boundary of the Teichmüller space $\mathcal{T}(F)$ of a surface F with negative Euler characteristic by an appropriate space of "projectivized measured foliations of compact support" in F. Specifically [37,9,32], let $\mathcal{MF}(F)$ denote the space of all isotopy classes rel ∂F of Whitehead equivalence classes of measured foliations in F, and let $\mathcal{MF}_0(F)$ denote the subspace comprised of those foliations with "compact support", i.e., leaves are disjoint from a neighborhood of the punctures and boundary, and no simple closed leaf is puncture- or boundary-parallel.

Define the projectivized spaces

$$\mathcal{PF}_{0}(F) = [\mathcal{MF}_{0}(F) - \{\vec{0}\}]/\mathbf{R}_{>\mathbf{0}}$$
$$\subseteq [\mathcal{MF}(F) - \{\vec{0}\}]/\mathbf{R}_{>\mathbf{0}} = \mathcal{PF}(F).$$

where $\vec{0}$ denotes the empty foliation and $\mathbf{R}_{>0}$ acts by homothety on transverse measures. Thus, $\mathcal{MF}(F_{g,r-1}^{s+1}) \subseteq \mathcal{MF}(F_{g,r}^s)$ and $\mathcal{MF}_0(F_{g,r}^s) \approx \mathcal{MF}_0(F_{g,0}^{r+s})$ with corresponding statements also for projectivized foliations.

A basic fact (as follows from the density of simple closed curves in $\mathcal{PF}(F)$) is that the action of MC(F) or PMC(F) on $\mathcal{PF}_0(F)$ has dense orbits, so the quotients are non-Hausdorff. (The action is actually minimal in the sense that every orbit is dense, and in fact, the action is ergodic for a natural measure class as independently shown by Veech [43] and Masur[44].)

We shall say that a measured foliation or its projective class fills F if every essential simple closed curve has positive transverse measure and that it quasi fills F if every essential curve with vanishing transverse measure is puncture-parallel.

Define the pre-arc complex $\mathcal{A}'(F)$ to be the subspace of $\mathcal{PF}(F)$ where each leaf in the underlying foliation is required to be an arc connecting punctures or boundary components, and define the open subspace $\mathcal{A}'_{\#}(F) \subseteq \mathcal{A}'(F)$ where the foliations are furthermore required to fill F.

In particular in the punctured case when r = 0, $s \ge 1$, and F has negative Euler characteristic the product $\mathcal{T}(F) \times \Delta^{s-1}$ of Teichmüller space with an open (s-1)-dimensional simplex Δ^{s-1} is canonically isomorphic to $\mathcal{A}'_{\#}(F)$, and in fact this descends to an isomorphism between the filling arc complex $\mathcal{A}_{\#}(F) = \mathcal{A}'_{\#}(F)/PMC(F)$ and the product $\mathcal{M}(F) \times \Delta^{s-1}$ of Riemann's (pure) moduli space $\mathcal{M}(F) = \mathcal{T}(F)/PMC(F)$ with the simplex [12,15,36,27] . Thus, the arc complex $\mathcal{A}(F) = \mathcal{A}'(F)/PMC(F)$ forms a natural combinatorial compactification of $\mathcal{A}_{\#}(F) \approx \mathcal{M}(F) \times \Delta^{s-1}$. (In fact, this is not the most useful combinatorial compactification when s > 1, cf. [30,46], where one chooses from among the punctures a distinguished one.) Another special case r, s of interest here is the case of *bordered surfaces* when $r \geq 1$ and $s \geq 0$. Choose one distinguished point on each boundary component, and define the analogous complexes $Arc'(F) \subseteq \mathcal{PF}(F)$, where leaves are required to be asymptotic to the distinguished points on the boundary (and may not be asymptotic to punctures) with its quasi filling subspace $Arc'_{\#}(F) \subseteq Arc'(F)$ and quotients $Arc_{\#}(F) = Arc'_{\#}(F)/PMC(F) \subseteq Arc(F) = Arc'(F)/PMC(F)$. In analogy to the punctured case, Arc(F) is proper homotopy equivalent to Riemann's moduli space of F (as a bordered surface with one distinguished point in each geodesic boundary component) modulo a natural action of the positive reals [31]. Furthermore [30], Arc-complexes occur as virtual links of simplices in the \mathcal{A} -complexes, and the local structure of the \mathcal{A} -complexes is thus governed by the global topology of Arc-complexes. In fact, the Arc-complexes are stratified spaces of a particular sort as explained in §5.

There are other geometrically interesting subspaces and quotients of $\mathcal{MF}(F)$ or $\mathcal{PF}(F)$, for instance the curve complex of Harvey [14] or the complex of pants decompositions of Hatcher-Thurston [13], which are surely discussed elsewhere in this volume.

As was noted before, the quotients $\mathcal{PF}_0(F)/PMC(F) \subseteq \mathcal{PF}(F)/PMC(F)$ are maximally non-Hausdorff, and yet for r > 0, $\mathcal{PF}(F)/PMC(F)$ contains as an open dense subset the (Hausdorff) stratified space Arc(F) = Arc'(F)/PMC(F); in particular, for the surface $F = F_{0,r}^s$ with $r + s \leq 3$, $PMC(F_{0,r}^s)$ is the free abelian group generated by Dehn twists on the boundary, $Arc(F_{0,r}^s)$ is piecewise-linearly homeomorphic to a sphere of dimension 3r + 2s - 7, and it is not difficult to understand the non-Hausdorff space $\mathcal{PF}(F)/PMC(F)$ with its natural foliation. This leads to our first problem:

Problem 1 Understand either classically or as quantum geometric objects the non-Hausdorff quotients of $\mathcal{PL}_0(F)$ or $\mathcal{PL}(F)$ by MC(F) or PMC(F).

 $\mathcal{T}(F)$ has been quantized in [5] and [17] as surveyed in [6] and [35], respectively, and $\mathcal{PL}_0(F_{1,0}^1)$ has been quantized in [6]. In interesting contrast, [21,54] has described a program for studying real quadratic number fields as quantum tori limits of elliptic curves, thus quantizing limiting curves rather than Teichmüller space.

2. Dehn-Thurston coordinates

Fix a surface $F = F_{g,r}^s$. In this section, we introduce global Dehn-Thurston coordinates from [23] for $\mathcal{MF}_0(F)$ and $\mathcal{MF}(F)$.

Define a pants decomposition of F to be a collection \mathcal{P} of curves disjointly embedded in F so that each component of $F - \cup \mathcal{P}$ is a generalized pair of pants $F_{0,r}^s$ with r + s = 3. One easily checks using Euler characteristics that there are 3g - 3 + 2r + s curves in a pants decomposition of F and 2g - 2 + r + s generalized pairs of pants complementary to \mathcal{P} .

Given a measured foliation \mathcal{F} in a generalized pair of pants P, where ∂P has components

 ∂_i , define the triple m_i of *intersection numbers* given by the transverse measure of ∂_i of \mathcal{F} , for i = 1, 2, 3.

Dehn-Thurston Lemma Isotopy classes (not necessarily the identity on the boundary) of Whitehead equivalence classes of non-trivial measured foliations in the pair of pants P with no closed leaves are uniquely determined by the triple (m_1, m_2, m_3) of non-negative real intersection numbers, which are subject only to the constraint that $m_1 + m_2 + m_3$ is positive.



Figure 1 Constructing measured foliations in pants

Proof The explicit construction of a measured foliation realizing a putative triple of intersection numbers is illustrated in Figure 1 in the two representative cases that m_1, m_2, m_3 satisfy all three possible (weak) triangle inequalities in Figure 1a or perhaps $m_1 \ge m_2 + m_3$ in Figure 1b. The other cases are similar, and the projectivization of the positive orthant in (m_1, m_2, m_3) -space is illustrated in Figure 2. Elementary topological considerations show that any measured foliation of P is isotopic to a unique such foliations keeping endpoints of arcs in the boundary of P, completing the proof. q.e.d.



Figure 2 Measured foliations in pants

In order to refine the Dehn-Thurston Lemma and keep track of twisting around the boundary, we shall introduce in each component ∂_i of the boundary ∂F an arc $w_i \subseteq \partial_i$ called a window, for i = 1, 2, 3. We require that the support of the restriction to ∂P of \mathcal{F} lie in the union of the windows, so-called windowed measured foliations. (Collapsing each window to a point gives a surface with a distinguished point in each boundary component, so a windowed measured foliation in P gives rise to an element of Arc(P) in the sense of §1.) We seek the analogue of the Dehn-Thurston Lemma for windowed isotopy classes.

To this end, there are two conventions to be made:

1) when a leaf of \mathcal{F} connects a boundary component to itself, i.e., when it is a loop, then it passes around a specified leg, right or left, of P as illustrated in Figure 3a-b, i.e., it contains a particular boundary component or puncture in its complementary component with one cusp;

2) when a leaf is added to the complementary component of a loop in P with two cusps, then it either follows or precedes the loop as illustrated in Figure 3c-d.

For instance in Figure 2, the conventions are: 1) around the right leg for loops; and 2) on the front of the surface. We shall call these the *standard twisting conventions*.



Figure 3c Arc follows loop.

Figure 3d Arc precedes loop.

Figure 3 Twisting conventions.

Upon making such choices of convention, we may associate a twisting number $t_i \in \mathbf{R}$ to \mathcal{F} as follows. Choose a regular neighborhood of ∂P and consider the sub-pair of pants $P_1 \subseteq P$ complementary to this regular neighborhood. Given a weighted arc family α in P, by the Dehn-Thurston Lemma, we may perform an isotopy in P supported on a neighborhood of P_1 to arrange that $\alpha \cap P_1$ agrees with a conventional windowed arc family in P_1 (where the window in ∂P_1 arises from that in ∂P in the natural way via a framing of the normal bundle to ∂P in P).

For such a representative of \mathcal{F} , we finally consider its restriction to each annular neighborhood A_i of ∂_i . Choose another arc *a* whose endpoints are *not* in the windows (and again such an arc is essentially uniquely determined up to isotopy rel windows from a framing

of the normal bundle to ∂P in P in the natural way); orient a and each component arc of $(\cup \alpha) \cap A_i$ from ∂P_1 to ∂P , and let t_i be the signed (weighted) intersection number of a with the (weighted) arc family $(\cup \alpha) \cap A_i$, for i = 1, 2, 3.



4a Right twisting for $t_i \ge 0$ **4b** Left twisting for $t_i \le 0$

Figure 4 Windowed measured foliations in the annulus

As illustrated in Figure 4, all possible real twisting numbers $-m_i \leq t_i \leq m_i$ arise provided $m_i \neq 0$, where again in this figure, the indicated "weight" of a component arc is the width of a band of leaves parallel to the arc. By performing Dehn twists along the core of the annulus, it likewise follows that every real twisting number t_i occurs provided $m_i \neq 0$. Again, elementary topological considerations show that each windowed isotopy class of a windowed measured foliation is uniquely determined by its invariants:

Lemma 1 Points of $\mathcal{MF}(P)$ are uniquely determined by the triple $(m_i, t_i) \in \mathbf{R}_{\geq 0} \times \mathbf{R}$, which are subject only to the constraint that $\forall i = 1, 2, 3(m_i = 0 \Rightarrow t_i \geq 0)$.

One difference between the Dehn-Thurston Lemma and Lemma 1 is that closed leaves are permitted in the latter (but not in the former), where the coordinates $m_i = 0$ and $t_i = |t_i| > 0$ correspond to the class of a foliated annulus of width t_i whose leaves are parallel to ∂_i . In the topology of projective measured foliations, extensive twisting to the right or left about ∂_i approaches the curve parallel to ∂_i . One imagines identifying in the natural way the ray $\{0\} \times \mathbf{R}_{\geq 0}$ with the ray $\{0\} \times \mathbf{R}_{\leq 0}$ in the half plane $\mathbf{R}_{\geq 0} \times \mathbf{R}$ and thinks therefore of (m_i, t_i) as lying in the following quotient homeomorph of \mathbf{R}^2 :

$$R = (\mathbf{R}_{>0} \times \mathbf{R})/\text{antipodal map.}$$

We shall also require the subspace

$$Z = (\mathbf{Z}_{>0} \times \mathbf{Z})/\text{antipodal map},$$

which corresponds to the collection of all disjointly embedded weighted curves and arcs in F with endpoints in the windows.

Arguing as above with an annular neighborhood of a pants decomposition, one concludes:

Theorem 2 [23,32] Given an isotopy class of pants decomposition \mathcal{P} of $F = F_{g,r}^s$, where each pants curve is framed, there is a homeomorphism between $\mathcal{MF}(F)$ and the space of all

pairs $(m_i, t_i) \in R$ as *i* ranges over the elements of \mathcal{P} . Likewise, there is a homeomorphism between $\mathcal{MF}_0(F)$ and the space of all pairs $(m_i, t_i) \in R$ as *i* ranges over the elements of $\mathcal{P} - \partial F$. In particular, $\mathcal{PF}_0(F) \approx S^{3g-3+r+s}$ and $\mathcal{PF}(F) \approx S^{3g-3+2r+s}$.

There is the following "standard problem" about which not much is known (on the torus, it devolves to greatest common divisors, cf. [6], and see [11] for genus two):

Problem 2 Given a tuple $\times_{i=1}^{N}(m_i, t_i) \in \mathbb{Z}^N$, give a tractable expression in terms of Dehn-Thurston or other coordinates for the number of components of the corresponding weighted family of curves and arcs.

There is an algorithm which leads to a multiply weighted curve from an integral measure on a general train track akin to that on the torus gotten by serially "splitting" the track, cf. [6], but we ask in Problem 2 for a more closed-form expression. See also Problem 3. A related problem which also seems challenging is to describe $\mathcal{A}'(F)$ or Arc'(F) in Dehn-Thurston coordinates on $\mathcal{MF}(F)$.

This class of curve and arc component counting problems might be approachable using the quantum path ordering techniques of [5,6] or with standard fermionic statistical physics [57,29].

3. Mapping class action on Dehn-Thurston coordinates

As already observed by Max Dehn [8] in the notation of Theorem 2, a Dehn twist on the i^{th} pants curve in a pants decomposition of $F = F_{g,r}^s$ acts linearly, leaving invariant all coordinates (m_j, t_j) , for $j \neq i$, and sending

$$\begin{array}{l} (\dagger) & m_i \mapsto m_i, \\ t_i \mapsto t_i \pm m_i \end{array}$$

As proved by Hatcher-Thurston [13] using Cerf theory, the two elementary transformations or moves illustrated in Figure 5 act transitively on the set of all pants decompositions of any surface F. Thus, any Dehn twist acts on coordinates by the conjugate of a linear map, where the conjugating transformation is described by compositions of "elementary transformations" on Dehn-Thurston coordinates corresponding to the elementary moves. More explicitly, it is not difficult [23] to choose a finite collection of pants decompositions of F whose union contains all the curves in Lickorish's generating set [20] and calculate the several compositions of elementary moves relating them.

In any case, the calculation of the action of Dehn twist generators for MC(F) thus devolves to that of the two elementary moves on Dehn-Thurston coordinates. This problem was suggested in [8], formalized in [38], and solved in [23] as follows.

Let \lor and \land respectively denote the binary infimum and supremum.

Given an arc family in the pair of pants, we introduce the notation ℓ_{ij} for the arc connecting boundary components *i* and *j*, for i, j = 1, 2, 3, where the windowed isotopy class of ℓ_{ii} depends upon the choice of twisting conventions. Given a weighted arc family in the pair of pants, the respective weights λ_{ij} , for $\{i, j, k\} = \{1, 2, 3\}$, on these component arcs are given in terms of the intersection numbers m_1, m_2, m_3 by the following formulas:

$$2\lambda_{ii} = (m_i - m_j - m_k) \lor 0$$

$$2\lambda_{ij} = (m_i + m_j - m_k) \lor 0, \text{ for } i \neq j,$$

and the intersection numbers are in turn given by $m_i = 2\lambda_{ii} + \lambda_{ij} + \lambda_{ik}$.



Figure 5 The elementary transformations.

Theorem 3 [23,32] Adopt the standard twisting conventions, the enumeration of curves indicated in Figure 5, and let (m_i, t_i) denote the Dehn-Thurston coordinates of a measured foliation with respect to the pants decomposition \mathcal{P} .

First Elementary Transformation Let λ_{ij} denote the weight of ℓ_{ij} with respect to \mathcal{P} and λ'_{ij} the weight with respect to \mathcal{P}' , so in particular, $r = \lambda_{12} = \lambda_{13}$. Then the first elementary transformation is given by the following formulas.

$$\begin{split} \lambda'_{11} &= (r - |t_1|) \lor 0, \\ \lambda'_{12} &= \lambda'_{13} = L + \lambda_{11}, \\ \lambda'_{23} &= |t_1| - L, \\ t'_2 &= t_2 + \lambda_{11} + (L \land t_1) \lor 0, \\ t'_1 &= -sgn(t_1) \ (\lambda_{23} + L), \end{split}$$

where $L = r - \lambda'_{11}$ and $sgn(x) \in \{\pm 1\}$ is the sign of $x \in \mathbf{R}$, with sgn(0) = -1.

(The formulas [24] above correct a typographical error in [32].)

Second Elementary Transformation Let λ_{ij} denote the weight of ℓ_{ij} in the bottom pair of pants for \mathcal{P} and κ_{ij} the weight in the top pair or pants, and let λ'_{ij} denote the weight in the left pair of pants for \mathcal{P}' and κ'_{ij} in the right pair or pants. The second elementary transformation is given by the following formulas.

$$\begin{split} \kappa_{11}' &= \kappa_{22} + \lambda_{33} + (L - \kappa_{13}) \lor 0 + (-L - \lambda_{12}) \lor 0, \\ \kappa_{22}' &= (L \land \lambda_{11} \land (\kappa_{13} - \lambda_{12} - L)) \lor 0, \\ \kappa_{33}' &= (-L \land \kappa_{11} \land (\lambda_{12} - \kappa_{13} + L)) \lor 0, \\ \kappa_{23}' &= (\kappa_{13} \land \lambda_{12} \land (\kappa_{13} - L) \land (\lambda_{12} + L)) \lor 0, \\ \kappa_{12}' &= -2\kappa_{22}' - \kappa_{23}' + \kappa_{13} + \kappa_{23} + 2\kappa_{33}, \\ \kappa_{13}' &= -2\kappa_{33}' - \kappa_{23}' + \lambda_{12} + \lambda_{23} + 2\lambda_{22}, \\ \lambda_{11}' &= \lambda_{22} + \kappa_{33} + (K - \lambda_{13}) \lor 0 + (-K - \kappa_{12}) \lor 0, \\ \lambda_{22}' &= (K \land \kappa_{11} \land (\lambda_{13} - \kappa_{12} - K)) \lor 0, \\ \lambda_{33}' &= (-K \land \lambda_{11} \land (\kappa_{12} - \lambda_{13} + K)) \lor 0, \\ \lambda_{23}' &= (\lambda_{13} \land \kappa_{12} \land (\lambda_{13} - K) \land (\kappa_{12} + K)) \lor 0, \\ \lambda_{12}' &= -2\lambda_{22}' - \lambda_{23}' + \lambda_{13} + \lambda_{23} + 2\lambda_{33}, \\ \lambda_{13}' &= -2\lambda_{33}' - \lambda_{23}' + \kappa_{12} + \kappa_{23} + 2\kappa_{22}, \\ t_{2}' &= t_{2} + \lambda_{33} + ((L + \kappa_{33}' - \kappa_{22}') \lor (\kappa_{23}' + 2\kappa_{33}' - \lambda_{12})) \land 0, \\ t_{3}' &= t_{3} - \kappa_{33}' + ((L + \kappa_{33}' - \kappa_{22}') \lor (\kappa_{23}' + 2\kappa_{33}' - \lambda_{12})) \land 0, \\ t_{4}' &= t_{4} - \lambda_{33}' + ((K + \lambda_{33}' - \lambda_{22}') \lor (\lambda_{23}' + 2\lambda_{33}' - \kappa_{12})) \land 0, \\ t_{5}' &= t_{5} + \kappa_{33} + ((\kappa_{13} - \kappa_{23}' - 2\kappa_{22}') \land (L + \kappa_{33}' - \kappa_{22}')) \lor 0, \\ t_{1}' &= \kappa_{22} + \lambda_{22} + \kappa_{33} + \lambda_{33} - (\lambda_{11}' + \kappa_{11}' + (t_{2}' - t_{2}) + (t_{5}' - t_{5})) \\ &+ [(sgn(L + K + \lambda_{33}' - \lambda_{22}' + \kappa_{33}' - \kappa_{22}')] (t_{1} + \lambda_{33}' + \kappa_{33}'), \\ \end{split}$$

where $L = \lambda_{11} + t_1$, $K = \kappa_{11} + t_1$, and $sgn(x) \in \{\pm 1\}$ is the sign of $x \in \mathbf{R}$, with

$$sgn(0) = \begin{cases} +1, & if \ \lambda_{12} - 2\kappa'_{33} - \kappa'_{23} \neq 0; \\ -1, & otherwise. \end{cases}$$

These formulas are derived in [23], in effect, by performing explicit isotopies of arcs in certain covers of $F_{1,1}^0$ and $F_{0,4}^0$. Their computer implementation has been useful to some for analyzing specific mapping classes, e.g., fibered knot monodromies given by Dehn twists.

It is notable that the formulas are piecewise-*integral*-linear or PIL, cf. [37]. Furthermore, all of the "corners in the PIL structure actually occur", so the formulas are non-redundant in this sense; on the other hand, a given word in Lickorish's generators (i.e., the composition of conjugates of linear mappings (†) by specific PIL transformations given by finite compositions of the two elementary transformations) may not have "all possible corners occur". Insofar as continuous concave PIL functions are in one-to-one correspondence with tropical polynomials [40,41], we are led to the following problem:

Problem 3 Give a useful (piecewise) tropical description of the two elementary transformations. One thus immediately derives a (piecewise) tropical polynomial representation of the mapping class groups. What properties does it have, for instance under iteration?

As alternative coordinates, [9] describes a family of curves whose intersection numbers alone coordinatize measured foliations of compact support (but there are relations), and presumably these intersection numbers could be computed using Theorem 3.

We also wonder what are further applications or consequences of all these formulas.

4. Pseudo-Anosov maps and the length spectrum of moduli space

Thurston's original construction of pseudo-Anosov (pA) mappings [37] (cf. [9]) was generalized in [23] (cf. [25,39]) to give the following recipe for their construction:

Theorem 4 [23,25] Suppose that C and D are each families of disjointly embedded essential simple closed curves so that each component of $F - (C \cup D)$ is either disk, a once-punctured disk, or a boundary-parallel annulus. Let w be any word consisting of Dehn twists to the right along elements of C and to the left along elements of D so that the Dehn twist on each curve of C or D occurs at least once in w. Then w represents a pseudo-Anosov mapping class.

Problem 4 Does the recipe in Theorem 4 give virtually all pA maps? That is, given a pA map f, is there some iterate f^n , for $n \ge 1$, so that f^n arises from the recipe?

(This question from [23,25] is related [10] to the Ehrenpreis Conjecture, our Problem 14.)

In relation to Problem 4, let us mention that there are still other descriptions of pA maps up to iteration, for instance by Mosher [42] and in joint work of the author with Papadopoulos [45]; these descriptions are combinatorial rather than in terms of Dehn twists.

For a fixed surface F, consider the set of logarithms of dilatations of all pA maps supported

on F. This characteristic "spectrum" $\Sigma(F) \subseteq \mathbf{R}_{>0}$ of F is precisely the Teichmüller geoedsic length spectrum of Riemann's moduli space $\mathcal{M}(F)$. The spectrum $\Sigma(F)$ is discrete. (In fact, dilatations occur as spectral radii of integral-linear Perron-Frobenius symplectomorphisms in a range of dimensions bounded above and below in terms of the topological type of F.)

Problem 5 For a given surface F, calculate $\Sigma(F)$. More modestly, calculate the least element of $\Sigma(F)$ or the least gap among elements of $\Sigma(F)$. Characterize the number fields arising as dilatations of pA maps on F.

Problems 4 and 5 are clearly related. For instance, the recipe in Theorem 4 allows one to give estimates on least elements in Problem 5, cf. [26,2]. McMullen [50] has also given estimates and examples.

5. Arc complexes

Refining the discussion in §1, let $F_{g,\vec{\delta}}^s$ denote a bordered surface of type $F_{g,r}^s$, with r > 0and $\vec{\delta} = (\delta_1, \ldots, \delta_r)$ an *r*-dimensional vector of natural numbers $\delta_i \ge 1$, where there are δ_i distinguished points on the *i*th boundary component of $F_{g,\vec{\delta}}^s$, for $i = 1, \ldots, r$. Construct an arc complex Arc(F) as before as the PMC(F)-orbits of isotopy classes of families of disjointly embedded essential and non-boundary parallel arcs connecting distinguished points on the boundary.

Given two bordered surfaces S_1, S_2 , we consider inclusions $S_1 \subseteq S_2$, where the distinguished points and punctures of S_1 map to those of S_2 , and S_1 is a complementary component to an arc family in S_2 (possibly an empty arc family if $S_1 = S_2$).

Define the type 1 surfaces to be the following: $F_{1,(1,1)}^0$, $F_{0,(1,1,1)}^0$, $F_{0,(1,1,1)}^1$, $F_{0,(1,1)}^2$.

Theorem 5 [46] The arc complex Arc(F) of a bordered surface F is PL-homeomorphic to the sphere of dimension $6g - 7 + 3r + 2s + \delta_1 + \delta_2 + \cdots + \delta_r$ if and only if $M \not\subseteq F$ for any type 1 surface M. In other words, Arc(F) is spherical only in the following cases: polygons (g = s = 0, r = 1), multiply punctured polygons (g = 0, r = 1), "generalized" pairs of pants (g = 0, r + s = 3), the torus-minus-a-disk (g = r = 1, s = 0), and the once-punctured torus-minus-a-disk (g = r = s = 1). Only the type 1 surfaces have an arc complex which is a PL-manifold other than a sphere.

Problem 6 Calculate the topological type (PL-homeomorphism, homotopy, homology... type) of the *Arc*-complexes.

The first non-trivial case is the calculation of the topological type of the PL-manifolds Arc(M) for the four type 1 surfaces M.

Arc complexes as stratified spaces conjecturally have specific singularities and topology described recursively as follows. A PL sphere is a type zero space. A closed, connected, and simply connected manifold is a type one space provided it occurs among a list of four specific such (non-spherical) manifolds of respective dimensions 5,7,7, and 9, namely, the arc complexes of the four type one surfaces. For n > 1, define a type n space to be a finite polyhedron, defined up to PL-isomorphism, so that the link of each vertex in any compatible triangulation is PL isomorphic to an iterated suspension of the join of at most two spaces of type less than n.

By Theorem 5, many links of simplices are indeed of this type, and we conjecture that any arc complex is of some finite type. (As explained in [46], a specific collapsing argument in the "calculus of mapping cylinders" would give a proof a of this conjecture.)

The non-Hausdorff space $\mathcal{PF}(F)/PMC(F)$ thus contains the stratified space Arc(F) as an open dense subset, explaining one classical (i.e., non-quantum) aspect to Problem 1. In light of the stratification of Arc-complexes in general, one might hope to apply techniques such as [1,34] to address parts of Problem 6.

Problem 7 Devise a matrix model (cf. [28]) for the calculation of the Euler characteristics of *Arc*-complexes.

Problem 8 [Contributed by the referee] Does Theorem 5 say anything about the structure of the end of Riemann's moduli space? For instance, what is the homology near the end?

Take the one-point compactification F^{\times} of $F = F_{g,r}^s$, where all of the $s \ge 0$ punctures of F are identified to a single point in F^{\times} , so $F^{\times} = F$ if and only if $s \le 1$.

Let ∂ denote the boundary mapping of the chain complex $\{C_p(Arc) : p \geq 0\}$ of Arc = Arc(F). Suppose that α is an arc family in F with corresponding cell $\sigma[\alpha] \in C_p(Arc)$. A codimension-one face of $\sigma[\alpha]$ of course corresponds to removing one arc from α , and there is a dichotomy on such faces $\sigma[\beta]$ depending upon whether the rank of the first homology of $F^{\times} - \bigcup \beta$ agrees with or differs by one from that of $F^{\times} - \bigcup \alpha$. This dichotomy decomposes ∂ into the sum of two operators $\partial = \partial^1 + \partial^2$, where ∂^2 corresponds to the latter case.

The operators ∂^1, ∂^2 are a pair of anti-commuting differentials, so there is a spectral sequence converging to $H_*(Arc)$ corresponding to the bi-grading

$$E_{u,v}^0 = \{ \text{chains on } \sigma[\alpha] \in C_p(Arc) : v = -\text{rank}(H_1(F_\alpha)) \text{ and } u = p - v \},\$$

where $\partial_1 : E^0_{u,v} \to E^0_{u-1,v}$ and the differential of the E^0 term is $\partial_2 : E^0_{u,v} \to E^0_{u,v-1}$.

It is not quite fair to call it a problem, nor a theorem since the argument is complicated and has not been independently checked, but we believe that this spectral sequence collapses in its E^1 -term to its top horizontal row except in dimension zero. Thus, the homology of *Arc* is the ∂_1 -homology of the ∂_2 -kernels in the top row, and on the other hand, it follows from [31] that the ∂_1 -homology of the top row itself agrees with that of uncompactified Riemann's moduli space $\mathcal{M}(F)$ As discussed in [30,46], the stratified structure of the arc complexes for bordered surfaces gives a corresponding stratified structure to $\mathcal{A}(F)$ for punctured F. This may be enough to re-visit the calculations of [19] and [16,22] with an eye towards avoiding technical difficulties with the Deligne-Mumford compactification $\overline{\mathcal{M}}(F)$.

6. Cell decompositions of $\mathcal{M}(F)$ and $\overline{\mathcal{M}}(F)$

For the next several sections unless otherwise explicitly stated, surfaces F will be taken to be once-punctured and without boundary. This is done for simplicity in order for instance that the moduli space $\mathcal{M}(F)$ itself, rather than some "decoration" of it, comes equipped with an ideal cell decomposition. Nevertheless, the theory extends, the discussion applies, and the problems and conjectures we articulate are intended in the more general setting (of multiply punctured surfaces with a distinguished puncture and no boundary).

The basic combinatorial tool for studying moduli space $\mathcal{M}(F)$ is the MC(F)-invariant ideal cell decomposition of the Teichmüller space $\mathcal{T}(F)$, and there are two effective constructions: from combinatorics to conformal geometry using Strebel coordinates on fatgraphs [36,12,15], and from hyperbolic geometry to combinatorics using the convex hull construction and simplicial coordinates [27]. (See [31] for further details.)

Problem 9 [Bounded Distortion Conjecture] Given a hyperbolic structure on F, associate its combinatorial invariant, namely, an ideal cell decomposition of F together with the projective simplicial coordinate assigned to each edge. Take these projective simplicial coordinates as Strebel coordinates on the dual fatgraph to build a conformal structure on F. The underlying map on Teichmüller space is of bounded distortion in the Teichmüller metric.

As posed by Ed Witten to the author in the early 1990's, a compelling problem at that time was to find an orbifold compactification of $\mathcal{M}(F)$ which comes equipped with a cellular description in terms of suitably generalized fatgraphs. Calculations such as [16,19,22] and more might then be performed using matrix models derived from the combinatorics of this putative compactification. Perhaps the desired compactification was the Deligne-Mumford compactification or perhaps another one. The combinatorial compactification of the previous section fails to provide an orbifold but rather another stratified generalization of manifold.

Guidance from Dennis Sullivan has recently led to the following solution:

Theorem 6 Suppose that F has only one puncture. Then $\overline{\mathcal{M}}(F)$ is homeomorphic to the geometric realization of the partially ordered set of MC(F)-orbits of pairs (α, \mathcal{A}) , where α fills F, and the screen \mathcal{A} is a collection of subsets of α so that:

[Fulton-MacPherson nest condition] for any two $A, B \in \mathcal{A}$ which are not disjoint, either $A \subseteq B$ or $B \subseteq A$; [Properness] $\cup A$ is a proper subset of α , and furthermore for any $A \in A$, we likewise have $\cup \{B \in A : A \neq B \subseteq A\}$ is a proper subset of A;

[Recurrence] for any $a \in A \in A$, there is an essential simple closed curve in F, meeting $\cup \alpha$ a minimal number of times in its isotopy class, meeting only the arcs in A, and crossing a,

where inclusion of ideal cell decompositions induces the partial ordering on the set of pairs (α, \mathcal{A}) .

In effect, the elements of the screen \mathcal{A} detect how quickly hyperbolic lengths of arcs in α diverge. It is a concise new description of a combinatorial structure on $\mathcal{M}(F)$, which bears similarities to renormalization in physics, and it would be interesting to make this precise. Let us also remark that the proof of the previous theorem depends upon working in the hyperbolic category, where the "pinch curves" in a nearly degenerate structure can be detected using the coordinates. (In fact, most of the required ideas and estimates are already described in [30].)

Problem 10 Though the (virtual) Euler characteristics are already known [28,56], devise a matrix model using screens to calculate these invariants for $\overline{M}(F)$.

7. Torelli groups

Recall [52],[48] the Torelli group $\mathcal{I}_k(F)$, defined as those mapping classes on F fixing a basepoint (taken to be a puncture) that act trivially on the kth nilpotent quotient of the fundamental group of F. Since the ideal cell decomposition of $\mathcal{T}(F)$ is invariant under MC(F), it is in particular invariant under each $\mathcal{I}_k(F)$, and so the quotient "Torelli space" $T_k(F) = \mathcal{T}(F)/\mathcal{I}_k(F)$ is a manifold likewise admitting an ideal cell decomposition.

Recent work [48] with Shigeyuki Morita studies the "Torelli tower"

$$\mathcal{T}(F) \to \cdots \to T_{k+1}(F) \to T_k(F) \to \cdots \to T_1(F) \to \mathcal{M}(F)$$

of covers of Torelli spaces, each $T_k(F)$ a manifold covering the orbifold $\mathcal{M}(F)$. In particular, one essentially immediately derives infinite presentations for all of the higher Torelli groups as well as a finite presentation for instance of the "level N (classical) Torelli groups", i.e., the subgroup of MC(F) which acts identically on homology with \mathbf{Z}/N coefficients.

Problem 11 [Level N Torelli Franchetta Problem] What is the second cohomology group of the level N Torelli group?

The cell decomposition of $\overline{\mathcal{M}}(F)$ described before is compatible with the ideal cell decompositions of Torelli spaces, i.e., the fatgraph dual to an ideal cell decomposition of F admits a "homology marking" in the sense of [48] as well as admitting the structure of screens. There are thus "DM type" boundaries of each Torelli space replete with an ideal cell decomposition. It is natural to try to understand the topology of these DM-type bordifications of Torelli spaces to approach the following class of problems:

Problem 12 Calculate various group-theoretic boundaries of mapping class and Torelli groups, for instance, Tits boundaries.

Further results in [48] arise by writing an explicit one cocyle representing the first Johnson homomorphism. In effect, one checks that the putative cycle represents a crossed homomorphism by verifying the several essentially combinatorial constraints imposed by the cell decomposition, and then compares with known values of the first Johnson homomorphism. As discussed more fully in [48], one might realistically hope to find similar canonical cocycles for the higher Johnson homomorphisms as well. In particular by work of Morita [53], the Casson invariant of a homology 3-sphere is algorithmically calculable from the second Johnson homomorphism, so this is of special interest.

More explicitly, there is a combinatorial move, a "Whitehead move", that acts transitively on ideal triangulations of a fixed surface (where one removes an edge e from the triangulation to create a complementary quadrilateral and then replaces e with the other diagonal of this quadrilateral). One seeks invariants of sequences of Whitehead moves lying in an MC(F)-module satisfying three explicit combinatorial conditions just as for the Johnson homomorphism. There is thus a kind of "machine" here for producing cocycles with values in various modules by solving for expressions that satisfy certain explicit combinatorial constraints; several such invariant one cocycles have been produced in this way on the computer but so far without success for constructing the higher Johnson homomorphisms.

Very recent work with Shigeyuki Morita and Alex Bene solves a related problem: the Magnus representations (which are closely related to the Johnson homomorphisms [52]) in fact lift directly to the groupoid level of [48] as the Fox Jacobians of appropriately enhanced Whitehead moves. One works in the free fundamental group of the punctured surface with basepoint distinct from the puncture, as discussed in the closing remarks of [48], and the corresponding dual fatgraph comes equipped with a canonical maximal tree defined by greedily adding edges to the tree while traversing a small circle around the puncture starting from the basepoint. Perhaps these explicit calculations of the Magnus representations might be of utility for instance to address the following standard question:

Problem 13 What are the kernels of the Magnus representations?

Finally in [48] by taking contractions of powers of our canonical one cocycle, new combinatorially explicit cycles and cocycles on $\mathcal{M}(F)$ are constructed which on the other hand generate the tautological algebra. It is natural to wonder about the extension of these classes to the screen model for $\overline{\mathcal{M}}(F)$ and to the combinatorial compactification, and to revisit [16,19,22] in this context.

8. Closed/open string theory

This section describes recent work [49] with Ralph Kaufmann, where it turns out that the material of Section 2 together with a further combinatorial elaboration describes a reasonable model for the phenomenology of interactions of open and closed strings.

Returning now to multiply punctured surfaces $F = F_{g,\vec{\delta}}^s$ with at least one distinguished point in each boundary component as in Section 5, let D denote the set of distinguished points in the boundary, and let S denote the set of punctures. Fix a set \mathcal{B} of "brane labels", let $\mathcal{P}(\mathcal{B})$ denote its power set, and define a *brane labeling* to be a function β : $D \cup S \to \mathcal{P}(\mathcal{B})$ subject to the unique constraint that if the value $\emptyset \in \mathcal{P}(\mathcal{B})$ is taken at a point on the boundary of F, then this point is the only distinguished point in its boundary component. Complementary components to the distinguished points in the boundary are called "windows". Given a brane labeling β on a windowed surface F, define the set $D(\beta) = \{d \in D : \beta(d) \neq \emptyset\}$, and consider proper isotopy classes rel D of arc families with endpoints in windows, where the arcs are required to be non-boundary parallel in $F - D(\beta)$. Say that such an arc family is *exhaustive* if each window has at least one incident arc, and construct the space $\widetilde{Arc}(F, \beta)$ of positive real weights on exhaustive arc families.

There are various geometric operations on the spaces $\{Arc(F,\beta)\}\$ induced by gluing together measured foliations along windows provided the total weights agree on the windows to be glued (taking unions of brane labels when combining distinguished points). It is shown in [49] that these geometric operations descend to the level of suitable chains on the spaces $Arc(F,\beta)$ and finally to the level of the integral homology groups of $Arc(F,\beta)$. These algebraic operations on homology satisfy the expected "operadic" equations of open/closed string theory, and new equations can be discovered as well.

In effect, [49] gives the string field theory of a single point, and natural questions are already discussed in detail in [49] including the "passage to conformal field theory" which seems to be provided by corresponding operations not for exhaustive arc families but rather for quasi filling arc families; an obvious challenge is to perform meaningful calculations in CFT. Calculate the homology groups of exhaustive or quasi filling arc families in brane labeled windowed surfaces, and calculate the homology groups of their combinatorial or DM-type compactifications (the former problem already articulated as our Problem 6 and the latter surely also posed elsewhere in this volume). Organize and understand the many algebraic relations of open/closed strings on the level of homology. Find the BRST operator. Introduce excited strings, i.e., extend to string field theory of realistic targets. Is the Torelli structure of any physical significance?

9. Punctured solenoid

A problem well-known in the school around Dennis Sullivan is:

Problem 15 [Ehrenpreis Conjecture] Given two closed Riemann surfaces, there are finite unbranched covers with homeomorphic total spaces which are arbitrarily close in the

Teichmüller metric.

This section describes joint work with Dragomir Šarić on related universal constructions in Teichmüller theory [47].

As a tool for understanding dynamics and geometry, Sullivan defined the hyperbolic solenoid [33] as the inverse limit of the system of finite-sheeted unbranched pointed covers of any fixed closed oriented surface of negative Euler characteristic. Following Ahlfors-Bers theory, he developed its Teichmüller theory and studied the natural dynamics and geometry, in particular introducing two principal mapping class groups, the continuous "full mapping class group" and the countable "baseleaf preserving mapping class group". He furthermore showed that the Ehrenpreis Conjecture is equivalent to the conjecture that the latter group has dense orbits in the Teichmüller space of the solenoid. [55] contains many basic results and open problems about Sullivan's solenoid.

Following this general paradigm, one defines the punctured hyperbolic solenoid S as the inverse limit over all finite-index subgroups Γ of $PSL_2(\mathbf{Z})$ of the system of covers \mathcal{U}/Γ over the modular curve $\mathcal{U}/PSL_2(\mathbf{Z})$, where \mathcal{U} denotes the upper half-space. In effect, branching is now permitted but only at the missing punctures covering the three orbifold points of the modular curve.

Let us build a particular model space homeomorphic to the punctured solenoid. Take $\Gamma = PSL_2(\mathbf{Z})$ acting by fractional linear transformations on \mathcal{U} , and let $\hat{\Gamma}$ denote its profinite compleion. Thus, $\gamma \in \Gamma$ acts naturally on $\mathcal{U} \times \hat{\Gamma}$ by $\gamma \cdot (z, t) = (\gamma z, t\gamma^{-1})$, and the quotient is homeomorphic to S.

Truly the entire decorated Teichmüller theory of a punctured surface of finite type [27] extends appropriately to S: there are global coordinates, there is an ideal cell decomposition of the decorated Teichmüller space, and there is an explicit non-degenerate two-form, the latter of which are invariant under the action of the baseleaf preserving subgroup $Mod_{BLP}(S)$ of the full mapping class group Mod(S); generators for $Mod_{BLP}(S)$ are provided by appropriate equivariant Whitehead moves, and a complete set of relations has recently been derived as well in further recent work with Dragomir Šarić and Sylvain Bonnot.

There are a number of standard questions (again see [55]): Is $Mod_{BLP}(S)$ finitely generated? Do the "mapping class like" elements generate $Mod_{BLP}(S)$? It seems to be a deep question which tesselations of the disk, other than the obvious so-called "TLC" ones, arise from the convex hull construction applied to the decorated solenoid in [47]. Is there any relationship between Mod(S) and the absolute Galois group? (This cuts both ways since only partial information is known about either group; notice that the action of the latter on $\hat{\Gamma}$ is explicit as a subgroup of the Grothendieck-Teichmüller group [51].)

Bibliography

[1] Nils Andreas Baas, "Bordism theories with singularities", Proceedings of the Advanced Study Institute on Algebraic Topology (Aarhus Univ., Aarhus, 1970), Vol. I, pp. 1-16. Various Publ. Ser., No. 13, Mat. Inst., Aarhus Univ., Aarhus, 1970.

[2] Max Bauer, "An upper bound for the least dilatation", *Trans. Amer. Math. Soc.* **330** (1992), 361-370.

[3] Moira Chas and Dennis P. Sullivan, "String topology", to appear Ann. Math., preprint math.GT/9911159,

[4] —, "Closed string operators in topology leading to Lie bialgebras and higher string algebra", The legacy of Niels Henrik Abel, Springer, Berlin (2004), 771-784, Preprint math.GT/0212358.

[5] L. Chekhov and V. Fock, "A quantum Techmüller space", *Theor. Math. Phys.*, **120** (1999) 1245-1259; "Quantum mapping class group, pentagon relation, and geodesics" *Proc. Steklov Math. Inst.* **226** (1999) 149-163.

[6] Leonid Chekhov and R. C. Penner, "On the quantization of Teichmüller and Thurston theories", to appear: Handbook of Teichmüller theory, European Math Society, ed. A. Papadopoulos, math.AG/0403247.

[7] Ralph L. Cohen and John D.S. Jones "A homotopy theoretic realization of string topology", *Math. Ann.* **324** (2002), 773-798.

[8] Max Dehn, Lecture notes from Breslau, 1922, Archives of the University of Texas at Austin.

[9] A. Fathi, F. Laudenbach, V. Poenaru, "Travaux de Thurston sur les Surfaces", Asterisques **66-67** (1979), Sem. Orsay, Soc. Math. de France.

[10] Tim Gendron, "The Ehrenpreis conjecture and the moduli-rigidity gap", Complex manifolds and hyperbolic geometry (Guanajuato, 2001), *Contemp. Math.* **311**, 207-229.

[11] Andrew Haas and Perry Susskind, "The connectivity of multicurves determined by integral weight train tracks", *Trans. Amer. Math. Soc.* **329** (1992), 637-652.

[12] John L. Harer, "The virtual cohomological dimension of the mapping class group of an orientable surface", *Invent. Math.* **84** (1986), 157-176.

[13] A. Hatcher and W. Thurston, "A presentation for the mapping class group of a closed orientble surface", *Topology* **19** (1980), 221-137.

[14] William J. Harvey, "Boundary structure of the modular group", Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud.97, Princeton Univ. Press, Princeton, N.J., (1981), 245-251.

[15] John H. Hubbard and Howard Masur, "Quadratic differentials and foliations", Acta

Math. 142 (1979), 221-274.

[16] Kyoshi Igusa, "Combinatorial Miller-Morita-Mumford classes and Witten cycles", *Algebr. Geom. Topol.* **4** (2004), 473-520.

[17] Rinat M. Kashaev, "Quantization of Teichmüller spaces and the quantum dilogarithm", *Lett. Math. Phys.*, **43**, (1998), 105-115; q-alg/9705021.

[18] Ralph Kaufman, Muriel Livernet, R. C. Penner, "Arc operads and arc algebras", *Geom. Topol.* 7 (2003), 511-568.

[19] Maxim Kontsevich, "Intersection theory on the moduli space of curves and the matrix Airy function", *Comm. Math. Phys.* **147** (1992), 1-23.

[20] W. B. R. Lickorish, "A finite set of generators for the homeotopy group of a twomanifold" *Proc. Camb. Phil. Soc.* **60** (1964), 769-778; Corrigendum, *Proc. Camb. Phil. Soc.* **62** (1966), 679-681.

[21] Yuri I. Manin, "Von Zahlen und Figuren", Talk at the International Conference "Gomtrie au vingtime cicle: 1930–2000", Paris, Institut Henri Poincar, Sept. 2001 27 pp., preprint math.AG/0201005.

[22] Gabriel Mondello, "Combinatorial classes on $\overline{M}_{g,n}$ are tautological", Int. Math. Res. Not. (2004), 2329-2390.

[23] R. C. Penner, "The action of the mapping class group on isotopy classes of curves and arcs in surfaces", thesis, Massachusetts Institute of Technology (1982), 180 pages.

[24] —, "The action of the mapping class group on curves in surfaces", *L'Enseignement Mathematique* **30** (1984), 39-55.

[25] —, "A construction of pseudo-Anosov homeomorphisms", *Proceedings of the American Math Society* **104** (1988), 1-19.

[26] —, "Bounds on least dilatations", Trans. Amer. Math. Soc. 113 (1991), 443-450.

[27] —, "The decorated Teichmüller space of punctured surfaces", *Communications in Mathematical Physics* **113** (1987), 299-339.

[28] —, "Perturbative series and the moduli space of Riemann surfaces", *Journal of Differential Geometry* **27** (1988), 35-53.

[29] , "An arithmetic problem in surface geometry", *The Moduli Space of Curves*, Birkhäuser (1995), eds. R. Dijgraaf, C. Faber, G. van der Geer, 427-466.

[30] —, "The simplicial compactification of Riemann's moduli space", Proceedings of the 37th Taniguchi Symposium, World Scientific (1996), 237-252.

[31] —, "Decorated Teichmüller space of bordered surfaces", *Communications in Analysis and Geometry* **12** (2004), 793-820, math.GT/0210326.

[32] R. C. Penner with John L. Harer *Combinatorics of Train Tracks*, Annals of Mathematical Studies **125**, Princeton Univ. Press (1992); second printing (2001).

[33] Dennis Sullivan, Linking the universalities of Milnor-Thurston, Feigenbaum and Ahlfors-Bers, Milnor Festschrift, Topological methods in modern mathematics (L. Goldberg and A. Phillips, eds.), Publish or Perish, 1993, 543-563. preprint.

[34] —, "Combinatorial invariants of analytic spaces", Proc. of Liverpool Singularities– Symposium, I (1969/70) Springer, Berlin (1971), 165-168.

[35] Joerg Teschner "An analog of a modular functor from quantized Teichmller theory", to appear Handbook of Teichmüller theory, European Math Society (ed. A. Papadopoulos), math.QA/0405332.

[36] Kurt Strebel, Quadratic Differentials, *Ergebnisse der Math. und ihrer Grenzgebiete*, Springer-Verlag, Berlin (1984).

[37] William P. Thurston, "On the geometry and dynamics of diffeomorphisms of surfaces", Bull. Amer. Math. Soc., **19** (1988) 417–431.

[38] —, "Three dimensional manifolds, Kleinian groups and hyperbolic geometry", Bull. Amer. Math. Soc. 6(1986), 357-381.

[39] Albert Fathi, "Dmonstration d'un thorme de Penner sur la composition des twists de Dehn", *Bull. Soc. Math. France* **120** (1992), 467-484.

[40] Anatol N. Kirillov, "Introduction to tropical combinatorics", Physics and combinatorics, 2000 (Nagoya), 82-150, World Sci. Publishing, River Edge, NJ, 2001.

[41] Jürg Richter-Gebert, Bernd Sturmfels, Thorsten Theobald, "First steps in tropical geometry", to appear in Proc. of COnf. on Idempotent Mathematicals and Mathematical Physics, *Contemp. Math.*, math.AG/0306366.

[42] Lee Mosher, "The classification of pseudo-Anosovs" in Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984), 13-75, London Math. Soc. Lecture Note Ser., **112**, Cambridge Univ. Press, Cambridge, 1986.

[43] William A. Veech, "Dynamics over Teichmller space", *Bull. Amer. Math. Soc.* 14 (1986), 103-106.

[44] Howard Masur, "Ergodic actions of the mapping class group", *Proc. Amer. Math. Soc* **94** (1985), 455-459.

[45] Athanase Papadopoulos and R. C. Penner, "A construction of pseudo-Anosov homeomorphisms", *Proc. Amer. Math Soc.* **104** (1988), 1-19.

[46] R. C. Penner, "The structure and singularities of arc complexes", preprint (2004) math.GT/0410603.

[47] R. C. Penner and Dragomir Šarić, "Teichmüller theory of the punctured solenoid",

preprint (2005), math.DS/0508476.

[48] S. Morita and R. C. Penner, "Torelli groups, extended Johnson homomorphisms, and new cycles on the moduli space of curves", preprint (2006), math.GT/0602461.

[49] Ralph M. Kaufmann and R. C. Penner, "Closed/open string diagrammatics" (2006), to appear in *Nucl. Phys B*, math.GT/0603485.

[50] Curt McMullen, "Billiards and Teichmüller curves on Hilbert modular surfaces", J. Amer. Math. Soc. 16 (2003), 857885.

[51] Leila Schneps, "The Grothendieck-Teichmüller group \widehat{GT} : a survey", Geometric Galois Actions (L. Schneps and P. Lochak, eds.) London Math Society Lecture Notes **242** (1997), 183-203.

[52] Shigeyuki Morita, "Abelian quotients of subgroups of the mapping class group of surfaces", *Duke Math. J.* **70** (1993), 699–726.

[53] Shigeyuki Morita, "Casson's invariant for homology 3-spheres and characteristic classes of surface bundles I" *Topology* **28** (1989), 305-323.

[54] Yuri I. Manin, "Real multiplication and noncommutative geometry", *The legacy of Niels Henrik Abel*, eds. O. A. Laudal and R. Piene, Springer Verlag, Berlin 2004, 685-727, preprint math.AG/0202109.

[55] Chris Odden, "The baseleaf preserving mapping class group of the universal hyperbolic solenoid", *Trans. A.M.S.* **357** (2004), 1829-1858.

[56] John L. Harer and Don Zagier, "The Euler characteristic of the moduli space of curves", *Invent. Math.* **85** (1986), 457-485.

[57] Claude Itzykson and Jean-Michel Drouffe, Statistical Field Theory, volume 1, Cambridge Monographs on Mathematical Physics, Cambridge Univ. Press (1989).