# DYNKIN DIAGRAM SEQUENCES AND STABILIZATION PHENOMENA

SANKARAN VISWANATH

ABSTRACT. We continue the study of stabilization phenomena for Dynkin diagram sequences initiated in the earlier work of Kleber and the present author. We consider a more general class of sequences than that of this earlier work, and isolate a condition on the weights that gives stabilization of tensor product and branching multiplicities. We show that all the results of the previous article can be naturally generalized to this setting. We also prove some properties of the partially ordered set of dominant weights of indefinite Kac-Moody algebras, and use this to give a more concrete definition of a stable representation ring. Finally, we consider the classical sequences  $B_n, C_n, D_n$  that fall outside the purview of the earlier work, and work out some easy-to-describe conditions on the weights which imply stabilization.

### 1. INTRODUCTION

In this article, we will consider sequences of Dynkin diagrams and study the behavior of representations of the associated Kac-Moody algebras. The sequences of Dynkin diagrams considered are of the form

$$Z_k := \qquad \begin{pmatrix} X_1 & & & \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & &$$

where  $X_1$  and  $X_2$  are fixed Dynkin diagrams and the string of intermediate nodes has length k. The article [4] considered the Dynkin diagram sequences  $Z_k$  arising in the special case when  $X_2 = A_1$  (the diagram with just a single node).

For most choices of  $X_1, X_2$ , the associated Kac-Moody algebra  $\mathfrak{g}(Z_k)$  is infinite dimensional, non-affine (i.e of *indefinite* type) and very little is known about such Lie algebras in general. So, rather than study representations of the individual  $\mathfrak{g}(Z_k)$ , we study them in the limit as  $k \to \infty$ .

We consider dominant integral weights  $\lambda, \mu, \nu$  which are *supported* on the two ends of the Dynkin diagram of  $Z_k$ . In [4], the primary object of interest was the multiplicity  $c_{\lambda\mu}^{\nu}(k)$  of the irreducible highest weight representation  $L(\nu^{(k)})$  of  $\mathfrak{g}(Z_k)$  in the tensor product  $L(\lambda^{(k)}) \otimes L(\mu^{(k)})$ . Specifically under some additional conditions, it was shown that these multiplicities become constant (*stabilize*) for large k; this generalizes the classical stabilization results for the  $A_n$  diagram which more or less follow from the Littlewood-Richardson rule.

In this article, we consider the tensor product multiplicity  $c_{\lambda\mu}^{\nu}(k)$  as well as the branching multiplicities  $b_{\lambda\beta}(k)$ . The question we ask is this :

<sup>1991</sup> Mathematics Subject Classification. 17B67.

Key words and phrases. Tensor product multiplicity, branching multiplicity, Littelmann path model, partial order on dominant weights.

Under what conditions on the weights  $\lambda, \mu, \nu, \beta$  do  $c^{\nu}_{\lambda\mu}(k)$  and  $b_{\lambda\beta}(k)$  stabilize ?

In section 2 we provide a sufficient condition that ensures stabilization. Having formulated this criterion for stabilization, we follow two distinct threads:

First, we recall that the goal of the previous article [4] was slightly different; it sought to find conditions on the diagram  $X_1$  ( $X_2 = A_1$  there) which would ensure stabilization of  $c_{\lambda\mu}^{\nu}(k)$  for all weights  $\lambda, \mu, \nu$ . The condition on  $X_1$  which made this work was called *extensibility*. In sections 3 and 4, we generalize this approach and define a notion of extensibility for pairs ( $X_1, X_2$ ). This new notion is strong enough to ensure stabilization for all  $\lambda, \mu, \nu, \beta$  while still admitting enough interesting examples of diagrams. In particular, this notion subsumes the earlier notion of [4].

It was also shown in [4] that one could use the stable values of the  $c_{\lambda\mu}^{\nu}(k)$  to define an operation \*, which mimics the limit as  $k \to \infty$  of the tensor product. A very surprising fact discovered there was the associativity of \*. A notion of a stable representation ring was formulated as a consequence. We derive all these results for extensible pairs  $(X_1, X_2)$  and define a more concrete, modified version of the stable representation ring in this case. This appears in section 5.

As our second thread, we turn to the classical sequences of Dynkin diagrams  $B_n, C_n, D_n$ . These were notable exceptions to the extensibility condition of [4]. So, while nice stabilization results hold for the  $A_n$ , nothing much could be said about these other classical types. We remedy this in section 6. For these types, our methods do not imply stabilization for all choices of  $\lambda, \mu, \nu, \beta$ , but we work out some easy to describe conditions on the weights under which they do.

While this article was in preparation, Webster [9] has shown that a more general version of our stabilization result (Theorem 1) can be proved using quiver varieties and their connections with representations of Kac-Moody algebras. The results and formulation in [9] and in section 2 of the present article overlap substantially. Webster's approach also proves a 'polynomiality of weight multiplicities' result for these kinds of Dynkin diagram sequences.

Acknowledgements: I'd like to thank Michael Kleber for his constant encouragement and many valuable suggestions on an earlier draft of this article. I'd also like to thank Ben Webster for explaining many aspects of his approach to me.

## 2. A CRITERION FOR STABILIZATION

2.1. Notations. We begin with some notations concerning Kac-Moody algebras and Dynkin diagrams. Let X be a Dynkin diagram (in the sense of [3, Chapter 4]) associated to a symmetrizable generalized Cartan matrix C(X). Using the data of X, one constructs  $\mathfrak{g}(X)$ , the Kac-Moody algebra associated to X. Let  $\mathfrak{h}(X)$  denote the Cartan subalgebra of  $\mathfrak{g}(X)$ , with dual  $\mathfrak{h}^*(X)$ . We let N(X) denote the set of nodes of X and det  $X := \det C(X)$ . By abuse of notation we will usually write  $p \in X$  to mean that  $p \in N(X)$ . For each  $p \in X$ , let  $\check{\alpha}_p, \alpha_p, \omega_p$  respectively denote the simple coroot, the simple root and the fundamental weight corresponding to p. So for example  $\omega_p(\check{\alpha}_q) = \delta_{pq}$  for  $p, q \in X$ .

We let Q(X), P(X) be the root and weight lattices of  $\mathfrak{g}(X)$ . The set  $\{\alpha_p, p \in X\}$ forms a  $\mathbb{Z}$  basis of Q(X), and when det  $X \neq 0$  the set  $\{\omega_p, p \in X\}$  forms a  $\mathbb{Z}$  basis of P(X). Let  $Q^+(X)$  be the set of  $\mathbb{Z}^{\geq 0}$  linear combinations of the  $\alpha_p$  and  $P^+(X)$ denote the set of  $\mathbb{Z}^{\geq 0}$  linear combinations of the  $\omega_p$ . When det X = 0, the fundamental weights are not uniquely defined and we pick one of the possible  $\omega_p$ 's for each node p. We will usually not run into diagrams with det X = 0 (see section 6.3 for an exception).

2.2. Dynkin diagram sequences. In this article, we will be interested in *sequences* of Dynkin diagrams; for example, we have the classical Dynkin diagram sequence  $A_k$ ,  $k \ge 1$ 

$$\begin{array}{c|c} \bullet & \bullet \\ 1 & 2 & 3 \end{array} \cdots \begin{array}{c} \bullet \\ k \end{array}$$

We will let  $A_0$  denote the empty diagram.

By a marked Dynkin diagram, we will mean the data  $(X,\xi)$  where X is a Dynkin diagram and  $\xi \in X$  is a distinguished node.

Given a marked Dynkin diagram  $(X, \xi)$  and an integer  $m \ge 0$ , one can construct the Dynkin diagram X(m) obtained by "attaching" the diagram  $A_m (m \ge 0)$  to the node  $\xi$  as follows:



In the notation of [4], this is  $X_{d+m}$  where d is the number of nodes in X. We let  $\xi(m)$  denote the end node (labeled m in the figure) and consider  $(X(m), \xi(m))$  as a marked Dynkin diagram.

One can consider the sequence of symmetrizable Kac-Moody algebras  $\mathfrak{g}(X(m))$ associated with the X(m). For most choices of X and for most values of m, these turn out to be infinite dimensional non affine Kac-Moody algebras (i.e of *indefinite type*), but one can still study their integrable highest weight representations. The objective of [4] was to study how multiplicities in tensor product decompositions of such representations of  $\mathfrak{g}(X(m))$  change with m.

2.3. **Pairs.** We now consider a broader class of sequences of Dynkin diagrams; these will be obtained starting with *pairs*  $(X_1, X_2)$  of marked Dynkin diagrams, rather than with a single diagram X.

For i = 1, 2 let  $(X_i, \xi_i)$  be given marked Dynkin diagrams such that  $C(X_i)$  are symmetrizable. For each  $k \ge 1$  we form a Dynkin diagram  $Z_k = Z_k(X_1, X_2)$  by taking the diagram  $A_k$  and attaching its two ends to  $\xi_1$  and  $\xi_2$  as shown in figure



The figure doesn't show the rest of the nodes of  $X_1$  and  $X_2$ . The matrix  $C(Z_k)$  is clearly symmetrizable. The associated Kac-Moody algebra will be denoted  $\mathfrak{g}(Z_k)$ . If  $X_2 = A_1$ , the diagram with a single node, this construction coincides with the earlier one;  $Z_k = X_1(k+1)$ .

In the rest of section 2, we will often need to refer to the following subdiagrams of the  $Z_k$ 's.

- (1)  $X_{12} := X_1 \cup X_2$ .
- (2) For  $l \ge 0$ , we identify  $X_1(l)$  with the subdiagram of  $Z_k$  formed by  $X_1$  and the intermediate nodes labeled  $1, \dots, l$  in the above figure.

(3) Similarly given  $r \ge 0$ , identify  $X_2(r)$  with the subdiagram formed by  $X_2$  and the intermediate nodes labeled  $k, k-1, \dots, k-r+1$ .

Each of these is a subset of  $Z_k$  for all but finitely many values of k. When these subsets are encountered, the specific value of k being used will be clear from context.

(4) With l, r as above, let  $Y_k(l, r) \subset Z_k$  denote the subdiagram isomorphic to  $A_{k-l-r}$  formed by the nodes numbered  $l+1, l+2, \dots, k-r$ . So for instance,  $Y_k(0,0)$  is the diagram  $A_k$  in the middle.

2.4. **Representations of**  $\mathfrak{g}(Z_k)$ . We now consider integrable highest weight representations of the Kac-Moody algebra  $\mathfrak{g}(Z_k)$ . These are indexed by dominant integral weights of  $\mathfrak{g}(Z_k)$ . Our immediate interest will be in dominant integral weights which are "supported" on  $X_1 \cup X_2$ . To make this more precise :

Let  $W(X_1, X_2)$  be the set of all functions  $f : N(X_1) \cup N(X_2) \to \mathbb{Z}$  and  $W^+(X_1, X_2) := \{f \in W(X_1, X_2) : image(f) \subset \mathbb{Z}^{\geq 0}\}$ . Similarly for each  $s \in \mathbb{Z}$ , let  $R_s(X_1, X_2)$  be the set of all functions  $f : N(X_1) \cup N(X_2) \to \mathbb{Z}$  such that  $f(\xi_1) = f(\xi_2) = s$  and  $R_s^+(X_1, X_2) := \{f \in R_s(X_1, X_2) : image(f) \subset \mathbb{Z}^{\geq 0}\}$ . We will let the elements of  $W(X_1, X_2)$  and  $R_s(X_1, X_2)$  define elements of the weight and root lattices of  $\mathfrak{g}(Z_k)$  as follows: given  $\lambda \in W(X_1, X_2)$ , define  $\lambda^{(k)} \in P(Z_k)$  by

$$\lambda^{(k)} := \sum_{p \in X_{12} \subset Z_k} \lambda(p) \, \omega_p$$

We define  $P_0(Z_k) := \{ \mu \in P(Z_k) : \mu(\check{\alpha}_p) = 0 \ \forall p \in Y_k(0,0) \}$  and  $P_0^+(Z_k) := P_0(Z_k) \cap P^+(Z_k)$ . These are the weights that are supported on  $X_1 \cup X_2$ . It is clear that  $\lambda^{(k)} \in P_0(Z_k)$  and that  $\lambda \in W^+(X_1, X_2) \Leftrightarrow \lambda^{(k)} \in P_0^+(Z_k)$ .

Similarly given  $\gamma \in R_s(X_1, X_2)$  define

$$\gamma^{(k)} := \sum_{p \in X_{12} \subset Z_k} \gamma(p) \, \alpha_p + s \sum_{p \in Y_k(0,0)} \alpha_p$$

It is easily seen that (i)  $\gamma^{(k)} \in Q(Z_k)$  (ii)  $\gamma^{(k)} \in Q^+(Z_k) \Leftrightarrow \gamma \in R_s^+(X_1, X_2)$  (iii)  $\gamma^{(k)} \in P_0(Z_k)$  since  $\gamma(\xi_1) = \gamma(\xi_2) = s$ .

We now consider two important representation theoretic notions:

Given  $\lambda \in W^+(X_1, X_2)$ , it defines an integrable highest weight representation  $L(\lambda^{(k)})$  of  $\mathfrak{g}(Z_k)$  for each  $k \geq 1$ . Assume  $\mu, \nu \in W^+(X_1, X_2)$ . We can consider the tensor product  $L(\lambda^{(k)}) \otimes L(\mu^{(k)})$ ; this is an integrable representation in category  $\mathcal{O}$  and hence decomposes into a direct sum of integrable highest weight representations. This is usually an infinite direct sum, but each direct summand occurs with finite multiplicity. We let  $c_{\lambda\mu}^{\nu}(k)$  be the multiplicity of the representation  $L(\nu^{(k)})$  in the decomposition of  $L(\lambda^{(k)}) \otimes L(\mu^{(k)})$ .

Similarly given  $\beta \in W(X_1, X_2)$ , we consider the branching multiplicity of the weight  $\beta^{(k)}$  for the action of the  $\mathfrak{g}(A_k)$  corresponding to the subdiagram  $A_k$  in the middle. This number, denoted  $b_{\lambda\beta}(k)$  is defined to be the dimension of the space  $\{v \in L(\lambda^{(k)})_{\beta^{(k)}} : \mathfrak{n}^+(A_k) v = 0\}$  i.e vectors of weight  $\beta^{(k)}$  annihilated by the positive root spaces of  $\mathfrak{g}(A_k)$ .

Both the tensor product and branching multiplicities are functions of k. If  $f : \mathbb{Z}^{>0} \to \mathbb{Z}$  is a function, we say that f stabilizes if there exists K such that f(k) =

f(k') for all  $k, k' \ge K$ . In this case, we set  $f(\infty) := f(K)$ . We will be interested in conditions under which the  $c_{\lambda\mu}^{\nu}(k)$  and  $b_{\lambda\beta}(k)$  stabilize.

**Definition 1.** Given  $\lambda, \mu \in W(X_1, X_2)$ , let  $\lambda \sim \mu$  if  $\lambda^{(k)} - \mu^{(k)} \in Q(Z_k)$  for infinitely many values of k.

**Remark 1.** It is clear that ~ is an equivalence relation. Further if  $(\lambda + \mu) \nsim \nu$ , then  $c_{\lambda\mu}^{\nu}(k)$  clearly stabilizes with  $c_{\lambda\mu}^{\nu}(\infty) = 0$ . Similarly  $\lambda \nsim \beta$  implies  $b_{\lambda\beta}(k)$ stabilizes with  $b_{\lambda\beta}(\infty) = 0$ .

2.5. **Example: the pair**  $(X(m), A_n)$ . Let  $(X, \xi)$  be a given marked Dynkin diagram. Fix  $m \ge 0$  and set  $X_1 = X(m)$  with  $\xi(m)$  distinguished. Fix  $n \ge 0$  and take  $X_2 = A_n$  with the end node being distinguished. Then  $Z_k = X(m+n+k)$ . This is the configuration considered in [4]. Weights of the form  $\lambda^{(k)}$  for  $\lambda \in W^+(X_1, X_2)$  were called *double headed weights* there. Under a further hypothesis on X (the so called *extensibility condition* [4, definition 2.4], see also section 3.1 below), [4] obtained a stabilization result for tensor product multiplicities  $c_{\lambda\mu}^{\nu}(k)$ . We state this and a related result in our present notation.

**Proposition 1.** Let X be an extensible marked Dynkin diagram. For  $m, n \ge 0$ , let  $X_1 := X(m), X_2 := A_n$ , and  $Z_k := Z_k(X_1, X_2)$ . Take  $\eta, \delta \in W(X_1, X_2)$  such that  $\eta \sim \delta$ . Then there exists  $s \in \mathbb{Z}$  and  $\gamma \in R_s(X_1, X_2)$  such that

(2.1) 
$$\eta^{(k)} - \delta^{(k)} = \gamma^{(k)} \quad \forall k \ge 1$$

This is essentially proposition 4.1 of [4]. It is here that weights of the form  $\gamma^{(k)}$ ,  $\gamma \in R_s(X_1, X_2)$  appear naturally.

**Proposition 2.** Let X be an extensible marked Dynkin diagram. For  $m, n \ge 0$ , let  $X_1 := X(m), X_2 := A_n$ , and  $Z_k := Z_k(X_1, X_2)$ . Suppose  $\lambda, \mu, \nu \in W^+(X_1, X_2)$  are such that there exists  $s \in \mathbb{Z}$  and  $\gamma \in R_s(X_1, X_2)$  such that

(2.2) 
$$\lambda^{(k)} + \mu^{(k)} - \nu^{(k)} = \gamma^{(k)} \quad \forall k \ge 1$$

Then for all k, k' > 2s,  $c_{\lambda\mu}^{\nu}(k) = c_{\lambda\mu}^{\nu}(k')$ . Thus  $c_{\lambda\mu}^{\nu}(k)$  stabilizes.

Observe that if  $\gamma \notin R_s^+(X_1, X_2)$ , then  $\gamma^{(k)} \notin Q^+(Z_k)$  for all  $k \ge 1$ . This implies that  $c_{\lambda\mu}^{\nu}(k) = 0 \ \forall k \ge 1$ . So we may as well assume  $s \ge 0$  and  $\gamma \in R_s^+(X_1, X_2)$ . The proof that  $c_{\lambda\mu}^{\nu}(k)$  stabilizes, crucially depends on the fact that  $\lambda^{(k)} + \mu^{(k)} - \nu^{(k)}$  has this specific form with  $\gamma \in R_s^+(X_1, X_2)$ . This proof appears as part of the proof of theorem 4.5 of [4] in sections 4.3-4.4 there.

2.6. Main theorem. We now extract the crux of the argument in [4] that proves Proposition 1 and formulate a more general theorem concerning tensor product as well as branching multiplicities. This works for arbitrary pairs of marked Dynkin diagrams, but we impose a condition on our weights that is analogous to equation (2.2).

**Theorem 1.** Let  $X_1, X_2$  be arbitrary marked Dynkin diagrams and  $\lambda, \mu, \nu \in W^+(X_1, X_2)$ ,  $\beta \in W(X_1, X_2)$ .

- (1) Suppose for some  $s \ge 0$ ,  $\exists \gamma \in R_s^+(X_1, X_2)$  with  $\lambda^{(k)} + \mu^{(k)} \nu^{(k)} = \gamma^{(k)}$ for all  $k \ge 1$ . Then for all k, k' > 2s,  $c_{\lambda\mu}^{\nu}(k) = c_{\lambda\mu}^{\nu}(k')$ .
- (2) Suppose for some  $s \ge 0$ ,  $\exists \gamma \in R_s^+(X_1, X_2)$  such that  $\lambda^{(k)} \beta^{(k)} = \gamma^{(k)}$  for all  $k \ge 1$ , then for all k, k' > 2s,  $b_{\lambda\beta}(k) = b_{\lambda\beta}(k')$ .

As remarked above, the essential ideas of this proof are the similar to those of proposition 2; We summarize the main steps below (mostly referring to [4] for the proofs). We also deduce the statement regarding the  $b_{\lambda\beta}$  which does not appear in [4].

2.7. Littelmann path model. As a first step we use the explicit combinatorial description of  $b_{\lambda\beta}(k)$  and  $c_{\lambda\mu}^{\nu}(k)$  given by Littelmann's path model. For  $k \geq 1$ , we let  $\Pi^{(k)}$  denote the set of piecewise linear paths  $\pi : [0,1] \to \mathfrak{h}^*(Z_k)$  such that  $\pi(0) = 0$ . To each node  $p \in N(Z_k)$ , we associate raising and lowering operators  $e_p$ ,  $f_p$  on  $\mathbb{Z}\Pi^{(k)}$  defined as follows: let  $\pi \in \Pi^{(k)}$  and  $\pi_p(t) := \pi(t)(\check{\alpha}_p^{(n)})$  for  $0 \leq t \leq 1$ . We consider the function  $a : [0,1] \to [0,1]$  defined by  $a(t) = \min\{1, \pi_p(s) - m_p | t \leq s \leq 1\}$ , where  $m_p = \min\{\pi_p(t) | 0 \leq t \leq 1\}$ . Note that a is an increasing function. If a(1) < 1,  $f_p\pi := 0$ . Otherwise,  $f_p\pi$  is the path defined by

(2.3) 
$$f_p \pi(t) := \pi(t) - a(t)\alpha_p$$

Similarly we consider the increasing function  $b : [0,1] \to [0,1]$  with  $b(t) = \max\{0, 1 - (\pi_p(s) - m_p) | 0 \le s \le t\}$ . If b(0) > 0, we set  $e_p \pi = 0$ ; otherwise

$$(2.4) e_p \pi(t) := \pi(t) + b(t)\alpha_p$$

See Littelmann's papers [5], [7], [6] for a more pictorial description of these operators, and Stembridge[8] for an axiomatic formulation.

Let  $\pi_{\lambda^{(k)}}(t) := t\lambda^{(k)}$  be the straight line path with  $\pi_{\lambda^{(k)}}(1) = \lambda^{(k)}$ . Paths that are obtained by repeated action of the  $f_p$ ,  $p \in Z_k$  on  $\pi_{\lambda^{(k)}}$  are called Lakshmibai-Seshadri (L-S) paths of shape  $\lambda^{(k)}$ . Consider the following sets:

$$\mathcal{P}_{k}(\lambda,\beta) := \{ \text{ L-S paths } \pi \text{ of shape } \lambda^{(k)} \text{ with } \pi(1) = \beta^{(k)} \}$$
$$\mathcal{P}_{k}^{+}(\lambda,\beta) := \{ \pi \in \mathcal{P}_{k}(\lambda,\beta) : e_{p}\pi = 0 \ \forall p \in Y_{k}(0,0) \}$$
$$\mathcal{P}_{k}^{(\mu)}(\lambda,\beta) := \{ \pi \in \mathcal{P}_{k}(\lambda,\beta) : \pi \text{ is } \mu^{(k)} \text{ dominant } \}$$

In the last equation,  $\mu^{(k)}$  dominance of  $\pi$  means that the shifted path  $\mu^{(k)} + \pi$  lies completely in the dominant Weyl chamber. We now have

## **Theorem 2.** (Littelmann [7])

(1)  $b_{\lambda\beta}(k) = \# \mathcal{P}_k^+(\lambda,\beta)$ (2)  $c_{\lambda\mu}^\nu(k) = \# \mathcal{P}_k^{(\mu)}(\lambda,\nu-\mu)$ 

2.8. **Proof of theorem 1.** We define an auxiliary set. For k > 2s let

$$\Sigma_k := \{ \pi \in \Pi^{(k)} : \pi(t)(\check{\alpha}_p) = 0 \ \forall t \in [0,1] \ \forall p \in Y_k(s,s) \}$$

These are all (not just L-S) paths that are supported on the complement of  $Y_k(s, s)$ . For all k, k' > 2s, we have bijections  $\phi_{kk'} : \Sigma_k \to \Sigma_{k'}$ ; given  $\pi \in \Sigma_k$ , there exist functions  $f_p(t)$  such that

(2.5) 
$$\pi(t) = \sum_{p \in X_1(s) \cup X_2(s)} f_p(t) \,\omega_p$$

We define  $\phi_{kk'}(\pi)$  by the same formula as on the right hand side except that we now interpret  $X_1(s)$  and  $X_2(s)$  as subdiagrams of  $Z_{k'}$ . It is clear that  $\phi_{kk'}$  and  $\phi_{k'k}$ are inverses of each other.

 $\mathbf{6}$ 

The following is the important proposition which details the relationship between these sets.

**Proposition 3.** Let k, k' > 2s. Then

(2.6) 
$$\mathcal{P}_k^{(\mu)}(\lambda,\beta) \subset \mathcal{P}_k^+(\lambda,\beta) \subset \Sigma_k$$

Further,  $\phi_{kk'}: \Sigma_k \to \Sigma_{k'}$  preserves these subsets i.e,

(2.7) 
$$\phi_{kk'}(\mathcal{P}_k^+(\lambda,\beta)) = \mathcal{P}_{k'}^+(\lambda,\beta)$$

(2.8) 
$$\phi_{kk'}(\mathcal{P}_k^{(\mu)}(\lambda,\beta)) = \mathcal{P}_{k'}^{(\mu)}(\lambda,\beta)$$

Proof: The first inclusion in (2.6) follows directly from the definition of  $\mu^{(k)}$ dominance and the fact that  $\mu^{(k)}(\check{\alpha}_p) = 0$  for all  $p \in Y_k(0,0)$ . The proof of the inclusion  $\mathcal{P}_k^+(\lambda,\beta) \subset \Sigma_k$  requires a careful argument with Littelmann paths; this appears in Section 4.4 of [4]. This latter argument also proves the following fact:

(2.9) 
$$\pi \in \mathcal{P}_k^+(\lambda,\beta) \Rightarrow \phi_{kk'}(\pi) \text{ is an L-S path of shape } \lambda^{(k')}$$

 $\sum_{\substack{p \in X_1(s) \cup X_2(s)}} \int_{f_p(t) \omega_p} f_p(t) \omega_p \text{ as in equation (2.5). We also write } \mu^{(k)} = 0.$  Then Given this fact (2.9) it is easy to prove (2.7) and (2.8): we let  $\pi \in \mathcal{P}_k^0(\lambda,\beta)$ with  $\pi(t) =$ 

$$\sum_{p \in X_{12}} o_p \, \omega_p$$
. Then

(2.10) 
$$\pi \in \mathcal{P}_{k}^{+}(\lambda,\beta) \Leftrightarrow f_{p}(t) \geq 0 \,\forall t \in [0,1] \,\forall p \in Y_{k}(0,0)$$
  
(2.11) 
$$\pi \in \mathcal{P}_{k}^{(\mu)}(\lambda,\beta) \Leftrightarrow \pi \in \mathcal{P}_{k}^{+}(\lambda,\beta) \text{ and } b_{p} + f_{p}(t) \geq 0 \,\forall t \in [0,1] \,\forall p \in X_{12}$$

These very same conditions ensure that  $\pi \in \mathcal{P}_{k'}^+(\lambda,\beta)$  or  $\pi \in \mathcal{P}_{k'}^{(\mu)}(\lambda,\beta)$  as the case may be. 

**Corollary 1.** From equation (2.7), we have  $b_{\lambda\beta}(k) = b_{\lambda\beta}(k')$ . Taking  $\beta = \nu - \mu$ , equation (2.8) shows  $c_{\lambda\mu}^{\nu}(k) = c_{\lambda\mu}^{\nu}(k')$  for all k, k' > 2s. This proves our main theorem 1. 

## 3. Extensible pairs

We briefly revisit the situation considered in section 2.5. Let X be an extensible marked Dynkin diagram,  $m, n \ge 0$  and set  $X_1 := X(m), X_2 = A_n$  so that  $Z_k(X_1, X_2) = X(m+n+k)$ . Given  $\lambda, \mu, \nu \in W^+(X_1, X_2)$ , we have: Case 1:  $(\lambda + \mu) \nsim \nu$ 

Remark 1 implies that  $c_{\lambda\mu}^{\nu}(k)$  stabilizes with  $c_{\lambda\mu}^{\nu}(\infty) = 0$ . Case 2:  $(\lambda + \mu) \sim \nu$ 

Proposition  $1 \Rightarrow \exists \gamma \in R_s(X_1, X_2)$  such that  $\lambda^{(k)} + \mu^{(k)} - \nu^{(k)} = \gamma^{(k)}$ . Proposition 2 then implies stabilization of  $c_{\lambda\mu}^{\nu}(k)$ . Thus  $c_{\lambda\mu}^{\nu}(k)$  stabilizes for **all** triples  $\lambda, \mu, \nu \in W^+(X_1, X_2)$ .

For arbitrary  $(X_1, X_2)$ , Remark 1 still applies. The conclusion of Proposition 1 however fails in general (see equation (6.1) for the  $B_n$ ). So, we can deduce stabilization of  $c_{\lambda\mu}^{\nu}(k)$  and  $b_{\lambda\beta}(k)$  only when  $\lambda, \mu, \nu, \beta$  have the special form of Theorem 1.

Our next goal will be to define a class of pairs  $(X_1, X_2)$  for which Proposition 1 holds. These will be called *extensible pairs*. They include the earlier situation as a special case; when X is an extensible diagram,  $(X(m), A_n)$  will turn out to be extensible pairs for  $m, n \ge 0$ . When  $(X_1, X_2)$  is an extensible pair, one again gets stabilization of  $c_{\lambda\mu}^{\nu}(k)$  and  $b_{\lambda\beta}(k)$  for all  $\lambda, \mu, \nu \in W^+(X_1, X_2)$ ,  $\beta \in W(X_1, X_2)$ .

Sections 3 - 5 below will be concerned exclusively with extensible pairs. We study their properties, define their associated *number of boxes* functions and finally use the stable values of  $c_{\lambda\mu}^{\nu}(k)$  to define a notion of a *stable representation ring*.

Among the classical types A, B, C, D, only type A falls within the class of extensible pairs. Readers interested in the B, C, D types may skip directly to section 6.

3.1. Review of results from [4]. First, we review some relevant notions from [4]. Given a marked Dynkin diagram  $(X,\xi)$  with d nodes, by a *numbering* of X we will mean a bijection  $\epsilon : N(X) \to \{1, 2, \dots, d\}$  such that  $\epsilon(\xi) = d$ . For the classical Dynkin diagram  $A_k, k \ge 1$  we fix the numbering  $\epsilon_A : N(A_k) \to \{1, 2, \dots, k\}$  shown below:

$$\begin{array}{c} \bullet & \bullet \\ 1 & 2 & 3 \end{array} \cdots \begin{array}{c} \bullet \\ k \end{array}$$

Given a numbering  $\epsilon$  of  $(X,\xi)$ , the diagram X(m) inherits a natural numbering j defined by  $j(p) = \epsilon(p)$  for  $p \in X \subset X(m)$  and  $j(p) = \epsilon_A(p) + d$  for  $p \in A_m \subset X(m)$ .

$$\begin{array}{c|c} \mathbf{X} & \bullet \\ \hline & d \\ d \\ d \\ d + 1 \\ d + 2 \\ \end{array} & \cdots \\ \hline & d \\ d \\ d \\ d \\ m \\ \end{array}$$

These numberings of X(m) are compatible for different *m*'s, i.e., for m' < m, the numbering *j* of X(m) restricted to  $X(m') \subset X(m)$  gives the numbering *j* of X(m').

Recall that for any Dynkin diagram X for which det  $X \neq 0$ , P(X)/Q(X) is a finite abelian group of order  $|\det X|$ . For any  $\eta \in P(X)$ , we let  $[\eta]$  denote its image in P(X)/Q(X).

Given a marked Dynkin diagram X define

 $\Delta_X := \det X - \det X(-1)$ 

where X(-1) is the diagram obtained from X by deleting the node  $\xi$  and all edges incident on it. We then have the following formula from Equation (2.2) of [4]:

(3.1) 
$$\det X(m) = \det X + m\Delta_X \quad \forall m \ge -1$$

If  $X = A_1$ , then X(-1) is empty; in this case we set det X(-1) := 1.

**Definition 2.** The marked Dynkin diagram X is *extensible* if det  $X \neq 0, \Delta_X \neq 0$  and  $gcd(det X, \Delta_X) = 1$ .

It was shown in [4] that the extensibility of X has many pleasant consequences. Notably, from Lemma (3.1) of [4]:

**Proposition 4.** Let X be extensible and suppose  $m \ge 0$  is such that det  $X(m) \ne 0$ . Then P(X(m))/Q(X(m)) is a cyclic group. Further if we let  $\bar{\omega}^{(m)}$  denote the fundamental weight corresponding to the end node of X(m) (i.e. p s.t j(p) = d+m), then  $[\bar{\omega}^{(m)}]$  generates this cyclic group.

More importantly, Proposition (3.3) of [4] gives

**Proposition 5.** Let X be extensible. There exists a unique sequence  $(a_i)_{i=1}^{\infty}$  of integers such that  $\forall m \geq 0$  with det  $X(m) \neq 0$  and  $\forall p \in X(m)$ , the relation

(3.2) 
$$-\Delta_X \,\omega_p \equiv a_{j(p)} \,\bar{\omega}^{(m)} \pmod{Q(X(m))}$$

holds in P(X(m)).

Recall that j is the node numbering introduced above. It is clear by proposition 4 that  $[-\Delta_X \omega_p]$  must be a multiple of  $[\bar{\omega}^{(m)}]$ ; the content of proposition 5 is that a single sequence  $(a_i)$  makes equation (3.2) hold for all values of m under consideration.

We will also need the following lemma which is essentially Equation (4.2) of [4]. Lemma 1. For fixed  $m \ge 0$ , and  $p \in X(m)$ , if we write

$$-\Delta_X \,\omega_p - a_{j(p)} \,\bar{\omega}^{(m)} = \sum_{q \in X(m)} c_{j(q)} \,\alpha_q$$

with  $c_i \in \mathbb{Z}$ , then  $c_{d+m} = -a_{j(p)}$ .

3.2. **Definition of extensible pairs.** The goal of this subsection is to generalize the notion of extensibility to pairs of marked Dynkin diagrams.

Let  $(X_1, X_2)$  be a given pair of marked Dynkin diagrams. We let  $\Delta_i := \Delta_{X_i} = \det X_i - \det X_i(-1)$ . We have

Lemma 2. For all 
$$k \ge 1$$

(3.3) 
$$\det Z_k = (k-1)\Delta_1 \Delta_2 + (\det X_1 \Delta_2 + \det X_2 \Delta_1)$$

*Proof:* We first prove this for k = 1.

Put  $\overline{X}_i := X_i(-1)$  for i = 1, 2. Now, the matrix  $C(Z_1)$  looks like

$$C(Z_1) = \begin{bmatrix} C(X_1) & & \\ & -1 & \\ \hline & -1 & 2 & -1 \\ \hline & & -1 & \\ \hline & & C(X_2) \end{bmatrix}$$

Expanding along the middle column we get

(3.4) 
$$\det Z_1 = 2 \det X_1 \det X_2 + \det P_1 + \det P_2$$

where  $P_1$  is a matrix of the form

$$P_1 = \begin{bmatrix} C(\bar{X}_1) & * & & \\ & * & & \\ \hline & -1 & -1 & \\ \hline & & & \\ \hline \end{array} \right]$$

The upper triangularity of  $P_1$  gives det  $P_1 = -\det \bar{X}_1 \det X_2$ . A similar calculation shows that  $P_2$  is a lower triangular matrix with det  $P_2 = -\det \bar{X}_2 \det X_1$ . Putting these back in Equation(3.4) completes the proof for k = 1.

If k > 1, then consider  $X_1(k-1)$ . We can think of  $Z_k = Z_k(X_1, X_2)$  as being the same thing as  $Z_1(X_1(k-1), X_2)$ . In terms of a picture:



Applying our k = 1 result gives

 $\det Z_k = \det X_1(k-1)\,\Delta_2 + \det X_2\,(\det X_1(k-1) - \det X_1(k-2))$ 

Lemma 2 implies det  $X_i(m) = \det X_i + m\Delta_i \quad \forall m \ge -1$ . Putting this back we get:

$$\det Z_k = \left(\det X_1 + (k-1)\Delta_1\right)\Delta_2 + \det X_2\Delta_1$$

proving our lemma.

**Definition 3.** A pair of marked Dynkin diagrams  $(X_1, X_2)$  is called an *extensible* pair if

- (1) Each  $X_i$  is an extensible diagram, i.e.,  $gcd(\det X_i, \Delta_i) = 1, \ \Delta_i \neq 0, det X_i \neq 0, i = 1, 2.$
- (2)  $\operatorname{gcd}(\Delta_1, \Delta_2) = 1.$

**Remark 2.** (1) Observe that  $A_1$  is an extensible Dynkin diagram since  $\Delta_{A_1} = 1$ . It is clear that X is extensible  $\iff (X, A_1)$  is an extensible pair.

(2) If  $k_0$  is such that det  $Z_{k_0} = 0$ , then many of our familiar statements for  $\mathfrak{g}(Z_{k_0})$  break down; for instance  $P(Z_{k_0})/Q(Z_{k_0})$  is no longer a finite group, the  $\omega_p$ 's do not span  $\mathfrak{h}^*(Z_{k_0})$  etc. But if  $(X_1, X_2)$  is an extensible pair,  $\Delta_1 \Delta_2 \neq 0$ . By Lemma 2 there can be at most one  $k_0$  for which det  $Z_{k_0} = 0$ . So this exceptional situation can occur for at most one value of k. Most of our later results will only hold for  $k \neq k_0$ .

**Lemma 3.** If  $(X_1, X_2)$  is an extensible pair, then  $gcd(\Delta_1 \Delta_2, det Z_k) = 1 \ \forall k \ge 1$ 

**Proof:** Let P be a prime such that  $P | \Delta_1 \Delta_2$ . Then  $P | \Delta_1$  or  $P | \Delta_2$ . Suppose  $P | \Delta_1$ , definition 3 implies that P does not divide either det  $X_1$  or  $\Delta_2$ . From Equation (3.3), this means that P cannot divide det  $Z_k$ . The  $P | \Delta_2$  case is similar.

**Remark 3.** It is easily seen that if  $(X_1, X_2)$  is an extensible pair, then so is  $(X_1(m), X_2(n))$  for almost all  $m, n \ge 0$ . The only exceptional values are those which make  $\det(X_1(m)) = 0$  or  $\det(X_2(n)) = 0$ .

**Example 1.** Let  $\mathcal{U}$  be the set of marked Dynkin diagrams U with det  $U \neq 0$  and  $\Delta_U = \pm 1$ . From Table 1 of [4], it is clear that Types  $A, E, F^{(1)}, F^{(2)}, G^{(1)}, G^{(2)}$  are all in  $\mathcal{U}$ .

Given  $X_1, X_2 \in \mathcal{U}$ , it is clear that  $(X_1, X_2)$  is an extensible pair. Some of the sequences  $Z_k$  obtained thus are :



3.3. Node numbering. Let  $(X_1, X_2)$  be an extensible pair and assume  $X_i$  has  $d_i$  nodes i = 1, 2. We assume we are given numberings  $\epsilon_i$  of  $X_i$  (i = 1, 2) as in section 3.1. We use the numberings  $\epsilon_1, \epsilon_2$  and  $\epsilon_A$  to construct two numberings  $i, \bar{i}$  of  $Z_k$ . Define  $i, \bar{i}: N(Z_k) \to \{1 \le m \le d_1 + d_2 + k\}$  by

$$i(p) := \begin{cases} \epsilon_1(p) & \text{if } p \in N(X_1) \\ \epsilon_A(p) + d_1 & \text{if } p \in N(A_k) \\ (d_1 + d_2 + k + 1) - \epsilon_2(p) & \text{if } p \in N(X_2) \end{cases}$$

$$\overline{i}(p) := \begin{cases} \epsilon_2(p) & \text{if } p \in N(X_2) \\ (d_2 + k + 1) - \epsilon_A(p) & \text{if } p \in N(A_k) \\ (d_1 + d_2 + k + 1) - \epsilon_1(p) & \text{if } p \in N(X_1) \end{cases}$$

Our earlier figure showed  $Z_k$  with the numbering given by  $i(\cdot)$ . The figure below shows the numbering via  $\bar{i}(\cdot)$ .



3.4.  $P(Z_k)/Q(Z_k)$  is cyclic. Let  $\widehat{Y}_k := Y_k(0,0) \cup \{\xi_1,\xi_2\}$ . Our new notion of *extensible pairs* has the same nice consequence as the previous notion of *extensible diagrams*:

**Lemma 4.** Let  $(X_1, X_2)$  be an extensible pair. For  $k \ge 1$ , if det  $Z_k \ne 0$ , then  $P(Z_k)/Q(Z_k)$  is a cyclic group. Further, if  $u, v \in \widehat{Y}_k \subset N(Z_k)$  are any two adjacent nodes then  $[\omega_u - \omega_v]$  generates this group.

**Proof:** Let  $p, p', p'' \in \widehat{Y}_k$  be 3 adjacent nodes as in figure



Iterating this argument, it is clear that the elements  $[\omega_u - \omega_v]$  with  $u, v \in \widehat{Y}_k$  and i(v) = i(u) + 1 are all equal. Further if  $[\omega_u - \omega_v]$  is a generator of the group, then so is  $[\omega_v - \omega_u]$ . It is thus enough to prove the lemma for a fixed choice of  $u, v \in \widehat{Y}_k$  with i(v) = i(u) + 1.

We write  $\omega_u - \omega_v = \sum_{p \in Z_k} c_p \alpha_p$  with  $c_p \in \mathbb{Q}$ . We now compute  $c_u$  and  $c_v$ . Set  $a = i(u) - d_1 - 1$  and  $b = \overline{i}(v) - d_2 - 1$ , so a + b = k - 2.

The coefficient of  $\alpha_u$  in  $\omega_u$  is the  $(u, u)^{th}$  element of  $C(Z_k)^{-1}$ , the inverse of the generalized Cartan matrix. This coefficient is thus:

$$\frac{\operatorname{cofactor of} (u, u)^{th} \text{ element of } C(Z_k)}{\det Z_k} = \frac{1}{\det Z_k} \det X_1(a) \det X_2(b+1)$$

Similarly the coefficient of  $\alpha_v$  in  $\omega_u$  is the  $(v, u)^{th}$  element of  $C(Z_k)^{-1}$ , and is thus  $-\det A/\det Z_k$  where

$$A = \begin{bmatrix} C(X_1(a)) & & \\ & -1 & \\ \hline & & -1 & -1 \\ \hline & & & C(X_2(b)) \end{bmatrix}$$

The upper triangularity of A gives det  $A = -\det X_1(a) \det X_2(b)$ . We now switch the roles of u and v; the coefficient of  $\alpha_u$  in  $\omega_v$  is det  $X_1(a) \det X_2(b)/\det Z_k$  and coefficient of  $\alpha_v$  in  $\omega_v$  is det  $X_1(a + 1) \det X_2(b)/\det Z_k$ . Finally we also have Equation (3.1) which gives det  $X_i(r) = \det X_i + r\Delta_i$ . Putting these all together, we get

(3.5) 
$$c_u - c_v = \frac{\det X_1 \Delta_2 + \det X_2 \Delta_1 + (k-2)\Delta_1 \Delta_2}{\det Z_k}$$

Lemma 3 implies that the numerator and denominator of Equation (3.5) are relatively prime. So, the least  $n \in \mathbb{Z}^{\geq 0}$  such that  $n(c_u - c_v) \in \mathbb{Z}$  is  $n = |\det Z_k|$ . In particular this implies that the order of the element  $[\omega_u - \omega_v]$  in  $P(Z_k)/Q(Z_k)$  is  $\geq |\det Z_k|$ . Since  $\#P(Z_k)/Q(Z_k) = |\det Z_k|$ , this completes the proof of Lemma 4.

Now fix  $u, v \in \widehat{Y}_k \subset Z_k$  with i(v) = i(u) + 1. Our next goal is to explicitly write each  $[\omega_p]$ ,  $p \in Z_k$  as a multiple of  $[\omega_u - \omega_v]$ . for each  $p \in Z_k$ . This is analogous to Proposition 5. In fact we will use this latter proposition to deduce our result below.

Consider the subdiagrams  $\widetilde{X}_1 := X_1(i(u))$  and  $\widetilde{X}_2 := X_2(\overline{\imath}(v))$  of  $Z_k$ . The numberings  $i, \overline{\imath}$  of  $Z_k$  can be restricted to  $\widetilde{X}_i$ , i = 1, 2 to give functions from  $\widetilde{X}_i$ into the set  $\{1, 2, \dots, d_1+d_2+k\}$ . Since  $(X_1, X_2)$  is an extensible pair, each  $X_i$  is an extensible diagram. For fixed  $i \in \{1, 2\}$ , let  $\{\widetilde{\omega}_p : p \in \widetilde{X}_i\}$  and  $\{\widetilde{\alpha}_p : p \in \widetilde{X}_i\}$  denote the fundamental weights and simple roots of  $\widetilde{X}_i$ . We can now apply Proposition 5 above to the  $X_i, i = 1, 2$ . This gives:

**Proposition 6.** There exist infinite sequences  $(a_i(X_1))_{i=1}^{\infty}$  and  $(a_i(X_2))_{i=1}^{\infty}$  (determined uniquely by  $X_1$  and  $X_2$ ) such that

(3.6) 
$$-\Delta_1 \ \tilde{\omega}_p \equiv a_{i(p)}(X_1) \ \tilde{\omega}_u \pmod{Q(X_1)} \quad \forall p \in X_1$$

(3.7) 
$$-\Delta_2 \ \tilde{\omega}_p \equiv a_{\bar{\imath}(p)}(X_2) \tilde{\omega}_v \pmod{Q(X_2)} \quad \forall p \in X_2$$

Additionally,

Lemma 5. (1) For 
$$p \in \tilde{X}_1$$
, let  
 $\tilde{\beta}_p := -\Delta_1 \ \tilde{\omega}_p - a_{i(p)}(X_1) \ \tilde{\omega}_u = \sum_{q \in \tilde{X}_1} \tilde{c}_{q,p} \ \tilde{\alpha}_q$ 

with  $\tilde{c}_{q,p} \in \mathbb{Z}$ . Then  $\tilde{c}_{u,p} = -a_{i(p)}(X_1)$ .

(2) For 
$$p \in \tilde{X}_2$$
, let  
 $\tilde{\beta}_p := -\Delta_2 \ \tilde{\omega}_p - a_{\bar{\imath}(p)}(X_2) \ \tilde{\omega}_v = \sum_{q \in \tilde{X}_2} \tilde{c}_{q,p} \ \tilde{\alpha}_q$ 
with  $\tilde{c}_{q,p} \in \mathbb{Z}$ . Then  $\tilde{c}_{v,p} = -a_{\bar{\imath}(p)}(X_2)$ .

*Proof:* Follows from lemma 1 gives.

We now consider the original diagram  $Z_k$ . Fix  $p \in \widetilde{X}_1$ , take the coefficients  $\tilde{c}_q$  given by Lemma 5 and consider the element  $\beta_p \in Q(Z_k)$  given by

$$\beta_p := \sum_{q \in \widetilde{X}_1} \widetilde{c}_{q,p} \, \alpha_q$$

The  $\alpha_q$  are now the simple roots of  $Z_k$ . Looking at how  $\widetilde{X}_1$  sits inside  $Z_k$  as a subdiagram, it is clear that the following relations hold:

(3.8) 
$$\beta_p(\tilde{\alpha}_r) = \beta_p(\alpha_r) \qquad \forall r \in X_1 \subset Z_k$$
$$\beta_p(\alpha_v) = -\tilde{c}_{u,p} = a_{i(p)}(X_1)$$
$$\beta_p(\alpha_q) = 0 \qquad \forall q \notin \widetilde{X}_1 \cup \{v\}$$

These follow from the fact that u is the only node connected to v and from Lemma 5.

Equation (3.8) and the definition of  $\hat{\beta}_p$  imply the following expression for  $\beta_p$  as a linear combination of the  $\omega_r$ ,  $r \in Z_k$ 

(3.9) 
$$\beta_p = -\Delta_1 \,\omega_p - a_{i(p)}(X_1)(\omega_u - \omega_v) \quad \forall p \in Z_k \text{ with } i(p) \le i(u)$$

The corresponding picture for  $p \in \widetilde{X}_2$  is obtained similarly; we define  $\beta'_p \in Q(Z_k)$  by

$$\beta_p' := \sum_{q \in \widetilde{X}_2} \widetilde{c}_{q,p} \alpha_q$$

The earlier argument for  $\beta_p$  can be carried out with obvious modifications and gives the following expression for  $\beta'_p$ :

(3.10) 
$$\beta'_p = -\Delta_2 \,\omega_p - a_{\bar{\imath}(p)}(X_2)(\omega_v - \omega_u) \quad \forall p \in Z_k \text{ with } \bar{\imath}(p) \le \bar{\imath}(v)$$

We now have the following proposition which follows from Equations (3.9), (3.10) and the fact that  $\beta_p, \beta'_p \in Q(Z_k)$ .

**Proposition 7.** Let  $p \in N(Z_k)$ . Then

(1) If  $i(p) \leq i(u)$ , then  $-\Delta_1 \omega_p \equiv a_{i(p)}(X_1)(\omega_u - \omega_v) \pmod{Q(Z_k)}$ (2) If  $\overline{i}(p) \leq \overline{i}(v)$ , then  $-\Delta_2 \omega_p \equiv a_{\overline{i}(p)}(X_2)(\omega_v - \omega_u) \pmod{Q(Z_k)}$ 

If 
$$i(p) \leq i(v)$$
, then  $-\Delta_2 \omega_p = d_{\overline{i}(p)}(\Lambda_2)(\omega_v - \omega_u) \pmod{Q(Z_k)}$ 

4. Stabilization of 
$$c^{\nu}_{\lambda\mu}(k)$$
 and  $b_{\lambda\beta}(k)$ 

4.1. Two sided dominant weights. We will now index dominant integral weights of  $\mathfrak{g}(Z_k)$  as in [4], in a two-sided (or *double headed*) fashion. Unlike in section 2.4, we consider weights whose supports are not necessarily contained in  $X_1 \cup X_2$ . Let

 $\mathcal{H}_1 := \{ (x_1, x_2, \cdots) : x_i \in \mathbb{Z} \,\forall i \text{ and } x_i \neq 0 \text{ for only finitely many } i \}$ 

and

$$\mathcal{H}_{1}^{+} = \{ (x_{1}, x_{2}, \cdots) \in \mathcal{H}_{1} : x_{i} \ge 0 \ \forall i \}$$

Given  $x = (x_1, x_2, \dots) \in \mathcal{H}_1$  we define the *length* of x to be:  $\ell(x) := \max\{i : x_i \neq 0\}$ . Let

$$\mathcal{H}_2 := \mathcal{H}_1 \times \mathcal{H}_1$$
 and  $\mathcal{H}_2^+ := \mathcal{H}_1^+ \times \mathcal{H}_1^+$ 

Given elements  $x = (x_1, x_2, \cdots)$ ,  $y = (y_1, y_2, \cdots)$  of  $\mathcal{H}_1$ , let  $\gamma = (x, y) \in \mathcal{H}_2$ . We define the *left support* of  $\gamma$  to be  $ls(\gamma) := max(\ell(x), d_1)$ , the *right support* as  $rs(\gamma) := max(\ell(y), d_2)$  and the *support*  $\ell(\gamma) := ls(\gamma) + rs(\gamma)$ . Recall that  $d_i$  here is the number of nodes in  $X_i$ . Now,  $\gamma$  can be used to define a weight of  $\mathfrak{g}(Z_k)$  for all large k; specifically for  $k \geq \ell(\gamma) - d_1 - d_2$  as follows. Define

$$\gamma^{(k)} := \sum_{p \in Z_k} (x_{i(p)} + y_{\bar{\imath}(p)}) \,\omega_p$$

Note that since k is large, at most one of  $x_{i(p)}$  or  $y_{\overline{i}(p)}$  can be nonzero for each  $p \in N(Z_k)$ . The figure below shows each node p of  $Z_k$  labeled by the corresponding  $x_i$  or  $y_i$ .

4.2. The number of boxes function. Given an extensible Dynkin diagram X, section 3.3 of [4] introduced the so called number of boxes function  $|\cdot|_X : \mathcal{H}_2 \to \mathbb{Z}$ . For an extensible pair  $(X_1, X_2)$ , we now define an analogous function  $|\cdot|_{(X_1, X_2)}$ . We shall refer to this also as the number of boxes function.

**Definition 4.** Given  $\gamma = (x, y) \in \mathcal{H}_2$  as above, define

$$|\gamma|_{(X_1,X_2)} := \Delta_2 \left(\sum_i x_i a_i(X_1)\right) - \Delta_1 \left(\sum_i y_i a_i(X_2)\right)$$

**Remark 4.** When  $X_2 = A_1$  we have  $\Delta_2 = 1$  and  $a_i(X_2) = i$ . Then  $|\gamma|_{(X_1,A_1)}$  coincides with the function  $|\gamma|_{X_1}$  in definition (3.12) of [4].

To see the significance of  $|\gamma|_{(X_1,X_2)}$ , we consider the following situation. Let  $\gamma \in \mathcal{H}_2$  and  $k \geq \ell(\gamma) - d_1 - d_2$  be fixed. Pick  $u, v \in \widehat{Y}_k$  such that  $i(u) \geq \operatorname{ls}(\gamma)$ ,  $\overline{i}(v) \geq \operatorname{rs}(\gamma)$  and i(v) = i(u) + 1. Consider  $\gamma^{(k)} \in P(Z_k)$ . Recall from Lemma 4 that  $\gamma^{(k)}$  must be congruent modulo  $Q(Z_k)$  to an integer multiple of  $\omega_u - \omega_v$ . To calculate this integer explicitly,

$$(4.1) \qquad \qquad -\Delta_{1} \Delta_{2} \gamma^{(k)} - |\gamma|_{(X_{1}, X_{2})} (\omega_{u} - \omega_{v}) = \Delta_{2} \left( \sum_{\substack{p \in Z_{k} \\ i(p) \leq i(u)}} x_{i(p)} \left[ -\Delta_{1} \omega_{p} - a_{i(p)} (X_{1}) (\omega_{u} - \omega_{v}) \right] \right) + \Delta_{1} \left( \sum_{\substack{p \in Z_{k} \\ \overline{i}(p) \leq \overline{i}(v)}} y_{\overline{i}(p)} \left[ -\Delta_{2} \omega_{p} - a_{\overline{i}(p)} (X_{2}) (\omega_{v} - \omega_{u}) \right] \right)$$

Proposition 7 shows that both terms within square brackets in the above equation are elements of  $Q(Z_k)$ . So, we have

Proposition 8. With notation as above,

(4.2) 
$$-\Delta_1 \Delta_2 \gamma^{(k)} \equiv |\gamma|_{(X_1, X_2)} (\omega_u - \omega_v) \pmod{Q(Z_k)}$$

Observe that this equation enables us to write  $\gamma^{(k)}$  itself as a multiple of  $(\omega_u - \omega_v)$ modulo  $Q(Z_k)$ . To see this, note that Lemma 3 gives  $gcd(\Delta_1 \Delta_2, \det Z_k) = 1$ . Since  $\#P(Z_k)/Q(Z_k) = |\det Z_k|$ , we have

(4.3) 
$$\gamma^{(k)} \equiv (\Delta_1 \, \Delta_2)^{-1} |\gamma|_{(X_1, X_2)} \; (\omega_u - \omega_v) \pmod{Q(Z_k)}$$

where  $(\Delta_1 \Delta_2)^{-1}$  denotes the inverse of  $\Delta_1 \Delta_2$  in  $\frac{\mathbb{Z}}{(\det Z_k)\mathbb{Z}}$ . We also have the following corollary to the above proposition:

Corollary 2. Let  $\gamma \in \mathcal{H}_2$ . Then

$$\begin{split} |\gamma|_{(X_1,X_2)} &= 0 \iff \gamma^{(k)} \in Q(Z_k) \text{ for all large } k \\ \iff \gamma^{(k)} \in Q(Z_k) \text{ for infinitely many values of } k \end{split}$$

**Proof:** Since  $\omega_u - \omega_v$  generates the cyclic group  $P(Z_k)/Q(Z_k)$ , Equation (4.3) implies that  $\gamma^{(k)} \in Q(Z_k)$  *iff* det  $Z_k$  divides  $|\gamma|_{(X_1,X_2)}$ . But as  $k \to \infty$ ,  $|\det Z_k| \to \infty$  since  $\Delta_1 \Delta_2 \neq 0$ ; this follows from Equation (3.3). The corollary follows.  $\Box$ 

4.3. **Depth.** In this subsection, we introduce an important statistic called the *depth* on the set  $\{\gamma : |\gamma|_{(X_1,X_2)} = 0\}$ .

Let  $\gamma = (x, y) \in \mathcal{H}_2$  with  $|\gamma|_{(X_1, X_2)} = 0$ . Fix  $k \ge \ell(\gamma) - d_1 - d_2$  and pick u, v as in previous the subsection.

Consider Equation (4.1). Imposing the condition  $|\gamma|_{(X_1,X_2)} = 0$  reduces the left hand side to  $-\Delta_1 \Delta_2 \gamma^{(k)}$ . As already remarked, the sum on the right hand side of (4.1) is an element of  $Q(Z_k)$ . We now calculate the coefficient of  $\alpha_u$  in this sum. Using Lemma 5 and Equation (3.9), the coefficient of  $\alpha_u$  on the RHS equals

$$-\Delta_2 \left(\sum_{\substack{p \in Z_k \\ i(p) \le i(u)}} x_{i(p)} a_{i(p)}(X_1)\right) = -\Delta_2 \sum_i x_i a_i(X_1)$$

Similarly, employing Equation (3.10), the coefficient of  $\alpha_v$  on the RHS of (4.1) becomes

$$-\Delta_1\left(\sum_{\substack{p\in Z_k\\\bar{\imath}(p)\leq\bar{\imath}(u)}}y_{\bar{\imath}(p)}a_{\bar{\imath}(p)}(X_2)\right) = -\Delta_1\sum_i y_i a_i(X_2)$$

Observe now that since  $|\gamma|_{(X_1,X_2)} = 0$ , the coefficients of  $\alpha_u$  and  $\alpha_v$  are in fact equal ! Putting everything together we get

**Lemma 6.** If  $|\gamma|_{(X_1,X_2)} = 0$ , then we have:

coefficient of 
$$\alpha_u$$
 in  $-\Delta_1 \Delta_2 \gamma^{(k)} = \text{ coefficient of } \alpha_v \text{ in } -\Delta_1 \Delta_2 \gamma^{(k)}$ 

(4.4) 
$$= -\Delta_2 \sum_i x_i a_i(X_1) = -\Delta_1 \sum_i y_i a_i(X_2)$$

This motivates the following

**Definition 5.** If  $\gamma = (x, y) \in \mathcal{H}_2$  with  $|\gamma|_{(X_1, X_2)} = 0$ , the *depth* of  $\gamma$  is defined to be

$$dep(\gamma) := \frac{\sum_{i} x_i a_i(X_1)}{\Delta_1} = \frac{\sum_{i} y_i a_i(X_2)}{\Delta_2}$$

**Remark 5.** (1) Observe that  $dep(\gamma) \in \mathbb{Z}$  since  $gcd(\Delta_1, \Delta_2) = 1$ .

(2) From equation (4.4), it is clear that dep( $\gamma$ ) is just the coefficient of  $\alpha_u$  (or  $\alpha_v$ ) in  $\gamma^{(k)}$ .

(3) When  $X_2 = A_1$ , our notion of depth reduces to the notion introduced in [4], Definition (4.3).

Corollary 2 guarantees that if  $|\gamma|_{(X_1,X_2)} = 0$ , then  $\gamma^{(k)}$  can be written as an integer linear combination of  $\{\alpha_p : p \in Z_k\}$  for all large k. In this subsection, we study how the coefficients of this linear combination change with k.

In order to be able to refer more easily to nodes of  $Z_k$  which occur to the left/right/middle of the diagram, we define some notation; for  $l, r \in \mathbb{Z}^{\geq 0}$  we let

$$L_{k}(l) := \{ p \in Z_{k} : i(p) \le l \}$$
  

$$R_{k}(r) := \{ p \in Z_{k} : \bar{\imath}(p) \le r \}$$
  

$$M_{k}(l,r) := Z_{k} - (L_{k}(l) \cup R_{k}(r))$$

We now have the following proposition which generalizes Proposition (4.1) of [4].

**Proposition 9.** Suppose  $\gamma \in \mathcal{H}_2$  with  $|\gamma|_{(X_1,X_2)} = 0$ ; let  $l := ls(\gamma), r := rs(\gamma)$ . Then there exist integers  $b_i \ (1 \le i \le l-1), c_i \ (1 \le i \le r-1)$  and s such that for all  $k \ge l + r - d_1 - d_2$  we have

(4.5) 
$$\gamma^{(k)} = \sum_{p \in L_k(l-1)} b_{i(p)} \alpha_p + \sum_{p \in M_k(l-1,r-1)} s \alpha_p + \sum_{p \in R_k(r-1)} c_{\bar{\imath}(p)} \alpha_p$$

**Proof:** Let k be as in the statement; the given assertion easily reduces to the following: there exists  $s \in \mathbb{Z}$  such that when we write  $\gamma^{(k)}$  as a linear combination of simple roots, the coefficients of  $\alpha_p, p \in M_k(l-1, r-1)$  are all equal to s.

Let  $s := \operatorname{dep}(\gamma)$ ; if  $\gamma^{(k)} = \sum c_p \alpha_p$ , remark 5 gives  $c_u = c_v = s$ . Now, all this holds for any u, v which satisfy  $i(u) \ge \operatorname{ls}(\gamma)$ ,  $\overline{i}(v) \ge \operatorname{rs}(\gamma)$  and i(v) = i(u) + 1. This concludes the proof.

The figure now shows the nodes of  $Z_k$  labeled by  $b_i, c_j, s$ :



4.4. Main theorem. Our main theorem is the following generalization of Theorem (4.5) of [4].

**Theorem 3.** Let  $(X_1, X_2)$  be an extensible pair.

- (1) Given  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ , there exists an integer  $K = K(\lambda, \mu, \nu)$  such that  $c_{\lambda\mu}^{\nu}(k) = c_{\lambda\mu}^{\nu}(k')$  for all  $k, k' \geq K$ .
- (2) Given  $\lambda \in \mathcal{H}_2^+$  and  $\beta \in \mathcal{H}_2$ , there exists K such that  $b_{\lambda\beta}(k) = b_{\lambda\beta}(k')$  for all  $k, k' \geq K$ .

**Proof:** Let l be the maximum of the left supports of  $\lambda, \mu, \nu, \beta$  and r be the maximum of their right supports. Let  $X'_1 := X_1(l - d_1)$  be the diagram formed by the nodes  $p \in L_k(l)$  and  $X'_2 := X_2(r - d_2)$  be that formed by  $p \in R_k(r)$ . We can view  $\lambda, \mu, \nu, \beta$  as elements of  $W(X'_1, X'_2)$  as in section 2.4.

Define  $\gamma := \lambda + \mu - \nu$ . Corollary 2 implies that if  $(\lambda + \mu) \sim \nu$ , then  $|\gamma|_{(X_1, X_2)} = 0$ . Proposition 9 now applies and we deduce stabilization of  $c^{\nu}_{\lambda\mu}(k)$  by applying Theorem 1.

The second assertion regarding the  $b_{\lambda\beta}(k)$  follows analogously. We set  $\gamma = \lambda - \beta$  and use Proposition 9 and Theorem 1.

### 5. The stable representation ring

5.1. **Partial orders.** Having established that the multiplicities  $c_{\lambda\mu}^{\nu}(k)$  stabilize for extensible pairs  $(X_1, X_2)$ , we shall now use the stable values  $c_{\lambda\mu}^{\nu}(\infty)$  as structure constants to define an associative multiplication \*. First, we take a small detour and define a partial order on the set  $\mathcal{H}_2^+$ .

Let  $(X_1, X_2)$  be an extensible pair of marked Dynkin diagrams as before. The set  $P^+(Z_k)$  of dominant weights of  $\mathfrak{g}(Z_k)$  is partially ordered with

$$\tau_1, \tau_2 \in P^+(Z_k), \quad \tau_1 \ge \tau_2 \iff \tau_1 - \tau_2 \in Q^+(Z_k)$$

We can also make  $\mathcal{H}_2^+$  into a poset. Let  $\lambda_1 = (x, y), \lambda_2 = (z, w) \in \mathcal{H}_2^+$ be such that  $|\lambda_1|_{(X_1, X_2)} = |\lambda_2|_{(X_1, X_2)}$ . Let  $l = \max(\operatorname{ls}(\lambda_1), \operatorname{ls}(\lambda_2))$  and  $r = \max(\operatorname{rs}(\lambda_1), \operatorname{rs}(\lambda_2))$ . Proposition 9 implies that there exist integers  $b_i$   $(1 \le i \le l-1)$ ,  $c_j$   $(1 \le j \le r-1)$  and s such that for  $k \ge l+r-d_1-d_2$ 

(5.1) 
$$\lambda_1^{(k)} - \lambda_2^{(k)} = \sum_{p \in L_k(l-1)} b_{i(p)} \alpha_p + \sum_{p \in M_k(l-1,r-1)} s \, \alpha_p + \sum_{p \in R_k(r-1)} c_{\bar{\imath}(p)} \alpha_p$$

**Definition 6.** Given  $\lambda_1, \lambda_2 \in \mathcal{H}_2^+$ , define  $\lambda_1 \succeq \lambda_2$  iff  $|\lambda_1|_{(X_1, X_2)}| = |\lambda_2|_{(X_1, X_2)}$ and the  $b_i, c_j, s$  which occur in Equation (5.1) are all non-negative.

It is easy to check that  $\succeq$  is a partial order on  $\mathcal{H}_2^+$  and that  $\lambda_1 \succeq \lambda_2$  implies that  $\lambda_1 + \mu \succeq \lambda_2 + \mu$  for all  $\mu \in \mathcal{H}_2^+$ . We also have these equivalent conditions which follow easily:

(5.2) 
$$\lambda_1 \succcurlyeq \lambda_2 \Leftrightarrow \lambda_1^{(k)} \ge \lambda_2^{(k)} \quad \forall k \ge l + r - d_1 - d_2$$

(5.3)  $\Leftrightarrow \lambda_1^{(k)} \ge \lambda_2^{(k)} \text{ for infinitely many values of } k$ 

where  $\geq$  is the partial order on  $P^+(Z_k)$ . We also have:

Lemma 7. (1) If  $\lambda_1 \succeq \lambda_2$ , then  $dep(\lambda_1 - \lambda_2) = s \ge 0$ (2) If  $\lambda, \mu, \nu \in \mathcal{H}_2^+$  are such that  $c_{\lambda\mu}^{\nu}(\infty) > 0$ , then  $\lambda + \mu \succeq \nu$ .

**Proof:** The proof of (1) is trivial. For (2), observe that  $c_{\lambda\mu}^{\nu}(\infty) > 0$  implies that  $c_{\lambda\mu}^{\nu}(k) > 0$  for all large k. In particular  $\lambda^{(k)} + \mu^{(k)} \ge \nu^{(k)}$  for all large k. Equation (5.3) completes the proof.

5.2. One sided intervals. The goal of this subsection is to analyze the partial orders on  $\mathcal{H}_2^+$  and  $P^+(Z_k)$ .

Assume  $(X_1, X_2)$  is an extensible pair. For each  $k \ge 1$ , we know that  $\mathfrak{g}(Z_k)$  is a symmetrizable Kac-Moody algebra. Suppose  $(X_1, X_2) \ne (A_{d_1}, A_{d_2})$  and k is such that det  $Z_k \ne 0$ , then  $Z_k$  cannot be a *finite* type Dynkin diagram; to see this we note that among the series of finite type Dynkin diagrams, only  $A_n$  is extensible while the B/C/D types are not (see Table 1 of [4]). The det  $Z_k \ne 0$  condition also implies that  $Z_k$  cannot be affine. So, it must be of *indefinite* type.

We first derive a result about the poset of dominant integral weights in any Kac-Moody algebra of *indefinite* type. Let  $\mathfrak{g}$  be a symmetrizable Kac-Moody algebra of indefinite type. Let A be the corresponding  $n \times n$  generalized Cartan matrix. Assume A is indecomposable and that det  $A \neq 0$ . We use the usual notation for roots, weights etc. Let (.|.) denote a nondegenerate, symmetric, bilinear,  $\mathbb{C}$ -valued form on the Cartan subalgebra  $\mathfrak{h}$ . This form exists because  $\mathfrak{g}$  is symmetrizable. Fix  $\gamma \in P^+(\mathfrak{g})$ . Consider the set:

$$U(\gamma) := \{\lambda \in P^+(\mathfrak{g}) : \lambda \ge \gamma\}$$

Here again  $\lambda \geq \gamma$  is defined by the condition  $\lambda - \gamma \in Q^+(\mathfrak{g})$ . The set  $U(\gamma)$  is the one-sided interval in the poset  $P^+(\mathfrak{g})$ , bounded below by  $\gamma$ .

**Proposition 10.**  $U(\gamma)$  is a finite set.

**Proof:** The definition of an *indefinite type* Kac-Moody algebra [3, Chapter 4] implies that there exists  $\xi = \sum_{i=1}^{n} u_i \alpha_i \in \mathfrak{h}^*$  which satisfies the conditions (a)  $u_j > 0 \forall j$  (b)  $\xi(\check{\alpha}_j) < 0 \forall j$ . Condition (b) implies that  $(\xi | \alpha_j) < 0 \forall j$  since  $(\xi | \alpha_j) = \frac{(\alpha_j | \alpha_j)}{2} \xi(\check{\alpha}_j)$  and  $(\alpha_j | \alpha_j) > 0$ .

Condition (a) together with the fact that  $\omega_j(\check{\alpha}_i) = \delta_{ij}$  implies that  $(\xi | \omega_j) > 0 \forall j$ . To see this observe:

$$(\xi|\omega_j) = \sum_{i=1}^n u_i(\alpha_i|\omega_j)$$
$$= u_j(\alpha_j|\alpha_j)/2$$

For any  $\eta \in \mathfrak{h}^*$ , define  $ht(\eta) := (\eta | \xi)$ . The conclusions of the above paragraph imply:

Fact 1:  $ht(\alpha) \leq 0$  for all  $\alpha \in Q^+(\mathfrak{g})$ . Fact 2:  $ht(\lambda) \geq 0$  for all  $\lambda \in P^+(\mathfrak{g})$ .

If  $\lambda \in U(\gamma)$ , then  $\lambda - \gamma \in Q^+(\mathfrak{g})$ . Hence  $\operatorname{ht}(\lambda) - \operatorname{ht}(\gamma) = \operatorname{ht}(\lambda - \gamma) \leq 0$ . So, we have:

**Fact 3**: If  $\lambda \in U(\gamma)$ , then

$$0 \le \operatorname{ht}(\lambda) \le \operatorname{ht}(\gamma)$$

We can now show that  $U(\gamma)$  is finite. Let  $\lambda = \sum_{i=1}^{n} a_i \omega_i$  with  $a_i \in \mathbb{Z}^{\geq 0} \forall i$ . Then  $\operatorname{ht}(\lambda) = \sum_{i=1}^{n} a_i (u_i(\alpha_i | \alpha_i)/2) \leq \operatorname{ht}(\gamma)$ . Since each term in the sum is nonnegative, this means that

$$a_i \le \frac{2\operatorname{ht}(\gamma)}{u_i(\alpha_i|\alpha_i)} \ \forall i$$

Since  $a_i$  is a nonnegative integer, there are only finitely many choices for  $a_i$ . Thus  $U(\gamma)$  is a finite set.

**Remark 6.** The above Proposition is false if  $\mathfrak{g}$  is of Finite or Affine type. For instance if  $\mathfrak{g} = \mathfrak{g}(A_1) = sl_2\mathbb{C}$ , and  $\gamma = \alpha_1$ , then  $U(\gamma) = \{n\alpha_1 : n \ge 1\}$ . Similarly for the rank 2 affine Lie algebra  $\mathfrak{g} = A_1^{(1)}$  (affine  $sl_2$ ), if  $\gamma = \alpha_1 + \alpha_2$ , then  $U(\gamma) = \{n(\alpha_1 + \alpha_2) : n \ge 1\}$ .

We now consider the poset  $(\mathcal{H}_2^+, \succeq)$  and ask the analogous question. Fix  $\gamma \in \mathcal{H}_2^+$ and let

(5.4) 
$$U(\gamma) := \{\lambda \in \mathcal{H}_2^+ : \lambda \succcurlyeq \gamma\}$$

We use the same notation  $U(\cdot)$  that we used before, but the context will resolve any confusion. The following example shows that the poset  $(\mathcal{H}_2^+, \succeq)$  is unlike the poset  $P^+(\mathfrak{g})$ ; even if most (or all) members of the set  $\{\mathfrak{g}(Z_k)\}$  are of indefinite type, the set  $U(\gamma)$  may still be infinite.

**Example 2.** Let  $X_1 = E_{10}$  with the node numbering and distinguished vertex shown below



and take  $X_2 = A_1$ . We let  $\epsilon_i$  denote the element  $(0, 0, \dots, 0, 1, 0, 0, \dots)$  of  $\mathcal{H}_1^+$ , where the 1 occurs in the *i*<sup>th</sup> position. Take  $\gamma := (\epsilon_{10}, 0)$ . For k > 0, consider the element  $\lambda_k := (\epsilon_{10+k}, \epsilon_k)$  of  $\mathcal{H}_2^+$ . We claim that  $\lambda_k \in U(\gamma)$  for all k > 0. The proof follows by actual computation of the  $b_i, c_j, s$ :

$$\begin{aligned} b_i &= 0 \quad (i \leq 10), \\ s &= k \end{aligned} \qquad b_i &= i - 10 \quad (10 < i < 10 + k) \\ c_i &= i \quad (1 \leq i < k) \end{aligned}$$

Thus  $U(\gamma)$  is an infinite set.

However, a weaker finiteness assertion is true for the poset  $\mathcal{H}_2^+$ . Given  $s \in \mathbb{Z}$ , define

$$U(\gamma, s) := \{\lambda \in U(\gamma) : \operatorname{dep}(\lambda - \gamma) = s\}$$

Lemma 7 implies that  $U(\gamma, s) = \emptyset$  if s < 0. Clearly

$$U(\gamma) = \bigsqcup_{s=0}^{\infty} U(\gamma, s)$$

We have:

**Proposition 11.**  $U(\gamma, s)$  is a finite set for each  $s \in \mathbb{Z}^{\geq 0}$ .

**Proof:** Let  $\gamma = (x, y)$ , and suppose  $\lambda = (z, w) \in U(\gamma, s)$  with  $x, y, z, w \in \mathcal{H}_1^+$ . Write  $x = (x_i)_{i \ge 1}$ ,  $y = (y_i)_{i \ge 1}$ ,  $z = (z_i)_{i \ge 1}$ ,  $w = (w_i)_{i \ge 1}$ . Let  $l = \max(\operatorname{ls}(\gamma), \operatorname{ls}(\lambda))$ ,  $r = \max(\operatorname{rs}(\gamma), \operatorname{rs}(\lambda))$ .

First assume that  $(X_1, X_2) = (A_{d_1}, A_{d_2})$ . Then  $\Delta_1 = \Delta_2 = 1$  and  $a_i(X_1) = a_i(X_2) = i \forall i$ . We have

$$s = \sum_{j} jz_j - \sum_{j} jx_j = \sum_{j} jw_j - \sum_{j} jy_j$$

Clearly, there are only finitely many choices for (z, w) such that this condition holds. Hence the number of  $\lambda = (z, w)$  in  $U(\gamma, s)$  is finite and we are done.

So, assume now that  $(X_1, X_2) \neq (A_{d_1}, A_{d_2})$ . Since  $\lambda \geq \gamma$ , there are non-negative integers  $b_i$   $(1 \leq i \leq l-1)$ ,  $c_j$   $(1 \leq j \leq r-1)$  and s such that for all large k

(5.5) 
$$\lambda^{(k)} - \gamma^{(k)} = \sum_{p \in L_k(l-1)} b_{i(p)} \alpha_p + \sum_{p \in M_k(l-1,r-1)} s \, \alpha_p + \sum_{p \in R_k(r-1)} c_{\bar{\imath}(p)} \alpha_p$$

 ${\bf Claim \ 1.} \ {\rm ls}(\lambda) \leq {\rm ls}(\gamma) + s \ and \ \ {\rm rs}(\lambda) \leq {\rm rs}(\gamma) + s.$ 

**Proof:** Suppose  $ls(\lambda) > ls(\gamma) + s$ . Then  $l = ls(\lambda)$ . Take  $p \in Z_k$  such that i(p) = l. Then  $(\lambda^{(k)} - \gamma^{(k)})(\check{\alpha}_p) > 0$ . From Equation (5.5), we get:

$$(\lambda^{(k)} - \gamma^{(k)})(\check{\alpha}_p) = s - b_{l-1}$$

Thus

(a) 
$$b_{l-1} < s$$

Again for all  $q \in Z_k$  such that  $ls(\gamma) < i(q) < ls(\lambda)$ , we have  $(\lambda^{(k)} - \gamma^{(k)})(\check{\alpha}_q) = \lambda^{(k)}(\check{\alpha}_q) \ge 0$ . Using Equation (5.5) again, we get:

$$(\lambda^{(k)} - \gamma^{(k)})(\check{\alpha}_q) = 2b_{i(q)} - b_{i(q)+1} - b_{i(q)-1}$$

Hence

(**b**) 
$$b_{i(q)-1} \leq 2b_{i(q)} - b_{i(q)+1}$$

Facts (a) and (b) imply that  $\{s, b_{l-1}, b_{l-2}, \dots, b_{ls(\gamma)}\}$  is a strictly decreasing sequence of nonnegative integers. But the number of terms in the sequence is  $l - ls(\gamma) + 1 > s + 1$ , which is a contradiction. The proof for the right support is

analogous. Our claim is thus true.  $\Box$ So, the support  $\ell(\lambda) \leq \ell(\gamma) + 2s$ . Choose k large such that (i)  $\ell(\gamma) + 2s \leq d_1 + d_2 + k$  and (ii)  $\mathfrak{g}(Z_k)$  is a Kac-Moody algebra of indefinite type. This is possible, since we started with the assumption that  $(X_1, X_2) \neq (A_{d_1}, A_{d_2})$ .

By our claim above, given any  $\lambda \in U(\gamma, s)$ ,  $\lambda^{(k)}$  is a well defined dominant weight of  $\mathfrak{g}(Z_k)$ . Further we also have  $\lambda^{(k)} \geq \gamma^{(k)}$  in  $P^+(Z_k)$ , i.e,  $\lambda^{(k)} \in U(\gamma^{(k)}) \subset P^+(Z_k)$ . But Proposition 10 implies that the set  $U(\gamma^{(k)})$  is finite. This implies that the number of possible choices for  $\lambda$  is also finite.  $\Box$ 

5.3. In this subsection, we will define a commutative (associative)  $\mathbb{C}$  algebra  $\mathcal{R}(X_1|X_2)$  whose multiplication operation has the  $c_{\lambda\mu}^{\nu}(\infty)$  as structure constants.

**Definition 7.** If  $\lambda = (x, y) \in \mathcal{H}_2$ , we define

$$(\lambda)_{X_1} := \frac{\sum_i x_i a_i(X_1)}{\Delta_1}$$

and

$$(\lambda)_{X_2} := \frac{\sum_i y_i a_i(X_2)}{\Delta_2}$$

Clearly  $|\lambda|_{(X_1,X_2)} = \Delta_1 \Delta_2 ((\lambda)_{X_1} - (\lambda)_{X_2})$  and if  $|\lambda|_{(X_1,X_2)} = 0$ , then dep $(\lambda) = (\lambda)_{X_1} = (\lambda)_{X_2}$ .

Let  $\mathcal{R}$  be the  $\mathbb{C}$  vector space with basis  $\{v_{\lambda} : \lambda \in \mathcal{H}_{2}^{+}\}$  and let  $\widehat{\mathcal{R}}$  denote its formal completion; so a typical element of  $\widehat{\mathcal{R}}$  is an infinite sum  $\xi := \sum_{\lambda \in \mathcal{H}_{2}^{+}} c_{\lambda} v_{\lambda}$ .

**Definition 8.** Let  $\mathcal{R}(X_1|X_2)$  be the set of all  $\xi := \sum_{\lambda \in \mathcal{H}_2^+} c_\lambda v_\lambda \in \widehat{\mathcal{R}}$  which satisfy

the following two conditions:

- (1)  $\{|\lambda|_{(X_1,X_2)} : c_\lambda \neq 0\}$  is a finite set
- (2)  $\{(\lambda)_{X_1} : c_{\lambda} \neq 0\} \subset \frac{1}{\Delta_1} \mathbb{Z}$  is bounded above.

Remark 7. Since

$$(\lambda)_{X_2} = (\lambda)_{X_1} - \frac{|\lambda|_{(X_1, X_2)}}{\Delta_1 \Delta_2}$$

the two conditions above imply that  $\{(\lambda)_{X_2} : c_\lambda \neq 0\} \subset \frac{1}{\Delta_2}\mathbb{Z}$  is also bounded above.

5.4. The operation \* on  $\mathcal{R}(X_1|X_2)$ . STEP 1: Observe that  $v_{\lambda}, v_{\mu} \in \mathcal{R}(X_1|X_2)$  for all  $\lambda, \mu \in \mathcal{H}_2^+$ . Define

$$v_{\lambda} * v_{\mu} := \sum_{\nu \in \mathcal{H}_2^+} c_{\lambda\mu}^{\nu}(\infty) \, v_{\nu}$$

The right hand side is in  $\mathcal{R}(X_1|X_2)$  since  $c_{\lambda\mu}^{\nu}(\infty) > 0$  implies  $\lambda + \mu \succeq \nu$ , i.e,

(5.6) 
$$|\nu|_{(X_1,X_2)} = |\lambda|_{(X_1,X_2)} + |\mu|_{(X_1,X_2)}$$
 and

(5.7) 
$$\deg(\lambda + \mu - \nu) = (\lambda)_{X_1} + (\mu)_{X_1} - (\nu)_{X_1} \ge 0$$

**STEP 2:** Given  $\xi := \sum_{\lambda \in \mathcal{H}_2^+} c_\lambda v_\lambda$  and  $\eta := \sum_{\mu \in \mathcal{H}_2^+} d_\mu v_\mu$  in  $\mathcal{R}(X_1|X_2)$ , we define  $\xi * \eta$  by bilinearity, i.e,

(5.8) 
$$\xi * \eta := \sum_{\lambda} \sum_{\mu} c_{\lambda} d_{\mu} (v_{\lambda} * v_{\mu})$$
$$= \sum_{\lambda} \sum_{\mu} \sum_{\nu} c_{\lambda} d_{\mu} c_{\lambda\mu}^{\nu}(\infty) v_{\nu}$$

We need to show that (5.8) is well defined. More precisely, we need to show the following:

**Proposition 12.** Let  $\xi := \sum_{\lambda \in \mathcal{H}_2^+} c_\lambda v_\lambda$  and  $\eta := \sum_{\mu \in \mathcal{H}_2^+} d_\mu v_\mu$  be elements of  $\mathcal{R}(X_1|X_2)$ . Given  $\nu \in \mathcal{H}_2^+$ , the set

$$F_{\nu} := \{ (\lambda, \mu) \in \mathcal{H}_2^+ \times \mathcal{H}_2^+ : c_{\lambda} \neq 0, d_{\mu} \neq 0, c_{\lambda\mu}^{\nu}(\infty) \neq 0 \}$$

is finite.

**Proof:** As before,  $c_{\lambda\mu}^{\nu}(\infty) > 0$  gives us Equations (5.6), (5.7). Since  $\xi, \eta \in \mathcal{R}(X_1|X_2)$ , the sets  $\{(\lambda)_{X_1} : c_{\lambda} \neq 0\}$  and  $\{(\mu)_{X_1} : d_{\mu} \neq 0\}$  are bounded above. Together with Equation (5.7) this implies that these sets are also bounded below; so in fact both sets must be finite. So  $dep(\lambda + \mu - \nu) = (\lambda)_{X_1} + (\mu)_{X_1} - (\nu)_{X_1}$  takes only finitely many values  $s_1, s_2, \cdots, s_p$  say. This means that

$$\lambda + \mu \in \bigcup_{i=1}^p U(\nu, s_i)$$

By Proposition 11 the right hand side is a finite set. Thus  $F_{\nu}$  is finite too. Thus \* is a well defined operation on  $\mathcal{R}(X_1|X_2)$ .

**Remark 8.** Let X be an extensible diagram; definition (5.5) of [4] introduced a  $\mathbb{C}$  algebra  $\Lambda^X$  which is a subspace of  $\widehat{\mathcal{R}}$ . From its definition, it is clear that  $\Lambda^X$  is just the subalgebra of  $\mathcal{R}(X|A_1)$  generated by the  $\{v_{\lambda} : \lambda \in \mathcal{H}_2^+\}$ .

5.5. Associativity. In order to show that \* is associative, we need to show that for all  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ ,  $(v_\lambda * v_\mu) * v_\nu = v_\lambda * (v_\mu * v_\nu)$ . This was proved in [4] (see equation (5.10) there) for the X(m) series. The proof in [4] uses the so called *Interval stabilization lemma*. We state the corresponding lemma for our context:

**Lemma 8.** Let  $\lambda_1, \lambda_2 \in \mathcal{H}_2^+$  with  $\lambda_1 \succcurlyeq \lambda_2$ . Let  $I(\lambda_1, \lambda_2) := \{\gamma \in \mathcal{H}_2^+ : \lambda_1 \succcurlyeq \gamma \succcurlyeq \lambda_2\}$  and  $I^{(k)}(\lambda_1, \lambda_2) := \{\beta \in P^+(Z_k) : \lambda_1^{(k)} \ge \beta \ge \lambda_2^{(k)}\}$  for k large. Then

- (1)  $I(\lambda_1, \lambda_2)$  is a finite set
- (2) There exists N such that for all  $k \ge N$ ,  $I^{(k)}(\lambda_1, \lambda_2) = \{\gamma^{(k)} : \gamma \in I(\lambda_1, \lambda_2)\}$



FIGURE 1.  $B_n, C_n, D_n$ 

The proof is exactly the same as the proof of Lemma (5.1) of [4]; instead of equation (5.1) of [4], we use our equation (5.1).  $\Box$ 

Now, given  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ , we fix  $\gamma \in \mathcal{H}_2^+$  and let  $c_{\lambda\mu\nu}^{\gamma}(k)$  denote the multiplicity of the representation  $L(\gamma^{(k)})$  in the triple tensor product  $L(\lambda^{(k)}) \otimes L(\mu^{(k)}) \otimes L(\nu^{(k)})$ . If this multiplicity becomes a constant for large k, we denote the constant value by  $c_{\lambda\mu\nu}^{\gamma}(\infty)$ . It is easily seen that the associativity of \* is implied by the following lemma:

**Lemma 9.** The multiplicity  $c_{\lambda\mu\nu}^{\gamma}(k)$  does become constant for large k and we have

$$c_{\lambda\mu\nu}^{\,\gamma}(\infty) = \sum_{\delta \in \mathcal{H}_2^+} c_{\lambda\mu}^{\,\delta}(\infty) \, c_{\delta\nu}^{\,\gamma}(\infty) = \sum_{\delta \in \mathcal{H}_2^+} c_{\lambda\delta}^{\,\gamma}(\infty) \, c_{\mu\nu}^{\,\delta}(\infty)$$

The above relation is just equation (5.8) of [4]. The proof given there carries over with no change.  $\Box$ 

We call  $\mathcal{R}(X_1|X_2)$  with the operation \* the *stable representation ring* for the pair  $(X_1, X_2)$ .

## 6. BCD DIAGRAMS

Having dealt with extensible pairs in sections 3 - 5, we now turn to the classical  $B_n, C_n, D_n$  sequences. These fall outside the class of extensible sequences. Our goal is to give simple characterizations of weights  $\lambda, \mu, \nu, \beta$  which are of the form of Theorem 1 and hence exhibit stabilization behavior.

Let X be one of the diagrams  $B_3, C_3, D_3$  below :

- (1)  $B_3: \begin{array}{c} & \bullet \\ 1 & 2 & 3 \\ & \bullet \\ & \bullet \end{array}$
- (2)  $C_3$ :
- (3)  $D_3$ : <sup>1</sup> • • • •

The nodes marked by the extra circles are taken to be distinguished. Then  $X(m) = Y_{m+3}$  where  $Y \in \{B, C, D\}$  (see figure 1).

It is an easy fact that  $\det(B_n) = 2 = \det(C_n)$  and  $\det(D_n) = 4$  for all  $n \ge 3$ . So  $\Delta_X = 0$  and X is not an extensible diagram. In fact the proofs of the results of [4] often used the fact that for extensible X,  $\Delta_X \neq 0$  and hence  $|\det X(m)| \to \infty$  as  $m \to \infty$ .

Recall the definitions of the sets  $\mathcal{H}_1, \mathcal{H}_1^+, \mathcal{H}_2, \mathcal{H}_2^+$  etc from section 4.1. For  $Y \in \{B, C, D\}$ , we let  $\omega_i^{(n)}$  (resp  $\alpha_i^{(n)}$ ) be the fundamental weight (resp. simple root) corresponding to the  $i^{th}$  node of  $Y_n$  in figure 1. We let  $\bar{\omega}_i^{(n)} := \omega_{n-i+1}^{(n)}$  and  $\bar{\alpha}_i^{(n)} :=$ 

$$\alpha_{n-i+1}^{(n)}$$
. Given  $\gamma = (x, y) \in \mathcal{H}_2$  and  $n \ge \ell(x) + \ell(y)$ , let  
 $\gamma^{(n)} := \sum_i x_i \, \omega_i^{(n)} + \sum_i y_i \, \bar{\omega}_i^{(n)}$ 

For the series  $B_n, C_n, D_n$ , our aim is to characterize  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ ,  $\beta \in \mathcal{H}_2$  which have the specific form of Theorem 1.

Let  $\gamma = (x, y) \in \mathcal{H}_2$ ; set  $|\gamma| := \sum_i i y_i$ . We let  $l := \max\{i : x_i \neq 0\}$  and  $r := \max\{i : y_i \neq 0\}$ .

6.1.  $B_n.\ {\rm First},$  we state the following lemma concerning the fundamental weights of  $B_n$ 

Lemma 10. (1) Let 
$$1 \le i < n$$
. If  $\bar{\omega}_i^{(n)} = \sum_{j=1}^n c_{ij}(n) \bar{\alpha}_j^{(n)}$ , then  $c_{ij}(n) = i$  for  
 $i \le j \le n$ .  
(2)  $2\omega_1^{(n)} = \sum_{j=1}^n j \bar{\alpha}_j^{(n)}$ .  
(3) Let  $1 < i \le n$ . If  $\omega_i^{(n)} = \sum_{j=1}^n d_{ij}(n) \bar{\alpha}_j^{(n)}$ , then  $d_{ij}(n) = j$  for  $1 \le j \le (n-i+1)$ .

**Proof:** Follows by direct computation.

For  $\gamma$  as above, define

$$\operatorname{ht}_B(\gamma) := \frac{x_1}{2} + \sum_{i>1} x_i$$

Suppose we write

$$\gamma^{(n)} = \sum_{i=1}^{n} p_i(n) \,\bar{\alpha}_i^{(n)} \text{ with } p_i(n) \in \mathbb{Z},$$

lemma 10 implies

(6.1) 
$$p_i(n) = \sum_j j y_j + i \operatorname{ht}_B(\gamma) \quad \text{for } r \le i \le (n - l + 1)$$

We note that  $\gamma^{(n)} \in Q^+(B_n)$  for all large *n* implies that  $\operatorname{ht}_B(\gamma) \in \mathbb{Z}^{\geq 0}$ . Observe that  $p_i(n)$  grows linearly as a function of *i* unless  $\operatorname{ht}_B(\gamma) = 0$ , in which case

$$p_i(n) = p_j(n) = |\gamma| \text{ for } r \le i, j \le (n-l+1)$$

This is exactly the requirement of Theorem 1. To rephrase this in the notations of section 2.4, identify  $B_l$  and  $A_r$  with the subdiagrams of  $B_n$  formed by the leftmost l and the rightmost r nodes (see figure 1). Then equation (6.1) shows that

 $\exists s \in \mathbb{Z}$  such that  $\gamma \in R_s(B_l, A_r) \Leftrightarrow \operatorname{ht}_B(\gamma) = 0$ 

In this case  $s = |\gamma|$ . Theorem 3 now implies:

**Proposition 13.** Consider the sequence  $B_n$ ; let  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ ,  $\beta \in \mathcal{H}_2$ .

- (1) If  $ht_B(\lambda) + ht_B(\mu) = ht_B(\nu)$ ,  $c^{\nu}_{\lambda\mu}(n)$  stabilizes.
- (2) If  $ht_B(\lambda) = ht_B(\beta)$ ,  $b_{\lambda\beta}(n)$  stabilizes.

**Remark 9.** From the remarks of the above paragraph we conclude that if  $c_{\lambda\mu}^{\nu}(n) > 0$ (resp.  $b_{\lambda\beta}(n) > 0$ ) for all large n, then  $\operatorname{ht}_B(\lambda) + \operatorname{ht}_B(\mu) \ge \operatorname{ht}_B(\nu)$  (resp.  $\operatorname{ht}_B(\lambda) \ge \operatorname{ht}_B(\beta)$ ). Thus our theorem above deals with the  $\nu$  and  $\beta$  which have the maximum allowed height, and shows that stabilization holds in this case.

**Corollary 3.** Let  $\lambda$  be a weight supported on the  $A_r$  portion of the diagram of  $B_n$ , i.e, let  $\lambda = (0, y)$  for some  $y \in \mathcal{H}_1^+$ . Then  $ht_B(\lambda) = 0$ . If  $\mu, \nu, \beta$  are also of this form, then the height compatibility conditions of the above proposition are trivially satisfied. So, for all such weights of the  $B_n$ 's, tensor product and branching multiplicities stabilize.

**Example 3.** For this example, we will use the following indexing scheme: let  $[m_1 m_2 \cdots m_n]$  denote the irreducible representation of  $B_n$  with highest weight  $\sum_{i=1}^n m_i \omega_i^{(n)}$ .

Consider the representation  $[100\cdots01]$  of  $B_n$ . Its highest weight is obtained as  $\lambda^{(n)}$  where  $\lambda := (x, y)$  with  $x_1 = y_1 := 1$  and  $x_i = y_i = 0$   $\forall i > 1$ . Observe that  $\operatorname{ht}_B(\lambda) = 1/2$ . We give below the decomposition of the tensor square of this representation in  $B_3, B_4$  and  $B_5$ . Note that our theorem above guarantees stabilization of multiplicities for  $\nu$  for which  $\operatorname{ht}_B(\nu) = 1/2 + 1/2 = 1$ . The data below was generated using the program LiE:

(1) In 
$$B_3$$
:

$$(6.2) \qquad [101] \otimes [101] = 1.[202] + 1.[210] + 2.[201] + 2.[200] + 1.[012] + 2.[011] + 2.[010] + 1.[001] + 1.[020] + 1.[003] + 1.[002] + 1.[000]$$

(2) In  $B_4$ :

$$[1001] \otimes [1001] = 1.[2002] + 1.[2010] + 2.[2001] + 2.[2000] (6.3) + 1.[0102] + 1.[0110] + 2.[0101] + 2.[0100] + 1.[0012] + 2.[0011] + 2.[0010] + 1.[0020] + 1.[0003] + 1.[0002] + 1.[0001] + 1.[0000]$$

(3) In  $B_5$ :

$$\begin{split} [10001] \otimes [10001] &= 1.[20002] + 1.[20010] + 2.[20001] + 2.[20000] \\ (6.4) \\ &+ 1.[01002] + 1.[01010] + 2.[01001] + 2.[01000] \end{split}$$

$$+1.[00102] + 2.[00101] + 2.[00100] + 1.[00110]$$

+2.[00011] + 1.[00002] + 2.[00010] + 1.[00012] + 1.[00020] + 1.[00003] + 1.[00001] + 1.[00000]

In each case, the right hand side is arranged so that the weights  $\nu$  in the first three rows satisfy  $\operatorname{ht}_B(\nu) = 1$ . Suppose we look at such  $\nu \in \mathcal{H}_2^+$  which make sense in  $B_3, B_4$  and  $B_5$ ; from the data we do see that the multiplicities of these are the same for  $B_n$ , n = 3, 4, 5. The last rows consists of  $\nu$  for which  $\operatorname{ht}_B(\nu) = 0$ .

6.2.  $C_n$ ,  $D_n$ . Define  $\operatorname{ht}_C(\gamma) = \sum_i x_i$ . The analogous statements for  $C_n$  are contained in the following:

**Proposition 14.** (1)  $\exists s \in \mathbb{Z}$  such that  $\gamma \in R_s(C_l, A_r) \Leftrightarrow ht_C(\gamma) = 0$ , in which case  $s = |\gamma|$ .

- (2) Given  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ ,  $\operatorname{ht}_C(\lambda) + \operatorname{ht}_C(\mu) = \operatorname{ht}_C(\nu) \Rightarrow c_{\lambda\mu}^{\nu}(n)$  stabilizes.
- (3) Given  $\beta \in \mathcal{H}_2$  with  $\operatorname{ht}_C(\lambda) = \operatorname{ht}_C(\beta)$ , then  $b_{\lambda\beta}(n)$  stabilizes.

**Proof:** This again follows from the expression for the fundamental weights  $\omega_i^{(n)}$  and  $\bar{\omega}_i^{(n)}$  of  $C_n$  in the basis of simple roots.

For type D, the height function turns out to be :

$$ht_D(\gamma) := \frac{x_1}{2} + \frac{x_2}{2} + \sum_{i>2} x_i$$

The three assertions of proposition 14 are true for  $D_n$  if we replace  $C_l$  by  $D_l$  and  $ht_C$  by  $ht_D$ . We leave the remaining details to the reader.

- **Remark 10.** (1) For one-sided weights  $\lambda, \mu, \nu, \beta$  as in corollary 3, we evidently obtain stabilization in types C and D as well.
  - (2) We saw above that the p<sub>i</sub>(n) in general grows linearly with i. Our proof of stabilization however only works when the p<sub>i</sub>(n)'s are constant. Hence we only deduce stabilization under the height compatibility condition of propositions 13 and 14.
  - (3) However, from computer generated data of multiplicities in tensor products, it appears that stabilization still holds, even when the  $p_i(n)$ 's are non constant in the middle. Our method of proof however fails in this situation.

The last remark leads us to the following conjecture:

**Conjecture:** For types  $B, C, D, c_{\lambda\mu}^{\nu}(n)$  stabilizes for all triples  $\lambda, \mu, \nu \in \mathcal{H}_2^+$ .

It seems likely that a similar statement holds for the branching multiplicities as well.

6.3. Final remarks. We briefly mention two other papers that are related to the work of the present article.

In [1], Benkart, Kang, Lee, Misra and Shin considered the affine diagrams  $A_n^{(1)}$ and looked at integral weights of a fixed level that are parametrized by  $\mathcal{H}_2 \times \mathbb{Z}$  (the extra degree of freedom comes from the fact that the Cartan subalgebra of  $A_n^{(1)}$ is n+2 dimensional, and so a weight is not uniquely determined by specifying its values on the simple coroots alone). Given a level zero element  $\gamma \in \mathcal{H}_2 \times \mathbb{Z}$ , they worked out the condition that  $\gamma$  must satisfy so that  $\gamma^{(n)} \in Q(A_n^{(1)})$  for all large n. They show that such  $\gamma$ 's must have the form as in Theorem 1 (see proposition 1.22 of [1]).

Our proof of stabilization then applies (even though  $A_n^{(1)}$  is not strictly of the form  $Z_k(X_1, X_2)$  for any  $X_1, X_2$ ). So for  $A_n^{(1)}$  we obtain stabilization of  $c_{\lambda\mu}^{\nu}(n)$  and  $b_{\lambda\beta}(n)$  for all  $\lambda, \mu, \nu, \beta$ , subject to a compatibility condition on their levels.

In the present work, we have not mentioned an important representation theoretic statistic - weight multiplicities. The main objective of [1] (and its follow up paper [2]) was to analyze the behavior of these numbers as  $k \to \infty$ . In our notation, given  $\lambda \in \mathcal{H}_2^+$  and  $\beta \in \mathcal{H}_2$ , let  $m_{Z_k}(\lambda, \beta)$  be the dimension of the weight space with weight  $\beta^{(k)}$  in the representation  $L(\lambda^{(k)})$  of  $\mathfrak{g}(Z_k)$ . These papers show that for the case when  $Z_k$  is either  $A_n$  or  $A_n^{(1)}$ ,  $m_{Z_k}(\lambda, \beta)$  is a polynomial in k. They also consider other classical and affine sequences and establish this polynomiality under the assumption that the weights  $\lambda$  and  $\beta$  are "one-sided" in the sense of corollary 3.

As mentioned in the introduction, Webster [9] has recently proved more general versions of the stabilization results of the present article using quiver varieties and their connections to representation theory. He has also proved the polynomiality of weight multiplicities for more general  $Z_k$ 's, thereby extending the result of [2]. In fact [9] uses the result of [2] for the  $A_k$  to prove it for all  $Z_k$ 's. However the method originally used in [2] is very different in flavor than that in [9].

## References

- G. Benkart, S-J. Kang, H. Lee, K.C. Misra, and D-U. Shin, The polynomial behavior of weight multiplicities for the affine Kac-Moody algebras A<sub>r</sub><sup>(1)</sup>, Compositio Math. **126** (2001), no. 1, 91–111.
- G. Benkart, S-J. Kang, H. Lee, and D-U. Shin, *The polynomial behavior of weight multiplicities for classical simple Lie algebras and classical affine Kac-Moody algebras*, Recent Developments in Quantum Affine Algebras and Related Topics (Providence) (N. Jing and K. C. Misra, eds.), Contemp. Math., vol. 248, Amer. Math. Soc., 1999, pp. 1–29.
- 3. V.G. Kac, Infinite dimensional Lie algebras, third ed., Cambridge University Press, 1990.
- M. Kleber and S. Viswanath, Tensor product stabilization in Kac-Moody algebras, Advances in Mathematics 201 (2006), no. 1, 1–35.
- P. Littelmann, The path model for representations of symmetrizable Kac-Moody algebras, Proceedings of the International Congress of Mathematicians (Zürich 1994), vol. 1,2, Birkhäuser, Basel, 1995, pp. 298–308.
- A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras, Invent. Math. 116 (1994), no. 1-3, 329–346.
- 7. \_\_\_\_, Paths and root operators in representation theory, Ann. of Math 142 (1995), no. 3, 499–525.
- J. R. Stembridge, Combinatorial models for Weyl characters, Adv. Math. 168 (2002), no. 1, 96–131.
- B. Webster, Stabilization phenomena in Kac-Moody algebras and quiver varieties, arXiv:math.RT/0505619, 2005.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CA 95616, USA *E-mail address:* svis@math.ucdavis.edu