

## FREE CONSTRUCTION OF CAT(1) SPACES

KOICHI NAGANO

ABSTRACT. We construct several non-trivial examples of CAT(1) spaces by using the idea of free construction.

## 1. INTRODUCTION

A spherical (resp. Euclidean) building  $G$  of dimension  $n$  has plenty of apartments; more precisely, for any two points in  $G$  there exists an isometric embedding from the standard unit  $n$ -sphere (resp. the Euclidean  $n$ -space) to  $G$  whose image contains the given two points. Spherical or Euclidean buildings satisfy other rigid metric properties in the context of the geometry of CAT(1) spaces (cf. [KL]). In [CL, L, BL], we see several metric characterizations of buildings.

We say that a CAT(1) space is  $n$ -round if any two points are contained in an isometrically embedded unit  $n$ -sphere. For instance, every spherical building of dimension  $n$  is  $n$ -round. Every  $\pi_{\mathbb{R}}$ -gon in the sense of [B] is 1-round. Every hemispherex of dimension  $n \geq 2$  in [BB] is  $(n - 1)$ -round (cf. Example 2.3). In general, every  $n$ -round space is geodesically complete and  $(n - 1)$ -round, and it has diameter  $\pi$ .

A geodesically complete  $\mathbb{R}$ -tree with no graph structure has plenty of lines. This example suggests that only from the property of plenty of apartments one can not derive the rigid structure of buildings. Nevertheless, it seems to be non-trivial to see whether an  $n$ -round space of dimension  $n$  without building structure exists. The main purpose of this paper is to construct an  $n$ -round space of dimension  $n$  that is not a spherical building.

The Reshetnyak gluing theorem helps us to obtain various examples of CAT(1) spaces. For our purpose, we need another idea. Roughly speaking, the purpose is achieved by mixing the idea of gluing with that of free construction. The idea of free construction was used by Tits [T] to obtain generalized polygons with large automorphism group.

For a CAT(1) space  $X$ , we denote by  $\dim X$  the geometric dimension of  $X$  in the sense of [K]. By using the idea of free construction, we obtain the following:

---

*Date:* May 28, 2005.

*1991 Mathematics Subject Classification.* [.

*Key words and phrases.* CAT(1) space.

Partially supported by the 2004 JSPS Postdoctoral Fellowships for Research Abroad.

**Theorem 1.1.** *Let  $(X_\lambda)_{\lambda \in \Lambda}$  be a family of CAT(1) spaces of diameter  $\leq \pi$ . Then for any  $n$  there exists an  $n$ -round space  $Y$  such that*

- (1)  $\dim Y = \max\{n, \sup_{\lambda \in \Lambda} \dim X_\lambda\}$ ;
- (2) *all points in  $Y$  are non-manifold points*;
- (3) *each  $X_\lambda$  is isometrically embedded in  $Y$ .*

*In particular, there exists an  $n$ -round space of dimension  $n$  that is not a spherical building.*

By our free construction, we also see the following. For a fixed  $l \geq \pi$ , let  $(X_\lambda)_{\lambda \in \Lambda}$  be a family of CAT(1) spaces of diameter  $\leq l$ . Then for any  $n$  there exists a CAT(1) space in which any two points are contained in an isometrically embedded standard  $n$ -sphere of diameter  $l$ , and in which each  $X_\lambda$  is isometrically embedded (Theorem 3.3).

Next we consider the following condition. We say that a CAT(1) space is  $n$ -rolling if it has at least one pair of antipodes and if any antipodes are contained in an isometrically embedded unit  $n$ -sphere, where antipodes mean two points that have distance  $\geq \pi$ . Every  $n$ -round space is  $n$ -rolling. The unit  $n$ -hemisphere is  $(n-1)$ -rolling. Every  $n$ -rolling space is geodesic and  $(n-1)$ -rolling, and it has diameter  $\pi$ . If an  $n$ -rolling space has dimension  $n$ , then it is geodesically complete (cf. [BL]). It was proved in [BL] that if an  $n$ -rolling space of dimension  $n$  contains a relatively compact open set, then it is a spherical building.

A CAT(1) space has dimension 1 if and only if it is locally an  $\mathbb{R}$ -tree of injectivity radius  $\geq \pi$ . A CAT(1) space of dimension 1 is 1-rolling if and only if it is 1-round. In general, we have:

**Theorem 1.2.** *For any  $n \geq 2$ , there exists an  $n$ -rolling space of dimension  $n$  that is not  $n$ -round.*

In fact, for any  $n \geq 2$ , such  $n$ -rolling spaces of dimension  $n$  can be obtained abundantly (Proposition 3.2).

We remark that the free construction discussed below also works in the CAT( $\kappa$ ) setting. The details are omitted in this paper.

We mention the relation between our notion and the asymptotic geometry of CAT(0) spaces. If  $H$  is a locally compact, geodesically complete CAT(0) space of dimension  $n$ , and if its Tits ideal boundary is  $(n-1)$ -rolling, then  $H$  is a Euclidean building. This is substantially a corollary of the works of [CL, L, BL] (cf. Section 7 in [BL]).

## 2. PRELIMINARIES

**2.1. CAT(1) spaces.** A minimizing geodesic in a metric space means a length-minimizing curve joining two points. Any single point is also assumed to be a minimizing geodesic. A metric space is said to be *geodesic* if any two points can be joined by a minimizing geodesic. A locally geodesic space is *geodesically complete* if any minimizing geodesic is contained in a bi-infinite locally minimizing geodesic.

We denote by  $\mathbb{S}^n$  the standard unit  $n$ -sphere.

A complete metric space  $X$  with distance  $d$  is said to be CAT(1) if any two points  $x, y \in X$  with  $d(x, y) < \pi$  can be joined by a minimizing geodesic and if any geodesic triangle in  $X$  with perimeter  $< 2\pi$  is thinner than its comparison triangle in  $\mathbb{S}^2$ . If a CAT(1) space  $X$  is geodesically complete, then for any  $x \in X$  we have a point  $y \in X$  with  $d(x, y) \geq \pi$ . We refer to [BH] for the basics of CAT(1) spaces.

Let  $X$  be a CAT(1) space. For  $x, y, z \in X$  with  $d(x, y), d(x, z) < \pi$ , the angle at  $x$  between  $y$  and  $z$  is denoted by  $\angle_x(y, z)$ . The upper semi-continuity of angles holds for not only the usual convergences but the ultralimits. For  $x \in X$ , the space of directions at  $x$  equipped with the interior distance induced from  $\angle_x$  is denoted by  $\Sigma_x X$ . This is also a CAT(1) space. The *geometric dimension*  $\dim X$  of  $X$  is defined inductively as follows: if  $X$  is discrete, then  $\dim X := 0$ ; in the other case,  $\dim X := 1 + \sup_{x \in X} \dim \Sigma_x X$ . Some geometric study of the geometric dimension is seen in [K].

We say that a point  $x \in X$  is *strongly singular in  $X$*  if  $\Sigma_x X$  has at least two connected components. We denote by  $S(X)$  the set of all non-manifold points in  $X$ . Every manifold point in a CAT(1) space has the space of directions consisting of a single connected component. If  $x$  is strongly singular in  $X$ , then  $x \in S(X)$ .

**2.2. Poles.** Let  $X$  be a geodesically complete CAT(1) space of dimension  $\geq 2$ . A point  $p \in X$  with  $\dim \Sigma_p X \geq 1$  is called a *pole* if the diameter of  $\Sigma_p X$  is greater than  $\pi$ . A point  $p \in X$  with  $\dim \Sigma_p X \geq 1$  is a pole if and only if for some  $\epsilon > 0$  we have a minimizing geodesic of positive length  $< \pi$  through  $p$  with endpoints  $x$  and  $y$  such that for any  $q \in X$  with  $d(p, q) < \pi$  either (1)  $\pi + \epsilon \leq \angle_p(x, q) + \angle_p(y, q)$ , or (2)  $\angle_p(x, q) = \pi$  or  $\angle_p(y, q) = \pi$ . If  $X$  is  $n$ -round for  $n \geq 2$ , then it has no pole.

**Example 2.1.** For  $n \geq 2$ , the spherical suspension over a standard  $(n - 1)$ -sphere of diameter  $> \pi$  is a geodesically complete CAT(1) space of dimension  $n$  and diameter  $\pi$ . The suspension points are poles.

**Example 2.2.** Let  $n \geq 2$ , and let  $p, q \in \mathbb{S}^n$  be antipodes. Take some antipodes  $\bar{p}, \bar{q}$  in another unit  $n$ -sphere  $\mathbb{S}^n$ . Let  $X$  be the quotient metric space  $\mathbb{S}^n \sqcup \mathbb{S}^n / \sim$  made by attaching  $\mathbb{S}^n$  to  $\mathbb{S}^n$  at  $p = \bar{p}$  and  $q = \bar{q}$ . This is a geodesically complete CAT(1) space of dimension  $n$  and diameter  $\pi$ . The antipodes  $p$  and  $q$  in  $X$  are poles.

**Example 2.3.** Let  $n \geq 2$ . A spherical simplicial complex is called an  *$n$ -hemispherex* if it is obtained from  $\mathbb{S}^n$  by attaching unit  $n$ -hemispheres along great-hyperspheres such that no pair of antipodes in the original  $\mathbb{S}^n$  belongs to all hyperspheres ([BB]). Every  $n$ -hemispherex is  $(n - 1)$ -round, not  $n$ -rolling, and has a pole.

**2.3. Inductive limits and their ultraproducts.** For two metric spaces  $X$  and  $Y$ , we write  $X < Y$  if and only if we have an isometric embedding from  $X$  to  $Y$ . We say that a map  $f: X \rightarrow Y$  is a  $\pi$ -truncated isometric embedding if for any  $x, y \in X$  with  $d(x, y) \leq \pi$  we have  $d(f(x), f(y)) = d(x, y)$ . We write  $X \prec Y$  if and only if we have a  $\pi$ -truncated isometric embedding from  $X$  to  $Y$ .

We fix a non-principal ultrafilter  $\omega$  on  $\mathbb{N}$ . Let  $Y$  be a metric space. We denote by  $Y^\omega$  the ultraproduct of  $Y$  defined as the ultralimit of  $(Y_i)$  with respect to  $\omega$  for the constant sequence  $Y_i = Y$ . Note that  $Y^\omega$  is complete, and  $Y < Y^\omega$ . If  $Y$  is geodesically complete, then so is  $Y^\omega$ . If  $Y$  is a CAT(1) space of  $\dim Y = n$ , then  $\dim Y^\omega = n$  (cf. [L]).

Let  $Y_0 \prec Y_1 \prec \cdots \prec Y_{i-1} \prec \cdots$  be an inductive system of metric spaces  $Y_{i-1}$  with distance  $d_{i-1}$ ,  $i \in \mathbb{N}$ , and  $\pi$ -truncated isometric embeddings from  $Y_{i-1}$  to  $Y_i$ . We denote by  $Y_\infty := \bigcup_{i \in \mathbb{N}} Y_{i-1}$  the inductive limit space with the distance  $d$  defined by  $d|_{Y_{i-1} \times Y_{i-1}} := \min\{\pi, d_{i-1}\}$ . We call  $Y_\infty$  the  $\pi$ -truncated inductive limit. If  $Y_i$  is geodesically complete for each  $i$ , then so is  $Y_\infty$ . We need the following later on, which follows from a standard argument.

**Lemma 2.4.** *For a sequence of CAT(1) spaces  $Y_{i-1}$  of  $\dim Y_{i-1} \leq m$  with  $Y_{i-1} \prec Y_i$ ,  $i \in \mathbb{N}$ , let  $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$  be the  $\pi$ -truncated inductive limit, and  $(Y_\infty)^\omega$  the ultraproduct of  $Y_\infty$ . Then  $(Y_\infty)^\omega$  is a CAT(1) space of  $\dim(Y_\infty)^\omega \leq m$ . Moreover, if in addition  $Y_i$  is geodesically complete for each  $i$ , then we have the following:*

- (1) *if each point in  $Y_\infty$  is strongly singular in some  $Y_i$ , then each point in  $(Y_\infty)^\omega$  is again strongly singular;*
- (2) *if any two points in  $Y_\infty$  are contained in an isometrically embedded unit  $n$ -sphere, then  $(Y_\infty)^\omega$  is  $n$ -round;*
- (3) *if any antipodes in  $Y_\infty$  are contained in an isometrically embedded unit  $n$ -sphere, then  $(Y_\infty)^\omega$  is  $n$ -rolling.*

### 3. CONSTRUCTION

We prove Theorems 1.1 and 1.2.

**3.1. Round spaces.** For a CAT(1) space  $X$ , we denote by  $\Gamma_X$  the set of all minimizing geodesics in  $X$  of length  $\leq \pi$ .

By using the idea of free construction, we first show the following:

**Proposition 3.1.** *Let  $X$  be a CAT(1) space of diameter  $\leq \pi$ . Then for any  $n$  there exists an  $n$ -round space  $Y$  with  $\dim Y = \max\{n, \dim X\}$ ,  $Y = S(Y)$ , and  $X < Y$ .*

*Proof.* For a fixed  $n$ , let  $m := \max\{n, \dim X\}$ . We construct a CAT(1) space  $Y$  inductively as follows. Put  $Y_0 := X$ . Let  $Y_{i-1}$  be a CAT(1) space of  $\dim Y_{i-1} \leq m$ . Set  $\Gamma_{i-1} := \Gamma_{Y_{i-1}}$ . For each  $\gamma \in \Gamma_{i-1}$ , we take a minimizing geodesic  $\bar{\gamma}$  in  $\mathbb{S}^n = \mathbb{S}^n(\gamma)$  of the same length. We define  $Y_i$  as the quotient metric space  $Y_{i-1} \sqcup (\bigsqcup_{\gamma \in \Gamma_{i-1}} \mathbb{S}^n(\gamma)) / \sim$  made by gluing

$\mathbb{S}^n(\gamma)$  to  $Y_{i-1}$  along  $\gamma = \bar{\gamma}$ . For each  $i \in \mathbb{N}$ ,  $Y_i$  is a geodesically complete, geodesic space. By the Reshetnyak gluing theorem,  $Y_i$  is CAT(1). It follows that  $\dim Y_i = m$ . Let  $f_i: Y_{i-1} \rightarrow Y_i$  be the inclusion. This is a  $\pi$ -truncated isometric embedding, and hence  $Y_{i-1} \prec Y_i$ . For any  $x, y \in Y_{i-1}$ ,  $f_i(x)$  and  $f_i(y)$  are contained in an embedded unit  $n$ -sphere. For any  $z \in Y_{i-1}$ ,  $f_i(z)$  is strongly singular in  $Y_i$ . Let  $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$  be the  $\pi$ -truncated inductive limit. Each point in  $Y_\infty$  is strongly singular in some  $Y_i$ , and any two points are contained in an embedded unit  $n$ -sphere. Let  $Y := (Y_\infty)^\omega$  be the ultraproduct of  $Y_\infty$ . By Lemma 2.4,  $Y$  is an  $n$ -round space with  $\dim Y = m$ ,  $Y = S(Y)$ , and  $X < Y$ .  $\square$

We remark that each  $Y_i$  has diameter  $> \pi$ . The space  $Y$  does not depend on the choices of minimizing geodesics and antipodes in  $\mathbb{S}^n$ .

Now we prove Theorem 1.1.

*Proof.* Let  $(X_\lambda)_{\lambda \in \Lambda}$  be a family of CAT(1) spaces of diameter  $\leq \pi$ . We fix  $n$ . For each  $X_\lambda$ , by Proposition 3.1 we have an  $n$ -round space  $X'_\lambda$  with  $\dim X'_\lambda = \max\{n, \dim X_\lambda\}$ ,  $X'_\lambda = S(X'_\lambda)$ , and  $X_\lambda < X'_\lambda$ .

Let  $m := \max\{n, \sup_{\lambda \in \Lambda} \dim X_\lambda\}$ . We construct a space  $Y$  inductively in the following way. Put  $Y_0 := X'_{\lambda_0}$  for some  $\lambda_0$ . Let  $Y_{i-1}$  be a geodesically complete, geodesic CAT(1) space of  $\dim Y_{i-1} \leq m$ . Set  $\Gamma_{i-1} := \Gamma_{Y_{i-1}}$ . For each  $(\gamma, \lambda) \in \Gamma_{i-1} \times \Lambda$ , take a minimizing geodesic  $\bar{\gamma}$  in  $X'_\lambda = X'_\lambda(\gamma)$  of the same length. Let

$$Y_i := Y_{i-1} \sqcup \left( \bigsqcup_{(\gamma, \lambda) \in \Gamma_{i-1} \times \Lambda} X'_\lambda(\gamma) \right) / \sim$$

be the quotient metric space made by gluing  $X'_\lambda(\gamma)$  to  $Y_{i-1}$  along  $\gamma = \bar{\gamma}$  for each  $\lambda$ . The space  $Y_i$  is a geodesically complete, geodesic CAT(1) space of  $\dim Y_i = m$ . Then  $Y_{i-1} \prec Y_i$  by the inclusion  $f_i: Y_{i-1} \rightarrow Y_i$ , and  $X'_\lambda < Y_i$  for each  $\lambda$ . For any  $x, y \in Y_{i-1}$ ,  $f_i(x)$  and  $f_i(y)$  are contained in an embedded unit  $n$ -sphere. For any  $z \in Y_{i-1}$ ,  $f_i(z)$  is strongly singular in  $Y_i$ . Let  $Y := (Y_\infty)^\omega$  be the ultraproduct of the  $\pi$ -truncated inductive limit  $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$ . By Lemma 2.4,  $Y$  is an  $n$ -round space with  $\dim Y = m$  and  $Y = S(Y)$ . Moreover,  $X_\lambda < Y$  for each  $\lambda$ .  $\square$

**3.2. Rolling spaces.** For a CAT(1) space  $X$ , we denote by  $A_X$  the subset of  $X \times X$  consisting of all pairs of antipodes in  $X$ .

To prove Theorem 1.2, it suffices to show:

**Proposition 3.2.** *Let  $n \geq 2$ , and let  $X$  be a geodesically complete, geodesic CAT(1) space of dimension  $n$  and diameter  $\pi$ . Assume that  $X$  has a pole. Then there exists an  $n$ -rolling space  $Y$  of dimension  $n$  such that  $Y = S(Y)$ ,  $X < Y$ , and  $Y$  is not  $n$ -round.*

*Proof.* We first construct a space  $Y$  as follows. Put  $Y_0 := X$ . Let  $Y_{i-1}$  be a geodesically complete, geodesic CAT(1) space of dimension  $n$ , and set  $A_{i-1} := A_{Y_{i-1}}$ . For each  $(p, q) \in A_{i-1}$ , take some antipodes  $\bar{p}, \bar{q}$  in  $\mathbb{S}^n = \mathbb{S}^n(p, q)$ . Define  $Y_i$  as  $Y_{i-1} \sqcup \left( \bigsqcup_{(p, q) \in A_{i-1}} \mathbb{S}^n(p, q) \right) / \sim$  made

by attaching  $\mathbb{S}^n(p, q)$  to  $Y_{i-1}$  at  $p = \bar{p}$  and  $q = \bar{q}$ . The space  $Y_i$  is a geodesically complete, geodesic CAT(1) space of dimension  $n$ . Then  $Y_{i-1} \prec Y_i$  by the inclusion  $f_i: Y_{i-1} \rightarrow Y_i$ . For each  $(x, y) \in A_{i-1}$ , the points  $f_i(x)$  and  $f_i(y)$  are contained in an embedded unit  $n$ -sphere. For each  $i$  and for each  $z \in Y_i$ ,  $f_{i+1}(z)$  is strongly singular in  $Y_{i+1}$ . Let  $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$  be the  $\pi$ -truncated inductive limit. Each point in  $Y_\infty$  is strongly singular in some  $Y_i$ , and any antipodes in  $Y_\infty$  are contained in an embedded unit  $n$ -sphere. Let  $Y := (Y_\infty)^\omega$  be the ultraproduct of  $Y_\infty$ . By Lemma 2.4,  $Y$  is an  $n$ -rolling space of dimension  $n$  with  $Y = S(Y)$  and  $X < Y$ . It remains to show that  $(Y_\infty)^\omega$  is not  $n$ -round.

Let  $p_0 \in X$  be a pole. For some  $\epsilon > 0$ , we have a minimizing geodesic of positive length  $< \pi$  through  $p_0$  with endpoints  $x$  and  $y$  such that for any  $q \in X$  with  $d(p_0, q) < \pi$  either (1)  $\pi + \epsilon \leq \angle_{p_0}(x, q) + \angle_{p_0}(y, q)$ , or (2)  $\angle_{p_0}(x, q) = \pi$  or  $\angle_{p_0}(y, q) = \pi$ . For the points  $p_0, x, y \in X < Y_\infty$ , the point  $(p_0) \in (Y_\infty)^\omega$  lies on the minimizing geodesic  $\gamma^\omega$  in  $(Y_\infty)^\omega$  with endpoints  $(x)$  and  $(y)$ , where  $(p_0), (x), (y)$  are the constant sequences.

Suppose that  $(Y_\infty)^\omega$  is  $n$ -round. Then  $\gamma^\omega$  is contained in an embedded unit  $n$ -sphere. Choose a point  $(q_j)$  in the unit  $n$ -sphere such that  $0 < d((p_0), (q_j)) < \pi/2$ , and both  $\angle_{(p_0)}((x), (q_j))$  and  $\angle_{(p_0)}((y), (q_j))$  are equal to  $\pi/2$ . Consider  $q_j$  with  $0 < d(p_0, q_j) < \pi/2$  for almost all  $j$ . Set  $\alpha_j := \angle_{p_0}(x, q_j)$  and  $\beta_j := \angle_{p_0}(y, q_j)$ . By the assumption and the construction, either (1)  $\pi + \epsilon \leq \alpha_j + \beta_j$ , or (2)  $\alpha_j = \pi$  or  $\beta_j = \pi$ . The upper semi-continuity of angles with respect to the ultralimits leads to a contradiction. Therefore  $(Y_\infty)^\omega$  is not  $n$ -round.  $\square$

The CAT(1) spaces in Examples 2.1, 2.2, and 2.3 satisfy the assumption in Proposition 3.2. Hence we have proved Theorem 1.2.  $\square$

**3.3. Modification.** For  $l \geq \pi$ , we say that a CAT(1) space is  $(n, l)$ -round if any two points are contained in an isometrically embedded standard  $n$ -sphere  $\mathbb{S}_l^n$  of diameter  $l$ .

Let  $X$  be a CAT(1) space. For  $l \geq \pi$ , we denote by  $\Gamma_X^l$  the set of all minimizing geodesics in  $X$  of length  $\leq l$ . In the proof of Theorem 1.1, by replacing  $\mathbb{S}^n$  (resp.  $\Gamma_X$ ) with  $\mathbb{S}_l^n$  (resp.  $\Gamma_X^l$ ), and by considering  $l$ -truncated isometric embeddings and  $l$ -truncated inductive limits instead of  $\pi$ -truncated ones, we obtain:

**Theorem 3.3.** *For a fixed  $l \geq \pi$ , let  $(X_\lambda)_{\lambda \in \Lambda}$  be a family of CAT(1) spaces of diameter  $\leq l$ . Then for any  $n$  there exists an  $(n, l)$ -round space  $Y$  such that  $\dim Y = \max\{n, \sup_{\lambda \in \Lambda} \dim X_\lambda\}$ ,  $Y = S(Y)$ , and  $X_\lambda < Y$  for each  $\lambda \in \Lambda$ .*

**Acknowledgments.** I am grateful to Alexander Lytchak for his valuable comment, to Werner Ballmann for his comment on this work, and to Linus Kramer for his information on [B]. This paper was written during my stay at the University of Bonn. I would like to thank all geometers in Bonn for their great hospitality.

## REFERENCES

- [BB] W. Ballmann and M. Brin, *Diameter rigidity of spherical polyhedra*, Duke Math. J. **122** (1999), 597–609.
- [BL] A. Balser and A. Lytchak, *Building-like spaces*, preprint, 2004; Math. ArXiv., math.MG/0410437.
- [B] C. D. Bennet, *Twin trees and  $\lambda_\Lambda$ -gons*, Trans. Amer. Math. Soc. **349** (1997), Number 5, 2069–2084.
- [BH] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Math. Wiss., Volume 319, Springer-Verlag, 1999.
- [CL] R. Charney and A. Lytchak, *Metric characterizations of spherical and Euclidean buildings*, Geom. Topol. **5** (2001), 521–550.
- [K] B. Kleiner, *The local structure of length spaces with curvature bounded above*, Math. Z. **231** (1999), 409–456.
- [KL] B. Kleiner and B. Leeb, *Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings*, Inst. Hautes Études Sci. Publ. Math. **86** (1997), 115–197 (1998).
- [L] A. Lytchak, *Rigidity of spherical buildings and spherical joins*, preprint, to appear in Geom. Funct. Anal.
- [T] J. Tits, *Endliche Spiegelungsgruppen, die als Weylgruppen auftreten*, Invent. Math. **43**, (1977), 283–295.

*E-mail address:* nagano@math.tohoku.ac.jp / nagano@math.uni-bonn.de

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY

AOBA, SENDAI, MIYAGI, 980-8578, JAPAN

*Current address:* Mathematisches Institut, Universität Bonn

Beringstraße 1, D-53115 Bonn, Germany