

On the periodic properties of self-decimated generators of pseudorandom numbers

Sergey Agievich, Oleg Solovey

National Research Center for Applied Problems of Mathematics and Informatics

Belarusian State University

Fr. Skorina av. 4, 220050 Minsk, Belarus

agievich@bsu.by, solovey@bsu.by

Abstract

We consider a self-decimated generator of pseudorandom numbers and examine the preperiod λ and the period μ of its state sequence. We obtain the expectations and variances of λ and μ for the case when decimation steps are chosen randomly and independently from the set $\{1, 2\}$.

1 Results

Let \mathbb{N} be the set of all positive integers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $A \subset \mathbb{N}_0$ be some finite alphabet. We denote by A^* the set of all finite words in the alphabet A (including the empty word ε) and by A^ω the set of all one-way infinite words (see [1, 2] for further details). Given a word $a \in A^*$, let $l(a)$ be the length of a and $w(a)$ be the sum of letters of a . Let c^n denote n successive instances of the letter c and ab denote the concatenation of the words a and b .

Given $s = s_0s_1\dots \in A^\omega$ and $T \in \mathbb{N}$, determine the numbers $\lambda = \lambda(s, T) \in \mathbb{N}_0$ and $\mu = \mu(s, T) \in \mathbb{N}$ such that

$$T \mid (s_\lambda + s_{\lambda+1} + \dots + s_{\lambda+\mu-1}) \quad (1)$$

and T does not divide any of the sums $s_t + s_{t+1} + \dots + s_\tau$, $0 \leq t < \tau < \lambda + \mu - 1$. For example, if $s = 2212221\dots = 2^212^31\dots$, then

$$\lambda(s, 1) = 0, \mu(s, 1) = 1, \quad \lambda(s, 2) = 0, \mu(s, 2) = 1, \dots \quad \lambda(s, 8) = 2, \mu(s, 8) = 5, \dots$$

The characteristics λ and μ describe periodic properties of self-decimated generators of pseudorandom numbers (see [3, 4]). Consider a generator G with the set of internal states \mathcal{S} , $|\mathcal{S}| = T$, and the state-transition function $\varphi: \mathcal{S} \rightarrow \mathcal{S}$. Let φ be a full cycle substitution,

usual way of G functioning, one chooses an initial state $S_0 \in \mathcal{S}$, calculates the sequence

$$S_{t+1} = \varphi(S_t), \quad t = 0, 1, \dots, \quad (2)$$

and uses the current internal state S_t to determine the current output pseudorandom number. The *self-decimation* (of internal states of G) means using an additional function $d: \mathcal{S} \rightarrow A$ and replacing (2) by the rule

$$S_{t+1} = \varphi^{d(S_t)}(S_t), \quad t = 0, 1, \dots \quad (3)$$

Now, if the word s consists of the successive letters $s_t = d(S_t)$, then $\lambda(s, T)$ and $\mu(s, T)$ are respectively the preperiod and period of the state sequence (3).

If the letters of s are chosen randomly, then λ and μ become random variables. The expectations $\mathbf{E} \lambda$, $\mathbf{E} \mu$ and variances $\mathbf{D} \lambda$, $\mathbf{D} \mu$ of these variables are of interest in connection with the estimation of the preperiod and period of self-decimated generators. One of the well-known results in the theory of random mappings can be stated as follows: If the letters of s are chosen randomly, independently and uniformly from the alphabet $A = \{0, 1, \dots, T-1\}$, then $\mathbf{E} \lambda(s, T)$ and $\mathbf{E} \mu(s, T)$ have the form $\sqrt{\pi T/8} + O(1)$ as $T \rightarrow \infty$ (see, for example, [5]). Unfortunately, obtaining similar asymptotic (or exact) expressions for arbitrary alphabet A seems to be a more complex task.

We consider the frequent choice $A = \{1, 2\}$ and obtain the following result.

Theorem. *If the letters of the word $s = s_0 s_1 \dots \in \{1, 2\}^\omega$ are chosen randomly, independently with the probabilities*

$$\mathbf{P} \{s_t = 1\} = 1 - p, \quad \mathbf{P} \{s_t = 2\} = p, \quad 0 < p < 1, \quad t = 0, 1, \dots,$$

then as $T \rightarrow \infty$ it holds that

$$\mathbf{E} \lambda(s, T) = \frac{p}{(1-p)(1+p)^2} + O(p^T T), \quad \mathbf{E} \mu(s, T) = \frac{T}{1+p} + O(p^T T), \quad (4)$$

$$\mathbf{D} \lambda(s, T) = \frac{p + p^3 + p^4}{(1-p)^2(1+p)^4} + O(p^T T^2), \quad \mathbf{D} \mu(s, T) = \frac{T p (1-p)}{(1+p)^3} + O(p^T T^2). \quad (5)$$

Note that (4), (5) also hold for the random word $s = s_0 s_1 \dots \in \{q, 2q\}^\omega$, where $q \in \mathbb{N}$ is coprime to T and $\mathbf{P} \{s_t = q\} = 1 - p$, $\mathbf{P} \{s_t = 2q\} = p$.

Note also that R. Rueppel in the paper [4] examined the case when the word s is determined from a linear recurrence sequence $\sigma_0, \sigma_1, \dots$ over the field of order 2 by the rule

$$s_t = \begin{cases} 1, & \sigma_t = 0, \\ 2, & \sigma_t = 1, \end{cases} \quad t = 0, 1, \dots$$

Rueppel showed that if the characteristic polynomial of the sequence $\sigma_0, \sigma_1, \dots$ is primitive of degree k , then

$$\mu(s, T) = \left\lfloor \frac{2T}{3} \right\rfloor, \quad T = 2^k - 1,$$

where $\lfloor z \rfloor$ is the largest integer $\leq z$. As we can see from Theorem, this estimation agrees with the expectation $\mathbf{E} \mu(s, T)$ for the random word s with $p = 1/2$.

Given s and T , determine the words $s_0 \dots s_{\lambda-1}$ and $s_\lambda \dots s_{\lambda+\mu-1}$, which we call the *prefix* and *cyclic part* of s , respectively. Conversely, any nonempty word $a \in A^*$ is a cyclic part of some word s . Given a , we can determine the possible values of $T = T(a)$ and the set $B(a)$ of the possible prefixes of s , using the following restrictions:

P1) $w(a) \equiv 0 \pmod{T}$;

P2) the residues

$$0, \quad a_0, \quad a_0 + a_1, \dots, \quad a_0 + a_1 + \dots + a_{l(a)-2} \pmod{T} \quad (6)$$

are pairwise distinct;

P3) if $b \in B(a)$ and $b \neq \varepsilon$, then any of the residues

$$-b_{l(b)-1}, \quad -b_{l(b)-1} - b_{l(b)-2}, \dots, \quad -b_{l(b)-1} - b_{l(b)-2} - \dots - b_0 \pmod{T}$$

differs from the residues (6).

Indeed, the condition P1 follows from the definition of a cyclic part. If the condition P2 fails, then

$$a_t + \dots + a_\tau \equiv 0 \pmod{T}$$

for some t, τ , $0 \leq t < \tau < l(a) - 1$ and a cannot be a cyclic part. Finally, if the condition P3 fails, then one of the congruences

$$b_t + \dots + b_{l(b)-1} \equiv 0 \pmod{T}, \quad b_t + \dots + b_{l(b)-1} + a_0 + \dots + a_\tau \equiv 0 \pmod{T}$$

holds for some $t \leq l(b) - 1$, $\tau < l(a) - 1$ and a cannot be a cyclic part again.

Suppose that $A = \{1, 2\}$ and let $\mathbf{x} = (x_1, x_2)$, $\mathbf{y} = (y_1, y_2)$. Given $\Omega \subseteq A^*$, define the generating function

$$G_\Omega(\mathbf{x}, \mathbf{y}, z) = \sum_{\substack{n_1, n_2, m_1, m_2 \geq 0 \\ t \geq 1}} g(n_1, n_2, m_1, m_2, t) x_1^{n_1} x_2^{n_2} y_1^{m_1} y_2^{m_2} z^t, \quad (7)$$

where $g(n_1, n_2, m_1, m_2, t)$ is the number of the words s with the cyclic part $a \in \Omega$ and prefix $b \in B(a)$ such that a contains n_1 letters 1 and n_2 letters 2, b contains m_1 letters 1 and m_2 letters 2, and $T(a) = t$.

Let us divide A^* into the subsets $\Omega_1, \Omega_2, \Omega_3, \Omega_4$ (which will be defined below) and determine the generating function of the form (7) for each subset.

1. $\Omega_1 = \{2\}^*1$. If $a \in \Omega_1$, then $T \mid w(a)$ according to P1 and $T \geq l(a) > w(a)/2$ according to P2. Therefore, $T(a) = w(a)$. Now, if $a = 2^m 1$, then $B(a) = \{2^k : k = 0, \dots, m\}$

$$\begin{aligned}
G_{\Omega_1}(\mathbf{x}, \mathbf{y}, z) &= \sum_{m \geq 0} x_1 x_2^m z^{2m+1} \sum_{k=0}^m y_2^k \\
&= \frac{x_1 z}{1-y_2} \sum_{m \geq 0} (x_2 z^2)^m - \frac{x_1 y_2 z}{1-y_2} \sum_{m \geq 0} (x_2 y_2 z^2)^m \\
&= \frac{x_1 z}{(1-y_2)(1-x_2 z^2)} - \frac{x_1 y_2 z}{(1-y_2)(1-x_2 y_2 z^2)} \\
&= \frac{x_1 z}{(1-x_2 z^2)(1-x_2 y_2 z^2)}.
\end{aligned}$$

2. $\Omega_2 = \{1, 2\}^* 1 \{2\}^* 1$. Again, $T(a) = w(a)$ according to P1, P2. Let $a = \alpha 1 2^m 1$, $\alpha \in A^*$. Using P3, we obtain $B(a) = \{2^k : k = 0, \dots, m\}$. Therefore,

$$\begin{aligned}
G_{\Omega_2}(\mathbf{x}, \mathbf{y}, z) &= \sum_{l \geq 0} (x_1 z + x_2 z^2)^l \sum_{m \geq 0} x_1^2 x_2^m z^{2m+2} \sum_{k=0}^m y_2^k \\
&= \frac{x_1^2 z^2}{(1-x_1 z - x_2 z^2)(1-x_2 z^2)(1-x_2 y_2 z^2)}.
\end{aligned}$$

3. $\Omega_3 = \{1, 2\}^* 1 2 \{2\}^*$. In this case $T(a) = w(a)$ and $B(a) = \{\varepsilon\} \cup \{2^k 1 : k = 0, \dots, m\}$ for $a = \alpha 1 2 2^m$. Therefore,

$$\begin{aligned}
G_{\Omega_3}(\mathbf{x}, \mathbf{y}, z) &= \sum_{l \geq 0} (x_1 z + x_2 z^2)^l \sum_{m \geq 0} x_1 x_2^{m+1} z^{2m+3} \left(1 + \sum_{k=0}^m y_1 y_2^k \right) \\
&= \frac{x_1 x_2 z^3 (1 + y_1 - x_2 y_2 z^2)}{(1-x_1 z - x_2 z^2)(1-x_2 z^2)(1-x_2 y_2 z^2)}.
\end{aligned}$$

4. $\Omega_4 = \{2\}^*$. If $a = 2^m$, then $T(a) = 2m$ and $B(a) = \{\varepsilon\} \cup \{2^k 1 : k = 0, \dots, m-1\}$. In addition, $T(a) = m$ and $B(a) = \{\varepsilon\}$ for odd m . The generating function has the form

$$\begin{aligned}
G_{\Omega_4}(\mathbf{x}, \mathbf{y}, z) &= \sum_{m \geq 1} x_1^m z^{2m} \left(1 + \sum_{k=0}^{m-1} y_1 y_2^k \right) + \sum_{m \geq 0} x_2^{2m+1} z^{2m+1} \\
&= \frac{1 + x_2 y_1 z^2 - x_2 y_2 z^2}{(1-x_2 z^2)(1-x_2 y_2 z^2)} + \frac{x_2 z}{1-x_2^2 z^2}.
\end{aligned}$$

Finally, we get

$$\begin{aligned}
G_{A^*}(\mathbf{x}, \mathbf{y}, z) &= G_{\Omega_1}(\mathbf{x}, \mathbf{y}, z) + G_{\Omega_2}(\mathbf{x}, \mathbf{y}, z) + G_{\Omega_3}(\mathbf{x}, \mathbf{y}, z) + G_{\Omega_4}(\mathbf{x}, \mathbf{y}, z) \\
&= \frac{1 + x_2 y_1 z^2 + x_1 x_2 y_2 z^3 - x_2 y_2 z^2}{(1-x_2 y_2 z^2)(1-x_1 z - x_2 z^2)} + \frac{x_2 z}{1-x_2^2 z^2}.
\end{aligned}$$

To find the expectation $\mathbf{E} \lambda(s, T)$, introduce the operator $E_{\mathbf{y}} = y_1 \frac{\partial}{\partial y_1} + y_2 \frac{\partial}{\partial y_2}$ and denote $\boldsymbol{\pi} = (1-p, p)$. We have

$$\mathbf{E} \lambda(s, T) = \sum_{n_1, n_2, m_1, m_2 \geq 0} \frac{g(n_1, n_2, m_1, m_2, T)(m_1 + m_2)}{(1-p)^{n_1+m_1} p^{n_2+m_2}} = [z^T] E_{\mathbf{y}} G_{A^*}(\boldsymbol{\pi}, \mathbf{y}, z)|_{\mathbf{y}=\boldsymbol{\pi}},$$

of (4) follows from

$$\begin{aligned}
E_{\mathbf{y}} G_{A^*}(\boldsymbol{\pi}, \mathbf{y}, z)|_{\mathbf{y}=\boldsymbol{\pi}} &= \frac{(1-p)pz^2}{(1-z)(1-p^2z^2)^2} \\
&= \frac{p}{(1-p)(1+p)^2(1-z)} - \frac{1}{4(1-pz)^2} - \frac{p}{4(1-p)(1-pz)} \\
&\quad + \frac{1-p}{4(1+p)(1+pz)^2} - \frac{(1-p)p}{4(1+p)^2(1+pz)} \\
&= \frac{p}{(1-p)(1+p)^2} \sum_{k \geq 0} z^k - \frac{1}{4} \sum_{k \geq 0} (k+1)p^k z^k - \frac{p}{4(1-p)} \sum_{k \geq 0} p^k z^k \\
&\quad + \frac{1-p}{4(1+p)} \sum_{k \geq 0} (k+1)(-p)^k z^k - \frac{(1-p)p}{4(1+p)^2} \sum_{k \geq 0} (-p)^k z^k.
\end{aligned}$$

Similarly, introducing the operator $E_{\mathbf{x}} = x_1 \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_2}$ and using the equalities

$$\begin{aligned}
\mathbf{E} \mu(s, T) &= [z^T] E_{\mathbf{x}} G_{A^*}(\mathbf{x}, \boldsymbol{\pi}, z)|_{\mathbf{x}=\boldsymbol{\pi}}, \\
\mathbf{D} \lambda(s, T) &= [z^T] E_{\mathbf{y}}^2 G_{A^*}(\boldsymbol{\pi}, \mathbf{y}, z)|_{\mathbf{y}=\boldsymbol{\pi}} - (\mathbf{E} \lambda(s, T))^2, \\
\mathbf{D} \mu(s, T) &= [z^T] E_{\mathbf{x}}^2 G_{A^*}(\mathbf{x}, \boldsymbol{\pi}, z)|_{\mathbf{x}=\boldsymbol{\pi}} - (\mathbf{E} \mu(s, T))^2,
\end{aligned}$$

we prove the remaining parts of (4), (5).

References

1. Goulden I. P., Jackson D. M. (1983). *Combinatorial Enumeration*. N.Y.: John Wiley.
2. Choffrut Ch., Karhumaki J. (1997). *Combinatorics of Words, in Handbook of Formal Languages (Rozenberg G., Salomaa A. eds.)*. Berlin, Heidelberg: Springer-Verlag.
3. Chambers W., Gollmann D. (1998). Generators for sequences with nearmaximal linear equivalence. *IEE Proc. E*. Vol. **135**, pp. 67-69.
4. Rueppel R. (1988). When shift registers clock themselves. *Lecture Notes in Computer Science, Advances in Cryptology – Eurocrypt 87*. Vol. **304**. Springer-Verlag, pp. 53-64.
5. Flajolet P., Odlyzko A. (1990). Random mapping statistics. *Lecture Notes in Computer Science, Advances in Cryptology – Eurocrypt 89*. Vol. **434**. Springer-Verlag, pp. 329-354.