p-ADIC MULTIPLE ZETA VALUES II — TANNAKIAN INTERPRETATIONS

HIDEKAZU FURUSHO

ABSTRACT. We establish a tannakian formalism of p-adic multiple polylogarithms and p-adic multiple zeta values introduced in our previous paper via a comparison isomorphism between a de Rham fundamental torsor and a rigid fundamental torsor of the projective line minus three points and also discuss its Hodge and étale analogues. As an application we give a way to erase log poles of p-adic multiple polylogarithms and introduce overconvergent p-adic multiple polylogarithms which might be p-adic multiple analogue of Zagier's single-valued complex polylogarithms.

Contents

0. Introduction	1
1. Preliminaries	4
1.1. de Rham setting	4
1.2. rigid setting	6
1.3. Betti setting	8
1.4. étale setting	9
2. Tannakian interpretations	10
2.1. Berthelot-Ogus side — main results	11
2.2. Hodge side — analogous results	19
2.3. Artin side — reviews	23
3. Motivic views	24
3.1. Drinfel'd's context	24
3.2. Racinet's context	27
3.3. Deligne-Goncharov's context	28
References	32

0. INTRODUCTION

This paper is the continuation of our previous paper [F1]. Let p be a prime number. Let $m, k_1, \dots, k_m \in \mathbf{N}$. In [F1] we constructed p-adic multiple polylogarithm $Li_{k_1,\dots,k_m}(z)$ ($z \in \mathbf{C}_p$), that is, a p-adic analogue of the (one-variable) complex multiple polylogarithm defined locally near 0 by

(0.1)
$$Li_{k_1,\dots,k_m}(z) = \sum_{\substack{0 < n_1 < \dots < n_m \\ 1}} \frac{z^{n_m}}{n_1^{k_1} \cdots n_m^{k_m}}$$

and p-adic multiple zeta value $\zeta_p(k_1, \dots, k_m)$, that is, a p-adic analogue of multiple zeta value

(0.2)
$$\zeta(k_1, \cdots, k_m) = \sum_{0 < n_1 < \cdots < n_m} \frac{1}{n_1^{k_1} \cdots n_m^{k_m}} \qquad (k_m > 1).$$

This was achieved by giving a *p*-adic Drinfel'd associator $\Phi_{KZ}^p(A, B)$ of the *p*-adic KZ equation and its fundamental solution $G_0(A, B)(z)$ ($z \in \mathbf{C}_p$). The purpose of this paper is to relate these constructions to the motivic fundamental torsors of the projective line minus three points. To do this we will give a tannakian interpretation of the constructions of [F1]. Using the Frobenius action on the rigid realization of the above torsor, we will define the overconvergent *p*-adic multiple polylogarithm. This function can be expressed by the *p*-adic multiple polylogarithm constructed in [F1]. Using Hodge and *l*-adic étale realizations we will recover the complex multiple polylogarithm and the *l*-adic polylogarithms of Wojtkowiak.

Our main tool is a comparison isomorphism (Lemma 2.2)

(0.3)
$$\pi_1^{\mathrm{DR}}(\mathbf{P}^1_{\mathbf{Q}_p} \setminus \{0, 1, \infty\} : \overrightarrow{01}, z) \cong \pi_1^{p, \mathrm{rig}}(\mathbf{P}^1_{\mathbf{F}_p} \setminus \{0, 1, \infty\} : \overrightarrow{01}, z_0)$$

between the de Rham (§1.1) and the rigid (§1.2) fundamental torsor where $z_0 \in \mathbf{P}^1(\mathbf{F}_p) \setminus \{0, 1, \infty\}$ is the reduction of $z \in \mathbf{P}^1(\mathbf{Q}_p) \setminus \{0, 1, \infty\}$ and $\overrightarrow{01}$ is a tangential basepoint. There is a canonical de Rham path d_z (Lemma 1.2) in LHS constructed from the canonical extension of unipotent connections and a canonical rigid path c_{z_0} (Lemma 1.8) in RHS, Besser-Vologodsky's Frobenius invariant path [Bes, Vol]. Identifying these two torsors by (0.3) we get a de Rham loop $d_z^{-1}c_{z_0} \in \pi_1^{\mathrm{DR}}(\mathbf{P}_{\mathbf{Q}_p}^1 \setminus \{0, 1, \infty\} : \overrightarrow{01})(\mathbf{Q}_p)$. Our tannakian interpretation of *p*-adic multiple polylogarithms is

Theorem 2.3. By the embedding $i : \pi_1^{\mathrm{DR}}(\mathbf{P}^1_{\mathbf{Q}_p} \setminus \{0, 1, \infty\} : \overrightarrow{01})(\mathbf{Q}_p) \hookrightarrow \mathbf{Q}_p \langle \langle A, B \rangle \rangle$ (2.1), the loop $d_z^{-1}c_{z_0}$ corresponds to the non-commutative formal power series $G_0(A, B)(z)$ of the p-adic multiple polylogarithms.

Similarly by letting z to be a tangential basepoint 10 we get a tannakian interpretation of p-adic multiple zeta values:

Theorem 2.5. By the embedding *i* the loop $d_{\overrightarrow{10}}^{-1}c_{\overrightarrow{10}}$ corresponds to the noncommutative formal power series $\Phi_{KZ}^{p}(A, B)$ of the *p*-adic multiple zeta values.

RHS of (0.3) admits the Frobenius action ϕ_p (Definition 1.7). Deligne [De2] introduced a variant of *p*-adic multiple zeta values to be a coefficient of the series $\Phi_{\text{De}}^p(A, B) = i(d_{\overrightarrow{10}}^{-1}\phi_p(d_{\overrightarrow{10}}))$ which we call the *p*-adic Deligne associator.

Theorem 2.8. There is the following explicit relationship between the p-adic Drinfel'd associator and the p-adic Deligne associator:

$$\Phi^p_{\mathrm{KZ}}(A,B) = \Phi^p_{\mathrm{De}}(A,B) \cdot \Phi^p_{\mathrm{KZ}}\left(\frac{A}{p}, \Phi^p_{\mathrm{De}}(A,B)^{-1}\frac{B}{p}\Phi^p_{\mathrm{De}}(A,B)\right).$$

This gives formulae expressing our *p*-adic multiple zeta values in terms of Deligne's *p*-adic multiple zeta values and vice versa. Namely his *p*-adic multiple zeta values are equivalent to ours. We introduce a pro-unipotent overconvergent iso-crystal $(\mathcal{V}_{\text{KZ}}^{\dagger}, \nabla_{\text{KZ}}^{\dagger})$ on $\mathbf{P}_{\mathbf{F}_{p}}^{1} \setminus \{0, 1, \infty\}$ associated with the *p*-adic KZ equation and show

Proposition 2.11. The pro-object $(\mathcal{V}_{KZ}^{\dagger}, \nabla_{KZ}^{\dagger})$ naturally admits a Frobenius structure.

The canonical section 1 of \mathcal{V}^{\dagger} under the Frobenius action will map to a noncommutative power series $G_0^{\dagger}(z)$ whose coefficient will be denoted by $Li_{k_1,\dots,k_m}^{\dagger}(z)$. This series satisfies the differential equation in [U2, Y]

$$dg = \left(\frac{A}{z} + \frac{B}{z-1}\right)gdz - g\left(\frac{dz^p}{z^p}\frac{A}{p} + \frac{dz^p}{z^p-1}\Phi_{\mathrm{De}}^p(A,B)^{-1}\frac{B}{p}\Phi_{\mathrm{De}}^p(A,B)\right)$$

and $Li_{k_1,\dots,k_m}^{\dagger}(z)$ are an overconvergent analogue of our *p*-adic multiple polylogarithms. The special value of these functions at z = 1 will be the Deligne's *p*-adic multiple zeta values. The relation between $G_0^{\dagger}(A, B)(z)$ and our $G_0(A, B)(z)$ is given by the following:

Theorem 2.14. The overconvergent p-adic multiple polylogarithm $Li_{k_1,\dots,k_m}^{\dagger}(z)$ is expressed as a combination of our p-adic multiple zeta values, p-adic multiple polylogarithms and the p-adic logarithm by

$$G_0^{\dagger}(A,B)(z) = G_0(A,B)(z) \cdot G_0\left(\frac{A}{p}, \Phi_{\rm De}^p(A,B)^{-1}\frac{B}{p}\Phi_{\rm De}^p(A,B)\right)(z^p)^{-1}.$$

The function $Li_{k_1,\dots,k_m}(z)$ is not overconvergent but a Coleman function with log poles around z = 1 and ∞ . The above formulae is a way to erase these log poles. We will give calculations in Example 2.16.

The organization of this paper is as follows. §1 is preliminary for the rest of paper. We will review the definition of various (de Rham, rigid, Betti and étale) fundamental groups, torsors with their various additional structures, i.e. the Frobenius action, the infinity Frobenius action and the Galois group action. We also discuss tangential basepoints. The Deligne's canonical path in the de Rham fundamental torsor in Lemma 1.2 and Besser-Vologodsky's Frobenius invariant path in the rigid fundamental torsor in Lemma 1.8 will be recalled.

§2 is devoted to tannakian formalisms which contains our main results. In Theorem 2.3 and Theorem 2.5 we will clarify tannakian origins of the *p*-adic multiple polylogarithms and p-adic multiple zeta values. An explicit relationship between our *p*-adic multiple zeta values and Deligne's *p*-adic multiple zeta values [De2] will be stated in Theorem 2.8. A formula to express the overconvergent p-adic multiple polylogarithms in terms of our *p*-adic multiple polylogarithms will be stated in Theorem 2.14. In $\S2.2$ the Hodge analogue of these results will be discussed. The tannakian origin of complex multiple polylogarithms will be discussed in Proposition 2.20 and for multiple zeta values in Proposition 2.21. The infinity Frobenius action on the Betti fundamental torsor of $\mathbf{P}^1(\mathbf{C}) \setminus \{0, 1, \infty\}$ will be used to introduce another variant of multiple polylogarithms (2.20). In Theorem 2.27 it will be shown that these functions are single-valued and real-analytic. In Proposition 2.26 we show a formula analogous to Theorem 2.14 to express them in terms of complex multiple polylogarithms. §2.3 is a brief explanation of analogous story in the étale side. We discuss an étale analogue of polylogarithm (2.22) expressed in terms of Wojtkowiak's l-adic polylogarithm and an étale analogue of Riemann zeta values (Example 2.33) expressed in term of Soulé characters and cyclotomic characters following [NW, I].

§3 is motivic. The algebra generated by *p*-adic multiple zeta values will be related to Drinfel'd's [Dr] pro-algebraic group \underline{GRT}_1 , Racinet's [R] pro-algebraic group \underline{DMR}_0 and the motivic Galois group $\pi_1(\mathcal{MT}(\mathbf{Z}))$. In §3.1 we recall Drinfel'd's pro-algebraic bi-torsor of Grothendieck-Teichmüller. In §3.2, we recall Racinet's

[R] pro-algebraic bi-torsor made by double shuffle relations and briefly explain a partial analogous story to §3.1. §3.3 is a review of the formalism of mixed Tate motives in [DG]. The torsor of motivic Galois group is related with the torsors in §3.1 and §3.2. We give some motivic interpretations on Zagier's conjecture on multiple zeta values and Ihara's conjecture on Galois image. We also explain a motivic way of proving double shuffle relations for *p*-adic multiple zeta values using [DG] and [Y].

Acknowledgments . This work was supported by the NSF grants DMS-0111298. The author would like to express particular thanks to Prof. Deligne for many suggestions. He is also grateful to Amir Jafari and Go Yamashita for many useful discussion. He is also grateful to the referee for comments.

1. Preliminaries

This section is preliminaries for the next section. We give materials which are indispensable to our main results by recalling the definitions of various (de Rham, rigid, Betti and étale) fundamental groups and torsors with tangential base points and discussing additional structures there.

In this section, we concentrate on a curve $X_K = \overline{X}_K - D_K$, where \overline{X}_K is a proper smooth, geometrically connected curve \overline{X}_K over a field K and D_K is its divisor over K (sometimes we omit K).

1.1. **de Rham setting.** In this subsection, we will recall the definitions of the de Rham fundamental group, torsor, path space, the tangential base point and the canonical base point and see how we obtain the canonical de Rham path in the de Rham fundamental torsor by the canonical base point, all of which are developed in [De1]§12.

Notation 1.1. In this subsection we assume that K is a field of characteristic 0. We denote the category of the nilpotent part ¹ of the category of the pair (\mathcal{V}, ∇) of a coherent \mathcal{O}_X -module sheaf \mathcal{V} on X_K and an integrable (i.e. $\nabla \circ \nabla = 0$) connection $\nabla : \mathcal{V} \to \mathcal{V} \otimes \Omega^1_{X_K}$ by $\mathcal{NC}^{\mathrm{DR}}(X_K)$, It forms a neutral tannakian category (for the basics, consult [DM]) by [De1]. For our quick review of these materials, see [S1].

In the de Rham realization, we consider three fiber functors. Suppose that x is a K-valued point of X, i.e. $x : SpecK \to X_K$. The first one is $\omega_x : \mathcal{NC}^{DR}(X_K) \to Vec_K$ (Vec_K : the category of finite dimensional K-vector spaces) which is the pullback of $\mathcal{NC}^{DR}(X_K)$ by x. Actually this forms a fiber functor (consult [DM] for its definition) by [S1]§3.1. The second fiber functor is the one introduced in [De1]§15 (and developed to higher dimensional case in [BF]): Let $s \in D$ and put $T_s = T_s \overline{X}_K$ and $T_s^{\times} = T_s \setminus \{0\} (\cong \mathbf{G}_m)$, where $T_s \overline{X}_K$ is the tangent vector space of \overline{X}_K at s. By the construction of the extension of unipotent integrable connections to tangent bundles in [De1]§15.28-15.36, we have a morphism of tannakian categories (see also [BF])

(1.1)
$$Res_s : \mathcal{NC}^{\mathrm{DR}}(X_K) \to \mathcal{NC}^{\mathrm{DR}}(T_s^{\times})$$

which we call the tangential morphism. By pulling back, each K-valued point t_s of T_s^{\times} determines a fiber functor $\omega_{t_s} : \mathcal{NC}^{\mathrm{DR}}(X_K) \to \operatorname{Vec}_K$ which we call the

 $^{^1\}mathrm{It}$ means the full subcategory consisting of objects which are iterated extensions of the unit objects.

tangential base point. The last fiber functor is the special one under the assumption

(1.2)
$$H^1(\overline{X}, \mathcal{O}_{\overline{X}}) = 0.$$

(i.e. \overline{X} is a curve with genus 0): Let $\omega_{\Gamma} : \mathcal{NC}^{\mathrm{DR}}(X_K) \to \operatorname{Vec}_K$ be a functor sending each object $(\mathcal{V}, \nabla) \in \mathcal{NC}^{\mathrm{DR}}(X_K)$ to $\Gamma(\overline{X}_K, \mathcal{V}_{\mathrm{can}})$, the global section of its canonical extension $(\mathcal{V}_{\mathrm{can}}, \nabla_{\mathrm{can}})$ into \overline{X}_K [De1] §12.2. This functor forms a fiber functor ([De1] §12.4) because $\mathcal{V}_{\mathrm{can}}$ forms a trivial bundle over \overline{X}_K due to the condition (1.2) by [De1] Proposition 12.3. We call ω_{Γ} the canonical base point. We note that the assumption (1.2) is necessary. The functor ω_{Γ} would be no longer a fiber functor if \overline{X} was a proper smooth curve with genus g > 0.

Lemma 1.2 ([De1]§15.52). Let ω_* and $\omega_{*'}$ be any fiber functors above. Then under the assumption (1.2) there is a canonical isomorphism $d_{**'}: \omega_* \to \omega_{*'}$.

Proof. It is because \mathcal{V}_{can} forms a trivial bundle for $(\mathcal{V}, \nabla) \in \mathcal{NC}^{DR}(X_K)$ that we get a canonical isomorphism

$$\omega_x(\mathcal{V},\nabla) = \mathcal{V}_{(x)} = \mathcal{V}_{\operatorname{can},(x)} \cong \mathcal{O}_{\overline{X}_K,(x)} \underset{K}{\otimes} \Gamma(\overline{X}_K, \mathcal{V}_{\operatorname{can}}) \cong \mathcal{O}_{\overline{X}_K,(x)} \underset{K}{\otimes} \omega_{\Gamma}(\mathcal{V}, \nabla) \cong \omega_{\Gamma}(\mathcal{V}, \nabla)$$

and similarly

$$\omega_{t_s}(\mathcal{V}, \nabla) \cong \mathcal{O}_{T_s^{\times}, (t_s)} \underset{K}{\otimes} \Gamma(\overline{X}_K, \mathcal{V}_{\operatorname{can}}) \cong \omega_{\Gamma}(\mathcal{V}, \nabla).$$

Here (x) stands for the fiber of sheaves at x. By factoring through ω_{Γ} , we obtain a canonical isomorphism between any two of them.

The fiber functors ω_x and ω_{t_s} (and ω_{Γ} under the assumption (1.2)) play a role of 'base points'.

Definition 1.3 ([De1]). Let ω_* and $\omega_{*'}$ be any fiber functors above. The de Rham fundamental group $\pi_1^{\mathrm{DR}}(X_K:*)$ is <u>Aut</u>^{\otimes}(ω_*) and the de Rham fundamental torsor $\pi_1^{\mathrm{DR}}(X_K:*,*')$ is <u>Isom</u>^{\otimes}($\omega_*, \omega_{*'}$) (for <u>Aut</u>^{\otimes} and <u>Isom</u>^{\otimes}, see [DM]).

The former is a pro-algebraic group over K and the latter is a pro-algebraic bitorsor over K where $\pi_1^{\text{DR}}(X_K:*)$ acts from the right and $\pi_1^{\text{DR}}(X_K:*)$ acts from the left ² and by definition $\pi_1^{\text{DR}}(X_K:*,*) = \pi_1^{\text{DR}}(X_K:*)$. Particularly under the assumption (1.2), we have a canonical de Rham path $d_{*,*'} \in \pi_1^{\text{DR}}(X_K:*,*')(K)$ by Lemma 1.2. This path is compatible with the composition: $d_{*,*'} \cdot d_{*,*'} = d_{*,*''}$. We also note that for any field extension K'/K the category $\mathcal{NC}^{\text{DR}}(X_{K'})$ is equivalent to $\mathcal{NC}^{\text{DR}}(X_K) \otimes_K K'$ which is the category whose set of morphisms, K'-linear space, is replaced by the extension of the set of morphisms of $\mathcal{NC}^{\text{DR}}(X_K)$, K-linear space, by K' ([De1]§10.41), whence $\pi_1^{\text{DR}}(X_{K'}:*) \cong \pi_1^{\text{DR}}(X_K:*) \times_K K'$ and $\pi_1^{\text{DR}}(X_{K'}:*,*') \cong \pi_1^{\text{DR}}(X_K:*,*') \times_K K'$.

To establish the notion of the de Rham path space, we recall the notion of affine group \mathcal{T} -schemes for a tannakian category \mathcal{T} ([De1]§5.4 and [DG]§2.6). The category of affine \mathcal{T} -schemes is the dual of the category of ind-objects of \mathcal{T} which are unitary commutative algebras, that is, each object is an ind-object A of \mathcal{T} with a product $A \otimes A \to A$ and a unit $1 \to A$ verifying usual axioms. We denote the corresponding affine \mathcal{T} -schemes with A by SpecA. An affine group \mathcal{T} -scheme is a

²For any two loops γ and γ' , the symbol $\gamma' \cdot \gamma$ stands for the composition of two which takes γ at first and takes γ' next. Throughout this paper, we keep this direction.

group object of the category of affine \mathcal{T} -schemes, i.e. the spectrum of a commutative Hopf algebra.

Definition 1.4. The de Rham path space ³ $\mathcal{P}_{X_K}^{\mathrm{DR}}$ is an affine group $\mathcal{NC}^{\mathrm{DR}}(X_K) \otimes \mathcal{NC}^{\mathrm{DR}}(X_K)$ ⁴-scheme introduced in [De1]§6.13 and [DG]§4, which satisfies

$$(\omega \otimes \omega')(\mathcal{P}_{X_K}^{\mathrm{DR}}) = \underline{Isom}^{\otimes}(\omega, \omega')(K)$$

for any fiber functors $\omega, \omega' : \mathcal{NC}^{\mathrm{DR}}(X_K) \to \operatorname{Vec}_K$ and is represented by an indobject $\mathcal{A}^{\mathrm{DR}}$ of $\mathcal{NC}^{\mathrm{DR}}(X_K) \otimes \mathcal{NC}^{\mathrm{DR}}(X_K)$ i.e. $\mathcal{P}_{X_K}^{\mathrm{DR}} = \operatorname{Spec}\mathcal{A}^{\mathrm{DR}}$. The de Rham fundamental path space $\mathcal{P}_{X_K,\omega}^{\mathrm{DR}}$ with the base point ω is its pull-back by $\omega \otimes id$, which is represented by the ind-object $\mathcal{A}_{\omega}^{\mathrm{DR}} := (\omega \otimes id)(\mathcal{A}^{\mathrm{DR}})$ of $\mathcal{NC}^{\mathrm{DR}}(X_K)$.

By definition, we have

(1.3)
$$(\omega_* \otimes \omega_{*'}) \mathcal{P}_{X_K}^{\mathrm{DR}} = \omega_{*'} (\mathcal{P}_{X_K,\omega_*}^{\mathrm{DR}}) = \pi_1^{\mathrm{DR}} (X_K : *, *')(K).$$

1.2. **rigid setting.** In this paper we take rigid fundamental groups (torsors) as a crystalline realization of motivic fundamental groups (torsors). We will recall the definition of the rigid fundamental group [CLS, S2], torsor, path space, the tangential base point [BF, De1] and the canonical Frobenius invariant path [Bes, Vol].

Notation 1.5. In this subsection we assume that K is a non-archimedean local field of characteristic 0. We denote V to be its valuation ring and k to be the residue field of characteristic p > 0 with $q(=p^r)$ -elements. We also assume that \overline{X}_K with D_K is a generic fiber of a proper smooth geometrically connected curve \mathcal{X} with a divisor \mathcal{D} over V. We denote its special fiber over k by \overline{X}_0 with D_0 . Put $X_0 = \overline{X}_0 \setminus D_0$ and its natural embedding $j : X_0 \hookrightarrow \overline{X}_0$. We denote the category of the nilpotent part of the category of (overconvergent ⁵) isocrystals, which consists of the pair $(\mathcal{V}^{\dagger}, \nabla^{\dagger})$ of a coherent $j^{\dagger}\mathcal{O}_{]\overline{X}_0[}^{}$ -module sheaf \mathcal{V}^{\dagger} and an integrable connection $\nabla^{\dagger} : \mathcal{V}^{\dagger} \to \mathcal{V}^{\dagger} \otimes_{\mathcal{O}_{]\overline{X}_0[}} \Omega^1_{]\overline{X}_0[}$, by $\mathcal{NI}^{\dagger}(X_0)$. Here $]\overline{X}_0[$ stands for the tubular neighborhood [Ber] of \overline{X}_0 . This category depends only on X_0 and forms a neutral tannakian category and depends only on X_0 (cf. [Ber, CLS, S2]). For our quick review of these materials, see [Bes].

In the rigid realization, we consider three fiber functors. Suppose that x_0 is a k-valued point of X_0 , i.e. $x_0: Spec \ k \to X_0$. The first one is $\omega_{x_0}: \mathcal{NI}^{\dagger}(X_0) \to Vec_{K_0}$ which is the pull-back of $\mathcal{NI}^{\dagger}(X_0)$ by x_0 . This forms a fiber functor and is described by $\omega_{x_0}(\mathcal{V}^{\dagger}, \nabla^{\dagger}) = \mathcal{V}^{\dagger}(]x_0] \nabla^{\dagger} = \{v \in \mathcal{V}^{\dagger}(]x_0]) \ | \nabla^{\dagger}(v) = 0\}$ for $(\mathcal{V}^{\dagger}, \nabla^{\dagger}) \in \mathcal{NI}^{\dagger}(X_0)$ on the tubular neighborhood of $]x_0[$ [Bes]. The second fiber functor is $\omega_{s_0}: \mathcal{NI}^{\dagger}(X_0) \to Vec_K \ (s_0 \in D_0)$ introduced in [BF] Definition 1.3: This is described by $\omega_{s_0}(\mathcal{V}^{\dagger}, \nabla^{\dagger}) = \{v \in \mathcal{V}^{\dagger}(]s_0] [\log^a z_{s_0}] \ | \nabla^{\dagger}(v) = 0\}$, where \log^a is a branch of p-adic logarithm such that $\log^a p = a \in K$ and z_{s_0} is a local parameter of $]s_0[$ (cf.[F1] §2.1). The last fiber functor is the rigid version of the tangential base point: Put $T_{s_0} = T_{s_0} \overline{X_0}$ and $T_{s_0}^{\times} = T_{s_0} \setminus \{0\} (\cong \mathbf{G}_m)$ where $T_{s_0} \overline{X_0}$ is the tangent

³In [De1] §6.13, it is called the fundamental groupoid.

⁴That is the tensor product of tannakian categories [De1]§5.18.

 $^{^5\}mathrm{We}$ can omit it because it is shown in [CLS] that unipotent (nilpotent) isocrystals are overconvergent.

⁶We do not call \dagger 'plus' but 'dagger'. For the definition of $j^{\dagger}\mathcal{O}_{|\overline{X}_0|}$, see [Bes].

space of \overline{X}_0 at s_0 . By the construction in [BF]§2, we have a morphism of tannakian categories

(1.4)
$$\operatorname{Res}_{s_0} : \mathcal{NI}^{\dagger}(X_0) \to \mathcal{NI}^{\dagger}(T_{s_0}^{\times}),$$

which we call the tangential morphism. Thus each k-valued point t_{s_0} of $T_{s_0}^{\times}$ determines a fiber functor $\omega_{t_{s_0}} : \mathcal{NI}^{\dagger}(X_0) \to Vec_K$ by pulling back. Shortly the construction of Res_{s_0} developed in [BF] is the transmission of the morphism (1.1) into the rigid setting by the categorical equivalences $\mathcal{NC}^{\mathrm{DR}}(X_K) \cong \mathcal{NI}^{\dagger}(X_0)$ and $\mathcal{NC}^{\mathrm{DR}}(T_s^{\times}) \cong \mathcal{NI}^{\dagger}(T_{s_0}^{\times})$ in [CLS, S2], where s_0 is the reduction of s.

The fiber functors, ω_{x_0} , ω_{s_0} and $\omega_{t_{s_0}}$, play a role of 'base points'.

Definition 1.6. Let ω_* and $\omega_{*'}$ be any fiber functors above. The rigid fundamental group $\pi_1^{p,\mathrm{rig}}(X_0:*)$ is $\underline{Aut}^{\otimes}(\omega_*)$ and the rigid fundamental torsor $\pi_1^{p,\mathrm{rig}}(X_0:*,*')$ is $\underline{Isom}^{\otimes}(\omega_*,\omega_{*'})$ (cf. [CLS, S2]). The rigid path space $\mathcal{P}_{X_0}^{\dagger}$ is an affine group $\mathcal{NI}^{\dagger}(X_0) \otimes \mathcal{NI}^{\dagger}(X_0)$ -scheme which satisfies

$$(\omega \otimes \omega')(\mathcal{P}_{X_0}^{\dagger}) = \underline{Isom}^{\otimes}(\omega, \omega')(K)$$

for any fiber functors $\omega, \omega' : \mathcal{NI}^{\dagger}(X_0) \to Vec_K$ and is represented by an ind-object \mathcal{A}^{\dagger} of $\mathcal{NI}^{\dagger}(X_0) \otimes \mathcal{NI}^{\dagger}(X_0)$, i.e. $\mathcal{P}_{X_0}^{\dagger} = Spec\mathcal{A}^{\dagger}$. The rigid path space $\mathcal{P}_{X_0,\omega}^{\dagger}$ with the base point ω is its pull-back by $\omega \otimes id$, which is represented by the ind-object $\mathcal{A}_{\omega}^{\dagger} := (\omega \otimes id)(\mathcal{A}^{\dagger})$ of $\mathcal{NI}^{\dagger}(X_0)$.

We note that $\pi_1^{p,\operatorname{rig}}(X_0:*,*')$ is a pro-algebraic bitorsor over K where $\pi_1^{p,\operatorname{rig}}(X_0:*)$ acts from the right and $\pi_1^{p,\operatorname{rig}}(X_0:*')$ acts from the left. By definition we have

$$(\omega_* \otimes \omega_{*'}) \mathcal{P}_{X_0}^{\dagger} = \omega_{*'}(\mathcal{P}_{X_0,\omega_*}^{\dagger}) = \pi_1^{p, \mathrm{rig}}(X_0 : *, *')(K).$$

Let F_q be r-th power of the absolute Frobenius automorphism on X_0 and its induced automorphism on $T_{s_0}^{\times}$ ($s_0 \in D_0$). The pull-back functors $F_q^* : \mathcal{NI}^{\dagger}(X_0) \to \mathcal{NI}^{\dagger}(X_0)$ and $F_q^* : \mathcal{NI}^{\dagger}(T_{s_0}^{\times}) \to \mathcal{NI}^{\dagger}(T_{s_0}^{\times})$ determine equivalences of tannakian categories. We note that they are compatible with (1.4) ([BF] §2). They induce isomorphisms ([De1] §10.44)

(1.5)
$$F_q: \mathcal{P}_{X_0} \to F_q^* \mathcal{P}_{X_0},$$

(1.6)
$$F_q: \mathcal{P}_{X_0,\omega} \to F_q^*(\mathcal{P}_{X_0,F_q^*\omega})$$

(1.7)
$$F_q: \pi_1^{p, \operatorname{rig}}(X_0: *, *') \to \pi_1^{p, \operatorname{rig}}(X_0: *, *').$$

Definition 1.7. ([De1] §10.44) The Frobenius actions ϕ_q stands for the inverse ⁷ F_q^{-1} in (1.5) ~ (1.7).

By (1.5) and (1.6) the ind-objects \mathcal{A}^{\dagger} and $\mathcal{A}^{\dagger}_{\omega}$ in Definition 1.6 naturally admit structures of *ind*-overconvergent *F*-isocrystals, where an overconvergent *F*-isocrystal means an overconvergent isocrystal *V* with a Frobenius structure ϕ (i.e. a horizontal isomorphism $\phi : F_q^* V \to V$). The description of the Frobenius structure in case of $X = \mathbf{P}^1 \setminus \{0, 1, \infty\}$ is one of our main topics in this paper. The following lemma which takes a place of Lemma 1.2 in our rigid setting is indispensable to our study.

⁷We take the inverse to follow [De1] §13.6.

Lemma 1.8. Let ω_* and $\omega_{*'}$ be any fiber functors defined above. Then there exists a unique path $c_{*,*'} \in \pi_1^{p,\operatorname{rig}}(X_0 : *, *')(K)$ which is invariant under the Frobenius action, i.e. $F_q(c_{*,*'}) = c_{*,*'}$.

This result is shown by Besser [Bes] and Vologodsky [Vol] for usual base points (more generally in higher dimensional setting) but their proofs also work for tangential base points and points on D_0 directly. We call $c_{*,*'}$ the Frobenius invariant path. We have compatibilities, such as functorialities and $c_{*',*''} \cdot c_{*,*'} = c_{*,*''}$ (for more details, see [Bes]).

1.3. Betti setting. We will briefly recall the definition of the Betti fundamental group, torsor and the tangential base point in [De1]§15.

Notation 1.9. In this subsection, we assume $k = \mathbf{C}$. We denote the category of the nilpotent part of the category of local systems ⁸ of finite dimensional **Q**-vector spaces over the topological space $X(\mathbf{C})$ by $\mathcal{NL}^{\text{Be}}(X(\mathbf{C}))$. It also forms a neutral tannakian category over **Q** by [De1]§10.

In the Betti realization, we consider two fiber functors. Suppose that $x \in X(\mathbf{C})$. The first one is $\omega_x : \mathcal{NL}^{\text{Be}}(X(\mathbf{C})) \to Vec_{\mathbf{Q}}$ which associates each nilpotent local system V with the pullback V(x). Actually this forms a fiber functor of $\mathcal{NL}^{\text{Be}}(X(\mathbf{C}))$ by [De1]§10. The second one is the tangential basepoint: Let $s \in D(\mathbf{C})$ and put $T_s = T_s \overline{X}(\mathbf{C})$ and $T_s^{\times} = T_s \setminus \{0\} (\cong \mathbf{G}_m)$, where $T_s \overline{X}(\mathbf{C})$ is the tangent vector space of $\overline{X}(\mathbf{C})$ at s. By [De1]§15.3-15.12, we have a morphism of tannakian categories

(1.8)
$$\operatorname{Res}_s : \mathcal{NL}^{\operatorname{Be}}(X(\mathbf{C})) \to \mathcal{NL}^{\operatorname{Be}}(T_s^{\times}(\mathbf{C})).$$

which we call the tangential morphism. By pulling back, each point $t_s \in T_s^{\times}(\mathbf{C})$ determines a fiber functor $\omega_{t_s} : \mathcal{NL}^{\text{Be}}(X(\mathbf{C})) \to Vec_{\mathbf{Q}}$.

The above fiber functors, ω_x and ω_{t_s} , play a role of 'base points' and the following is a reformulation of the Betti fundamental group and torsor in [De1]§13.

Definition 1.10. Let ω_* and ω'_* be any fiber functors above. The Betti fundamental group $\pi_1^{\text{Be}}(X(\mathbf{C}):*)$ is <u>Aut</u>^{\otimes}(ω_*) and the Betti fundamental torsor $\pi_1^{\text{Be}}(X(\mathbf{C}):*,*')$ is <u>Isom</u>^{\otimes}($\omega_*, \omega_{*'}$).

The former is a pro-algebraic group over \mathbf{Q} and the latter is a pro-algebraic bitorsor over \mathbf{Q} where $\pi_1^{\text{Be}}(X(\mathbf{C}):*)$ acts on the right and $\pi_1^{\text{Be}}(X(\mathbf{C}):*')$ acts on the left. We note that we have a natural morphism from the topological fundamental group $\pi_1^{\text{top}}(X(\mathbf{C}):*)$ (resp. torsor $\pi_1^{\text{top}}(X(\mathbf{C}):*,*')$) to the set of \mathbf{Q} -valued points on the Betti fundamental group $\pi_1^{\text{Be}}(X(\mathbf{C}):*)$ (resp. torsor $\pi_1^{\text{Be}}(X(\mathbf{C}):*,*')$), which induces an isomorphism between $\pi_1^{\text{Be}}(X(\mathbf{C}):*)$ and the Malcev (unipotent) completion [De1]§9 over \mathbf{Q} of $\pi_1^{\text{top}}(X(\mathbf{C}):*)$ ([De1] §10.24). As for the topological fundamental groups and torsors with tangential base points, see [De1]§§15.3-15.12. Although we have a canonical path, both in the de Rham setting $d_{*,*'}$ Lemma 1.2 and in the rigid setting $c_{*,*'}$ Lemma 1.8, the author do not know whether we have any canonical path connecting any two base points * and *' in the Betti setting or not.

⁸We mean covariant functors $V : \Pi_{X(\mathbf{C})} \to Vec_{\mathbf{Q}}$ where $\Pi_{X(\mathbf{C})}$ stands for the category whose objects are points on $X(\mathbf{C})$ and whose set of morphisms between two points * and *' on $X(\mathbf{C})$ is the usual topological fundamental torsor $\pi_1^{\text{top}}(X(\mathbf{C}):*,*')$.

Assumption 1.11. Further we assume that (\overline{X}, D) has a real structure $(\overline{X'}, D')$, i.e. $(\overline{X}, D) = (\overline{X'} \times_{\mathbf{R}} \mathbf{C}, D' \times_{\mathbf{R}} \mathbf{C})$,

Let F_{∞} be an involution of $X(\mathbf{C})$ induced from the complex conjugation $z \in \mathbf{C} \mapsto \bar{z} \in \mathbf{C}$ and its induced automorphism on T_s^{\times} ($s \in D$). The pull-back functor $F_{\infty}^{*} : \mathcal{NL}^{\operatorname{Be}}(X(\mathbf{C})) \to \mathcal{NL}^{\operatorname{Be}}(X(\mathbf{C}))$ and $F_{\infty}^{*} : \mathcal{NL}^{\operatorname{Be}}(T_s^{\times}(\mathbf{C})) \to \mathcal{NL}^{\operatorname{Be}}(T_s^{\times}(\mathbf{C}))$ determine equivalences of tannakian categories. We note that they are compatible with (1.8). They induce the isomorphism

$$F_{\infty}: \pi_1^{\operatorname{Be}}(X(\mathbf{C}):*,*') \to \pi_1^{\operatorname{Be}}(X(\mathbf{C}):\bar{*},\bar{*}')$$

where $\bar{*}$ (resp. $\bar{*}'$) means $F_{\infty}(*)$ (resp. $F_{\infty}(*')$).

Definition 1.12. The infinity Frobenius action ϕ_{∞} stands for the inverse of F_{∞} above.

The description of the infinity Frobenius action in case of $X = \mathbf{P}^1 \setminus \{0, 1, \infty\}$ will be discussed in §2.

1.4. **étale setting.** We will briefly recall the definition of the étale fundamental group, torsor and the tangential base point in [De1]§15.

Notation 1.13. In this subsection, we assume that K is a field of characteristic 0. We denote \overline{K} to be its algebraic closure and $X_{\overline{K}}$ to be the scalar extension of X_K by \overline{K} . Let l be a prime (we do not need to assume that l is inequal to the prime p in §1.2). We denote the category of the nilpotent part of the category of étale local systems ⁹ of finite dimensional \mathbf{Q}_l -vector spaces over $X(\overline{K})$ by $\mathcal{NE}^l(X_{\overline{K}})$. It also forms a neutral tannakian category over \mathbf{Q}_l .

In the étale realization, we consider two fiber functors. Suppose that $x \in X(\overline{K})$. The first one is the pullback $\omega_x : \mathcal{NE}^l(X_{\overline{K}}) \to Vec_{\mathbf{Q}_l}$ which associates each nilpotent étale local system V with the pullback V(x). The second one is the tangential base point: Let $s \in D(\overline{K})$ and put $T_s = T_s \overline{X}_{\overline{K}}$ and $T_s^{\times} = T_s \setminus \{0\} (\cong \mathbf{G}_m)$, where $T_s \overline{X}_{\overline{K}}$ is the tangent vector space of $\overline{X}_{\overline{K}}$ at s. By [De1]§§15.13-15.26, we have a morphism of tannakian categories

(1.9)
$$Res_s : \mathcal{NE}^l(X_{\overline{K}}) \to \mathcal{NE}^l(T_s^{\times})$$

which we call the tangential morphism. By pulling back, each \overline{K} -valued point t_s in T_s^{\times} determines a fiber functor $\omega_{t_s} : \mathcal{NE}^l(X_{\overline{K}}) \to Vec_{\mathbf{Q}_l}$.

The above fiber functors, ω_x and ω_{t_s} , play a role of 'basepoints' and the following is a reformulation of the *l*-adic étale fundamental group and torsor in [De1] §13.

Definition 1.14. Let ω_* and $\omega_{*'}$ be any fiber functors above. The *l*-adic étale fundamental group $\pi_1^{l,\text{\'et}}(X_{\overline{K}}:*)$ is $\underline{Aut}^{\otimes}(\omega_*)$ and the *l*-adic étale fundamental torsor $\pi_1^{l,\text{\'et}}(X_{\overline{K}}:*,*')$ is $\underline{Isom}^{\otimes}(\omega_*,\omega_{*'})$.

The former is a pro-algebraic group over \mathbf{Q}_l and the latter is a pro-algebraic bitorsor over \mathbf{Q}_l where $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*)$ acts on the right and $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*')$ acts on the left. Let $\overline{K'}$ be an algebraically closed field containing \overline{K} . The scalar extension induces a categorical equivalence $\mathcal{NE}^l(X_{\overline{K'}}) \cong \mathcal{NE}^l(X_{\overline{K}})$, from which we

⁹We mean covariant functors $V : \hat{\Pi}_{X_{\overline{K}}} \to Vec_{\mathbf{Q}_l}$ where $\hat{\Pi}_{X_{\overline{K}}}$ stands for the category whose objects are points on $X(\overline{K})$ and whose set of morphisms between two points * and *' on $X(\overline{K})$ is the pro-finite fundamental torsor $\hat{\pi}_1(X_{\overline{K}}:*,*')$ in [G].

get isomorphisms $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{K'}}:*) \cong \pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*)$ and $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{K'}}:*,*') \cong \pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*,*') \cong \pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*') \cong \pi_1^{l,\text{\acute{e}t}}(X_$

Suppose that * and *' are K-valued points on X or T_s^{\times} for $s \in D$ (this assumption is attained by enlarging the base field K). Let $\sigma \in Gal(\overline{K}/K)$. We get Galois action

$$\sigma: \pi_1^{l,\text{\'et}}(X_{\overline{K}}:*,*') \to \pi_1^{l,\text{\'et}}(X_{\overline{K}}:*,*')$$

from the Galois action $id \otimes \sigma : X_{\overline{K}} \to X_{\overline{K}}$ in a similar way to our construction of F_{q*} in §1.2 and $F_{\infty*}$ in §1.3. We note that the pro-finite fundamental group $\hat{\pi}_1(X_{\overline{K}}:*)$ (resp. torsor $\hat{\pi}_1(X_{\overline{K}}:*,*')$) also admits Galois action and we have a morphism from $\hat{\pi}_1(X_{\overline{K}}:*) \to \pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*)(\mathbf{Q}_l)$ (resp. $\hat{\pi}_1(X_{\overline{K}}:*,*') \to \pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*,*')(\mathbf{Q}_l)$) which is compatible with its group (resp. torsor) structure and their Galois group actions. Actually this morphism induces an isomorphism between $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{K}}:*)$ and the Malcev (unipotent) completion over \mathbf{Q}_l of $\hat{\pi}_1(X_{\overline{K}}:*)$ [De1] §10.

Lemma 1.15. Assume that K is a number field with ring of integers \mathcal{O}_K . Let * and *' be any K-valued points. Let \mathfrak{p} be a prime ideal, coprime to $l\mathcal{O}_K$, where the triple (X, *, *') has a proper smooth model over $\mathcal{O}_{K,\mathfrak{p}}$ with a good reduction. Then there exists a unique path $e_{*,*'}^{\mathfrak{p}} \in \pi_1^{l,\acute{e}t}(X_{\overline{K}}:*,*')(\mathbf{Q}_l)$ which is invariant under the action of Frobenius element $Frob_{\mathfrak{p}}$ of \mathfrak{p} , i.e. $Frob_{\mathfrak{p}}(e_{*,*'}^{\mathfrak{p}}) = e_{*,*'}^{\mathfrak{p}}$.

Proof. By the above assumption $\pi_1^{l,\text{\'et}}(X_{\overline{K}}:*,*')$ is unramified at \mathfrak{p} , whence we have an action of the geometric Frobenius $Frob_{\mathfrak{p}}$ of \mathfrak{p} . Since the eigenvalues of the $Frob_{\mathfrak{p}}$ -action on $H^1_{\text{\'et}}(X_{\overline{K}}, \mathbf{Q}_l)$ are positive weight Weil numbers, we obtain a unique Frobenius invariant path $e^{\mathfrak{p}}_{*,*'}$ by an inductive construction same to [Bes]§3 (or [Vol]§4.3).

We note that this Frobenius invariant path $e_{*,*'}^{\mathfrak{p}}$ equips nice compatibilities such as functorialities and $e_{*',*''}^{\mathfrak{p}} \cdot e_{*,*'}^{\mathfrak{p}} = e_{*,*''}^{\mathfrak{p}}$ similar to $d_{*,*'}$ in §1.1 and $c_{*,*'}$ in §1.2. As Besser [Bes] gave the Frobenius invariant path $c_{*,*'}$ an interpretation of Coleman's *p*-adic iterated integration in the rigid setting, the author thinks that there may be also a significance on the above special path $e_{*,*'}^{\mathfrak{p}}$ in the *l*-adic étale setting.

2. TANNAKIAN INTERPRETATIONS

This section is divided into three sides. The first side is the Berthelot-Ogus side (§2.1) where we shall give our main results. A tannakian interpretation of p-adic MPL's and p-adic MZV's (p: prime)¹⁰ is given in Theorem 2.3 and Theorem 2.5. By describing Frobenius action on the rigid fundamental group of the projective line minus three points we deduce an explicit formula between our p-adic MZV's and Deligne's p-adic MZV's [De2, DG] in Theorem 2.8. Further by considering Frobenius action on a rigid path space we introduce an overconvergent version of p-adic MPL in Definition 2.13. Its expression in terms of our p-adic MPL's is given in Theorem 2.14. The second side is the Hodge side (§2.2), where results analogous to §2.1 are presented. A tannakian reinterpretation of MPL's and MZV's are given in Proposition 2.20 and Proposition 2.21 following [BD]. Using the infinity Frobenius action on the Betti fundamental torsor of the projective line minus three points we introduce a new version of MPL (2.20). Its monodromy-free property is shown in Theorem 2.27. Its description is given in Proposition 2.26. The third side is the

 $^{^{10}}$ MPL stands for multiple polylogarithm and MZV stands for multiple zeta values.

Artin side ($\S2.3$). We discuss *l*-adic étale analogues of MPL's and MZV's defined in a way that Wojtkowiak [W] introduced the *l*-adic polylogarithms.

Throughout this and next section, we fix $X = \mathbf{P}^1 \setminus \{0, 1, \infty\} = Spec \mathbf{Q}[t, \frac{1}{t}, \frac{1}{t-1}] \subset \overline{X} = \mathbf{P}^1$ and $D = \{0, 1, \infty\}$. We denote the local parameter of T_0^{\times} induced from t by t_0 and the point $t_0 = 1$ on T_0^{\times} by $\overrightarrow{01}$. Similarly we denote the local parameter of T_1^{\times} induced from 1 - t by t_1 and the point $t_1 = 1$ on T_1^{\times} by $\overrightarrow{10}$.

2.1. Berthelot-Ogus side — main results. In this side, we shall use a Berthelot-Ogus-type comparison isomorphism (Lemma 2.2) between a de Rham fundamental torsor and a rigid fundamental torsor for X. Our first results are a tannakian interpretation of p-adic MPL's and p-adic MZV's in Theorem 2.3 and Theorem 2.5 respectively. Secondly we consider Frobenius action on a rigid fundamental group. The definition of Deligne's p-adic MZV's [De2, DG] which describes the Frobenius action is recalled. By using a tangential morphism we give an explicit formula between his and ours in Theorem 2.8. Thirdly we consider Frobenius action on a rigid path space. Overconvergent p-adic MPL is introduced in Definition 2.13. It describes the Frobenius action and takes Deligne's p-adic MZV as 'a special value' at z = 1. A formula to express it in terms of our p-adic MPL's is given in Theorem 2.14.

Before we state our first results we give notations and basic lemma below which will be frequently employed in this subsection.

Notation 2.1. We denote the special path $d_{\overrightarrow{01},z}$ (§1.1) in the de Rham fundamental torsor $\pi_1^{\mathrm{DR}}(X:\overrightarrow{01},z)(\mathbf{Q}_p)$ for $z \in X(\mathbf{Q}_p)$ by d_z . In particular we simply denote $d_{\overrightarrow{01},\overrightarrow{10}}$ in by d. Put $X_0 = \mathbf{P}_{\mathbf{F}_p}^1 \setminus \{0,1,\infty\} \subset \overline{X_0} = \mathbf{P}_{\mathbf{F}_p}^1$. Similarly we denote the special path $c_{\overrightarrow{01},z_0}$ (§1.2) in the rigid fundamental torsor $\pi_1^{p,\mathrm{rig}}(X_0:\overrightarrow{01},z_0)(\mathbf{Q}_p)$ for $z_0 \in \overline{X_0}(\mathbf{F}_p)$ by c_{z_0} . In particular we simply denote $c_{\overrightarrow{01},\overrightarrow{10}}$ by c. The composition of the embedding $\pi_1^{\mathrm{DR}}(T_0^{\times}:\overrightarrow{01}) \to \pi_1^{\mathrm{DR}}(X:\overrightarrow{01})$ by (1.1) and the exponential map $H_1^{\mathrm{DR}}(T_0^{\times},\mathbf{Q}) \to \pi_1^{\mathrm{DR}}(T_0^{\times},\overrightarrow{01})(\mathbf{Q})$ sends the dual of $\frac{dt_0}{t_0}$ to a loop denoted by x. Similarly a composite morphism to $\pi_1^{\mathrm{DR}}(T_0^{\times},\overrightarrow{10})(\mathbf{Q})$ Then it is easy to see that x and y are free generators of the pro-algebraic group $\pi_1^{\mathrm{DR}}(X:\overrightarrow{01})$. In this section, we consider the embedding into the non-commutative formal power series with two variables A and B

(2.1)
$$i: \pi_1^{\mathrm{DR}}(X:\overline{01})(\mathbf{Q}) \hookrightarrow \mathbf{Q}\langle\langle A, B \rangle\rangle$$

which sends $x \mapsto e^A := 1 + \frac{A}{1!} + \frac{A^2}{2!} + \cdots$ and $y \mapsto e^B := 1 + \frac{B}{1!} + \frac{B^2}{2!} + \cdots$. We note that this morphism is isomorphic to the embedding of $\pi_1^{\mathrm{DR}}(X:\overrightarrow{01})(\mathbf{Q})$ into the universal embedding algebra of $Lie\pi_1^{\mathrm{DR}}(X:\overrightarrow{01})(\mathbf{Q})$. By abuse of notation we also employ the same symbol i when we mention its scalar extension.

The following Berthelot-Ogus type comparison isomorphism for fundamental torsors is our basics in this subsection.

Lemma 2.2. Let z be a \mathbf{Q}_p -valued point of X or T_s^{\times} (for $s \in D$) and z_0 be its modulo p reduction.

(1) If z_0 lies on $X_0(\mathbf{F}_p)$ or $T_{s_0}^{\times}(\mathbf{F}_p)$,

$$T_1^{\mathrm{DR}}(X_{\mathbf{Q}_p}:\overrightarrow{01},z) \cong \pi_1^{p,\mathrm{rig}}(X_0:\overrightarrow{01},z_0).$$

(2) If z_0 does not lie on $X_0(\mathbf{F}_p)$ nor $T_{s_0}^{\times}(\mathbf{F}_p)$,

$$\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}_p}:\overline{01},z) \times \mathbf{Q}_{p,st} \cong \pi_1^{p,\mathrm{rig}}(X_0:\overline{01},z_0) \times \mathbf{Q}_{p,\mathrm{st}}$$

where $\mathbf{Q}_{p,st}$ stands for the polynomial algebra $\mathbf{Q}_p[l(p)]$ with one variable l(p).

Proof . An equivalence of tannakian categories

(2.2)
$$\mathcal{NC}^{\mathrm{DR}}(X_{\mathbf{Q}_p}) \cong \mathcal{NI}^{\dagger}(X_{\mathbf{F}_p})$$

is shown in [CLS, S2] in more general situation. In case of $z_0 \in X_0(\mathbf{F}_p)$ it gives rise to a natural isomorphism

$$\omega_z(\mathcal{V}, \nabla) \cong \omega_{z_0}(\mathcal{V}^{\dagger}, \nabla^{\dagger})$$

for $(\mathcal{V}, \nabla) \in \mathcal{NC}^{\mathrm{DR}}(X_{\mathbf{Q}_p})$, where $(\mathcal{V}^{\dagger}, \nabla^{\dagger}) \in \mathcal{NI}^{\dagger}(X_{\mathbf{F}_p})$ is associated with (\mathcal{V}, ∇) via (2.2). The equivalences (2.2) and $\mathcal{NC}^{\mathrm{DR}}(T_{s,\mathbf{Q}_p}^{\times}) \cong \mathcal{NI}^{\dagger}(T_{s_0}^{\times})$ are compatible with tangential morphisms (cf. [BF]). They also give rise to an natural isomorphism

$$\omega_z \Big(Res(\mathcal{V}, \nabla) \Big) \cong \omega_{z_0} \Big(Res(\mathcal{V}^{\dagger}, \nabla^{\dagger}) \Big)$$

for $(\mathcal{V}, \nabla) \in \mathcal{NC}^{\mathrm{DR}}(X_{\mathbf{Q}_p})$ in case of $z_0 \in T_0^{\times}(\mathbf{F}_p)$. The equivalence (2.2) and the above two isomorphisms of fiber functors follow the isomorphism in (1). In the rest case a natural isomorphism

$$\omega_z(\mathcal{V},\nabla)\otimes\mathbf{Q}_p[l(p)]\cong\omega_{z_0}(\mathcal{V}^{\dagger},\nabla^{\dagger})\otimes\mathbf{Q}_p[l(p)]$$

is explained in [Vol] $\S4.4$, which follows the isomorphism in (2).

We note that this variable l(p) reflects a branch parameter $a \in \mathbf{Q}_p$ of the *p*-adic polylogarithm \log^a . By identifying the above two fundamental torsors by Lemma 2.2 we get a de Rham loop $d_z^{-1}c_{z_0}$, which lies in $\pi_1^{\mathrm{DR}}(X:\overline{01})(\mathbf{Q}_p)$ in case of (1) and in $\pi_1^{\mathrm{DR}}(X:\overline{01})(\mathbf{Q}_{p,\mathrm{st}})$ in case of (2).

Theorem 2.3. Let z be a \mathbf{Q}_p -valued point of X and z_0 be its modulo p reduction on $\overline{X_0}$. The de Rham loop $d_z^{-1}c_{z_0}$ corresponds to the special value of the fundamental solution G_0^{-11} at z of the p-adic KZ equation by the embedding i:

$$i(d_z^{-1}c_{z_0}) = G_0(z).$$

Proof. Our main tools to prove the theorem are a tannakian description of the KZ equation and Besser's tannakian interpretation of Coleman's p-adic iterated integration theory.

We start with the KZ equation. Let $(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}})$ be a pro-object of $\mathcal{NC}^{\mathrm{DR}}(X)$ associated with the KZ equation. Namely $\mathcal{V}_{\mathrm{KZ}} = \mathcal{O}_X \langle \langle A, B \rangle \rangle$ and $\nabla_{\mathrm{KZ}}(g) = dg - (\frac{A}{t} + \frac{B}{t-1})gdt$ for local section g. By [De1] Proposition 12.10, the Lie algebra $\mathfrak{p}^{\mathrm{DR}}$ of $\pi_1^{\mathrm{DR}}(X:\Gamma)$ is free generated by two representatives $(\frac{dt}{t})^*$ and $(\frac{dt}{t-1})^*$ in $H_1^{\mathrm{DR}}(X, \mathbf{Q})$ which are dual basis of $\frac{dt}{t}$ and $\frac{dt}{t-1}$ in $H_{\mathrm{DR}}^1(X, \mathbf{Q}) \cong H^0(\overline{X}, \Omega\frac{1}{\overline{X}}(\log D))$ respectively

¹¹In case of $z_0 \in X_0(\mathbf{F}_p)$ it means the fundamental solution of the *p*-adic KZ (Knizhnik-Zamolodchikov) equation in $\mathbf{Q}_p\langle\langle A, B\rangle\rangle$ constructed in [F1]Theorem 3.3 and in case of $z_0 \notin X_0(\mathbf{F}_p)$ it means the series in $\mathbf{Q}_{p,st}\langle\langle A, B\rangle\rangle$ whose specialization at l(p) = a is $G_0^a(z)$ in loc.cit.

because $H^1(\overline{X}, \mathcal{O}_{\overline{X}}) = 0$. By the basic theorem [DM] of tannakian categories, the fiber functor ω_{Γ} induces an equivalence of categories

(2.3)
$$\mathcal{NC}^{\mathrm{DR}}(X_K) \cong \mathcal{N}Rep \ \mathfrak{p}^{\mathrm{DR}}$$

where the right hand side stands for the category of nilpotent representations of the Lie algebra on finite dimensional \mathbf{Q} -vector spaces.

Lemma 2.4. By (2.3) the pro-object $(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}})$ corresponds to the vector space $\mathbf{Q}\langle\langle A, B \rangle\rangle$ with the left multiplication $L : \mathfrak{p}^{\mathrm{DR}} \to \mathfrak{gl}(\mathbf{Q}\langle\langle A, B \rangle\rangle)$ such that $L\left(\left(\frac{dt}{t}\right)^*\right) = L_A$ and $L\left(\left(\frac{dt}{t-1}\right)^*\right) = L_B$. Here L_F for $F \in \mathbf{Q}\langle\langle A, B \rangle\rangle$ means the left multiplication by F.

Proof. In [De1]§12.5, Deligne give a recipe of the correspondence (2.3) (in more general situation): Let $(\mathcal{V}, \nabla) \in \mathcal{NC}^{\mathrm{DR}}(X)$. We put $V = \Gamma(\overline{X}, \mathcal{V}_{\mathrm{can}}) = \omega_{\Gamma}(\mathcal{V}, \nabla)$ and decompose as $\nabla_{\mathrm{can}} = d + \omega$ where $d : V \otimes \mathcal{O}_{\overline{X}} \to V \otimes \Omega_{\overline{X}}^{1}(\log D)$ is the differential induced from $d : \mathcal{O}_{\overline{X}} \to \Omega_{\overline{X}}^{1}(\log D)$ and $\omega \in H^{0}(\overline{X}, \Omega_{\overline{X}}^{1}(\log D)) \otimes EndV$. By the integrability of ∇ , we have the Lie algebra homomorphism $\rho : \mathfrak{p}^{\mathrm{DR}} \to \mathfrak{gl} V$ such that $\rho|_{H_{1}^{\mathrm{DR}}(X,\mathbf{Q})} = -\omega$, which gives an object $(V,\rho) \in \mathcal{N}Rep \ \mathfrak{p}^{\mathrm{DR}}$. Conversely let $(V,\rho) \in \mathcal{N}Rep \ \mathfrak{p}^{\mathrm{DR}}$. We put $\omega = -\rho|_{H_{1}^{\mathrm{DR}}(X,\mathbf{Q})}(\alpha) \in H_{\mathrm{DR}}^{1}(X,\mathbf{Q}) \otimes EndV \cong H^{0}(\overline{X},\Omega_{\overline{X}}^{1}(\log D)) \otimes EndV$ where $\alpha \in H_{\mathrm{DR}}^{1}(X,\mathbf{Q}) \otimes H_{1}^{\mathrm{DR}}(X,\mathbf{Q})$ is a canonical tensor representing the identity on $H_{1}^{\mathrm{DR}}(X,\mathbf{Q})$. This gives $(\mathcal{V},\nabla) = (V \otimes \mathcal{O}_X, d_X + \omega|_X) \in \mathcal{N}C^{\mathrm{DR}}(X)$.

Following his recipe we see that L corresponds to

$$\omega = \frac{dt}{t} L_A + \frac{dt}{t-1} L_B \in H_1^{\mathrm{DR}}(X, \mathbf{Q}) \widehat{\otimes} \mathfrak{gl}(\mathbf{Q} \langle \langle A, B \rangle \rangle)$$

which gives the claim.

By abuse of notation, we denote its induced group homomorphism by

 $L: \pi_1^{\mathrm{DR}}(X:\Gamma) \to Aut \mathbf{Q}\langle\langle A, B\rangle\rangle.$

By the constructions of i and L above, we have

Here Int $d_{\overrightarrow{01},\Gamma} : \pi_1^{\mathrm{DR}}(X:\overrightarrow{01})(\mathbf{Q}) \to \pi_1^{\mathrm{DR}}(X:\Gamma)(\mathbf{Q})$ is the automorphism sending $\gamma \mapsto d_{\overrightarrow{01},\Gamma} \cdot \gamma \cdot d_{\overrightarrow{01},\Gamma}^{-1}$ and $ev_1 : Aut\mathbf{Q}\langle\langle A, B \rangle\rangle \to \mathbf{Q}\langle\langle A, B \rangle\rangle$ is the evaluation at 1 sending $\sigma \mapsto \sigma(1)$.

Our second tool is Besser's tannakian interpretation of the Coleman's *p*-adic iterated integration theory [C] in [Bes]: He called a set $v = \{v_{x_0} \in \omega_{x_0}(\mathcal{V}^{\dagger}, \nabla^{\dagger}) | x_0 \in \overline{X_0}(\overline{\mathbf{F}_p})\}$ for $(\mathcal{V}^{\dagger}, \nabla^{\dagger}) \in \mathcal{NI}^{\dagger}(X_0)$ a collection of analytic continuation along Frobenius of horizontal section of $(\mathcal{V}^{\dagger}, \nabla^{\dagger})$ if $c_{x_0, x'_0}(v_{x_0}) = v_{x'_0}$ for all $x_0, x'_0 \in \overline{X_0}(\overline{\mathbf{F}_p})$ where c_{x_0, x'_0} is a Frobenius invariant path in Lemma 1.8 and showed that each Coleman function of $X(\mathbf{C}_p)$ (in the sense of [Bes] §5) is expressed as a set $\{\theta(v_{x_0})\}_{x_0}$ with a $j^{\dagger}\mathcal{O}_{]\overline{X_0}[} \otimes \mathbf{C}_p$ -module homomorphism $\theta : \mathcal{V}^{\dagger} \otimes \mathbf{C}_p \to j^{\dagger}\mathcal{O}_{]\overline{X_0}[} \otimes \mathbf{C}_p$ and a collection $v = \{v_{x_0}\}_{x_0}$ of analytic continuation along Frobenius of a pair $(\mathcal{V}^{\dagger}, \nabla^{\dagger}) \in \mathcal{NI}^{\dagger}(X_0)$. In [BF] we further extend Coleman functions to normal bundles by making use of Frobenius invariant path to tangential basepoints.

In our previous paper [F1] we consider the *p*-adic KZ equation and construct its fundamental solution $G_0(t)$, a two variable non-commutative formal power series

with Coleman function coefficients. In his terminologies it is a collection $\{G_0|_{]x_0}[\}_{x_0}$ of analytic continuation along Frobenius of horizontal section of $(\mathcal{V}_{\text{KZ}}^{\dagger}, \nabla_{\text{KZ}}^{\dagger})$ (the associated pro-object of $\mathcal{NI}^{\dagger}(X_0)$ with $(\mathcal{V}_{\text{KZ}}, \nabla_{\text{KZ}}))$, i.e. $c_{x_0,y_0}(G_0|_{]x_0}] = G_0|_{]y_0}[$ for x_0 and y_0 . And since in our case the tangential morphism (1.1) for s = 0sends $(\mathcal{V}_{\text{KZ}}, \nabla_{\text{KZ}})$ to $(\mathcal{O}_{T_0^{\times}}\langle\langle A, B\rangle\rangle, d - A\frac{dt_0}{t_0}), t_0^A = 1 + \frac{\log^a t_0}{1!}A + \frac{(\log^a t_0)^2}{2!}A^2 + \frac{(\log^a t_0)^3}{3!}A^3 + \cdots$ might be a collection of analytic continuation along Frobenius of horizontal section of $Res_0(\mathcal{V}_{\text{KZ}}^{\dagger}, \nabla_{\text{KZ}}^{\dagger})$. By our tangential base point interpretation ([BF] Proposition 2.11) of the notion of the constant term ([BF] Definition 2.1) of horizontal sections, we see that t_0^A is an analytic continuation along Frobenius of $G_0(t)$ to T_0^{\times} because both the constant term of $G_0(t)$ at t = 0 and the constant term of t_0^A at $t_0 = 0$ are equal to 1. Namely $c_{\overrightarrow{01},y_0}(t_0^A|_{|\overrightarrow{01}|}) = G_0(t)|_{]y_0}[$.

Let z be a \mathbf{Q}_p -valued point X with reduction z_0 . We regard c_{z_0} to be a de Rham path via the identification in Lemma 2.4. The isomorphism $d_{*,\Gamma} : \omega_*(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}}) \to \omega_{\Gamma}(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}}) = \mathbf{Q}_p \langle \langle A, B \rangle \rangle$ for $* = \overrightarrow{01}$ and z let us rewrite

$$(2.5) d_{z,\Gamma} \cdot c_{z_0} \cdot d_{\Gamma,\overrightarrow{01}}(1) = G_0(z).$$

By (2.4), (2.5) and
$$d_{z,\Gamma} = d_{\overrightarrow{01},\Gamma} \cdot d_z^{-1}$$
, we prove Theorem 2.3.

In our previous paper [F1] Theorem 3.15 we showed that in each coefficient of $G_0(z)$ there appears *p*-adic MPL $Li_{k_1,\dots,k_m}(z)$, which is a Coleman function admitting the expansion (0.1) on $|z|_p < 1$. In precise the coefficient of $A^{k_m-1}B\cdots A^{k_1-1}B$ in $G_0(z)$ is $(-1)^m Li_{k_1,\dots,k_m}(z)$:

 $G_0(z) = 1 + \log zA - Li_1(z)B + \dots + (-1)^m Li_{k_1,\dots,k_m}(z)A^{k_m-1}B \cdots A^{k_1-1}B + \dots$

By Theorem 2.3 we may say that the loop $d_z^{-1}c_{z_0}$ is a tannakian origin of *p*-adic MPL's.

In case when z is also a tangential base point $\overline{01}$, similarly we have

Theorem 2.5. The de Rham loop $d^{-1}c \in \pi_1^{DR}(X : \overrightarrow{01})(\mathbf{Q}_p)$ corresponds to the *p*-adic Drinfel'd associator $\Phi_{KZ}^p \in \mathbf{Q}_p(\langle A, B \rangle)$ by the embedding *i*, *i.e.*

$$i(d^{-1}c) = \Phi_{\mathrm{KZ}}^p.$$

Proof. The *p*-adic Drinfel'd associator Φ_{KZ}^p is the series constructed in [F1] Definition 3.12 which is equal to the limit value ([F1] Lemma 3.27)

(2.6)
$$\lim_{\epsilon \to 0}' \exp\left(-\log^a \epsilon \cdot B\right) \cdot G_0^a(1-\epsilon) \in \mathbf{Q}_p(\langle A, B \rangle)$$

In other word it is the constant term of $G_0^a(z)$ at z = 1. Then by a similar argument to the proof of Theorem 2.3 we have $c_{\overrightarrow{01},\overrightarrow{10}}(t_0^A|_{|\overrightarrow{01}|}) = \Phi_{\text{KZ}}^p \cdot t_1^B|_{|\overrightarrow{10}|}$, whence

(2.7)
$$d_{\overrightarrow{10},\Gamma} \cdot c \cdot d_{\Gamma,\overrightarrow{01}}(1) = \Phi_{\mathrm{KZ}}^p$$

By (2.4), (2.16) and $d_{\overrightarrow{10},\Gamma} = d_{\overrightarrow{01},\Gamma} \cdot d^{-1}$, we prove Theorem 2.5.

In our previous paper [F1] Theorem 3.30 we showed that in each coefficient of the *p*-adic Drinfel'd associator Φ_{KZ}^p there appears *p*-adic MZV $\zeta_p(k_1, \dots, k_m)$ $(k_m > 1)$ introduced in [F1], which is a *p*-adic analogue of (0.2). In precise the coefficient of $A^{k_m-1}B \cdots A^{k_1-1}B$ in Φ_{KZ}^p is $(-1)^m \zeta_p(k_1, \dots, k_m)$:

$$\Phi_{\rm KZ}^p = 1 + \dots + (-1)^m \zeta_p(k_1, \dots, k_m) A^{k_m - 1} B \dots A^{k_1 - 1} B + \dots$$

By Theorem 2.5 we may say that the loop $d^{-1}c$ is a tannakian origin of *p*-adic MZV's.

Note 2.6. The *p*-adic Drinfel'd associator Φ_{KZ}^p is group-like [F1], that means $\Delta \Phi_{KZ}^p = \Phi_{KZ}^p \widehat{\otimes} \Phi_{KZ}^p$ where $\Delta : \mathbf{Q}_p \langle \langle A, B \rangle \rangle \to \mathbf{Q}_p \langle \langle A, B \rangle \rangle$ is the linear map induced from $\Delta(A) = A \otimes 1 + 1 \otimes A$ and $\Delta(B) = B \otimes 1 + 1 \otimes B$. Hence each coefficient of Φ_{KZ}^p must satisfy (integral) shuffle product formula (cf. [F1], [FJ]). We can recover and express general coefficients of Φ_{KZ}^p in terms of $\zeta_p(k_1, \dots, k_m)$'s $(k_m > 1)$ by the following method: Let W be a word, i.e. a monic and monomial element in $\mathbf{Q}_p \langle \langle A, B \rangle \rangle$. Put C_W to be its coefficient of Φ_{KZ}^p . We have $C_A = C_B = 0$. For a convergent word W written as $A^{k_m - 1}B \cdots A^{k_1 - 1}B$ $(k_m > 1)$, $C_W = (-1)^m \zeta_p(k_1, \dots, k_m)$. For a divergent word W' written as $B^r W$ (r > 0) with W: convergent, shuffle product formula gives

$$C_B^r \cdot C_W = r! C_{B^r W}$$
 + other terms.

By an induction with respect to r, $C_{W'}$ is calculated and is expressed in terms of C_W 's with W: convergent. For a word $W'' = B^r W A^s$ (s > 0) with W: convergent, a similar induction argument let us able to calculate $C_{W''}$.

Our second result is on a description of a Frobenius action. We consider a Frobenius action F_p on $\pi_1^{\text{DR}}(X:\overrightarrow{01},\overrightarrow{10})(\mathbf{Q}_p)$ by transmitting that on $\pi_1^{p,\text{rig}}(X_0:\overrightarrow{01},\overrightarrow{10})(\mathbf{Q}_p)$ by Lemma 2.2. It gives a new path $F_p(d)$ in $\pi_1^{\text{DR}}(X:\overrightarrow{01},\overrightarrow{10})(\mathbf{Q}_p)$ which is our central object to discuss here. In Arizona Winter School 2002, Deligne [De2] introduced another version of *p*-adic MZV which has a different tannakian origin from ours (see also [DG] §5).

Definition 2.7. The *p*-adic Deligne associator Φ_{De}^p is the series in $\mathbf{Q}_p\langle\langle A, B\rangle\rangle$ which corresponds to $d^{-1}\phi_p(d)$:

$$i(d^{-1}\phi_p(d)) = \Phi_{\mathrm{De}}^p.$$

To adapt the notations to ours above, we denote $\zeta_p^{\text{De}}(k_1, \dots, k_m)$ $(k_m > 1)$ to be the coefficient of $A^{k_m-1}B \cdots A^{k_1-1}B$ in Φ_{De}^p multiplied by $(-1)^m$:

$$\Phi_{\rm De}^p = 1 + \dots + (-1)^m \zeta_p^{\rm De}(k_1, \dots, k_m) A^{k_m - 1} B \dots A^{k_1 - 1} B + \dots$$

and call it Deligne's *p*-adic MZV. Since Φ_{De}^p is group-like, we can recover and express general coefficients of Φ_{De}^p in terms of $\zeta_p^{De}(k_1, \dots, k_m)$'s $(k_m > 1)$ as Note 2.6. Deligne's *p*-adic MZV's come from the loop $d^{-1}\phi_p(d)$ while ours come from the loop $d^{-1}c$. So his are different from ours. But we can give a close relationship between them:

Theorem 2.8.

(2.8)
$$\Phi_{\rm KZ}^p(A,B) = \Phi_{\rm De}^p(A,B) \cdot \Phi_{\rm KZ}^p\left(\frac{A}{p}, \Phi_{\rm De}^p(A,B)^{-1}\frac{B}{p}\Phi_{\rm De}^p(A,B)\right).$$

Here the last term means the series substituting $(\frac{A}{p}, \Phi_{\text{De}}^p(A, B)^{-1}\frac{B}{p}\Phi_{\text{De}}^p(A, B))$ in $\Phi_{\text{KZ}}^p(A, B)$.

This formula looks nasty. But according to the convention ¹² in [F0] of the Grothendieck-Teichmüller group [Dr] we rewrite simpler $(p, \Phi_{KZ}^p) = (p, \Phi_{De}^p) \circ$

¹²Let k be a field with characteristic 0. Let $c_1, c_2 \in k$ and $g_1(A, B), g_2(A, B) \in k\langle\langle A, B \rangle\rangle$. The product $(c_2, g_2) \circ (c_1, g_1)$ is defined by $(c_1c_2, g_2(A, B)g_1(\frac{A}{c_2}, g_2(A, B)^{-1}\frac{B}{c_2}g_2(A, B)))$.

 $(1, \Phi_{\rm KZ}^p)$, i.e.

(2.9)
$$(p, \Phi_{\text{De}}^p) = (p, \Phi_{\text{KZ}}^p) \circ (1, \Phi_{\text{KZ}}^p)^{-1}$$

Proof. We extends the Frobenius action ϕ_p on $\pi_1^{DR}(X, \overrightarrow{01})(\mathbf{Q}_p)$ to $\mathbf{Q}_p\langle\langle A, B \rangle\rangle$ by *i*.

Lemma 2.9. $\phi_p(A) = \frac{A}{p}, \ \phi_p(B) = \Phi_{\text{De}}^p - \frac{1}{p} \Phi_{\text{De}}^p.$

Proof. By the compatibility of the Frobenius action on the tangential morphism $\mathbf{Q}_p(1) = \pi_1^{p,\mathrm{rig}}(T_0^{\times}:\overrightarrow{01})(\mathbf{Q}_p) \to \pi_1^{p,\mathrm{rig}}(X_0:\overrightarrow{01})(\mathbf{Q}_p)$ we have $\phi_p(x) = x^{\frac{1}{p}}$. Similarly we have $\phi_p(y') = y'^{\frac{1}{p}}$. Since Frobenius action is compatible with the torsor structure we have $\phi_p(y) = \phi_p(d^{-1}y'd) = (d^{-1}\phi_p(d))^{-1}d^{-1}y'^{\frac{1}{p}}d(d^{-1}\phi_p(d)) = (d^{-1}\phi_p(d))^{-1}y^{\frac{1}{p}}(d^{-1}\phi_p(d))$.

By
$$d^{-1}\phi_p(d) = d^{-1}c\phi_p(d^{-1}c)^{-1}$$
 we have
 $\Phi^p_{\text{De}} = \Phi^p_{\text{KZ}} \cdot \phi_p(\Phi^p_{\text{KZ}})^{-1}.$

By Lemma 2.9 we get Theorem 2.8.

By expanding our explicit formula, we can express our *p*-adic MZV's arising from Frobenius invariant path in terms of Deligne's *p*-adic MZV's describing Frobenius action very explicitly and vice versa. The following are the easiest examples.

Examples 2.10. (1)
$$\zeta_p^{\text{De}}(k) = (1 - \frac{1}{p^k})\zeta_p(k)$$
 $(k > 1).$
(2) $\zeta_p^{\text{De}}(a,b) = (1 - \frac{1}{p^{a+b}})\zeta_p(a,b) - (\frac{1}{p^b} - \frac{1}{p^{a+b}})\zeta_p(a)\zeta_p(b) - \sum_{r=0}^{a-1} (-1)^r (\frac{1}{p^{a-r}} - \frac{1}{p^{a+b}}) {\binom{b-1+r}{b-1}} \zeta_p(a-r)\zeta_p(b+r) - (-1)^a \sum_{s=0}^{b-1} (\frac{1}{p^{b-s}} - \frac{1}{p^{a+b}}) {\binom{a-1+s}{a-1}} \zeta_p(a+s)\zeta_p(b-s)$ $(b > 1).$

We may say that Deligne's *p*-adic MZV's are not equal to but equivalent to ours.

Our third result is on Frobenius structures. We give a Frobenius structure on $(\mathcal{V}_{\text{KZ}}^{\dagger}, \nabla_{\text{KZ}}^{\dagger})$. We introduce and discuss an overconvergent variant of the *p*-adic MPL which describes this structure.

Proposition 2.11. The pro-object $(\mathcal{V}_{KZ}^{\dagger}, \nabla_{KZ}^{\dagger})$ naturally admits a Frobenius structure

(2.10)
$$\phi: F_p^*(\mathcal{V}_{\mathrm{KZ}}^\dagger, \nabla_{\mathrm{KZ}}^\dagger) \to (\mathcal{V}_{\mathrm{KZ}}^\dagger, \nabla_{\mathrm{KZ}}^\dagger).$$

Proof. We just derive it from the Frobenius structure on $\mathcal{A}_{\overrightarrow{01}}^{\dagger}$ (Definition 1.6). The following arguments in the de Rham setting would help our understandings.

Lemma 2.12. The pro-object $(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}})$ is isomorphic to the pro-object $(\mathcal{A}_{\overline{01}}^{\mathrm{DR}})^{\vee}$, the dual of the ind-object $\mathcal{A}_{\overline{01}}^{\mathrm{DR}}$ (Definition 1.4).

Proof. By the categorical equivalence (2.3), $(\mathcal{A}_{\Gamma}^{\mathrm{DR}})^{\vee}$ corresponds to a natural representation ρ_{Γ} of $Lie \ \pi_1^{\mathrm{DR}}(X : \Gamma)$ on the universal enveloping algebra $U\pi_1^{\mathrm{DR}}(X : \Gamma)$ of $Lie \ \pi_1^{\mathrm{DR}}(X : \Gamma)$ (we note that $U\pi_1^{\mathrm{DR}}(X : \Gamma)$ is isomorphic to the dual H of the coordinate ring $\omega_{\Gamma}(\mathcal{A}_{\Gamma}^{\mathrm{DR}})$ of $\pi_1^{\mathrm{DR}}(X : \Gamma)$ by [A] Theorem 2.5.3 because $Lie \ \pi_1^{\mathrm{DR}}(X : \Gamma)$ is the set of primitive elements in H by loc.cit.§4.3). By (1.3) this representation ρ_{Γ} is induced from the left action of $\pi_1^{\mathrm{DR}}(X : \Gamma)$ on $\pi_1^{\mathrm{DR}}(X : \Gamma, \Gamma)$, thus $U\pi_1^{\mathrm{DR}}(X : \Gamma)$ is a free $U\pi_1^{\mathrm{DR}}(X : \Gamma)$ -module of rank 1 by ρ_{Γ} . Therefore we say that the

corresponding representations ρ_{Γ} and L (Lemma 2.4) are equivalent, whence we have $(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}}) \cong (\mathcal{A}_{\Gamma}^{\mathrm{DR}})^{\vee}$. On the other hand by the canonical isomorphism $d_{\overrightarrow{01},\Gamma}: \omega_{\overrightarrow{01}} \to \omega_{\Gamma}$ it is easy to see $(\mathcal{A}_{\overrightarrow{01}}^{\mathrm{DR}})^{\vee} \cong (\mathcal{A}_{\Gamma}^{\mathrm{DR}})^{\vee}$. Whence we get the claim. \Box

By the categorical equivalence (2.2) we deduce $(\mathcal{V}_{\text{KZ}}^{\dagger}, \nabla_{\text{KZ}}^{\dagger}) \cong (\mathcal{A}_{\vec{0}\vec{1}}^{\dagger})^{\vee}$ from Lemma 2.12. Consequently we get the Frobenius structure on $(\mathcal{V}_{\text{KZ}}^{\dagger}, \nabla_{\text{KZ}}^{\dagger})$ by transmitting from $\phi : \mathcal{A}_{\vec{0}\vec{1}}^{\dagger} \to F_p^* \mathcal{A}_{\vec{0}\vec{1}}^{\dagger}$. We get Proposition 2.11.

In the following section we introduce single-valued real-analytic MPL $Li_{k_1,\dots,k_m}^-(z)$. Here we introduce the following *p*-adic analogue $Li_{k_1,\dots,k_m}^{\dagger}(z)$. From now on we fix a lift of Frobenius by $\tilde{F}_p(t) = t^p$. By $\mathcal{V}_{\mathrm{KZ}} = \mathcal{O}_X\langle\langle A, B \rangle\rangle$, we have $\mathcal{V}_{\mathrm{KZ}}^{\dagger} = j^{\dagger}\mathcal{O}_{|\overline{X}_0|}\langle\langle A, B \rangle\rangle$.

Definition 2.13. The overconvergent *p*-adic MPL $Li_{k_1,\dots,k_m}^{\dagger}(z) \in j^{\dagger}\mathcal{O}_{]\overline{X}_0[}(]\overline{X}_0[)$ is the coefficient of $A^{k_m-1}B\cdots A^{k_1-1}B$ in $G_0^{\dagger}(z)$ multiplied by $(-1)^m$:

$$G_0^{\dagger}(z) = 1 + \dots + (-1)^m Li_{k_1,\dots,k_m}^{\dagger}(z) A^{k_m - 1} B \dots A^{k_1 - 1} B + \dots$$

where $G_0^{\dagger}(z) \in \mathcal{V}_{\mathrm{KZ}}^{\dagger}(]\overline{X}_0[)$ is the image of $1 \in \tilde{F}_p^*\mathcal{V}_{\mathrm{KZ}}^{\dagger}(]\overline{X}_0[) = \mathcal{V}_{\mathrm{KZ}}^{\dagger}(]\overline{X}_0[) = j^{\dagger}\mathcal{O}_{|\overline{X}_0[}(]\overline{X}_0[)\langle\langle A, B \rangle\rangle$ by (2.10).

We note that since $G_0^{\dagger}(z)$ is group-like we can recover and express general coefficients of $G_0^{\dagger}(z)$ in terms of $Li_{k_1,\dots,k_m}^{\dagger}(z)$ as Note 2.6. Our choice of \tilde{F}_p is a good lifting of the Frobenius of $\mathbf{P}_{\mathbf{F}_p}^{\dagger} \setminus \{1\}$. So the overconvergent *p*-adic MPL is analytic on]0[and] ∞ [, i.e. $Li_{k_1,\dots,k_m}^{\dagger}(z) \in j'^{\dagger}\mathcal{O}_{]\overline{X}_0[}(]\overline{X}_0[)$ with $j': \mathbf{P}_{\mathbf{F}_p}^{1} \setminus \{1\} \hookrightarrow \overline{X}_0$ while our *p*-adic MPL $Li_{k_1,\dots,k_m}(z)$ is a Coleman function, i.e. belong to the ring A_{Col} of Coleman functions of X which contains $j^{\dagger}\mathcal{O}_{]\overline{X}_0[}(]\overline{X}_0[)$. In Lemma 2.15 we see that overconvergent *p*-adic MPL's have a different tannakian origin from ours, so that they are different from ours. But we have a close relationship between them in $A_{\text{Col}}\langle\langle A, B \rangle\rangle$:

Theorem 2.14.

(2.11)
$$G_0^{\dagger}(A,B)(z) = G_0(A,B)(z) \cdot G_0\left(\frac{A}{p}, \Phi_{\mathrm{De}}^p(A,B)^{-1}\frac{B}{p}\Phi_{\mathrm{De}}^p(A,B)\right)(z^p)^{-1}.$$

Proof. Let z be a \mathbf{Q}_p -valued point of X whose modulo p reduction z_0 lies on $X(\mathbf{F}_p)$ (If p = 2, we need to enlarge the base field \mathbf{Q}_p). We have a de Rham path d_z in $\pi_1^{\mathrm{DR}}(X:\overrightarrow{01},z)(\mathbf{Q}_p)$ while we also have a de Rham path $\phi_p(d_{z^p})$ by $\pi_1^{\mathrm{DR}}(X:\overrightarrow{01},z) \times \mathbf{Q}_p \cong \pi_1^{p,\mathrm{rig}}(X_0:\overrightarrow{01},z_0) \cong \pi_1^{\mathrm{DR}}(X:\overrightarrow{01},z^p) \times \mathbf{Q}_p$ in Lemma 2.2. Identifying them we get a de Rham loop $d_z^{-1}\phi_p(d_{z^p})$ in $\pi_1^{\mathrm{DR}}(X:\overrightarrow{01})(\mathbf{Q}_p)$.

Lemma 2.15. $i(d_z^{-1}\phi_p(d_{z^p})) = G_0^{\dagger}(z).$

Proof. By the categorical equivalence (2.2), we identify $\mathcal{P}_{\vec{01}}^{\mathrm{DR}} \times \mathbf{Q}_p$, $\mathcal{A}_{\vec{01}}^{\mathrm{DR}} \otimes \mathbf{Q}_p$ and $\mathcal{V}_{\mathrm{KZ}} \otimes \mathbf{Q}_p$ with $\mathcal{P}_{\vec{01}}^{\dagger}$, $\mathcal{A}_{\vec{01}}^{\dagger}$ and $\mathcal{V}_{\mathrm{KZ}}^{\dagger}$ respectively. Because $\mathcal{P}_{\vec{01}}^{\mathrm{DR}} = Spec\mathcal{A}_{\vec{01}}^{\mathrm{DR}}$, we have a natural injection $\mathcal{P}_{\vec{01}}^{\mathrm{DR}} \to (\mathcal{A}_{\vec{01}}^{\mathrm{DR}})^{\vee}$ of pro-objects. By Lemma 2.12 we get a morphism $\mathcal{P}_{\vec{01}}^{\mathrm{DR}} \to \mathcal{V}_{\mathrm{KZ}}$, whose associated morphism $\mathcal{P}_{\vec{01}}^{\dagger} \to \mathcal{V}_{\mathrm{KZ}}^{\dagger}$ commutes with Frobenius actions, i.e.

$$\begin{split} \tilde{F}_{p}^{*} \mathcal{P}_{\overrightarrow{01}}^{\dagger} & \xrightarrow{\phi = F_{p}^{-1}} \mathcal{P}_{\overrightarrow{01}}^{\dagger} \\ \downarrow & \downarrow \\ \tilde{F}_{p}^{*} \mathcal{V}_{\mathrm{KZ}}^{\dagger} & \xrightarrow{\phi = F_{p}^{-1}} \mathcal{V}_{\mathrm{KZ}}^{\dagger} \end{split}$$

commutes. By taking the fiber ω_z , we easily see that $d_{z^p} \in \pi_1^{\mathrm{DR}}(X:\overrightarrow{01},z^p)(\mathbf{Q}_p) = \omega_{z^p}(\mathcal{P}_{\overrightarrow{01}}^{\dagger}) = \omega_z(\tilde{F}_p^*\mathcal{P}_{\overrightarrow{01}}^{\dagger})$ corresponds to $1_{(z^p)} \in \omega_{z^p}(\mathcal{V}_{\mathrm{KZ}}^{\dagger}) = \omega_z(\tilde{F}_p^*\mathcal{V}_{\mathrm{KZ}}^{\dagger})$. Hence the image $\phi_p(d_{z^p}) \in \pi_1^{\mathrm{DR}}(X:\overrightarrow{01},z)(\mathbf{Q}_p) = \omega_z(\mathcal{P}_{\overrightarrow{01}}^{\dagger})$ is mapped to $G_0^{\dagger}(z) \in \mathcal{V}_{\mathrm{KZ},(z)}^{\dagger} = \omega_z(\mathcal{V}_{\mathrm{KZ}} \otimes \mathbf{Q}_p)$. The second row induces morphism $\omega_{\overrightarrow{01}}(\mathcal{P}_{\overrightarrow{01}}^{\mathrm{DR}}) = \pi_1^{\mathrm{DR}}(X:\overrightarrow{01})(\mathbf{Q}_p) \to \omega_{\overrightarrow{01}}(\mathcal{V}_{\mathrm{KZ}}) \cong \mathbf{Q}_p \langle \langle A, B \rangle \rangle$, so that the loop $d_z^{-1}\phi_p(d_{z^p})$ corresponds to $G_0^{\dagger}(z)$.

By $d_z^{-1}\phi_p(d_{z^p}) = d_z^{-1}c_{z_0} \cdot \phi_p(d_{z^p}^{-1}c_{z_0})^{-1}$ we get the equality (2.11) whenever we fix each \mathbf{Q}_p -point z. By letting z varies for all \mathbf{Q}_p -points on $]z_0[$ and restricting G_0^{\dagger} and G_0 into $]z_0[$, we see that the equality (2.11) holds in $\mathbf{Q}_p[[t_{z_0}]]\langle\langle A, B\rangle\rangle$ (t_{z_0} : a local parameter of $]z_0[\bigcap X(\mathbf{Q}_p))$, hence for $A_{Col}|_{z_0[}$. By the uniqueness principle of Coleman functions [C], the equality (2.11) holds for the whole space. After all we get Theorem 2.14.

By Lemma 2.15 we may say that p-adic MZV à la Deligne is 'a special value ' of the overconvergent p-adic MPL. By Theorem 2.14 the overconvergent p-adic MPL is is described very explicitly as a combination of our p-adic MPL's. The following are the easiest examples.

Examples 2.16. (1)
$$Li_k^{\dagger}(z) = Li_k(z) - \frac{1}{p^k}Li_k(z^p) = \sum_{\substack{(n,p)=1\\ (n,p)=1}} \frac{z^n}{n^k}.$$

(2) $Li_{a,b}^{\dagger}(z) = Li_{a,b}(z) - \frac{1}{p^{a+b}}Li_{a,b}(z^p) - (\frac{1}{p^b} - \frac{1}{p^{a+b}})\zeta_p(a)Li_b(z^p)$
 $-\sum_{r=0}^{a-1} (-1)^r \frac{1}{p^{a-r}} {b-1+r \choose r}Li_{a-r}(z^p) \{Li_{b+r}(z) - \frac{1}{p^{b+r}}Li_{b+r}(z^p)\}$
 $-(-1)^a \sum_{s=0}^{b-1} {a-1+s \choose a-1} (\frac{1}{p^{b-s}} - \frac{1}{p^{a+b}})\zeta_p(a+s)Li_{b-s}(z^p).$

We note that the coefficient of A of $G_0(z)$ is $\log^a z$ while in the overconvergent side the coefficient of $G_0^{\dagger}(z)$ is $\log^a z - \frac{1}{p}\log^a z^p = 0$ and $Li_k^{\dagger}(z)$ has been studied by Coleman [C]. Unver nearly got the same formula (2) in [U2] 5.16. In Example 2.28, we will give a Hodge analogue of the above two formulae. Our *p*-adic MPL has log poles around 0 and ∞ however the overconvergent *p*-adic MPL does not. The formula (2.11) is an algorithm to erase log poles at ∞ of our *p*-adic MPL. We remark that the overconvergent *p*-adic MPL is rigid analytic on an open rigid analytic subspace $]\overline{X}_0[$ containing $]\mathbf{P}^1 \setminus \{1\}[$.

Remark 2.17. The *p*-adic KZ equation

$$dg = (\frac{A}{z} + \frac{B}{z-1})gdz$$

does not have a solution with overconvergent function coefficients but have a solution with Coleman function coefficients (for instance $G_0(z)$). However if we modify it as follows

(2.12)
$$dg = \left(\frac{A}{z} + \frac{B}{z-1}\right)gdz - g\left(\frac{dz^p}{z^p}\frac{A}{p} + \frac{dz^p}{z^p-1}\Phi_{\rm De}^p(A,B)^{-1}\frac{B}{p}\Phi_{\rm De}^p(A,B)\right),$$

it has a solution with overconvergent function coefficients (for instance $G_0^{\dagger}(z)$). We note that the similar differential equation for the loop $d_{z^p}^{-1}F_p(d_z)$ is introduced in [U2, Y] before.

2.2. Hodge side — analogous results. In this side, we use a Hodge-type comparison isomorphism (Lemma 2.19) between a de Rham fundamental torsor and a Betti fundamental torsor for X. A tannakian interpretation of MPL's and MZV's are given in Proposition 2.20 and Proposition 2.21. By using the infinity Frobenius action on a Betti path space we introduce a new version of MPL's (2.20). Its single-valuedness is shown in Theorem 2.27. A formula to express them in terms of usual (multi-valued) MPL's is shown in Proposition 2.26.

Notation 2.18. Let z be a point in $X(\mathbf{C})$. We fix a topological path $b_z \in \pi_1^{\text{top}}(X(\mathbf{C}) : \overrightarrow{01}, z)$. By abuse of notations we also denote b_z to be the corresponding Betti path in $\pi_1(X(\mathbf{C}) : \overrightarrow{01}, z)(\mathbf{Q})$. For $z = \overrightarrow{10}$ we consider a special path $b \in \pi_1(X(\mathbf{C}) : \overrightarrow{01}, \overrightarrow{10})(\mathbf{Q})$ which comes from a one-point set $\pi_1^{\text{top}}(X(\mathbf{R}) : \overrightarrow{01}, \overrightarrow{10})$.

The following Hodge type comparison isomorphism for fundamental torsors is our basics in this subsection.

Lemma 2.19. Let z be a C-valued point of X. Then we have

(2.13)
$$\pi_1^{\mathrm{DR}}(X:0\hat{1},z) \times \mathbf{C} \cong \pi_1^{\mathrm{Be}}(X:0\hat{1},z) \times \mathbf{C}$$

This follows from an equivalence of tannakian categories

(2.14)
$$\mathcal{NC}^{\mathrm{DR}}(X_{\mathbf{C}}) \cong \mathcal{NL}^{\mathrm{Be}}(X(\mathbf{C}))$$

(see also [De1]). By identifying the above two fundamental torsors by (2.13) we get a de Rham loop $d_z^{-1}b_z$, which lies in $\pi_1^{\text{DR}}(X:\overline{01})(\mathbf{C})$.

Proposition 2.20. Let z be a **C**-valued point of X. The de Rham loop $d_z^{-1}b_z$ corresponds to the special value at z of the analytic continuation along b_z of the fundamental solution G_0^{-13} of the KZ equation by the embedding i:

$$i(d_z^{-1}b_z) = G_0(z).$$

Proof. The proof is given in a similar way to the proof of Theorem 2.3. By [De1] §12.15-12.16 we have a tannakian interpretation of horizontal sections of $\mathcal{P}_{\overline{01}}^{\mathrm{DR}}$, By Lemma 2.12 we have a tannakian interpretation of horizontal sections of $(\mathcal{V}_{\mathrm{KZ}}, \nabla_{\mathrm{KZ}})$. Because both the constant term of $G_0(t)$ at t = 0 and the constant term of t_0^A at $t_0 = 0$ are equal to 1, we have

$$b_{\overrightarrow{0}1,z}(t_0^A|_{\overrightarrow{0}1})$$
 = the analytic continuation of $G_0(t)$ along b_z

which implies

(2.15)
$$d_{z,\Gamma} \cdot b_{\overrightarrow{01},z} \cdot d_{\Gamma,\overrightarrow{01}}(1) = G_0(z).$$

By (2.4), (2.15) and $d_{z,\Gamma} = d_{\overrightarrow{01},\Gamma} \cdot d_z^{-1}$, we get Proposition 2.20.

In each coefficient of $G_0(z)$ there appears MPL (0.1). In precise the coefficient of $A^{k_m-1}B \cdots A^{k_1-1}B$ in $G_0(z)$ is $(-1)^m Li_{k_1,\cdots,k_m}(z)$:

$$G_0(z) = 1 + \log zA - Li_1(z)B + \dots + (-1)^m Li_{k_1,\dots,k_m}(z)A^{k_m-1}B \cdots A^{k_1-1}B + \dots$$

 $^{^{13}}$ It means the fundamental solution of the KZ equation satisfying $G_0(z)\approx z^A$ constructed by Drinfel'd [Dr].

By Proposition 2.20 we may say that the loop $d_z^{-1}b_z$ is a tannakian origin of MPL's. We remark that a choice of the Betti path b_z corresponds to a choice of each branch of analytic continuation of MPL.

For $z = \overline{10}$ similarly we have

Proposition 2.21. The de Rham loop $d^{-1}b \in \pi_1^{DR}(X:\overrightarrow{01})(\mathbf{C})$ corresponds to the Drinfel'd associator $\Phi_{KZ} \in \mathbf{C}\langle\langle A, B \rangle\rangle$ by the embedding *i*, *i.e.*

$$i(d^{-1}b) = \Phi_{\mathrm{KZ}}.$$

Proof. The Drinfel'd associator Φ_{KZ} is the series constructed in [De1], which is also equal to the limit value

$$\lim_{\epsilon \to 0} \exp\left(-\log \epsilon \cdot B\right) \cdot G_0(1-\epsilon) \in \mathbf{C}\langle\langle A, B \rangle\rangle$$

as (2.6). In other word it is the constant term of $G_0(z)$ at z = 1. Then by a similar argument to the proof of Theorem 2.5 we have $b_{\vec{0}\vec{1},\vec{10}}(t_0^A|_{\vec{0}\vec{1}}) = \Phi_{\text{KZ}} \cdot t_1^B|_{\vec{10}}$, whence

(2.16)
$$d_{\overrightarrow{10},\Gamma} \cdot b \cdot d_{\Gamma,\overrightarrow{01}}(1) = \Phi_{\mathrm{KZ}}.$$

By (2.4), (2.16) and $d_{\overrightarrow{10},\Gamma} = d_{\overrightarrow{01},\Gamma} \cdot d^{-1}$, we prove Proposition 2.21.

In each coefficient of the Drinfel'd associator Φ_{KZ} there appears MZV (0.2). In precise the coefficient of $A^{k_m-1}B \cdots A^{k_1-1}B$ in Φ_{KZ} is $(-1)^m \zeta(k_1, \cdots, k_m)$:

$$\Phi_{\rm KZ} = 1 + \dots + (-1)^m \zeta(k_1, \dots, k_m) A^{k_m - 1} B \dots A^{k_1 - 1} B + \dots$$

By Proposition 2.21 we may say that the loop $d^{-1}b$ is a tannakian origin of MZV's. The above proposition enables us to calculate the period map

(2.17)
$$p: \pi_1^{\operatorname{Be}}(X(\mathbf{C}): \overrightarrow{01})(\mathbf{C}) \to \pi_1^{\operatorname{DR}}(X: \overrightarrow{01})(\mathbf{C}).$$

In order to do that, as in §2.1 we fix generator of $\pi_1^{\text{Be}}(X(\mathbf{C}): \vec{01})(\mathbf{Q})$ and its parameterization.

Notation 2.22. We denote the loop in $\pi_1^{\text{Be}}(X(\mathbf{C}) : \overrightarrow{01})(\mathbf{Q})$ (resp. in $\pi_1^{\text{Be}}(X(\mathbf{C}) : \overrightarrow{10})(\mathbf{Q})$) which comes from a generator in $\pi_1^{\text{top}}(T_0^{\times}(\mathbf{C}) : \overrightarrow{01})$ (resp. in $\pi_1^{\text{top}}(T_1^{\times}(\mathbf{C}) : \overrightarrow{10})$) going around the point $0 \in T_0(\mathbf{C})$ (resp. $0 \in T_1(\mathbf{C})$) counterclockwisely by x_{Be} (resp. y'_{Be}). Put $y_{\text{Be}} = b^{-1}y'_{\text{Be}}b$ (for b see Notation 2.18). Easily we see that two loops x_{Be} and y_{Be} are free generators of the pro-algebraic group $\pi_1^{\text{Be}}(X(\mathbf{C}) : \overrightarrow{01})(\mathbf{Q})$. Wojtkowiak [W] consider the embedding

(2.18)
$$j: \pi_1^{\operatorname{Be}}(X(\mathbf{C}): \overrightarrow{01})(\mathbf{Q}) \hookrightarrow \mathbf{Q}\langle\langle A, B \rangle\rangle$$

which sends $x_{\text{Be}} \mapsto e^A := 1 + \frac{A}{1!} + \frac{A^2}{2!} + \cdots$ and $y_{\text{Be}} \mapsto e^B := 1 + \frac{B}{1!} + \frac{B^2}{2!} + \cdots$. We note that this morphism is isomorphic to the embedding of $\pi_1^{\text{Be}}(X(\mathbf{C}):\vec{01})(\mathbf{Q})$ into the universal embedding algebra of $Lie\pi_1^{\text{Be}}(X(\mathbf{C}):\vec{01})(\mathbf{Q})$.

We note that this embedding j is not compatible with i in (2.1) under the Hodge comparison isomorphism (2.13). As is similar to the proof of Theorem 2.8, we extend the period map (2.17) to

$$p: \mathbf{C}\langle\langle A, B\rangle\rangle \to \mathbf{C}\langle\langle A, B\rangle\rangle$$

via the embedding i (2.1) and j (2.18).

Lemma 2.23. $p(A) = 2\pi i A$, $p(B) = \Phi_{KZ}^{-1}(2\pi i B) \Phi_{KZ}$.

Proof. By the compatibility of the period map on the tangential morphisms $\mathbf{Q}_{\mathrm{Be}}(1) \stackrel{\mathrm{exp}}{=} \pi_1^{\mathrm{Be}}(T_0^{\times}(\mathbf{C}):\overrightarrow{01}) \rightarrow \pi_1^{\mathrm{Be}}(X(\mathbf{C}):\overrightarrow{01}) \text{ and } \mathbf{Q}_{\mathrm{DR}}(1) \stackrel{\mathrm{exp}}{=} \pi_1^{\mathrm{DR}}(T_0^{\times}:\overrightarrow{01}) \rightarrow \pi_1^{\mathrm{DR}}(X:\overrightarrow{01}),$ we have $p(x_{\mathrm{Be}}) = x^{2\pi i}$. Similarly we have $p(y_{\mathrm{Be}}) = y'^{2\pi i}$. Since the period map is compatible with the torsor structure we have $p(y_{\mathrm{Be}}) = p(b^{-1}y'_{\mathrm{Be}}b) = b^{-1}y'^{2\pi i}b = (d^{-1}b)^{-1}y^{2\pi i}d^{-1}b$. By Proposition 2.21, we get the claim.

Our next work is on a description of a Frobenius action. To begin with we consider the infinity Frobenius action ϕ_{∞} on $\pi_1^{\text{DR}}(X:\vec{01},\vec{10})(\mathbf{C})$ by transmitting the infinity Frobenius action on $\pi_1^{\text{Be}}(X(\mathbf{C}):\vec{01},\vec{10})(\mathbf{C})$ by Lemma 2.19. It gives a new path $\phi_{\infty}(d)$ in $\pi_1^{\text{DR}}(X:\vec{01},\vec{10})(\mathbf{C})$. We denote Φ_{KZ}^- to be the series in $\mathbf{C}\langle\langle A, B \rangle\rangle$ which corresponds to $d^{-1}\phi_{\infty}(d)$:

$$i(d^{-1}\phi_{\infty}(d)) = \Phi_{\mathrm{KZ}}^{-}.$$

It is a Hodge counterpart of the *p*-adic Deligne associator Φ_{De}^p . By same arguments to the proof of Lemma 2.9 and Theorem 2.8 the extension of the infinity Frobenius action on $\pi_1^{\text{DR}}(X:\vec{01})(\mathbf{C})$ into $\mathbf{C}\langle\langle A, B\rangle\rangle$ by *i* is described as follows:

Lemma 2.24.
$$\phi_{\infty}(A) = -A$$
, $\phi_{\infty}(B) = \Phi_{\text{KZ}}^{-1}(-B)\Phi_{\text{KZ}}^{-1}$.

This gives the following formula analogous to (2.8).

Lemma 2.25.

$$\Phi_{\rm KZ}(A,B) = \Phi_{\rm KZ}^{-}(A,B) \cdot \Phi_{\rm KZ}\left(-A, \Phi_{\rm KZ}^{-}(A,B)^{-1}(-B)\Phi_{\rm KZ}^{-}(A,B)\right).$$

This formula looks nasty. But according to the convention in [F0] of the Grothendieck-Teichmüller group [Dr] we rewrite simpler

$$(-1, \Phi_{\mathrm{KZ}}^{-}) = (-1, \Phi_{\mathrm{KZ}}) \circ (1, \Phi_{\mathrm{KZ}})^{-1}.$$

Let z be a **C**-valued point of X. By a same argument to above, we get a de Rham loop $d_z^{-1}\phi_{\infty}(d_{\bar{z}}) \in \pi_1^{\text{DR}}(X:\overline{01})(\mathbf{C})$. Put $G_0^-(z) := i(d_z^{-1}\phi_{\infty}(d_{\bar{z}}))$. The following is an analogue to Theorem 2.14.

Proposition 2.26.

$$(2.19) \ G_0^-(A,B)(z) = G_0(A,B)(z) \cdot G_0\left(-A, \Phi_{\mathrm{KZ}}^-(A,B)^{-1}(-B)\Phi_{\mathrm{KZ}}^-(A,B)\right)(\bar{z})^{-1}.$$

Proof. By $d_z^{-1}\phi_{\infty}(d_{\bar{z}}) = d_z^{-1}b_z \cdot \phi_{\infty}(d_{\bar{z}}^{-1}\overline{b_z})^{-1}$ and Lemma 2.24 we get the equality.

We introduce a new version of MPL $Li_{k_1,\dots,k_m}^-(z)$ to be the coefficient of $A^{k_m-1}B$ $\cdots A^{k_1-1}B$ in $G_0^-(z)$ multiplied by $(-1)^m$:

(2.20)
$$G^{-}(z) =: 1 + \dots + (-1)^{m} Li^{-}_{k_{1},\dots,k_{m}}(z) A^{k_{m}-1} B \dots A^{k_{1}-1} B + \dots$$

We note that since $G^{-}(z)$ is group-like we can recover and express general coefficients of $G^{-}(z)$ in terms of $Li^{-}_{k_{1},\cdots,k_{m}}(z)$ and $\log |z|$ as Note 2.6.

Theorem 2.27. The coefficient $Li_{k_1,\dots,k_m}^-(z)$ is single-valued and real-analytic on $X(\mathbf{C}) = \mathbf{P}^1(\mathbf{C}) \setminus \{0, 1, \infty\}.$

Proof. The real-analyticity follows from (2.19) because both $G_0(z)$ and $G_0(\bar{z})$ are real-analytic. The single-valuedness is immediate by definition.

The single-valued MPL's are not equal to usual MPL's because they have a different tannakian origins. But by Proposition 2.26 the single-valued MPL is described very explicitly as a combination of usual MPL's. The following are the easiest examples:

Examples 2.28. (1)
$$Li_{k}^{-}(z) = Li_{k}(z) - \sum_{a=0}^{k-1} (-1)^{k-a} \frac{(\log|z|^{2})^{a}}{a!} Li_{k-a}(\bar{z})$$

(2) $Li_{a,b}^{-}(z) = Li_{a,b}(z) - \sum_{r=0}^{a-1} \sum_{s=0}^{r} [(-1)^{a+r+s} {b-1+s \choose s} \frac{(\log|z|^{2})^{r-s}}{(r-s)!} Li_{a-r}(\bar{z}) \cdot {Li_{b+s}(z) - \sum_{w=0}^{b+s-1} (-1)^{b+s+w} \frac{(\log|z|^{2})^{w}}{w!} Li_{b+s-w}(\bar{z})}] - \sum_{u=0}^{b-1} \frac{(\log|z|^{2})^{u}}{u!} \cdot [(-1)^{a+b+u} Li_{a,b-u}(\bar{z}) + \{(-1)^{b+u} - (-1)^{a+b+u}\}\zeta(a)Li_{b-u}(\bar{z}) + \sum_{w=0}^{b-u-1} \{(-1)^{a+b+u+v} - (-1)^{b+u}\} {a+v-1 \choose a-1} \zeta(a+v)Li_{b-u-v}(\bar{z})].$

The coefficient of A of $G_0(z)$ is $\log z$ while the coefficient of $G_0^-(z)$ is $\log \overline{z} + \log z = \log |z|^2$. We also remark that Zhao [Zh] constructed a single-valued version of several variable MPL $Li_{k_1,\dots,k_m}(z_1,\dots,z_m) = \sum_{0 < n_1 < \dots < n_m} \frac{z_1^{n_1\dots z_m^{n_m}}}{n_1^{k_1\dots n_m^{k_m}}}.$

Remark 2.29. In [Za] Zagier studied another variant of polylogarithm

$$P_k(z) = \Re_k \sum_{a=0}^{k-1} \frac{B_a}{a!} (\log |z|^2)^a Li_{k-a}(z)$$

where \Re_k denote \Re or \Im depending whether k is odd or even and $\frac{t}{e^t-1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}$. He showed that its single-valuedness and real-analyticity. Beilinson and Deligne [BD] give its interpretations in terms of variants of mixed Hodge structures. Their computations [BD] §1.5 says that $P_k(z)$ appears as the coefficient of $A^{k-1}B$ of $\log G_0^-(z)$ multiplied by $-\frac{1}{2}$, whence the tannakian origin of $P_k(z)$ is also $d_z^{-1}\phi_{\infty}(d_{\bar{z}})$. We notice that $P_k(z)$ and $Li_k^-(z)$ are related with each other by

(2.21)
$$P_k(z) = \frac{1}{2} \sum_{i=0}^{k-1} \frac{B_i}{i!} \{ \log |z|^2 \}^i Li_{k-i}^-(z)$$

and

$$Li_k^{-}(z) = 2\sum_{i=0}^{k-1} \frac{(\log |z|^2)^i}{(i+1)!} P_{k-i}(z).$$

Our MPL has monodromies around 0, 1 and ∞ however the variant does not. The formula (2.19) gives an algorithm to erase monodromies of our MPL at 0, 1 and ∞ .

Remark 2.30. The KZ equation

$$dg = (\frac{A}{z} + \frac{B}{z-1})gdz$$

have a solution with multi-valued complex analytic function coefficients (for instance $G_0(z)$) whereas the following modification analogous to (2.12)

$$dg = \left(\frac{A}{z} + \frac{B}{z-1}\right)gdz - g\left(\frac{d\bar{z}}{\bar{z}}(-A) + \frac{d\bar{z}}{\bar{z}-1}\Phi_{\mathrm{KZ}}^-(A,B)^{-1}(-B)\Phi_{\mathrm{KZ}}^-(A,B)\right)$$

does have a solution with single-valued real-analytic function coefficients (for instance $G_0^-(z)$).

2.3. Artin side — reviews. In this side we will discuss *l*-adic étale analogues of MPL's and MZV's which we call *l*-adic MPL's and *l*-adic multiple Soulé elements respectively and whose definitions are imitations of Wojtkowiak's *l*-adic polylogarithms [W].

Let l be a prime (we do not assume whether l is equal to the previous prime p in §2.1 or not). Let $\overline{\mathbf{Q}}$ be the algebraic closure of \mathbf{Q} in \mathbf{C} . The following Artin-type comparison isomorphism take place of Lemma 2.2 and Lemma 2.19.

Lemma 2.31. Suppose that x and y are $\overline{\mathbf{Q}}$ -valued points of X or T_s^{\times} ($s \in \{0, 1, \infty\}$) then there exists an isomorphism

$$\pi_1^{\text{Be}}(X(\mathbf{C}):x,y) \times_{\mathbf{Q}} \mathbf{Q}_l \cong \pi_1^{l,\acute{e}t}(X_{\overline{\mathbf{Q}}}:x,y).$$

It follows from the comparison isomorphism [G] XII Corollary 5.2 (see also [De1] §13.11).

Let $i : \pi_1^{l,\text{\acute{e}t}}(X_{\overline{\mathbf{Q}_l}}: \overrightarrow{01})(\mathbf{Q}_l) \hookrightarrow \mathbf{Q}_l \langle \langle A, B \rangle \rangle$ be the embedding associated with j(2.18). Suppose that $z \in X(\mathbf{Q})$ and $b_z \in \pi_1^{\text{top}}(X(\mathbf{C}): \overrightarrow{01}, z)$. It determines a Betti path in $\pi_1^{\text{Be}}(X(\mathbf{C}): \overrightarrow{01}, z)(\mathbf{Q})$ and we regard this to be an *l*-adic étale path (which we denote b_z by abuse of notation) in $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{\mathbf{Q}}}: \overrightarrow{01}, z)(\mathbf{Q}_l)$ Lemma 2.31. Suppose that $\sigma \in Gal(\overline{\mathbf{Q}}/\mathbf{Q})$. We obtain another path $\sigma(b_z)$ by the Galois group action on $\pi_1^{l,\text{\acute{e}t}}(X_{\overline{\mathbf{Q}}}: \overrightarrow{01}, z)(\mathbf{Q}_l)$. Combining $\sigma(b_z)$ with the inverse b_z^{-1} , we get an *l*-adic étale loop $b_z^{-1}\sigma(b_z) \in \pi_1^{l,\text{\acute{e}t}}(X_{\overline{\mathbf{Q}}}: \overrightarrow{01})(\mathbf{Q}_l)$. Wojtkowiak [W] introduced *l*-adic polylogarithm $\ell_k^z(\sigma)$ $(k \ge 1)$ to be the coefficient of $A^{k-1}B$ of $\log j(b_z^{-1}\sigma(b_z))$ multiplied by $(-1)^{k-1}$ (cf. [NW]). In accordance with expressions in §2.1 and §2.2, we consider the following *l*-adic étale analogues of MPL's and MZV's.

Definition 2.32. Let $m, k_1, \dots, k_m \ge 1, z \in X(\mathbf{Q})$ and $\sigma \in Gal(\overline{\mathbf{Q}}/\mathbf{Q})$. The *l*-adic multiple polylogarithm (shortly the *l*-adic MPL) $Li_{k_1,\dots,k_m}^l(z)(\sigma)$ denotes the coefficient of $A^{k_m-1}B \cdots A^{k_1-1}B$ in $j(b_z^{-1}\sigma(b_z))$ multiplied by $(-1)^m$ and *l*-adic multiple Soulé element (shortly the *l*-adic MSE) $\zeta_l(k_1,\dots,k_m)(\sigma)$ $(k_m > 1)$ denotes the coefficient of $A^{k_m-1}B \cdots A^{k_1-1}B$ in the *l*-adic Ihara associator $\Phi_{\sigma}^l(A,B) = j(b^{-1}\sigma(b))$ multiplied by $(-1)^m$.

By making a same computations similar to [BD] §1.5, we have

(2.22)
$$\ell_k^z(\sigma) = (-1)^k \sum_{i=0}^{k-1} \frac{B_i}{i!} \{\rho_z(\sigma)\}^i Li_{k-i}^l(z)(\sigma).$$

Here $\rho_z(\sigma)$ is the generalized *l*-adic Kummer 1-cocycle in [NW] Definition 3. By comparing it with (2.21), we may say that $Li_k^l(z)(\sigma)$ is a *l*-adic analogue of the usual (Leibnitz's) polylogarithm $Li_k(z) = \sum_{n=1}^{\infty} \frac{z^n}{n!}$ and Wojtkowiak's $\ell_k^z(\sigma)$ is a *l*adic analogue of the Beilinson-Deligne's polylogarithm $D_k(z)$. By asking for *p*adic Hodge theory, we may give new *p*-adic MPL's valued on the ring of B_{crys} of Fontaine's *p*-adic periods, which might unify *p*-adic MPL's in §2.1 and those above (for l = p).

The following may explain why we call the multiple Soulé elements.

Examples 2.33 ([I] §6.3.Theorem). (1) When k is odd, $\zeta_l(k)(\sigma) = \frac{\kappa_k^{(l)}(\sigma)}{(1-l^{k-1})(k-1)!}$

- Here $\kappa_k^{(l)}(\sigma)$ is *l*-adic Soulé element (see [I] §6.2 for its definition). (2) When k is even, $\zeta_l(k)(\sigma) = \frac{B_k}{2(k!)} \{1 \chi_l(\sigma)^k\}$. Here B_k is the Bernoulli number defined by $\frac{t}{e^t-1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}$ and $\chi_l(\sigma)$ is the *l*-adic cyclotomic character.

The author expects that MSE is a kind of 'multiple Euler system' which helps our understanding of multiple zeta functions.

We end this section by giving a brief observation on *l*-adic behavior of $\zeta_l(k)(\sigma)$ and p-adic behavior of $\zeta_p(k)$ with respect to weights.

Remark 2.34. Let $\sigma \in Gal(\overline{\mathbf{Q}}/\mathbf{Q})$. By the expression $\kappa_k^l(\sigma) = \int_{\mathbf{Z}_k^{\times}} x^{k-1} d\mu^{\sigma}$ (μ^{σ}) : the measure associated with the Kummer distribution [NW] §2) we see that $(k-1)!(1-l^{k-1})\zeta_l(k)(\sigma)$ admits a nice *l*-adic behavior with respect to k, that is, for $k \equiv k' \mod (l-1)l^M$ $(M \in \mathbf{N})$ we have

(2.23)
$$(k-1)!(1-l^{k-1})\zeta_l(k)(\sigma) \equiv (k'-1)!(1-l^{k'-1})\zeta_l(k')(\sigma) \mod l^{M+1}\mathbf{Z}_l.$$

On the other hand in the *p*-adic setting, by $L_p(1-k,\omega^{1-k}) = -(1-p^{k-1})\frac{B_k}{k} = (1-p^{k-1})\zeta(1-k)$ and $L_p(k,\omega^{1-k}) = (1-\frac{1}{p^k})\zeta_p(k)$ we see that $(1-\frac{1}{p^k})\zeta_p(k)$ admits a nice p-adic behavior with respect to k, that is, for $k \equiv k' \mod (p-1)p^M$ $(M \in \mathbb{N})$ we have

(2.24)
$$(1 - \frac{1}{p^k})\zeta_p(k) \equiv (1 - \frac{1}{p^{k'}})\zeta_p(k') \mod p^{M+1}\mathbf{Z}_p.$$

The author is not sure whether we have 'multiple analogues' of (2.23) and (2.24). He is grateful to the referee who suggested him that Coleman's integral formula ([C] Lemma 7.2) $Li_k^{\dagger}(z) = \int_{\mathbf{Z}_p^{\times}} x^{-k} d\mu_z(x)$ for $z \in \mathbf{C}_p \setminus \{|z-1|_p < 1\}$ might help look for such relations. Here μ_z is a measure on \mathbf{Z}_p given by $\mu_z(a+p^n\mathbf{Z}_p)=\frac{z^a}{1-z^{p^n}}$ with $n \in \mathbf{N}$ and $0 \leq a < p^n$.

3. MOTIVIC VIEWS

This section is complementary. We will consider three boxes related to the algebra generated by p-adic MZV's. In §3.1 our special box is Drinfel'd's [Dr] proalgebraic bi-torsor where the Grothendieck-Teichmüller pro-algebraic groups act. In §3.2 our special box is Racinet's [R] pro-algebraic torsor defined by double shuffle relations. In §3.3 our special box is pro-algebraic bi-torsor of Deligne-Goncharov's [DG] motivic Galois group. The relationship between these three box will be discussed.

3.1. Drinfel'd's context. In this subsection we will add the Berthelot-Ogus part into the story of [F0] owing Unver's result [U1] and discuss special elements in Ihara's stable derivation algebra which are *p*-adic analogues of the Drinfel'd's element ψ in [Dr] Proposition 6.3.

In his celebrated paper [Dr], Drinfel'd introduced the Grothendieck-Teichmüller torsor, a triple $(\underline{GRT}_1, \underline{M}_1, \underline{GT}_1)$, where \underline{M}_1 is the pro-algebraic bi-torsor with a left (resp. right) action of the Grothendieck-Teichmüller pro-algebraic group GRT_1 (resp. GT_1) (for their presentations see also [F0]§2). In [F0]§4 we explained that

for a prime *l* the *l*-adic Galois image pro-algebraic group $\underline{Gal}_{\mathbf{Q}_{l}}^{(l)}$ (which is defined in loc.cit. to be the Zariski closure of the image of the Galois group $Gal(\overline{\mathbf{Q}}/\mathbf{Q}(\mu_{l^{\infty}}))$ -action in $\underline{Aut}\pi_{1}^{l,\text{ét}}(X_{\overline{\mathbf{O}}}:\overrightarrow{01})(\mathbf{Q}_{l}))$ lies in $\underline{GT}_{1} \times \mathbf{Q}_{l}$, i.e. we have

(3.1)
$$\Phi_{\mathbf{Q}_{l}}^{(l)} : \underline{Gal}_{\mathbf{Q}_{l}}^{(l)} \hookrightarrow \underline{GT}_{1} \times \mathbf{Q}_{l},$$

whereas in [F0] §3 we also saw that the spectrum of the **Q**-algebra $\mathfrak{Z}(\subset \mathbf{C})$ generated by $(2\pi i)^{-(k_1+\cdots+k_m)}\zeta(k_1,\cdots,k_m)$ $(m,k_i \in \mathbf{N}, k_m > 1)$ is embedded into \underline{M}_1 , i.e. we have

We note that (3.1) and (3.2) are expected to be isomorphism [F0]. In [F1] we announced to make up still lacking Berthelot-Ogus part in our next upcoming paper '*p*-adic multiple zeta values III'; we planned to construct an embedding

(3.3)
$$\Phi_{\text{crys}}^{(p)} : Spec\mathfrak{Z}^{(p)} \hookrightarrow \underline{GRT}_1$$

where $\mathfrak{Z}^{(p)}$ is a subalgebra of \mathbf{Q}_p generated by all *p*-adic MZV's. But we would cancel it and give an explanation below instead because a recent Unver's paper [U1] nearly achieved it.

Proposition 3.1. The *p*-adic Drinfel'd associator $\Phi_{KZ}^p(A, B)$ satisfies the following defining equations of <u>*GRT*</u>₁:

Here $\mathbb{L}^{\wedge}_{\mathbf{Q}_p}$ stands for the completed free Lie algebra generated by A and B and $\underline{U\mathfrak{P}}^{(5)}_{\cdot}$ stands for the universal enveloping algebra of the pure sphere 5-braid Lie algebra $\mathfrak{P}^{(5)}$ with standard generators $X_{i,j}$ $(1 \leq i, j \leq 5)$.

Proof. By $(p, \Phi_{De}^p) = (p, 1) \circ (1, \Phi_{De}^p(pA, pB))$ and $(p, \Phi_{KZ}^p) = (p, 1) \circ (1, \Phi_{KZ}^p(pA, pB))$ in our convention of <u>*GRT*</u>₁, (2.9) is rewritten into

(3.4)
$$(1, \Phi_{\text{De}}^p(pA, pB)) = (1, \Phi_{\text{KZ}}^p(pA, pB)) \circ (1, \Phi_{\text{KZ}}^p(A, B))^{-1}$$

In [U1] Unver showed that the *p*-adic Deligne associator $\Phi_{\text{De}}^p(A, B)$ satisfies the above relations, in other words $(1, \Phi_{\text{De}}^p(A, B)) \in \underline{GRT}_1(\mathbf{Q}_p)$. Thus $(1, \Phi_{\text{De}}^p(p^n A, p^n B)) \in \underline{GRT}_1(\mathbf{Q}_p)$ for $n \in \mathbf{N}$. Put

$$(1, a_n) = (1, \Phi_{\mathrm{De}}^p(pA, pB))^{-1} \circ (1, \Phi_{\mathrm{De}}^p(p^2A, p^2B))^{-1} \circ \dots \circ (1, \Phi_{\mathrm{De}}^p(p^nA, p^nB))^{-1}$$

Because <u>*GRT*</u>₁ is closed under the multiplication, $(1, a_n)$ must belongs to <u>*GRT*</u>₁(\mathbf{Q}_p). By (3.4),

$$(1, \Phi_{\mathrm{De}}^p(p^n A, p^n B)) = (1, \Phi_{\mathrm{KZ}}^p(p^n A, p^n B)) \circ (1, \Phi_{\mathrm{KZ}}^p(p^{n-1} A, p^{n-1} B))^{-1}.$$

So we have $(1, a_n) = (1, \Phi_{KZ}^p(A, B)) \circ (1, \Phi_{KZ}^p(p^n A, p^n B))^{-1}$. As $n \operatorname{goes} \infty, \Phi_{De}^p(p^n A, p^n B)$ converges to 1 in $\mathbf{Q}_p \langle \langle A, B \rangle \rangle$ which is a projective limit of finite dimensional topological \mathbf{Q}_p -vector space. Hence the series a_n $(n \ge 1)$ should converge to $\Phi_{KZ}^p(A, B)$ in $\mathbf{Q}_p \langle \langle A, B \rangle \rangle$. By the locally-compactness of the topological group $\underline{GRT}_1(\mathbf{Q}_p)$ the limit $(1, \Phi_{KZ}^p(A, B))$ must lie on $\underline{GRT}_1(\mathbf{Q}_p)$, whence we get the claim. \Box

By Proposition 3.1 we get a morphism (3.3). The injectivity is easy to check.

Since there appear *p*-adic MZV's in each coefficient of the *p*-adic Drinfel'd associator Φ_{KZ}^p , we get algebraic relations among them. It is remarkable that we get $\zeta_p(2k) = 0 \ (k \ge 1)$ from (0), (i) and (ii) (proved in the same way to [De1]§18.16), that is, we have a geometric proof for $L_p(2k, \omega^{1-2k}) = 0$.

In [F0] §6 by discussing a weight filtration on our torsor, we associated (3.1) and (3.2) with two surjections between their coordinate rings, $Gr\Phi_{\mathbf{Q}_l}^{(l)\sharp} : \mathcal{O}(\underline{GRT}_1) \cdot \widehat{\otimes} \mathbf{Q}_l \twoheadrightarrow Gr_{\mathbf{Q}_l}^W \mathcal{O}(\underline{Gal}_{\mathbf{Q}_l}^{(l)})$ and $Gr\Phi_{\mathrm{Hod}}^{\sharp} : \mathcal{O}(\underline{GRT}_1) \cdot \twoheadrightarrow (Z_{\mathbf{Q}_l}^{-}/(\pi^2)Z_{\mathbf{Q}_l})$ where Z is a graded **Q**-algebra whose degree w-component $(w \ge 1)$ is a **Q**-vector space $Z_w (\subset \mathbf{R})$ generated by all MZV's with weight w and $Z_0 = \mathbf{Q}$. By (3.3), we also obtain

$$\Phi_{\mathrm{crys}}^{(p)\sharp}: \mathcal{O}(\underline{GRT}_1)_{\bullet} \twoheadrightarrow Z_{\bullet}^{(p)}$$

where $Z_{\star}^{(p)}$ is a graded **Q**-algebra whose degree *w*-component is a **Q**-vector space $Z_{w}^{(p)}(\subset \mathbf{Q}_{p})$ generated by all *p*-adic MZV's with weight *w* and $Z_{0}^{(p)} = \mathbf{Q}$.

Remark 3.2. (1) In [F0]§6.6.1, we introduced an invertible element Φ_{GRT} in $\mathcal{O}(\underline{GRT}_1)\langle\langle A, B \rangle\rangle$ and showed $Gr\Phi_{\text{Hod}}^{\sharp}(\Phi_{GRT}) = \Phi_{\text{KZ}} \mod \pi^2$ and $Gr\Phi_{\mathbf{Q}_l}^{(l)\sharp}(\Phi_{GRT}) = Gr\Phi_{\text{Ih}}^{(l)}$ (a series associated with the *l*-adic Ihara associator Φ_{σ}^l in loc.cit). By (3.3) we get their Berthelot-Ogus counterpart

$$\Phi_{\rm crys}^{(p)\sharp}(\Phi_{GRT}) = \Phi_{\rm KZ}^p$$

(2) In loc.cit §6.6.2, we consider the meta-abelian quotient $B_{GRT} \in \mathcal{O}(\underline{GRT}_1)[[A, B]]$ of Φ_{GRT} and showed $Gr\Phi_{\text{Hod}}^{\sharp}(B_{GRT}) = \frac{\Gamma(1-A)\Gamma(1-B)}{\Gamma(1-A-B)} \mod \pi^2$ and $Gr\Phi_{\mathbf{Q}_l}^{(l)\sharp}(B_{GRT})$ $= GrB^{(l)}(A, B)$ where $\Gamma(1-z) = \exp\{\gamma z + \sum_{n=2}^{\infty} \zeta(n) \frac{z^n}{n}\}$ (γ : Euler constant) is the classical gamma function and $GrB^{(l)}(A, B)$ is the associated graded

quotient of the Ihara's universal power series for Jacobi sums [I]. By (3.3) we get their Berthelot-Ogus counterpart

$$\Phi_{\rm crys}^{(p)\sharp}(B_{GRT}) = \prod_{k=1}^{\infty} \frac{\Gamma_p(1+p^k A)\Gamma_p(1+p^k B)}{\Gamma_p(1+p^k (A+B))}$$

where $\Gamma_p(1-z) = \exp\{\gamma_p z + \sum_{n=2}^{\infty} L_p(n, \omega^{1-n}) \frac{z^n}{n}\}$ (γ_p : a *p*-adic analogue of Euler constant) is the Morita's *p*-adic gamma function ([M] Theorem 1).

Euler constant) is the Morita's p-aute gamma function ([M] Theorem 1).

Next we discuss a special element of Drinfel'd's $\mathfrak{grt}_1 = \bigoplus_{n \in \mathbb{N}} \mathfrak{grt}_1^{n-14}$ which is the graded Lie algebra of \underline{GRT}_1 [Dr] §5. The original Drinfel'd associator $\Phi_{\mathrm{KZ}}(A, B)$ satisfies the defining equations of \underline{M} [Dr] which is similar to but different from those of \underline{GRT}_1 [Dr] however we say the following.

Lemma 3.3. The '- ' part $\Phi_{KZ}^{-}(A, B)$ of the Drinfel'd associator satisfies the same equations to those of Proposition 3.1.

Proof. Drinfel'd showed $(1, \Phi_{\mathrm{KZ}}(\frac{A}{2\pi i}, \frac{B}{2\pi i})) \in \underline{M}_1(\mathbf{C})$ in [Dr]. By the complex conjugate action we get $(1, \Phi_{\mathrm{KZ}}(\frac{-A}{2\pi i}, \frac{-B}{2\pi i})) \in \underline{M}_1(\mathbf{C})$. By the torsor structure of $\underline{M}_1(\mathbf{C})$, there is a unique element $(1, \varphi) \in \underline{GRT}_1(\mathbf{C})$ such that $(1, \varphi) \circ$

¹⁴Actually it is isomorphic to the graded completion of Ihara's [I] stable derivation algebra \mathfrak{D} .

$$(1, \Phi_{\mathrm{KZ}}(\frac{-A}{2\pi i}, \frac{-B}{2\pi i})) = (1, \Phi_{\mathrm{KZ}}(\frac{A}{2\pi i}, \frac{B}{2\pi i})), \text{ equivalently}$$
$$\Phi_{\mathrm{KZ}}\left(\frac{A}{2\pi i}, \frac{B}{2\pi i}\right) = \varphi(A, B) \cdot \Phi_{\mathrm{KZ}}\left(\frac{-A}{2\pi i}, \varphi(A, B)^{-1}(\frac{-B}{2\pi i})\varphi(A, B)\right).$$

By comparing the equations in Lemma 2.25, we get $\varphi = \Phi_{\text{KZ}}^-(\frac{A}{2\pi i}, \frac{B}{2\pi i})$. So $(1, \varphi) = (1, \Phi_{\text{KZ}}^-(\frac{A}{2\pi i}, \frac{B}{2\pi i})) \in \underline{GRT}_1$, from which we get the claim.

Remark 3.4. By Lemma 3.3, $(1, \Phi_{KZ}^-) \in \underline{GRT}_1(\mathbf{R})$. By the logarithmic morphism $Log: \underline{GRT}_1(\mathbf{R}) \to \mathfrak{grt}_1(\mathbf{R})$, Drinfel'd obtained in [Dr] Proposition 6.3 a canonical element $\mathfrak{f}(\infty)$ in $\mathfrak{grt}_1(\mathbf{R})$, the image of $(1, \Phi_{KZ}^-)$, with the following presentations:

(3.5)
$$f(\infty) = \sum_{m \ge 3: \text{ odd}} f(\infty)_m \text{ where } f(\infty)_m = 2\zeta(m)(adA)^{m-1}(B) + \dots \in \mathfrak{grt}_1^m(\mathbf{R}).$$

On the other hand, $(1, \Phi_{\text{KZ}}^p) \in \underline{GRT}_1(\mathbf{Q}_p)$ by Proposition 3.2. By the logarithmic morphism $Log : \underline{GRT}_1(\mathbf{Q}_p) \to \mathfrak{grt}_1(\mathbf{Q}_p)$, we also get a canonical element \mathfrak{f}_p in $\mathfrak{grt}_1(\mathbf{Q}_p)$, the image of $(1, \Phi_{\text{KZ}}^p)$, with the following presentation:

$$\mathfrak{f}(p) = \sum_{m \ge 3} \mathfrak{f}(p)_m \quad \text{where} \quad \mathfrak{f}(p)_m = \zeta_p(m)(adA)^{m-1}(B) + \dots \in \mathfrak{grt}_1^m(\mathbf{Q}_p).$$

(N.B. $\mathfrak{f}(\infty)_m = 0$ for m = 1 or even.) We stress that the summation in $\mathfrak{f}(\infty)$ is taken for odd number m greater than or equal to 3 while the summation in $\mathfrak{f}(p)$ is taken for natural number m greater than or equal to 3.

3.2. Racinet's context. We will recall Racinet's [R] pro-algebraic torsor \underline{DMR}_1 which is defined by two shuffle relations and discuss a story for \underline{DMR}_1 analogous to previous section.

MZV $\zeta(k_1, \dots, k_m)$ $(m, k_1, \dots, k_m \in \mathbf{N}, k_m > 1)$ satisfies two types of so called shuffle product formula, expressing a product of two MZV's as a linear combination of other such values. The first type, known as series shuffle product formula and for which the easiest example is the relation

$$\zeta(k_1) \cdot \zeta(k_2) = \zeta(k_1, k_2) + \zeta(k_2, k_1) + \zeta(k_1 + k_2) ,$$

is easily obtained from the expression (0.2). The second type of shuffle product formula, known as integral shuffle product formula, come from their iterated integral expressions. The easiest example is the formula

$$\zeta(k_1) \cdot \zeta(k_2) = \sum_{i=0}^{k_1-1} \binom{k_2-1+i}{i} \zeta(k_1-i,k_2+i) + \sum_{j=0}^{k_2-1} \binom{k_1-1+j}{j} \zeta(k_2-j,k_1+j) + \sum_{j=0}^{k_2-1} \binom{k_2-1+j}{j} \zeta(k_2-j,k_2+j) + \sum_{j=0}^{k_2-1+j} \binom{k_2-1+j}{j} \zeta(k_2-j,k_2+j) + \sum_{j=0}^{k_2-1} \binom{k_2-1+j}{j} \zeta(k_2-j,k_2+j) - \sum_{$$

The double shuffle relations are linear relations combining series shuffle product formula and integral shuffle product formula. The above two formulae give the simplest example

$$\begin{aligned} \zeta(k_1,k_2) &+ \zeta(k_2,k_1) + \zeta(k_1+k_2) \\ &= \sum_{i=0}^{k_1-1} \binom{k_2-1+i}{i} \zeta(k_1-i,k_2+i) + \sum_{j=0}^{k_2-1} \binom{k_1-1+j}{j} \zeta(k_2-j,k_1+j) \end{aligned}$$

We have two extended notions of MZV's to non-admissible indices, the case when the last component k_m is equal to 1, which we call series regularization and integral regularization here. The usual and series regularized MZV's satisfy series shuffle product formulae and the usual and integral regularized MZV's satisfy integral shuffle product formulae. These two kinds of regularized MZV's are related by so called regularized relations.

Racinet constructed a pro-algebraic variety ¹⁵ <u>DMR</u> whose set <u>DMR(k)</u> of kvalued points (k: a field of characteristic 0) consists of a series $\varphi(A, B) \in k \langle \langle A, B \rangle \rangle$ such that $\varphi(A, -B)$ satisfies (3.2.1.1) and (3.2.1.2) in [R], which are relations reflecting the above relations. Let $\lambda \in k$. He denoted <u> $DMR_{\lambda}(k)$ </u> to be the subset consisting of the series in $\underline{DMR}(k)$ whose coefficient of AB is equal to $\frac{\lambda^2}{24}$. He showed in [R] Theorem 1 that $\underline{DMR}_1(k)$ (resp. $\underline{DMR}_0(k)$) has a structure of the pro-algebraic torsor (resp. pro-algebraic group) where $\underline{DMR}_{0}(k)$ acts from the left (resp. whose group law is defined by the same way to the Grothendieck-Teichmüller group \underline{GRT}_1).

We embed $\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})(k)$ into $k\langle\langle A,B\rangle\rangle$ via i (2.1) and $\pi_1^{\mathrm{Be}}(X(\mathbf{C}):\overrightarrow{01})(k)$ into $k\langle\langle A,B\rangle\rangle$ via j (2.18). The group automorphism of $\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})(k)$ sending $A \mapsto A$ and $B \mapsto \varphi^{-1} B \varphi$ induces an embedding of both \underline{DMR}_0 and \underline{GRT}_1 into $\underline{Aut}\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\vec{01})$ as pro-algebraic groups. And the group isomorphism from $\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\vec{01})(k)$ to $\pi_1^{\mathrm{Be}}(X(\mathbf{C}):\vec{01})(k)$ sending $A \mapsto A$ and $B \mapsto \varphi^{-1}B\varphi$ induces an embedding of both \underline{DMR}_1 and \underline{M}_1 into $\underline{Isom}(\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01}), \pi_1^{\mathrm{Be}}(X(\mathbf{C}):\overrightarrow{01}))$ as pro-algebraic torsors. From now on we regard \underline{DMR}_0 and \underline{GRT}_1 to be sub-proalgebraic groups of $\underline{Aut}\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})$ and regard \underline{DMR}_1 and \underline{M}_1 to be sub-proalgebraic torsors of $\underline{Isom}(\pi_1^{\text{DR}}(X_{\mathbf{Q}}:\overrightarrow{01}), \pi_1^{\text{Be}}(X(\mathbf{C}):\overrightarrow{01})).$ We note that $\Phi_{\text{KZ}}\left(\frac{A}{2\pi i}, \frac{B}{2\pi i}\right)$ belongs to $\underline{DMR}_1(\mathbf{C})$, which gives a morphism

$$(3.6) \qquad \qquad \Psi_{\text{Hod}} : Spec\mathfrak{Z} \hookrightarrow \underline{DMR}_1$$

analogous to (3.2) and conjectured to be isomorphic [R]. The paper [BF] is a trial to give a morphism

(3.7)
$$\Psi_{\text{crys}}^{(p)} : Spec\mathfrak{Z}^{(p)} \hookrightarrow \underline{DMR}_0,$$

which is an analogous map to (3.3). We showed double shuffle relations, i.e. series and integral shuffle product formulae, for usual p-adic MZV's but not for extended values. In [FJ] we will give a complete proof to give an embedding $\Psi_{\rm crvs}^{(p)}$, namely show series and integral shuffle product formulae and regularized relations for regularized *p*-adic MZV's.

3.3. Deligne-Goncharov's context. We will discuss the pro-algebraic bi-torsor of the motivic Galois group in [DG] by recalling the category of mixed Tate motives over \mathbf{Z} . This torsor is related with the Drinfel'd's torsor (§3.1) in Proposition 3.8 and the Racinet's torsor $(\S3.2)$ in Proposition 3.9. We also explain how the motivic formalism fits into our story in Note 3.10: In Hodge side we give a motivic interpretation of Zagier's dimension conjecture on MZV's. We recall how the upperbounding part of this conjecture follows as a consequence. In Artin side we see how Ihara's conjecture [I2] on Galois image is motivically related with Zagier's conjecture via Problem 3.7. In Berthelot-Ogus side we deduce another motivic way, different from [FJ], of proving double (series and integral) shuffle relations and regularized relations for (extended) p-adic multiple zeta values by using Yamashita's [Y] fiber functor of the crystalline realization.

¹⁵For our convenience slightly we change the definition.

Let k be a field with characteristic 0. Levine [L2] and Voevodsky [Voe] constructed a triangulated category of mixed motives over k. Levine [L2] showed an equivalence of these two categories. This category denoted by $DM(k)_{\mathbf{Q}}$ has Tate objects $\mathbf{Q}(n)$ $(n \in \mathbf{Z})$. Let $DMT(k)_{\mathbf{Q}}$ be the triangulated sub-category of $DM(k)_{\mathbf{Q}}$ generated by $\mathbf{Q}(n)$ $(n \in \mathbf{Z})$. Levine [L1] extracted a neutral tannakian \mathbf{Q} -category $MT(k)_{\mathbf{Q}}$ of mixed Tate motives over k from $DMT(k)_{\mathbf{Q}}$ by taking a heart with respect to a t-structure under the Beilinson-Soulé vanishing conjecture which says $gr_i^{\gamma}K_n(k) = 0$ for n > 2i. Here LHS is the graded quotient of the algebraic K-theory for k with respect to γ -filtration.

From now on we assume that k is a number field. In this case the Beilinson-Soulé vanishing conjecture holds and we have $MT(k)_{\mathbf{Q}}$. This category satisfies the following expected properties: Each object M has an increasing filtration of subobjects called weight filtration, $W : \cdots \subseteq W_{m-1}M \subseteq W_mM \subseteq W_{m+1}M \subseteq \cdots$, whose intersection is 0 and union is M. The quotient $Gr_{2m+1}^WM := W_{2m+1}M/W_{2m}M$ is trivial and $Gr_{2m}^WM := W_{2m}M/W_{2m+1}M$ is a direct sum of finite copies of $\mathbf{Q}(m)$ for each $m \in \mathbf{Z}$. Morphisms of $MT(k)_{\mathbf{Q}}$ are strictly compatible with weight filtration. The extension group is related to K-theory as follows

$$Ext^{i}_{MT(k)\mathbf{Q}}(\mathbf{Q}(0),\mathbf{Q}(m)) = \begin{cases} K_{2m-i}(k)\mathbf{Q} & \text{for } i=1, \\ 0 & \text{for } i>1. \end{cases}$$

There are realization fiber functors ([L2] and [H]) corresponding to usual cohomology theories.

Let S be a finite set of finite places of k. Let \mathcal{O}_S be the ring of S-integers in k. Deligne and Goncharov [DG] defined the full subcategory $MT(\mathcal{O}_S)$ of mixed Tate motives over \mathcal{O}_S , whose objects are mixed Tate motives M in $MT(k)_{\mathbf{Q}}$ such that for each subquotient E of M which is an extension of $\mathbf{Q}(n)$ by $\mathbf{Q}(n+1)$ for $n \in \mathbf{Z}$, the extension class of E in $Ext^1_{MT(k)_{\mathbf{Q}}}(\mathbf{Q}(n), \mathbf{Q}(n+1)) = Ext^1_{MT(k)_{\mathbf{Q}}}(\mathbf{Q}(0), \mathbf{Q}(1)) = k^{\times}_{\mathbf{Q}}$ lies in $\mathcal{O}_S^{\times} \otimes \mathbf{Q}$. In this category the following hold:

$$Ext^{1}_{MT(\mathcal{O}_{S})}(\mathbf{Q}(0),\mathbf{Q}(m)) = \begin{cases} 0 & \text{for } m < 1, \\ \mathcal{O}_{S}^{\times} \otimes \mathbf{Q} & \text{for } m = 1, \\ K_{2m-1}(k)_{\mathbf{Q}} & \text{for } m > 1, \end{cases}$$
$$Ext^{2}_{MT(\mathcal{O}_{S})}(\mathbf{Q}(0),\mathbf{Q}(m)) = 0.$$

Let $\omega_{\text{can}} : MT(\mathcal{O}_S) \to Vect_{\mathbf{Q}}$ ($Vect_{\mathbf{Q}}$: the category of \mathbf{Q} -vector spaces) be the fiber functor which sends each motive M to $\bigoplus_n Hom(\mathbf{Q}(n), Gr_{-2n}^W M)$. Let G_{can} be the motivic Galois group $\underline{Aut}^{\otimes}(MT(\mathbf{Z}) : \omega_{\text{can}})$. The action of G_{can} on $\omega_{\text{can}}(\mathbf{Q}(1)) = \mathbf{Q}$ defines a surjection $G_{\text{can}} \to \mathbf{G}_m$ and its kernel U_{can} is the unipotent radical of G_{can} . There is a canonical splitting $\tau : \mathbf{G}_m \to G_{\text{can}}$ which gives a negative grading on the Lie algebra $LieU_{\text{can}}$ (consult [De1] §8 for the full story). The above computations of Ext-groups follows

Proposition 3.5. The graded Lie algebra $LieU_{can}$ is free and its degree n-part of $LieU_{can}^{ab} = U_{can}^{ab}$ is isomorphic to the dual of $Ext_{MT(\mathcal{O}_S)}^1(\mathbf{Q}(0), \mathbf{Q}(-n))$.

Proof . See [De1] §8 and [DG] §2.

Let us restrict in the case of $k = \mathbf{Q}$, $S = \emptyset$, $\mathcal{O}_S = \mathbf{Z}$. Let $\omega_* : M \mapsto M_*$ (* = Be, DR) be the fiber functor which associates each mixed Tate motive $M \in MT(\mathbf{Z})$ with the underlying vector space of its Betti, De Rham realization respectively.

For $*, *' = \{Be, DR\}$ we denote the corresponding tannakian fundamental group <u>Aut</u>^{\otimes}($\mathcal{MT}(\mathbf{Z})$: ω_*) by G_* and the corresponding tannakian fundamental torsor <u>Isom</u>^{\otimes}($\mathcal{MT}(\mathbf{Z})$: $\omega_*, \omega_{*'}$) by $G_{**'}$. Note that the latter is a $(G_*, G_{*'})$ -bi-torsor and $G_{**} = G_*$. Let U_* be the sub-pro-algebraic group of G_* whose action on $\omega_*(\mathbf{Q}(1)) = \mathbf{Q}$ is trivial and $U_{**'}$ be the sub-pro-algebraic torsor of $G_{**'}$ which induces a trivial map from $\omega_*(\mathbf{Q}(1)) = \mathbf{Q}$ to $\omega_{*'}(\mathbf{Q}(1)) = \mathbf{Q}$. Then $U_{**'}$ is a $(U_*, U_{*'})$ -bi-torsor and $U_{**} = U_*$. Since $M_{\rm DR} = \omega_{\rm can}(M)$ we have $G_{\rm can} = G_{\rm DR}$. By Proposition 3.5 the Lie algebra $LieU_{DR}$ of the unipotent part U_{DR} of G_{DR} should be a graded free Lie algebra generated by one element in each degree -m ($m \ge 3$: odd).

Remark 3.6. In [De1] §8.12, Deligne constructed a free basis of the Lie algebra $LieU_{\rm DR}(\mathbf{C})$ as follows: The infinity Frobenius action ϕ_{∞} on $\omega_{\rm Be}(M)$ for $M \in$ $\mathcal{MT}(\mathbf{Z})$ determines a point in $G_{\text{Be}}(\mathbf{Q})$ and by the Hodge comparison isomorphism (the period map) $\omega_{\text{Be}}(M) \otimes \mathbf{C} \simeq \omega_{\text{DR}}(M) \otimes \mathbf{C}$ it gives a point in $G_{\text{DR}}(\mathbf{C})$ which we denote by the same symbol ϕ_{∞} . Put $\psi_{\infty} = \phi_{\infty} \circ \tau(-1)$. Then $\psi_{\infty} \in U_{\text{DR}}(\mathbf{C})$ because $\phi_{\infty} \in G_{\mathrm{DR}}(\mathbf{C})$ goes to $-1 \in \mathbf{G}_m$. The element $\mathfrak{e}_m(\infty)$ $(m \ge 3:$ odd) generates freely the Lie algebra $LieU_{\rm DR}({f C})$ where $\mathfrak{e}_m(\infty)$ means a degree -mpart of the image $\mathfrak{e}(\infty) = \sum_m \mathfrak{e}_m(\infty)$ of ψ_∞ by the logarithmic morphism Log: $U_{\rm DR}(\mathbf{C}) \rightarrow LieU_{\rm DR}(\mathbf{C}).$ (N.B. $\mathfrak{e}_m(\infty) = 0$ for m = 1 or even.)

In [DG] §4 they constructed the motivic fundamental group $\pi_1^{\mathcal{M}}(X:\overline{01})$ with $X = \mathbf{P}^1 \setminus \{0, 1, \infty\}$, which is an ind-object of $MT(\mathbf{Z})$. This is an affine group $MT(\mathbf{Z})$ -scheme (cf. §1.1), whose de Rham realization and Betti realization agree with the de Rham fundamental group $\pi_1^{\text{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})$ in §1.1 and the Betti fundamental group $\pi_1^{\text{Be}}(X(\mathbf{C}):\vec{01})$ in §1.3 respectively. Namely $\omega_{\text{DR}}(\pi_1^{\mathcal{M}}(X:\vec{01})) =$ $\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})$ and $\omega_{\mathrm{Be}}(\pi_1^{\mathcal{M}}(X:\overrightarrow{01})) = \pi_1^{\mathrm{Be}}(X(\mathbf{C}):\overrightarrow{01})$. Since all the structure morphism of $\pi_1^{\mathcal{M}}(X:01)$ belong to the set of morphisms of $MT(\mathbf{Z})$ we have

(3.8)
$$\varphi: G_{**'} \to \underline{Isom}(\omega_*(\pi_1^{\mathcal{M}}(X:01)), \omega_{*'}(\pi_1^{\mathcal{M}}(X:01)))$$

for $*, *' \in \{Be, DR\}$. On this map φ the following is one of the basic problems.

Problem 3.7. Is φ injective?

This might be said a question which asks a validity of a unipotent variant of the so-called 'Belyi's theorem' in [Bel] in the pro-finite setting. Equivalently this asks if the motivic fundamental group $\pi_1^{\mathcal{M}}(X:\overline{01})$ is a generator of the tannakian category $MT(\mathbf{Z})$.

The $(U_{\rm DR}, U_{\rm Be})$ -bi-torsor $U_{\rm DR, Be}$ is related to the $(\underline{GRT}_1, \underline{GT}_1)$ -bi-torsor \underline{M}_1 in $\S3.1$ as follows.

Proposition 3.8. $\varphi(U_{\rm DR}) \subset \underline{GRT}_1, \ \varphi(U_{\rm DR,Be}) \subset \underline{M}_1, \ \varphi(U_{\rm Be}) \subset \underline{GT}_1.$

Proof . By Lemma 2.24 the automorphism which corresponds to $\phi_{\infty}\tau(-1)$ is described by $(1, \Phi_{KZ}^{-})$. So the induced homomorphism $\Phi : LieU_{DR} \to DerLie\pi_{1}^{DR}(X_{Q})$ $\overrightarrow{01}$) sends $\mathfrak{e}(\infty)$ to $\mathfrak{f}(\infty)$ (cf. Remark 3.4). Since $\mathfrak{f}(\infty)_m$ is the image of the free generator $\mathfrak{e}(\infty)_m$, the image $\Phi(LieU_{\rm DR})$ must lies in $\mathfrak{grt}_1(=Lie\underline{GRT}_1) \subset$ $DerLie\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})$. So $\varphi(U_{\mathrm{DR}})$ should lie in $\underline{GRT}_1 \subset \underline{Aut}\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})$. The morphism $\tau(2\pi i)^{-1}p$ lies in $U_{\mathrm{DR,Be}}(\mathbf{C})$. Its image $\varphi(\tau(2\pi i)^{-1}p)$, which

is a morphism from $\pi_1^{\mathrm{DR}}(X_{\mathbf{Q}}:\overrightarrow{01})(\mathbf{C})$ to $\pi_1^{\mathrm{Be}}(X(\mathbf{C}):\overrightarrow{01})(\mathbf{C})$, is described by

 $(1, \Phi_{\mathrm{KZ}}(\frac{A}{2\pi i}, \frac{B}{2\pi i}))$ by Lemma 2.23. By [Dr] this element belongs to $\underline{M}_1(\mathbf{C})$, which implies $\varphi(U_{\mathrm{DR,Be}}) \subset \underline{M}_1$. It is because φ is a morphism from the U_{DR} -torsor $U_{\mathrm{DR,Be}}$ to the \underline{GRT}_1 -torsor \underline{M}_1 , which sends $\tau(2\pi i)^{-1}p \in U_{\mathrm{DR,Be}}(\mathbf{C})$ to $(1, \Phi_{\mathrm{KZ}}(\frac{A}{2\pi i}, \frac{B}{2\pi i})) \in \underline{M}_1(\mathbf{C})$.

The $(U_{\text{DR}}, U_{\text{Be}})$ -bi-torsor $U_{\text{DR,Be}}$ is also related to the <u>DMR</u>₀-torsor <u>DMR</u>₁ in §3.2 as follows.

Proposition 3.9. $\varphi(U_{\text{DR}}) \subset \underline{DMR}_0, \ \varphi(U_{\text{DR},\text{Be}}) \subset \underline{DMR}_1.$

Proof. The proof is similar to the previous proposition. As for the proof for $\varphi(U_{DR}) \subset \underline{DMR}_0$, we use Racinet's result [R] 5.3.2, which is the same argument to [Dr] Proposition 6.3, that the Lie algebra \mathfrak{dmr}_0 of his \underline{DMR}_0 contains the Drinfel'd's element $\mathfrak{f}(\infty)_m$ (3.5).

At present we do not know the relationship between Drinfel'd's <u> GRT_1 </u> and Racinet's <u> DMR_0 </u> although it might be expected that they are equal. However the recent Terasoma's work (to appear) might suggests a direction <u> $DMR_0 \subset GRT_1$ </u>.

Below we explain how the motivic formalism fits in the scheme of our subject in the previous section.

Note 3.10. (1) In Hodge side: as is shown in the proof of Proposition 3.8 the pair $(1, \Phi_{KZ}(\frac{A}{2\pi i}, \frac{B}{2\pi i}))$ is the image of $\tau(2\pi i)^{-1}p$ by φ , that is, $(1, \Phi_{KZ}(\frac{A}{2\pi i}, \frac{B}{2\pi i})) \in \varphi(U_{DR,Be})(\mathbf{C})$. Since $(2\pi i)^{-(k_1+\cdots+k_m)}\zeta(k_1, \ldots, k_m)$ $(m, k_i \in \mathbf{N}, k_m > 1)$ appears among each coefficient of $\Phi_{KZ}(\frac{A}{2\pi i}, \frac{B}{2\pi i})$, it gives rise a morphism

(3.9)
$$\Gamma_{\text{Hod}} : Spec\mathcal{Z} \hookrightarrow \varphi(U_{\text{DR,Be}})$$

analogous to (3.2) and (3.6). There is a conjecture on the dimension of the vector space of MZV's at each weight which is called Zagier conjecture (see [Za] and also [F0]) and partly proved (for upper-bounding part) by Terasoma [T]. By using the embedding (3.9) and Proposition 3.5, Deligne and Goncharov also get a partial (upper-bounding part) proof of Zagier conjecture in [DG]. We also get to know that to say Zagier conjecture holds is equivalent to say the surjectivity of Γ_{Hod} and the injectivity of φ (i.e. the validity of Problem 3.7).

(2) In Artin side: we consider the absolute Galois group $Gal(\overline{\mathbf{Q}}/\mathbf{Q})$ action on the *l*-adic (*l*: a prime) étale fundamental group $\varphi_1 : Gal(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \underline{Aut}\pi_1^{l,\text{ét}}(X_{\overline{\mathbf{Q}}}: \overrightarrow{01})$ and the *l*-adic Galois image pro-algebraic group $\underline{Gal}_{\mathbf{Q}_l}^{(l)}$ defined to be the Zariski closure of $\varphi_1(Gal(\overline{\mathbf{Q}}/\mathbf{Q}(\mu_l^\infty)))$ in [F0] §4. The category $MT(\mathbf{Z})$ has a fiber functor $\omega_{l,\text{ét}} : M \mapsto M_{l,\text{ét}}$ of the *l*-adic étale realization which associates each motives M with its underlying \mathbf{Q}_l -vector space of its *l*-adic étale realization. The absolute Galois group $Gal(\overline{\mathbf{Q}}/\mathbf{Q})$ acts functorially on $\omega_{l,\text{ét}}(M)$ and hence it induces a morphism $Gal(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow G_{l,\text{ét}}(\mathbf{Q}_l)$. Hereafter we fix an embedding $\overline{\mathbf{Q}} \hookrightarrow \mathbf{C}$. Then there is the Artin comparison isomorphism $\omega_{l,\text{ét}}(M) \simeq \omega_{\text{Be}}(M) \otimes \mathbf{Q}_l$ functorial with respect to M, which gives $G_{l,\text{ét}} = G_{\text{Be}} \times \mathbf{Q}_l$. We might have a morphism $\varphi_2 : Gal(\overline{\mathbf{Q}}/\mathbf{Q}) \to G_{\text{Be}}(\mathbf{Q}_l)$. Because of $\pi_1^{l,\text{ét}}(X_{\overline{\mathbf{Q}}}: \overline{01}) = \omega_{l,\text{ét}}(\pi_1^{\mathcal{M}}(X:\overline{01}))$ $= \omega_{\text{Be}}(\pi_1^{\mathcal{M}}(X:\overline{01})) \times \mathbf{Q}_l$, the map φ_1 might factor through G_{Be} action on $\omega_{Be}(\pi_1^{\mathcal{M}}(X:\overline{01}))$, that means $\varphi_1 = \varphi \circ \varphi_2$ (consult [De1] §8 for full story). In loc.cit. §8.14 it is also shown that the image of φ_2 is open in the topology of $G_{\text{Be}}(\mathbf{Q}_l)$. So by restricting into unipotent parts we get an isomorphism

(3.10)
$$\Gamma_{l,\text{\acute{e}t}} : \underline{Gal}_{\mathbf{Q}_{l}}^{(l)} \xrightarrow{=} \varphi(U_{\text{Be}}) \times \mathbf{Q}_{l}$$

which might be said an analogue of (3.1). This is the way how the common **Q**-structure of the *l*-adic Galois image pro-algebraic group for all prime *l* is detected in [DG] Remark 6.13. The graded Lie algebra $\mathfrak{g}_{.}^{(l)}$ (cf.[I2] and also [F0]) associated with <u>Gal_{Q_l}^{(l)}</u> is conjectured by Ihara (loc.cit.) to be a free Lie algebra generated by one element in each degree 3,5,7,9,... As in Hodge case we can say that this Ihara's conjecture is equivalent to the injectivity of φ (i.e. the validity of Problem 3.7).

(3) In Berthelot-Ogus side: Yamashita will construct in [Y] the fiber functor $\omega_{p,\mathrm{crys}}$: $M \to M_{p,\mathrm{crys}}$ of the crystalline realization of $MT(\mathbf{Z})$ which is compatible with $\omega_{\rm DR}$. It associates each mixed Tate motive M with its underlying \mathbf{Q}_l -vector space of its crystalline realization by using Fontaine's functor to give a conjectured dimension bounding for $Z_w^{(p)}$. It admits a functorial crystalline Frobenius action $\phi_{p,crys}$ on $\omega_{p,crys}(M)$ and a functorial comparison isomorphism $\omega_{p,crys}(M) = \omega_{DR}(M) \otimes \mathbf{Q}_p$ for $MT(\mathbf{Z})$, which gives a point in $G_{\rm DR}(\mathbf{Q}_p)$ denoted by the same symbol $\phi_{p,{\rm crys}}$. Since $\phi_{p,\mathrm{crys}}$ goes to $\frac{1}{p} \in \mathbf{G}_m$, the element $\psi_p = \phi_{p,\mathrm{crys}} \circ \tau(p)$ must lie in $U_{\rm DR}(\mathbf{Q}_p)$. As is similar to Remark 3.4 and Remark 3.6 we also get in the p-adic setting elements $e(p)_m$ $(m \ge 3)$ which is on a degree -mpart of $LieU_{\rm DR}(\mathbf{Q}_p)$. The action of ψ_p on $\pi_1^{\rm DR}(X_{\mathbf{Q}}:\vec{01})$ is described as $A \mapsto A$ and $B \mapsto \Phi_{\rm De}^p(A, B)^{-1}B\Phi_{\rm De}^p(A, B)$ by Lemma 2.9. Therefore $(1, \Phi_{\mathrm{De}}^{p}) \in \varphi(U_{\mathrm{DR}}(\mathbf{Q}_{p}))$. By Proposition 3.9, $\Phi_{\mathrm{De}}^{p} \in \underline{DMR}_{0}(\mathbf{Q}_{p})$. Since Deligne's *p*-adic MZV's $\zeta_{p}^{\mathrm{De}}(k_{1}, \cdots, k_{m})$ are coefficients of $\Phi_{\mathrm{De}}^{p}(A, B)$, they must satisfy double shuffle relations. By the same argument to Proposition 3.1, we get $(1, \Phi_{KZ}^p) \in \varphi(U_{DR}(\mathbf{Q}_p))$. It follows that we also get an embedding

(3.11)
$$\Gamma_{\text{crvs}}^{(p)} : Spec\mathfrak{Z}^{(p)} \hookrightarrow \varphi(U_{\text{DR}}).$$

Combining (3.11) with Proposition 3.9, we get the embedding (3.7). Therefore our *p*-adic MZV's $\zeta_p(k_1, \dots, k_m)$ must satisfy (series and integral) double shuffle relations and regularization relations, which are defining relations of <u>*DMR*</u>₀. This is a motivic proof of double shuffle relations for *p*-adic MZV's, which was a project posed in [De2].

References

- [A] Abe, E.; Hopf Algebras, Cambridge Univ. Press, 1980.
- [BD] Beïlinson, A. and Deligne, P.; Interprétation motivique de la conjecture de Zagier reliant polylogarithmes et regulateurs, Motives (Seattle, WA, 1991), 97–121, Proc. Sympos. Pure Math., 55, Part 2, Amer. Math. Soc., Providence, RI, 1994.
- [Bel] Belyĭ, G. V., Galois extensions of a maximal cyclotomic field, (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 43 (1979), no. 2, 267–276, 479.
- [Ber] Berthelot, P.; Cohomologie rigide et cohomologie rigide à support propre, Première partie, Prépublication IRMAR 96-03, 89 pages (1996).
- [Bes] Besser, A.; Coleman integration using the Tannakian formalism, Mathematische Annalen 322 (2002) 1, 19-48.
- [BF] _____and Furusho, H.; The double shuffle relations for p-adic multiple zeta values, math.NT/0310177, to appear in AMS Contemporary Math.

- [C] Coleman, R.; Dilogarithms, regulators and p-adic L-functions, Invent. Math. 69 (1982), no. 2, 171–208.
- [CLS] Chiarellotto, B. and Le Stum, B.; F-isocristaux unipotents, Compositio Math. 116 (1999),no. 1, 81–110.
- [De1] Deligne, P.; Le groupe fondamental de la droite projective moins trois points, Galois groups over Q (Berkeley, CA, 1987), 79–297, Math. S. Res. Inst. Publ., 16, Springer, New York-Berlin, 1989.
- [De2] _____Periods for the fundamental group, a short note on Arizona Winter School 2002, available from http://swc.math.arizona.edu~swcenter/notes/files/02DelignePD.dvi.
- [DG] _____and Goncharov, A.; Groupes fondamentaux motiviques de Tate mixte, Ann. Sci. Ecole Norm. Sup. (4) 38 (2005), no. 1, 1-56.
- [DM] _____and Milne, J.; Tannakian categories, in Hodge cycles, motives, and Shimura varieties (P.Deligne, J.Milne, A.Ogus, K.-Y.Shih editors), Lecture Notes in Mathematics 900, Springer-Verlag, 1982.
- [Dr] Drinfel'd, V. G.; On quasitriangular quasi-Hopf algebras and a group closely connected with $\operatorname{Gal}(\overline{Q}/Q)$, Leningrad Math. J. 2 (1991), no. 4, 829–860.
- [F0] Furusho, H.; Multiple zeta values and Grothendieck-Teichmüller groups, to appear in AMS Contemporary Math.
- [F1] _____p-adic multiple zeta values I p-adic multiple polylogarithms and the p-adic KZ equation, Inventiones Mathematicae, Vol. 155, No.2, 253-286(2004).
- [FJ] _____and Jafari, A.; Double shuffle relations for regularized p-adic multiple zeta values, available at math.NT/0510681, preprint in 2005.
- [G] Grothendieck, A.: Revêtement Étales et Groupe Fondamental (SGA 1), Lecture Notes in Math. 224, Springer-Verlag 1971.
- [H] Huber, H.; Realization of Voevodsky's motives, J.Alg.Geom. 9. 2000, 755-799.
- Ihara, Y.; Braids, Galois groups, and some arithmetic functions, Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990), 99–120, Math. Soc. Japan, Tokyo, 1991.
- [I2] _____; Some arithmetic aspects of Galois actions on the pro-*p* fundamental group of $\mathbb{P}^1 \{0, 1, \infty\}$, Arithmetic fundamental groups and noncommutative algebra (Berkeley, CA, 1999), 247–273, Proc. Sympos. Pure Math., 70, Amer. Math. Soc., Providence, RI, 2002.
- [L1] Levine, M.; Tate motives and the vanishing conjectures for algebraic K-theory, in: Algebraic K-theory and algebraic topology (Lake Louis, 1991), 167-188, NATO Adv. Sci, but. Ser.C Math. Phys. 407, Kluwer, 1993.
- [L2] _____; Mixed Motives, Math surveys and Monographs 57, AMS, 1998.
- [M] Morita, Y.; A p-adic integral representation of the p-adic L-function. J. Reine Angew. Math. 302 (1978), 71–95.
- [NW] Nakamura, H. and Wojtkowiak, Z.; On explicit formulae for *l*-adic polylogarithms, Arithmetic fundamental groups and noncommutative algebra (Berkeley, CA, 1999), 285–294, Proc. Sympos. Pure Math., 70, Amer. Math. Soc., Providence, RI, 2002.
- [R] Racinet, G.; Doubles melanges des polylogarithmes multiples aux racines de l'unite, Publ. Math. Inst. Hautes Etudes Sci. No. 95 (2002), 185–231.
- [S1] Shiho, A.; Crystalline fundamental groups. I. Isocrystals on log crystalline site and log convergent site, J. Math. S. Univ. Tokyo 7(2000), no. 4, 509–656.
- [S2] ____; Crystalline fundamental groups. II. Log convergent cohomology and rigid cohomology, J. Math. S. Univ. Tokyo 9 (2002), no. 1, 1–163.
- [T] Terasoma, T., Mixed Tate motives and multiple zeta values, Invent. Math. 149 (2002), no. 2, 339–369.
- [U1] Unver, S.; Drinfel'd-Ihara Relations for the Crystalline Frobenius, math.AG/0310386, preprint on 2003.
- [U2] _____; p-adic multi-zeta values, J. Number Theory 108 (2004), no. 1, 111–156.
- [Voe] Voevodsky, V.; Triangulated categories of motives over a field, in: Cycles, transfer and motivic homology theories, 188-238, Ann of Math Studies 143, Princeton University Press, 1993.
- [Vol] Vologodsky, V.; Hodge structure on the fundamental group and its application to p-adic integration, Mosc. Math. J. 3 (2003), no. 1, 205–247.

- [W] Wojtkowiak, W.; On l-adic polylogarithms, Prépublication no.549, Universite de Nice-Sophia Antipolis, Juin 1999.
- [Y] Yamashita, G.; Bounds for the dimensions of the *p*-adic multiple *L*-value spaces, in preparation.
- [Za] Zagier, D.; Polylogarithms, Dedekind zeta functions and the algebraic K-theory of fields. Arithmetic algebraic geometry (Texel, 1989), 391–430, Progr. Math., 89, Birkhauser Boston, Boston, MA, 1991
- [Zh] Zhao, J.; Multiple polylogarithms: analytic continuation, monodromy, and variations of mixed Hodge structures. Contemporary trends in algebraic geometry and algebraic topology (Tianjin, 2000), 167–193, Nankai Tracts Math., 5, World Sci. Publishing, River Edge, NJ, 2002.

Graduate School of Mathematics, Nagoya University, Chikusa-ku, Furo-cho, Nagoya, 464-8602, Japan

 $E\text{-}mail\ address:\ \texttt{furusho@math.nagoya-u.ac.jp}$