

Linear relations among holomorphic quadratic differentials and induced Siegel's metric on \mathcal{M}_g

Marco Matone and Roberto Volpato

Dipartimento di Fisica "G. Galilei" and Istituto Nazionale di Fisica Nucleare
Università di Padova, Via Marzolo, 8 – 35131 Padova, Italy

We derive the $(g-2)(g-3)/2$ linearly independent relations among the products of pairs in a basis of holomorphic abelian differentials in the case of compact non-hyperelliptic Riemann surfaces of genus $g \geq 4$. By the Kodaira-Spencer map this leads to the modular invariant metric on the moduli space induced by the Siegel metric.

Let Σ be a compact non-hyperelliptic Riemann surface of genus $g \geq 4$ and $\{\omega_1, \dots, \omega_g\}$ a basis of $H^0(\Sigma, K)$, with K the canonical line bundle of Σ . We denote by $\hat{\mathcal{M}}_g$ the corresponding locus in the moduli space \mathcal{M}_g of compact Riemann surfaces. Each element of $H^0(\Sigma, K^2)$ can be written as a linear combination of the $M := g(g+1)/2$ elements in

$$\mathcal{S} := \{\omega_i \omega_j \mid i \leq j = 1, \dots, g\} .$$

Since $N := H^0(\Sigma, K^2) = 3g - 3$, there are $M - N = (g-2)(g-3)/2$ independent linear relations among the quadratic differentials $\omega_i \omega_j$. In this paper we explicitly express such relations.

Let p_1, \dots, p_g and q_1, \dots, q_{2g-2} be two sets of arbitrary points on Σ . Choose a local trivialization of the canonical line bundle and set

$$a_{ij,r} := \det \omega(p_1, p_2, \dots, p_{i-1}, q_r, p_{i+1}, \dots, p_g) \det \omega(p_1, p_2, \dots, p_{j-1}, q_r, p_{j+1}, \dots, p_g) ,$$

where $\det \omega(x_1, \dots, x_g) := \det \omega_i(x_j)$. Let us define the matrix

$$A(k, l) := \begin{pmatrix} a_{12,1} & \dots & a_{1g,1} & a_{23,1} & \dots & a_{2g,1} & a_{kl,1} \\ a_{12,2} & \dots & a_{1g,2} & a_{23,2} & \dots & a_{2g,2} & a_{kl,2} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{12,2g-2} & \dots & a_{1g,2g-2} & a_{23,2g-2} & \dots & a_{2g,2g-2} & a_{kl,2g-2} \end{pmatrix} ,$$

$$3 \leq k < l \leq g, \quad g \geq 4.$$

Theorem 1.

$$\det A(k, l) = 0 ,$$

$$3 \leq k < l \leq g.$$

Let $\Delta_{i,2g-2}$ be the minor associated to $A(k, l)_{i,2g-2}$ and D_{mn} , $1 \leq m, n \leq g$, be the minor of the matrix $\omega_i(p_j)$ associated to the element in the position (m, n) . Let us denote by $A_{ij,r}(k, l)$, $1 \leq i, j \leq g$, $1 \leq r \leq 2g-2$, $3 \leq k < l \leq g$, the matrix obtained from $A(k, l)$ by replacing the column ${}^t(a_{11,r}, \dots, a_{kl,r})$ with the column ${}^t(D_{1i}D_{1j}, \dots, D_{ki}D_{lj})$.

Corollary. For each r , $1 \leq r \leq 2g-2$, the following relations

$$\sum_{i,j=1}^g \frac{\det A_{ij,r}(k, l)}{\Delta_r 2g-2} \omega_i \omega_j = 0 ,$$

$3 \leq k < l \leq g$, provide $(g-2)(g-3)/2$ linearly independent conditions on \mathcal{S} which do not depend on the points q_j , $1 \leq j \leq 2g-2$.

In particular, these relations hold for the standard basis of the holomorphic abelian differentials.

Theorem 2. Fix a local trivialization of K and let p_1, \dots, p_g be a set of points in Σ such that

$$\det \omega(p_1, \dots, p_g) \neq 0 .$$

Then

$$\sigma_i(z) := \frac{\det \omega(p_1, p_2, \dots, p_{i-1}, z, p_{i+1}, \dots, p_g)}{\det \omega(p_1, \dots, p_g)} , \quad (1)$$

$i = 1, \dots, g$, determines a basis of $H^0(\Sigma, K)$ which does not depend on the choice of the basis $\{\omega_1, \dots, \omega_g\}$ and, up to normalization, of the trivialization.

Proof. The matrix $[\omega]_{ij} := \omega_i(p_j)$ is non-singular. Then

$$\sigma_i(z) = \sum_j [\omega]_{ij}^{-1} \omega_j(z) ,$$

$i = 1, \dots, g$ is a basis of $H^0(\Sigma, K)$. □

Note that the σ_i satisfy the following properties

$$\sigma_i(p_j) = \delta_{ij} ,$$

$$\det \sigma_i(p_1, p_2, \dots, p_{j-1}, z, p_{j+1}, \dots, p_g) = \sigma_j(z) .$$

Petri basis. Let us assume that the points p_1, \dots, p_g are in “general position” and denote by η_i the holomorphic 1-differential vanishing at $p_1, \dots, \check{p}_i, \dots, p_g$. By Riemann-Roch theorem the set $\{\eta_1, \dots, \eta_g\}$ is a basis of 1-differentials. Moreover, we assume that the effective divisor $(\eta_1) + (\eta_2) - \sum_{i=3}^g p_i$ consists of $3g-3$ distinct points. The quadratic differentials

$$\eta_1^2 , \quad \eta_1 \eta_2 , \quad \eta_2^2 , \quad \eta_1 \eta_i , \quad \eta_2 \eta_i , \quad \eta_i^2 ,$$

$i = 3, \dots, g$ introduced by Petri [1], provide a basis of $H^0(\Sigma, K^2)$.

Let us introduce the set $\{v_1, \dots, v_M\} \subset H^0(\Sigma, K^2)$ determined by

$$v_k := \begin{cases} \sigma_k^2, & k = 1, \dots, g, \\ \sigma_{i+j}\sigma_j, & k = i + j(2g - j + 1)/2, \end{cases}$$

where $j = 1, \dots, g-1, i = 1, \dots, g-j$.

Proposition. $\{v_j\} := \{v_1, \dots, v_N\}$ is a basis of $H^0(\Sigma, K^2)$.

Proof. Set $D := \sum_{i=3}^g p_i$ and let us consider the effective divisors $D_i := (\sigma_i) - D, i = 1, 2$. Let us first prove that σ_i is the unique 1-differential, up to normalization, vanishing at $D_i, i = 1, 2$. Any 1-differential $\sigma'_i \in H^0(\Sigma, K)$ vanishing at D_i , corresponds to an element σ'_i/σ_i of $H^0(\Sigma, \mathcal{O}(D))$, the space of meromorphic functions f on Σ such that $(f) + D$ is an effective divisor. Suppose that there exists σ'_i such that σ'_i/σ_i is not a constant, so that $h^0(\Sigma, \mathcal{O}(D)) \geq 2$. By Riemann-Roch theorem

$$h^0(\Sigma, K \otimes \mathcal{O}(-D)) = h^0(\Sigma, \mathcal{O}(D)) - \deg D - 1 + g \geq 3,$$

there exist at least 3 linearly independent 1-differentials vanishing at D and, in particular, there exists a linear combination of such differentials vanishing at p_1, \dots, p_g . This implies that $\det \eta(p_1, \dots, p_g) = 0$ for an arbitrary basis $\{\eta\}$ of $H^0(\Sigma, K)$, contradicting the hypotheses. Fix $\zeta_i, \zeta_{1i}, \zeta_{2i} \in \mathbb{C}$ such that

$$\sum_{i=3}^g \zeta_i \sigma_i^2 + \sum_{i=1}^g \zeta_{2i} \sigma_1 \sigma_i + \sum_{i=2}^g \zeta_{1i} \sigma_2 \sigma_i = 0.$$

Evaluating this relation at the point $p_j, 3 \leq j \leq g$ yields

$$\zeta_j = 0.$$

Set

$$t_1 := - \sum_{j=2}^g \zeta_{1j} \sigma_j, \quad t_2 := \sum_{j=1}^g \zeta_{2j} \sigma_j, \quad (2)$$

so that

$$\sigma_1 t_2 = \sigma_2 t_1.$$

Since D, D_1 and D_2 consist of pairwise distinct points, t_i vanishes at $D_i, i = 1, 2$ and then, by the former remarks, $t_1/\sigma_1 = t_2/\sigma_2 = \zeta \in \mathbb{C}$. By (2),

$$\zeta \sigma_1 + \sum_{j=2}^g \zeta_{1j} \sigma_j = 0, \quad \zeta \sigma_2 - \sum_{k=1}^g \zeta_{2k} \sigma_k = 0,$$

and, by linear independence of $\{\sigma_i\}$, $1 \leq i \leq g$,

$$\zeta = \zeta_{1j} = \zeta_{2k} = 0 ,$$

$$2 \leq j \leq g, 1 \leq k \leq g. \quad \square$$

Let $W(P)$ be the Wronskian $W(v_1, \dots, v_N)(P)$ of the basis $\{v_j\}$ at a generic point $P \in \Sigma$ and $\hat{W}_{kl}(P)$ the Wronskian $W(v_1, \dots, v_{k-1}, v_l, v_{k+1}, \dots, v_N)(P)$ at P .

Lemma. *The $(g-2)(g-3)/2$ linearly independent relations*

$$v_l(z)W(P) = \sum_{k=1}^N v_k(z)\hat{W}_{kl}(P) , \quad (3)$$

$l = N+1, \dots, M, k = 1, \dots, N$, hold $\forall z \in \Sigma$.

Proof. Consider the $(N+1) \times (N+1)$ matrix obtained by adding to the matrix associated to $W(P)$ the column $(v_1(z), \dots, v_N(z))^t$ and the row $(v_l(P), \dots, \partial^{N-1}v_l(P), v_l(z))$. Expanding v_l over the basis $\{v_j\}$, we get

$$(v_l(P), \dots, \partial^{N-1}v_l(P), v_l(z)) = \sum_{j=1}^N C_{lj}(v_j(P), \dots, \partial^{N-1}v_j(P), v_j(z)) .$$

Then the determinant of this matrix vanishes, which is the content of the Lemma. \square

Remarks.

1. The ratio

$$\frac{\hat{W}_{kl}(P)}{W(P)} ,$$

does not depend on P .

2. Since for $i = 1, \dots, g$

$$\begin{aligned} v_j(p_i) &= \delta_{ji} , & j &= 1, \dots, g , \\ v_j(p_i) &= 0 , & j &= g+1, \dots, M , \end{aligned} \quad (4)$$

it follows that setting $z = p_i$ in (3) we obtain

$$\hat{W}_{il}(P) = 0 ,$$

for $i = 1, \dots, g$ and $l = N + 1, \dots, M$.

3. The Wronskians in the expansion (3) can be replaced by the corresponding determinant $\det v_j(x_i)$, where x_1, \dots, x_{3g-3} are arbitrary points on Σ .

Proof of Theorem 1. Assume that $\det \omega(p_1, \dots, p_g) \neq 0$. Let us define

$$x_j := \begin{cases} p_j, & j = 1, \dots, g, \\ q_{j-g}, & j = g + 1, \dots, N + 1. \end{cases}$$

Fix k, l with $3 \leq k < l \leq g$ and consider the matrix

$$\begin{pmatrix} v_1(x_1) & \dots & v_N(x_1) & \sigma_k(x_1)\sigma_l(x_1) \\ \vdots & \ddots & \vdots & \vdots \\ v_1(x_{N+1}) & \dots & v_N(x_{N+1}) & \sigma_k(x_{N+1})\sigma_l(x_{N+1}) \end{pmatrix}.$$

By (4), this matrix has $\text{diag}(1, \dots, 1)$ in the $g \times g$ upper left corner, 0 in the $g \times (2g - 2)$ upper right corner and $(\det \omega(p_1, \dots, p_g))^{-2} A(k, l)$ in the $(2g - 2) \times (2g - 2)$ lower right corner. On the other hand, by the lemma and the last remark the determinant of this matrix vanishes. Since such relations hold for (p_1, \dots, p_g) in a dense subset in Σ^g , they hold $\forall (p_1, \dots, p_g) \in \Sigma^g$ and the theorem follows. \square

Proof of the Corollary. Divide the relations of Theorem 1 by $\Delta_{i, 2g-2}$ and note that

$$a_{mn,r} = \sum_{i,j=1}^g D_{mi} D_{nj} \omega_i(q_r) \omega_j(q_r).$$

Independence of the points q_j , $1 \leq j \leq 2g - 2$, follows by noting that the coefficients of $\omega_i \omega_j$ in the relations are functions of q_j with no zeroes or poles.

We express the v_k in the matrix form

$$v_k = \sigma^t \cdot C_k \cdot \sigma,$$

where

$$(C_k)_{ij} := \begin{cases} \delta_{ik} \delta_{jk}, & k = 1, \dots, g, \\ \frac{1}{2}(\delta_{im} \delta_{j(l+m)} + \delta_{i(l+m)} \delta_{jm}), & k = l + m(2g - m + 1)/2, \\ & m = 1, \dots, g - 1, \quad l = 1, \dots, g - m. \end{cases}$$

Let $\hat{W}_{k,ij}(P)$ be the Wronskian $W(v_1, \dots, v_{k-1}, \omega_i \omega_j, v_{k+1}, \dots, v_N)(P)$ at P and set

$$(ij|lm) := \sum_{k=1}^N \frac{\hat{W}_{k,ij}}{W} ([\omega]^{-1t} \cdot C_k \cdot [\omega])_{lm} - \frac{1}{2}(\delta_{il}\delta_{jm} + \delta_{im}\delta_{jl}) .$$

Theorem 3. *The following relations*

$$\sum_{l,m=1}^g (ij|lm) \omega_l(z) \omega_m(z) = 0 , \quad (5)$$

$1 \leq i, j \leq g, \forall z \in \Sigma$, hold. Furthermore, if the basis $\omega_1, \dots, \omega_g$ depends only on the marking of the Riemann surface, then the coefficients $(ij|lm)$ depend only on the Riemann period matrix.

Proof. Eq.(5) is equivalent to the relations

$$\omega_i(z) \omega_j(z) = \sum_{k=1}^N \frac{\hat{W}_{k,ij}}{W} v_k(z) , \quad (6)$$

$i, j = 1, \dots, M, \forall z \in \Sigma$, which follow by the lemma. □

The induced measure. Let $\{\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g\}$ be a symplectic basis of the first homology group $H_1(\Sigma, \mathbb{Z})$ and let $\{\omega_1, \dots, \omega_g\}$ be the dual basis of $H^0(\Sigma, K)$, i.e. the basis satisfying the following condition

$$\oint_{\alpha_i} \omega_j = \delta_{ij} ,$$

$1 \leq i, j \leq g$. Let τ be the Riemann period matrix of Σ , defined by

$$\tau_{ij} := \oint_{\beta_i} \omega_j ,$$

$1 \leq i, j \leq g$. A different symplectic basis $\{\tilde{\alpha}_1, \dots, \tilde{\alpha}_g, \tilde{\beta}_1, \dots, \tilde{\beta}_g\}$ of $H_0(\Sigma, \mathbb{Z})$ satisfies

$$\begin{pmatrix} \tilde{\alpha} \\ \tilde{\beta} \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} ,$$

where $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(2g, \mathbb{Z})$, and the corresponding Riemann period matrices are related by the modular transformation

$$\tilde{\tau} = (A\tau + B) \cdot (C\tau + D)^{-1} .$$

Note that the basis $\{\sigma_i\}$ defined in Eq.(1) does not depend on the choice of the basis of $H_0(\Sigma, K)$ and therefore is modular invariant.

It can be proved that $\tau \in \mathcal{H}_g$, where

$$\mathcal{H}_g := \{Z \in M_g(\mathbb{C}) \mid {}^t Z = Z, \operatorname{Im} Z > 0\} ,$$

is the Siegel upper half space. The line element ds on \mathcal{H}_g

$$ds^2 := \operatorname{Tr} [(\operatorname{Im} Z)^{-1} dZ (\operatorname{Im} Z)^{-1} d\bar{Z}] ,$$

defines the volume element

$$\frac{|\wedge_{i \leq j}^g dZ_{ij}|^2}{(\det \operatorname{Im} Z)^{g+1}} .$$

Let k be the Kodaira-Spencer map identifying the quadratic differentials on Σ with the fiber of the cotangent of the Teichmüller space at the point representing Σ . We have

$$k(\omega_i \omega_j) = (2\pi i)^{-1} d\tau_{ij} ,$$

where $\{\omega_1, \dots, \omega_g\}$ is the standard basis of the holomorphic abelian differentials. By (5) we have

$$\sum_{l,m=1}^g (ij|lm) d\tau_{lm} = 0 ,$$

$i, j = 1, \dots, g$. Let us set

$$d\Xi_i := 2\pi i k(v_i) , \quad i = 1, \dots, N .$$

By Eq.(6) we have

$$d\tau_{ij} = \sum_{k=1}^N \frac{\hat{W}_{k,ij}}{W} d\Xi_k , \tag{7}$$

$i, j = 1, \dots, g$.

Theorem 4. *The line element $ds|_{\hat{\mathcal{M}}_g}$ on $\hat{\mathcal{M}}_g$ induced by the Siegel metric is*

$$ds^2|_{\hat{\mathcal{M}}_g} := \sum_{i,j=1}^N g_{ij} d\Xi_i d\bar{\Xi}_j ,$$

where

$$g_{ij} := |W|^{-2} \operatorname{Tr} [(\operatorname{Im} \tau)^{-1} \hat{W}_i (\operatorname{Im} \tau)^{-1} \bar{\hat{W}}_j] .$$

It follows that the Siegel induced modular invariant volume form on $\hat{\mathcal{M}}_g$ is

$$d\nu := (\det g)^{\frac{1}{2}} d\Xi_1 \wedge \dots \wedge d\bar{\Xi}_N .$$

Remarks.

1. Relations between the τ_{ij} follow by integrating Eq.(7). Such relations should appear in studying the structure of the spectrum of the Laplacian acting on degree zero bundles. We may expect that in the case of period matrices of branched covering of the torus, considered in [2], which in turn correspond to Jacobians with a particular CM, the derived relations lead to identities of number theoretical interest.
2. The relationships for the quadratic differentials lead to θ -identities that will be considered in a forthcoming publication. We expect that the constraints on the Riemann period matrix naturally lead to the construction of modular functions.
3. The measure $d\nu$ and permutations of products of determinants of abelian differentials appear in the superstring amplitudes [3].

Acknowledgements. Work partially supported by the European Community's Human Potential Programme under contract MRTN-CT-2004-005104 "Constituents, Fundamental Forces and Symmetries of the Universe".

References

- [1] E. Arbarello, M. Cornalba, P.A. Griffiths and J. Harris, “Geometry of Algebraic Curves,” Vol.1. Berlin, Heidelberg, New-York, Tokyo, 1985.
- [2] M. Matone, “Eigenfunctions of the Laplacian Acting on Degree Zero Bundles over Special Riemann Surfaces,” *Trans. Am. Math. Soc.* **356**, 2989 (2004) [math.ag/0105051](#).
- [3] M. Matone and R. Volpato, “Higher genus superstring amplitudes from the geometry of moduli spaces,” *Nucl. Phys. B* **732**, 321 (2006) [hep-th/0506231](#).