

Families of Painlevé VI equations having a common solution

Bassem BEN HAMED *

Faculté des Sciences de Sfax, Département de Mathématiques
BP 802, route Soukra km 4 Sfax 3018, Tunisia

Lubomir GAVRILOV
Université Paul Sabatier, MIG
Laboratoire Emile Picard, UMR 5580
31062 Toulouse Cedex 9, France

November 11, 2018

Abstract

We classify all functions satisfying non-trivial families of \mathbf{PVI}_α equations. It turns out that, up to an Okamoto equivalence, there are exactly four families parameterized by affine planes or lines. Each affine space is generated by points of "geometric origin", associated either to deformations of elliptic surfaces with four singular fibers, or to deformations of three-sheeted covers of \mathbb{P}^1 with branching locus consisting of four points.

*The first author was partially supported by a grant of the Faculty of Sciences of Sfax, Tunisia.

1 Introduction

Consider the Painlevé VI (\mathbf{PVI}_α) equation

$$\begin{aligned} \frac{d^2\lambda}{dt^2} = & \frac{1}{2}\left(\frac{1}{\lambda} + \frac{1}{\lambda-1} + \frac{1}{\lambda-t}\right)\left(\frac{d\lambda}{dt}\right)^2 - \left(\frac{1}{t} + \frac{1}{t-1} + \frac{1}{\lambda-t}\right)\frac{d\lambda}{dt} \\ & + \frac{\lambda(\lambda-1)(\lambda-t)}{t^2(t-1)^2}\left[\alpha_0 - \alpha_1\frac{t}{\lambda^2} + \alpha_2\frac{t-1}{(\lambda-1)^2} + \left(\frac{1}{2} - \alpha_3\right)\frac{t(t-1)}{(\lambda-t)^2}\right]. \end{aligned}$$

parameterized by $\alpha = (\alpha_0, \alpha_1, \alpha_2, \alpha_3) \in \mathbb{C}^4$. Although any solution of \mathbf{PVI}_α , for generic α_i , is transcendental (and even a "new transcendental function") there is a large amount of solutions which are algebraic in t . Their general classification is still an open problem (e.g. [15, Manin]), except in the particular case $\alpha_1 = \alpha_2 = \alpha_3 = 0$ [4, Dubrovin, Mazzocco]. The present paper addresses the question of classifying *families* of algebraic solutions. The simplest case occurs when a given algebraic solution satisfies each member of a non-trivial family of \mathbf{PVI}_α equations. By a non-trivial family of \mathbf{PVI}_α equations we mean a set $\{\mathbf{PVI}_\alpha\}_\alpha$ containing at least two distinct elements corresponding to, say, α' and α'' . Then this solution satisfies the \mathbf{PVI}_α equations corresponding to the affine line containing α' and α'' . It follows that each non-trivial family as above corresponds to an affine plane in the parameter space $\mathbb{C}^4\{\alpha\}$. We classify all such affine spaces, together with their associated algebraic solutions (Theorem 1). The proof of Theorem 1 does not use the notion of Picard-Fuchs equation. It turns out that the solutions which we obtain coincide surprisingly with the solutions obtained earlier by Doran, who used deformations of elliptic surfaces with four singular fibers and the related Picard-Fuchs equations, see Theorem 2.

The second purpose of the paper is to give a (partial) explanation of the above coincidence. It is well known that each solution $(\lambda(t), \alpha)$ of a given \mathbf{PVI}_α equation governs the isomonodromy deformation of a 2×2 Fuchsian system with four singular points. We say that such a deformation is "geometric", if there is a fundamental matrix of solutions whose entries are Abelian integrals depending algebraically on the deformation parameter, see [3, Doran]. A geometric deformation of a Fuchsian system is isomonodromic and the associated solution of the corresponding Schlesinger system (or \mathbf{PVI}_α equation) is an algebraic function in t . When this holds true, we say that the algebraic solution $(\lambda(t), \alpha)$ of \mathbf{PVI}_α is of geometric origin. The solutions found by Doran are of geometric origin and they correspond to special values

of α which belong to the affine spaces of \mathbf{PVI}_α equations described in Theorem 1. We shall prove that the same list of solutions can be obtained from deformations of ramified covers of \mathbb{P}^1 with four ramification points. The corresponding values of the parameters α are different and are shown on Table 5, see Theorem 3. Finally we note that all these points of geometric origin generate the affine planes of \mathbf{PVI}_α equations, described in Theorem 1.

Acknowledgments. The solution 1A, as well its relation to the family 2A was pointed out to us by Philip Boalch. We acknowledge his interest and critical remarks.

2 Families of Painlevé VI equations having a common solution

Let $\lambda = \lambda(t)$ be a solution of the equations $\mathbf{PVI}_{\alpha'}$, $\mathbf{PVI}_{\alpha''}$, $\alpha' \neq \alpha''$. Then $\lambda(t)$ satisfies the implicit equation

$$\beta_0 - \beta_1 \frac{t}{\lambda^2} + \beta_2 \frac{t-1}{(\lambda-1)^2} - \beta_3 \frac{t(t-1)}{(\lambda-t)^2} = 0 \quad (1)$$

where $\beta = \alpha' - \alpha'' = (\beta_0, \beta_1, \beta_2, \beta_3)$, and hence it is an algebraic function. The function $\lambda(t)$ satisfies, moreover, the family $\{\mathbf{PVI}_\alpha\}_\alpha$, where α belongs to the affine line

$$\{\alpha' + s(\alpha' - \alpha'') : s \in \mathbb{C}\} \subset \mathbb{C}^4.$$

It is seen from this that the set of all α such that \mathbf{PVI}_α is satisfied by the function $\lambda(t)$ form an affine subspace of \mathbb{C}^4 . We refer to the set of these \mathbf{PVI}_α equations as to a *family of Painlevé VI equations having a common solution*.

Theorem 1 *The list of all families of Painlevé VI equations having a common solution, together with the corresponding solution, is shown in Table 1.*

Remark 1 *Each solution $\lambda(t)$ is defined by a relation $P(\lambda(t), t) \equiv 0$ where P is an irreducible polynomial given in Table 1. The solutions in each of the 5 series of families on Table 1 are equivalent up to a S_4 -symmetry of Painlevé VI equation (see section 2.1).*

Moreover, the solutions $1A, 1B, \dots, 1F$ are Okamoto equivalent to the solutions $2A, 2B, 2C$. More precisely, the solution $1A$

$$a\lambda^2 - bt = 0, \alpha = (a, b, \frac{1}{8}, \frac{1}{8})$$

is equivalent, after applying the transformation w_2 of Okamoto [16, p.363], to the solution $\lambda^2 - t = 0$ where the parameter α equals to

$$\left(\frac{\sqrt{2a} - \sqrt{2b}}{2\sqrt{2}}\right)^2, \left(\frac{\sqrt{2a} + \sqrt{2b}}{2\sqrt{2}}\right)^2, \left(1 - \frac{\sqrt{2a} - \sqrt{2b}}{2\sqrt{2}}\right)^2, \left(1 - \frac{\sqrt{2a} + \sqrt{2b}}{2\sqrt{2}}\right)^2.$$

Thus, up to Okamoto equivalence, the families of Painlevé VI equations having a common solution are represented (for instance) by the four families $2A, 3A, 4A, 5A$ on Table 1.

Remark 2 The functions $0, 1, t$ are not considered as solutions of Painlevé VI equation, and therefore are excluded from Table 1.

Outline of the Proof. Denote by Γ_β the compactified and normalized algebraic curve defined by (1), with affine model

$$\Gamma_\beta^{aff} = \{(\lambda, t) \in \mathbb{C}^2 : N(\lambda, t) = 0\} \quad (2)$$

where

$$\begin{aligned} N(\lambda, t) &= \beta_0 \lambda^2 (\lambda - 1)^2 (\lambda - t)^2 - \beta_1 t (\lambda - 1)^2 (\lambda - t)^2 \\ &+ \beta_2 (t - 1) \lambda^2 (\lambda - t)^2 - \beta_3 t (t - 1) \lambda^2 (\lambda - 1)^2 = 0. \end{aligned}$$

In the case when Γ_β is irreducible, the relation $\{N(\lambda, t) = 0\}$ defines an algebraic function $\lambda(t)$. If this function were a solution of some \mathbf{PVI}_α equation, then the only ramification points of $\lambda(t)$ would be at $t = 0, 1, \infty$ (because \mathbf{PVI}_α satisfies the so called Painlevé property [10]). Equivalently, the pair (Γ_β, t) is a Belyi pair, which means that the only possible critical values of the map

$$\pi : \Gamma_\beta \rightarrow \mathbb{CP}^1 : (\lambda, t) \rightarrow t \quad (3)$$

are $0, 1$ or ∞ . This means also that if $\Delta(t)$ is the discriminant of $N(\lambda, t)$ with respect to λ , then it is a polynomial whose only roots are at $t = 0$ and $t = 1$. A direct computation shows that this is impossible. The more difficult case is when $N(\lambda, t)$ is reducible over \mathbb{C} . Then Γ_β defines several algebraic

Name	Solution of \mathbf{PVI}_α equation	\mathbf{PVI}_α equation
1A	$a\lambda^2 - bt$	$(a, b, \frac{1}{8}, \frac{1}{8})$
1B	$a(\lambda - 1)^2 + b(t - 1)$	$(a, \frac{1}{8}, b, \frac{1}{8})$
1C	$a(\lambda - t)^2 - bt(t - 1)$	$(a, \frac{1}{8}, \frac{1}{8}, b)$
1D	$-at(\lambda - 1)^2 + b(t - 1)\lambda^2$	$(\frac{1}{8}, a, b, \frac{1}{8})$
1E	$a(\lambda - t)^2 + b(t - 1)\lambda^2$	$(\frac{1}{8}, a, \frac{1}{8}, b)$
1F	$a(\lambda - t)^2 - bt(\lambda - 1)^2$	$(\frac{1}{8}, \frac{1}{8}, a, b)$
2A	$\lambda^2 - t$	(a, a, b, b)
2B	$\lambda^2 - 2\lambda + t$	(a, b, a, b)
2C	$\lambda^2 - 2\lambda t + t$	(b, a, a, b)
3A	$\lambda^4 - 6\lambda^2 t + 4\lambda t + 4\lambda t^2 - 3t^2$	$(a, 9a, a, a)$
3B	$3\lambda^4 - 4\lambda^3 - 4\lambda^3 t + 6\lambda^2 t - t^2$	$(9a, a, a, a)$
3C	$\lambda^4 - 4\lambda^3 + 6t\lambda^2 - 4t^2\lambda + t^2$	$(a, a, 9a, a)$
3D	$\lambda^4 - 4t\lambda^3 + 6t\lambda^2 - 4t\lambda + t^2$	$(a, a, a, 9a)$
4A	$\lambda^4 - 2t\lambda^3 - 2\lambda^3 + 6t\lambda^2$ $- 2t^2\lambda - 2t\lambda + t^3 - t^2 + t$	$(a, \frac{1}{8}, a, a)$
4B	$\lambda^4 - 2t\lambda^3 + 2t^2\lambda - t^3$	$(a, a, \frac{1}{8}, a)$
4C	$\lambda^4(t^2 - t + 1) - 2\lambda^3 t(t + 1) + 6t^2\lambda^2$ $- 2\lambda t^2(t + 1) + t^3$	$(\frac{1}{8}, a, a, a)$
4D	$\lambda^4 - 2\lambda^3 + 2t\lambda - t$	$(a, a, a, \frac{1}{18})$
5A	$-2\lambda^3 + 3t\lambda^2 + 3\lambda^2 - 6t\lambda + t^2 + t$	$(4a, \frac{1}{8}, a, a)$
5B	$\lambda^3 - 3\lambda^2 + 3t\lambda - 2t^2 + t$	$(a, \frac{1}{18}, 4a, a)$
5C	$\lambda^3 - 3t\lambda^2 + 3t\lambda + t^2 - 2t$	$(a, \frac{1}{18}, a, 4a)$
5D	$2\lambda^3 - 3t\lambda^2 + t^2$	$(4a, a, \frac{1}{18}, a)$
5E	$\lambda^3 - 3t\lambda + 2t^2$	$(a, 4a, \frac{1}{18}, a)$
5F	$\lambda^3 - 3t\lambda^2 + 3t\lambda - t^2$	$(a, a, \frac{1}{18}, 4a)$
5G	$\lambda^3(2 - t) - 3t\lambda^2 + 3t^2\lambda - t^2$	$(\frac{1}{18}, a, 4a, a)$
5H	$\lambda^3(t + 1) - 6t\lambda^2 + 3t(t + 1)\lambda - 2t^2$	$(\frac{1}{18}, 4a, a, a)$
5I	$(1 - 2t)\lambda^3 + 3t\lambda^2 - 3t\lambda + t^2$	$(\frac{1}{18}, a, a, 4a)$
5J	$\lambda^3 - 3\lambda^2 + 3t\lambda - t$	$(a, a, 4a, \frac{1}{18})$
5K	$\lambda^3 - 3t\lambda + 2t$	$(a, 4a, a, \frac{1}{18})$
5L	$\lambda^3 - 3\lambda^2 + t$	$(4a, a, a, \frac{1}{18})$

Table 1: List of all algebraic solutions satisfying families of \mathbf{PVI}_α equations

functions and we have to apply the above to each of them. Finally we have to check whether the obtained function is actually a solution of some \mathbf{PVI}_α equation. To check whether a given polynomial $N(\lambda, t)$ is reducible over \mathbb{C} is a difficult task in general. We shall make use of the action of the symmetric group \mathcal{S}_4 (see section 2.1) on the set of curves Γ_β , parameterized by $\beta \in \mathbb{CP}^3$.

It turns out that, first, curves Γ_β with a trivial stabilizer under the action of \mathcal{S}_4 can not produce a solution of \mathbf{PVI}_α . The stabilizer of a curve acts on it as a group of automorphisms (symmetries) which imposes additional restrictions on β .

The second ingredient of the proof is the study of the Puiseux expansion of $\lambda(t)$ in a neighborhood of $t = 0, 1, \infty$ (section 2.2). These expansions depend on the stabilizer of Γ_β only and imply the possible topological types of the solution $\lambda(t)$. Equivalently, to each solution $\lambda(t)$ we associate a Belyi pair and the Puiseux expansions determine their possible *dessin d'enfant*. The algebraic functions which we obtain in this way are, *a posteriori*, the solutions of the \mathbf{PVI}_α presented in Table 1.

Proof of Theorem 1.

2.1 The action of \mathcal{S}_4

The set of automorphisms of the projective line \mathbb{CP}^1 which send four distinct points $(0, 1, t, \infty)$ to the points $(0, 1, \tilde{t}, \infty)$ ($\tilde{t} = \tilde{t}(t)$ is uniquely defined) form a group isomorphic to \mathcal{S}_4 generated by the transpositions

$$x^1 : s \mapsto 1 - s, x^2 : s \mapsto \frac{1}{s}, x^3 : s \mapsto \frac{t - s}{t - 1}. \quad (4)$$

Each x^i sends an isomonodromic family of Fuchsian systems with singular points at $0, 1, t, \infty$ to an isomonodromic family of such systems with singular points at $0, 1, \tilde{t}, \infty$. Therefore x^i induce an action of \mathcal{S}_4 on the set of \mathbf{PVI}_α equations, and hence on the set of curves Γ_β . Explicitly we have

$$x^i : \Gamma_\beta \rightarrow \Gamma_{x_*^i(\beta)} : (\lambda, t) \rightarrow (x^i(\lambda), x^i(t)), i = 1, 2 \quad (5)$$

and

$$x^3 : \Gamma_\beta \rightarrow \Gamma_{x_*^3(\beta)} : (\lambda, t) \rightarrow (x^3(\lambda), x^3(0)) = \left(\frac{t - \lambda}{t - 1}, \frac{t}{t - 1} \right) \quad (6)$$

where

$$\begin{cases} x_*^1 : (\beta_0, \beta_1, \beta_2, \beta_3) & \rightarrow (\beta_0, \beta_2, \beta_1, \beta_3) \\ x_*^2 : (\beta_0, \beta_1, \beta_2, \beta_3) & \rightarrow (\beta_1, \beta_0, \beta_2, \beta_3) \\ x_*^3 : (\beta_0, \beta_1, \beta_2, \beta_3) & \rightarrow (\beta_0, \beta_3, \beta_2, \beta_1) \end{cases} \quad (7)$$

which is the standard representation of \mathcal{S}_4 on \mathbb{C}^4 (upon identifying $\infty, 0, 1, t$ to $\beta_0, \beta_1, \beta_2, \beta_3$ respectively). The proof of the above facts is straightforward, see [10] for details.

2.2 The topological type of the projection $\Gamma_\beta \rightarrow \mathbb{CP}^1$ in a neighborhood of the pre-image of $t = 0, 1, \infty$

Let Γ_β be the compactified and normalized curve defined by (2) (it is a disjoint union of Riemann surfaces). In this section we determine the topological type of the projection (3)

$$\Gamma_\beta \rightarrow \mathbb{CP}^1$$

in a neighborhood of the pre-image of $t = 0, 1, \infty$ in Γ_β . In the projective space \mathbb{CP}^3 with coordinates $[\beta_0 : \beta_1 : \beta_2 : \beta_3]$ consider the complex polyhedron W formed by the ten planes (2-faces)

$$W = \cup_{i \neq j} \{\beta_i = \beta_j\} \cup_k \{\beta_k = 0\}. \quad (8)$$

It has also 45 1-faces (projective lines) and 120 0-faces (points). We shall see in the process of the proof that the topological type of the projection in a neighborhood of the pre-image of $t = 0, 1, \infty$ is one and the same when β belongs to a given i -face, but does not belong to any other j -face with $j < i$. For this reason we shall use, until the end of this paper, the following convention. *When we say that a point β belongs to a given face (satisfies some set of relations (8)), then this will mean that it does not belong to any other face of smaller dimension (does not satisfy any other relation from the list (8)).* The topological type in a neighborhood of the pre-image of any point is determined by a partition of the degree of the map which is 6. Thus a partition $(1 + 1 + 1 + 1 + 2)$ means that we have 5 pre-images, and that the multiplicity of π at each pre-image is 1, 1, 1, 1, 2 respectively. Similarly, a partition $(1 + 1 + 2 + 2)$ means that have 4 pre-images with multiplicities 1, 1, 2, 2 respectively etc. To formulate the result we note that the symmetric group \mathcal{S}_4 acts on the polyhedron W by its standard representation (7), as well on the set of curves Γ_β by (5). The subgroup \mathcal{S}_3 generated by x^1, x^2 permutes the ramification points $0, 1, \infty$ according to (4) without changing the topological type of the projection π over each of these points.

Proposition 1 *The topological type of the projection (3) in a neighborhood of the pre-image of $t = 0, 1, \infty$ is one and the same when β belongs to a given*




Figure 1: The Newton polygon of $N(\lambda, t)$ and $N^0(\lambda, t)$

face of the polyhedron W or it does not belong to W . This topological type is shown on Table 2 (one representative for each orbit of $\mathcal{S}_3 = \langle x^1, x^2 \rangle$)

Proof The bi-rational transformations x^1, x^2 defined by (5) are compatible with the projection π and permute the points $t = 0, 1, \infty$. Therefore it suffices to consider the pre-image of 0. Let us consider in detail the "generic" case, when $\beta \notin W$. It follows from the Newton polygon of $N(\lambda, t)$, shown on fig. 1, that there are at least three Puiseux series in a neighborhood of $(0, 0)$ (for the terminology see for instance [12, Kirwan]). The first two correspond to the line segment $[(3, 0), (1, 2)]$ and have non-equivalent leading terms

$$\lambda = c_1 t + \dots, \lambda = c_2 t + \dots$$

where

$$(\beta_3 - \beta_1)c_{1,2}^2 + 2\beta_1 c_{1,2} - \beta_1 = 0$$

provided that

$$\beta_1 \neq \beta_3, \beta_1^2 + \beta_1(\beta_3 - \beta_1) \neq 0.$$

The third one corresponds to the line segment $[(1, 2), (0, 4)]$ and has leading term

$$\lambda = c_3 t^{1/2} + \dots$$

where

$$(\beta_0 - \beta_2)c_3^2 + \beta_3 - \beta_1 = 0,$$

provided that

$$\beta_0 \neq \beta_2, \beta_1 \neq \beta_3.$$

Taking into consideration that

$$N(\lambda, 0) = \lambda^4(\beta_0\lambda^2 - 2\beta_0\lambda + \beta_0 - \beta_2)$$

we conclude that we have at least five pre-images of multiplicities at least 1, 1, 2, 1, 1 respectively. As the degree of the map π is six, then its topological type is exactly $(1 + 1 + 1 + 1 + 2)$. The topological type of the projection π over 0 and 1 is obtained by acting with the group \mathcal{S}_3 generated by x^1, x^2 .

In a similar way one verifies that when $\beta_0 = \beta_2$, or $\beta_1 = \beta_3$ the multiplicities are $(1 + 1 + 1 + 3)$. If $\beta_0 = \beta_2$ and $\beta_1 = \beta_3$ the multiplicities are $(1 + 1 + 2 + 2)$. The case $\beta_0 = \beta_1 = \beta_3$ is the same as $\beta_0 = \beta_2$ and the multiplicity is $(1 + 1 + 1 + 3)$. The case $\beta_0 = \beta_1 = \beta_2 = \beta_3$ is of the same type as $\beta_0 = \beta_2$ and $\beta_1 = \beta_3$. The multiplicities of π over 1 and ∞ are obtained as before. This completes the study of faces of W for which $\beta_i \neq 0$. In the case $\beta_3 = 0$ we consider the curve

$$\Gamma_\beta^0 = \{(\lambda, t) \in \mathbb{C}^2 : \beta_0 - \beta_1 \frac{t}{\lambda^2} + \beta_2 \frac{t-1}{(\lambda-1)^2} = 0, \lambda \neq 0, 1\}.$$

The polynomial $N(\lambda, t)$ is replaced by

$$N^0(\lambda, t) = \beta_0\lambda^2(\lambda-1)^2 - \beta_1t(\lambda-1)^2 + \beta_2(t-1)\lambda^2$$

whose Newton polygon is shown on fig. 1. It follows that there is at least one Puiseux expansion with leading term

$$\lambda = ct^{1/2} + \dots, \beta_1 + (\beta_2 - \beta_0)c^2 = 0$$

provided that $\beta_1 \neq 0, \beta_2 \neq \beta_0$. As

$$N^0(\lambda, 0) = \lambda^2(\beta_0(\lambda-1)^2 - \beta_2)$$

then $t = 0$ has at least three pre-images, provided that $\beta_0\beta_2 \neq 0$. We conclude that $t = 0$ has exactly three pre-images with multiplicities 2, 1, 1 respectively, provided that β belongs to the 2-face $\beta_3 = 0$. The remaining 1-faces and 0-faces are studied in the same way. The result is summarized on Table 2. It worth noting that in all cases the computing of the leading term of the Puiseux expansion suffices to deduce the result.

face of W	stabilizer	$t = 0$	$t = 1$	$t = \infty$
$\beta_i \neq \beta_j$	S_4	(1+1+1+1+2)	(1+1+1+1+2)	(1+1+1+1+2)
$\beta_0 = \beta_2$	$S_2 \times S_2$	(1+1+1+3)	(1+1+1+1+2)	(1+1+1+1+2)
$\beta_0 = \beta_2, \beta_1 = \beta_3$	D_4	(1+1+2+2)	(1+1+1+1+2)	(1+1+1+1+2)
$\beta_0 = \beta_1 = \beta_2$	S_3	(1+1+1+3)	(1+1+1+3)	(1+1+1+3)
$\beta_0 = \beta_1 = \beta_2 = \beta_3$	S_4	(1+1+2+2)	(1+1+2+2)	(1+1+2+2)
$\beta_3 = 0$	S_3	(1+1+2)	(1+1+2)	(1+1+2)
$\beta_0 = \beta_2, \beta_3 = 0$	S_2	(1+3)	(1+1+2)	(1+1+2)
$\beta_0 = \beta_1 = \beta_2, \beta_3 = 0$	S_3	(1+3)	(1+3)	(1+3)
$\beta_2 = \beta_3 = 0$	$S_2 \times S_2$	(2)	(1+1)	(2)
$\beta_2 = \beta_3 = 0, \beta_0 = \beta_1$	$S_2 \times S_2$	(2)	(1+1)	(2)

Table 2: Multiplicity of π at the pre-images of $t = 0, 1, \infty$.

We conclude this section by the following elementary claim which will be often useful in the computations

Proposition 2 *Let $N_1(\lambda, t)$ be a polynomial of non-zero degree with respect to λ and of non-zero degree with respect to t , which divides $N(\lambda, t)$, and $\beta_1\beta_2\beta_3 \neq 0$. Then*

$$N_1(0, t) = c_0 t^{n_0}, c_0 \neq 0, 1 \leq n_0 \leq 3, N_1(1, t) = c_1 (t-1)^{n_1}, c_1 \neq 0, 1 \leq n_1 \leq 3$$

and

$$N_1(t, t) = c_2 t^{m_0} (t-1)^{m_1}, c_2 \neq 0, 1 \leq m_0, 1 \leq m_1, m_0 + m_1 \leq 3.$$

Proof. We have $N(0, t) = -\beta_1 t^3$. For a fixed $\lambda = c \sim 0$ the polynomial $N(c, t) \in \mathbb{C}[t]$ has exactly three roots which tend to zero when c tends to zero. Therefore the polynomial $N_1(c, t) \in \mathbb{C}[t]$ has at least one and at most three roots which tends to zero when c tends to zero, which proves the claim concerning $N_1(0, t)$. The claim concerning $N_1(1, t)$ is proved in the same way. As $N_1(0, 0) = N_1(1, 1) = 0$ then $N_1(t, t)$ is divided by $t(t-1)$ but also divides $N(t, t) = -\beta_3 t^3 (t-1)^3$. \square

We are ready to compute the solutions of \mathbf{PVI}_α corresponding to the faces of W . Let Γ be the Riemann surface of an irreducible component of Γ_β , which defines a solution of some \mathbf{PVI}_α equation. Then the only ramification points of the induced map

$$\pi : \Gamma \rightarrow \mathbb{CP}^1 : (\lambda, t) \rightarrow t \tag{9}$$




Figure 2: Dessins of degree 1, 2 and 3

are at $0, 1, \infty$, and Γ is connected. The pair (Γ, π) is called a Belyi pair, to which we associate a *dessins d'enfant*, which is the graph obtained as a pre-image of the segment $[0, 1]$ under the map π . The degree of the dessin is the degree of π (see [18]). The dessin d'enfant will be useful when describing the topological type of the projection π .

2.3 The case $\beta \notin W$

We suppose that $\lambda(t)$ is an algebraic function, such that $N(\lambda(t), t) \equiv 0$, and consider the corresponding Belyi pair (Γ, π) , (9). Let $\{\lambda_1, \dots, \lambda_d\} = \pi^{-1}(t_0)$ where $t_0 \neq 0, 1, \infty$. The loops originating from t_0 and going clockwise once around $0, 1$ and ∞ induce permutations σ_0, σ_1 and σ_∞ of the points $\lambda_1, \dots, \lambda_d$, such that $\sigma_0 \sigma_1 \sigma_\infty = 1$. According to Table 2 σ_0, σ_1 and σ_∞ are transpositions, unless one of them is the identity permutation. In the former case $(\sigma_0 \sigma_1)^2 = 1$ and hence $\sigma_0 \sigma_1 = \sigma_1 \sigma_0$. Thus σ_∞ is a product of two disjoint transpositions, which contradicts to Table 2. On the other, if one of the permutations $\sigma_0, \sigma_1, \sigma_\infty$ is the identity, the group generated by them is either \mathbf{Z}_2 or is trivial. This shows that the degree of π is either two (because the covering (9) is connected), or one. The corresponding dessin d'enfants are shown on fig.2 (i) and (ii) (in particular $N(\lambda, t) \in \mathbb{C}[\lambda, t]$ is reducible). If the dessin is of degree one, then the solution is defined as $\lambda = P(t)$, where P is a polynomial (the coefficient of λ^6 in the polynomial $N(\lambda, t)$ is $\beta_0 \neq 0$). By Proposition 2 we conclude that either $\lambda = t$, or $\lambda = t^2$ or $\lambda = t^3$. But $\lambda - t$, $\lambda - t^2$, $\lambda - t^3$ can not divide $N(\lambda, t)$, provided that $\beta_i \neq 0$. If the dessin is of degree two, then $\lambda(t)$ is defined by $\lambda^2 + 2p(t)\lambda + q(t) = 0$. The functions p, q are polynomials in t , because the coefficient of λ^6 in the polynomial $N(\lambda, t)$ is $\beta_0 \neq 0$. Further, we may suppose (acting with an appropriate symmetry x^i on Γ , see section 2.1) that $\lambda(t)$ is ramified over 0 and ∞ only. By Proposition

2 $q(t)$ is a non-constant polynomial which divides t^3 . As $p(t)^2 - q(t)$ is a non-constant monomial of odd degree, then $p(t) \equiv 0$. Proposition 2 implies that we have either $\lambda^2 = t$ or $\lambda^2 = t^3$. The polynomial $\lambda^2 - t^3$ can not divide $N(\lambda, t)$ while $\lambda^2 - t$ divides $N(\lambda, t)$ if and only if $\beta_0 = \beta_1$ and $\beta_2 = \beta_3$ (this case is excluded, as $\beta \notin W$). *The curve Γ_β does not define a solution.*

2.4 The face $\beta_1 = \beta_2$

The possible dessins d'enfant are determined as above. Namely, when one of the permutations $\sigma_0, \sigma_1, \sigma_\infty$ is identity, the dessin is of degree one or two. Up to a symmetry it is equivalent to the one shown on fig. 2 (i) or (ii). Reasoning as in the case $\beta \notin W$ we conclude that Γ_β does not define a solution.

If, on the other hand, σ_0, σ_1 are non-trivial transpositions we have one more case compared to section 2.3: σ_3 is cyclic of order three, and σ_0, σ_1 are non-disjoined permutations. Taking into account that the covering (9) is connected, we conclude that the dessin is of degree three. Up to a symmetry, it is shown on 2 (iii) and $\lambda(t)$ satisfies $N_1(\lambda(t), t) \equiv 0$ where N_1 is an irreducible polynomial of degree three in λ dividing the polynomial $N(\lambda, t)$, defined after formula (2). We denote

$$N = N_1 N_2, \Gamma_1^{aff} = \{N_1(\lambda, t) = 0\}, \Gamma_2^{aff} = \{N_2(\lambda, t) = 0\}, \Gamma_\beta^{aff} = \Gamma_1^{aff} \cup \Gamma_2^{aff}.$$

As before, let $\Gamma_1, \Gamma_2, \Gamma_\beta$ be the corresponding compactified and normalized curves. The symmetry x^1 is an automorphism of Γ_β and hence it is either an automorphism of the curves Γ_1 and Γ_2 , or it permutes these curves (we used that Γ_1 is irreducible). Suppose first that x^1 is an automorphism of Γ_1 . Then the rational function

$$\frac{N_1(\lambda, t)}{\lambda(\lambda - 1)(\lambda - t)} = 1 + \frac{A_1}{\lambda} + \frac{B_1}{\lambda - 1} + \frac{C_1}{\lambda - t}$$

is invariant under the action of x^1 too. Here A_1, B_1, C_1 are polynomials in t which divide $t, t - 1$ and $t(t - \lambda)$ respectively (see (1)), and hence we have

$$A_1(1 - t) = -B_1(t), B_1(1 - t) = -A_1(t), C_1(1 - t) = -C_1(t).$$

Similarly, if

$$\frac{N_2(\lambda, t)}{\lambda(\lambda - 1)(\lambda - t)} = 1 + \frac{A_2}{\lambda} + \frac{B_2}{\lambda - 1} + \frac{C_2}{\lambda - t}$$

then

$$A_2(1-t) = -B_2(t), B_2(1-t) = -A_2(t), C_2(1-t) = -C_2(t).$$

We conclude that $C_1(t) = c_1t(t-1)$, $C_2(t) = c_2t(t-1)$, which contradicts to $C_1C_2 = -\beta_3t(t-1)/\beta_1$, $\beta_1, \beta_3 \neq 0$. Suppose now that the map x^1 exchanges the curves Γ_1 and Γ_2 . Then we have

$$A_1(1-t) = -B_2(t), B_1(1-t) = -B_2(t), C_1(1-t) = -C_2(t).$$

The polynomial $N(\lambda, t)$ is of degree three with respect to t and

$$A_1A_2 = -\frac{\beta_1}{\beta_0}t, B_1B_2 = -\frac{\beta_2}{\beta_0}(t-1), C_1C_2 = -\frac{\beta_3}{\beta_0}t(t-1).$$

Therefore without loss of generality we may suppose that $C_1(t) = c_1t$, $C_2 = c_1(t-1)$ and $A_1(t) = a_1t$, $B_2 = b_2(t-1)$ or $A_2(t) = a_2t$, $B_1 = b_1(t-1)$, where $a_i, b_j \neq 0$. In both cases the polynomials $N_1(\lambda, t)$, $N_2(\lambda, t)$ are of degree two in t , in contradiction to the fact that the degree of $N(\lambda, t)$ with respect to t is three. *We conclude that the curve Γ_β does not define a solution.*

2.5 The face $\beta_0 = \beta_2$, $\beta_1 = \beta_3$.

We have the identity

$$N(\lambda, t) = (\lambda^2 - 2\lambda + t)(\beta_0\lambda^2(\lambda - t)^2 - \beta_1t^2(\lambda - 1)^2).$$

Indeed,

$$\lambda^2 - 2\lambda + t = 0 \tag{10}$$

defines a solution of \mathbf{PVI}_α , e.g. [3], Table 2, solution 2B. Its dessin is equivalent to the one on fig. 2 (ii). The function $\lambda(t)$ defined by

$$\lambda(\lambda - t) - ct(\lambda - 1) = 0, c = \pm\sqrt{\frac{\beta_1}{\beta_0}}$$

is ramified over $0, 1, \infty$ only provided that $c = 0, \pm 1$. This is, however, impossible as $\beta_0 \neq \beta_1, \beta_i \neq 0$. *The curve Γ_β defines the solution (10).*

2.6 The face $\beta_0 = \beta_1 = \beta_2$.

According to Table 2 each of the permutations $\sigma_0, \sigma_1, \sigma_\infty$ is either the identity, or is a cycle of length three.

If one of the permutations $\sigma_0, \sigma_1, \sigma_\infty$ is the identity then the group generated by $\sigma_0, \sigma_1, \sigma_\infty$ is either \mathbf{Z}_3 or the trivial one $\{\mathbf{1}\}$, and hence the degree of the corresponding dessin is one or three. The case of degree one does not lead to a solution (see section 2.3). The case of degree three is studied as in section 2.4 and does not lead to a solution too (provided that $\beta_i \neq 0$).

If neither of the permutations $\sigma_0, \sigma_1, \sigma_\infty$ is the identity, then they are disjoint three-cycles. As the symmetric group \mathcal{S}_3 contains only two three-cycles we conclude that the degree of the projection π is at least four. Suppose that $\lambda(t)$ is defined by the polynomial N_1 , $N_1(\lambda(t), t) \equiv 0$, where $N_1 \in \mathbb{C}[\lambda, t]$ is irreducible of degree four in λ . Then x^1, x^2 are automorphisms of

$$\Gamma_1 = \{N_1(\lambda, t) = 0\}.$$

It follows that the curve

$$\Gamma_2 = \{N_2(\lambda, t) = 0\}$$

defined by the polynomial $N_2 = N/N_1$ is also invariant. We have

$$\frac{N_2(\lambda, t)}{\lambda(\lambda - 1)} = 1 + \frac{A}{\lambda} + \frac{B}{\lambda - 1}$$

where A, B are polynomials in t of degree at most three. The $x^{1,2}$ invariance of the above expression implies

$$A(t)A(1/t) = 1, A(1/t)B(t) = -B(1/t), B(t) = -A(1 - t).$$

with solutions

$$A(t) = t, B(t) = t - 1; A(t) = t^3, B(t) = (t - 1)^3; A(t) = -t^2, B(t) = (t - 1)^2.$$

The case $A(t) = t^3, B(t) = (t - 1)^3$ does not lead to a solution as $N_1(\lambda, t)$ does depend on t . The case $A(t) = -t^2, B(t) = (t - 1)^2$ implies $N_2(\lambda, t) = (\lambda - t)^2$, and hence $\beta_3 = 0$. Finally, in the case $A(t) = t, B(t) = t - 1$ we have $N_2(\lambda, t) = \lambda^2 - 2\lambda + 2\lambda t - t$ (which does not define a solution). The condition that $N_2(\lambda, t)$ divides $N(\lambda, t)$ leads to $\beta_3 = 9\beta_0$ and we get

$$N(\lambda, t) = (\lambda^2 - 2\lambda + 2\lambda t - t) (t^2 - 4\lambda^3 t + 6\lambda^2 t - 4\lambda t + \lambda^4).$$

The function $\lambda(t)$ defined by

$$t^2 - 4\lambda^3t + 6\lambda^2t - 4\lambda t + \lambda^4 = 0 \quad (11)$$

is indeed a solution of \mathbf{PVI}_α , e.g. [3], Table 2, solution 3D. In the case when the dessin corresponding to $\lambda(t)$ is of degree five we conclude that the polynomial $N_2(\lambda, t)$ is linear in λ . By Proposition 2 we get $N_2(\lambda, t) = \lambda - t^2$ which implies $\beta_1 = 0$. To resume, *the curve Γ_β defines a solution, provided that $\beta_0 = \beta_1 = \beta_2 = \beta_3/9$.*

2.7 The face $\beta_0 = \beta_1 = \beta_2 = \beta_3$.

We have

$$N(\lambda, t) = \beta_0(\lambda^2 - 2\lambda + t)(\lambda^2 - 2\lambda t + t)(\lambda^2 - t)$$

and the three algebraic functions defined by $N(\lambda, t) = 0$ are solutions of suitable \mathbf{PVI}_α equations, e.g. [3], Table 2, solutions 2B, 2C, 2A respectively. *The curve Γ_β defines three solutions*

2.8 The face $\beta_3 = 0$.

Recall that in this case $N(\lambda, t) = (\lambda - t)^2 N^0(\lambda, t)$ where

$$N_0(\lambda, t) = \beta_0\lambda^2(\lambda - 1)^2 - \beta_1t(\lambda - 1)^2 + \beta_2(t - 1)\lambda^2.$$

The same arguments as in section 2.3 show that the corresponding dessin d'enfant is of degree one or two.

If the degree is one, then the solution is $\lambda = P(t)$ for some non-constant polynomial P . Therefore $N^0(P(t), t) \not\equiv 0$ and $P(t)$ can not be a solution.

If the degree is two, then $\lambda(t)$ has exactly two ramification points. Without loss of generality we suppose that these points are 0 and ∞ , and as in section 2.3 we conclude that $\lambda(t)$ is defined by $\lambda^2 + 2p(t)\lambda + q(t) = 0$ for some $p, q \in \mathbb{C}[t]$. The polynomial $\lambda^2 + 2p(t)\lambda + q(t)$ divides

$$N_0(\lambda, t) = t(\beta_2\lambda^2 - \beta_1(\lambda - 1)^2) + \beta_0\lambda^2(\lambda - 1)^2 - \beta_2\lambda^2$$

and hence $p(t) = c_1$ and $q(t) = c_2t$ for some constants c_1, c_2 . Without loss of generality we suppose that the ramification points of $\lambda(t)$ are 0 and ∞ and hence the discriminant $4(p^2 - q)$ is a power of t . This implies that $c_1 = 0$. Finally, a direct computation shows that the identity $N^0(\sqrt{-c_2t}, t) \equiv 0$ implies $\beta_2 = 0$ which is not true. *The curve Γ_β does not define a solution.*

2.9 The face $\beta_3 = 0, \beta_0 = \beta_2$.

It is easier to analyze the face $\beta_3 = 0, \beta_1 = \beta_2$, which is equivalent to $\beta_3 = 0, \beta_0 = \beta_2$ after applying the transformation x^2 . Suppose for a moment that $\beta_3 = 0, \beta_1 = \beta_2$. The dessin is of degree at most three, and hence $N^0(\lambda, t)$ is reducible. It follows that $\lambda - c$ divides $N^0(\lambda, t)$ for some constant c . As $N^0(\lambda, t)$ is linear in t , then $\lambda - c$ is deduced from the coefficient of t which equals to $\beta_1(1 - 2\lambda)$. Thus $c = 1/2$ and the condition that $1 - 2\lambda$ divides $N^0(\lambda, t)$ leads to $\beta_0 = 4\beta_1 = 4\beta_2$, in which case

$$N^0(\lambda, t) = \beta_0 (2\lambda - 1) (2\lambda^3 - 3\lambda^2 + t).$$

The function $\lambda(t)$ defined by $(2\lambda^3 - 3\lambda^2 + t) = 0$ is indeed a solution, see [3], Table 2, solutions 5L. Applying the transformation $(x^2)^{-1} = x^2$ of section 2.1 we get the solution (see [3], Table 2, solution 5K

$$\lambda^3 - 3\lambda t + 2t = 0$$

defined by Γ_β with $\beta_3 = 0, \beta_1 = 4\beta_0 = 4\beta_2$. The curve Γ_β defines a solution provided that $\beta_3 = 0, \beta_1 = 4\beta_0 = 4\beta_2$.

2.10 The face $\beta_3 = 0, \beta_0 = \beta_1 = \beta_2$.

The polynomial $N^0(\lambda, t)$ is irreducible and defines a solution, see [3], Table 2, solution 4D.

2.11 The face $\beta_3 = 0, \beta_2 = 0$.

A direct computation shows that the relation $\beta_0\lambda^2 - \beta_1 t = 0$ defines a solution.

2.12 The face $\beta_3 = 0, \beta_2 = 0, \beta_0 = \beta_1$.

The solution is $\lambda^2 = t$.

The results are summarized in Table 1. Theorem 1 is proved.

3 Algebraic solutions of PVI_α and Picard-Fuchs equations

It was noted in the Remark after Theorem 1 that the families 1A, 1B, ..., 1F are Okamoto equivalent to the families 2A, 2B, 2C. To this end we consider

the remaining 23 families $2A - 5L$, see Table 1. To each of them corresponds an affine plane or line in the parameter space $\mathbb{C}^4\{\alpha\}$ which, as we shall prove bellow, is generated by special points α of geometric origin, see Tables 4 and 5. Indeed, observe that exactly the same 23 solutions $2A - 5L$ were already obtained by Doran, see Theorem 2 bellow, by making use of deformations of elliptic surfaces with four singular fibers. The corresponding special values of the parameter α are given in Table 4. The main result of this section is that exactly the same list of solutions can be obtained from deformations of ramified covers of \mathbb{P}^1 with four ramification points. The corresponding values of the parameters α are different and are shown on Table 5, see Theorem 3.

Recall that an elliptic surface is a complex compact analytic surface S with a projection $S \rightarrow \mathbb{P}^1$, such that the general fiber $f^{-1}(z) = \Gamma_z$ is an elliptic curve. Two elliptic surfaces are equivalent, if there is a bi-analytic map compatible with the projections, see [14, Kodaira].

We may suppose that the fiber Γ_z is written in the Weierstrass form

$$\Gamma_z = \{(x, y) \in \mathbb{C}^2 : y^2 = 4x^3 - g_2(z)x - g_3(z)\}$$

and consider the complete elliptic integrals of first and second kind

$$\eta_1 = \int_{\gamma(z)} \frac{dx}{y}, \quad \eta_2 = \int_{\gamma(z)} \frac{xdx}{y}$$

where $\gamma(z) \subset \Gamma_z$ is a continuous family of closed loops (representing a locally constant section $z \mapsto H_1(\Gamma_z, \mathbb{Z})$ of the associated homology bundle). Then η_1, η_2 satisfy the following Picard-Fuchs system (this goes back at least to [6, Griffiths], see [17, Sasai])

$$\Delta(z) \frac{d}{dz} \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} = \begin{pmatrix} -\frac{\Delta'_z}{12} & -\frac{3\delta}{2} \\ -\frac{g_2\delta}{8} & \frac{\Delta'_z}{12} \end{pmatrix} \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} \quad (12)$$

where

$$\Delta(g_2, g_3) = g_2^3 - 27g_3^2.$$

and

$$\delta(z) = 3g_3 \frac{dg_2}{dz} - 2g_2 \frac{dg_3}{dz}.$$

The singular points of the system correspond to singular fibers of the surface. The elliptic surfaces with four singular fibers were classified by [9, Herfurter] who obtained 50 distinct case, but only 5 of them contain an additional

parameter, see Table 3. They lead to non-trivial isomonodromic deformations of the above Picard-Fuchs system with four regular singular points. If we renormalize the singular points to be $0, 1, \infty, t$ then the zero λ of $\delta(z)$, considered as a function in t is a solution of an appropriate \mathbf{PVI}_α equation, see [16] for details. The result is summarized as follows

Theorem 2 [3, Theorem 3.13] *All algebraic solutions $(\lambda(t), \alpha)$ of \mathbf{PVI}_α equation coming from moduli of elliptic surfaces with four singular fibers are shown on Table 1. The corresponding values of α together with the stabilizer of the solution and the \mathbf{PVI}_α equation under the action of the symmetric group S_4 are listed on Table 4.*

The Picard-Fuchs system (12) has infinite monodromy group. We shall deduce similar families of Picard-Fuchs systems having a finite monodromy. Consider a ramified covering $\Gamma \rightarrow \mathbb{P}^1$ of degree three with branching locus consisting of four points, where Γ is a Riemann surface. We choose an affine model $\Gamma^{aff} = \{(x, z) \in \mathbb{C}^2 : f(x, z) = 0\}$ of Γ , where $f(x, z) = 4x^3 - g_2x - g_3$, and $g_2 = g_2(z), g_3 = g_3(z)$ are suitable polynomials. Moreover, without loss of generality, we suppose that the covering $\Gamma \rightarrow \mathbb{P}^1$ is induced from the projection

$$\{(x, z) \in \mathbb{C}^2 : f(x, z) = 0\} \rightarrow \mathbb{C} : (x, z) \mapsto z. \quad (13)$$

Let $x_1(z), x_2(z)$ be two distinct roots of f . Then $\gamma(z) = x_1(z) - x_2(z)$ is a 0-cycle of the fiber $\{x : f(x, z) = 0\}$ and the Abelian integrals above are replaced by the algebraic functions

$$\eta_1(z) = \int_{\gamma(z)} x = x_1(z) - x_2(z), \eta_2(z) = \int_{\gamma(z)} x^2 = x_1^2(z) - x_2^2(z).$$

A straightforward computation shows that η_1, η_2 satisfy the following Picard-Fuchs system (see [5])

$$\Delta(z) \frac{d}{dz} \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} = \begin{pmatrix} \frac{\Delta'_z}{6} & -3\delta \\ -\frac{g_2\delta}{2} & \frac{\Delta'_z}{3} \end{pmatrix} \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}. \quad (14)$$

As before, if we renormalize the system (14) to have singular points at $0, 1, t, \infty$, then the root $\lambda(t)$ of $\delta(z)$ is a solution of a suitable \mathbf{PVI}_α equation, provided that the deformation is isomonodromic. The last property holds, strictly speaking, in the case when the system is non-resonant. In our case it holds too, because the deformation is isoprincipal in the sense of [11]. This

can be also checked by a direct computation. Thus, if we consider the fibration (13) and take for g_2, g_3 the expressions found by Herfurtner, see Table 3, we get the same 23 algebraic solutions shown on Table 1. The corresponding values for α are of course different and are computed in Table 5. In this way we prove

Theorem 3 *The algebraic solutions $(\lambda(t), \alpha)$ of \mathbf{PVI}_α equation coming from deformations of the covering (13) with g_2, g_3 as on the Herfurtner list, Table 3, are shown on Table 1. The corresponding values of α together with the stabilizer of the solution and the \mathbf{PVI}_α equation under the action of the symmetric group S_4 are listed on Table 5.*

Remark 3 *As we already noted, the monodromy group of the Picard-Fuchs system (14) is finite. Particular cases of this system, in a more or less explicit way, were considered by many authors, e.g. [2, Boalch], [4, Dubrovin-Mazzocco], [7, 8, Hitchin], [13, Kitaev].*

To this end we present, for convenience of the reader, one example of such computation. Using the Picard-Fuchs system (14), the Abelian integral of first kind η_1 satisfies the following equation

$$p_0(z, a)\eta_1'' + p_1(z, a)\eta_1' + p_2(z, a)\eta_1 = 0 \quad (15)$$

where

$$\begin{aligned} p_0(z, a) &= 144\delta\Delta^2 \\ p_1(z, a) &= 144\Delta\left(\delta\frac{d\Delta}{dz} - \Delta\frac{d\delta}{dz}\right) \\ p_2(z, a) &= 12\delta\frac{d^2\Delta}{dz^2} - 216\delta^3g_2 - 12\Delta\frac{d\delta}{dz}\frac{d\Delta}{dz} - \delta\left(\frac{d\Delta}{dz}\right)^2. \end{aligned}$$

Consider, for instance, the deformation 2 from the Herfurtner list (Table 3)

$$\begin{aligned} g_2 = g_2(z, a) &= 3z^3(z + a) \\ g_3 = g_3(z, a) &= z^5(z + 1). \end{aligned}$$

We have

$$\begin{aligned} \Delta &= \Delta(g_2, g_3) = 27z^9((3a - 2)z^2 + (3a^2 - 1)z + a^3) \\ \delta &= \delta(z, a) = -3z^7((3a - 2)z + a). \end{aligned}$$

name	deformation
1	$g_2(z, a) = 3(z - 1)(z - a^2)^3$ $g_3(z, a) = (z - 1)(z - a^2)^4(z + a)$
2	$g_2(z, a) = 12z^2(z^2 + az + 1)$ $g_3(z, a) = 4z^3(2z^3 + 3az^2 + 3az + 2)$
3	$g_2(z, a) = 12z^2(z^2 + 2az + 1)$ $g_3(z, a) = 4z^3(2z^3 + 3(a^2 + 1)z^2 + 6az + 2)$
4	$g_2(z, a) = 3z^3(z + a)$ $g_3(z, a) = z^5(z + 1)$
5	$g_2(z, a) = 3z^3(z + 2a)$ $g_3(z, a) = z^4(z^2 + 3az + 1)$

Table 3: The Herfurtner list of "deformable" elliptic surfaces with four singular fibers

The Picard-Fuchs equation (15) takes the form

$$\begin{aligned}
& 144z^2((3a - 2)z + a)((3a - 2)z^2 + (3a^2 - 1)z + a^3)^2 \eta_1'' \\
& + 144z((3a - 2)z^2 + (3a^2 - 1)z + a^3)(3(3a - 2)^2 z^3 \\
& + 2(3a - 2)(3a - 1)(a + 1)z^2 + a(3a^3 + 7a^2 - 3)z + 2a^4) \eta_1' \\
& + [135(3a - 2)^3 z^5 + (3a - 2)^2(468a^2 + 267a - 164)z^4 \\
& + 2(3a - 2)(189a^4 + 522a^3 - 48a^2 - 208a + 10)z^3 \\
& - 2a(270a^5 - 1269a^4 + 252a^3 + 460a^2 - 70)z^2 \\
& - a^4(243a^3 - 666a^2 + 176)z + 27a^7] \eta_1 = 0
\end{aligned}$$

and has four regular singular points at ∞ and the roots of $(3a - 2)z^2 + (3a^2 - 1)z + a^3$ (the roots of Δ), as well one apparent singularity at the root of $(3a - 2)z + a$ (which is a root of δ). Re-normalizing the singular points to $0, 1, t, \infty$ we get

$$\lambda = \frac{a^2 - a + 1}{a^2(2 - a)}, t = \frac{2a - 1}{a^3(2 - a)}, a \in \mathbb{C}. \quad (16)$$

The parameter a defines an algebraic isomonodromic deformation of the Picard-Fuchs equation (15) with Riemann schema

$$\begin{pmatrix} 0 & 1 & t & \lambda & \infty \\ -\frac{3}{4} & \frac{1}{4} & \frac{1}{4} & 0 & \frac{5}{4} \\ -\frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} & 2 & \frac{3}{4} \end{pmatrix}.$$

Therefore the algebraic function $\lambda = \lambda(t)$ determined implicitly by (16) is an algebraic solution of \mathbf{PVI}_α equation with

$$\alpha_0 = \frac{1}{8}, \quad \alpha_1 = \frac{1}{8}, \quad \alpha_2 = \frac{1}{8}, \quad \alpha_3 = \frac{1}{8}.$$

(see [10] for details). Eliminating a from (16) we get

$$\lambda^4 - 2t\lambda^3 - 2\lambda^3 + 6t\lambda^2 - 2t^2\lambda - 2t\lambda + t^3 - t^2 + t = 0$$

which is an equation for the solution 4A with $\alpha = (1/8, 1/8, 1/8, 1/8)$. Together with the Doran's point $\alpha = (0, 1/8, 0, 0)$, this implies that the solution $\lambda(t)$ satisfies also the implicit equation (1)

$$1 + \frac{t-1}{(\lambda-1)^2} - \frac{t(t-1)}{(\lambda-t)^2} = 0$$

corresponding to the affine line through $(1/8, 1/8, 1/8, 1/8)$ and $(0, 1/8, 0, 0)$ described in Table 1, 4A. The solution (16) with $\alpha = (1/8, 1/8, 1/8, 1/8)$ was found by Hitchin [7, section 6.1], [8, (34)].

Remark 4 *If we repeat the same computation, but making use of the Picard-Fuchs system (12) then of course we obtain the same algebraic solution but with $\alpha = (1/8, 0, 0, 0)$. This value has been erroneously computed by Doran [3] to be $(1/18, 0, 0, 0)$. This led him to the wrong conclusion that the solution 4C is equivalent by an Okamoto transformation to the "cubic" solution B_3 of Dubrovin-Mazzocco [4, p.140] with $\alpha = (25/18, 0, 0, 0)$, see [3, Remark 7]. As the Okamoto transformations of \mathbf{PVI}_α act within the ring $\mathbb{Z}[1/2, \sqrt{2\alpha_1}, \sqrt{2\alpha_1}, \sqrt{2\alpha_1}, \sqrt{2\alpha_1}]$, then no solution of $\mathbf{PVI}_{(25/18, 0, 0, 0)}$ is equivalent to a solution of $\mathbf{PVI}_{(1/8, 0, 0, 0)}$. M. Mazzocco kindly informed us for a missprint in the formula for the B_3 -solution, [4, p.140]. The corrected formula is reproduced in [3, formula (3.1)].*

Stabilizer of the solution	Name of the solution	\mathbf{PVI}_α equation	Stabilizer of \mathbf{PVI}_α equation
D_4	$2A$	$(0, 0, \frac{1}{18}, \frac{1}{18})$ $(\frac{1}{18}, \frac{1}{18}, 0, 0)$	$S_2 \times S_2$
	$2B$	$(\frac{1}{18}, 0, \frac{1}{18}, 0)$ $(0, \frac{1}{18}, 0, \frac{1}{18})$	
	$2C$	$(\frac{1}{18}, 0, 0, \frac{1}{18})$ $(0, \frac{1}{18}, \frac{1}{18}, 0)$	
D_4	$2A$	$(0, 0, 0, 0)$	S_4
	$2B$		
	$2C$		
S_3	$3A$	$(0, 0, 0, 0)$	S_4
	$3B$		
	$3C$		
	$3D$		
S_3	$4A$	$(0, \frac{1}{8}, 0, 0)$ $(0, 0, \frac{1}{8}, 0)$ $(\frac{1}{8}, 0, 0, 0)$ $(0, 0, 0, \frac{1}{8})$	S_3
	$4B$		
	$4C$		
	$4D$		
S_2	$5A$	$(0, \frac{1}{18}, 0, 0)$	S_3
	$5B$		
	$5C$		
	$5D$	$(0, 0, \frac{1}{18}, 0)$	
	$5E$		
	$5F$		
	$5G$	$(\frac{1}{18}, 0, 0, 0)$	
	$5H$		
	$5I$		
	$5J$		
$5K$	$(0, 0, 0, \frac{1}{18})$		
$5L$			

Table 4: Solutions $(\lambda(t), \alpha)$ of \mathbf{PVI}_α equations related to the Picard-Fuchs system (12).

Stabilizer of the solution	Name of the solution	\mathbf{PVI}_α equation	Stabilizer of \mathbf{PVI}_α equation
D_4	$2A$	$(\frac{1}{8}, \frac{1}{8}, \frac{1}{18}, \frac{1}{18})$ $(\frac{1}{18}, \frac{1}{18}, \frac{1}{8}, \frac{1}{8})$	$S_2 \times S_2$
	$2B$	$(\frac{1}{18}, \frac{1}{8}, \frac{1}{18}, \frac{1}{8})$ $(\frac{1}{8}, \frac{1}{18}, \frac{1}{8}, \frac{1}{18})$	
	$2C$	$(\frac{1}{18}, \frac{1}{8}, \frac{1}{8}, \frac{1}{18})$ $(\frac{1}{8}, \frac{1}{18}, \frac{1}{18}, \frac{1}{8})$	
D_4	$2A$	$(\frac{1}{8}, \frac{1}{8}, \frac{1}{2}, \frac{1}{2})$ $(\frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8})$	$S_2 \times S_2$
	$2B$	$(\frac{1}{2}, \frac{1}{8}, \frac{1}{2}, \frac{1}{8})$ $(\frac{1}{8}, \frac{1}{2}, \frac{1}{8}, \frac{1}{2})$	
	$2C$	$(\frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{1}{2})$ $(\frac{1}{8}, \frac{1}{2}, \frac{1}{2}, \frac{1}{8})$	
S_3	$3A$	$(\frac{1}{8}, \frac{9}{8}, \frac{1}{8}, \frac{1}{8})$	S_3
	$3B$	$(\frac{9}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8})$	
	$3C$	$(\frac{1}{8}, \frac{1}{8}, \frac{9}{8}, \frac{1}{8})$	
	$3D$	$(\frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{9}{8})$	
S_3	$4A$	$(\frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8})$	S_4
	$4B$		
	$4C$		
	$4D$		
S_2	$5A$	$(\frac{1}{2}, \frac{1}{18}, \frac{1}{8}, \frac{1}{8})$	S_3
	$5B$	$(\frac{1}{8}, \frac{1}{18}, \frac{1}{2}, \frac{1}{8})$	
	$5C$	$(\frac{1}{8}, \frac{1}{18}, \frac{1}{8}, \frac{1}{2})$	
	$5D$	$(\frac{1}{2}, \frac{1}{8}, \frac{1}{18}, \frac{1}{8})$	
	$5E$	$(\frac{1}{8}, \frac{1}{2}, \frac{1}{18}, \frac{1}{8})$	
	$5F$	$(\frac{1}{8}, \frac{1}{8}, \frac{1}{18}, \frac{1}{2})$	
	$5G$	$(\frac{1}{18}, \frac{1}{8}, \frac{1}{2}, \frac{1}{8})$	
	$5H$	$(\frac{1}{18}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8})$	
	$5I$	$(\frac{1}{18}, \frac{1}{8}, \frac{1}{8}, \frac{1}{2})$	
	$5J$	$(\frac{1}{8}, \frac{1}{8}, \frac{1}{2}, \frac{1}{18})$	
	$5K$	$(\frac{1}{8}, \frac{1}{2}, \frac{1}{8}, \frac{1}{18})$	
	$5L$	$(\frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{1}{18})$	

Table 5: Solutions $(\lambda(t), \alpha)$ of \mathbf{PVI}_α equations related to the Picard-Fuchs system (14).

References

- [1] B. Ben Hamed, Connexion de Gauss-Manin et Solutions Algébriques de P_{VI} , mémoire de DEA, Université de Toulouse III, 2003, see also arxiv.org/abs/math.CA/0405308.
- [2] Ph. Boalch, From Klein to Painlevé via Fourier, Laplace and Jimbo, Proc. London Math. Soc. (3) **90** (2005) 167-208
- [3] Charles F. Doran, Algebraic and geometric isomonodromic deformations, J. Diff. Geometry, **59** (2001) 33-85.
- [4] B. Dubrovin, M. Mazzocco, Monodromy of certain Painlevé-VI transcendents and reflection groups, Invent. Math. **141**, 55-147 (2000).
- [5] L. Gavrilov, H. Movasati, The infinitesimal 16th Hilbert problem in dimension zero, arxiv.org/abs/math.CA/0507061 (2005).
- [6] Griffiths, P.A.; The residue calculus and some transcendental results in algebraic geometry, I, Proc. Nat. Acad. Sci. U.S.A., **55** (1966), 1303-1309.
- [7] N. J. Hitchin, Poncelet polygons and the Painlevé equations, in *Geometry and Analysis*, (S. Ramanan, ed.) Oxford University Press, Bombay, 1995.
- [8] N. J. Hitchin, A lecture on the octahedron, Bull. London. Math. Soc. **35** (2003) 577-600.
- [9] S. Herfurtner, Elliptic surfaces with four singular fibres. Math. Ann. **291** (1991), no. 2, 319–342.
- [10] K. Iwazaki, H. Kimura, S. Shimomura, M. Yoshida, *From Gauss to Painlevé*, Aspects of Mathematics, Vieweg 1991.
- [11] V. Katsnelson, D. Volok, Deformations of fuchsian systems of linear differential equations and the Schlesinger system, preprint, Weizmann Institute of Science, 2005.
- [12] F. Kirwan, *Complex Algebraic Curves*, London Math. Soc. Student Texts **23**, Cambridge University Press, 1992.

- [13] A.V. Kitaev, Grothendieck's dessins d'enfants, their deformations, and algebraic solutions of the sixth Painlevé and Gauss hypergeometric equations, *Algebra i Analiz* **17** (2005), no. 1, 224–275.
- [14] K. Kodaira, On compact analytic surfaces. II, *Ann. of Math. (2)* **77** (1963), 563–626.
- [15] Yu. I. Manin, Sixth painlevé equation, Universal elliptic curve and Mirror of \mathbb{P}^2 , *Amer. Math. Soc. Transl. (2)* Vol. **186**, 1998 p. 131-151.
- [16] K. Okamoto, Studies on the Painlevé Equations I, *Ann. Mat. Pura Appl.*, **14**(1987) 337-381.
- [17] Sasai T., Monodromy representation of homology of certain elliptic surfaces, *J. Math. Soc. Japan*, **26**, No2 (1974) 296-305.
- [18] L. Schneps, Dessins d'enfants on the Riemann sphere. The Grothendieck theory of dessins d'enfants (Luminy, 1993), 47–77, *London Math. Soc. Lecture Note Ser.*, **200**, Cambridge Univ. Press, Cambridge, 1994.